The adaptive landscapes of three global *Escherichia coli* transcriptional regulators

Authors

Cauã Antunes Westmann[1,2], Leander Goldbach[1,2], Andreas Wagner[1,2,3*]

[1] – Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstrasse 190, Zurich CH-8057, Switzerland

[2] – Swiss Institute of Bioinformatics, Quartier Sorge-Batiment Genopode, 1015 Lausanne,

Switzerland

[3] – The Santa Fe Institute, Santa Fe, NM 87501, USA

* - Corresponding author: andreas.wagner@ieu.uzh.ch

Abstract

The evolution of new gene regulation is an important source of evolutionary adaptations and innovations, especially when organisms encounter new environments. At its heart is the process by which new and strong DNA binding sites of transcription factors (TFs) originate in evolution. Here we study the potential of Darwinian evolution to create strong binding sites for the three *Escherichia coli* global transcription factors CRP, Fis, and IHF. Using a massively parallel reporter assay, we measure for each TF the ability of more than 30,000 binding sites to regulate gene expression. We use the resulting data to map the adaptive landscape of each TF. We find that all three landscapes are rugged, epistatic, and harbor multiple peaks. The highest peaks are widely scattered throughout the landscape, indicating that strong regulation can be achieved by very different binding sites. Landscape ruggedness does not prevent the evolution of strong regulation, because more than 10% of evolving populations can attain one of the highest peaks. Adaptive evolution starting from the same DNA sequence can attain more than one high peak, and some high peaks are more likely to be reached than others. Our experiments show that de novo adaptive evolution of new gene regulation is feasible. It is also subject to a blend of chance, historical contingency, and evolutionary biases that favor some peaks and evolutionary paths over others.

Keywords: adaptive landscapes, navigability, gene regulation, bacteria, massively parallel reporter assay, global regulators

INTRODUCTION

Transcriptional regulation controls the ability of RNA polymerase to initiate the transcription

of a gene into mRNA[1]. It is crucial in the life of all organisms, orchestrating the expression of

genes in response to environmental cues and cellular states[2,3]. Transcriptional regulation is

mediated by transcription factors (TFs), proteins that bind DNA near a gene at short DNA

words known as transcription factor binding sites (TFBSs). The binding of a TF to its binding

site on DNA can either hinder or facilitate transcription initiation by RNA polymerase[1,4,5].

The stronger the binding of a TF to its TFBS is, the more strongly the TF can regulate a

nearby gene[6–9]. In bacteria, TFs operate within a hierarchically organized gene regulatory

network. At the bottom of this hierarchy are *local* TFs that regulate the expression of one or

few genes and modulate specific biological processes [10–12] . At the top are *global* TFs that

may regulate hundreds of genes and control many cellular functions [11,13–15]. Consistent with

their broader role, global regulators are typically more highly expressed and bind to a broader

range of TFBSs than local regulators[12].

Both TFs and their TFBSs evolve, but TFBSs evolve more rapidly. The reason is that a

mutation in a TF can affect the expression of many genes, whereas a mutation in a TFBS may

affect only one gene and is less likely to be deleterious [13,16–18]. A special case of TFBS

evolution is the evolution of a strong TFBS from a DNA word with weak or no regulatory

activity. Such *de novo* evolution of TFBSs can create new regulatory interactions and change

the structure of gene regulatory networks[19–22]. Unfortunately, we know little about the ability

of Darwinian evolution to create TFBSs de novo. A strong TFBS may have to arise from a

non-binding site through an evolutionary path of multiple mutational steps. Darwinian

evolution can favor this process only if strong binding is adaptive and if each mutational step

in a path increases binding strength, i.e., if the path is *accessible* to Darwinian evolution. To

find whether such paths exist may require the analysis of multiple paths. This is challenging

because sequence space contains an astronomical number of potential TFBSs for any one TF.

The number of evolutionary paths to strong transcriptional regulation is thus also

astronomical. For each such path, the strength of each TFBS along the path has to be

measured experimentally[23–26].

Building on our previous work on a local TF[27], we take a step to address this challenge for

three global regulators in *Escherichia coli*. Specifically, we perform three independent and

massively parallel experiments[23] using a plasmid-based expression system for each TF. Our

synthetic biology approach allows us to explore the regulatory effects of tens of thousands of

TFBS variants in a high-throughput manner. We quantify how strongly each of more than

30,000 binding sites for each TF can regulate the expression of a nearby reporter gene.

The first TF we study is the cAMP receptor protein (CRP). In the absence of glucose, CRP

modulates the expression of genes mostly involved in carbon metabolism. It allows *E. coli* to

efficiently switch between sources of carbon and energy [28–31]. The second TF is the factor for

inversion stimulation (Fis), which helps to regulate the expression of genes involved in

growth phase transitions. It also modulates the supercoiling of DNA[32–34] and influences DNA

replication, recombination, and repair[32–34]. The third factor is the integration host factor

(IHF). IHF regulates genes involved in stress responses and stationary phase survival[35,36].

Similar to Fis, it is involved in DNA compaction, replication, and recombination, but also in

the assembly of complex nucleoprotein structures[35,36]. We chose these factors because they

are the most global regulators in *E. coli*, and they are diverse, belonging to different protein

families.

We use the regulatory data for each TF's binding sites to map a genotype-phenotype

landscape for these binding sites [37,38]. Such a landscape is a conceptual analog of a physical

landscape, where each location corresponds to one genotype in sequence space, and the

elevation at this location corresponds to a phenotype – the ability to regulate gene expression

[6,27,39]. One of our two main aims is to provide the first characterization of such landscapes for

global bacterial regulators, and to find out whether these landscapes are different or similar.

When strong regulation of a gene is adaptive, a regulatory landscape becomes an adaptive

landscape or fitness landscapes [38,40–42]. Our second aim is to understand how populations

would evolve on each of our three landscapes when they are viewed as fitness landscapes.

Specifically, we study how evolving populations would traverse each landscape through

individual mutational steps that change a TFBS its ability to regulate a gene via its cognate

TF. A peak in such a landscape is a TFBS conveying stronger regulation than all neighboring

TFBSs in sequence space. If such a landscape is rugged (has multiple peaks), natural

selection alone may not enable a population to discover the highest peaks, i.e., the strongest

TFBSs. The reason is that the peaks may be separated by valleys of low regulation strength

that cannot be traversed by natural selection alone [43–45 38]. In other words, high peaks may not

be easily accessible through Darwinian evolution – they may be reachable by few or no

evolutionary paths of single DNA mutations where each mutational step increases TFBS

strength[6]. One factor that can reduce peak accessibility is epistasis – non-additive effects of

two or more mutations on a phenotype. In addition, epistasis can reduce the predictability of

evolutionary trajectories towards a peak [37,46].

To accomplish both aims, we first studied the topography of the three landscapes, and then

simulated the dynamics of evolving populations on them. All three landscapes are highly

rugged. They contain more than 2,000 peaks that are scattered through genotype space and

are also rife with epistatic interactions. Despite these features, evolving populations can reach

the strongest TFBSs in all three landscapes. Which of several high peak is reached is

contingent on chance events during adaptive evolution.

RESULTS

Landscape mapping

We constructed a plasmid system (**Supplementary Table S1**, **Supplementary Figures S1-S2**

and Supplementary Methods 2-3) to inducibly express each of the three global TFs CRP,

Fis, and IHF in *E. coli*. We expressed each factor in a host strain in which its encoding

chromosomal gene had been knocked out (*Δcrp*, *Δfis* and *Δihf,* see **Supplementary Table**

S2). This ensures that each TF is expressed only upon induction. Expression of each TF can

be induced with anhydrotetracycline (atc), which allowed us to control TF expression levels

and measure how the binding of the TF to DNA can influence transcriptional regulation by

measuring the expression of a green fluorescent reporter gene (*gfp*) (**Figure 1a**). In our

plasmid system, stronger binding of a TF to DNA leads to lower GFP expression. The reason

is that the TF's binding site is located directly upstream of the *gfp* gene, and TF-DNA binding

thus downregulates GFP expression by interfering with *gfp* transcription. Our three plasmid-

strain pairs attained similar cell densities during late exponential or early stationary phase

(**Supplementary Figure S3**) where we performed all our measurements.

For each plasmid expressing one of our three TFs, we cloned an entire library of TFBSs

upstream of the *gfp* reporter. Specifically, in this library, we randomized each of the eight

most important (information-rich) base pair positions within the TFBS (see **Supplementary**

Methods 4 and Supplementary Table S3). We obtained these most information-rich

positions for each TF from the alignment of known TFBSs in the RegulonDB database[47].

Each library comprised 4^8=65,536 unique sequences. We constructed three replicates of each

library (**Supplementary Methods 4 and 5**).

We then mapped these TFBS genotypes to their respective regulatory phenotypes using a

well-established technique known as sort-seq[23,39,48–50] (**Figure 1b**), which combines

fluorescence-activated cell sorting (FACS) with high-throughput sequencing (**Figure 1c**). In

sort-seq, one first sorts cells into multiple fluorescence "bins" depending on the level of GFP

expression. A cell's GFP fluorescence serves as a proxy for GFP expression levels[51] and

quantifies how strongly the TFBS library member in this cell can regulate GFP expression

(**Methods**, **Supplementary Methods 5**).

The results of our sort-seq experiments are three maps – one for each transcription factor –

from each of more than 30,000 genotypes (TFBS variants) to regulatory phenotypes

(regulation strength). One can view each map as a genotype-phenotype landscape that we

also call a regulatory landscape[8,9,27,39,52–56] (**Figure 1d**). Whenever strong regulation entails

high fitness, such a landscape is also an adaptive landscape or fitness landscape[6,38]. For the

purpose of analyzing the landscape quantitatively, we represent it as a network of genotypes

(TFBS variants). Each node in this network corresponds to a genotype. Edges link

neighboring genotypes, which differ in a single nucleotide (**Figure 1d**).

Figure 1. Mapping TFBS genotype-phenotype landscapes. a-c) Sort-Seq procedure.

We utilized a plasmid-based fluorescence reporter system followed by sort-seq to map TFBS genotype-phenotype landscapes. **a. Library generation.** For generating our *E. coli* libraries, we cloned TFBS sequence variants (2^8 = 65,536), represented as black circles, into our plasmid, between a sigma70 constitutive promoter (shown as a black right-facing arrow) and a *gfp* gene (shown as a green right-facing arrow), to measure transcriptional repression through fluorescence intensity. When a TF binds to the TFBS, it blocks RNA polymerase activity through steric hindrance, reducing *gfp* transcription. Thus, lower fluorescence levels (light green) represent stronger TF-TFBS binding. The TFBS variants produce a range of regulation strengths resulting, in variable green fluorescence intensities in bacterial cells (green-colored rectangles). **b. Sorting procedure.** We sorted libraries into expression bins based on fluorescence intensity using a fluorescence-activated cell sorting (FACS). **c. Sequencing.** We then sequenced the TFBSs from each bin and mapped their sequence (genotype) to their respective regulation strengths, depicted through a color gradient from

dark green (low-affinity TFBS, high fluorescence) to light green (high-affinity TFBS, low

fluorescence). **d) Genotype-phenotype mapping.** To construct a regulatory landscape, we

connected TFBS genotypes (colored circles) that differed by a single nucleotide via edges,

thereby establishing an interconnected genotype network. Each genotype conveys a

regulatory phenotype (strength of regulation, heatmap colors), which can be viewed as the

elevation dimension (z-axis) in a landscape.

Landscapes exhibit diverse regulation strengths and distribution breadths

To evaluate the ability of our library to regulate gene expression, we first measured the

distribution of GFP fluorescence intensities across the bins in two conditions, i.e., in the

presence or absence of the TF (with or without the atc inducer, **Supplementary figures S4-**

S6). In the presence of the TF, GFP expression was lower on average and showed a broader

distribution than in the absence of the TF (**Supplementary figures S4-S6**). This indicates that

the TFBSs in each library can indeed downregulate GFP expression, but some library variants

convey stronger regulation than others, hence the broader fluorescence distribution.

We then pooled barcoded DNA sequences extracted from cells in each bin and biological

replicate. We sequenced at least 250 unique TFBS genotypes from each bin, with an average

of 3992.3±4293.9 unique sequences per bin for the three TFs (as detailed in **Methods**,

Supplementary Methods 7, and **Supplementary Figures S4-S6**). The resulting sequences

covered 95%, 90%, and 93% of the total library sizes (N = 65,536 genotypes) for CRP, Fis

and IHF, respectively. To ensure the reliability of our data, we excluded sequences with low

read coverage and sequences that were not present in all triplicates (**Methods,**

Supplementary Methods 7.3). This quality filtering step resulted in library sizes of 31,975

genotypes for CRP (49% of all 4^8 genotypes), 43,222 genotypes for Fis (66%), and 41,325

genotypes for IHF (63%). The correlation in read counts for each variant across replicates

was very high for this quality-filtered data (Pearson's R= 0.98-0.99) (**Supplementary**

Figures S7-S9). We used this data for all further analyses.

The fluorescence and sequence data from each bin allowed us to quantify the variant's ability

to regulate gene expression for each TFBS library variant. We refer to the resulting metric as

the *regulation strength S* conveyed by the variant (**Methods**, **Supplementary Methods 7.2**,

Figure 1d). It ranges between S=0 (no regulation) to S=1 (strongest regulation among all

variants in the library). Although our experiments directly quantify expression regulation, and

not the affinity or binding strength of a TF to a TFBS variant, we also use *binding strength* as

a proxy for regulation strength, because TF-DNA binding is necessary for regulation[1,4,11,57–59].

To each of our three TFs, we also assigned a reference TFBS that is naturally occurring and

conveys strong regulation by the TF, as proven by previous experimental work[31,60–62]. We

refer to this TFBS as the wild-type (WT_{CRP}[60,63], WT_{Fis}[61,64] and WT_{IHF}[62,65], **Supplementary**

Table S4). We refer to TFBSs with regulation strengths below and above the wild-type

(WT_{CRP}: $S_{WT}=0.71$, WT_{Fis}: $S_{WT}=0.97$, WT_{IHF}: $S_{WT}=0.95$) as weak and strong, respectively.

We observed a broad range of regulation strengths S for each TF landscape, with varying

spreads. The CRP landscape exhibited the lowest average regulation strength and a narrower

distribution of S compared to Fis and IHF, which had similar distributions (mean±s.d.,

0.37±0.1 for CRP, 0.57±0.13 for Fis, and 0.52±0.14 for IHF; see **Figure 2a, Supplementary**

Figure S10). Next, we analyzed the strongly regulating TFBSs to identify nucleotides that

may be particularly frequent and thus, potentially important for strong regulation (**Figure**

2b). We discovered a moderate association between the most frequent nucleotides

(highlighted in yellow in **Figure 2b**) and the most informative nucleotides from the available

position weight matrices (PWMs) for these TFs[66]. (Pearson correlation coefficients: R=0.51,

R=0.43, R=0.47 for Fis, CRP and IHF, respectively, with p-values smaller than 10^{-16},

rejecting the null hypothesis of an absence of association). This observation suggests that our

logos capture different information compared to available PWMs. This is expected, because

our approach allows us to filter sequences by regulatory strength thresholds to construct our

matrices, unlike the traditional method of aligning genomic TFBSs without considering their

regulation strengths[67,68].

To further validate the data from our sort-seq experiments, we isolated cells harboring 10

different TFBS variants from each of 13 bins of each library (i.e., 10×13=130 variants per

library), and determined their regulation strength more directly by quantifying GFP

expression with a microplate reader (**Methods** and **Supplementary Figure S11**). This

comparison validates the sort seq approach by revealing a strong association of the two

independent quantifications of regulation strength (Pearson's R=-0.81 for CRP, R=-0.74 for

Fis, and R=-0.73 for IHF). (**Methods** and **Supplementary Figure S11**).

All three regulatory landscapes are highly rugged

The study of our landscapes in a network framework (**Figure 1d**) can help to quantify

different aspects of landscape topography (**Supplementary Table S5**). One of them is the

ruggedness of each landscape. It can be quantified by the number of peaks[69–72]. In the

network framework, a peak is a TFBS whose neighbors are all weaker regulators (with lower

S) than the TFBS itself. We find that all three landscapes are highly rugged, with 2,154,

2,312, and 2,453 peaks for the CRP, Fis, and IHF landscapes, respectively (**Figure 2c,**

Supplementary Figure S12, Supplementary Table S5). Not surprisingly, peak genotypes

generally are stronger regulators than non-peak genotypes (**Figure 2c**). Only a small fraction

of peak genotypes regulate expression more strongly than the wild-type (**Figure 2c**; 61 for

CRP, 172 for Fis, and 199 for IHF). We refer to such peaks as high peaks and distinguish

them from low peaks ($S<S_{WT}$).

We compared the ruggedness of our landscapes with that of a well-established theoretical

model of uncorrelated random landscapes, in which each sequence is assigned a fitness at

random from the same fitness distribution, and neighbors have uncorrelated fitness values[72–74]

Such landscapes are maximally rugged [72–74]. We created 100 uncorrelated random landscapes

for each of our three TF landscapes by randomly shuffling the measured regulation strengths

among all genotypes (**Supplementary Methods 7.6**). The number of peaks in our TF

landscapes lies within 93%, 96%, and 98% percent of that of a maximally rugged random

landscape, which, on average [mean±s.d.], has 2,308±133, 2,405±89 and 2,373±97 peaks for

CRP, Fis and IHF, respectively. This analysis underlines that our landscapes are indeed highly

rugged.

Because we use only quality-filtered genotype data, our landscapes lack regulatory

information for about 40% of the 4^8 genotypes. While we cannot exclude that this

undersampling of genotypes has led to systematic biases in our determination of landscape

ruggedness, we note that the sampling of landscape genotypes by our experiments was not

strongly biased with respect to regulation strength. Specifically, when we analyzed the

relative connectivity of genotypes in our landscapes – the fraction of each genotype's 24

possible neighbors for which our experiments yielded regulatory data – we found that it is

only weakly correlated with regulation strength (R=-0.1, -0.1, 0.01 for the CRP, Fis, and IHF

landscapes, **Figures S13-S15**). Similarly, the relative connectivity of peak genotypes is only

weakly correlated with their regulation strength (R=-0.05, -0.04, 0.06 for the CRP, Fis, and

IHF landscapes). Furthermore, the prevalence of sign epistasis (**Supplementary Table S5**)

supports the notion that our landscapes are indeed rugged (see **Supplementary Methods 7.5**

for further details on epistasis and its evolutionary consequences).

Figure 2. a. Genotypes in each landscape vary broadly in their regulation strength. The

violin plots show the distribution of regulation strengths S (vertical axis) for each TF

landscape (horizontal axis). The width of a plot at a given value of S represents the frequency

of TFBSs at this value. The vertical length of the box in each violin plot covers the range

between the first and third quartiles (IQR). The horizontal line within the box represents the

median value, and whiskers span 1.5 times the IQR. The white circle shows the regulation

strength of the wild type for each landscape (CRP: 0.71, Fis: 0.97, IHF: 0.95). The landscape

sizes are as follows. CRP: N=31,975 TFBS variants; Fis: N= 43,222; IHF: N= 41,325. **b.**

Peak genotypes are stronger regulators than non-peak genotypes. Dual violin plots show

the distribution of regulation strength (vertical axis) for the three TF landscapes (horizontal axis), stratified by non-peak genotypes (grey, CRP, N=29,821, Fis, N= 40,910, IHF, N= 38,872), and peak genotypes (red, CRP, N= 2,154, Fis, N= 2,312, IHF, N= 2,453). The black tick-mark in each plot indicates the mean regulation strength of both non-peak and peak genotypes taken together (mean ± standard deviation, CRP: 0.37 ±0.1, Fis: 0.57±0.13, IHF: 0.52±0.14). The white circle on each plot marks the regulation strength of the wild-type (CRP: 0.71, Fis: 0.97, IHF: 0.95). **c. Sequence logos and nucleotide frequency matrices for strong CRP, Fis and IHF binding sites.** Each sequence logo [67,68] is based on an alignment of TFBSs with greater regulation strength than the wild-type for each of the three TFs IHF, Fis, and CRP. Each logo also shows the non-varying position (grey) of the TFBS genotype from which our libraries were created. In each logo, the height of each letter at each TFBS position indicates the information content at that nucleotide position – the taller the letter, the more frequent the nucleotide is in strongly regulating TFBSs [67,68]. Similar information is conveyed by the frequency matrices displayed as heat maps below each logo. They represent the variability of each nucleotide at each position (horizontal axis) through a color gradient (see color legend). Tall letters in the sequence logo and yellow letters in the frequency matrix indicate frequent, and thus likely important nucleotides for strong regulation.

Landscape peaks are widely scattered in genotype space.

In a rugged adaptive landscape, reaching high fitness peaks can be challenging, because such peaks are separated from other genotypes by valleys of low fitness that cannot be traversed by natural selection alone[38]. If natural selection favors strong gene regulation, the ruggedness of our regulatory landscapes may thus present a challenge for adaptive evolution. To better understand the potential magnitude of this challenge, we next analyzed our landscapes' topography in greater detail. We began by studying the distribution of peaks in genotype

space. If a landscape's highest peaks are widely scattered through genotype space, then they

may be accessible via fitness-increasing evolutionary paths from diverse non-peak genotypes.

This may facilitate adaptive evolution compared to a landscape where peaks are clustered in a

small region of genotype space.

For each of our three landscapes, we determined the genetic distance between peaks, i.e., the

minimum number of mutations needed to transition from one peak to another, regardless of

their effect on regulation strength. We compared the distribution of these distances to the

distribution of genetic distances for an equal number of randomly selected non-peak variants.

In all three landscapes, the distances between peaks are almost indistinguishable from those

of random genotypes, differing on average by fewer than 0.1 mutational steps. In other

words, peaks are about as widely dispersed in each landscape as random genotypes

(**Supplementary Figure S16**). This is the case for both low and high peaks (**Supplementary**

Figures S17-S18). A principal component analysis further underscores this dispersion

(**Supplementary Methods 7.7, Supplementary Figures S19-S21**).

Accessible paths to a peak are often not the shortest possible paths

Next, we focused on mutational paths to peaks that are evolutionarily accessible, i.e., a series

of mutational steps where each step increases regulation strength. Specifically, we

enumerated all accessible paths that exist from each non-peak genotype to each peak

genotype for all three landscapes. We found that these paths are generally longer than the

shortest genetic distance between a non-peak and the peak genotypes. Specifically, the mean

length of accessible paths exceeded the shortest distance by 1.5, 1.8 and 1.8 steps for the

CRP, Fis and IHF landscapes (**Supplementary Figure S22**). In other words, accessible

mutational paths are often not the most direct paths.

The existence of indirect paths implies that some mutational steps are evolutionarily

prohibited because they decrease regulation strength. The reason is closely linked to epistasis

– non-additive effects of two or more mutations on regulation strength. (see **Supplementary**

Methods 7.5 for further details on epistasis). More specifically, such inaccessible steps are a

result of sign epistasis. In this kind of epistasis, a double mutant of a TFBS regulates

expression more strongly than the TFBS itself, even though one or both constituent single

mutants regulate expression more weakly than the TFBS[37,75–77.] Indeed, epistasis is prevalent

in all three landscapes **(Table S5)**. Specifically, we observe sign epistasis in 62%, 66%, and

65% percent of interactions between single mutant pairs in the CRP, Fis, and IHF landscapes.

The existence of accessible paths alone does not tell us how easily a high peak can be found

through Darwinian evolution. The reason is that only a very small fraction of all paths to that

peak may be accessible, and an evolving population may not find any one of those paths. To

quantify the fraction of accessible paths, we determined the total number of paths from each

non-peak genotype to each high fitness peak, and computed the fraction of those paths that

are accessible. **Figure 3a** shows how this fraction depends on the number of mutational steps

in a path. It behaves similarly for all three landscapes (**Figure 3a**). The majority of two-step

paths (a fraction greater than 80%) are accessible in all three landscapes (**Figure 3a**), but the

fraction of accessible paths dwindles rapidly with an increasing number of mutational steps.

It reaches a minimum below 1% for all three landscapes at eight mutational steps (**Figure**

3a).

High peaks have large basins of attraction that share many TFBS genotypes

In the following analysis, we computed the basin of attraction of each peak, defined as the set

of non-peak genotypes from which accessible paths to the peak exist[70,78,79]. In other words, a

peak's basin of attraction comprises all genotypes from which adaptive evolution can access

the peak. The peaks in our landscapes have widely different basin sizes. They include peaks

accessible from a mere fraction of genotypes to those accessible by a majority of genotypes

(**Figure 3b**). Notably, high peaks generally had larger basins of attraction than low peaks

(**Figure 3c, Supplementary Figure S23**). In addition, we found that many genotypes are part

of multiple basins of attraction. Adaptive evolution may reach different high peaks starting

from any such genotype. For all pairs of high peaks, we computed the pairwise overlap

(intersection) of the basins of attraction, i.e., the fraction of genotypes that are part of both

basins of attraction. (**Figure 3d, Supplementary Figures S24-S25**). The number of

genotypes in this intersection varies widely, but the basins of attraction of high peaks

generally share a substantial fraction of genotypes (mean shared genotypes: 48%, 42%, and

36% among all pairs of high peaks for the CRP, Fis, and IHF landscapes, **Figure 3d**)

Figure 3. Peaks and their basins of attractions. a. The fraction of accessible paths

declines with increasing path length. The figure illustrates how the fraction of accessible

paths to high peaks (vertical axis, logarithmic scale) decreases with the length of the shortest

accessible path (horizontal axis). Accessible paths are defined as paths where each step

increases regulation, starting from a specified initial genotype. Each colored line corresponds

to data from a different TF landscape: CRP (green), Fis (orange), and IHF (blue). Circles

indicate the mean fraction of accessible paths for a given path length. **b. Higher peaks have**

larger basins of attraction. The split-violin plots display the distribution of basin sizes

(vertical axis) for high (red) and low (blue) peaks in the three TF landscapes CRP, Fis, and

IHF (horizontal axis). High peaks have significantly larger basins of attraction (Welch Two

Sample t-tests; CRP: t-value = 9.0898, df = 63.509, and p-value = 4.22×10^{-13}, with mean

basin sizes of x_{high} = 9528.426 and x_{low} = 5655.035. Fis: t-value = 7.617, df = 172.35, and p-

value = 1.645×10^{-12}, with mean basin sizes of x_{high} = 14206.70 and x_{low} = 10931.94. IHF: t-

value = 6.1777, df = 201.99, and p-value = 3.521×10^{-9}, with mean basin sizes of x_{high} =

10965.872 and x_{low} = 8664.953.) The violin plots show the distribution of basin sizes for

each of the two kinds of peaks. Their width represents the frequency of a given basin size.

The vertical length of the box in each plot covers the range between the first and third

quartiles (IQR). The horizontal line within the box represents the median value, and whiskers

span 1.5 times the IQR. **c.** The regulation strength of peak genotypes (horizontal axis) is

plotted against the size of their basins of attraction (vertical axis, shown as a percentage of all

(non-peak genotypes) for peaks in all three landscapes (color legend). Color-coded values of

R at the top of the graph indicate Pearson correlation coefficients for each landscape (color

legend). Curved lines are based on a linear regression analysis (in linear space), with grey

shading indicating 95% confidence intervals (CRP: R^2=0.28, N = 2,154; Fis: R^2=0.24, N =

2,312; IHF: R^2=0.23, N = 2,434). **d. Basins of attractions share many TFBS genotypes.**

The violin plots with embedded boxplots illustrate the fraction of genotypes shared between

basins of attractions (vertical axis) for all pairs of high peaks in the CRP (green), Fis

(orange), and IHF (purple) TF landscapes (horizontal axis). Specifically, we computed basin

overlap as one minus the Jaccard coefficient (**Supplementary Methods 7.5**). A value of one

(zero) indicates that the basin of attraction of two peaks share all (no) genotypes (CRP: 61

high peaks , N=1,830 comparisons, Fis: 172 high peaks, N=14,706 comparisons, IHF: 199

high peaks, N=19,701 comparisons). Their width represents the frequency of a given basin

overlap. The vertical length of the box in each box covers the range between the first and

third quartiles (IQR). The horizontal line within the box represents the median value, and

whiskers span 1.5 times the IQR.

Genetic drift facilitates and clonal interference reduces the attainment of high peaks

Taken together, our analyses thus far indicate that high peaks in the CRP, Fis, and IHF

landscapes are more accessible than low peaks (**Figure 3b**), and their basins of attraction also

share more genotypes (**Figure 3d**). However, the accessible evolutionary paths to high peaks

are often indirect. In addition, the fraction of accessible paths to any one high peak dwindles

rapidly with the distance from that peak (**Figure 3a**).

To understand how these topographical features jointly affect adaptive evolution, we

simulated the evolutionary dynamics on our three landscapes under the assumption that

natural selection would favor strong regulation and that fitness is proportional to regulation

strength.

Because high peaks constitute only a small fraction of all peaks (2.8%, 0.4%, and 0.5% in the

CRP, Fis, and IHF landscapes), we hypothesized that most evolving populations would reach

only low peaks. To test this hypothesis, we took advantage of the fact that *E.coli* has a low

genomic mutation rate of 2.2×10^{-10} per base pair per generation [43], and that we study

evolution only within an 8-nucleotide segment of a TFBS. In addition, *E. coli* has a large

effective population size ($>10^8$ individuals[43]), which means that genetic drift is weak and

even minor differences in fitness are visible to natural selection [43,80]. These conditions imply

that a population on our landscape would evolve in the well-studied strong-selection weak-

mutation regime (SSWM)[39,81–83]. In this regime, a population is monomorphic most of the

time, until a beneficial mutation arises, which usually sweeps rapidly to fixation. In other

words, adaptive evolution can be modeled as an adaptive walk, in which each mutational step

is taken with a fixation probability that has been derived by Kimura [80]. We thus call the

resulting adaptive walks Kimura walks (**Supplementary Methods 8**). We simulated 10^3 such

adaptive walks starting from each of 15,000 randomly and uniformly selected starting (non-

peak) TFBS genotypes from each landscape. Each random walk comprised up to 25

mutational fixation steps, unless it reached a fitness peak earlier. Each adaptive walk also

accounted for known *E. coli* mutational biases [84–86] (**Supplementary Methods 8**).

As we hypothesized, most adaptive walks indeed reach only low peaks (90%, 85%, and 87%

percent for the CRP, Fis, and IHF landscapes.). However, the percentage of walks that

terminate at high peaks is several-fold greater than the proportion of high peaks. Specifically,

10 percent of walks terminate at high peaks in the CRP landscape, which is 3.6-fold higher

than the percentage (2.8%) of high peaks in this landscape. In the Fis and IHF landscape,

adaptive walks are 2 and 1.6-fold more likely to terminate at high peaks than expected from

the proportion of these peaks among all peaks (7.4% and 8.1% percent) (**Figure 4a**). In other

words, adaptive walks reach high fitness peaks more often than one might expect.

The adaptive walks that reached a high peak were short (**Figure 4b**). On average, they were

also no more than half a mutational step longer than the shortest genetic distance between

each starting genotype and peak (**Figure 4b**, CRP landscape: path length 3.2 ± 1.6

[mean±s.d.] vs. genetic distance 2.8 ± 1.2; Fis landscape: path length 3.1 ± 1.5 vs. genetic

distance 2.7 ± 1.2; IHF landscape: path length 3.1 ± 1.5 vs. genetic distance 2.7 ± 1.2).

These short paths can be explained by our previous observation that peaks are widely

distributed across the genotype spaces of the three landscapes. This distribution makes it

easier for populations starting from any (non-peak) genotype to find a local peak through few

mutations. Additionally, path lengths realized during adaptive walks tend to be shorter than

the average lengths of accessible paths. (**Figure 4b**) The reason is that the average length of

accessible paths takes all accessible paths into consideration, whereas adaptive walks tend to

stop at the first peak they encounter, which results in shorter realized path lengths.

Because genetic drift can cause evolving populations to escape a low fitness peak and attain a

higher fitness peak, we also asked how small population sizes affect the likelihood that a

population attains a high fitness peak (**Supplementary Material 8, Supplementary Figures**

S26-28). When simulating adaptive evolution in populations of N=15,000 individuals, we

found that the likelihood that a population reaches a high peak increased by at least 10%

percent (to 18%, 20%, and 21% for the CRP, Fis, and IHF landscapes, **Supplementary**

Figures S26-S28).

We also studied how deviations from the strong selection weak mutation regime would affect

evolutionary dynamics. In large populations or at high mutation rates, populations tend to be

polymorphic and are subject to clonal interference, where the most advantageous of several

co-occurring mutations typically dominates and becomes fixed [87–89]. To approximate this

dynamic, we conducted simulations using "greedy" adaptive walks (**Supplementary**

Methods 8) [38,90]. In such an adaptive walk, it is always a genotype's mutational neighbor

with the highest increase in regulation strength that becomes fixed [82,90–92]. In other words, a

greedy walk starting from a given genotype is deterministic. We, therefore, simulated only

one walk for each of the 15,000 randomly chosen (non-peak) starting genotypes. We found

that the fraction of greedy walks reaching high regulation peaks is slightly lower than in the

SSWM regime, with 1%, 2%, and 5% fewer walks achieving such peaks in the CRP, Fis, and

IHF landscapes, respectively (**Supplementary Figure S29**). This outcome is expected,

because greedy walks prioritize immediate fitness gains at the expense of the ability to

discover higher fitness peaks[90].

The attainment of peaks is highly contingent on chance events

Because different basins of attraction share many TFBS genotypes (**Figure 3d,**

Supplementary Figures S24-S25), it is possible that adaptive evolution starting from any

one genotype can reach different peaks, depending on which mutational paths it takes. In

other words, the structure of the landscapes we study may lead to evolutionary contingency

[93,94] – the dependence of an outcome of evolution on unpredictable chance events[26,93,95].

Indeed, non-peak genotypes can access on average around 30% of the total number of high

peaks in each landscape (**Supplementary Figure S25).** To assess the prevalence of

evolutionary contingency, we determined how many different peaks are attained by 10^3

adaptive walks starting from the same randomly chosen genotype. We performed this analysis

on 15,000 starting genotypes, focusing on the subset of those starting genotypes from which

at least one high peak is reached during the 10^3 walks. Specifically, 28.6%, 23%, and 24.5%

of starting genotypes for the CRP, Fis, and IHF landscapes, reached at least one high peak.

Notably, these percentages represent the fraction of starting genotypes that can access at least

one high peak, not the total fraction of adaptive walks that lead to a high peak, which is lower

(as shown in **Figure 4a**). For instance, while adaptive walks starting from 28.6% of

genotypes reached at least one high peak in the CRP landscape only 10% of all adaptive

walks in this landscape end at a high peak (**Figure 4a**). Importantly, for any one starting

genotype where random walks reached at least one high peak, they typically reached more

than one high peak (**Figure 4c**, median±IQR, 4±4, 4±8, 6±6 for the CRP, Fis, and IHF

landscapes, respectively). The number of high peaks attained from any one starting genotype

was as large as 21, 30, and 33 for the three respective landscapes. Thus, evolutionary

contingency is pervasive in all three landscapes. In addition, not only the end point of

evolution but also the evolutionary trajectories leading to this end point show contingency.

That is, adaptive walks starting from a specific genotype and ending at a specific high peak

often take multiple paths to this peak (**Supplementary Figure S30**).

Biases towards specific evolutionary outcomes have been documented in both empirical and

computational studies [26,27,70,93,96–98]. They also exist in our landscapes (**Figure 4d,**

Supplementary Figure S31). **Figure 4d** illustrates both contingency and bias for 10^3

adaptive walks starting from each of 10 different genotypes in the CRP landscape (thus a total

of 10^4 adaptive walks). Depending on the starting variant, the adaptive walks reached

between 1 and 16 high peaks. Whenever different peaks are reached, they are reached by

markedly different proportions of adaptive walks. The same holds in the Fis and IHF

landscapes (**Supplementary Figure S32**). Additionally, while multiple routes to each peak

can be traversed (**Supplementary Figure S30**), we observed in all three landscapes biases

towards specific paths that are taken more frequently than others (**Supplementary Figure**

S33).

Figure 4. Peak accessibility, contingency and biases in adaptive walks. a. More than

10% percent of adaptive walks lead to high peaks. The graph shows the cumulative

distribution function (CDF, vertical axis) of regulation strength (horizontal axis) attained at

the end of 15 million adaptive walks (15,000 starting genotypes × 1,000 adaptive walks each)

for each TF landscape (color legend, CRP=: green; Fis: orange; IHF: purple). Dashed lines

intersect the CDF at points equivalent to the wild-type regulation strength for each TF (color

legend). They help to infer the proportion y_{TF} of walks that terminate at high peaks ($S > S_{WT}$),

which is indicated by the numerical values at the graph's top for each of the landscapes (color

legend). **b. Paths to high peaks are not much longer than genetic distances from the**

starting genotype. Colored boxplots summarize, for each of the three landscapes (horizontal

axis), the distribution of shortest genetic distances (blue) and actual path lengths (green) for

adaptive walks terminating at high peaks. Path lengths are typically less than one mutational

step longer than genetic distances. Each box covers the range between the first and third

quartiles (IQR). The horizontal line within each box represents the median value, and

whiskers span 1.5 times the IQR. Numbers atop each plot indicate means ± one standard

deviation. **c. Different numbers of high peaks are reached from different starting**

variants. Violin plots integrated with boxplots show the distribution in the number of high

peaks reached (vertical axis) from each starting variant that attained any high peak during

10,000 adaptive walks. The number of these variants is shown as an integer on top of the

panel, and its percentage among all 15,000 starting variants is shown in parentheses (color

legend). Underneath it, the panel shows the mean ± 1 s.d. of the number of attained high

peaks. **a. d. Some high peaks are reached more often than others.** We randomly and

uniformly sampled 10 starting genotypes from the CRP landscape, started 10^3 adaptive walks

from each, and recorded the number and frequency of distinct high peaks attained in these

random walks. Results for each starting genotype are symbolized by a vertical bar. The

number of stacks within each bar (delineated by horizontal lines, also indicated by an integer

above each bar) indicates the number of high peaks reached by the 10^3 adaptive walks.

Starting variants are ordered in ascending order based on this number of attained peaks. Stack

height indicates the fraction of walks that reached the same peak, and is indicated in red,

orange, and yellow for the three most frequently attained peaks. **See Supplementary Figure**

S32 for the Fis and IHF landscapes.

DISCUSSION

We evaluated the ability of three *E. coli* global regulators to control gene expression through

each of more than 30,000 TFBSs for each regulator. To achieve this, we utilized a synthetic

plasmid-based system that facilitates high-throughput fluorescence measurements. This

system allowed us to quantify the regulatory strength of individual TFBSs by measuring gene

expression through GFP fluorescence [51]. Additionally, the system insulates the control of

transcription from any direct effects library sequences might have on transcription, such as

the transcriptional and translational impacts of 5' untranslated regions on gene expression

[99,100]. Recent studies have investigated large empirical datasets of cis-regulatory genotypes,

examining the impact of sequence variation on gene regulation in both eukaryotes and

prokaryotes[6,39,49,59,101–104]. However, few studies have focused on these interactions from an

evolutionary perspective, as we do here [6,39,103].

We showed that the regulatory landscapes of all three TFs are highly rugged and have

multiple peaks. The ruggedness of all three landscapes is also supported by the prevalence of

epistasis in them (**Supplementary Table S5**). A particularly important form of epistasis is

sign epistasis[37,75,76], because it can lead to multiple adaptive peaks [37,75,76] (see **Supplementary**

Methods 7.5) Our landscapes contain up to 65% of mutation pairs with sign epistasis, a value

that is especially high compared to the almost exclusively additive interactions of mutations

in eukaryotic TFs[6,105]. However, the TFs we study are not exceptions among other

prokaryotic TFs. Recent studies have shown that epistatic interactions are common in binding

sites of local prokaryotic regulators, such as AraC (sign epistasis in more than 50% of 20

mutants[106]) and Cl (sign epistasis in 85% of 113 mutants[107]). We attribute this greater

incidence of epistasis to the nature of prokaryotic TFBSs. Specifically, prokaryotic TFBSs

are at approximately 20bps, twice as long as eukaryotic ones[57,108] and exhibit symmetries

reflecting the dimeric state of their cognate TFs[109–111]. These factors may increase the

likelihood of intramolecular epistasis.

Despite high ruggedness and epistasis, we found that a modest proportion of evolving

populations can access the highest peaks of each landscape. Specifically, in the large

populations that are typical of *E. coli,* at least 10% percent of evolving populations reach one

of the highest peaks. The clonal interference that may occur in even larger populations or in

populations with high mutation rates only modestly reduces this percentage to 5%

(**Supplementary Figure S29**) Conversely, in small populations where genetic drift can help

a population escape from a low peak, this percentage increases to 18%. Numbers like these

render the de novo evolution of strong transcription factor binding sites plausible for our

three global regulators. Once such binding site have originated in one population, they can

also spread to others through horizontal gene transfer[112,113].

It is well known that epistasis can influence the trajectories of evolving populations on

adaptive landscapes[37,76,114]. The regulation strength of a TFBS is partially determined by the

combined interactions of the nucleotides within the TFBS[115–118], which in turn affects TF-

TFBS binding affinities. Different combinations of mutations can result in similar regulation

strengths and create multiple evolutionary pathways to achieve optimal or near-optimal

regulation[6,119]. This can lead to overlapping basins of attraction among different peak TFBSs

conveying strong regulation, such that even evolution starting from the same genotypes can

take different paths and reach different peaks with similar regulation strengths

(**Supplementary Figure S30**). Indeed, we observed such overlapping basins in our

landscapes (**Figure 3d and Supplementary Figure S24**). Moreover, epistasis can create

plateaus of regulation strength where various combinations of nucleotides yield TFBSs with

intermediate regulation strengths. These plateaus serve as common evolutionary

intermediates that multiple starting sequences can traverse on their way to different peaks,

and thus contribute further to the shared basins of attraction we observe (**Supplementary**

Figure S24).

In a landscape with substantial epistasis, the sequence of mutations that occurs in an evolving

population can render adaptive evolution highly contingent on this sequence[93,94]. For

example, some beneficial mutations within a TFBS might only be accessible after other

specific mutations have occurred. Different starting sequences or early mutations can also

change the spectrum of accessible mutations, leading to different peaks with similarly strong

regulation [93,94]. Indeed, we observed in all three landscapes that different evolving

populations starting from the same genotypes in the landscape attain different peaks (**Figures**

3d, 4d and Supplementary Figure S32) [93,95,120,121]. Such contingency renders evolution less

predictable [26,93,122].

Despite the prevalence of contingency in all three landscapes, evolving populations starting

from the same genotype more often attain some peaks than others (**Supplementary Figure**

S32). Moreover, for each attained peak with multiple possible evolutionary routes, we also

observed that some paths are more frequently transversed than their alternatives

(**Supplementary Figure S33**). That is, evolution is biased towards traversing some paths and

attaining some peaks more often than others[123,124]. These observations emphasize the

complex interplay between chance, contingency, and evolutionary biases in shaping the

outcomes of adaptive evolution.[26,93,95,97].

The three TFs we studied here belong to different protein families, yet they have regulatory

landscapes with similar topography. All three landscapes highly rugged, highly epistatic, and

harbor multiple peaks that are widely scattered, and this holds for peaks of low, intermediate,

and high regulation strength. This might help to explain how TFBSs with intermediate and

high regulation strength can be easily attained during evolution. In addition, peak

accessibility in all three landscapes increases with peak height (**Supplementary Figure S31**).

On the one hand, these commonalities may be caused by common biological or biochemical

properties of the three TFs. For example, CRP has been suggested to possess nucleoid-

associated protein properties similar to Fis and IHF, due to its ability to bend and loop

DNA[125]. On the other hand, the commonalities may reflect general characteristics of global

gene regulation. One of them is that global TFs often bind unspecifically to multiple

TFBSs[11,13]. Also, they may bind DNA with a broad range of different affinities centered

around intermediate affinity, rather than bind few sites but very strongly [13]. This possibility is

supported by comprehensive analyses of in vitro eukaryotic TF binding affinities for

thousands of TFBS variants, showing essentially continuous binding affinity distributions for

hundreds of sites[6].

Despite broad similarities among the three landscapes we study, we also found some

differences. Most notable is the narrower distribution of regulation strengths for the binding

sites of CRP compared to those of Fis and IHF (**Figure 2a**). This is not unexpected, given

that CRP binding sites are known for being quasi-symmetric and less degenerate than those

of Fis and IHF[28,31]. Previous studies have also shown that Fis and IHF binding sites are less

conserved in their nucleotide structure and more biased toward AT-rich content[22,126,127]. One

possible explanation for this observation is the nature of the binding dynamics of these

regulators. CRP, mostly restricted to gene regulatory interactions, may have evolved to finely

tune its regulatory output, resulting in a narrower distribution of regulation strengths. In

contrast, Fis and IHF, which are involved in functions beyond gene regulation, such as DNA

replication and genomic structural maintenance[128,129], might encompass a more diverse range

of regulation strengths.

The landscapes of our global regulators also differ in two respects from that of the previously

characterized local regulator TetR [27]. First, the peaks of the TetR landscape are more

clustered in genotype space due to its lower diversity of strongly binding TFBSs [27]. Second,

the distribution of regulation strengths for its TFBSs is narrower and shows more strongly

binding TFBSs [27]. A possible explanation may lie in the fact that TetR regulates only two

transcription units[130].

Some of the properties of our landscapes are similar to what would be expected from

theoretical models, such as uncorrelated random landscapes[72,73,131,132]. First, both kinds of

landscapes have many peaks. Specifically, the number of peaks of the CRP, Fis, and IHF

landscapes is 93%, 96%, and 98% of that expected in a random landscape of the same size

(**Supplementary methods 7.6**). Second, the peaks of our landscapes are widely scattered in

genotype space. This scattering may reflect a biological property, namely that global

regulators can strongly bind multiple and highly diverse sites[11–13]. In addition, high peaks in

our empirical landscapes also have large basins of attractions, an observation that is

consistent with predictions from simple theoretical models [132].

One limitation of our work stems from our rigorous quality filtering of sort-seq data. As a

result, we lack regulatory data for some 40% of the TFBSs in each landscape. This limited

diversity of reliable data is a common feature in mutational library studies. It has several

technical causes, such as biases in library synthesis [133], PCR amplification [134], cloning, and

loss of sequence diversity after cell sorting [102]. In future work, this limitation could be

overcome by a combination of strategies, such as subsampling complete genotype spaces or

combining different molecular methods to overcome biases and diversity loss during PCR

amplification[135,136], sorting[48,137,138], and high-throughput sequencing[103].

A second limitation comes from our use of the sort-seq method, which is best-suited for our

work, because it allows the high-throughput measurement and sorting of millions of

individual cells in a standardized and straightforward manner. However, the method's

accuracy depends on the binning procedure. Other studies have used between two and several

dozen bins, with or without unbiased sampling [48,50,59,101,102,137,139]. Recommendations

emerging from this work include sorting of cells into at least four logarithmically ($\log_2$)

equally-spaced bins, each covering approximately 12.5-15% of the fluorescence distribution,

[59,102,140]. We followed these recommendations, using 13 bins. In addition, we computed a

weighted average to calculate expression values from sequences appearing in multiple bins, a

straightforward method validated by robust studies in transcriptional regulation analysis [39,49].

In addition, we validated individual regulation strengths with an independent method to

demonstrate its reliability.

A third limitation of our study is that we only examined variation at eight positions for each

TF. We selected these positions based on their importance for DNA binding and regulation by

our TFs, as evidenced by their high information content. We cannot exclude the possibility

that selecting a different set of positions could yield different landscape topographies.

However, we speculate that less information-rich position would not reduce but rather

increase the potential for the de novo evolution of strong TFBSs. For example, they may

facilitate landscape navigability through extradimensional bypasses[78,141] or provide small-

incremental changes in regulation strengths. These small changes help populations reach

peaks via diminishing returns effects, meaning that as a population gets closer to a peak, each

subsequent mutation contributes progressively smaller improvements to regulation strength

[142,143]. Investigating a wider array of positions and larger regulatory landscapes remains an

important task for future work.

Fourth, we use a simplified empirical system for the study of gene regulation—a promoter

followed by a single TFBS. Although this regulatory architecture exists for some genes[144],

global regulators often form part of more complex, combinatorial architectures[4,144,145] that are

influenced by environmental factors[2,146,147], concentrations of active TFs[148–150], chromosome

structure[151–155], methylation states[156], and more. Studies on more complex promoter

architectures may reveal regulatory landscapes with different topographies.

Lastly, we also analyzed the adaptive evolution of TFBSs in a simplified manner, performing

data-driven evolutionary simulations rather than experimental evolution. Although many

studies assume that regulation strength and fitness are correlated, this is not always the case.

Low binding affinities can be adaptive during development [157], and many genes exhibit a

nonlinear fitness-expression function [158] with a plateau of maximal fitness across a wide

range of expression levels [159–161]. For instance, most mutations and polymorphisms in the

promoter of the yeast gene TDH3 do not significantly affect fitness in a glucose-rich medium

[160]. For these reasons we focused our observations and interpretations on regulatory

phenotypes (regulation strength). Our simplified approach allowed us to model population

evolution in a large genotype space and avoid monitoring thousands of evolving regulatory

sequences simultaneously in vivo[162,163]. However, it cannot replace experimental evolution of

TFBSs, which also remains an important challenge for future work.

Transcription factor binding sites are among the simplest units of biological organization.

Our work provides the first large-scale analysis of the regulatory landscapes formed by such

sites for global transcriptional regulators. It shows that strong binding sites can readily evolve

de novo, even though prokaryotic transcription factor binding sites are much larger than their

eukaryotic counterparts and have more rugged regulatory landscapes. In addition, the

evolution of these simple sequences also displays phenomena that have been characterized in

much more complex systems, such as evolutionary contingency and evolutionary biases.

MATERIALS AND METHODS

The Supplementary Materials contain extended details of experimental procedures and data

analysis.

Strains and plasmids

Bacterial strains and plasmids used in this work are listed in Table 1. We obtained

electrocompetent *E. coli* cells of strain SIG10-MAX® from Sigma Aldrich (CMC0004). We

used this strain for molecular cloning and library generation due to its high transformation

efficiency. The genotype of this strain (**Supplementary Table S4**) is similar to DH5α

(Sigma Aldrich commercial information, see **Supplementary Table S4**). The strain is

resistant to the antibiotic streptomycin.

We amplified plasmid libraries in SIG10-MAX®, extracted their DNA, and transformed them

into mutants derived from *E. coli* K-12 strain BW25113 that harbor chromosomal deletions

of the *crp*, *fis*, or *ihfa* gene. We obtained these mutant strains from the KEIO collection [164]

and used them for sort-seq experiments. The design, genetic parts, and assembly of the

plasmid vectors we used in this study are available in the Supplementary material, as are all

primers, TFBS sequences/libraries, strains, and plasmids.

Sort-Seq procedure

To explore the regulatory effects of each transcription factor on binding sites in the

corresponding library, we constructed three plasmids, each of which enables the inducible

expression of one of our three TFs. These are plasmids pCAW-Sort-Seq-V2-CRP, pCAW-

Sort-Seq-V2-Fis, and pCAW-Sort-Seq-V2-IHF (**Supplementary Methods 3** and

Supplementary Table S1). We cloned the TFBS libraries (**Supplementary Table S3**) into

their respective plasmids and then transformed them into mutant strains lacking the

corresponding TFs (*Δcrp*, *Δfis*, and *Δihf*, as listed in **Supplementary Table S2**). We induced

TF expression using anhydrotetracycline (Atc) and, after overnight growth, performed cell

sorting for cell populations. During sorting, we distributed cells into 13 equally-spaced

logarithmic bins based on their fluorescence levels. We replicated each sort-seq experiment

three times from three separate library transformations for each TF. To mitigate the impact of

extrinsic noise (gene expression variation among cells [165]), we adhered to standard protocols

by normalizing our GFP fluorescence measurements against mScarlet-I fluorescence values

obtained from flow-cytometry assays [7,48,166,167]. We subsequently recovered the sorted cells

from each bin in 50mL Falcon tubes containing 10 mL of LB medium supplemented with

chloramphenicol and incubated them overnight at 37°C with shaking at 220 rpm. After this

growth period, we re-sorted cell cultures from each bin to eliminate potential contaminants

and ensure that the cell populations had preserved their fluorescence distributions. Following

re-sorting, we extracted plasmids from the cell population of each bin. We amplified and

barcoded the TFBS region from each population through a polymerase chain reaction (PCR).

Lastly, we sequenced barcoded amplicons containing TFBS sequences, and used the

sequencing results to calculate the regulation strengths of each TF to its TFBSs in the

corresponding library. More details are provided in the Supplementary Material.

Regulation strengths

Due to gene expression and measurement noise, individual TFBS variants in a sort-seq

experiment usually appear in more than a single bin, and their read count (frequency) varies

among bins[48,49,101,138]. Following established practice[27,49,103], we used a weighted average of

these frequencies for each variant to represent the mean expression level driven by the

variant. To facilitate the interpretation of this quantity, we converted this expression level

into a regulation strength relative to the highest observed regulation strength for a given TF,

to which we assigned a value of one. From each library, we selected a single naturally

occurring binding site for each TF that was previously characterized in the literature as a

strong binder. We called this TFBS the WT sequence and used it as a baseline to separate

weakly (higher GFP expression) from strongly (lower GFP expression) binding TFBSs.

Validating regulation strengths with plate reader measurements

To further validate our regulation strength data, we chose 10 DNA binding sites from each

bin (10 variants ×13 bins = 130 variants in total per TF library, **Supplementary File 2**),

covering a wide range of measured regulation strengths. We cloned these sequences into the

appropriate vector pCAW-Sort-Seq-V2-TF (TF: CRP, Fis or IHF), and transformed them into

the appropriate mutant strain. We picked individual colonies and grew them overnight (16

hours, 37°C, 220 rpm) in liquid LB supplemented with 50 μg/mL of chloramphenicol and

anhydrotetracycline. We diluted the cultures to 1:10 (v/v) in cold Dulbecco's PBS (Sigma-

Aldrich #D8537) to a final volume of 1 mL. We transferred 200 μl of the diluted cultures into

individual wells in 96-well plates and measured GFP fluorescence (emission:

485nm/excitation: 510nm, bandpass: 20nm, gain: 50), as well as the optical density at 600nm

(OD_{600}) as an indicator of cell density. We then normalized fluorescence by the measured

OD_{600} value to account for differences in cell density among cultures and compared the

obtained ratios to the previously inferred regulation strengths for the 130 selected variants.

We performed all such measurements in biological and technical triplicates (three colonies

per sample, and three wells per colony, respectively).

Code Availability and Data Analysis

All code used for processing data and plotting, as well as the final processed data, plasmid

sequences, and primer sequences are available in our GitHub repository.

Supplementary Information is linked to the online version of the paper
829

Acknowledgements: We gratefully acknowledge financial support from the Swiss National
Science Foundation grant 310030_208174. We also extend our thanks to the UZH University
Priority Research Program in Evolutionary Biology, the UZH flow cytometry facility, and the
Functional Genomics Center Zurich for their technical support. We are especially thankful to
Andrei Papkou for his guidance with computational analysis and for engaging in theoretical
discussions.

Author contributions: C.A.W. and A.W conceived the study and designed experiments.
C.A.W. carried out experiments. C.A.W and A.W. analyzed data. C.A.W wrote computer
code to carry out bioinformatic work and analysis. C.A.W generated figures. L.G. wrote
computer code to carry out simulations. C.A.W. and A.W. wrote the paper, which was edited
by all authors.
841

842   **Data availability:** Sequencing data has been deposited in the NCBI database under the
843   BioProject accession code: PRJNA1162449. The data generated in this study have been
844   deposited in the Zenodo public repository and are accessible via the following DOI:
845   10.5281/zenodo.13838265
846

847   **Computer code:** The computer code generated in this study has been deposited in the
848   Zenodo public repository and is accessible via the following DOI: 10.5281/zenodo.13838265
849

850   **Competing interests:** The authors declare no competing interest
851

852   **Corresponding authors:** Correspondence to andreas.wagner@ieu.uzh.ch
853

S17. Genetic diversity among peaks across TF landscapes
S18. Genetic diversity among high peaks across TF landscapes
S19. Principal component analysis of the CRP landscape
S20. Principal component analysis of the Fis landscape
S21. Principal component analysis of the IHF landscape
S22. Accessible paths to high peaks are less than two mutational steps longer than the
shortest genetic distances to the peak
S23. Moderate positive correlation between peak regulation strength and basin size for
the three regulatory landscapes
S24. The basins of attractions of high peaks overlap
S25. Distribution of the fraction of accessible high peaks per TFBS variant across TF
landscapes
S26. Kimura adaptive walks on the CRP landscape
S27. Kimura adaptive walks on the Fis landscape
S28. Kimura adaptive walks on the IHF landscape
S29. Fraction of greedy adaptive walks attaining high peaks for different TFs
S30. Diversity of alternative shortest paths for pairs of starting variants and their
respective attained peaks
S31. Peak genotypes conveying stronger regulation are also reached more often during
adaptive evolution.
S32. The frequency distribution of attained high peaks is biased towards some
frequently attained peaks.
S33. The frequency distribution of transversed paths to high peaks is biased towards
certain frequently traversed paths.
928

SUPPLEMENTARY METHODS

1. General procedures:

Although all general procedures have been previously described [27], we describe them again

below for completeness.

1.1 Media and reagents

To prepare SOB medium, we mixed 25.5g of solid medium stock (VWR J906) with 960 ml

of purified water and subjected the resulting suspension to autoclaving. To prepare the SOC

medium, we dissolved 20 ml of 1 M D-glucose (Sigma G8270) and 20 ml of 1 M magnesium

sulfate (Sigma 230391) in 960 ml of pre-prepared SOB solution. For the LB medium, we

combined 25g of solid medium stock (Sigma-Aldrich L3522) with 1 liter of purified water

and then autoclaved it. To prepare the M9 minimal medium, we diluted M9 minimal salt

sourced from Sigma (M6030) in distilled water as per the manufacturer's guidelines,

autoclaved the solution, and added 0.4% glucose (Sigma G8270), 0.2% casamino acid (Merk

Millipore, 2240), 2 mM magnesium sulfate (Sigma 230391), and 0.1 mM calcium chloride

(Sigma C7902). Where required, we supplemented growth media with chloramphenicol (50

µg/mL), anhydrotetracycline (100 ng/mL, Cayman-chemicals #10009542), and/or glucose

(0.4% w/v final concentration). We prepared anhydrotetracycline by diluting the dried

chemical in absolute ethanol, from a stock concentration (1000X) of 100 µg/mL to a working

concentration of 100 ng/mL.

1.2 Overnight incubation of cultures in liquid and solid medium

We cultivated bacteria in liquid LB medium (using either 15mL or 50mL Falcon tubes),

enriched with chloramphenicol at a concentration of 50 µg/mL. We incubated these cultures

for a period of 16 hours at a temperature of 37°C, with a shaking speed of 200rpm and 50

mm orbital motion, using an Infors HT Multitron Incubator Shaker. Similarly, for cultures in

solid medium, we grew bacterial colonies on LB-agar plates (using sterile plastic Petri dishes

of 90mm × 15mm dimensions), also supplemented with chloramphenicol at a concentration

of 50 µg/mL, and incubated them for the same time and at the same temperature.

1.3 PCR Reactions

Except where specifically mentioned, we amplified DNA fragments through a polymerase

chain reactions (PCR), employing Q5® high-fidelity polymerase (NEB #M0491L) to

minimize mutation introduction into the amplicons. We followed the protocol recommended

by NEB, aiming for a final reaction mixture of 50uL. We conducted each PCR reaction twice,

and combined the products from these duplicates after the completion of the reaction. We

determined the primer melting temperatures (Tm) using the NEB Tm calculator (accessible at

https://tmcalculator.neb.com/#!/main), with primers at a concentration of 500nM.

1.4 Verifying PCR products through gel electrophoresis

Unless indicated otherwise, we verified successful PCR amplification via gel electrophoresis

to ensure the presence of singular-band amplicons and the absence of non-specific bands.

This process involved separating PCR products in a 0.8% agarose Tris-EDTA (TAE) gel. We

conducted electrophoresis for 45 minutes at 120V, or until the bands had progressed beyond

the halfway point of the gel's total length.

1.5 DNA purification with commercial kits

Once we had confirmed a successful PCR through gel electrophoresis, we purified the PCR

products utilizing the Monarch® DNA PCR/Gel Extraction Kit (NEB #T1020L), adhering to

the manufacturer's protocol. In situations requiring further purification (such as the

occurrence of non-specific bands post-PCR), we performed gel purification. This involved

mixing 10 µL of 6x NEB DNA dye with each 50 µL PCR product, then loading all 60 µL onto

a 1% agarose gel. We carried out electrophoresis for 45 minutes at 120V, or until the bands

had moved more than halfway through the gel. We then excised the DNA band corresponding

to the amplified sequence using a scalpel. For extracting DNA from the gel, we used the

Monarch® DNA Gel Extraction Kit (NEB #T1020L).

1.6 Gibson assembly[168]*:*

We assembled PCR-amplified fragments using the NEBuilder-HiFi® DNA Assembly Master

Mix kit (NEB #E2621L). We determined the molarity required for assembly based on the

guidelines outlined by the Barrick Lab (details available at

https://barricklab.org/twiki/bin/view/Lab/ProtocolsGibsonCloning). We incubated the

assembly mix for one hour at 50°C in a dry bath incubator, followed by cooling on ice for

subsequent steps.

1.7 Preparation of electrocompetent cells

We utilized glycerol/mannitol step centrifugation to prepare electrocompetent cells [169]. To this

end, we first cultured the appropriate *E. coli* strain (**Supplementary Table S4**) in 5 mL SOB

medium at 37°C with a shaking speed of 250 rpm overnight. The next day, we transferred 3

mL of the culture to 300 mL SOB medium, and incubated under the same conditions until the

OD600 reached a value between 0.4 and 0.6 (measured at an optical path length of 1 cm),

which took approximately 2-4 hours. After cooling the culture on ice for 15 minutes, we

centrifuged the cells at 4°C and 1,500 g for 15 minutes. We then resuspended the cells in 60

mL of ice-cold distilled water (dH$_2$O) and divided them into three 50 mL tubes. Gradually, we

added 10 mL of an ice-cold glycerol/mannitol solution (consisting of 20% glycerol (w/v) and

1.5% mannitol (w/v)) to each tube using a 10 mL pipette. We centrifuged the tubes at 1,500 g

and 4°C for 15 minutes in an Eppendorf 5810/5810 R centrifuge with

acceleration/deceleration set to zero. After discarding the supernatant, we resuspended cells

in 3.0 mL of the same glycerol/mannitol solution. We transferred these suspensions to pre-

cooled 1.5 mL tubes, and incubated them in a dry ice-ethanol bath for about 1 minute.

Finally, we stored the suspensions at -80°C for future transformation experiments.

1011

1.8 Electroporation

In all transformation experiments described in this study, we used 100 µL of

electrocompetent cells for electroporation, employing 0.2 cm cuvettes (EP202, Cell Projects,

UK) and a Micropulser electroporator (Bio-Rad) set to the EC3 setting (15k V/cm). Post-

electroporation, we recovered cells in 1 mL of SOC media, warmed in advance in 15 mL

Falcon tubes. This recovery step lasted for 1.5 hours at 37°C with a shaking speed of 220

rpm. Unless specified otherwise, we spread 300 µL of each recovered culture on an LB agar

plate that contained 50 µg/mL chloramphenicol. We then incubated this plate overnight for 16

hours at 37 °C. Subsequently, we confirmed the identity of the clones on the plate via Sanger

sequencing.

1022

2. The design of plasmid pCAW-Sort-Seq-V2

The plasmid pCAW-Sort-Seq-V2 is a derivative of the plasmid pCAW-Sort-Seq[27]. Briefly,

plasmid pCAW-Sort-Seq harbours a pBBR1 replication origin, which ensures a broad host

range and a low copy number—typically between 5 to 10 copies per cell[170].It also harbours a

chloramphenicol resistance gene, a TetR repression system[171] and a TFBS measuring module.

The latter consists of an interchangeable TFBS that lies between a constitutive promoter and

a superfolder GFP (*sfgfp*[172]) reporter gene. In this system, TF-TFBS interactions can decrease

GFP production by physically obstructing the bacterial RNA polymerase, a phenomenon

known as steric hindrance. The reporter gene *sfgfp* is insulated by a transcriptional insulator

named RiboJ, a synthetic ribozyme that removes 5'UTR interferences from variable TFBS

sequences in the mRNA by self-cleavage[173]. More details and features have been previously

described in ref. [27]. Here, we have integrated a bicistronic expression cassette into pCAW-

Sort-Seq to enable the regulated expression of a TF of choice. This cassette harbors the gene

encoding one of our focal TFs situated upstream of a *mscarlet-I* [174] reporter gene to monitor

the expression of this bicistronic operon via fluorescence. Regulation is achieved through the

pLtetO-1 [171] promoter, a synthetic promoter tightly repressed by TetR. Expression is initiated

by the addition of anhydrotetracycline (Cayman Chemicals, catalog #10009542), which

relieves TetR-mediated repression and activates the expression of the entire cassette.

1041

3. Construction of the plasmid pCAW-Sort-Seq-V2 and its variants

We initiated the construction of pCAW-Sort-Seq-V2 by synthesizing (Twist Biosciences,

Calfornia, USA) and cloning a codon-optimized *mscarlet-I* gene expression cassette into the

pCAW-Sort-Seq plasmid. This gene is oriented antiparallel to the tetracycline resistance gene

(*tetr*). This cloning step resulted in the formation of the intermediate pCAW-Sort-Seq-

mScarlet plasmid. Subsequently, we amplified for each of our three TFs the encoding gene

(*crp*, *fis* and *ihf*) from the genome of the *E. coli* strain BW25113 and inserted it upstream of

the *mscarlet-I* gene in a bicistronic operon configuration using Gibson assembly. This process

produced TF-specific expression plasmid variants. The final phase involved removing the

core promoter to create negative controls for each TF and cloning libraries into each plasmid

variant upstream of the sf*gfp* gene. This yielded the plasmids used in our experiments

(**Supplementary Table S1**). The specific sets of primers used for each PCR reaction are

listed in **Supplementary Table S6**.

4. Library design, synthesis, and cloning
1056

We based the design of the TFBS library for each TF on consensus sequences available in the

RegulonDB database[47], the most comprehensive database for *E. coli* transcriptional studies.

We designed each library by randomizing the eight most important TFBS positions, such that

each of the four nucleotides had an equal probability (0.25) to occur at each position. The

expected library size for this approach is $(4^8) = 65,536$ sequences. Library compositions can

be found in **Supplementary Table S3**. We designed libraries in the Snapgene® software

(snapgene.com), and had them synthesized by IDT (Coralville, USA) as single-stranded DNA

Ultramers® of 140bp (4nmol). We resuspended each library in nuclease-free distilled water

and serially diluted it to a concentration of 50ng/uL. We used Ultramers® as templates in a

PCR reaction for the formation of dsDNA fragments and library amplification. We amplified

the Ultramer® DNA molecules with the following PCR program: 98°C/30 s; 25 cycles of

98°C/10 s, 60°C/15 s and 72°C/80 s; and 1 cycle of 72°C/5 min. We opted for a maximum of

20 cycles to reduce amplification biases. We analyzed amplification products through gel

electrophoresis to confirm that only a single product band with the expected size of around

140bp was present for each PCR reaction. After confirming the presence of single bands, we

purified the products using the Monarch® DNA gel extraction kit (NEB #T1020L). Whenever

unspecific bands appeared during electrophoresis, we gel-purified the PCR products using the

same kit.

1075

For the construction of each plasmid-based library, we digested 1μg of the purified library

with HindIII-HF (NEB #R3104) and BamHI (NEB #R3136) restriction enzymes in a 100μL

reaction, followed by overnight incubation at 37°C. We isolated the appropriate cloning

plasmid from its host strain using the QIAprep spin miniprep kit (Qiagen, Germany), and

digested it with the same enzymes. Post-digestion, we added 3μL of Quick CIP (calf

intestinal alkaline phosphatase) to prevent self-ligation by dephosphorylating DNA ends. We

purified the ligated DNA using the Monarch® DNA gel extraction kit (NEB #T1020L).

1083

We performed the ligation with a 10:1 molar ratio of insert-to-vector, using 100ng of vector,

10 units of T4 DNA ligase (NEB #M0202L), and 2 µL of 10X ligation buffer in a 20 µL

reaction. We incubated the mixture at 20-22°C for approximately 16 hours, followed by 15-

minute inactivation of the ligase at 65°C. We purified the ligation product, resulting in 10 µL

of DNA resuspended in dH_2O. We transformed *E. coli* SIG10-MAX® cells by

electroporation with this purified product.

1090

Post-transformation, we plated 50 µL of each recovered culture on LB agar for colony-

forming unit counting (cfu) and transformation efficiency estimation. From the agar plates,

we selected 30 colonies for colony PCR and Sanger sequencing to assess library diversity

(NightSeq® service, Microsynth, Switzerland). Our mean transformation efficiency was 10^6

cells per transformation. We diluted the remaining 950 µL of transformants in 9 mL of LB

medium with chloramphenicol and cultured it overnight. We then aliquoted the overnight

culture into 1 mL cryotubes with 20% glycerol and stored at -80°C.

1098

After transforming plasmid libraries into the SIG10-MAX® strains for reasons of

transformation efficiency and plasmid maintenance, we extracted the plasmids using a

QIAprep spin miniprep kit (Qiagen, Germany), and transformed them into the appropriate

host mutant host strains *Δcrp, Δfis and Δihf* (see the strain genotypes in **Supplementary**

Table S2). We cultured each library-transformed strain overnight, aliquoted it in 1 mL

cryotubes with 20% glycerol, and stored it at -80°C for further experimentation.

1105

5. Analysing and sorting cells

In preparation for cell sorting, we cultivated cells harboring each library in liquid LB medium

enriched with chloramphenicol. Specifically, we cultured 1mL of transformed cell aliquots

and a streak of cells with a control plasmid (a promoterless pCAW-Sort-Seq-V2 plasmid

lacking sfGFP expression) in 50ml Falcon tubes with 9mL LB medium (containing 50 μg/mL

chloramphenicol) overnight. Subsequently, we diluted these overnight cultures at a 1:100

ratio (v/v) in LB medium with chloramphenicol and divided them into two aliquots. We

supplemented one aliquot with the anhydrotetracycline (Atc) inducer to express the plasmid-

encoded TF. The other aliquot remained unchanged. We incubated both cultures for 5 hours

until late-exponential/early-stationary phase (200RPM, 37°C). Subsequently, we diluted

20μL of the cultures in 1mL of cold filtered Dulbecco's PBS (Sigma-Aldrich #D8537) in 15

mL FACS tubes.

1118

We performed FACS-sorting on a FACS Aria III flow cytometer (BD Biosciences, San Jose,

CA) using a 70 μm nozzle. We utilized a 488 nm laser for detecting forward scatter (FSC)

and side scatter (SSC) with a 488nm/10nm band-pass filter. We set the flow rate to 1.0,

adjusting sample dilution as necessary to achieve no more than ≈10,000 events/second.

Considering the small size of bacterial cells, we reduced the particle detection threshold to

the lowest feasible setting (200 arbitrary units on FSC and SSC channels), increasing it to a

maximum of 500 units if background noise was excessive. We then adjusted FSC-H and

SSC-H for cells with the negative control plasmid to center the bacterial population in the

cytometer's software (BD FACSDiva™ Software v9.0). We based the sorting and binning of

cells on both mScarlet-I (PE-Texas-Red channel, excitation laser: YellowGreen 561nm, LP

filter: 600 nm, BP filter: 610/20nm) and sfGFP fluorescence (FITC channel, excitation laser:

488nm , LP filter: 502nm, BP filter: 530nm/30nm), setting both the PE-Texas-Red and FITC

channels voltages so that the median fluorescence of the negative control was between 0 and

100 (arbitrary units) on the FITC-H and PE-Texas-Red-H axes.

1133

Initially, the sorting process involved establishing a gate for identifying cells that are red-

fluorescence-positive, i.e., reporter and thus TF-expressing. To establish this gate, we began

by measuring the autofluorescence of our negative control culture on the PE-Texas-Red-H

axis. Thereafter, we examined the fluorescence of a positive control that had been induced

with Atc and was expressing the mScarlet-I protein. We then configured a gate around the

positive mScarlet-I-expressing population, ensuring that all cells sorted through this gate

exhibited a level of reporter expression higher than the negative control.

1141

Next, we established the green-fluorescent sorting gates. For setting sorting gates on the

FITC-H axis, we first recorded the autofluorescence of the negative control culture. This

median autofluorescence defined the upper boundary of the lowest bin (B1) for the

experimental population. We then analyzed the fluorescence of 10^6 cells expressing sfGFP

and containing the library, without sorting, in order to establish the boundaries of the next

binning gates. We took the lower bound of the highest bin (B13) for the experimental

population to correspond to the 95th percentile of the fluorescence distribution of this

population. We chose boundaries between the remaining 11 intermediate bins with

equidistant spacing on a binary logarithmic ($\log_2$) scale. After we had determined the gates in

this way, we calculated the fractions of the previously 10^6 recorded cells that fell inside each

bin.

1153

To maintain statistical robustness and minimize sampling error in our downstream analyses,

we wanted to ensure that each of the 65,536 unique sequences in our library was represented

by at least 30 cells after sorting. In addition, it was crucial to account for an anticipated

diversity loss of up to 70% of the cell population during sorting, due to factors such as cell

death and the dilution of low-frequency and lower-fitness genotypes during the post-sorting

recovery phase [102]. To establish the number of cells required to be sorted initially, we thus

determined a multiplicative factor f based on the minimally needed number of 30 cells per

sequence and the expected cell retention rate post-sorting of 30%, i.e., $f = \frac{30}{0.30} = 100$

cells/sequence. We thus multiplied our initial library size by 100 for sorting purposes, i.e., we

sorted total of 6,553,600 cells. This calculation aims to ensure that despite a substantial

reduction in cell numbers post-sorting, each unique sequence remains adequately represented

in the surviving cell population. We sorted cells into 1.5 mL Eppendorf tubes, each

containing 500 µL of LB medium, and kept at 4 °C to prevent growth during sorting and

sample processing. We replicated the entire sorting procedure three times, each based on

independent library transformations.

1169

After sorting, we added 1mL of LB without antibiotics to each tube and removed 20 uL of the

resulting volume for serial dilutions. We transferred the remaining liquid culture (980 uL) to

50 mL falcon tubes and allowed each culture to recover for 2 hours (37°C, 220 rpm). After

recovery, we added 9mL of LB supplemented with chloramphenicol, and grew the cultures

overnight for freezing part of them in glycerol stock aliquots, for validating our binning

procedure (see below), and for extracting plasmid DNA for subsequent PCR and sequencing

steps. We used the 20 uL initially removed from each 1mL culture for preparing two serial

dilutions (10^{-4} and 10^{-6}) that we plated on LB-Cm agar plates (200uL per plate) to estimate

the post-sorting viability through colony forming unit (cfu) counting. By knowing how many

cells were sorted into each bin, we can estimate how many cells would be expected in our

dilutions and compare this number with the cfu counts we observed. In this way, we

estimated that on average (across bins) 77% of cells remained viable (standard deviation:

18.15%). We also estimated the genetic diversity of the library through Sanger sequencing of

DNA from individual colonies (NightSeq® service, Microsynth, Switzerland).

1184

To validate our binning procedure, we re-grew binned cultures from either overnight

recovered cultures or frozen aliquot stocks. We then measured their expression distributions

by flow cytometry, which reproduced the original expression measurements. This approach

allowed us to compare the fluorescence distributions of the re-grown cultures with the pre-

sorting distributions. Specifically, we assessed whether the geometric mean of the

fluorescence distribution for each sorting bin matched those recorded during the initial

sorting procedure. Next, we re-sorted cells derived from each bin in order to eliminate cell

cross-contaminations and other factors that could alter the bin distributions. The geometric

mean is often preferred over the arithmetic mean in this type of analysis, because it is less

affected by the presence of outliers in the data [175,176]. It is also a more accurate representation

of the central tendency of data that is log-normally distributed, which is often the case with

flow cytometry fluorescence measurements[175,176].

1197

6. DNA extraction and sequencing

We diluted 500 uL of individual glycerol stocks of each replicate subpopulation of sorted

cells (i.e., cells from each bin of fluorescence intensity) in 5mL of LB supplemented with

chloramphenicol in 15mL Falcon tubes, and grew the resulting cell culture overnight (16

hours, 37°C, 220 rpm). On the next day, we isolated plasmids from each culture using a

QIAprep® spin miniprep kit (Qiagen, Germany). In order to allow the sequencing of multiple

pooled samples (multiplexing), we barcoded our regulatory region through PCR with specific

HPLC-purified primers (**Supplementary Table S6**) provided by Eurofins (Konstanz,

Germany). We added barcodes to the 5'region of the amplicon through a PCR reaction. We

performed this PCR reaction with the Q5 high-fidelity polymerase, and did so in triplicate for

each miniprep-isolated plasmid library. To calculate primer melting temperatures (Tm), we

used the NEB Tm calculator (https://tmcalculator.neb.com/#!/main) for a primer

concentration of 500nM. We performed the PCR with the following program: 98°C/30 s; 25

cycles of 98°C/10 s, 64°C/30 s and 72°C/30 s; and 1 cycle of 72°C/2 min.

1212

After PCR amplification, we digested the reaction products with the restriction enzymes DpnI

(NEB #R0176L) and Exonuclease I (NEB #M0293L) in order to remove traces of genomic

DNA, plasmids, and single-stranded DNA that could interfere with sequencing. The Master

Mix we used for a single digestion harbored 1μL of 10x CutSmart® Buffer (NEB #B6004S),

1μL of Exonuclease I (NEB #M0293L), 1μL of DpnI restriction enzyme (NEB #R0176L),

and 7μL of distilled nuclease-free water. For each PCR product, we added 10μL of the Master

Mix. We incubated the reaction for 1 hour at 37°C, following 15 minutes at 80°C for

deactivation of the enzymes.

1221

We purified the digestion products using the Monarch® DNA PCR/Gel Extraction Kit (NEB

#T1020L) and fractionated them through gel electrophoresis to confirm that only a single

band with a size ≈150bp was present. After this confirmation, we pooled the purified PCR

products from the different bins of each replicate sorting equimolarly to a total mass of 2,600

ng and a volume of 100 μL (26ng/μL of DNA) in 1.5mL Eppendorf tubes. We then sent the

pooled purified amplicons for adapter ligation and sequencing at Eurofins (NGSelect

Amplicons® on Illumina HiSeq), obtaining 15 million paired-end reads (2 × 150 bp) for all

samples.

1230

1231 We quantified the total number of reads (*t*) required for sequencing based on several factors,

1232 including the number of bins (*b)*, biological replicates (*r)*, strains (*s)*, genotypes (*g)*, and the

1233 reads per genotype (*p)*. Using these variables, we first calculated the total number n_{bc} of

1234 needed barcodes as follows:

$$n_{bc} = b \times r \times s$$

1235 For each of our three TFs, the values of these parameters are *b*=13 bins, *r*=3 biological

1236 replicates, and *s*=1 strains, yielding n_{bc}=39. The total number of required sequence reads can

1237 be estimated through the equation:

$$t = r \times g \times p$$

1238 where we aimed for *p*=30 paired-reads per genotype. The number *g* of expected genotypes

1239 per library is the same (65,536) for each TF. These values lead to a total of $t = 3 \times 65,536 \times$

1240 30 = 7,667,712 required paired-end reads for each library.

1241

7. Data analysis:

7.1 Filtering and preparing sequencing reads

We processed the sequencing data with a blend of custom python and awk scripts,

complemented by established bioinformatics utilities. Firstly, we trimmed sequences by

computationally excising Illumina adapters, followed by merging paired-end reads. We then

organized the paired-end reads into distinct files, each tagged with a sequence barcode

identifying the sequence bin from which the corresponding reads originated.

1249

For the removal of Illumina adapter sequences from the paired-end reads, we employed

Cutadapt [177] with the following parameters:

*cutadapt -j 8 -e 0.1 --no-indels --overlap=8 --discard-untrimmed *

*-a "^\\$FWD...\\$REV_RC;max_error_rate=0.2;min_overlap=6" *

*-A "^\\$REV...\\$FWD_RC;max_error_rate=0.2;min_overlap=6" --pair-filter=any *

*-o 'Sample_${sample}_trimmed_1.fastq.gz' *

*-p 'Sample_${sample}_trimmed_2.fastq.gz' *

*--max-ee=2 -l=114 *

$reads

1259

Here, "ADAPTER_FWD" and "ADAPTER_REV_RC" are placeholders for the actual

adapter sequences used, which are 5'-

AGATCGGAAGAGCACACGTCTGAACTCCAGTCA-3' for read 1, and 5'-

AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT-3' for read 2.

Because our amplicons were short (153 bp) and each sequencing sample was divided into two

files with overlapping single-end reads, merging the two files (single-end reads) was

necessary, for which we used the FLASH software [178]. After the filtering and merging of

reads, we demultiplexed paired-end reads with the following FLASH parameters:

flash -t $ncores $reads -O -m 60 -M 140 -z -o 'Sample_${sample}_merged'

Next, we utilized the FastX toolkit (http://hannonlab.cshl.edu/fastx_toolkit/)[179] to discern and

preserve only those sequences that exceeded a high-quality threshold of Q = 33.

Subsequently, we refined our dataset by removing sequences that contained undesirable

mutations or insertions/deletions (indels) within the region extending from the start of the

core promoter to the end of the variable TFBS library. We performed this filtering step using

a custom R [180] script. Thereafter, we used custom awk and R[180] scripts to convert the data into

a table. Each row of this table contains data from one TFBS in the library, and each column

contains the number of reads for this TFBS from one of the thirteen bins into which we had

sorted cells. We estimated the average regulation strength of each TFBS from this data.

1278

7.2 Calculating regulation strengths:

We calculated regulation strengths as previously described[27]. Briefly, in sort-seq experiments,

sequences often appear in multiple fluorescence bins due to random mis-sorting[48,137] and the

stochastic nature of gene expression[165]. Following methods established in previous research

[39,49,103], we determined the reporter expression level driven from each TFBS variant by

calculating a weighted average of bins in which the variant occurred. This involved

multiplying the frequency of each sequence (x_i) in a given bin i by a numerical value

representing that bin (w_i=1,2,3,…,13), and then averaging these products over the total

sequence count. Mathematically, this weighted average calculates as

$e = \sum_{i=1}^{n} \frac{(x_i * w_i)}{\sum_{i=1}^{n} x_i},$

where e is the expression level driven by the sequence.

1290

1291   This approach yields a continuous spectrum of expression levels e within the range of 1 to 13.

1292   High expression values indicate robust GFP expression and, thus, weak binding of a TF to a

1293   TFBS variant. To facilitate interpretation, we inverted this scale so that higher values indicate

1294   stronger repression and lower GFP expression, i.e., we define the regulation strength b

1295   (strength of regulation) as follows:

1296

   $b = e_{max} + 1 - e$

1297

where e_{max}=13 is the maximal expression level (maximal bin value). We then normalized b by

the maximal regulation strength among all TFBSs we studied for a given TF. The resulting

normalized regulation strength scores b vary from 0 to 1, where low scores denote TFBS

variants with weak TF binding (weak reporter repression), and high scores indicate variants

with strong TF binding (strong reporter repression). A score of b =1 signifies the highest

regulation strength (repression) calculated among all TFBSs for a given TF.

1304

1305

7.3 Combining data from triplicates:

As a quality-filtering step, we eliminated TFBS variants that did not appear in all three

replicates or that were represented by less than 30 reads in total (summing their read counts

over the thirteen bins). Given that we have 13 bins and most sequences are typically found

within an average of 3.4 bins, we wanted a minimum of 10 reads per bin to prevent

misinterpretation of repression levels and also guide our threshold's choice. This procedure

reduced the fraction of TFBS variants for which we had sort-seq-based sequence data from

95%, 90%, and 93% of the total library size (65,536 sequences) for CRP, Fis and IHF,

respectively, to 49%, 66% and 63%.

1315

Subsequently, we calculated the regulation strengths b of the remaining TFBS variants

(**Supplementary Materials and Methods 7.2**). Then, we determined the coefficient of

variation of regulation strengths across replicates for each TFBS variant, which estimate the

consistency of measured regulation strengths among replicates. Notably, most TFBS variants

exhibited a coefficient of variation (CV) below 0.5, which was the threshold we set for

filtering sequences. Sequences with a CV above 0.5 were excluded from the analysis. This

threshold is commonly used in transcriptional studies employing fluorescent reporters to

account for transcriptional noise and measurement variability [165]. Finally, we averaged

regulation strengths for each TFBS variant across all replicates and normalized the resulting

averages by the maximal observed regulation strength among all TFBSs of a given TF.

1326

7.4 Frequency matrices and sequence logos:

We generated frequency matrices of nucleotides that occur in TFBSs from each fluorescence

bin by counting the frequency of each nucleotide at each variable position of the TFBS

library. From this data, we generated heatmaps and DNA sequence logos representing the

frequency matrices graphically. A sequence logo consists of a stack of the letters A, C, G, and

T, at each position of a DNA sequence, where the relative size of each letter indicates its

frequency in the sequence. The total height of the stack corresponds to the information

content of that position, in bits [68,181]. A sequence logo is a graphical representation of the

informational properties of a TFBS. When mutated, nucleotides with high information

content are more likely to lead to a loss of binding (repression) than nucleotides with low

information content.

7.5 Creation of genotype networks and determining network metrics:

We used in-house Python script and the Python package *igraph*[182] to generate directed

genotype networks. These are graphs in which TFBS variants are nodes (vertices, genotypes),

and variants that differ in a single nucleotide are connected by an edge. Each node of this

network is associated with the corresponding DNA sequence and the associated regulation

strength. Each edge is directed, i.e., it corresponds to a binding-score-increasing mutation,

and points from a TFBS variant with lower regulation strength to a neighbor with higher

regulation strength. We extracted the largest weakly connected subgraph (the"giant

component" [6]) of the network, and used it for all further analyses. This giant component

comprises the vast majority (99%) of sequenced genotypes for all of our TF landscapes. We

used in-house Python and R scripts for all network analyses described below.

Epistasis: Epistasis refers to non-additive interactions between two or more mutations. It can

impose severe constraints on molecular evolution, because the mutations that are beneficial in

one genetic background may be deleterious in another[37]. Epistasis between two mutations can

be classified as magnitude, simple sign, or reciprocal sign epistasis, depending on the sign

(i.e., positive or negative) of the fitness effect of individual mutations and their

combinations[37]. In magnitude epistasis, the effect of a mutation on regulation strength varies

depending on the genetic background but the sign of this effect (increasing or decreasing

regulation strength) does not. Simple sign epistasis occurs if one single mutant has a lower

regulation strength than both the wild type and the double mutant, while the other single

mutant has a regulation strength that is intermediate to the wild type and double mutant.

Reciprocal sign epistasis occurs when both mutations independently decrease regulation

strength, but their combination increases regulation strength. The presence of reciprocal sign

epistasis is a necessary condition for the existence of multiple peaks in an adaptive landscape

[75,76].

To determine the incidence of epistasis in our landscapes, we employed a method that

involves identifying all "squares" in a genotype network with the *motifs* function from the

igraph library in R. Each square consists of a "wild-type" sequence, a double-nucleotide

mutant, and the corresponding two single mutants. We assessed epistasis for each square

along a single axis by selecting the highest- regulation strength sequence as the double

mutant. Our analysis placed each square into one of three categories: no sign epistasis, simple

sign epistasis, and reciprocal sign epistasis. The no sign epistasis category included both

magnitude epistasis and additivity (no epistasis) without differentiating between them,

because neither affects the accessibility of landscape peaks through only binding-increasing

mutations[11,108]. We determined the proportion of all squares that fell into each category (see

Supplementary Table S5).

1374

1375 _Peaks:_ A peak is a genotype (TF binding site variant) whose neighbors all convey lower

1376 regulation strength than itself. Two peaks are connected if they are neighbors and convey the

1377 same regulation strength. We refer to the genotype with the highest regulation strength as the

1378 summit or global peak[6,183].

1379

Genotype connectivity: To evaluate the sparsity of our networks and the likelihood of peak

misclassification due to the incompleteness of our landscapes, we used a metric called

"genotype connectivity." This represents the number of immediate (1-mutant) neighbors of a

given genotype for which our experiment produced repression strength data. A perfect

experiment creating a complete landscape data set would provide repression data for each of

our 4^8 genotype and all of its 24 immediate neighbors. In practice, however, data for some

genotypes and some of their neighbors is unavailable. We quantified the "relative

connectivity" of any one genotype as the ratio of the actual number of adjacent genotypes for

which we have repression data to the theoretical maximum of 24 neighbors. This measure

helps us to address questions about the sparsity of our landscape and its implications.

Specifically, it allows us to determine if our sample is biased, with some regions or genotypes

being more connected than others. This is particularly important for assessing whether peaks

and high peaks are less connected than non-peaks, which could potentially lead to peak

misassignments.

Accessible Paths: In the context of our genotype networks, we call a mutational path

accessible if the regulation strength increases with each mutational step [6,183]. We

systematically enumerated all shortest accessible paths, distinguishing them by their path

length, i.e., by the number of mutational steps in a path. Notably, there can be multiple

alternative accessible paths with the same number of mutational steps from a single starting

genotype to a peak.

1400

1401 *Basins of attraction:* The basin of attraction of a peak comprises all TFBS variants from

1402 which accessible paths to the peak exist. We refer to the basin's size as the number of

1403 variants in the basin. We determined basin sizes by exhaustive enumeration.

1404

1405 Overlap between basins. The basins of attraction of different peaks may comprise

1406 overlapping sets of variants. To determine the overlap between two basins B_1 and B_2, we

1407 used the Jaccard index J [184,185], which is equal to the size of the intersection between two sets

1408 of variants divided by the size of their union:

$$J = \frac{B1 \cap B2}{B1 \cup B2}$$

7.6 Generating randomly shuffled landscapes

To generate uncorrelated random landscapes for our three TF landscapes, we performed a

random shuffling of regulation strength values from our genotypes, followed by landscape

analysis to identify peaks. To this end, we employed an ad-hoc python script to randomly

permute experimentally measured regulation strength values among all genotypes in each of

our landscape. This procedure preserves the distribution of regulation strengths in each

landscape, but also creates an uncorrelated random landscape [73,79]. Following random

shuffling, we analyzed each shuffled landscape's topography with the same methods we had

applied to the original (unshuffled) landscape to identify its peaks. To ensure statistical

robustness, we repeated the random shuffling and peak identification 100 times. We then

compared the number of peaks between the 100 shuffled landscapes and the original

(unshuffled) landscape.

7.7 Principal component analysis

In order to investigate which nucleotides in a TFBS are most important for changes in

regulation strengths, we performed a principal component analysis for each landscape

(**Supplementary Figures S19-S21**). To this end, we first one-hot-encoded our data. This

encoding represents each categorical value (nucleotide at each position of a DNA string) as a

binary vector of length 4. It thus converts an entire DNA string of length L into a 4xL binary

matrix. We used this binary matrix to perform PCA with the R base function *prcomp*.

1428

8. Simulated adaptive walks:

We simulated the adaptive evolution of a population on our landscapes by performing two

different types of random walks, "greedy" adaptive random walks, and random walks based

on Kimura's model of fixation probabilities.

For all simulations, we assumed that only point mutations occur and that the time it takes for

a point mutation to become fixed in a population is much shorter than the time it takes for a

new mutation to appear that will eventually go to fixation. This scenario is also called the

strong selection weak mutation (SSWM) scenario [39,81–83]. Under this scenario, evolving

populations are monomorphic most of the time, i.e., all individuals have the same genotype.

This scenario is realistic when the product of effective population size N and mutation rate μ

is small ($N\mu < 1$), which is the case for *E. coli* (N=1.8×10^8, μ=2×10^{-10}) [43]. It allows us to

model adaptive evolution as an adaptive random walk in our landscapes. Our simulations also

account for mutation bias, which means that different types of mutation have a different

probability of occurring in a population. We use mutation biases that were experimentally

determined for *Escherichia coli*[86].

In a greedy adaptive random walk, starting from any one genotype, only the mutational

neighbor that conveys the largest fitness advantage is fixed in a population. We initiated one

greedy random walk from each non-peak genotype and terminated the walk once a fitness

peak was reached, i.e., once every neighbor of the current genotype had lower fitness than the

genotype itself. Because every greedy random walk is deterministic in the absence of neutral

mutations, it is sufficient to initiate one greedy random walk per starting genotype.

Another well-established model for random walks allows for genetic drift. It uses fixation

probabilities computed by Kimura[80,186,187] i.e., $f_{ij} = (1 - e^{-2s}) / (1 - e^{-2Ns})$, where f_{ij} is the

probability of fixing mutation j in the background of genotype i, N is the effective population

size, and s is the selection coefficient, i.e. the difference in fitness between genotypes i and

j[80,186]. For a given pair of genotypes, the only parameter of this model is the effective

population size N, for which we explored values of N=10^8, N=10^5 and N=10^2. Generally, the

smaller the population size is, the larger is the probability that neutral or deleterious

mutations become fixed in a population.

For these "Kimura" adaptive walks, we chose 15,000 random starting genotypes for each of

the three values of N, and simulated 1,000 random walks for each of them. At each step, we

randomly picked a mutation j, generated a random number in the interval [0, 1], and

considered the mutation to become fixed if the random number fell within the interval [0, f_{ij}].

Computationally, we accelerated this process by precomputing all fixation probabilities for

all genetic backgrounds, and then used the *python* function *numpy.choice* to generate a

random sample from the multinomial distribution of fixation probabilities at each step[188].

Because of genetic drift, Kimura's random walks do not necessarily terminate when they

reach a fitness peak. For this reason, we simulated each such random walk for a maximum of

25 mutational steps.

SUPPLEMENTARY TABLES

1469

1470   **Table S1. Plasmids used in this study**

Name	Antibiotic to select for plasmid persistence (concentration in μg/ml)	Relevant features	Source
pCAW-Sort-Seq-V2	Chloramphenicol (50)	pBBR1, TetR, *sfgfp, mscarlet-I*, no TFBS, no TF	This study, derived from [27]
pCAW-Sort-Seq-V2-CRP_lib-CRP	Chloramphenicol (50)	pBBR1, TetR, *sfgfp, mscarlet-I,* , CRP operator, *crp*	This study
pCAW-Sort-Seq-V2-Fis_lib-Fis	Chloramphenicol (50)	pBBR1, TetR, *sfgfp, mscarlet-I,* , Fis operator, *fis*	This study
pCAW-Sort-Seq-V2-IHF_lib-IHF	Chloramphenicol (50)	pBBR1, TetR, *sfgfp, mscarlet-I,* , IHF operator, *ihf*	This study

1471

1472 **Table S2. Table of strains**

Strain	Genotype	Antibiotic resistance	Reference
SIG10-MAX from Sigma Aldrich *Cloning strain*	F- mcrA Δ(mrr-hsdRMS-mcrBC) endA1 recA1 Φ80dlacZΔM15 ΔlacX74 araD139 Δ(ara,leu)7697galU galK rpsL nupG λ- tonA (StrR)	Streptomycin	Sigma Aldrich
E. coli DH5α *Cloning strain*	F- endA1 gln V44 thi-1 recA1 relA1 gyrA96 deoR nupG Φ80dlacZΔM15 Δ(lacZYA-argF)U169, hsdR17(rK − mK +), λ−	None	189
E. coli JW5702-4 *Δcrp mutant strain.*	Δ(araD-araB)567, ΔlacZ4787(::rrnB-3), λ−, Δ*crp*-765::kan, rph-1, Δ(rhaD-rhaB)568, hsdR514.	Kanamycin	164
E. coli JW1702-1 Δ*ihfA* mutant strain.	Δ(araD-araB)567, ΔlacZ4787(::rrnB-3), λ−, ΔihfA786::kan, rph-1, Δ(rhaD-rhaB)568, hsdR514.	Kanamycin	164
E. coli JW3229-1 Δ*fis* mutant strain	bΔ(araD-araB)567, ΔlacZ4787(::rrnB-3), λ−, Δfis-779::kan, rph-1, Δ(rhaD-rhaB)568, hsdR514.	Kanamycin	164

1473

1474   **Supplementary Table S3. Libraries used in the present study.**

1475

Sequence Name	Sequence*	TF	Library size
CRP	TGT**NN**TCTAGA**NNNNN**T**N**TT	CRP-Fis	$4^8 = 65,536$
Fis	**N**CTC**N**TT**NNNN**AAG**NN**A	CRP-IHF	$4^8 = 65,536$
IHF	T**NNNNN**ATGAG**NNN**	Fis-IHF	$4^8 = 65,536$

1476

1477   *IUPAC nucleotide nomenclature for degenerate nucleotides: **N**: A, C, T or G

1478

1479 **Supplementary Table S4. TFBS reference ("wild-type", WT) sequences used in the**

1480 **present study.**

1481

Sequence Name	Sequence	TF	Reference
WT_{CRP}	TGTGATCTAGATCACATTTT	CRP	[63,190]
WT_{Fis}	GCTCATTTTTTAAGCAA	Fis	[61,191,192]
WT_{IHF}	TAGTTTATGAGGTA	IHF	[62,65]

1482

1483

1484 **Table S5. Global landscape properties**

	Number of genotypes[1]	Number of peaks[2]	Number of low peak[3]	Number of high peaks[4]	Number of squares[5]	Magnitude epistasis or additivity[6,7]	Simple sign epistasis[7]	Reciprocal sign epistasis[7]
CRP	31,975 (49%)[1]	2,154 (7%)	2,093 (97.2%)	61 (2.8%)	580,289	37%	34%	28%
Fis	43,222 (66%)[1]	2,312 (5%)	2,140 (92.6%)	172 (7.4%)	1,194,144	34%	34%	32%
IHF	41,325 (63%)[1]	2,453 (6%)	2,254 (91.9%)	199 (8.1%)	1,026,818	34%	33%	32%

1485

[1] Percentages indicate the proportion of the studied library relative to its expected size.

[2] Percentages indicate the proportion of peaks relative to the total number of genotypes

[3] Low peaks are peaks with regulation strengths below the wild sequence. Percentages refer

to the proportion of all peak genotypes.

[4] High peaks are peaks with regulation strengths above the wild-type sequence. Percentages

refer to the proportion of all peak genotypes.

[5] A square represents a quadruplet of sequences containing a focal sequence (ab), one double

mutant of this sequence (AB), and the two intermediate single mutants (Ab and aB).

[6] This category includes both magnitude epistasis and additivity (no epistasis) without

distinguishing them, because neither of the two subcategories affects peak accessibility[11,108]

[7] Percentages refer to the proportion of all squares.

1497

1498 **Table S6. Primers for engineering the pCAW-Sort-Seq-V2 plasmid and for cloning**

1499 **libraries.**

Name	Sequence	Usage
pCAW_frag1_F	5'-CGTCCGACTTACGGAAGGTAGATTTTACGGC-3'	Linearizing the pCAW-Sort-Seq fragment1 for Gibson Assembly
pCAW_frag1_R	5'-CTCGTGCCTAACGGAAGGTAGATTTTACGGC-3'	Linearizing the pCAW-Sort-Seq fragment1 for Gibson Assembly
pCAW_frag2_F	5'-TAAGATTGCCACGGAAGGTAGATTTTACGGC-3'	Linearizing the pCAW-Sort-Seq fragment2 for Gibson Assembly
pCAW_frag2_R	5'-AGGCCTGACTACGGAAGGTAGATTTTACGGC-3'	Linearizing the pCAW-Sort-Seq fragment2 for Gibson Assembly
Ultramer_ds_F	5'-TTCTCAAAAGCTTCCTGCAGTATTC-3'	Amplifying Ultramer® libraries
Ultramer_ds_R	5'-CGGAAAGCACATCCGGTGAC-3'	Amplifying Ultramer® libraries
TFBS_R	5'-CCGTTTGTAGCATCACCTTC-3'	Sequencing the TFBS region
pCAW_Gibs_Lib_F	5'-GTCTGATGAGTCCGTGAGGACG-3'	Linearizing the pCAW-Sort-Seq plasmid
pCAW_Gibs_Lib_R	5'-GAGAAAAGAAAACCGCCGATCCTG-3'	Linearizing the pCAW-Sort-Seq plasmid
Ultramer_Gibs_F	5'-GGTGGACAGGATCGGCGGTTTTCTTTTCTCTTCTCAAAAGCTTCCTGCAGTATTC-3'	Amplifying Ultramer® libraries for Gibson Assembly

| Ultramer_Gibs_R | 5'-GGCTGTTTCGTCCTCACGGACTCATCAGACCGGAAAGCACATCCGGTG-3' | Amplifying Ultramer® libraries for Gibson Assembly |

1500

1501

1502 **SUPPLEMENTARY FIGURES**

1503

1504

Supplementary Figure S1. Construction of pCAW-Sort-Seq-V2. The plasmid pCAW-Sort-

Seq-V2 is a derivative of plasmid pCAW-Sort-Seq[27]. The figure illustrates the sequential

cloning steps involved in the plasmid's construction, from the bottom upwards. For an in-

depth explanation, see **Supplementary Methods**. Briefly, we began with the synthesis and

cloning of a codon-optimized *mscarlet* gene expression cassette into the pCAW-Sort-Seq

plasmid, oriented antiparallel to the tetracycline resistance gene (*tetr*). This generated the

intermediate plasmid pCAW-Sort-Seq-mscarlet plasmid. Subsequently, we inserted each TF

gene, exemplified by the *crp* gene for the purpose of this representation, upstream of the

mscarlet gene in a bicistronic operon configuration. This step created the TF-specific

expression plasmid variants (one each for CRP, Fis and IHF). The final stage, not depicted

here, involved the incorporation of the appropriate wild-type TFBSs and libraries into each

plasmid variant upstream of the *gfp* gene, which yielded the library plasmids we utilized in

our sort-seq experiments.

1518

1519

1520

1521

1522

1523

1524

Supplementary Figure S2. Components of plasmid pCAW-Sort-Seq-V2. The plasmid

system **pCAW-Sort-Seq-V2** is a derivative of the plasmid pCAW-Sort-Seq[27]. It encodes the

broad-host, low-copy number replication origin pBBR1 [170] (5 to 10 copies per cell), together

with an origin of transfer (OriT) for conjugation (blue), and a chloramphenicol resistance

gene (yellow). The interchangeable regulatory region where the TFBS is located (cyan) is

placed between a constitutive promoter (BBa_J23110[193]) and a superfolder GFP (*sfgfp*, green)

fluorescent reporter gene[172]. The transcriptional insulator RiboJ[173] is located upstream of the

sfgfp gene. The *tetr* gene (purple) is derived from the original Tn10 transposon[51,73] under the

control of a low-strength constitutive promoter (pLac promoter variant developed in ref. [194]).

TetR regulates the expression of a bicistronic operon encoding the TF of interest (represented

by CRP, dark purple), and an mScarlet-I red fluorescent reporter[174] (red). Promoters and their

respective insulators[195] are represented as white arrows. Transcriptional terminators are

represented as black arrows[196]. Ribosome binding sites (RBSs) are represented as pink

arrows.

1539

1540

Supplementary Figure S3. Growth profile for strains used in this study. Growth profiles

for all strains used for landscape mapping are represented as optical densities (OD$_{600}$, y-axis)

measured over 7 hours of growth at 1-hour intervals (x-axis). We transformed all strains with

the plasmid pCAW-Sort-Seq-V2 and grew them in LB medium without inducers. Colors

(legend) represent the different strains. Wild-type (purple) refers to the BW25113 parent-

strain from which the mutants (Δ*crp*, *Δfis*, Δ*ihf*) were derived. Each circle represents the

mean of three biological replicates independently measured in a plate reader. Shaded areas

correspond to one standard deviation calculated from the three replicates.

1549

1550

1551

1552

1553

1554

1555

1556

1557

Supplementary Figure S4. Sorting procedure for CRP library. Each density plot is based

on a histogram representing the distribution of GFP fluorescence for 10^6 cells. The histogram

represents data from one replicate, and the same sorting strategy was applied to the other two

replicates. Specifically, the horizontal axis shows GFP fluorescence (FITC-H values, arbitrary

units, $\log_2$ scale), while the vertical axis indicates the relative frequency of cells with a given

fluorescence value. Heatmap colors indicate fluorescence, from low (dark purple) to high

(yellow). The vertical axis indicates the three experimental conditions under which we

measured fluorescence: 1) negative control ("Neg"), without GFP expression (promoterless

pCAW-v2); 2) CRP library in pCAW-v2 without TF expression ("Lib -CRP") and 3) CRP

library in pCAW-v2 with TF expression ("Lib +CRP"). To create each density plot from the

histogram data, we performed density smoothing using a Gaussian kernel function with the

aid of the *ggplot2* package[197]. The vertical dashed grey line marks the upper limit of cell

autofluorescence (negative control). We did not select cells with fluorescence below this

threshold for sorting. Rectangular boxes (1 to 13) represent fluorescence bins/gates into

which we sorted *E.coli* cells, spaced equally on a binary logarithmic ($\log_2$) scale.

1573

Supplementary Figure S5. Sorting procedure for Fis library. Each density plot represents the distribution of GFP fluorescence for 10^6 cells. The histogram represents data from one replicate, and the same sorting strategy was applied to the other two replicates. Specifically, the horizontal axis shows GFP fluorescence (FITC-H values, arbitrary units, $\log_2$ scale), while each histogram indicates the relative frequency of cells with a given fluorescence value. Heatmap colors indicate fluorescence, from low (dark purple) to high (yellow). The vertical axis indicates the three experimental conditions under which we measured fluorescence: 1) negative control ("Neg"), without GFP expression (promoterless pCAW-v2); 2) Fis library in pCAW-v2 without TF expression ("Lib -Fis") and 3) Fis library in pCAW-v2 with TF expression ("Lib +Fis"). We performed density smoothing using a Gaussian kernel function to create a smooth density plot for each histogram with the aid of the *ggplot2* package [197]. The vertical dashed grey line marks the upper limit of cell autofluorescence (negative control). We did not select cells with fluorescence below this threshold for sorting. Rectangular boxes (1 to 13) represent bins/gates for each sample, spaced equally on a binary logarithmic ($\log_2$) scale.

1589

1590
Supplementary Figure S6. Sorting procedure for IHF library. Each density plot

represents the distribution of GFP fluorescence for 10^6 cells. The histogram represents data

from one replicate, and the same sorting strategy was applied to the other two replicates.

Specifically, the horizontal axis shows GFP fluorescence (FITC-H values, arbitrary units, $\log_2$

scale), while each histogram indicates the relative frequency of cells with a given

fluorescence value. Heatmap colors indicate fluorescence, from low (dark purple) to high

(yellow). The vertical axis indicates the three experimental conditions under which we

measured fluorescence: 1) negative control ("Neg"), without GFP expression (promoterless

pCAW-v2); 2) IHF library in pCAW-v2 without TF expression ("Lib -IHF") and 3) IHF

library in pCAW-v2 with TF expression ("Lib +IHF"). We performed density smoothing

using a Gaussian kernel function to create a smooth density plot for each histogram with the

aid of the ggplot2 package156. The vertical dashed grey line marks the upper limit of cell

autofluorescence (negative control). We did not select cells with fluorescence below this

threshold for sorting. Rectangular boxes (1 to 13) represent bins/gates for each sample,

spaced equally on a binary logarithmic ($\log_2$) scale.

1606

1607

1608

1609

Supplementary Figure S7. Correlation of read counts among replicates of the CRP library. The figure presents a pairwise comparison of log-transformed read counts across three experimental replicates for CRP. Each circle represents a distinct genotype, plotted according to its read counts in two sequencing replicates (x-axis for one replicate and y-axis for the other). The density of genotypes within the plot is indicated in the legend, with purple representing lower-density regions and yellow representing higher-density regions. The red line across each plot depicts a linear model's fit to the data, illustrating the association between the read counts of genotypes across replicates. Pearson correlation coefficients (upper triangle) are indicated for each pair of replicates. Their values are close to 1, demonstrating high reproducibility.

Supplementary Figure S8. Correlation of read counts among replicates of the Fis library. The figure presents a pairwise comparison of log-transformed read counts across three experimental replicates for Fis. Each circle represents a distinct genotype, plotted according to its read counts in two sequencing replicates (x-axis for one replicate and y-axis for the other). The density of genotypes within the plot is indicated in the legend, with purple representing lower-density regions and yellow representing higher-density regions. The red line across each plot depicts a linear model's fit to the data, illustrating the association between the read counts of genotypes across replicates. Pearson correlation coefficients (upper triangle) are indicated for each pair of replicates. Their values are close to 1, demonstrating high reproducibility.

Supplementary Figure S9. Correlation of read counts among replicates of the IHF library. The figure presents a pairwise comparison of log-transformed read counts across three experimental replicates for IHF. Each circle represents a distinct genotype, plotted according to its read counts in two sequencing replicates (x-axis for one replicate and y-axis for the other). The density of genotypes within the plot is indicated in the legend, with purple representing lower-density regions and yellow representing higher-density regions. The red line across each plot depicts a linear model's fit to the data, illustrating the association between the read counts of genotypes across replicates. Pearson correlation coefficients (upper triangle) are indicated for each pair of replicates. Their values are close to 1, demonstrating high reproducibility.

1643

Supplementary Figure S10. Regulation strength distribution across the libraries a. CRP library. The histogram illustrates the distribution of regulation strengths (normalized to the maximally observed value) among 31,975 library variants. The average regulation strength across all variants is highlighted by a dashed blue line (mean = 0.37), while the wild type's (WT = 0.71) strength is denoted by a dashed red line. A total of N=84 variants surpass the WT in regulation strength. **B. Fis library.** The histogram illustrates the distribution of regulation strengths (normalized to the maximum observed value) among 43,222 library variants. The average regulation strength across all variants is highlighted by a dashed blue line (mean = 0.57), while the wild type's (WT = 0.97) strength is denoted by a dashed red line. A total of N=172 variants surpass the WT in regulation strength. **c. IHF library.** The histogram illustrates the distribution of regulation strengths (normalized to the maximum observed value) among 41,325 library variants. The average regulation strength across all variants is highlighted by a dashed blue line (mean = 0.52), while the wild type's (WT = 0.95) strength is denoted by a dashed red line. A total of N=199 variants surpass the WT in regulation strength.

1659

1660

1661

Supplementary Figure S11. Validating regulation strengths. In this analysis, we validated

the regulation strengths of 10 selected TFBSs for each bin and for each library (130

sequences per library). These TFBSs covered a wide range of regulation strengths determined

from sort-seq data. The horizontal axis of each panel shows the regulation strengths of the

130 binding sites as quantified through sort-seq. The vertical axis shows the fluorescence

levels of the same binding sites, as quantified by a plate reader (arbitrary units). The top of

each panel shows the result of a test of the null-hypothesis that the Pearson correlation

coefficient R between the two quantities is equal to zero, together with a 95% confidence

interval. The shaded grey area surrounding the regression line indicates the 95% confidence

interval, i.e., the probability that the true linear regression line of the population lies within

the confidence interval of the regression line calculated from the sample data at a confidence

level of 95%. R^2 indicates the coefficient of determination. Each circle in the plot represents

the mean of three independent (biological) replicate fluorescence measurements in a plate

reader after eight hours of growth. Circle colors indicate GFP fluorescence from low (dark

green) to high (light green **a. Association between plate reader fluorescence and**

regulation strengths for the CRP library. R = -0.81, P < 0.001; R² = 0.66, P < 0.001, N=

130. **b. Association between plate reader fluorescence and regulation strengths for the**

Fis library. R = -0.74, P < 0.001; R² = 0.54, P < 0.001, N=130. **c. Association between plate**

reader fluorescence and regulation strengths for the IHF library. R = -0.73, P < 0.001; R²

= 0.53, P < 0.001 N = 130

1682

1683

Supplementary Figure S12. Adaptive peaks and their regulation strengths across TF

landscapes. a. Adaptive peaks in the CRP landscape. The distribution of regulation

strengths across the landscape's 2,154 adaptive peaks is shown. The WT's regulation strength

(0.71) is marked by a horizontal dashed red line for reference. A total of N=61 peaks have

higher regulation strength than the WT. The color gradient indicates genotype density along

the regulation strength axis, with the histogram on the right detailing the regulation strength

distribution. **b)** like a) but for the **Fis landscape,** based on its 2,312 adaptive peaks. (WT

regulation strength: 0.97) N=172 peaks regulate expression more strongly than the WT. **c)**

like b) but for the IHF landscape, based on its 2,434 adaptive peaks. (WT regulation strength:

0.95) N=199 peaks regulate expression more strongly than the WT.

Supplementary Figure S13. Quantitative analysis of CRP landscape sparsity. a. Distribution of relative connectivity among genotypes. Violin plots augmented with embedded boxplots illustrate the distribution of relative connectivity, i.e,. the ratio of the

empirically observed number of adjacent genotypes to the theoretical maximum number of

24(=8×3) within a fully connected network. Genotypes are stratified into the three categories

"other" (non-peak genotypes), "peaks" (excluding high peaks), and "high peaks". Mean

relative connectivities and standard deviations, are shown above each categorical division,

revealing lower relative connectivity in peaks relative to non-peak genotypes (Two-sample

Welch's t-test, p-value = 4.59×10^{-77}, N1= 29,821; N2 = 2,093). High peaks exhibit

comparable or superior connectivity relative to non-peak genotypes (Two-sample Welch's t-

test, p-value = 1, N1 = 29,821; N2 = 61). **b. Weak association between connectivity and**

regulation strength. Scatterplot of relative connectivity against regulation strength. The

density of genotypes (circles) in the scatterplot area is represented by a color gradient from

purple (low density) to yellow (high density). The weak association (Pearson correlation, R =

-0.13; t = -23.888; degrees of freedom (df) = 31,973; p-value < 2.2×10^{-16}) indicates that

strongly regulating genotypes dare not necessarily densely connected. Marginal histograms

adjacent to the scatterplot quantify the distributions of regulation strength and relative

connectivity. **c. Visualization of CRP landscape based on regulation strength.** Visual

overview of the CRP landscape with genotypes color-coded according to regulation strength

(low strength: purple; medium: green; high: yellow; grey: genotypes with missing data). To

enhance clarity, edges between genotypes are not shown. The landscape is projected onto a

2D space through a force-directed algorithm[182], with axes representing arbitrary units. **d.**

Spatial distribution of regulation strengths. Each panel shows a density contour plot of the

distribution of regulation strength scores of genotypes within one of four regulation strength

categories (no data [missing data], low, medium, high). It indicates the density of genotypes

within each of the four categories through a color gradient from blue (low density) to yellow

(high density). **e. Visualization of CRP landscape based on relative connectivity.**

Analogous to panel c, but for relative connectivity. Color-codes indicate relative connectivity

(see color legend). **f. Spatial distribution of relative connectivity.** Like panel d, but for four

categories of relative connectivity (no data, low, medium, high).

Supplementary Figure S14. Quantitative analysis of the Fis landscape sparsity. a. Distribution of relative connectivity among genotypes. Violin plots augmented with

embedded boxplots illustrate the distribution of relative connectivity, i.e., the ratio of the empirically observed number of adjacent genotypes to the theoretical maximum number of 24 (=8×3) within a fully connected network. Genotypes are stratified into the three categories "other" (non-peak genotypes), "peaks" (excluding high peaks), and "high peaks." Mean relative connectivities and standard deviations are shown above each categorical division, revealing lower relative connectivity in peaks relative to non-peak genotypes (Two-sample Welch's t-test, p-value = 4.59×10^{-77}, N1 = 40,910; N2 = 2,312). High peaks exhibit comparable or superior connectivity relative to non-peak genotypes (Two-sample Welch's t-test, p-value = 9.31×10^{-13}, N1 = 40,910; N2 = 172). **b. Weak association between connectivity and regulation strength.** Scatterplot of relative connectivity against regulation strength. The density of genotypes (circles) in the scatterplot area is represented by a color gradient from purple (low density) to yellow (high density). The weak association (Pearson correlation, R = -0.13; t = -23.888; degrees of freedom (df) = 31,973; p-value $< 2.2 \times 10^{-16}$) indicates that strongly regulating genotypes are not necessarily densely connected. Marginal histograms adjacent to the scatterplot quantify the distributions of regulation strength and relative connectivity. **c. Visualization of Fis landscape based on regulation strength.** Visual overview of the Fis landscape with genotypes color-coded according to regulation strength (low strength: purple; medium: green; high: yellow; grey: genotypes with missing data). To enhance clarity, edges between genotypes are not shown. The landscape is projected onto a 2D space through a force-directed algorithm [182], with axes representing arbitrary units. **d. Spatial distribution of regulation strengths.** Each panel shows a density contour plot of the distribution of regulation strength scores of genotypes within one of four regulation strength categories (no data [missing data], low, medium, high). The density of genotypes within each category is indicated by a color gradient from blue (low density) to yellow (high density). **e. Visualization of Fis landscape based on relative connectivity.** Analogous to

1753   panel c, but for relative connectivity. Color codes indicate relative connectivity (see color

1754   legend). **f. Spatial distribution of relative connectivity.** Similar to panel d, but for four

1755   categories of relative connectivity (no data, low, medium, high).

Supplementary Figure S15. Quantitative analysis of the IHF landscape sparsity. a. Distribution of relative connectivity among genotypes. Violin plots augmented with

embedded boxplots illustrate the distribution of relative connectivity, i.e., the ratio of the empirically observed number of adjacent genotypes to the theoretical maximum number of 24 (=8×3) within a fully connected network. Genotypes are stratified into the three categories "other" (non-peak genotypes), "peaks" (excluding high peaks), and "high peaks." Mean relative connectivities and standard deviations are shown above each categorical division, revealing lower relative connectivity in peaks relative to non-peak genotypes (Two-sample Welch's t-test, p-value = 1.14×10^{-77}, N1 = 38,872; N2 = 2,254). High peaks exhibit comparable or superior connectivity relative to non-peak genotypes (Two-sample Welch's t-test, p-value = 9.31e-13, N1 = 38,872; N2 = 199). **b. Weak association between connectivity and regulation strength.** Scatterplot of relative connectivity against regulation strength. The density of genotypes (circles) in the scatterplot area is represented by a color gradient from purple (low density) to yellow (high density). The slight positive correlation (Pearson correlation, R = 0.017; t = 3.51; degrees of freedom (df) = 41,323; p-value = 0.0004485) indicates that genotypes with elevated regulation strength are only marginally more likely to be densely connected. Marginal histograms adjacent to the scatterplot quantify the distributions of regulation strength and relative connectivity. **c. Visualization of IHF landscape based on regulation strength.** Visual overview of the IHF landscape with genotypes color-coded according to regulation strength (low strength: purple; medium: green; high: yellow; grey: genotypes with missing data). To enhance clarity, edges between genotypes are not shown. The landscape is projected onto a 2D space through a force-directed algorithm [182], with axes representing arbitrary units. **d. Spatial distribution of regulation strengths.** Each panel shows a density contour plot of the distribution of regulation strength scores of genotypes within one of four regulation strength categories (no data [missing data], low, medium, high). The density of genotypes within each category is indicated by a color gradient from blue (low density) to yellow (high density). **e. Visualization of IHF landscape**

1784   **based on relative connectivity.** Analogous to panel c, but for relative connectivity. Color

1785   codes indicate relative connectivity (see color legend). **f. Spatial distribution of relative**

1786   **connectivity.** Similar to panel d, but for four categories of relative connectivity (no data, low,

1787   medium, high).

1788

Supplementary Figure S16. Peaks are spread across each landscape. The vertical axis

shows the genetic distance d between peaks and an equal number of randomly chosen non-

peak genotypes (color legend) for the CRP (N= 2,154), Fis (N= 2,312), and IHF (N= 2,434)

landscape (horizontal axis). (CRP: d_{peak}= 6, d_{random}= 5.94, t-statistic= 52.49, p-value <0.01;

Fis: d_{peak} = 6, d_{random} = 5.97, t-statistic= 30.96, p-value <0.01; IHF: d_{peak} = 6, d_{random} = 5.97, t-

statistic= 34.07, p-value <0.01; null-hypothesis: means do not differ). Although the

differences are statistically significant, peaks are almost as dispersed within genotype space

as non-peaks. Each box covers the range between the first and third quartiles (IQR). The

horizontal line within the box represents the median value, and whiskers span 1.5 times the

IQR.

1799

1800

Supplementary Figure S17. Genetic diversity among peaks across TF landscapes. a.

CRP landscape. Comparative analysis of genetic distances among nucleotide sequences

from 2,154 peaks (red) versus 2,154 non-peak, random variants (grey). Mean genetic

distances are displayed with vertical lines, showing a slight difference between peaks (d=5.6,

in red) and non-peaks (d=5.5, in grey), demonstrating large genetic diversity within the peaks

(Welch Two Sample t-test, t = 78.299, df = 4637305, p-value < 2.2×10^{-16}). **b. Fis**

landscape. Comparative analysis of genetic distances among nucleotide sequences from

2,312 peaks (red) versus 2,312 non-peak, random variants (grey). Mean genetic distances are

displayed with vertical lines, showing no significant difference between peaks (d=5.7, in red)

and non-peaks (d=5.7, in grey), demonstrating large genetic diversity within the peaks (t =

27.234, df = 5412460, p-value < 2.2×10^{-16}). **c. IHF landscape.** Comparative analysis of

genetic distances among nucleotide sequences from 2,434 peaks (red) versus 2,434 non-peak,

random variants (grey). Mean genetic distances are displayed with vertical lines, showing no

significant difference between peaks (d=6, in red) and non-peaks (d=6, in grey),

demonstrating large genetic diversity within the peaks (t = 54.689, df = 6014743, p-value <

2.2×10^{-16}).

1817

1818

1819

Supplementary Figure S18. Genetic diversity among high peaks across TF landscapes.

a. CRP landscape. The overlayed histograms compare the genetic diversity between 61 high

regulation strength peaks (red) and 61 non-peak, randomly selected variants (grey). No

significant difference exists (W) between high-regulation strength peaks (mean d=5.4, in red)

compared to randomly selected genotypes (mean d=5.5, in grey; Welch Two Sample t-test , t

= -0.78555, df = 3653.7, p-value = 0.4322). **c. Fis landscape.** The overlayed histograms

compare the genetic diversity between 172 high-regulation strength peaks (red) and 172 non-

peak, randomly selected genotypes (grey). No significant difference exists between high-

regulation strength peaks (mean d=5.4, in red) and random genotypes (mean d=5.8, in grey;

Welch Two Sample t-test, t = -16.634, df = 29357, p-value < 2.2×10^{-16}). **c. IHF landscape.**

The overlayed histograms compare the genetic diversity between 199 high regulation strength

peaks (red) and 199 non-peak, randomly selected genotypes (grey). The analysis reveals no

significant difference between high regulation strength peaks (mean d=5.8, in red) compared

to the random variants (mean d=5.8, in grey; Welch Two Sample t-test, t = 0.73365, df =

39366, p-value = 0.4632).

1835

1836

Supplementary Figure S19. Principal component analysis of the CRP landscape. a. PCA plot. A principal component analysis reveals that adaptive peaks are distributed widely within the genotype space (Supplementary Methods). The panels show principal components (PC) 1 and 2. Axes labels also indicate the percentage of variation explained by each component. Every circle corresponds to one among the 31,975 genotypes within the landscape. Light blue

and light red circles represent low and high peaks, respectively, while grey circles represent

non-peak variants. The clustered structure in the genotype space is due to the different

influence of individual nucleotides on the variation observed in the data (see panel **b**). **b. The**

contribution of individual nucleotides to principal component analysis. A PCA

contribution plot is a way to visualize the relative importance of different variables to the

variation observed in the data. In this plot, the contribution of each variable (nucleotide

identity at each position of a TFBS) is expressed as the square of the cosine of the angle

($\cos^2$) between the variable's vector (column representing the presence/absence of a base

letter at each position in the sequence) and each principal component axis. This quantity is

represented as an arrow that indicates the correlation of the variable with PC1 and PC2, the

two principal components that capture the largest amount of variation in the data. Both length

and color of the arrow represent the contribution of the variable to the variation observed in

the data. A high $\cos^2$ value (red) indicates that the variable is strongly correlated with the

principal component, and therefore makes a large contribution to the variation observed in the

data. A low $\cos^2$ value (blue) indicates that the variable is weakly correlated with the principal

component, and therefore makes a small contribution to the variation observed in the data.

Each arrow's length represents the importance of the variable's contribution relative to other

variables in the plot. Each nucleotide is represented with a base letter (A, T, C, G) followed

by a number that indicates the position of that base in the binding site sequence (e.g. G18

stands for a guanine at position 18 of the binding site).

1863

Supplementary Figure S20. Principal component analysis of the Fis landscape a. PCA plot. A principal component analysis reveals that adaptive peaks are distributed widely within the genotype space (Supplementary Methods). The panels show principal components (PC) 1 and 2. Axes labels also indicate the percentage of variation explained by each component. Every circle corresponds to one among the 43,222 genotypes within the landscape. Light blue and light red circles represent low and high peaks, respectively, while grey circles represent

non-peak variants. The clustered structure of the data in genotype space is due to the different

influence of individual nucleotides on the variation observed in the data (see panel b). **b. The**

contribution of individual nucleotides to principal component analysis. A PCA

contribution plot visualizes the relative importance of different variables to the variation

observed in the data. In this plot, the contribution of each variable (nucleotide identity at each

position of a TFBS) is expressed as the square of the cosine of the angle ($\cos^2$) between the

variable's vector (column representing the presence/absence of a base letter at each position

in the sequence) and each principal component axis. This quantity is represented as an arrow

that indicates the correlation of the variable with PC1 and PC2, the two principal components

that capture the largest amount of variation in the data. Both the length and color of the arrow

represent the contribution of the variable to the variation observed in the data. A high $\cos^2$

value (red) indicates that the variable is strongly correlated with the principal component and

therefore makes a large contribution to the variation observed in the data. A low $\cos^2$ value

(blue) indicates that the variable is weakly correlated with the principal component and

therefore makes a small contribution to the variation observed in the data. Each arrow's

length represents the importance of the variable's contribution relative to other variables in

the plot. Each nucleotide is represented with a base letter (A, T, C, G) followed by a number

that indicates the position of that base in the binding site sequence (e.g., G18 stands for a

guanine at position 18 of the binding site).

Supplementary Figure S21. Principal Component Analysis of the IHF Landscape

a. PCA Plot. A principal component analysis reveals that adaptive peaks are distributed widely within the genotype space (Supplementary Methods). The panels show principal components (PC) 1 and 2. Axes labels also indicate the percentage of variation explained by each component. Every circle corresponds to one among the 41,325 genotypes within the landscape. Light blue and light red circles represent low and high peaks, respectively, while grey circles represent non-peak variants. The clustered structure in the genotype space is due to the different influence of individual nucleotides on the variation observed in the data (see

panel b). **b. The contribution of individual nucleotides to principal component analysis**.

A PCA contribution plot is a way to visualize the relative importance of different variables to

the variation observed in the data. In this plot, the contribution of each variable (nucleotide

identity at each position of a TFBS) is expressed as the square of the cosine of the angle

($\cos^2$) between the variable's vector (column representing the presence/absence of a base

letter at each position in the sequence) and each principal component axis. This quantity is

represented as an arrow that indicates the correlation of the variable with PC1 and PC2, the

two principal components that capture the largest amount of variation in the data. Both the

length and color of the arrow represent the contribution of the variable to the variation

observed in the data. A high $\cos^2$ value (red) indicates that the variable is strongly correlated

with the principal component and therefore makes a large contribution to the variation

observed in the data. A low $\cos^2$ value (blue) indicates that the variable is weakly correlated

with the principal component and therefore makes a small contribution to the variation

observed in the data. Each arrow's length represents the importance of the variable's

contribution relative to other variables in the plot. Each nucleotide is represented with a base

letter (A, T, C, G) followed by a number that indicates the position of that base in the binding

site sequence (e.g., G18 stands for a guanine at position 18 of the binding site).

1916

1917

Supplementary Figure S22. Accessible paths to high peaks are less than two mutational steps longer than the shortest genetic distances to the peak. Each composite histogram shows the distribution of genetic distances (blue) and shortest accessible path length (green) for all pairs of high peaks and non-peak genotypes for one of the three landscapes. We used a t-test of the null hypothesis that the means of these distance distributions are statistically indistinguishable. This null hypothesis is rejected for all three landscapes. **a. CRP Landscape.** Mean genetic distance: 5.74; mean number of mutational steps in shortest accessible paths: 7.2; t = 500.17, df = 1,212,712, p < 2.2×10^{-16}. **b. Fis Landscape.** Mean genetic distance: 5.98; mean number of mutational steps in shortest accessible paths: 7.7; t = -1271, df = 5,639,001, p < 2.2×10^{-16}. **c. IHF Landscape.** Mean genetic distance: 5.78; mean number of mutational steps in shortest accessible paths: 7.6; t = -1365, df = 6,636,001, p < 2.2×10^{-16}.

1930

1931

1932 **Supplementary Figure S23. Moderate positive correlation between peak regulation**

1933 **strength and basin size for the three regulatory landscapes. a. CRP landscape.** The

1934 scatter plot shows the association between regulation strength conveyed by a peak variant

1935 (horizontal axis, colors from dark blue [low] to yellow [high]) and the size of its basin of

1936 attraction (absolute numbers on the left vertical axis, percentages on the right vertical axis,

1937 note the logarithmic scales). The dashed vertical line represents the regulation strength of the

1938 wild type. The black curve represents a linear regression line (in linear space). R is the

1939 Pearson correlation coefficient (R=0.39) and R^2 is the coefficient of determination for the

1940 linear regression model (R^2 =0.18, N = 2,154) **b. Fis landscape.** Like panel *a* (R=0.37, R^2

1941 =0.15, N = 2,312), **c. IHF landscape.** Like panel *a* (R=0.41, R^2 =0.19, N = 2,434).

1942

Supplementary Figure S24. The basins of attractions of high peaks overlap. Each heatmap matrix quantifies the proportion of shared TFBS variants in the basins of attractions of all pairs of high peaks (Methods). Each matrix entry quantifies the basin overlap for one pair of high peaks using the Jaccard index (**Supplementary Methods 7.5**), where row and column indices correspond to specific pairs of peaks. The maximum overlap of one (red) signifies that two basins share all variants, while the minimum value of zero (blue) indicates that they share no variants. **a. Basin overlap in the CRP landscape.** The average Jaccard index is 0.461 with a standard deviation of ±0.144, based on 3,721 pairwise comparisons. **b. Basin overlap in the Fis landscape**. The average Jaccard index is 0.426 with a standard deviation of ±0.136, derived from 29,584 pairwise comparisons. **c. Basin overlap in the IHF landscape.** The average Jaccard index is 0.364, with a standard deviation of ±0.145, based on 39,601 pairwise comparisons.

1957

1958 **Supplementary Figure S25. Distribution of the fraction of accessible high peaks per**

1959 **TFBS variant across TF landscapes.** The figure shows violin plots with embedded

1960 boxplots, for the distribution of high peak accessibility of each variant within the CRP

1961 (green), Fis (orange), and IHF (purple) landscape. The y-axis quantifies peak accessibility as

1962 the fraction of the total number of high peaks that can be accessed from each (non-peak)

1963 genotype within each TF landscape, revealing the extent to which individual variants can

1964 potentially access different high peaks. In the CRP landscape, each genotype can access, on

1965 average, a proportion of 0.314±0.315 (mean±s.d.) high peaks from the total number of high

1966 peaks, with a median value of 0.194. Notably, 5,152 out of 29,821 genotypes (17%) can

1967 access no high peak. Fis: mean high peak accessibility: 0.363±0.311 (median: 0.305), with

1968 4,304 out of 40,895 (10%) genotypes lacking access to any high peaks. IHF: mean peak

1969 accessibility: 0.301±0.289 (median: 0.205); 4,629 out of 38,872 genotypes (12%) show lack

1970 access to high peaks.

1971

1972

Supplementary Figure S26. Kimura adaptive walks on the CRP landscape a-c.
Fraction of adaptive walks attaining TFBS variants with strong regulation for different
population sizes. The vertical axis indicates the cumulative fraction of adaptive walks on the
CRP landscape that end at a genotype with a regulation strength shown on the horizontal
axes. Data is based on 15 million simulated adaptive walks (15,000 random starting
genotypes × 1,000 adaptive walks each). The vertical dashed line indicates the regulation
strength of the wild-type TFBS for CRP, and can be used to infer the proportion of walks that
terminate at lower regulatory peaks (Horizontal line, blue lettering). For example, a value of
y=0.82 implies that 82% of walks terminate at peaks lower than the wild-type. The remaining
18% reach higher peaks. **a. Population size 10^2.** 18% of walks terminate at peaks higher than
the wild-type. **b. Population size 10^5.** 10% of the walks terminate at higher peaks. **c.**
Population size 10^8. 10% of walks terminate at higher peaks. **d-f. Number of high peaks**
reached by a single adaptive walks. The histograms show the distribution of the number of
high peaks visited by a single adaptive walk, based on the same $15 × 10^6$ adaptive walks
analyzed in a-c. Note that a walk can visit multiple peaks when genetic drift is strong,

1988   because genetic drift can displace a population from an adaptive peak. The mean ± s.d. of the

1989   distribution is shown at the top of each plot. Population sizes: **d** 10^2, **e** 10^5, and **f** 10^8.

Supplementary Figure S27. Kimura adaptive walks on the Fis landscape. a-c. Fraction of adaptive walks attaining high peaks for different population sizes. The vertical axis indicates the cumulative percentage of adaptive walks on the Fis landscape that end at a genotype with a regulation strength shown on the horizontal axis. Data is based on 15 million simulated adaptive walks (15,000 starting genotypes × 1,000 adaptive walks each). The vertical dashed line indicates the regulation strength of the wild-type TFBS for Fis, and can be used to infer the proportion of walks that terminate at lower regulatory peaks (horizontal line, blue lettering). For example, a y-value of 0.9 implies that 90% of walks terminate at peaks lower than the wild-type, with the remaining 10% reaching higher peaks. **a. Population size 10^2.** 27% of walks terminate at peaks higher than the wild-type. **b. Population size 10^5.** 13% of walks terminate at higher peaks. **c. Population size 10^8.** 13% of walks terminate at higher peaks. **d-f. Number of high peaks reached by individual adaptive walks.** The histograms show the distribution of the number of high peaks visited by a single adaptive walk, based on the same 150 million adaptive walks analyzed in a-c. Note that a walk can visit multiple peaks when genetic drift is strong because genetic drift can

2006   displace a population from an adaptive peak. The mean $\pm$ s.d. of the distribution is shown at

2007   the top of each plot. Population sizes: **d** 10^2, **e** 10^5, and **f** 10^8.

2008

Supplementary Figure S28. Kimura adaptive walks on the IHF landscape. a-c. Fraction of adaptive walks attaining high peaks for different population sizes. The vertical axis indicates the cumulative percentage of adaptive walks on the IHF landscape that end at a genotype with a regulation strength shown on the horizontal axis. Data is based on 15 million simulated adaptive walks (15,000 starting points × 1,000 adaptive walks each). The vertical dashed line indicates the regulation strength of the wild-type TFBS for IHF, and can be used to infer the proportion of walks that terminate at lower regulatory peaks (horizontal line, blue lettering). For example, a y-value of 0.9 implies that 90% of walks terminate at peaks lower than the wild-type, with the remaining 10% reaching higher peaks. **a. Population size 10^2.** 31% of walks terminate at peaks higher than the wild-type. **b. Population size 10^5.** 15% of walks terminate at higher peaks. **c. Population size 10^8.** 15% of walks terminate at higher peaks. **d-f. Number of high peaks reached by individual adaptive walks.** The histograms show the distribution of the number of high peaks visited by a single adaptive walk, based on the same 150 million adaptive walks analyzed in a-c. Note that a walk can visit multiple peaks when genetic drift is strong because genetic drift can displace a population from an

2024 adaptive peak. The mean $\pm$ s.d. of the distribution is shown at the top of each plot. Population

2025 sizes: **d** 10^2, **e** 10^5, and **f** 10^8.

2026

2027

2028 **Supplementary Figure S29. Fraction of greedy adaptive walks attaining high peaks for**

2029 **different TFs. a-c. Fraction of greedy adaptive walks attaining high peaks for different**

2030 **TFs.** The vertical axis indicates the cumulative distribution function (CDF) of adaptive walks

2031 on each TF landscape that end at a genotype with a regulation strength shown on the

2032 horizontal axis. Data is based on 15,000 adaptive walks (15,000 starting points × 1 adaptive

2033 greedy walk for each) for each TF landscape. The vertical dashed line indicates the regulation

2034 strength of the wild-type TFBS for their respective TFs and can be used to infer the

2035 proportion of walks that terminate at lower regulatory peaks (horizontal line, blue lettering).

2036 For example, a value of y=0.9 implies that 90% of walks terminate at peaks lower than the

2037 wild-type, with the remaining 10% terminating at higher peaks. **a. CDF for adaptive walks**

2038 **in the CRP landscape.** Note the intersection at y= 0.89, meaning that 11% of the walks

2039 terminated on high peaks. **b. CDF for adaptive walks in the Fis landscape.** Note the

2040 intersection at y= 0.85, meaning that 15% of the walks terminated on high peaks. **c. CDF for**

2041 **adaptive walks in the IHF landscape.** Note the intersection at y= 0.82, meaning that 18% of

2042 the walks terminated on high peaks.

2043

Supplementary Figure S30. Diversity of alternative shortest paths for pairs of starting genotypes and their respective attained peaks. Each violin plot is based on data from 15,000 randomly (and uniformly) sampled starting (non-peak) genotypes from each TF landscape – CRP, Fis and IHF (see color legend) – and on 1'000 Kimura adaptive walks per starting genotypes. For each pair of a starting genotype and a high peak that is attainable from this starting genotype, we calculated the number of distinct shortest paths reaching the peak (vertical axis, note logarithmic scale). The width of the violin plot at any given y-axis value represents the density of data points at that value. Wider sections indicate a higher density of data points. The text at the top of each violin indicates the median number of shortest paths and the interquartile range (IQR) for each TF. For example, in the CRP landscape peaks are reached via a median number (± IQR) of 4 (± 4) shortest paths, with a sample size (number of starting genotypes × attainable peak pairs) of N = 345,224. The violin regions appear discretized because the number of shortest paths is a discrete variable, not a continuous one. This means that the data can only take on specific integer values, resulting in discrete regions within the violin plots with a non-zero density of data points.

2060

Supplementary Figure S31. Peak genotypes conveying stronger regulation are also reached more often during adaptive evolution. Each scatter plot represents the association between the strength of regulation conveyed by a peak genotype (horizontal axis) and the number of times the peak is attained (vertical axis, note the logarithmic scale). The data for each plot is based on 15,000 randomly (and uniformly) sampled starting (non-peak) genotypes from each TF landscape – CRP, Fis and IHF (see color legend) – and on 1'000 Kimura adaptive walks per starting genotypes (15×10^6 adaptive walks per landscape). The heatmap colors represent the density of peak genotypes from low (dark blue) to high (yellow). The dashed vertical line represents the regulation strength of the wild type TFBS. The red curve is derived from linear regression (in linear space). P-values in each panel result from a test of the null hypothesis that the two variables are not associated (Pearson correlation coefficient R=0; Coefficient of determination R^2=0). **a**. **CRP landscape** (R= 0.56; R^2 =0.34, N = 2,154). **b. Fis landscape.** (R=0.6, R^2 =0.36, N = 2,312), **c. IHF landscape.** (R=0.62, R^2 =0.4, N = 2,434).

Supplementary Figures S32. The frequency distribution of attained high peaks is biased towards some frequently attained peaks. We randomly and uniformly sampled 10 starting genotypes, started 10^3 adaptive walks from each, and recorded the number and frequency of distinct high peaks attained in these random walks. Each starting variant is symbolized by a vertical bar. The number of stacks within each bar (delineated by horizontal lines, also indicated by an integer above each bar) shows the number of high peaks reached by the 10^3 adaptive walks. Starting variants are ordered in ascending order based on this number of attained peaks. Stack height indicates the fraction of walks that reached the same peak, and is indicated in red, orange, and yellow for the three most frequently attained peaks. **a. Fis landscape. b. IHF landscape.**

2087

Supplementary Figure S33. The frequency distribution of traversed paths to high peaks is biased towards some paths that are frequently traversed. We randomly and uniformly sampled 20 starting genotypes, initiated 10^3 adaptive walks from each, and recorded the number and frequency of distinct paths traversed by these random walks for each attained peak. We then randomly and uniformly sampled a high peak attained by each starting variant for our analysis. Each starting variant is represented by a vertical bar. The number of stacks within each bar indicates the number of unique paths traversed to reach the sampled high peak. Stack height represents the fraction of walks that reached the high peak, with the three most frequently transversed paths in red, orange, and yellow. **a. CRP landscape. b. Fis landscape. c. IHF landscape.**

2098

REFERENCES:

1. Browning, D. F. & Busby, S. J. W. Local and global regulation of transcription initiation in bacteria. *Nat Rev Microbiol* **14**, 638–650 (2016).

2. Cases, I., De Lorenzo, V. & Ouzounis, C. A. Transcription regulation and environmental adaptation in bacteria. *Trends in Microbiology* vol. 11 248–253 Preprint at https://doi.org/10.1016/S0966-842X(03)00103-3 (2003).

3. Seshasayee, A. S., Bertone, P., Fraser, G. M. & Luscombe, N. M. Transcriptional regulatory networks in bacteria: from input signals to output responses. *Curr Opin Microbiol* **9**, 511–519 (2006).

4. Barnard, A., Wolfe, A. & Busby, S. Regulation at complex bacterial promoters: How bacteria use different promoter organizations to produce different regulatory outcomes. *Curr Opin Microbiol* **7**, 102–108 (2004).

5. Browning, D. F. D. F. & Busby, S. J. W. S. The regulation of bacterial transcription initiation. *Nat Rev Microbiol* **2**, 57–65 (2004).

6. Aguilar-Rodríguez, J., Payne, J. L. & Wagner, A. A thousand empirical adaptive landscapes and their navigability. *Nat Ecol Evol* **1**, 0045 (2017).

7. Sharon, E. *et al.* Inferring gene regulatory logic from high-throughput measurements of thousands of systematically designed promoters. *Nat Biotechnol* **30**, 521–530 (2012).

8. Shultzaberger, R. K., Malashock, D. S., Kirsch, J. F. & Eisen, M. B. The Fitness Landscapes of cis-Acting Binding Sites in Different Promoter and Environmental Contexts. *PLoS Genet* **6**, e1001042 (2010).

9. Haldane, A., Manhart, M. & Morozov, A. V. Biophysical Fitness Landscapes for Transcription Factor Binding Sites. *PLoS Comput Biol* **10**, 36–38 (2014).

10. Shen-Orr, S. S., Milo, R., Mangan, S. & Alon, U. Network motifs in the transcriptional regulation network of Escherichia coli. *Nat Genet* **31**, 64–68 (2002).

11. Martínez-Antonio, A. *et al.* Identifying global regulators in transcriptional regulatory networks in bacteria. *Curr Opin Microbiol* **6**, 482–489 (2003).

12. Lozada-Chávez, I., Angarica, V. E., Collado-Vides, J. & Contreras-Moreira, B. The Role of DNA-binding Specificity in the Evolution of Bacterial Regulatory Networks. *J Mol Biol* **379**, 627–643 (2008).

13. Kurafeiski, J. D., Pinto, P. & Bornberg-Bauer, E. Evolutionary potential of cis-regulatory mutations to cause rapid changes in transcription factor binding. *Genome Biol Evol* **11**, 406–414 (2019).

14. Browning, D. F., Butala, M. & Busby, S. J. W. Bacterial Transcription Factors: Regulation by Pick "N" Mix. *J Mol Biol* **431**, 4067–4077 (2019).

15. Visweswariah, S. S. & Busby, S. J. W. Evolution of bacterial transcription factors: How proteins take on new tasks, but do not always stop doing the old ones. *Trends Microbiol* **23**, 463–467 (2015).

16. Wray, G. A. The evolutionary significance of cis-regulatory mutations. *Nat Rev Genet* **8**, 206–216 (2007).

17. Signor, S. A. & Nuzhdin, S. V. The Evolution of Gene Expression in cis and trans. *Trends in Genetics* **34**, 532–544 (2018).

18. Wittkopp, P. J., Haerum, B. K. & Clark, A. G. Evolutionary changes in cis and trans gene regulation. *Nature* **430**, 85–88 (2004).

19. Yona, A. H., Alm, E. J. & Gore, J. Random sequences rapidly evolve into de novo promoters. *Nat Commun* **9**, 111880 (2018).

20. Fuqua, T. & Wagner, A. Mobile DNA is replete with hotspots for the de novo emergence of gene regulation. *bioRxiv* 2023.10.22.563463 (2023) doi:10.1101/2023.10.22.563463.

21. McAdams, H. H., Srinivasan, B. & Arkin, A. P. The evolution of genetic regulatory systems in bacteria. *Nature Reviews Genetics* vol. 5 169–178 Preprint at https://doi.org/10.1038/nrg1292 (2004).

22. Dorman, C. J., Bhriain, N. N. & Dorman, M. J. The Evolution of Gene Regulatory Mechanisms in Bacteria. in 125–152 (2018). doi:10.1007/978-3-319-69078-0_6.

23. Kinney, J. B. & McCandlish, D. M. Massively Parallel Assays and Quantitative Sequence–Function Relationships. *Annu Rev Genomics Hum Genet* **20**, annurev-genom-083118-014845 (2019).

24. de Visser, J. A. G. M., Elena, S. F., Fragata, I. & Matuszewski, S. The utility of fitness landscapes and big data for predicting evolution. *Heredity 2018 121:5* **121**, 401–405 (2018).

25. Draghi, J. A. & Ogbunugafor, C. B. Exploring the expanse between theoretical questions and experimental approaches in the modern study of evolvability. *J Exp Zool B Mol Dev Evol* **340**, 8–17 (2023).

26. Louis, A. A. Contingency, convergence and hyper-astronomical numbers in biological evolution. *Studies in History and Philosophy of Science Part C: Studies in History and Philosophy of Biological and Biomedical Sciences* **58**, 107–116 (2016).

27. Westmann, C. A., Goldbach, L. & Wagner, A. The highly rugged yet navigable regulatory landscape of the bacterial transcription factor TetR. *bioRxiv* 2023.08.25.554764 (2023) doi:10.1101/2023.08.25.554764.

28. Kim, D. *et al.* Systems assessment of transcriptional regulation on central carbon metabolism by Cra and CRP. *Nucleic Acids Res* (2018) doi:10.1093/nar/gky069.

29. Borirak, O. *et al.* Time-series analysis of the transcriptome and proteome of Escherichia coli upon glucose repression. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics* **1854**, 1269–1279 (2015).

30. Pal, A., Iyer, M. S., Srinivasan, S., Seshasayee, A. S. N. & Venkatesh, K. V. Global pleiotropic effects in adaptively evolved Escherichia coli lacking CRP reveal molecular mechanisms that define the growth physiology. *Open Biol* **12**, (2022).

31. Khankal, R., Chin, J. W., Ghosh, D. & Cirino, P. C. Transcriptional effects of CRP* expression in Escherichia coli. *J Biol Eng* **3**, 1–14 (2009).

32. Nowak-Lovato, K. *et al.* Binding of Nucleoid-Associated Protein Fis to DNA Is Regulated by DNA Breathing Dynamics. *PLoS Comput Biol* **9**, (2013).

33. Kahramanoglou, C. *et al.* Direct and indirect effects of H-NS and Fis on global gene expression control in Escherichia coli. *Nucleic Acids Res* **39**, 2073–2091 (2011).

34. Gawade, P., Gunjal, G., Sharma, A. & Ghosh, P. Reconstruction of transcriptional regulatory networks of Fis and H-NS in Escherichia coli from genome-wide data analysis. *Genomics* **112**, 1264–1272 (2020).

35. Prieto, A. I. *et al.* Genomic analysis of DNA binding and gene regulation by homologous nucleoid-associated proteins IHF and HU in Escherichia coli K12. *Nucleic Acids Res* **40**, 3524–3537 (2012).

36. Arfin, S. M. *et al.* Global gene expression profiling in Escherichia coli K12. The effects of integration host factor. *J Biol Chem* **275**, 29672–29684 (2000).

37. Bank, C. Epistasis and Adaptation on Fitness Landscapes.
https://doi.org/10.1146/annurev-ecolsys-102320-112153 **53**, 457–479 (2022).
38. Fragata, I., Blanckaert, A., Dias Louro, M. A., Liberles, D. A. & Bank, C.
Evolution in the light of fitness landscape theory. *Trends Ecol Evol* **34**, 69–82
(2019).
39. Vaishnav, E. D. *et al.* The evolution, evolvability and engineering of gene
regulatory DNA. *Nature 2022 603:7901* **603**, 455–463 (2022).
40. Wright, S. Evolution in Mendelian Populations. *Genetics* **16**, 97 (1931).
41. Taylor, L. R. & Provine, W. B. Sewall Wright and Evolutionary Biology. *J Anim*
Ecol (2006) doi:10.2307/5082.
42. Wright, S. The roles of mutation, inbreeding, crossbreeding and selection in
evolution. *Proc of the 6th International Congress of Genetics* Preprint at
(1932).
43. Lynch, M. *et al.* Genetic drift, selection and the evolution of the mutation rate.
Nature Reviews Genetics 2016 17:11 **17**, 704–714 (2016).
44. Iwasa, Y., Michor, F. & Nowak, M. A. Stochastic tunnels in evolutionary
dynamics. *Genetics* **166**, 1571–1579 (2004).
45. Taylor, C. F. & Higgs, P. G. A population genetics model for multiple
quantitative traits exhibiting pleiotropy and epistasis. *J Theor Biol* **203**, 419–
437 (2000).
46. de Visser, J. A. G. M., Cooper, T. F. & Elena, S. F. The causes of epistasis.
Proceedings of the Royal Society B: Biological Sciences **278**, 3617–3624
(2011).
47. Santos-Zavaleta, A. *et al.* RegulonDB v 10.5: Tackling challenges to unify
classic and high throughput knowledge of gene regulation in E. coli K-12.
Nucleic Acids Res (2019) doi:10.1093/nar/gky1077.
48. Peterman, N. & Levine, E. Sort-seq under the hood: Implications of design
choices on large-scale characterization of sequence-function relations. *BMC*
Genomics **17**, 1–17 (2016).
49. de Boer, C. G. *et al.* Deciphering eukaryotic gene-regulatory logic with 100
million random promoters. *Nature Biotechnology 2019 38:1* **38**, 56–65 (2019).
50. Feng, H. *et al.* Deep-learning-assisted Sort-Seq enables high-throughput
profiling of gene expression characteristics with high precision. *Sci Adv* **9**,
eadg5296 (2023).
51. Garcia, H. G., Lee, H. J., Boedicker, J. Q. & Phillips, R. Comparison and
Calibration of Different Reporters for Quantitative Analysis of Gene
Expression. *Biophys J* **101**, 535 (2011).
52. Friedlander, T., Prizak, R., Barton, N. H. & Tkačik, G. Evolution of new
regulatory functions on biophysically realistic fitness landscapes. *Nat Commun*
8, (2017).
53. Le, D. D. *et al.* Comprehensive, high-resolution binding energy landscapes
reveal context dependencies of transcription factor binding. *Proceedings of the*
National Academy of Sciences **115**, 201715888 (2018).
54. Maerkl, S. J. S. J. & Quake, S. R. A systems approach to measuring the
binding energy landscapes of transcription factors. *Science (1979)* **315**,
(2007).
55. Poelwijk, F. J., Kiviet, D. J., Weinreich, D. M. & Tans, S. J. Empirical fitness
landscapes reveal accessible evolutionary paths. *Nature* (2007)
doi:10.1038/nature05451.

56. Nghe, P. *et al.* Predicting Evolution Using Regulatory Architecture. *Annu Rev Biophys* **49**, 181–197 (2020).

57. Struhl, K. Fundamentally different logic of gene regulation in eukaryotes and prokaryotes. *Cell* **98**, 1–4 (1999).

58. Bintu, L. *et al.* Transcriptional regulation by the numbers: models. *Curr Opin Genet Dev* **15**, 116–124 (2005).

59. Barnes, S. L., Belliveau, N. M., Ireland, W. T., Kinney, J. B. & Phillips, R. Mapping DNA sequence to transcription factor binding energy in vivo. *PLoS Comput Biol* **15**, e1006226 (2019).

60. Kolb, A., Spassky, A., Chapon, C., Blazy, B. & Buc, H. On the different binding affinities of CRP at the lac, gal and malt promoter regions. *Nucleic Acids Res* (1983) doi:10.1093/nar/11.22.7833.

61. Shao, Y., Feldman-Cohen, L. S. & Osuna, R. Functional Characterization of the Escherichia coli Fis-DNA Binding Sequence. *J Mol Biol* **376**, 771–785 (2008).

62. Huo, Y. X. *et al.* IHF-binding sites inhibit DNA loop formation and transcription initiation. *Nucleic Acids Res* **37**, 3878–3886 (2009).

63. Gunasekera, A., Ebright, Y. W. & Ebright, R. H. DNA sequence determinants for binding of the Escherichia coli catabolite gene activator protein. *Journal of Biological Chemistry* **267**, 14713–14720 (1992).

64. Aiyar, S. E. *et al.* Architecture of fis-activated transcription complexes at the Escherichia coli rrnB P1 and rrnE P1 promoters. *J Mol Biol* **316**, 501–516 (2002).

65. Ho, S. Characteristics of IHF Binding to Holliday Junction DNA. (2013).

66. Salgado, H. *et al.* RegulonDB v12.0: a comprehensive resource of transcriptional regulation in E. coli K-12. *Nucleic Acids Res* **52**, D255–D264 (2024).

67. D'Haeseleer, P. What are DNA sequence motifs? *Nature Biotechnology 2006 24:4* **24**, 423–425 (2006).

68. Stormo, G. D. DNA binding sites: Representation and discovery. *Bioinformatics* vol. 16 16–23 Preprint at https://doi.org/10.1093/bioinformatics/16.1.16 (2000).

69. Song, S. & Zhang, J. Unbiased inference of the fitness landscape ruggedness from imprecise fitness estimates. *Evolution (N Y)* **75**, 2658–2671 (2021).

70. Papkou, A., Garcia-Pastor, L., Escudero, J. A. & Wagner, A. A rugged yet easily navigable fitness landscape. *Science (1979)* **382**, (2023).

71. Obolski, U., Ram, Y. & Hadany, L. Key issues review: evolution on rugged adaptive landscapes. *Reports on Progress in Physics* **81**, 012602 (2018).

72. Kauffman, S. & Levin, S. Towards a general theory of adaptive walks on rugged landscapes. *J Theor Biol* **128**, 11–45 (1987).

73. Weinberger, E. Correlated and uncorrelated fitness landscapes and how to tell the difference. *Biol Cybern* **63**, 325–336 (1990).

74. Orr, H. A. A Minimum on the Mean Number of Steps Taken in Adaptive Walks. *J Theor Biol* **220**, 241–247 (2003).

75. Saona, R., Kondrashov, F. A. & Khudiakova, K. A. Relation Between the Number of Peaks and the Number of Reciprocal Sign Epistatic Interactions. *Bull Math Biol* **84**, (2022).

76. Poelwijk, F. J., Tănase-Nicola, S., Kiviet, D. J. & Tans, S. J. Reciprocal sign epistasis is a necessary condition for multi-peaked fitness landscapes. *J Theor Biol* **272**, 141–144 (2011).

77. Kvitek, D. J. & Sherlock, G. Reciprocal sign epistasis between frequently experimentally evolved adaptive mutations causes a rugged fitness landscape. *PLoS Genet* (2011) doi:10.1371/journal.pgen.1002056.

78. Conrad, M. The geometry of evolution. *Biosystems* **24**, 61–81 (1990).

79. Krug, J. & Oros, D. Evolutionary accessibility of random and structured fitness landscapes. (2023).

80. KIMURA, M. ON THE PROBABILITY OF FIXATION OF MUTANT GENES IN A POPULATION. *Genetics* **47**, 713–719 (1962).

81. Bank, C., Matuszewski, S., Hietpas, R. T. & Jensen, J. D. On the (un)predictability of a large intragenic fitness landscape. *Proc Natl Acad Sci U S A* **113**, 14085–14090 (2016).

82. Orr, H. A. The population genetics of adaptation: the adaptation of DNA sequences. *Evolution* **56**, 1317–1330 (2002).

83. Gillespie, J. H. MOLECULAR EVOLUTION OVER THE MUTATIONAL LANDSCAPE. *Evolution (N Y)* **38**, 1116–1129 (1984).

84. Lind, P. A. & Andersson, D. I. Whole-genome mutational biases in bacteria. *Proceedings of the National Academy of Sciences* **105**, 17878–17883 (2008).

85. Horton, J. S. & Taylor, T. B. Mutation bias and adaptation in bacteria. *Microbiology (N Y)* **169**, (2023).

86. Lee, H., Popodi, E., Tang, H. & Foster, P. L. Rate and molecular spectrum of spontaneous mutations in the bacterium *Escherichia coli* as determined by whole-genome sequencing. *Proceedings of the National Academy of Sciences* **109**, (2012).

87. Melissa, M. J., Good, B. H., Fisher, D. S. & Desai, M. M. Population genetics of polymorphism and divergence in rapidly evolving populations. *Genetics* **221**, (2022).

88. Li, J., Amado, A. & Bank, C. Rapid adaptation of recombining populations on tunable fitness landscapes. *Mol Ecol* (2023) doi:10.1111/MEC.16900.

89. Stolyarova, A. V. *et al.* Complex fitness landscape shapes variation in a hyperpolymorphic species. *Elife* **11**, (2022).

90. Park, S.-C., Neidhart, J. & Krug, J. Greedy adaptive walks on a correlated fitness landscape. *J Theor Biol* **397**, 89–102 (2016).

91. Kauffman, S. & Levin, S. Towards a general theory of adaptive walks on rugged landscapes. *J Theor Biol* **128**, 11–45 (1987).

92. ORR, H. A. A Minimum on the Mean Number of Steps Taken in Adaptive Walks. *J Theor Biol* **220**, 241–247 (2003).

93. Blount, Z. D., Lenski, R. E. & Losos, J. B. Contingency and determinism in evolution: Replaying life's tape. *Science (1979)* **362**, (2018).

94. Nonoyama, T. & Chiba, S. Phenotypic determinism and contingency in the evolution of hypothetical tree-like organisms. *PLoS One* **14**, (2019).

95. Xie, V. C., Pu, J., Metzger, B. P. H., Thornton, J. W. & Dickinson, B. C. Contingency and chance erase necessity in the experimental evolution of ancestral proteins. *Elife* **10**, 1–87 (2021).

96. Spor, A. *et al.* Phenotypic and genotypic convergences are influenced by historical contingency and environment in yeast. *Evolution* **68**, 772 (2014).

97. Lenski, R. E. Convergence and Divergence in a Long-Term Experiment with Bacteria*. *https://doi.org/10.1086/691209* **190**, S57–S68 (2017).

98. Lind, P. A., Farr, A. D. & Rainey, P. B. Experimental evolution reveals hidden diversityin evolutionary pathways. *Elife* **2015**, (2015).

99. Evfratov, S. A. *et al.* Application of sorting and next generation sequencing to study 5´-UTR influence on translation efficiency in Escherichia coli. *Nucleic Acids Res* **45**, 3487–3502 (2017).

100. Cuperus, J. T. *et al.* Deep learning of the regulatory grammar of yeast 5 ' untranslated regions from 500 , 000 random sequences. *Genome Res* **163**, 1–10 (2017).

101. Belliveau, N. M. *et al.* A Systematic and Scalable Approach for Dissecting the Molecular Mechanisms of Transcriptional Regulation in Bacteria. *Biophys J* **114**, 151a (2018).

102. Kinney, J. B., Murugan, A., Callan, C. G. & Cox, E. C. Using deep sequencing to characterize the biophysical mechanism of a transcriptional regulatory sequence. *Proceedings of the National Academy of Sciences* **107**, 9158–9163 (2010).

103. Lagator, M. *et al.* Predicting bacterial promoter function and evolution from random sequences. *Elife* **11**, (2022).

104. Urtecho, G., Tripp, A. D., Insigne, K. D., Kim, H. & Kosuri, S. Systematic Dissection of Sequence Elements Controlling σ70 Promoters Using a Genomically Encoded Multiplexed Reporter Assay in Escherichia coli. *Biochemistry* **58**, 1539–1551 (2019).

105. Aguilar-Rodríguez, J. & Payne, J. L. Robustness and Evolvability in Transcriptional Regulation. *Evolutionary Systems Biology* 197–219 (2021) doi:10.1007/978-3-030-71737-7_9.

106. Lagator, M., Igler, C., Moreno, A. B., Guet, C. C. & Bollback, J. P. Epistatic interactions in the arabinose cis-regulatory element. *Mol Biol Evol* **33**, 761–769 (2016).

107. Lagator, M., Paixão, T., Barton, N. H., Bollback, J. P. & Guet, C. C. On the mechanistic nature of epistasis in a canonical cis-regulatory element. *Elife* **6**, 1–16 (2017).

108. Stewart, A. J., Hannenhalli, S., Plotkin, J. B., Hannenhalli, S. & Plotkin, J. B. Why Transcription Factor Binding Sites Are Ten Nucleotides Long. *Genetics* **192**, 973–985 (2012).

109. Babu, M. M. & Teichmann, S. A. Functional determinants of transcription factors in Escherichia coli: Protein families and binding sites. *Trends in Genetics* **19**, 75–79 (2003).

110. Huffman, J. L. & Brennan, R. G. Prokaryotic transcription regulators: more than just the helix-turn-helix motif. *Curr Opin Struct Biol* **12**, 98–106 (2002).

111. Perez-Rueda, E. *et al.* Abundance, diversity and domain architecture variability in prokaryotic DNA-binding transcription factors. *PLoS One* **13**, (2018).

112. Oren, Y. *et al.* Transfer of noncoding DNA drives regulatory rewiring in bacteria. *Proceedings of the National Academy of Sciences* **111**, 16112–16117 (2014).

113. Price, M. N., Dehal, P. S. & Arkin, A. P. Horizontal gene transfer and the evolution of transcriptional regulation in Escherichia coli. *Genome Biol* **9**, (2008).

114. Weinreich, D. M., Watson, R. A. & Chao, L. Perspective: Sign epistasis and genetic constraint on evolutionary trajectories. *Evolution* (2005).

115. Stormo, G. D. & Zhao, Y. Determining the specificity of protein–DNA interactions. *Nat Rev Genet* **11**, 751–760 (2010).

116. Gerland, U., Moroz, J. D. & Hwa, T. Physical constraints and functional characteristics of transcription factor-DNA interaction. *Proceedings of the National Academy of Sciences* **99**, 12015–12020 (2002).

117. Zhao, Y., Ruan, S., Pandey, M. & Stormo, G. D. Improved models for transcription factor binding site identification using nonindependent interactions. *Genetics* (2012) doi:10.1534/genetics.112.138685.

118. O'Flanagan, R. A., Paillard, G., Lavery, R. & Sengupta, A. M. Non-additivity in protein–DNA binding. *Bioinformatics* **21**, 2254–2263 (2005).

119. Payne, J. L. & Wagner, A. The Robustness and Evolvability of Transcription Factor Binding Sites. *Science (1979)* **343**, (2014).

120. Blount, Z. D., Borland, C. Z. & Lenski, R. E. Historical contingency and the evolution of a key innovation in an experimental population of Escherichia coli. *Proc Natl Acad Sci U S A* **105**, 7899–7906 (2008).

121. Palmer, A. C. *et al.* Delayed commitment to evolutionary fate in antibiotic resistance fitness landscapes. *Nature Communications 2015 6:1* **6**, 1–8 (2015).

122. Vermeij, G. J. Historical contingency and the purported uniqueness of evolutionary innovations. *Proceedings of the National Academy of Sciences* **103**, 1804–1809 (2006).

123. Lässig, M., Mustonen, V. & Walczak, A. M. Predicting evolution. *Nat Ecol Evol* **1**, 1–9 (2017).

124. De Visser, J. A. G. M. & Krug, J. Empirical fitness landscapes and the predictability of evolution. *Nat Rev Genet* **15**, 480–490 (2014).

125. Heyde, S. A. H., Frendorf, P. O., Lauritsen, I. & Nørholm, M. H. H. Restoring Global Gene Regulation through Experimental Evolution Uncovers a NAP (Nucleoid-Associated Protein)-Like Behavior of Crp/Cap. *mBio* **12**, (2021).

126. Monteiro, L. M. O., Sanches-Medeiros, A., Westmann, C. A. & Silva-Rocha, R. Unraveling the Complex Interplay of Fis and IHF Through Synthetic Promoter Engineering. *Front Bioeng Biotechnol* **8**, 510 (2020).

127. Dillon, S. C. & Dorman, C. J. Bacterial nucleoid-associated proteins, nucleoid structure and gene expression. *Nat Rev Microbiol* **8**, 185–95 (2010).

128. Dorman, C. J. Genome architecture and global gene regulation in bacteria: Making progress towards a unified model? *Nat Rev Microbiol* **11**, 349–355 (2013).

129. Dorman, C. J., Schumacher, M. A., Bush, M. J., Brennan, R. G. & Buttner, M. J. When is a transcription factor a NAP? *Curr Opin Microbiol* **55**, 26 (2020).

130. Meier, I., Wray, L. V. & Hillen, W. Differential regulation of the Tn10-encoded tetracycline resistance genes tetA and tetR by the tandem tet operators O1 and O2. *EMBO J* **7**, 567–572 (1988).

131. Kauffman, S. A. & Weinberger, E. D. The NK model of rugged fitness landscapes and its application to maturation of the immune response. *J Theor Biol* **141**, 211–245 (1989).

132. Kingman, J. F. C. A simple model for the balance between selection and mutation. *J Appl Probab* **15**, 1–12 (1978).

133. Chen, Y. J. *et al.* Quantifying molecular bias in DNA data storage. *Nature Communications 2020 11:1* **11**, 1–9 (2020).

134. Kebschull, J. M. & Zador, A. M. Sources of PCR-induced distortions in high-throughput sequencing data sets. *Nucleic Acids Res* **43**, e143–e143 (2015).

135. Wong, T. S., Zhurina, D. & Schwaneberg, U. The Diversity Challenge in Directed Protein Evolution. *Comb Chem High Throughput Screen* **9**, 271–288 (2006).

136. Aird, D. *et al.* Analyzing and minimizing PCR amplification bias in Illumina sequencing libraries. *Genome Biol* **12**, 1–14 (2011).

137. Trippe, B. L. *et al.* Randomized gates eliminate bias in sort-seq assays. *Protein Science* **31**, e4401 (2022).

138. Gilliot, P.-A. & Gorochowski, T. E. Effective design and inference for cell sorting and sequencing based massively parallel reporter assays. *bioRxiv* 2022.11.07.515414 (2022) doi:10.1101/2022.11.07.515414.

139. Boer, C. de, Sadeh, R., Friedman, N. & Regev, A. Deciphering cis-regulatory logic with 100 million random promoters. *bioRxiv* 224907 (2018) doi:10.1101/224907.

140. De Boer, C. G., Ray, J. P., Hacohen, N. & Regev, A. MAUDE: Inferring expression changes in sorting-based CRISPR screens. *Genome Biol* **21**, 1–16 (2020).

141. Wu, N. C., Dai, L., Olson, C. A., Lloyd-Smith, J. O. & Sun, R. Adaptation in protein fitness landscapes is facilitated by indirect paths. *Elife* **5**, (2016).

142. Diaz-Colunga, J. *et al.* Global epistasis on fitness landscapes. *Philosophical Transactions of the Royal Society B* **378**, (2023).

143. MacLean, R. C., Perron, G. G. & Gardner, A. Diminishing Returns From Beneficial Mutations and Pervasive Epistasis Shape the Fitness Landscape for Rifampicin Resistance in Pseudomonas aeruginosa. *Genetics* **186**, 1345–1354 (2010).

144. Rydenfelt, M., Garcia, H. G., Cox, R. S. & Phillips, R. The influence of promoter architectures and regulatory motifs on gene expression in Escherichia coli. *PLoS One* **9**, 1–31 (2014).

145. Ishihama, A. Prokaryotic genome regulation: multifactor promoters, multitarget regulators and hierarchic networks. *FEMS Microbiol Rev* **34**, 628–645 (2010).

146. Cases, I. & de Lorenzo, V. Promoters in the environment: Transcriptional regulation in its natural context. *Nature Reviews Microbiology* vol. 3 105–118 Preprint at https://doi.org/10.1038/nrmicro1084 (2005).

147. Shepherd, M. J., Reynolds, M., Pierce, A. P., Rice, A. M. & Taylor, T. B. Transcription factor expression levels and environmental signals constrain transcription factor innovation. *Microbiology (N Y)* **169**, (2023).

148. González Pérez, A., Espinosa Angarica, V., Collado-Vides, J. & Vasconcelos, A. T. R. From sequence to dynamics: The effects of transcription factor and polymerase concentration changes on activated and repressed promoters. *BMC Mol Biol* **10**, 92 (2009).

149. Azam, T. A., Iwata, A., Nishimura, A., Ueda, S. & Ishihama, A. Growth phase-dependent variation in protein composition of the Escherichia coli nucleoid. *J Bacteriol* (1999) doi:<p></p>.

150. Weinert, F. M. *et al.* The Transcription Factor Titration Effect Dictates Level of Gene Expression. *Biophys J* (2014) doi:10.1016/j.bpj.2013.11.4483.

151. Lagomarsino, M. C., Espéli, O. & Junier, I. From structure to function of bacterial chromosomes: Evolutionary perspectives and ideas for new experiments. *FEBS Letters* Preprint at https://doi.org/10.1016/j.febslet.2015.07.002 (2015).

152. Dorman, C. J. & Dorman, M. J. DNA supercoiling is a fundamental regulatory principle in the control of bacterial gene expression. *Biophys Rev* **8**, 209–220 (2016).

153. Houdaigui, B. El *et al.* Bacterial genome architecture shapes global transcriptional regulation by DNA supercoiling. *Nucleic Acids Res* **47**, 5648–5657 (2019).

154. Sobetzko, P., Travers, A. & Muskhelishvili, G. Gene order and chromosome dynamics coordinate spatiotemporal gene expression during the bacterial growth cycle. *Proceedings of the National Academy of Sciences* **109**, E42–E50 (2011).

155. Meyer, S., Reverchon, S., Nasser, W. & Muskhelishvili, G. Chromosomal organization of transcription: in a nutshell. *Curr Genet* **64**, 555–565 (2018).

156. Sánchez-Romero, M. A., Cota, I. & Casadesús, J. DNA methylation in bacteria: from the methyl group to the methylome. *Curr Opin Microbiol* **25**, 9–16 (2015).

157. Crocker, J., Preger-Ben Noon, E. & Stern, D. L. *The Soft Touch: Low-Affinity Transcription Factor Binding Sites in Development and Evolution. Current Topics in Developmental Biology* vol. 117 (Elsevier Inc., 2016).

158. Srivastava, M. & Payne, J. L. On the incongruence of genotype-phenotype and fitness landscapes. *PLoS Comput Biol* **18**, e1010524 (2022).

159. Rest, J. S. *et al.* Nonlinear fitness consequences of variation in expression level of a eukaryotic gene. *Mol Biol Evol* **30**, 448–456 (2013).

160. Duveau, F., Toubiana, W. & Wittkopp, P. J. Fitness effects of cis-regulatory variants in the saccharomyces cerevisiae TDH3 promoter. *Mol Biol Evol* **34**, (2017).

161. Bergen, A. C., Olsen, G. M. & Fay, J. C. Divergent MLS1 Promoters Lie on a Fitness Plateau for Gene Expression. *Mol Biol Evol* **33**, (2016).

162. Ornelas, M. Y., Cournoyer, J. E., Bram, S. & Mehta, A. P. Evolution and synthetic biology. *Curr Opin Microbiol* **76**, 102394 (2023).

163. Yi, X. & Dean, A. M. Adaptive Landscapes in the Age of Synthetic Biology. *Mol Biol Evol* **36**, 890–907 (2019).

164. Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: The Keio collection. *Mol Syst Biol* **2**, 2006.0008 (2006).

165. Elowitz, M. B., Levine, A. J., Siggia, E. D. & Swain, P. S. Stochastic Gene Expression in a Single Cell. *Science (1979)* **297**, 1183–6 (2007).

166. Sharon, E. *et al.* Probing the effect of promoters on noise in gene expression using thousands of designed sequences. *Genome Res* **24**, 1698–1706 (2014).

167. Rudge, T. J. *et al.* Characterization of Intrinsic Properties of Promoters. *ACS Synth Biol* **5**, 89–98 (2016).

168. Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* **6**, 343–5 (2009).

169. Warren, D. J. Preparation of highly efficient electrocompetent Escherichia coli using glycerol/mannitol density step centrifugation. *Anal Biochem* **413**, 206–207 (2011).

170. Jahn, M., Vorpahl, C., Hübschmann, T., Harms, H. & Müller, S. Copy number variability of expression plasmids determined by cell sorting and droplet digital PCR. *Microb Cell Fact* **15**, 211 (2016).

171. Lutz, R. & Bujard, H. Independent and tight regulation of transcriptional units in Escherichia coli via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. *Nucleic Acids Res* **25**, 1203–10 (1997).

172. Pédelacq, J. D., Cabantous, S., Tran, T., Terwilliger, T. C. & Waldo, G. S. Engineering and characterization of a superfolder green fluorescent protein. *Nat Biotechnol* **24**, 79–88 (2006).

173. Lou, C., Stanton, B., Chen, Y. J., Munsky, B. & Voigt, C. A. Ribozyme-based insulator parts buffer synthetic circuits from genetic context. *Nat Biotechnol* (2012) doi:10.1038/nbt.2401.

174. Bindels, D. S. *et al.* mScarlet: a bright monomeric red fluorescent protein for
cellular imaging. *Nature Methods 2016 14:1* **14**, 53–56 (2016).
175. Beal, J. Biochemical complexity drives log-normal variation in genetic
expression. *Engineering Biology* **1**, 55–60 (2017).
176. Beal, J. *et al.* Reproducibility of fluorescent expression from engineered
biological constructs in E. coli. *PLoS One* **11**, (2016).
177. Martin, M. Cutadapt removes adapter sequences from high-throughput
sequencing reads. *EMBnet J* **17**, 10–12 (2011).
178. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to
improve genome assemblies. *Bioinformatics* **27**, 2957–2963 (2011).
179. Hannon, G. J. FASTX-Toolkit.
180. Bunn, A. & Korpela, M. R: A language and environment for statistical
computing. (2013) doi:10.1016/j.dendro.2008.01.002.
181. Schneider, T. D. & Stephens, R. M. Sequence logos: a new way to display
consensus sequences. *Nucleic Acids Res* **18**, 6097 (1990).
182. Csardi, G. The Igraph Software Package for Complex Network Research.
(2014).
183. Khalid, F. *et al.* Genonets server-a web server for the construction, analysis
and visualization of genotype networks. *Nucleic Acids Res* **44**, W70–W76
(2016).
184. Jaccard, P. THE DISTRIBUTION OF THE FLORA IN THE ALPINE ZONE.1.
New Phytologist **11**, 37–50 (1912).
185. Papkou, A., Garcia-Pastor, L., Escudero, J. A. & Wagner, A. A rugged yet easily
navigable fitness landscape of antibiotic resistance. *bioRxiv*
2023.02.27.530293 (2023) doi:10.1101/2023.02.27.530293.
186. Kimura, M. *The Neutral Theory of Molecular Evolution.* (Cambridge University
Press, 1983). doi:10.1017/CBO9780511623486.
187. Crow, J. and Kimura, M. An Introduction to Population Genetics Theory
[Paperback]. 608 (2009).
188. Harris, C. R. *et al.* Array programming with NumPy. *Nature 2020 585:7825*
585, 357–362 (2020).
189. Grant, S. G., Jessee, J., Bloom, F. R. & Hanahan, D. Differential plasmid
rescue from transgenic mouse DNAs into Escherichia coli methylation-
restriction mutants. *Proceedings of the National Academy of Sciences* **87**,
4645–4649 (1990).
190. Ebright, R. H., Ebright, Y. W. & Gunasekera, A. Consensus DNA site for the
Escherichia coli catabolite gene activator protein (CAP): CAP exhibits a 450-
fold higher affinity for the consensus DNA site than for the E.coli lac DNA site.
Nucleic Acids Res **17**, 10295–10305 (1989).
191. Shao, Y., Feldman-Cohen, L. S. & Osuna, R. Biochemical Identification of
Base and Phosphate Contacts between Fis and a High-Affinity DNA Binding
Site. *J Mol Biol* **380**, 327–339 (2008).
192. Kahramanoglou, C. *et al.* Direct and indirect effects of H-NS and Fis on global
gene expression control in Escherichia coli. *Nucleic Acids Res* **39**, 2073–2091
(2011).
193. Kelly, J. R. *et al.* Measuring the activity of BioBrick promoters using an in vivo
reference standard. *J Biol Eng* **3**, 4 (2009).
194. Meyer, A. J., Segall-Shapiro, T. H., Glassey, E., Zhang, J. & Voigt, C. A.
Escherichia coli "Marionette" strains with 12 highly optimized small-molecule
sensors. *Nature Chemical Biology 2018 15:2* **15**, 196–204 (2018).

195. Carr, S. B., Beal, J. & Densmore, D. M. Reducing DNA context dependence in
bacterial promoters. *PLoS One* (2017) doi:10.1371/journal.pone.0176013.
196. Chen, Y. J. *et al.* Characterization of 582 natural and synthetic terminators and
quantification of their design constraints. *Nat Methods* (2013)
doi:10.1038/nmeth.2515.
197. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag
New York, 2016).
2599