Opposing Regulation of TNF Responses and IL-1β+ Macrophages by PGE2-cAMP and IFN-γ Signaling

Upneet K. Sokhi[1], Bikash Mishra[1,2]*, Ruoxi Yuan[1,3]*, Yuri Chinenov[1,3]*, Anvita Singaraju[1,2], Karmela K. Chan[4,5], Anne R. Bass[4,5], Laura Donlin[1,2,4,5] and Lionel B. Ivashkiv[1,2,4,5]

[1]HSS Research Institute and David Z. Rosensweig Genomics Research Center, Hospital for Special Surgery, New York, NY
[2]Immunology and Microbial Pathogenesis Program, Weill Cornell Medicine, New York, NY.
[3]Computational Biology Core, David Z. Rosensweig Genomics Research Center, Hospital for Special Surgery, New York, NY
[4]Division of Rheumatology, Department of Medicine, Hospital for Special Surgery, New York, NY
[5]Department of Medicine, Weill Cornell Medicine, New York, NY.

* co-second authors

Correspondence: Lionel B. Ivashkiv at ivashkivl@hss.edu

Abstract

IL-1β-expressing macrophages have been described in rheumatoid arthritis (RA), immune checkpoint inhibitor-induced inflammatory arthritis (ICI-arthritis), and pancreatic cancer and proposed to be pathogenic. In RA and pancreatic cancer IL-1β+ macrophages express a TNF+PGE2 (TP) gene expression signature induced by cooperation between PGE2 and TNF signaling, but mechanisms that induce these cells and the extent to which they contribute to arthritic phenotypes are not known. In this study we used an integrated transcriptomic and epigenomic analysis in primary human monocytes to study PGE2-TNF crosstalk, and how it is regulated by IFN-γ, as occurs in RA synovial macrophages. We identified a (TNF + PGE2)-induced gene expression signature that is enriched in an IL1β+ RA macrophage subset defined by scRNAseq and includes genes in pathogenic IL-1, Notch and neutrophil chemokine pathways. A similar gene expression signature was apparent in an IL-1β+ macrophage subset newly identified by scRNAseq in ICI-arthritis. TP signature genes are distinct from canonical inflammatory NF-κB target genes such as *TNF*, *IL6* and *IL12B* and are activated by cooperation of PGE2-induced AP-1, CEBP and NR4A family transcription factors with TNF-induced NF-κB activity. Unexpectedly, IFN-γ suppressed induction of AP-1, CEBP and NR4A activity to ablate induction of IL-1, Notch and neutrophil chemokine genes, while promoting expression of distinct inflammatory genes such as *TNF* and T cell chemokines like CXCL10. These results reveal the basis for synergistic induction of inflammatory genes by PGE2 and TNF, and a novel regulatory axis whereby IFN-γ and PGE2 oppose each other to determine the balance between two distinct TNF-induced inflammatory gene expression programs relevant for RA and ICI-arthritis.

Introduction

Prostaglandin E2 (PGE2) is a small lipid molecule with homeostatic functions in various tissues whose production is strongly upregulated after tissue injury or inflammatory signaling. PGE2 plays a key role in early acute inflammation by increasing vascular permeability and activating mast cells, with associated tissue edema and influx of immune cells such as neutrophils (Kawahara et al., 2015; Tsuge et al., 2019). PGE2 is also a key mediator of acute pain, in part by acting directly on nociceptors (Kawabata, 2011). Inhibition of these PGE2-mediated functions explains the effectiveness of NSAID drugs that suppress PGE2 production in alleviating acute pain and tissue swelling. In contrast to activating mesenchymal cells, PGE2 has generally been considered to have primarily suppressive effects on both innate and adaptive immune cells (Chen et al., 2012; Kawahara et al., 2015; Luan et al., 2014; MacKenzie et al., 2013; Perretti et al., 2017; Rodriguez et al., 2014; Sundberg et al., 2014; Yokoyama et al., 2013). This includes inhibition of Th1 cells and IFN-γ production, and inhibition of induction of inflammatory genes such as *TNF*, *IL12B* and *IFNB* in macrophages and dendritic cells. Such inhibition raises the possibility that expression of these genes increases after NSAID therapy, which would contribute to lesser effectiveness and lack of disease-modifying activity of NSAIDS in chronic inflammatory conditions such as rheumatoid arthritis (RA) (Smolen et al., 2016).

In myeloid cells, PGE2 signals primarily via EP2 and EP4 receptors, which are G protein coupled receptors (GPCRs) that elevate intracellular cAMP to directly activate protein kinase A (PKA) and signaling effector EPAC, which is coupled to small GTPases Rap1/2 (Yokoyama et al., 2013). The immune suppressive effects of PGE2 have been attributed to PKA-mediated activation of transcription factor CREB and histone deacetylases 4 and 5 (HDAC4/5). CREB induces expression of transcriptional repressors such as *CREM* and *ATF3*, and CREB can suppress NF-κB activity in a gene-specific manner by mechanisms that are not fully understood but include competition for transcription coactivator CBP (Altarejos and Montminy, 2011; Gerlo et al., 2011). HDACs 4/5 bind to inflammatory gene loci such as *Tnf* and *Il12b* and suppress transcription (Luan et al., 2014). Additionally, PGE2 suppresses TLR-induced autocrine IFN responses by suppressing TLR4 trafficking and signaling via TRIF (Perkins et al., 2018), preventing induction of *Ifnb1* and de novo enhancers by suppressing activity of transcription factor MEF2A (Cilenti et al., 2021), and attenuating induction of IRF8 target genes by suppressing IRF8 expression (Bayerl et al., 2023). In accord with suppression of immune cells, activation of cAMP signaling has shown therapeutic benefit in various inflammatory conditions (Tsuge et al., 2019; Yokoyama et al., 2013). Inhibitors of phosphodiesterase 4, which increase intracellular cAMP in immune cells, have been approved by the FDA for treatment of psoriatic arthritis, asthma and COPD (Li et al., 2018). However, the efficacy of these therapies may be limited by concomitant pro-inflammatory effects of cAMP signaling, such as cell-, gene- and context-specific augmentation of NF-κB activity (Gerlo et al., 2011). The importance of deciphering mechanisms by which PGE2 activates inflammatory gene expression in macrophages is highlighted by a recent study showing that cooperation between PGE2 and TNF induces IL-1β+ tumor associated macrophages that are associated with disease progression in pancreatic ductal adenocarcinoma (Caronni et al., 2023).

Activated monocytes and macrophages that infiltrate inflamed synovium (joint tissues) in rheumatoid arthritis (RA) and produce inflammatory mediators and cytokines have been strongly implicated in disease pathogenesis (Gravallese and Firestein, 2023; Kalliolias and Ivashkiv,

2016; McInnes and Schett, 2011). High level expression of inflammatory genes in RA synovitis has long been appreciated, and recent studies using high dimensional single cell profiling technologies have identified subsets of synovial monocytes/macrophages (hereafter termed MonoMacs), and highlighted a pervasive IFN signature and likely pathogenic myeloid subsets that coordinately express NF-κB target genes and interferon-stimulated genes (Alivernini et al., 2020; Kuo et al., 2019; Lewis et al., 2019; Mandelin et al., 2018; Orange et al., 2018; Zhang et al., 2023; Zhang et al., 2019). One RA synovial macrophage subset, termed cluster 1 in ref.(Kuo et al., 2019) was increased in RA relative to osteoarthritis samples, was proposed to be pathogenic based on elevated expression of inflammatory genes, and was notable for high *IL1B* expression. IL-1β+ monocytes and macrophages have also been observed in ICI-arthritis that occurred after anti-PD-1 therapy (Zhou et al., 2024). In RA, the macrophage cluster 1 gene expression signature could be modeled by coculture of monocytes with synovial fibroblasts in the presence of TNF, and a substantial fraction of the fibroblast + TNF effect was dependent upon fibroblast-produced PGE2 (Donlin et al., 2014; Kuo et al., 2019). Notably, TNF-induced and PGE2-mediated fibroblast-monocyte crosstalk cooperatively induced expression of potentially pathogenic cluster 1 genes such as *HBEGF*, *EREG*, *IL1B* and transcription factor *STAT4*. Other studies also found that early RA synovial biopsy tissues showed enrichment of 'eicosanoid' and 'cAMP mediated signaling' pathways (Lewis et al., 2019), and RNAseq of purified macrophages from synovial biopsies showed 6 gene modules, one of which showed enrichment for "G-protein coupled receptor signaling pathway" (Mandelin et al., 2018). Collectively, these studies establish activation of the PGE2-cAMP pathway in a likely pathogenic subset of RA synovial macrophages, and suggest that crosstalk of PGE2-cAMP with TNF and possibly IFN signaling is important for the pathogenic phenotype.

We wished to understand the full spectrum of pathogenic inflammatory genes induced by cooperation between PGE2 and TNF signaling, to gain insight into mechanisms underlying signaling crosstalk, and address the important question of how PGE2-TNF crosstalk is modulated by IFN-γ, as occurs in RA synovial macrophages. We performed an integrated transcriptomic and epigenomic analysis of gene expression and chromatin accessibility using primary human monocytes, which correspond directly to cells that migrate into inflamed synovium and provide an opportunity to model in vivo activation. We identified a 'TNF + PGE2' (TP) gene expression signature that is enriched in RA synovial macrophage subsets and includes genes in pathogenic IL-1, Notch and neutrophil chemokine pathways. A similar gene expression signature was also apparent in ICI-arthritis. Expression of the TP signature was driven by cooperation of PGE2-induced AP-1, CEBP and NR4A family transcription factors with TNF-induced NF-κB activity. IFN-γ suppressed induction AP-1, CEBP and NR4A activity and suppressed IL-1, Notch and neutrophil chemokine genes, while promoting expression of distinct inflammatory genes and T cell chemokines. These results reveal a novel regulatory axis whereby IFN-γ and PGE2 oppose each other to determine the balance between two distinct TNF-induced inflammatory gene expression programs.

Results

PGE2 augments expression of a large subset of TNF-inducible genes

We used RNAseq to perform a transcriptomic analysis of the effects of PGE2 on the TNF response in primary human monocytes; the experimental design is depicted in Fig. 1A. As expected, co-stimulation of monocytes with PGE2 + TNF induced expression of genes enriched in inflammatory and NF-κB pathways, largely reflecting a TNF response (Fig. 1B). Clustering of genes upregulated after 3 hours of stimulation (n = 747; fold change > 2, FDR < 0.05) based upon pattern of expression confirmed that PGE2 suppresses expression of a subset of TNF-inducible genes including *TNF* itself and interferon-stimulated genes *CXCL9*, *CXCL10* and *CXCL11* (Fig. 1C, group III, n = 86 suppressed genes). Surprisingly, PGE2 augmented TNF-mediated induction of a substantially larger number of genes (Fig. 1C, groups I and V, 395 co-stimulated genes). Pathway analysis revealed that genes costimulated by PGE2 + TNF were similarly enriched in inflammatory pathways (e.g. response to lipopolysaccharide, neutrophil migration) although the enriched pathways were partially distinct (Fig. 1D). We also observed 2 groups of genes that were induced by PGE2 alone, which included canonical cAMP signaling and CREB target genes (Zhang et al., 2005) such as *HBEGF* and *CREM* (Fig. 1C, groups 2 and IV). RT-qPCR analysis confirmed strong costimulation of *IL1B*, *IL1A*, and *CSF3*, induction of *HBEGF* by PGE2 alone, and suppression of TNF-induced expression of *TNF* and *CXCL10* by PGE2 (Fig. 1E). Similar results on regulation of TNF-mediated gene expression by PGE2 were observed using mouse bone marrow-derived macrophages (Supplementary Fig. 1). Costimulation of STAT4 was confirmed at the protein level (Fig. 1F). Analysis of RNAseq performed at the 24 hr time point to determine the effects of PGE2 on the late phase TNF response showed similar results and additionally revealed that PGE2 inhibits TNF-induced expression of cholesterol pathway genes, which we had previously shown to be induced by TNF with delayed kinetics (Kusnadi et al., 2019) (Supplementary Fig. 2). Collectively, these results indicate that PGE2 has a dichotomous effect on TNF-induced gene expression, superinducing select key inflammatory genes such as *IL1B*, while suppressing expression of distinct inflammatory genes including *TNF* and interferon-stimulated genes in line with previous reports (Chen et al., 2012; Cilenti et al., 2021; Gerlo et al., 2011; Kawahara et al., 2015; Luan et al., 2014; MacKenzie et al., 2013; Sundberg et al., 2014; Tsuge et al., 2019).

PGE2 and TNF costimulation model aspects of the RA and ICI-arthritis synovial macrophage phenotypes

The above-described results were similar to results we had previously observed using TNF-stimulated monocyte – synovial fibroblast cocultures (Donlin et al., 2014; Kuo et al., 2019), except that PGE2-inducible genes such as *HBEGF* were strongly induced by PGE2 alone, and expression was not further increased by TNF (Fig. 1C, groups 2 and 4). These differences may reflect delayed kinetics of induction of fibroblast-derived PGE2 in cocultures, or from the modulating activity of various factors and cytokines produced by activated fibroblasts in these cultures. To assess the extent to which TNF-stimulated monocyte-fibroblast cross talk could be recapitulated by stimulation of monocytes with purified PGE2 and TNF, we compared the current (PGE2+TNF)-costimulated gene set with the TNF/fibroblast-costimulated gene set previously reported (Donlin et al., 2014; Kuo et al., 2019). Strikingly, (PGE2+TNF)-costimulated genes recapitulated 61% of fibroblast-augmented TNF-inducible genes (Fig. 2A, hypergeometric p = 8.36e-128), which is in accord with our previous conclusions that PGE2 mediates a substantial fraction of the fibroblast effect on monocytes (Kuo et al., 2019). Since TNF-stimulated fibroblast-cocultured monocytes modeled the phenotype of RA synovial macrophage cluster 1 (C1, also termed HBEGF+ or IL1B+ macrophages) (Kuo et al., 2019; Zhang et al.,

2023; Zhang et al., 2019), we next tested whether (PGE2+TNF)-stimulated monocytes expressed the defining genes of cluster 1 (128 genes, as defined in ref.(Kuo et al., 2019)). Strikingly, (PGE2 + TNF)-costimulated genes (Fig. 2B, orange + yellow) highly significantly mimicked the RA C1 phenotype (p < 10e-9 by Monte Carlo simulation), and recapitulated the C1 phenotype more completely than TNF/fibroblast-costimulated genes (Fig. 2B, orange + red). The (PGE2 + TNF)-costimulated genes were parts of pathogenic gene modules related to IL-1-NF-κB and Jak-STAT signaling, neutrophil chemotaxis, and the Notch pathway that has been recently implicated in RA pathogenesis (Wei et al., 2020; Zack et al., 2024) (Fig. 2C, blue font = C1 genes). Additional examples of induction of RA macrophage C1 genes by PGE2 and TNF are depicted in Fig. 2D. These results link (PGE2 + TNF)-costimulated macrophages to a pathogenic RA macrophage phenotype, and reveal that, instead of global suppressive effects in human MonoMacs, PGE2 promotes induction of genes important in various pathogenic inflammatory pathways.

We wished to test whether macrophage subsets expressing TP-costimulated genes could be detected in other types of inflammatory arthritis and thus investigated synovial macrophages from patients with ICI-arthritis. While immune checkpoint inhibitors are effective in the treatment of malignancies, ICI can lead to autoimmune side effects including inflammatory arthritis (Dougan et al., 2021). Approximately 5% of ICI-treated patients can develop autoimmune inflammatory arthritis that persists despite cessation of ICI therapy (Braaten et al., 2020), and understanding mechanisms and pathways that activate immune cells in ICI-arthritis may yield insights into pathogenesis of spontaneous autoimmune arthritides (Cappelli et al., 2020). We performed single cell RNA sequencing analysis of mononuclear cells FACS-sorted from synovial fluids of five ICI-arthritis patients, and from two synovial tissues from one ICI-arthritis patient who underwent bilateral knee replacements. These patients had received anti-PD1 antibodies as monotherapy, or in combination with anti-CTLA4 therapy (clinical characteristics of patients are presented in Supplementary Table 1). Following subclustering of monocytes and macrophages, we defined eight clusters from 14,110 cells (Fig. 2E). Cluster 4, accounting for 15% of cells, selectively showed elevated expression of PGE2+TNF-costimulated genes and known PGE2 targets including *HBEGF*, *IL1B*, *CXCL2/3/8*, *CREM* and *PLAUR* (Fig. 2F, genes marked in red). Cluster 4 macrophages are distinct from those in the other clusters, such as clusters 0 and 1 that expresses interferon-stimulated and different inflammatory genes. These results suggest that costimulation of gene expression by TNF and PGE2 can occur in ICI-arthritis.

cAMP signaling has dichotomous suppressive and augmenting effects on the TNF-induced inflammatory response

We wished to test whether both the pro- and anti-inflammatory effects of PGE2 on TNF-induced gene expression were mediated by cAMP signaling. We addressed this question using selective agonists of PGE2 receptors EP2 and EP4 that signal predominantly via cAMP, and also cell membrane-permeable dibutyryl-cAMP that directly activates this pathway. Selective agonists of EP2 and EP4 augmented TNF-induced expression of *IL1B, IL1A, STAT4 and CXCL1*, while suppressing induction of the ISG *CXCL10* (Fig. 3A). Dibutyryl-cAMP showed similar results (Fig. 3B). These results implicate cAMP signaling in both anti- and pro-inflammatory effects in this experimental system.

PGE2 effects on TNF-induced changes in chromatin accessibility

We then performed ATACseq analysis of chromatin accessibility (Supplementary Fig. 3A) to gain insight into how PGE2 costimulates TNF-induced gene expression. In accord with different signaling pathways activated by these factors, TNF and PGE2 induced mostly distinct ATACseq peaks (Fig. 4A); indeed, out of 15,190 induced ATACseq peaks (FDR < 0.05, fold induction > 2), only 1547 peaks (10.2%) were commonly induced by individual stimulation by TNF or PGE2 (Fig. 4A, columns 4 + 7). As expected, transcription factor binding motifs most significantly enriched under TNF-induced ATACseq peaks corresponded to AP-1 and NF-κB binding sites; additional enriched motifs included RUNX and IRF1 sites, the latter of which is in accord with the role of IRF1 in the delayed TNF-induced autocrine IFN response (Yarilina et al., 2008) (Fig. 4B, left). Similar to TNF, PGE2 induced peaks enriched for AP-1 motifs, and additionally induced peaks with highly significant enrichment of CEBPD and NR4A1 motifs (Fig. 4B, middle), which is in accord with PGE2-medated induction of expression of transcription factors (TFs) belonging to these three TF families (Altarejos and Montminy, 2011; Zhang et al., 2005). In line with the largely distinct open chromatin regions (OCRs) induced by TNF and PGE2, GREAT analysis of genes associated with these peaks showed enrichment of genes in largely distinct pathways, with TNF-induced peaks associated with genes in immune and cytokine pathways, whereas PGE2-induced peaks were associated with genes in cell matrix interactions and chemotaxis (Supplementary Fig. 3B and 3C).

To examine how these disparate factors could interact to costimulate gene expression, we extended our analysis to peaks that were co-induced by both TNF and PGE2 (Fig. 4A, column 4; n = 1531) and peaks that were induced only under conditions of costimulation (Fig. 4B, 2nd column, n = 3337; termed T+P unique). TNF and PGE2 cooperated to increase chromatin accessibility at co-induced peaks (Fig. 4C, upper), and the most significantly enriched motifs were AP-1 (Fra1, also known as FOSL1) and CEBPE (Fig. 4C, lower). These co-induced peaks were associated with immune and inflammatory response genes (Supplementary Fig. 3D). The peaks that were uniquely induced by PGE2 + TNF were most significantly enriched for AP-1, CEBP and ISRE/IRF motifs, and also for GAPBA, KLF7 and PBX1 binding sites (Fig. 4B, right and data not shown), and were associated with immune response and cytokine genes (Supplementary Fig. 3E). Collectively, the results suggest that TNF and PGE2 signals converge on gene regulatory elements associated with inflammatory genes to increase chromatin accessibility and induce de novo enhancers, and highlight a potential role for AP-1 and CEBP transcription factors (TFs) in costimulating TNF-induced gene expression.

We then used ChromVAR (Schep et al., 2017) to assess induction of transcription factor activity by PGE2, TNF, or their combination (Fig. 4D). This analysis reinforced the notion that, relative to TNF alone, the PGE2 + TNF costimulation condition was characterized by increased AP-1 and CEBP activity, and also NR4A1 activity. We then performed a TOBIAS analysis that uses footprinting to measure actual occupancy of TF motifs within ATACseq peaks (Bentsen et al., 2020). TNF stimulation induced most significant occupancy of NF-κB and AP-1 motifs, whereas PGE2 stimulation induced AP-1 and CEBP occupancy (Supplementary Fig. 4F). A direct pairwise comparison of T+P to T revealed most significant increases in AP-1 and CEBP occupancy (Fig. 4E). Overall, all three complementary approaches show that, relative to TNF alone, PGE2 costimulation boosts AP-1 and adds CEBP activation, and these factors can

cooperate to increase chromatin accessibility at commonly targeted sites, and to induce de novo enhancer formation.

IFN-γ opposes the effects of PGE2 on TNF-induced gene expression

In inflamed RA synovium, macrophages that express a 'TP signature' are also exposed to IFN-γ (Kuo et al., 2019; Orange et al., 2018; Zhang et al., 2023; Zhang et al., 2021; Zhang et al., 2019), which not only induces ISGs but also augments TLR- and TNF-induced expression of various inflammatory NF-κB target genes including *TNF*, *IL6*, and *IL12* family members (Mishra and Ivashkiv, 2024). We next examined the effects of IFN-γ on expression of (PGE2 + TNF)-costimulated inflammatory genes. Surprisingly, IFN-γ essentially completely abolished induction of inflammatory genes *IL1B* and *CXCL2*, and the Notch target gene *HEY1* (Fig. 5A); as a specificity control, induction of *STAT4* was minimally affected. We next tested whether IFN-γ broadly suppressed (T+P)-costimulated genes by analyzing the effects of IFN-γ on induction of subsets of these genes in our RNAseq data set, as defined in Supplementary Fig. 4A. Interestingly, IFN-γ suppressed expression of only approximately 27% of genes that were most clearly synergistically induced by (T + P) (Fig. 5B; this group included IL-1 pathway genes *NLRP3* and *PTGS2*). Similarly, IFN-γ suppressed induction of only a subset of additional genes that were costimulated by (T + P) (Supplementary Fig. 4B; these suppressed genes included IL-1 family members *IL1B* and *IL1A*, neutrophil chemokine genes *CXCL1/2/3/5/8* and canonical cAMP signaling targets such as *EREG* and *NR4A1/3*). Thus, IFN-γ suppressed expression of genes in the key RA pathogenic pathways described in Figure 2. This opposition between IFN-γ and PGE2 was also apparent in the regulation of a subset of TNF-induced genes that were suppressed by PGE2, whose expression was substantially higher under IFN-γ-stimulated conditions (Fig. 5C and 5D). Genes more highly expressed under IFN-γ-stimulated conditions included inflammatory genes such as *TNF* and ISGs such as *CXCL10*. Collectively, the results support a biology whereby PGE2 and IFN-γ at least in part oppose each other by regulating TNF-induced gene expression in different directions, and suggest that IFN-γ inhibits select inflammatory pathways while promoting others.

To obtain a broader view of the effects of IFN-γ on the entire TP-induced gene response, we performed a general linear model with between-treatment interaction contrast using our RNAseq data set (Fig. 6A and 6B). The conclusion that IFN-γ opposes a large component of the (TNF + PGE2) response was supported by identification of 3 positive interaction clusters (ICs) comprised of genes regulated in opposing directions when IFN-γ was added to (T+P) (Fig. 6A, interaction clusters IC2, IC3 and IC6 where red and blue lines diverge). Genes whose suppression by TP was reversed by IFN-γ (IC2) were enriched in IFN pathways and regulated by IRF transcription factors, implicating IRFs in opposing negative regulation of gene expression by (T+P). In contrast, genes that were induced by (T+P) but strongly suppressed when IFN-γ was added to the TP condition were enriched in Myc targets, suggesting counter-regulation of metabolic pathways in these terminally differentiated and nonproliferating cells. Interestingly, IFN-γ minimally affected the (T+P)-induced NF-κB response (IC4). Genes that comprise the two negative interaction clusters are upregulated (IC1) or downregulated (IC5) by IFN-γ or (T+P), but in combination IFN-γ did not provide an additional effect.

To explore the regulatory drivers of interaction clusters we selected all interacting differentially expressed transcription factors (|logFC|>1 in the interaction contrast) and built a protein association network using stringDB (Figure 6C). The size of a node reflects the number of unique functional connections between a node and other nodes in the network and edges connect associated nodes. Notably, TFs were grouped together with their targets in several ICs, for example IC2 contains several IRF2 and IRF8 and their targets; IC4 contains NF-κB1,2 and RELB and their targets; IC6 contains Myc and its targets (Figure 6 B and C). Although RNA levels of Jun are regulated only weakly its ultimate activity and binding site repertoire is affected by relative levels of dimerization partners and interacting proteins (Chang et al., 2018). For example, BATF2 (IC2) and BATF (IC6) were up and down regulated by the combination of IFN-γ and (T+P) relative to (T+P), which potentially affects relative abundance of JUN/BATF/IRF ternary complexes. Similarly, CEBPA, CEBPE (IC3), and MYC (IC6) are regulated in a reciprocal fashion: IFN-γ relieves a (T+P) inhibition of CEBPA and CEBPE yet also concomitantly decreases MYC mRNA levels. A recent report describing negative regulation of Myc by CEBPA/E (Theilgaard-Mönch et al., 2022) suggests that a similar mechanism is involved in IFN-γ dependent inhibition of Myc and its targets. Overall, the interaction analysis indicates that IFN-γ does not alter the core NF-κB response (which is activated primarily by TNF) and suggest that instead IFN-γ regulates the PGE2 response, at least in part by modulating PGE2-induced AP-1 and CEBP factors.

IFN-γ inhibits PGE2-induced gene expression and chromatin accessibility

We tested this notion that IFN-γ regulates the PGE2 response using combined transcriptomic and chromatin accessibility analysis. Analysis of our RNAseq data (Supplementary Fig. 5A) revealed two clusters of genes C2 and C5 that were induced by PGE2 but not by IFN-γ (Fig. 7A); the great majority of these PGE2-inducible genes were inhibited by IFN-γ, and C2 genes were more strongly inhibited than C5 genes. Strikingly, IFN-γ suppressed induction of canonical PGE2-cAMP target genes such as *CREM, HBEGF, HIF1A, PLAUR, KDM6B, AREG, VEGFA*, and genes in RA pathogenic gene modules and expressed in RA synovial macrophage cluster 1, such as IL-1 and Notch pathway genes and neutrophil chemokines CXCL1/2/3/5/8 (Fig. 7A, 7B, and Supplementary Fig. 5B). Thus, IFN-γ broadly suppresses the PGE2 response, including genes important in RA pathogenesis.

We then analyzed our ATACseq data to gain insight into mechanisms underlying negative regulation of PGE2 responses by IFN-γ (Supplementary Fig. 5C and Fig. 7C). IFN-γ suppressed 1765 out of 2411 (72%) of PGE2-induced ATACseq peaks (Fig. 7C, column G2 depicts suppressed peaks). These IFN-γ-suppressed peaks were most significantly enriched in AP-1 and CEBP motifs (Fig. 7D). ChromVAR analysis confirmed that PGE2 induced AP-1 activity, and this activity was suppressed by IFN-γ (Fig. 7E). Additionally, NR4A1/2, NFE2L2 and MAF activity was induced by PGE2 and suppressed by IFN-γ. In contrast to the HOMER de novo motif analysis described above that focused on IFN-γ-suppressed peaks, ChromVAR analysis, which takes into account all peaks showed that both PGE2 and IFN-γ induced CEBP activity. Remarkably, at the mRNA expression level, PGE2 and IFN-γ both induced and repressed different members of the CEBP TF family, with coordinate induction of CEBPB, but opposing regulation of CEBPA, CEBPD and CEBPE (Fig. 7F). TOBIAS footprinting analysis showed that both PGE2 (see Fig. 4E above) and IFN-γ induced CEBP occupancy (Fig. 7G); the

latter confirms findings in our previous report that used a different independent data set (Mishra and Ivashkiv, 2024). Direct comparison of IFN-γ + PGE2 relative to PGE2 alone using TOBIAS showed striking suppression of AP-1 occupancy and further supported IFN-γ-mediated suppression of NR4A2, NFE2L2 and MAF (Fig. 7H). Collectively, these results show that IFN-γ inhibits PGE2 responses by suppressing AP-1 and a PGE2-induced transcriptional program mediated by AP-1, NR4A1/2, NFE2L2 and MAF transcription factors, and suggest that IFN-γ and PGE2 costimulation remodels the composition and genomic binding profile of CEBP proteins to be distinct from that induced by each stimulus alone.

Discussion

IL-1β-expressing macrophages have been implicated in pathogenesis of RA, ICI-arthritis, and pancreatic cancer but mechanisms that induce these cells and the extent to which they contribute to arthritic phenotypes are not known. In this study, we utilized transcriptomics to demonstrate a 'TNF + PGE2' (TP) signature in RA and ICI-arthritis IL-1β+ synovial macrophage subsets that were defined by single cell RNA sequencing, and investigated mechanisms of crosstalk between PGE2 and TNF that costimulated expression of inflammatory and pathogenic genes. Epigenomic analysis revealed cooperation of PGE2-induced AP-1, CEBP and NR4A family transcription factors with TNF-induced NF-κB activity to drive expression of pathogenic gene modules including IL-1, Notch and neutrophil chemokine pathways. The genes in these pathways are distinct from canonical inflammatory genes such as *TNF*, *IL6* and *IL12B* that are driven by core NF-κB and IRF signaling. Unexpectedly, IFN-γ suppressed AP-1 and NR4A, and altered CEBP activity to ablate induction of IL-1, Notch and neutrophil chemokine genes, while promoting expression of distinct inflammatory genes such as *TNF* and T cell chemokines such as CXCL10. These results reveal the basis for synergistic induction of inflammatory genes by PGE2 and TNF, and a novel regulatory axis whereby IFN-γ and PGE2 oppose each other to determine the balance between two distinct TNF-induced inflammatory gene expression programs (Fig. 8A and 8B).

TNF and microbial products that engage Toll-like receptors (TLRs) activate a core set of inflammatory genes such as *TNF*, *IL6*, *IFNB* and *IL12B* that are NF-κB targets and whose expression is amplified by IFNs via induction of IRF factors that cooperate with NF-κB (Mishra and Ivashkiv, 2024) (Fig. 8A). Expression of these 'signature' inflammatory genes is also augmented by innate immune training, which is mediated in many contexts by IFNs and IRFs (Mishra and Ivashkiv, 2024). In contrast to IFNs that are generally thought to be 'pro-inflammatory' (Ivashkiv, 2018), PGE2-cAMP signaling in innate immune cells is considered anti-inflammatory based upon inhibition of *TNF* and ISGs such as *CXCL10* (Kawahara et al., 2015; Tsuge et al., 2019; Yokoyama et al., 2013) (Fig. 8A). Recent reports by our and other laboratories showed that PGE2 + TNF drive expression of EGFR ligands such as *HBEGF* in myeloid cells, thereby promoting proliferation and invasive behavior of synovial fibroblasts (Kuo et al., 2019), expansion of pancreatic tumors (Caronni et al., 2023), and survival and reparative capacities of intestinal epithelial cells (Zhou et al., 2022). Thus, TP costimulation can have pathogenic or protective roles, depending on context. The current study further extends this paradigm by showing that TP costimulation promotes expression of additional pathogenic modules in RA including IL-1-related genes that promote tissue degradation, Notch ligands that can activate synovial fibroblasts (Wei et al., 2020; Zack et al., 2024), and neutrophil chemokines

that can contribute to initiation or flares of inflammatory arthritis (Gravallese and Firestein, 2023; Zec et al., 2023).

In contrast to cooperation of IRFs with NF-κB that drives signature inflammatory genes, our results suggest that the distinct TP gene expression program is promoted by cooperation of NF-κB with CEBP, AP-1 and NR4A1/2 transcription factors (Fig. 8A, left). CEBP is a pioneer TF that can open chromatin, and its induction helps explain the formation of de novo enhancers under costimulation conditions. The above-mentioned PGE2-induced TFs are well-established target genes for canonical cAMP-PKA-CREB signaling (Altarejos and Montminy, 2011; Gerlo et al., 2011; Zhang et al., 2005), which is in accord with our results implicating this pathway downstream of PGE2 and EP2 and EP4 receptors. A role for CREB was not identified by our experimental approaches, possibly because the number of direct CREB target genes was too small to pass statistical significance thresholds, or because CREB binds to its target genes under basal conditions and is activated by phosphorylation rather than changes in DNA-binding (Altarejos and Montminy, 2011). Interestingly, IFN-γ suppressed the PGE2-induced transcriptional program by inhibiting the activity of AP-1 and NR4A1/2 and altering the expression profile of CEBP, which was associated with abrogation of expression of IL-1, Notch and neutrophil chemokine gene modules. These results identify an unexpected dual role for IFN-γ and suggest that the inflammatory profile of TNF signaling is regulated by the local balance between PGE2 and IFN-γ activity (Fig. 8B).

Major sources of PGE2 in inflammatory arthritis and tumors are stromal cells such as fibroblasts, and tumor cells (Bayerl et al., 2023; Gong et al., 2023; Kawahara et al., 2015; Kuo et al., 2019; Tsuge et al., 2019). This argues that macrophages in tissue niches characterized by interaction with fibroblasts will strongly express the PGE2-driven gene expression program including expression of EGFR and Notch ligands that activate fibroblasts, suggestive of a positive feedback loop. Associated high expression of TP-induced neutrophil chemokines would promote neutrophil-mediated inflammation. IL-1 action on myeloid cells strongly induces COX2 and thus PG production, and an autocrine IL-1-PGE2 loop can further amplify the TP response, as shown in pancreatic tumors (Caronni et al., 2023). In contrast, the major sources of IFN-γ in RA are T cells, which are localized either to ectopic germinal center-like structures or are dispersed throughout inflamed synovium (Gravallese and Firestein, 2023). IFN-γ is also highly expressed by CD8+ T cells in ICI-arthritis (Wang et al., 2023). Our results suggest that macrophages exposed to high concentrations of IFN-γ, such as those in germinal center-like niches and in ICI-arthritis, will express an attenuated PGE2 program, which is in accord with previously reported synovial macrophage clusters that express a predominant ISG signature and lower expression of the IL-1 pathway (Kuo et al., 2019; Zhang et al., 2023; Zhang et al., 2021; Zhang et al., 2019). The segregation of macrophage clusters expressing TP signature genes and ISGs in our ICI-arthritis data set further supports this idea. IFN-γ-exposed macrophages express T cell chemokines such as CXCL10 instead of neutrophil chemokines, thereby promoting T cell mediated pathogenesis instead of stromal-neutrophil mediated pathogenic responses. Based upon our previous work (Mishra and Ivashkiv, 2024; Qiao et al., 2013) and the current study, these macrophages would be predicted to express signature NF-κB target genes such as *TNF*. These ideas will need to be tested more extensively in future work using spatial transcriptomics, anticipating that gradients of PGE2 and IFN concentrations across inflamed synovium will give rise to macrophages with mixed phenotypes, as suggested

by the reported scRNAseq data (Alivernini et al., 2020; Kuo et al., 2019; Zhang et al., 2023; Zhang et al., 2019). Additionally, distinct subsets of RA patients may exhibit fibroblast/PGE2 versus T cell/IFN-γ dominant phenotypes and thus be differentially responsive to therapeutics.

In RA, macrophages are major producers of TNF, which is a well-established therapeutic target. However, only a subset of RA patients show >50% improvement after TNF blockade therapy, and even these patients can become resistant to therapy (Perera et al., 2024). One current explanation for resistance to TNF blockade therapy is that in subsets of RA patients synovitis is driven by distinct pathogenic mechanisms; additionally these different mechanisms can emerge when TNF activity is blocked which can explain acquired resistance to TNF blockade. Such distinct mechanisms of RA pathogenesis include those driven by IL-6 and IL-1 instead of TNF (Gravallese and Firestein, 2023; McInnes and Schett, 2011). While IL-6 blockade therapy is highly effective in a subset of RA patients, IL-1 blockade with anakinra, although an FDA-approved therapy for RA that is effective at suppressing tissue degradation, appears to have limited efficacy in suppressing inflammatory symptoms. One explanation for potential low efficacy in suppressing symptoms of inflammation such as joint swelling is that IL-1 pathways are not primarily coupled with symptoms of inflammation, but instead with tissue degradation via their effects on synovial fibroblasts. Alternatively, our work raises the intriguing possibility that interrupting the IL-1-PGE2 axis using IL-1 blockade will release genes such as *TNF* and *CXCL10* from suppression and result in the emergence of stronger TNF- and lymphocyte-driven pathogenic pathways that maintain synovitis. If substantiated, this notion would promote the idea of combination therapy of using anakinra with either TNF blockade or T cell-targeted therapy.

Methods

Primary human monocyte isolation and culture. Deidentified buffy coats were purchased from the New York Blood Center following a protocol approved by the Hospital for Special Surgery Institutional Review Board. Peripheral blood mononuclear cells (PBMCs) were isolated using density gradient centrifugation with Lymphoprep (Accurate Chemical) and monocytes were purified with anti-CD14 magnetic beads from PBMCs immediately after isolation as recommended by the manufacturer (Miltenyi Biotec). Monocytes were cultured at 37°C, 5% CO_2 in RPMI-1640 medium (Invitrogen) supplemented with 10% heat-inactivated defined FBS (HyClone Fisher), penicillin-streptomycin (Invitrogen), L-glutamine (Invitrogen) and 20 ng/ml human M-CSF.

Mouse bone marrow-derived macrophage culture. Animal experiments were approved by the Weill Cornell Medicine IACUC Committee. Male C57BL/6J mice at 6 to 8 weeks old were purchased from the Jackson Laboratories and housed under specific pathogen-free conditions. Bone marrow cells were harvested after euthanasia by CO_2 asphyxiation, and cultured in RPMI-1640 medium (Invitrogen) supplemented with 10% heat-inactivated defined FBS (HyClone Fisher), penicillin-streptomycin (Invitrogen), L-glutamine (Invitrogen) and 20 ng/ml mouse M-CSF.

Analysis of mRNA amounts (qPCR). Total RNA was isolated using the RNeasy Mini Kit (QIAGEN) following the manufacturer's instructions. Reverse transcription of RNA into complementary DNA (cDNA) was performed using the RevertAid RT Reverse Transcription Kit (Thermo Fisher Scientific) according to the manufacturer's protocol, and the resulting cDNA was used for downstream analysis. For quantitative real-time PCR (qPCR), Fast SYBR Green Master Mix (Applied Biosystems) and a QuantStudio5 Real-time PCR system (Applied Biosystems) were used. CT values obtained from qPCR were normalized to the housekeeping gene *GAPDH*. Relative expression of target genes was calculated using the ΔCt method, where ΔCt represents the difference in threshold cycle values between the target gene and *GAPDH*. The results are presented as a percentage of *GAPDH* expression (100/2^ΔCt).

Western blotting. Primary human monocytes were washed with cold PBS after indicated treatments and harvested in 50uL cold lysis buffer containing Tris-HCl pH 7.4, NaCl, EDTA, Triton X-100, Na3VO4, phosSTOP EASYPACK, Pefabloc, and EDTA-free complete protease inhibitor cocktail. After a 10-minute ice incubation, cell debris was pelleted at 16,000xg at 4°C for 10 minutes. The soluble protein fraction was combined with 4× Laemmli Sample buffer containing 2-mercaptoethanol and subjected to SDS-PAGE (electrophoresis) on 4–12% Bis-Tris gels. Following transfer of gels to polyvinylidene difluoride membranes, membranes were blocked in 5% (w/v) Bovine Serum Albumin in TBS with Tween-20 (TBST) at room temperature for at least one hour. Incubation with primary antibodies (diluted 1:1000 in blocking buffer) was performed overnight at 4°C. Membranes were washed three times with TBST and probed with anti-rabbit IgG secondary antibodies conjugated to horseradish peroxidase (diluted 1:5000 in blocking buffer) for one hour at room temperature. Enhanced chemiluminescent substrates (ECL western blotting reagents (PerkinElmer, cat: NEL105001EA)) were used for detection, followed by visualization on autoradiography film (Thomas Scientific, cat: E3018). Restore PLUS western blotting stripping buffer (Thermo Fisher Scientific) was applied for membranes requiring sequential probing with different primary antibodies.

RNA sequencing and data analysis. Libraries for sequencing were prepared using mRNA that was enriched from total RNA using NEBNext® Poly(A) mRNA Magnetic Isolation Module, and enriched mRNA was used as an input for the NEBNext Ultra II RNA Library Prep Kit (New England Biolabs (NEB)), following the manufacturer's instructions. Quality of all RNA and library preparations was evaluated with BioAnalyser 2100 (Agilent). Libraries were sequenced by the Genomic Resources Core Facility at Weill Cornell Medicine, obtaining 50-bp single-end or paired-end reads to a depth of ~15 - 20 million reads per sample. Read quality was assessed and adapters trimmed using fastp. Reads were then mapped to the human genome (hg38) using STAR aligner and reads in exons were counted against Gencode v37 with Featurecount. Differential gene expression analysis was performed in R with edgeR using quasi-likelihood framework. Only genes with expression levels exceeding 4 counts per million reads in at least one group were used for downstream analysis. Benjamini-Hochberg false discovery rate (FDR) procedure was used to correct for multiple testing. Genes were categorized as upregulated if log2FC ≥ 1 and FDR ≤ 0.05 threshold was satisfied, downregulated if log2FC ≤ -1 and FDR ≤ 0.05. Heatmaps were obtained using Morpheus web application and replotted using the R package pheatmap. To analyze the regulatory factors that involve in IFN-γ - (T+P) interactions we selected all transcription factors based on ref. (Lambert et al., 2018) from the interacting clusters (Figure 6) and used them to build protein associating network with StringDB (Szklarczyk et al., 2019) limiting the associations to the highest interaction score (0.9) and visualized with R *tidygraph* framework using circular layout. RNAseq experiments were performed using different blood donors: 1. Monocytes stimulated by vehicle control or T, P or TP for 3 or 24 hr (3 donors). 2. Monocytes primed with or without IFN-γ and then stimulated with vehicle control, T, P or TP for 3 hr (3 different donors). Aliquots of cells from this experiment were used for ATACseq.

ATAC sequencing and data analysis. One million monocytes were lysed using cold lysis buffer (10 mM Tris-HCl, pH 7.4, 10 mM NaCl, 3 mM MgCl2, and 0.1% IGEPAL CA-630), and nuclei were immediately pelleted at 500xg for 10 min in a refrigerated centrifuge. The pellet was resuspended in a transposase reaction mix consisting of 25 µl 2× TD buffer, 2.5 µl transposase (Illumina), and 22.5 µl nuclease-free water. The transposition reaction was carried out for 30 min at 37°C. Following transposition, the sample was purified using a MinElute PCR Purification kit. Library fragments were amplified using 1× NEB next PCR master mix and 1.25 M custom Nextera PCR primers, with subsequent purification using a Qiagen PCR cleanup kit, yielding a final library concentration of ~30 nM in 20 µl. Libraries were amplified for a total of 10–13 cycles and subjected to high-throughput sequencing at the Genomic Resources Core Facility at Weill Cornell Medicine with 50-bp paired-end reads. Data on ATAC-seq experiments were derived from three independent experiments with different blood donors.

For ATACseq data analysis, read alignments were performed against the GRCh38/hg38 reference human genome. Peak calling was conducted using MACS2 with the following parameters: "macs2 callpeak -f BAMPE -g hs -q 0.01 --nomodel --shift 37 --extsize 76 ". A master consensus peak set was generated by merging the resulting peak files for each treatment condition, followed by merging peaks within 50bp of each other. Quantification of peaks to compare global ATACseq signal changes in the BAM files was conducted using the NCBI/BAMscale program. Raw count matrices were obtained utilizing the BAMscale program. Subsequent analysis utilized the HSS Genomic Center Bioinformatic Core's ATACseq analysis

pipeline (https://gitlab.com/hssgenomics/Shiny-ATAC) for peak filtering, annotation relative to genomic features, differential peak analysis, and enrichment of signal around specific motifs using ChromVAR (Schep et al., 2017). Footprint analyses were performed using TOBIAS (Bentsen et al., 2020) according to the user manual. Briefly, we filtered JASPAR2022-CORE_vertebrates transcription factor list based on the TF expressed in our RNAseq data and used this as TF input for motif footprinting. De novo transcription factor motif enrichment analysis was carried out using the motif finder program findMotifsGenome in the HOMER package, focusing on the given peaks. Peak sequences were compared to random genomic fragments of the same size and normalized G+C content to identify enriched motifs in the targeted sequences.

ICI-arthritis sample preparation for single cell RNA sequencing.
All research using patient samples at the Hospital for Special Surgery adhered to approved protocols (IRB protocols 2017-1898 and 2017-1871) with informed consent as required; samples investigated in this study correspond to a subset of those subjected to a distinct analysis in ref. (Wang et al., 2023). Synovial fluids were collected from ICI-arthritis as discarded fluid during patient visits. Tissue samples were collected during arthroplasty surgery. Mononuclear cells were isolated from synovial fluid by density centrifugation using Ficoll-Paque Plus (GE Healthcare), preserved in Cryostor CS10 (Stemcell Technologies), and placed in liquid nitrogen for long-term storage. Tissues received post-surgery were processed into fragments of 2-3 mm^3, preserved in Cryostor CS10, and placed in liquid nitrogen for long-term storage.

For sample preparation for single-cell RNA sequencing, tissues were thawed into pre-warmed media at 37ºC. The media composition was RPMI 1640 (Corning) containing 10% defined FBS (Cytiva) and 1% of 200mM L-Glutamine (Gibco). The tissues were then washed twice in RPMI alone for five minutes and chopped into fine pieces using a blade. Finely cut tissue was subjected to 30 minutes in a digestion buffer composed of RPMI, Liberase TL (100µg/ml; Roche) and DNaseI (100µg/ml; Roche). The fragments were digested in 5 ml polystyrene tubes (5ml/sample of digestion buffer), securely placed in a MACSmix tube rotator (Miltenyi Biotec) in an incubator at 37ºC with 5%CO2. The tissue preparation was filtered through a 70µM cell strainer (BD) and mashed using the back of a syringe plunger. The eluate containing the cells was washed with media and centrifuged at 1500rpm for 4 minutes at 4ºC. The pellet was resuspended in media and filtered again through a 40µM cell strainer (BD). Cells in the eluate were then counted and used in downstream experiments. Fluid samples were thawed from cold storage into pre-warmed media at 37ºC and then processed similarly to tissue samples. Live mononuclear cells were then FACS-sorted from five synovial fluid and two synovial tissue samples. Cells were sorted on a three-laser BD FACS Aria Fusion cell sorter at the Flow Cytometry Core Facility at Weill Cornell Medicine (WCM). Intact cells were gated according to forward scatter and side scatter area (FSC-A and SSC-A). Doublets were excluded by serial FSC-H/FSC-W and SSC-H/SSC-W gates (H, height; W, width). Non-viable cells were excluded based on DAPI uptake, sorted through a 70µM nozzle at 70 523 psi. Flow cytometric quantification of cell populations was performed using FlowJo v.10.0.7.

ICI-arthritis single cell RNA sequencing
3' gene expression (GEX) libraries of live synovial mononuclear cells were prepared using Chromium Single Cell 3' v3 Kit reagents and protocols provided by 10X Genomics. The pooled libraries at 10nM concentration were sequenced using NovaSeq6000 S2 Flow Cell using the

Illumina platform at Genomics Research Core Facility at WCM. 10x FASTQ files were processed with the Cellranger count 4.0 pipeline with default parameters. Reads were aligned to the human reference sequence GRCh38. Seurat package (v.4.0.0) was used to perform unbiased clustering. The 3' GEX dataset of sorted live synovial cells had QC performed to also remove cells with less than 150 genes, more than 7500 genes, or > 25% mitochondrial gene expression, resulting in a total of 44,959 cells and 26,424 genes. This dataset was then log-normalized using a scale factor of 10,000. Potential confounders such as percent mitochondrial gene expression and number of UMI per cell were regressed out during scaling (mean of 0 and variance of 1) for 3' GEX dataset.

Principal component analysis was used with the top 2000 highly variable genes. Elbow plot was used to determine the statistically significant principal components of 17 PCs for follow-up analysis. Harmony (v1.0) was performed to improve integration and correct for batch effects on our samples, with parameters of max.iter.cluster = 30, and max.iter.harmony = 20 and sample as the only covariate. Eleven clusters for the 3' dataset at 0.2 resolution were found, and their identity was annotated based on the expression of differentially expressed genes (DEG) using FindAllMarkers function using default parameters. Additional grouping of the 3' GEX dataset of select clusters at resolution 0.2 was used to define only monocytes and macrophages to create a new data object. These clusters contained 14,110 cells after going through linear dimensional reduction of the top 5 PCs with Harmony as described above. Eight clusters were defined at 0.3 resolution for the final analysis.

Statistical Analysis. Graphpad Prism was used for all statistical analysis. Information about the specific tests used, and number of independent experiments is provided in the figure legends.

Data and code availability.
Sequencing data from this study have been deposited at GEO (RNAseq: GSE272019; ATACseq: GSE272017) and will be publicly available from the date of publication. The scRNAseq data object is available on Gitub.

Acknowledgments
We thank the Weill Cornell Medicine Genomics Core for sequencing and the Computational Biology Core of the David Z. Rosensweig Genomics Center at HSS for data analysis.

AUTHOR CONTRIBUTIONS
U.S. performed most of the experiments, B.M., R.Y., and Y.C. performed bioinformatic analysis, A.S., K.C., A.B. and L.D. contributed the ICI-arthritis experiments, and L.D. provided intellectual input. L.I. conceptualized and oversaw the study. All authors wrote and then reviewed and provided input on the manuscript.

DECLARATION OF INTERESTS
The authors declare no relevant conflicts of interest.

References

Alivernini, S., L. MacDonald, A. Elmesmari, S. Finlay, B. Tolusso, M.R. Gigante, L. Petricca, C. Di Mario, L. Bui, S. Perniola, M. Attar, M. Gessi, A.L. Fedele, S. Chilaka, D. Somma, S.N. Sansom, A. Filer, C. McSharry, N.L. Millar, K. Kirschner, A. Nerviani, M.J. Lewis, C. Pitzalis, A.R. Clark, G. Ferraccioli, I. Udalova, C.D. Buckley, E. Gremese, I.B. McInnes, T.D. Otto, and M. Kurowska-Stolarska. 2020. Distinct synovial tissue macrophage subsets regulate inflammation and remission in rheumatoid arthritis. *Nature medicine* 26:1295-1306.

Altarejos, J.Y., and M. Montminy. 2011. CREB and the CRTC co-activators: sensors for hormonal and metabolic signals. *Nature reviews. Molecular cell biology* 12:141-151.

Bayerl, F., P. Meiser, S. Donakonda, A. Hirschberger, S.B. Lacher, A.M. Pedde, C.D. Hermann, A. Elewaut, M. Knolle, L. Ramsauer, T.J. Rudolph, S. Grassmann, R. Öllinger, N. Kirchhammer, M. Trefny, M. Anton, D. Wohlleber, B. Höchst, A. Zaremba, A. Krüger, R. Rad, A.C. Obenauf, D. Schadendorf, A. Zippelius, V.R. Buchholz, B.U. Schraml, and J.P. Böttcher. 2023. Tumor-derived prostaglandin E2 programs cDC1 dysfunction to impair intratumoral orchestration of anti-cancer T cell responses. *Immunity* 56:1341-1358.e1311.

Bentsen, M., P. Goymann, H. Schultheis, K. Klee, A. Petrova, R. Wiegandt, A. Fust, J. Preussner, C. Kuenne, T. Braun, J. Kim, and M. Looso. 2020. ATAC-seq footprinting unravels kinetics of transcription factor binding during zygotic genome activation. *Nature communications* 11:4267.

Braaten, T.J., J.R. Brahmer, P.M. Forde, D. Le, E.J. Lipson, J. Naidoo, M. Schollenberger, L. Zheng, C.O. Bingham, A.A. Shah, and L.C. Cappelli. 2020. Immune checkpoint inhibitor-induced inflammatory arthritis persists after immunotherapy cessation. *Annals of the rheumatic diseases* 79:332-338.

Cappelli, L.C., M.A. Thomas, C.O. Bingham, 3rd, A.A. Shah, and E. Darrah. 2020. Immune checkpoint inhibitor-induced inflammatory arthritis as a model of autoimmune arthritis. *Immunological reviews* 294:106-123.

Caronni, N., F. La Terza, F.M. Vittoria, G. Barbiera, L. Mezzanzanica, V. Cuzzola, S. Barresi, M. Pellegatta, P. Canevazzi, G. Dunsmore, C. Leonardi, E. Montaldo, E. Lusito, E. Dugnani, A. Citro, M.S.F. Ng, M. Schiavo Lena, D. Drago, A. Andolfo, S. Brugiapaglia, A. Scagliotti, A. Mortellaro, V. Corbo, Z. Liu, A. Mondino, P. Dellabona, L. Piemonti, C. Taveggia, C. Doglioni, P. Cappello, F. Novelli, M. Iannacone, L.G. Ng, F. Ginhoux, S. Crippa, M. Falconi, C. Bonini, L. Naldini, M. Genua, and R. Ostuni. 2023. IL-1β(+) macrophages fuel pathogenic inflammation in pancreatic cancer. *Nature* 623:415-422.

Chang, Y.K., Z. Zuo, and G.D. Stormo. 2018. Quantitative profiling of BATF family proteins/JUNB/IRF hetero-trimers using Spec-seq. *BMC molecular biology* 19:5.

Chen, X., I. Barozzi, A. Termanini, E. Prosperini, A. Recchiuti, J. Dalli, F. Mietton, G. Matteoli, S. Hiebert, and G. Natoli. 2012. Requirement for the histone deacetylase Hdac3 for the inflammatory gene expression program in macrophages. *Proceedings of the National Academy of Sciences of the United States of America* 109:E2865-2874.

Cilenti, F., G. Barbiera, N. Caronni, D. Iodice, E. Montaldo, S. Barresi, E. Lusito, V. Cuzzola, F.M. Vittoria, L. Mezzanzanica, P. Miotto, P. Di Lucia, D. Lazarevic, D.M. Cirillo, M. Iannacone, M. Genua, and R. Ostuni. 2021. A PGE(2)-MEF2A axis enables context-dependent control of inflammatory gene expression. *Immunity* 54:1665-1682.e1614.

Donlin, L.T., A. Jayatilleke, E.G. Giannopoulou, G.D. Kalliolias, and L.B. Ivashkiv. 2014. Modulation of TNF-induced macrophage polarization by synovial fibroblasts. *Journal of immunology (Baltimore, Md. : 1950)* 193:2373-2383.

Dougan, M., A.M. Luoma, S.K. Dougan, and K.W. Wucherpfennig. 2021. Understanding and treating the inflammatory adverse events of cancer immunotherapy. *Cell* 184:1575-1588.

Gerlo, S., R. Kooijman, I.M. Beck, K. Kolmus, A. Spooren, and G. Haegeman. 2011. Cyclic AMP: a selective modulator of NF-kappaB action. *Cellular and molecular life sciences : CMLS* 68:3823-3841.

Gong, Z., Q. Li, J. Shi, P. Li, L. Hua, L.D. Shultz, and G. Ren. 2023. Immunosuppressive reprogramming of neutrophils by lung mesenchymal cells promotes breast cancer metastasis. *Science immunology* 8:eadd5204.

Gravallese, E.M., and G.S. Firestein. 2023. Rheumatoid Arthritis - Common Origins, Divergent Mechanisms. *The New England journal of medicine* 388:529-542.

Ivashkiv, L.B. 2018. IFNgamma: signalling, epigenetics and roles in immunity, metabolism, disease and cancer immunotherapy. *Nature reviews. Immunology* 18:545-558.

Kalliolias, G.D., and L.B. Ivashkiv. 2016. TNF biology, pathogenic mechanisms and emerging therapeutic strategies. *Nature reviews. Rheumatology* 12:49-62.

Kawabata, A. 2011. Prostaglandin E2 and pain--an update. *Biological & pharmaceutical bulletin* 34:1170-1173.

Kawahara, K., H. Hohjoh, T. Inazumi, S. Tsuchiya, and Y. Sugimoto. 2015. Prostaglandin E2-induced inflammation: Relevance of prostaglandin E receptors. *Biochimica et biophysica acta* 1851:414-421.

Kuo, D., J. Ding, I.S. Cohn, F. Zhang, K. Wei, D.A. Rao, C. Rozo, U.K. Sokhi, S. Shanaj, D.J. Oliver, A.P. Echeverria, E.F. DiCarlo, M.B. Brenner, V.P. Bykerk, S.M. Goodman, S. Raychaudhuri, G. Ratsch, L.B. Ivashkiv, and L.T. Donlin. 2019. HBEGF(+) macrophages in rheumatoid arthritis induce fibroblast invasiveness. *Science translational medicine* 11:

Kusnadi, A., S.H. Park, R. Yuan, T. Pannellini, E. Giannopoulou, D. Oliver, T. Lu, K.H. Park-Min, and L.B. Ivashkiv. 2019. The Cytokine TNF Promotes Transcription Factor SREBP Activity and Binding to Inflammatory Genes to Activate Macrophages and Limit Tissue Repair. *Immunity* 51:241-257.e249.

Lambert, S.A., A. Jolma, L.F. Campitelli, P.K. Das, Y. Yin, M. Albu, X. Chen, J. Taipale, T.R. Hughes, and M.T. Weirauch. 2018. The Human Transcription Factors. *Cell* 172:650-665.

Lewis, M.J., M.R. Barnes, K. Blighe, K. Goldmann, S. Rana, J.A. Hackney, N. Ramamoorthi, C.R. John, D.S. Watson, S.K. Kummerfeld, R. Hands, S. Riahi, V. Rocher-Ros, F. Rivellese, F. Humby, S. Kelly, M. Bombardieri, N. Ng, M. DiCicco, D. van der Heijde, R. Landewe, A. van der Helm-van Mil, A. Cauli, I.B. McInnes, C.D. Buckley, E. Choy, P.C. Taylor, M.J. Townsend, and C. Pitzalis. 2019. Molecular Portraits of Early Rheumatoid Arthritis Identify Clinical and Treatment Response Phenotypes. *Cell reports* 28:2455-2470.e2455.

Li, H., J. Zuo, and W. Tang. 2018. Phosphodiesterase-4 Inhibitors for the Treatment of Inflammatory Diseases. *Frontiers in pharmacology* 9:1048.

Luan, B., M.O. Goodarzi, N.G. Phillips, X. Guo, Y.D. Chen, J. Yao, M. Allison, J.I. Rotter, R. Shaw, and M. Montminy. 2014. Leptin-mediated increases in catecholamine signaling reduce adipose tissue inflammation via activation of macrophage HDAC4. *Cell metabolism* 19:1058-1065.

MacKenzie, K.F., K. Clark, S. Naqvi, V.A. McGuire, G. Noehren, Y. Kristariyanto, M. van den Bosch, M. Mudaliar, P.C. McCarthy, M.J. Pattison, P.G. Pedrioli, G.J. Barton, R. Toth, A. Prescott, and J.S. Arthur. 2013. PGE(2) induces macrophage IL-10 production and a regulatory-like phenotype via a protein kinase A-SIK-CRTC3 pathway. *Journal of immunology (Baltimore, Md. : 1950)* 190:565-577.

Mandelin, A.M., 2nd, P.J. Homan, A.M. Shaffer, C.M. Cuda, S.T. Dominguez, E. Bacalao, M. Carns, M. Hinchcliff, J. Lee, K. Aren, A. Thakrar, A.B. Montgomery, S.L. Bridges, Jr., J.M. Bathon, J.P. Atkinson, D.A. Fox, E.L. Matteson, C.D. Buckley, C. Pitzalis, D. Parks, L.B. Hughes, L. Geraldino-Pardilla, R. Ike, K. Phillips, K. Wright, A. Filer, S. Kelly, E.M.

Ruderman, V. Morgan, H. Abdala-Valencia, A.V. Misharin, G.S. Budinger, E.T. Bartom, R.M. Pope, H. Perlman, and D.R. Winter. 2018. Transcriptional Profiling of Synovial Macrophages Using Minimally Invasive Ultrasound-Guided Synovial Biopsies in Rheumatoid Arthritis. *Arthritis & rheumatology (Hoboken, N.J.)* 70:841-854.

McInnes, I.B., and G. Schett. 2011. The pathogenesis of rheumatoid arthritis. *The New England journal of medicine* 365:2205-2219.

Mishra, B., and L.B. Ivashkiv. 2024. Interferons and epigenetic mechanisms in training, priming and tolerance of monocytes and hematopoietic progenitors. *Immunological reviews* 323:257-275.

Orange, D.E., P. Agius, E.F. DiCarlo, N. Robine, H. Geiger, J. Szymonifka, M. McNamara, R. Cummings, K.M. Andersen, S. Mirza, M. Figgie, L.B. Ivashkiv, A.B. Pernis, C.S. Jiang, M.O. Frank, R.B. Darnell, N. Lingampali, W.H. Robinson, E. Gravallese, V.P. Bykerk, S.M. Goodman, and L.T. Donlin. 2018. Identification of Three Rheumatoid Arthritis Disease Subtypes by Machine Learning Integration of Synovial Histologic Features and RNA Sequencing Data. *Arthritis & rheumatology (Hoboken, N.J.)* 70:690-701.

Perera, J., C.A. Delrosso, A. Nerviani, and C. Pitzalis. 2024. Clinical Phenotypes, Serological Biomarkers, and Synovial Features Defining Seropositive and Seronegative Rheumatoid Arthritis: A Literature Review. *Cells* 13:

Perkins, D.J., K. Richard, A.M. Hansen, W. Lai, S. Nallar, B. Koller, and S.N. Vogel. 2018. Autocrine-paracrine prostaglandin E(2) signaling restricts TLR4 internalization and TRIF signaling. *Nature immunology* 19:1309-1318.

Perretti, M., D. Cooper, J. Dalli, and L.V. Norling. 2017. Immune resolution mechanisms in inflammatory arthritis. *Nature reviews. Rheumatology* 13:87-99.

Qiao, Y., E.G. Giannopoulou, C.H. Chan, S.H. Park, S. Gong, J. Chen, X. Hu, O. Elemento, and L.B. Ivashkiv. 2013. Synergistic activation of inflammatory cytokine genes by interferon-gamma-induced chromatin remodeling and toll-like receptor signaling. *Immunity* 39:454-469.

Rodriguez, M., E. Domingo, C. Municio, Y. Alvarez, E. Hugo, N. Fernandez, and M. Sanchez Crespo. 2014. Polarization of the innate immune response by prostaglandin E2: a puzzle of receptors and signals. *Molecular pharmacology* 85:187-197.

Schep, A.N., B. Wu, J.D. Buenrostro, and W.J. Greenleaf. 2017. chromVAR: inferring transcription-factor-associated accessibility from single-cell epigenomic data. *Nature methods* 14:975-978.

Smolen, J.S., D. Aletaha, and I.B. McInnes. 2016. Rheumatoid arthritis. *Lancet (London, England)* 388:2023-2038.

Sundberg, T.B., H.G. Choi, J.H. Song, C.N. Russell, M.M. Hussain, D.B. Graham, B. Khor, J. Gagnon, D.J. O'Connell, K. Narayan, V. Dancik, J.R. Perez, H.C. Reinecker, N.S. Gray, S.L. Schreiber, R.J. Xavier, and A.F. Shamji. 2014. Small-molecule screening identifies inhibition of salt-inducible kinases as a therapeutic strategy to enhance immunoregulatory functions of dendritic cells. *Proceedings of the National Academy of Sciences of the United States of America* 111:12468-12473.

Szklarczyk, D., A.L. Gable, D. Lyon, A. Junge, S. Wyder, J. Huerta-Cepas, M. Simonovic, N.T. Doncheva, J.H. Morris, P. Bork, L.J. Jensen, and C.V. Mering. 2019. STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic acids research* 47:D607-d613.

Theilgaard-Mönch, K., S. Pundhir, K. Reckzeh, J. Su, M. Tapia, B. Furtwängler, J. Jendholm, J.S. Jakobsen, M.S. Hasemann, K.J. Knudsen, J.B. Cowland, A. Fossum, E. Schoof, M.B. Schuster, and B.T. Porse. 2022. Transcription factor-driven coordination of cell cycle exit and lineage-specification in vivo during granulocytic differentiation : In memoriam Professor Niels Borregaard. *Nature communications* 13:3595.

Tsuge, K., T. Inazumi, A. Shimamoto, and Y. Sugimoto. 2019. Molecular mechanisms underlying prostaglandin E2-exacerbated inflammation and immune diseases. *International immunology* 31:597-606.

Wang, R., A. Singaraju, K.E. Marks, L. Shakib, G. Dunlap, I. Adejoorin, S.R. Greisen, L. Chen, A.K. Tirpack, C. Aude, M.R. Fein, D.J. Todd, L. MacFarlane, S.M. Goodman, E.F. DiCarlo, E.M. Massarotti, J.A. Sparks, A.H. Jonsson, M.B. Brenner, M.A. Postow, K.K. Chan, A.R. Bass, L.T. Donlin, and D.A. Rao. 2023. Clonally expanded CD38(hi) cytotoxic CD8 T cells define the T cell infiltrate in checkpoint inhibitor-associated arthritis. *Science immunology* 8:eadd1591.

Wei, K., I. Korsunsky, J.L. Marshall, A. Gao, G.F.M. Watts, T. Major, A.P. Croft, J. Watts, P.E. Blazar, J.K. Lange, T.S. Thornhill, A. Filer, K. Raza, L.T. Donlin, C.W. Siebel, C.D. Buckley, S. Raychaudhuri, and M.B. Brenner. 2020. Notch signalling drives synovial fibroblast identity and arthritis pathology. *Nature* 582:259-264.

Yarilina, A., K.H. Park-Min, T. Antoniv, X. Hu, and L.B. Ivashkiv. 2008. TNF activates an IRF1-dependent autocrine loop leading to sustained expression of chemokines and STAT1-dependent type I interferon-response genes. *Nature immunology* 9:378-387.

Yokoyama, U., K. Iwatsubo, M. Umemura, T. Fujita, and Y. Ishikawa. 2013. The prostanoid EP4 receptor and its signaling pathway. *Pharmacological reviews* 65:1010-1052.

Zack, S.R., O. Alzoubi, N. Satoeya, K.P. Singh, S. Deen, W. Nijim, M.J. Lewis, C. Pitzalis, N. Sweiss, L.B. Ivashkiv, and S. Shahrara. 2024. ANOTHER NOTCH IN THE BELT OF RHEUMATOID ARTHRITIS. *Arthritis & rheumatology (Hoboken, N.J.)*

Zec, K., B. Schonfeldova, Z. Ai, E. Van Grinsven, G. Pirgova, H.L. Eames, D.L. Berthold, M. Attar, E.B. Compeer, T.I. Arnon, and I.A. Udalova. 2023. Macrophages in the synovial lining niche initiate neutrophil recruitment and articular inflammation. *The Journal of experimental medicine* 220:

Zhang, F., A.H. Jonsson, A. Nathan, N. Millard, M. Curtis, Q. Xiao, M. Gutierrez-Arcelus, W. Apruzzese, G.F.M. Watts, D. Weisenfeld, S. Nayar, J. Rangel-Moreno, N. Meednu, K.E. Marks, I. Mantel, J.B. Kang, L. Rumker, J. Mears, K. Slowikowski, K. Weinand, D.E. Orange, L. Geraldino-Pardilla, K.D. Deane, D. Tabechian, A. Ceponis, G.S. Firestein, M. Maybury, I. Sahbudin, A. Ben-Artzi, A.M. Mandelin, 2nd, A. Nerviani, M.J. Lewis, F. Rivellese, C. Pitzalis, L.B. Hughes, D. Horowitz, E. DiCarlo, E.M. Gravallese, B.F. Boyce, L.W. Moreland, S.M. Goodman, H. Perlman, V.M. Holers, K.P. Liao, A. Filer, V.P. Bykerk, K. Wei, D.A. Rao, L.T. Donlin, J.H. Anolik, M.B. Brenner, and S. Raychaudhuri. 2023. Deconstruction of rheumatoid arthritis synovium defines inflammatory subtypes. *Nature* 623:616-624.

Zhang, F., J.R. Mears, L. Shakib, J.I. Beynor, S. Shanaj, I. Korsunsky, A. Nathan, L.T. Donlin, and S. Raychaudhuri. 2021. IFN-γ and TNF-α drive a CXCL10+ CCL2+ macrophage phenotype expanded in severe COVID-19 lungs and inflammatory diseases with tissue inflammation. *Genome medicine* 13:64.

Zhang, F., K. Wei, K. Slowikowski, C.Y. Fonseka, D.A. Rao, S. Kelly, S.M. Goodman, D. Tabechian, L.B. Hughes, K. Salomon-Escoto, G.F.M. Watts, A.H. Jonsson, J. Rangel-Moreno, N. Meednu, C. Rozo, W. Apruzzese, T.M. Eisenhaure, D.J. Lieb, D.L. Boyle, A.M. Mandelin, 2nd, B.F. Boyce, E. DiCarlo, E.M. Gravallese, P.K. Gregersen, L. Moreland, G.S. Firestein, N. Hacohen, C. Nusbaum, J.A. Lederer, H. Perlman, C. Pitzalis, A. Filer, V.M. Holers, V.P. Bykerk, L.T. Donlin, J.H. Anolik, M.B. Brenner, and S. Raychaudhuri. 2019. Defining inflammatory cell states in rheumatoid arthritis joint synovial tissues by integrating single-cell transcriptomics and mass cytometry. *Nature immunology* 20:928-942.

Zhang, X., D.T. Odom, S.H. Koo, M.D. Conkright, G. Canettieri, J. Best, H. Chen, R. Jenner, E. Herbolsheimer, E. Jacobsen, S. Kadam, J.R. Ecker, B. Emerson, J.B. Hogenesch, T. Unterman, R.A. Young, and M. Montminy. 2005. Genome-wide analysis of cAMP-

response element binding protein occupancy, phosphorylation, and target gene activation in human tissues. *Proceedings of the National Academy of Sciences of the United States of America* 102:4459-4464.

Zhou, L., W. Zhou, A.M. Joseph, C. Chu, G.G. Putzel, B. Fang, F. Teng, M. Lyu, H. Yano, K.I. Andreasson, E. Mekada, G. Eberl, and G.F. Sonnenberg. 2022. Group 3 innate lymphoid cells produce the growth factor HB-EGF to protect the intestine from TNF-mediated inflammation. *Nature immunology* 23:251-261.

Zhou, Z., X. Zhou, X. Jiang, B. Yang, X. Lu, Y. Fei, L. Zhao, H. Chen, L. Zhang, X. Si, N. Liang, Y. Wang, D. Yang, Y. Peng, Y. Yang, Z. Yao, Y. He, X. Wu, W. Zhang, M. Wang, H. Yang, and X. Zhang. 2024. Single-cell profiling identifies IL1B(hi) macrophages associated with inflammation in PD-1 inhibitor-induced inflammatory arthritis. *Nature communications* 15:2107.

Figure Legends

Figure 1. Cooperative induction of a distinct subset of inflammatory genes by PGE2 and TNF. (A) Experimental design. Primary human monocytes were stimulated with PGE2 (280 nM) and/or TNF (20 ng/ml) and harvested 3 or 24 hr after stimulation for RNAseq analysis. n = 3 independent blood donors. (B) Gene set enrichment analysis of genes induced > 2-fold by PGE2 + TNF (TP) (FDR < 0.05). (C) K-means clustering of differentially upregulated genes in any pairwise comparison relative to resting control (> 2-fold induction, FDR < 0.05). 3 hr time point. k = 5. (D) Pathway analysis of gene clusters in panel C. (E) qPCR analysis of gene expression in an additional 5 blood donors. Mean +/- SEM. Statistical significance was assessed using one-way ANOVA and Sidak's test for multiple comparisons (*p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001). (F). Western blot analysis. Representative blot out of 3 independent experiments.

Supplementary Figure 1. Regulation of gene expression by PGE2 and TNF in mouse bone marrow-derived macrophages. BMDMs were stimulated with PGE2 (280 nM) and/or mTNF (20 ng/ml) and harvested 3 or 24 hr after stimulation for qPCR analysis of gene expression, n = 3. Mean +/- SEM. Statistical significance was assessed using 2-way ANOVA and Tukey's test for multiple comparisons (*p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001).

Supplementary Figure 2. Analysis of differentially regulated genes at the 24 hr time point. Monocytes were stimulated for 24 hr as described in legend to Fig. 1A. K means clustering (k = 5, genes with > 2-fold induction, FDR < 0.05) (left panel). Pathway analysis of gene clusters (right panel). n = 3.

Figure 2. PGE2 and TNF costimulation model aspects of the RA and ICI-arthritis synovial macrophage phenotype. (A) TP costimulation recapitulates expression of 61% of the genes whose induction by TNF was augmented by coculture with synovial fibroblasts (also termed fibroblast-like synoviocytes, FLS). FLS-augmented TNF-inducible genes from data in refs. (Donlin et al., 2014; Kuo et al., 2019) were compared to the TP-induced genes at the 24 hr time point in Supplementary Fig. 2, fold-change >2, FDR < 0.05. The green area indicates extent of overlap, hypergeometric p = 8.36e-128. (B) Recapitulation of the RA synovial macrophage cluster 1 phenotype by TP-costimulated genes. The defining 128 genes of the C1 phenotype were overlapped with TP-costimulated genes and TNF/FLS-costimulated genes. TP-costimulated genes = orange + yellow = 52% of C1 defining genes (p < 10e-9 by Monte Carlo simulation). TNF/FLS-costimulated genes = orange + red = 34% of C1 defining genes. (C) Heat maps depicting expression of genes in pathogenic pathways, based on RNAseq data shown in Fig. 1. Blue font = genes expressed in C1 RA macrophages. (D) Heat map depicting regulation of representative genes that are expressed in RA C1 macrophages by P, T or TP. (E) UMAP visualization of monocyte and macrophage clusters based on scRNAseq of 14,110 macrophages and monocytes from 5 synovial fluids and 2 synovial tissues of ICI-arthritis patients. (F) Heatmap showing expression of key genes for the eight clusters identified in panel E. TNF+PGE2 signature genes are shown in red.

Figure 3. cAMP signaling has dichotomous suppressive and augmenting effects on the TNF-induced inflammatory response. (A) RT-qPCR analysis of gene expression in primary human monocytes stimulated with TNF and selective agonists of PGE2 receptors EP2 and EP4

that signal predominantly via cAMP. EP2 agonist = butaprost (10 μM); EP4 agonist = CAY10598 (10 μM). n = 3 (B) RT-qPCR analysis of gene expression in primary human monocytes stimulated with TNF and increasing concentrations of cAMP analog dibutyryl cAMP (10 and 100 μM; labeled A). n = 3. Mean +/- SEM. Statistical significance was assessed using 2 way ANOVA and Sidak's test for multiple comparisons (*p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001).

Figure 4. PGE2 effects on TNF-induced changes in chromatin accessibility. (A-E). Analysis of ATACseq data obtained using monocytes from 3 independent donors. (A) UPSET plot of differentially upregulated ATACseq peaks in any pairwise comparison relative to resting control (> 2-fold induction, FDR < 0.05). (B) De novo motif analysis using HOMER of ATACseq peaks induced by TNF (left panel), PGE2 (middle panel) or uniquely induced only under conditions of TP costimulation (right panel). (C) Upper. Violin plots showing normalized counts of ATACseq peaks induced by both PGE2 and TNF. ****p < 0.0001 by Wilcoxon rank sum test with Holm's correction for multiple comparisons. Lower. HOMER de novo motif analysis of the peaks in the upper panel. (D) Heatmap of the differential TF activity scores derived from ChromVAR analysis of ATACseq data for P, T or TP treated monocytes, compared to resting control. (E) Volcano plot of differential binding analysis of ATACseq peaks between the TP and T conditions using TOBIAS.

Supplementary Figure 3. ATACseq analysis of PGE2 and TNF stimulated monocytes. (A-F) Additional analysis of the ATACseq data obtained using monocytes from 3 independent donors. (A) Principal component plot. (B-E) Pathway analysis of genes associated with ATACseq peaks induced by TNF (B), PGE2 (C), co-induced by TP (D) or uniquely induced by TP (E) Volcano plots of differential binding analysis of ATACseq peaks using TOBIAS.

Figure 5. IFN-γ opposes the effects of PGE2 on TNF-induced gene expression. (A, C) RT-qPCR analysis of gene expression in primary human monocytes that were primed overnight with IFN-γ (100 U/ml) and then stimulated for 3 hr with P, T or TP as in Fig. 1. n = 3. Mean +/-SEM. Statistical significance was assessed using 2 way ANOVA with Sidak's test for multiple comparisons (*p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001). (B, D) Gene groups defined in Supplementary Fig. 4A based on pattern of expression in RNAseq data (n = 3) were subjected to hierarchical clustering.

Supplementary Figure 4. IFN-γ opposes the effects of PGE2 on TNF-induced gene expression. (A) RNAseq data, n = 3. K-means clustering of differentially upregulated genes in any pairwise comparison relative to resting control (> 1.25-fold induction, FDR < 0.01). 3 hr time point. k = 6; 3 gene sets (Groups 4-6) that are costimulated by T and P are apparent. (B) Gene groups 4 and 5 defined in panel A were subjected to hierarchical clustering.

Figure 6. Interaction analysis of IFN-γ and the TP response. (A) Differentially expressed genes in ((IFN-γ + TP) – IFN-γ)) – (IFN-γ – Resting) contrast define statistical interactions between IFN-γ and TP treatments using the RNAseq data, n = 3, FDR < 0.05, fold change > 2. Hierarchical clustering of z-transformed gene expression values (cpm) reveals 6 interaction clusters (right). Violin plots showing relative gene expression between resting (R), TP, IFN-γ alone (I) and IFN-γ + TP (ITP) conditions (second from left). Interaction plot (left). (B) Pathway analysis of the genes in the interaction clusters defined in panel A. (C) STRING functional

protein association network of transcription factors from each cluster (fold change > 4, FDR < 0.05). Lines designate functional interactions between individual TFs. The size of nodes is proportional to the number of STRINGDB interactions.

Figure 7. IFN-γ inhibits PGE2-induced gene expression and chromatin accessibility. (A-B, F) Analysis of RNAseq data obtained using monocytes from 3 independent donors. (A) K-means clustering of differentially upregulated genes in any pairwise comparison relative to resting control (> 2-fold induction, FDR < 0.05). k = 5. (B) Heat maps depicting expression of genes in pathogenic pathways and expressed in C1 RA macrophages as defined in Fig. 2C and 2D. (C-E, G, H). Analysis of ATACseq data obtained using monocytes from 3 independent donors. (C) UPSET plot of differentially upregulated ATACseq peaks in any pairwise comparison relative to resting control (> 2-fold induction, FDR < 0.05). (D) De novo motif analysis using HOMER of ATACseq peaks induced uniquely by PGE2 (corresponding to G2 in panel C). (E) Heatmap of the differential TF activity scores derived from ChromVAR analysis of ATACseq data for P, IFN-γ or IFN-γ + PGE2 treated monocytes, compared to resting control. (F) Heat map depicting expression of CEBP genes in RNAseq data. (G, H) Volcano plots of differential binding analysis of ATACseq peaks of IFN-γ versus resting control (G) and IFN-γ + PGE2 versus PGE2 (H) conditions using TOBIAS. The IFN-γ versus resting results (G) reproduce results in (Mishra and Ivashkiv, 2024) that were obtained in independent experiments with different blood donors.

Supplementary Figure 5. (A) PCA plot of RNAseq data shown in Fig. 7. (B) Pathway analysis of gene clusters identified in Fig. 7A. (C). PCA plot of ATACseq data shown in Fig. 7.

Figure 8. Crosstalk between PGE2 and TNF signaling and its regulation by IFN-γ. (A) PGE2 costimulates TNF-induced expression of select inflammatory genes such as IL-1 and Notch pathway genes and neutrophil chemokines by inducing transcription factors including CEBP, AP-1 and NR4A1/2 that cooperate with TNF-activated NF-κB (left). IFN-γ induces IRF transcription factors that cooperate with TNF-activated NF-κB to costimulate distinct inflammatory genes such as *TNF* and T cell chemokines such as CXCL10, reviewed in ref. (Mishra and Ivashkiv, 2024) (right). IFN-γ inhibits induction of TP-costimulated genes by suppressing induction of AP-1 and NR4A1/2, and altering the pattern of expression of CEBP factors. PGE2 suppresses TNF-mediated induction of TNF and ISGs. As PGE2 is produced by stromal cells and IFN-γ is produced by lymphocytes, neighboring cells in inflamed tissues will help determine the macrophage response to TNF. Induction of a subset of CEBP factors by IFN-γ is not depicted. (B) IFN-γ and PGE2 oppose each other to regulate the balance between distinct TNF-induced inflammatory responses. PGE2 signaling promotes a response that activates stromal cells via IL-1, EGFR ligands and Notch pathways, and promotes recruitment of neutrophils. IFN-γ suppresses these pathways and instead promotes inflammation via TNF and recruitment of T cells.

Sokhi et al., Figure 1

A

IL1B

IL1A

STAT4

CXCL1

CXCL10

B

IL1B

IL1A

STAT4

CXCL10

EREG

Sokhi et al., Figure 4

A

B

TNF

rank	p value	tgt num	bgd num	most similar motif
1	1e-1069	2210 (35.03%)	2494 (5.78%)	BATF
2	1e-719	1612 (25.55%)	1910 (4.43%)	NFkB-p65
3	1e-263	2461 (39.01%)	8611 (19.96%)	PB0058.1
4	1e-173	1406 (22.29%)	4376 (10.14%)	MF0006.1
5	1e-60	273 (4.33%)	574 (1.33%)	IRF1
6	1e-41	425 (6.74%)	1409 (3.27%)	RUNX1

PGE

rank	p value	tgt num	bgd num	most similar
1	1e-804	1482 (46.01%)	3139 (6.99%)	Fra1
2	1e-410	1153 (35.80%)	3753 (8.35%)	CEBPD
3	1e-107	930 (28.87%)	6238 (13.88%)	FOSL1::JUI
4	1e-100	900 (27.94%)	6095 (13.56%)	PB0058.1
5	1e-73	132 (4.10%)	220 (0.49%)	FOSL2
6	1e-44	162 (5.03%)	597 (1.33%)	NR4A1

T+P unique

rank	p value	tgt num	bgd num	most similar motif
1	1e-412	883 (36.73%)	2989 (6.42%)	JunB
2	1e-170	885 (36.81%)	6495 (13.95%)	PB0058.1
3	1e-79	299 (12.44%)	1590 (3.42%)	CEBPA
4	1e-40	180 (7.49%)	1076 (2.31%)	ISRE
5	1e-37	123 (5.12%)	577 (1.24%)	GABPA
6	1e-25	140 (5.82%)	976 (2.10%)	Atf1

C

rank	p value	tgt num	bgd num	most similar motif
1	1e-434	598 (56.47%)	2748 (5.88%)	Fra1
2	1e-70	513 (48.44%)	10881 (23.30%)	CEBPE
3	1e-18	89 (8.40%)	1303 (2.79%)	ZNF768
4	1e-18	56 (5.29%)	566 (1.21%)	Pax2
5	1e-17	290 (27.38%)	7823 (16.75%)	ETV1
6	1e-16	123 (11.61%)	2354 (5.04%)	NFkb-p65

D

Differential Motifs

E

TP vs T

A

IL1B

CXCL2

HEY1

STAT4

**

****

***

% GAPDH

+IFNγ

B

co-stimulated genes
Group 6

+IFNγ

NLRP3
PTGS2
CCL22

C

CXCL10

TNF

% GAPDH

*

+IFNγ

+IFNγ

D

TNF-induced and
suppressed by PGE2
Group 2

+IFNγ

TNF
CXCL9
CXCL10
CXCL11

A

B

MSigDB Hallmarks

ChEA2022

IC1
- Interferon Gamma Res
- Complement
- Apoptosis
- Inflammatory Respons
- Hypoxia

RELA 24523406
FOXO1 25302145
STAT4 19710469
STAT3 20064451
NR1H3 23393188
E2A 27217539

IC2
- Interferon Alpha Res
- Interferon Gamma Res
- Inflammatory Respons
- Apoptosis
- Hypoxia

SPI1 23127762
IRF8 27001747
IRF1 21803131
SOX2 20726797
ESR2 21235772

IC3
- Reactive Oxygen Spec
- p53 Pathway
- IL-6/JAK/STAT3 Signa
- Apoptosis
- KRAS Signaling Up
- Allograft Rejection

PPARG 20176806
SMRT 22465074
ENL 26711339
AF4 26711339
IRF8 27001747

IC4
- TNF-alpha Signaling
- Inflammatory Respons
- Interferon Gamma Res
- IL-6/JAK/STAT3 Signa
- KRAS Signaling Up

RELB 30642670
RELA 24523406
KDM2B 26808549
JUN 26020271
NR3C1 27076634
NFKB1 27076634

IC5
- p53 Pathway
- Hedgehog Signaling
- UV Response Up
- Estrogen Response Ea
- Apical Junction

ZNF217 24962896
NR3C1 27076634
Nerf2 26677805
FOXO1 32281255
FOXM1 26456572

IC6
- Myc Targets V2
- Unfolded Protein Res
- IL-2/STAT5 Signaling
- Myc Targets V1
- mTORC1 Signaling

MYC 28411283
MYC 22102868
MYC 18358816
MYC 19030024
MYCN 28898695

- logAdjusted P value

- logAdjusted P value

C

Sokhi et al, Figure 8A

24h Upregulated

24h Upregulated genes
(n=811)

Sokhi et al., Supplementary Figure 3

Sokhi et al., Supplementary Figure 4

Supplemental table: Clinical characteristics of ICB-arthritis patients

Pt ID	Age	Sex	Primary Malignancy	ICB-therapy	Arthritis Pattern	ICB Initiation to arthritis Onset (wks)	Arthritis Onset to SF Collection (wks)	ICB Initiation to SF Collection (wks)	Most Recent ICB to SF Collection (wks)	Duration of ICI-therapy	CDAI
HSS101	65	M	Melanoma	PD-1	Polyarthritis	4.3	147.6	151.9	106.6	46.3	34.0
HSS110	50	F	Melanoma	PD-1 + CTLA-4	Polyarthritis	52.3	173.9	226.2	185.0	42.2	9.0
HSS180	57	F	Renal cell carcinoma	PD-1 + CTLA-4	Monoarthritis	40.1	0.7	40.8	17.4	24.4	12.0
HSS200	62	F	Urothelial carcinoma	PD-1	Oligoarthritis	27.7	8.6	36.3	1.5	35.8	9.0
HSS219	55	M	Urothelial carcinoma	PD-1	Monoarthritis	269.7	21.9	291.6	32.6	260.0	7.0
RA335L and RA335R	60	F	Melanoma	PD-1	Polyarthritis	182.0	N/A	N/A	N/A	N/A	N/A