Protein-protein interaction is a major source of epistasis in genetic interaction

networks

Xavier Castellanos-Girouard[1,2], Adrian W.R. Serohijos[1,2], Stephen W. Michnick[1,2,*]

1. Department of Biochemistry, Université de Montréal; Montreal, Quebec, Canada

2. Robert-Cedergren Center for Bioinformatics and Genomics, Université de Montréal; Montreal,

Quebec, Canada

*Correspondence: Stephen.michnick@umontreal.ca

Abstract

Genetic interaction (GI) and protein-protein interaction (PPI) networks have proved among the

most important tools for inference of gene function and genotype-phenotype relationship[1–5]. It

remains unclear, however, how these two networks are related to each other. Here we demonstrate

that the strengths of epistatic genetic interactions between two genes strongly correlate with the

binding free energies of interactions between their cognate proteins, for both yeast and human

genomes. Consequently, we show that GI and PPI networks can reciprocally predict each other.

Further, we observe that functional divergence and redundancy in duplicated genes (paralogs) are

reflected in the binding affinity of their interactions with other proteins and in their genetic

interaction strengths. Finally, we demonstrate that the overall topologies of GI and PPI networks

significantly overlap in two topological features: Modules, in which genes/proteins are organized

by common function, and connectors, which link modules to each other. This opens avenues for

new integrated network approaches in understanding cellular processes and flow of genomic

information.

Introduction

Determining how genes cooperate to shape phenotype is a central goal in molecular biology. At the cellular level, our understanding of functional relationships between genes is best reflected in genetic interaction (GI) and protein-protein interaction (PPI) networks[1,3]. These two networks are highly complementary; GI networks link broad functional relationships between genes via their contribution to a phenotype, whereas PPI networks are maps of physical contacts between proteins. However, how these two networks are related to each other remains an open question[1,6].

GI networks are based on systematic or pooled pairwise deletion or disruption of genes and a genetic interaction is scored as effects on some trait, such as cell proliferation rates, that are less than or greater than the product of the effects of deleting the individual genes, a phenomenon called epistasis[7,8]. Epistasis is a major source of complexity in predicting genotype-phenotype relationships, making the elucidation of its molecular mechanism a significant endeavor[2,9–12]. GI networks also serve important applications such as identifying unique genetic vulnerabilities in cancer cells and inferring gene function by organizing genes into functional groups[1,13–15]. PPIs are also useful for inferring gene functions, linking proteins of unknown, to those of known function[3,16]. Finally, about two-thirds of mutations that cause Mendelian human diseases also disrupt PPIs, linking the integrity of PPI networks to disease[17]. Considering their central role in understanding normal and pathological cellular processes, modelling the relationship between GI and PPI networks has been a longstanding goal in systems biology[6,18].

Recent quantitative proteomic studies provide clues of a deeper physical relationship between PPI and GI networks[19–21]. The number of complexes formed between two proteins is determined by their abundances and their binding affinity. Hein *et al.* used mass spectrometry to measure the ratios of the abundances of the two binding partners (abundance stoichiometry) and of total numbers of protein pairs that form complexes (interaction stoichiometry), a number proportional to binding dissociation constants (K_d)[19]. The ranges of these binding and abundance stoichiometries were remarkably broad, between $\sim 10^{-7}$- 10^1 (Figure 1d, regions 2 and 4)[19]. Additionally, well annotated and known stable protein complexes were shown to be enriched around 1:1 abundance and interaction stoichiometries, suggesting a balance of the relative

concentration of proteins that form such stable PPIs. The strong negative epistasis that can occur

among genes that code for the subunits of essential and stable protein complexes raises the question

of whether the broad range of binding stoichiometries could be proportional to epistatic

strengths[1,6].

In addition, Cho *et al.* demonstrated that a quantitative measure of cellular spatial organization

similarity, measured by spinning disk confocal microscopy, of interacting proteins is proportional

to interaction stoichiometry[20]. These results imply that higher order physical organization of

proteins in the cell is also determined by binding strength. This relationship also defines

topological features of cellular PPI networks in which high interaction stoichiometry PPIs formed

modules of functionally related proteins while the low interaction stoichiometry pairs formed

transient interactions that connect modules[19,20]. GI networks also show a highly modular

organization and the correspondence of modular structures between PPI and GI networks has been

established[1,6,22]. Is it possible that there is a similar correspondence of PPI protein and GI pairs in

connector regions of PPI and GI networks?

Finally, it has been demonstrated that common patterns of epistasis between pairs of mutations in

subunits of protein complexes and a set of gene knockout strains correlate with the distance

between the mutated residues in protein complexes[23–25]. Consistent with the notion that mutations

of residues within functional sites, including hydrophobic core, allosteric, catalytic or binding

sites, should produce similar phenotypes and equally, be more strongly energetically coupled to

each other[26–28].

Following from these three observations, here, we test the hypotheses that the strengths of PPI and

GI correlate and that PPI networks and GI networks overlap in their topological features (modules

and connectors).

Results

Epistasis varies with protein abundance and PPI stoichiometries

Since a complete systematic GI network has been constructed for the budding yeast *Saccharomyces cerevisiae* (henceforth "yeast), we chose to make our initial comparison of PPI binding and epistatic strengths for this organism[1]. We first constructed a protein abundance versus PPI binding protein stoichiometry map based on data from a recent affinity purification-mass spectrometry and a meta-analysis of protein abundance measurements[19–21,29]. The distribution of yeast PPIs in the stoichiometry plot is qualitatively similar to that of human proteins (Figure 1b, c), the major difference being a higher frequency of low stoichiometry interactions for human proteins (Figure 1a, b). This higher frequency of weak interactions could reflect relaxation of selection pressure that occurred in the transition from unicellular to multicellular organisms as a consequence of reduced population, allowing for the increased rate of emergence of new genomic features, such as gene duplicates and accrual of new gene functions[30–32]. It could also reflect the increase in requirements of connector PPIs to coordinate activities of the expanded number of functional modules in metazoans. Furthermore, we also found that region 1 (Figure 1d) in the yeast stoichiometry plot is enriched for stable complexes (Supplementary Figure 1), consistent with findings for human proteins[19]. Notwithstanding the evolutionary distance between the two organisms, the similarity in their stoichiometry maps may reflect the conservation of core cellular processes, particularly the molecular mechanisms that mediate them, across yeast and humans[33].

We next sought to explore how GIs distribute across the PPI stoichiometry map. To this end, we merged the GI and PPI networks using the criterion that if the genes comprising a GI also code for proteins that interact, the GI score was assigned to that PPI. Yeast GIs were taken from a near-complete digenic knockout screen[1] and human GIs were taken from the largest double knockdown study in human cell lines (Jurkat and K562) to date[34]. Of note, the measured GI network in human cell lines is only comprised of 472 genes and consequently only about 0.05% of all possible double knockouts were measured, a much smaller sample than the data for yeast. By visualizing average GI strengths throughout the stoichiometry plots, we observed that negative GIs are strongest for PPIs near 1:1 interaction and abundance stoichiometries (Figure 1e, left panel). This observation

is consistent with previous analyses in yeast showing that negative GIs are enriched between genes

whose proteins assemble into complexes[1]. Some examples are highlighted in Table S1. This

pattern also holds in human cells (Figure 2f, left panel; Supplementary Figure 2a), but to a lesser

extent, likely due to the smaller sample size. In contrast, the strength of positive GIs seems to be

distributed more uniformly across the stoichiometry map. This suggests that positive GIs are not

predominantly driven by PPI and may be better explained in combination with other mechanisms.

Indeed, suppression mechanisms involving mRNA decay and protein degradation, as well as

specific patterns of disruption in metabolic pathways have been proposed as important sources of

positive GIs[1,35–37].

To further investigate the distribution of GI strength throughout the stoichiometric landscape, we

divided interactions into 4 regions previously outlined by Hein *et al.* (Figure 2c): region 1 consists

of interacting protein pairs of equal abundances and 1:1 binding ratios (enriched for stable

complexes); region 2 consists of protein pairs of equal abundances but low binding ratios (enriched

for transient interactions); region 3 consists of protein pairs of unequal abundances and 1:1 binding

ratios, and region 4 consists of protein pairs of unequal abundances and low binding ratios[19]. As

suggested by the stoichiometry plots, the distribution of negative GIs within region 1 is

significantly higher than that of other regions ($P < 0.01$, Wilcoxon rank-sum test) (Figure 1f, g,

left panels; Supplementary Figure 2c). The distribution of positive GI values is also highest in

region 1 in yeast and human cell lines (Figure 1f, g, right panels), in agreement with previous

observations that complexes, composed mostly of non-essential genes, are enriched for positive

GIs (Table S2)[1]. Interestingly, region 3 that is characterized by high interaction stoichiometry and

an excess of affinity purified *versus* bait protein show the lowest number of GIs among the four

regions (Figure 1g and h; Supplementary figure 2c), reflecting that the number of protein

complexes formed is not the sole determinant of GI strength. Taken together, these results show

that there is robust overlap between GI and PPI networks.

Figure 1. Strong negative GIs map to PPIs of equal protein stoichiometry. a, Genetic
interactions are generated by comparing fitness defects in genotypes with single or double gene
knockout/knockdowns, and protein-protein interactions are defined as physical contacts between
proteins. **b, c,** Protein-protein interactions (grey points) are plotted according to two
stoichiometries: Interaction Stoichiometry and Abundance Stoichiometry for yeast (b) and human
cell lines (c). **d,** The stoichiometry plot can be separated into 4 regions representing PPIs with
different ranges of protein stoichiometry values. **e (yeast), f (human),** Mean negative and positive
genetic interaction scores for PPIs throughout the stoichiometry plot. GI scores in human were
taken from a screen in K562 cell lines. **g (yeast), h (human),** Distribution of GI scores within the
4 defined regions of the stoichiometry plot. Significance levels are annotated as: * for $p \leq 0.05$, **
for $p \leq 0.01$, and *** for $p \leq 0.001$.

Negative epistasis correlates with PPI binding strength

Substantial efforts have been made to model how the physicochemical properties of proteins such as folding stability and PPI binding energies drive epistasis[28,38,39]. Whilst these approaches explain how epistasis arises from point mutations for single proteins or pairs of proteins in PPIs, the extent to which the strengths of GIs for whole gene knockout (or knock down) could be explained by binding free energies is unknown.

Interaction stoichiometry can provide an indirect measure of PPI strength if protein pairs are of equal abundance, however, interaction stoichiometry is inherently limited by the abundance of the bait in an affinity purification-mass spectrometry (AP-MS) experiment[19,20]. This is reflected in the triangular shape of stoichiometry maps (Figure 1b-d), where protein pairs of very low abundance stoichiometry have a strong tendency to not form high interaction stoichiometry interactions (Figures 1a, b). Abundance is therefore a confounding factor when using interaction stoichiometry as a proxy for binding strength.

To obtain direct measures of PPI binding affinities, we devised a simple approach to estimate equilibrium dissociation constants (K_d) from protein stoichiometries and protein abundances measured by mass spectrometry (*Methods*). Briefly, the number of complexes formed by a pair of proteins was estimated using the measured interaction stoichiometry and cellular abundances. The number of complexes were then used to estimate the number of unbound copies of each interactor. Finally, copy numbers were converted to concentrations, from which second order equilibrium dissociation constants were calculated (K_d; *Methods*, equations 1 and 2). We validated our results by comparing the estimated values to *in vitro* measurements of dissociation constants from literature and found a moderate but significant correlation of 0.40 (Pearson's r, *p*-value = 5.2×10^{-4}; Supplementary Figure 3a), which was comparatively better than that of interaction stoichiometry (Pearson's r = -0.29, *p*-value = 0.016; supplementary figure 3b). The theoretically expected relationship between K_d and interaction stoichiometry is inversely proportional. It has been noted that important criteria to assess reliability (*i.*e., controlling for titration and demonstrating equilibrium) in equilibrium binding experiments often go unreported throughout literature, and different experimental approaches can sometimes yield non-overlapping values of

K_d[40,41]. Moreover, experiments are generally performed *in vitro* and under conditions that do not

reflect native cellular environments. Considering these important sources of variability, we

reasoned that our values fall within an acceptable range of precision. To our knowledge, this is the

first report of a proteome-wide estimate of *in vivo* binding affinities.

The density heat map relating GI scores and K_d shows a visible bias towards negative GIs for high

affinity interactions in yeast (Figure 2a). This bias is also apparent in the data from human proteins,

however, the sparsity of GI data for human genes makes this relationship less clear (Figure 2b).

To quantitatively evaluate the relationship between K_d and GI strength, we binned the PPIs into

quantiles according to K_d and calculated the mean GI strength within each bin. The mean strengths

of negative GIs correlate remarkably and significantly with K_d for both yeast (Figure 2c;

Spearman's ρ = -0.98, *p*-value = 2.2 $\times$ 10^{-5}) and human PPIs (Figure 2d; Spearman's ρ = 0.95,

p-value = 0.014). A weak but significant correlation is also observable for positive GIs in yeast,

which is likely driven by an enrichment of significant positive GIs within complexes composed of

non-essential genes (Supplementary Figure 4c; Table S2)[1,6]. At the molecular level, this trend

could be explained by a mechanism where the absence of a single protein (single gene knockout)

is enough to completely disrupt the assembly of a complex in which it is embedded[42]. Specifically,

in protein biochemistry, the presence of a binding partner can increase the folding stability of a

protein by providing nucleation sites for its unfolded species, an effect that is heightened for

stronger PPI[43]. The single gene knockout could destabilize the other gene, consequently, the

double knockout is not substantially more drastic than the single knockout.

The strong correlation of PPI and negative GI suggests that PPI binding affinity is a major driver

of negative epistasis between the genes that code for the interacting proteins. These results imply

that one network could be predicted from the other. To test this idea, we devised an approach to

reconstruct GI network matrices from binding data alone (*Methods*). We observed that the clusters

along the diagonal of a matrix reconstructed from PPI binding data in yeast qualitatively matches

that of experimental results in genetic screens (Figure 2ef) [1].Overall, these results indicate that

there exists a strong relationship between the quantitative features of GI and PPI networks, and

that the modular structure of quantitative PPI networks seem to match that observed in GI

networks.

Figure 2. Genetic interactions correlate with PPI binding affinity. a, b, Density heat map of
scores as a function of dissociation constants (K_d) for yeast PPIs (a), and the human K562 cell line
(b) data. **c, d,** Binning PPIs according to K_ds reveals a significant monotonic relationship between
K_d and negative GIs. Negative GIs versus K_d fit to a sigmoidal equation, and for positive GIs
versus K_d relationship, a linear equation. Blue dots and trend line correspond to negative GI scores,
and yellow to positive GI scores. *P*-values for Spearman rank correlations coefficients were
computed with the following alternative hypothesis: Mean GI increases as K_ds decrease.
Correlations for unbinned data are reported in Table S3. **e,** A section of the yeast GI network
matrix, generated experimentally by synthetic genetic array (SGA). **f,** A reconstructed GI network
matrix, generated computationally from a PPI network weighted by K_ds. Hierarchical clustering
was performed on the reconstructed matrix, and the resulting order of genes was applied to the
experimental matrix. Full matrices are available on Figshare[44,45].

Redundancy and divergence of paralogs are reflected in genetic epistasis and PPI affinities

In our results above, we hypothesized that PPI, specifically binding affinity (K_d), is an accurate proxy for cellular function and thus, it quantitatively reflects epistasis across the genome. Since the number of protein complexes is a function of abundance and K_d, this poses a specific prediction that genomic events resulting in a maintained decrease or increase in abundance over macro-evolutionary time scales will be accompanied by a change in K_d. In particular, gene duplication events, which could effectively double the number of cellular copies of a protein, could relax selection on K_d and still maintain the same number of complexes. Correspondingly, due to the relaxed selection on K_d, we also expect a reduction in genetic interaction.

To test this hypothesis, we identified paralogs in our dataset and grouped them into four distinct types: 1) PPIs in which neither protein has a paralog, 2) PPIs in which one of the two (but not both) proteins has a paralog, 3) PPIs in which both proteins have a paralog, and 4) PPIs in which a protein interacts with its own paralog (paralogous heteromers); (Figure 3). Consistent with our expectation and previous analyses of GIs in paralogs, duplicated genes show weaker GIs with other genes than non-duplicates genes (Figure 3b)[46]. This effect seems to be additive overall, as gene pairs where both genes have paralogs show weaker GIs (Figure 3b; green). Remarkably, a similar trend is observed for binding affinity, where proteins encoded by duplicated genes tend to form weaker (high K_d) PPIs (Figure 3c).

Intriguingly, two distinct modes are observable in the distribution of K_ds for interactions in which both proteins have a paralog (Figure 3c). This result is reminiscent of results from analyses of tri-genic GIs involving duplicated genes, in which functionally redundant paralogs form more trigenic interactions (as opposed to digenic interactions) than functionally divergent paralogs, thereby resulting in a distinct bimodal distribution[47]. To test whether the separation of paralogs into redundant and divergent groups drives the bimodality observed in the K_ds, we computed the sequence divergence (defined here as the number of non-synonymous mutations per non-synonymous site) between genes and their paralogs and separated them accordingly into "high" and "low" divergence paralogs (supplementary figure 7). For interactions in which both proteins have a paralog, highly divergent paralogs participate in stronger interactions (lower K_d) than

paralogs with similar amino acid sequences (Figure 4d); this is also true for interactions in which

only one protein has a paralog (Figure 4c).

It is known that some duplicated genes have conserved sequences and functions over evolutionary

time[48,49]. However, protein abundance is expected to increase in the short term after gene

duplication events, resulting in an imbalance of concentrations of protein complex components,

which in turn negatively impacts fitness[50,51]. To reconcile functional conservation and

concentration imbalance, it has been proposed that for short evolutionary time scales, post-

translational expression attenuation is an important mechanism for robustness against an abrupt

abundance increase following gene duplication and, over larger time scales, reduced expression

allows for maintenance of redundant duplicated genes[52,53].

However, reduced expression is not observed for all gene duplication events, suggesting other

mechanisms may be at play[53]. In light of our results, we propose a model to explain how, on longer

evolutionary time scales, protein complexes adapt to increased expression of their subcomponents

after gene duplication (Figure 4e). We hypothesize that following a duplication event, paralogous

genes that conserve their sequence will conserve their interaction partners but participate in weaker

interactions overall to balance the binding equilibrium such that the number of complexes stays

the same (Figure 4e, Fate 1). This can happen, for instance, when mutations occur in a few

important amino acids at the binding interface or allosteric sites of paralogs, or if their interacting

partners diverge such that their binding affinity for the duplicated genes is decreased. Furthermore,

the subset of highly divergent paralogs participating in high affinity interactions (Figure 4d) can

be explained by a second scenario, where one of the duplicated genes diverges and loses (or

drastically reduces the affinity for) interacting partners while the other maintains high affinity

interactions (Figure 4e; Fate 2). This model is both compatible with previous works showing that

abundance is decreased following gene duplication[52,53] and explains the tendency of highly

divergent paralogs to make high affinity PPIs.

Figure 3. Functionally redundant paralogs participate in low binding affinity PPIs. a, GI scores for gene pairs which have one or two paralogs are lower than those without paralogs. Distributions for individual GI type (positive or negative) are included in supplementary figure 6. **b,** Dissociation constants (K_d) between proteins involved in a PPI are higher (lower affinity) if one or both proteins have a paralog. **c, d,** PPIs in which one (c) or both (d) proteins have a paralog separated according to sequence divergence. Significance levels are annotated as: * for $p \leq 0.05$, ** for $p \leq 0.01$, and *** for $p \leq 0.001$. **e,** Hypothesized evolutionary model where duplicated genes with low sequence diverge (Fate 1) participate in weaker interactions over time. For duplicated genes with high sequence divergence (Fate 2), one paralog loses (or greatly reduces) their affinity for common interactors.

Compositions of module and connector regions of the PPI network overlap with corresponding GI network regions

The classical interpretation of GI and PPI overlap mainly focuses on GIs within and between modules[4,6,18]; commonly referred to as the "within pathway" and "between pathway" models, respectively[18]. Indeed, protein-protein interaction modules ("complexes") are reflected in the functional modules in GI networks[1,6,22]. However, not all proteins that participate in strong interactions form stable complexes of specific function; some participate in transient interactions that "connect" protein complexes together[54]. The degree to which these "connectors" in PPI networks are reflected in GI networks has not been explored. We hypothesize that the ability of certain proteins to connects complexes in PPI networks also holds for GI networks. In other words, if a protein connects two complexes (modules) in the PPI network, it will likely connect the same two functional modules in GI networks.

At the cellular level, functional modules involving proteins are typically metabolic pathways, signaling pathways or protein complexes[55]. However, we specifically limit our analysis of protein modules to solely include stable protein complexes in yeast, as defined by the ComplexPortal database[56]. Furthermore, GI modules were taken from an analysis of the double knockout GI network used in the previous sections[1]. For both GIs and PPIs, connectors were defined as the set of genes or proteins that did not fall into a module. As a preliminary step, we sought to determine the extent to which proteins/genes in modules overlapped between the two networks using this classification. We observe a significant overlap between the modules and for connectors for both networks (P-value $= 1 \times 10^{-49}$, hypergeometric test) (Figure 4b, left panels).

GI modules map directly to PPI modules

We next sought to determine whether GI modules could be mapped one-to-one with protein complexes. To this end, we evaluated the similarity in gene composition of all possible GI module and PPI module pairings and identified module pairs with the highest degree of overlap (*methods*). Strikingly, of the 112 GI modules, 88 could be mapped to a protein complex (Figure 4B, upper left panel), which is significantly more than would be expected by chance (hypergeometric test *P*-

value = 7×10^{-160}, Figure 4C). This analysis is complementary to that of Costanzo et al.[1], in that we

use a set of manually curated complexes, as opposed to GO enrichment, to demonstrate

correspondence between GI and PPI modules. The remaining GI modules are likely attributable to

pathways not involving stable complexes. For instance, a GI module composed of ERG13 and

MVD1 was determined to be enriched for steroid and isoprenoid synthesis but was not mapped to

any known complex. It is also important to note that the large fraction of non-overlapping protein

complexes reflect the redundancy inherent to the ComplexPortal database, in which several

variants or subunits of a complex are included as separate entries.

Connector proteins in PPI networks are analogous to connector genes in GI networks

The ability of connector proteins to bridge protein complexes has been highlighted in recent mass

spectrometry screens of protein-protein interactions in human cells[19,20,54]. We reasoned that if most

GI modules overlap strongly with protein complexes, some GI connectors may have topological

properties similar to that of PPI connectors. The ability to bridge protein complexes being the most

salient topological property of protein connectors, we focused on systematically identifying this

motif network-wide for both GI and PPIs. Towards this aim, we devised an algorithm consisting

of four main steps as illustrated in Fig. 4a for a single pair of modules from each network: 1)

Mapping GI and PPI modules 1:1 as described above, 2) Finding the shortest paths between all

pairs of modules; this was done independently in GI and PPI networks, 3) Identifying connector

nodes (genes or proteins) in the shortest paths that are common between both networks, and 4)

combining the common components into an integrated network. We identified 555 shortest path

connectors common to both networks (Figure 4b, lower right panel). Comparing this degree of

overlap to that of randomized networks, we find that our results are extremely unlikely to arise by

chance (hypergeometric test *P*-value = 4×10^{-53}) (Figure 1D). Overall, our results demonstrate that

connectors bridging modules are a topological feature shared by both GI networks and PPI

networks.

Integrating PPI and GI networks highlights the functionality of connector genes and proteins

Previous analyses combining GIs and protein complexes have focussed on GIs within and between complexes and the functional interpretations thereof[1,6,18]. Our approach to integrating PPI and GI networks into a "module-connector-module" structure could reveal novel functional relationships between the two. In fact, our integrated network reveals a core set of modules shared between both networks and highlights known functionality of well-studied connectors. For instance, a subnetwork of our integrated network correctly identifies the karyopherin kap123p as an important bridge between the nuclear pore and various complexes that are transported and/or localized in the nucleus (figure 4e). The integrated network also places *RAS2* (Ras2p) adjacent to the adenylyl cyclase complex and the polarisome, which aligns with their respective roles in cell growth[57–60]. Interestingly, the integrated network also indicates that *RAS*2 connects the adenylyl cyclase complex to mitochondrial complexes, supporting a recently proposed mechanism in which Ras2p in the mitochondria promotes apoptosis *via* the cAMP/PKA pathway[61]. These examples highlight the potential of analysing motifs in the connector region for generating new hypotheses on the functions of genes.

Figure 4. Integrating genetic interactions networks and protein-protein interaction networks using common topological structures. a, Brief description of the algorithm devised to identify connectors of interest. First, shortest paths between modules in GI and PPI networks are determined. Modules of GI and PPI networks are mapped 1:1 and their common nodes are identified; common shortest path nodes are also identified. The final integrated network is comprised of common components in the two networks. **b,** Summary statistics of different overlapping components in GI and PPI networks. **c,** A statistical validation of overlaps between modules in GI and PPI networks by comparing the observed overlap with that of randomized modules. **d,** A statistical validation of overlaps between shortest path in GI and PPI networks by comparing the number of observed shortest path overlaps to that of randomized networks. **e,** A subset of the integrated network centered around the nuclear pore complex and the karyopherin KAP123.

Discussion

Determining how GI and PPI networks are related is an outstanding problem in systems biology and its resolution has profound implications for our understanding of the genotype-phenotype relationship. While some relationship between binding affinity and GI strength could fit intuitive expectation, a quantitative relationship between binding affinity and the continuum of GI values has never been demonstrated. In this study, we examined the two properties that define biological networks (weights and topology) to establish a rigorous framework through which PPI and GI networks can be analysed, predicted, and integrated. Although we focus on networks generated using data from the model organism *Saccharomyces cerevisiae*, we find that the principles established in yeast translate well to human networks.

The relationship between binding energies and GIs was further examined by estimating PPI dissociation constants proteome-wide, which we found to be strongly correlated with values from literature. These K_d estimations were used as a baseline to construct new models relating protein binding to genetic interactions. Indeed, binning PPIs according to binding information enabled both a reconstruction of a yeast GI network which qualitatively matched experimental results and the identification of an equation relating the general quantitative relationship between GIs and K_d:

$$\varepsilon^- = \frac{a}{1 + e^{b \cdot \ln(K_d) + d}} + c \tag{1}$$

Where K_d is the dissociation constant of a PPI, ε^- is negative epistasis (multiplicative model) between the interacting proteins' genes, and *a*, *b*, *c* and *d* are free parameters. Importantly, these parameters have tangible biological and biophysical meanings. For instance, the parameter *a* sets the range of epistasis values. Further, the parameter *b* (replaced by kT below), the parameter *d*, and K_d form the Gibbs free energy of binding equation:

$$\Delta G = -kT \ln(K_d) + d \tag{2}$$

We note that equation 1, relating K_d and epistasis is a sigmoidal, having the same form as the equation relating K_d and fraction bound. The implication of this correspondence between GI and PPI networks is profound: Tracing non-additive effects (epistasis) to their molecular thermodynamic underpinnings implies the possibility of modelling effects of individual genetic variations on a given trait (genotype-to-phenotype relationship). A caveat of our analysis is that epistasis is defined specifically in the context of pairwise gene deletions (or temperature sensitive

mutant gene knock downs) as opposed to other type of genetic changes such as point mutations,

recombination, or structural variants. Moreover, our model for the reconstruction of a complete GI

network (Figure 2f) explains approximately 2% of the entire variance in the negative GI network,

including GIs between genes whose proteins do not interact with each other. Although a better

understanding of how genetic perturbations percolate through the PPI network would likely

increase explainability, it is likely that GIs are also strongly driven by other molecular mechanisms

such as substrate and product dependencies between enzymes in metabolic networks, and

transcription factor-promoter relations in gene regulatory networks.

Certain limitations exist as well regarding our estimates of dissociation constants. First, each PPI

is treated as completely independent of any other, leading to deviations from the actual values. For

instance, competitive binding will systematically lead to underestimations of binding strength,

especially for very strong interactions. Second, under our current model, interaction

stoichiometries over 1:1, which may occur if more than one prey protein binds to a single bait

protein, lead to negative K_d values which are physically impossible. The number of interactions

which are discarded as a result represents a non-negligible fraction of the interactome, which may

bias the distribution of our estimated K_d. Third, our ability to assess the precision of our estimates

is limited to the quality of reference binding affinity data in the literature. Fourth, we assume all

reported interactions in our datasets to be direct. In reality, AP-MS approaches also detect indirect

interactions which are not readily distinguishable from direct ones.

Previous approaches aiming to identify common topological features between GI networks and

PPI networks have mostly focused on GIs within or between protein modules[4,6,18]. However, not

all genes and proteins cluster into highly connected groups. Although it has long been known that

certain proteins serve as "connectors" between protein modules[62], the extent to which this

"module-connector-module" motif occurs in GI networks had not been explored. We demonstrate

that the common topological organization in GI and PPI networks extends further than overlap in

modules, but also the connectors that bridge these modules. By merging modules and connectors

from both GI and PPI networks an integrated network was constructed, where the functional role

of connector genes/proteins are highlighted and can be used a novel tool for functional inference

that will be valuable for predicting GIs in other organisms.

Finally, the quantitative and topological relationships we have established in this study could
provide means of interpreting and predicting fundamental relationships between genes underlying
normal and disease phenotypes, as well as identify specific vulnerabilities or changes in PPIs that
could predict potential therapeutic targets for diseases.
449

450 **Acknowledgments**
451

The authors thank Anastasia Baryshnikova for advice on preprocessing of synthetic genetic array
genetic interaction data. We also thank Christian R. Landry, Ben Lehner, Elena Kuzmin, and
Brenda Andrews for comments on the manuscript. We acknowledge funding from the following
agencies: Natural Sciences and Engineering Research Council of Canada – Masters Canada
Graduate Scholarship (XCG); Fonds de Recherche du Québec en Santé – Masters Training
Scholarship (XCG); Canadian Institute of Health Research grants MOP-G-408523 (AWRS);
Natural Sciences and Engineering Research Council (NSERC) of Canada (NSERC) grant RGPIN-
2016-06566 (AWRS); Canada Research Chairs (SWM, AWRS); CIHR grant PJT-185976 and
NSERC grant RGPIN-03216-2021 (SWM).
461

462 **Author Contributions**
463

SWM conceptualized this study. XCG generated the computational pipeline. XCG, AWRS and
SWM analyzed and interpreted data. XCG, AWRS and SWM wrote the manuscript.

Supplementary Figures

Supplementary Figure 1. Stable protein complexes in *Saccharomyces cerevisiae* are enriched

for 1:1 stoichiometries. Yeast PPIs occurring between proteins of stable complexes as annotated

by the ComplexPortal database[56].

Supplementary Figure 2. The PPI stoichiometry landscape of GIs in the Jurkat cell line. a,
b, Mean negative and positive genetic interaction scores for PPIs throughout the stoichiometry
plot. GI scores in human were taken from a screen in Jurkat cell lines. **c, d,** Distribution of negative
and positive GI scores within the 4 defined regions of the stoichiometry plot. Significance levels
are annotated as: ** for $p \leq 0.01$.

Supplementary Figure 3. Validation of dissociation constants estimated from quantitative
mass spectrometry measurements. Pearson correlation was used in all panels. **a,** Estimated PPI
dissociation constants estimated from quantitative mass spectrometry data in human cell lines
correlate moderately but very significantly with values from literature. **b,** Interaction
stoichiometries from quantitative mass spectrometry data in human cell lines correlate weakly with
values from literature. **c,** Dissociation constants estimated from quantitative mass spectrometry
data in yeast correlate moderately but very significantly with those estimated from protein-
fragment complementation (PCA) results.

Supplementary Figure 4. Correlations between binned genetic interactions and K_d for genes

pairs of different essentialities. P-values are computed from two tailed test. All trendlines are fit

using a linear regression. Pearson correlation coefficients between dissociation constants (K_d) and

GIs are computed for genes pairs where both genes are essential (**a**), only one gene is essential (**b**),

or both genes are non-essential (**c**).

Supplementary Figure 5. Fitting an equation relating binned PPI weights (derived from K_d values) to GI scores. a, Yeast PPIs are binned into 25 quantiles according to their network weights. The average weight and GI score is calculated for every quantile and an exponential equation is used to fit the data. **b,** The fitting procedure described in (a) was tested with different numbers of quantiles to visualize their root mean square errors (RMSE). **c,** A narrowing of RMSE spread and a dip in RMSE values can be observed at around 25 quantiles.

506 **Supplementary Figure 6. Distribution of positive and negative genetic interaction scores**

507 **according to PPIs separated by their paralog composition. a,** Distribution of estimated K_d and

508 GIs (yeast dataset) according to the PPI's paralog composition/type. **b,** Distribution of negative GI

509 scores for different interaction type. **c,** Distribution of positive GI scores for different interaction

510 type.

Supplementary Figure 7. Dividing protein-protein interactions according to paralog sequence divergence. a, Sequence divergence (dN) between paralogous protein/gene pairs retrieved from literature. A dN threshold of 0.2 separates the two visible modes in the distribution. **b, c,** The statistically significant separations of dissociation constants between high divergence and low divergence observed in Figure 3 (panels c and d) are robust to the chosen dN threshold. P-values are calculated using Wilcoxon rank sum test. Red line indicates a P-value of 0.05.

520 **Supplementary Table 1[a]: Essential Complexes Enriched for Negative GIs**

ComplexPortal ID	Common Name
CPX-1890	Exocyst
CPX-1364	SMC5/6 complex
CPX-1041	DASH complex
CPX-1656	Transcription factor TFIIIC complex
CPX-607	Arp2/3 protein complex
CPX-1186	MIND complex
CPX-1869	Nuclear condensin complex
CPX-1831	Eukaryotic translation initiation factor 3 complex
CPX-2262	20S proteasome core particle
CPX-1895	mRNA cleavage factor complex

521 **[a]Examples were taken from Costanzo et al. (2016)[1]**
522
523 **Supplementary Table 2[a]: Non-Essential Complexes Enriched for Positive GIs**

ComplexPortal ID	Common Name
CPX-779	Elongator holoenzyme complex
CPX-1840	Golgi transport complex
CPX-568	CAF-1 complex
CPX-1197	FAR complex
CPX-1342	Set3C
CPX-1884	HDA complex
CPX-1193	Vacuolar proton-transporting V-type ATPase
CPX-1412	Ubp3-Bre5 ubiquitin hydrolase complex
CPX-1655	i-AAA complex
CPX-1178	Cytoplasmic dynein complex

524 **[a]Examples were taken from Costanzo et al. (2016)[1]**
525

526 **Supplementary Table 3: Correlation coefficients between unbinned K_d and GI values**

Organism	GI Type	Correlation Measure	Coefficient	p-value
Yeast	Negative	Pearson	0.22	1.3×10^{-23}
		Spearman	0.24	1.1×10^{-26}
	Positive	Pearson	0.083	2.8×10^{-3}
		Spearman	0.091	1.0×10^{-3}
Human	Negative	Pearson	0.098	0.11
		Spearman	0.13	0.027
	Positive	Pearson	0.038	0.48
		Spearman	0.019	0.73

527
528

Methods

Collecting yeast and human genetic and protein-protein interaction data

For yeast, raw synthetic genetic array (SGA) genetic interaction data was downloaded from TheCellMap[1,63] in matrix format. Data from essential and non-essential strain crosses (Essential vs. Essential, Essential vs. Nonessential, Nonessential vs. Nonessential) were combined. In instances where multiple GIs were measured for a pair of genes, only the entry with the most significant *P*-value was retained. For human cell lines, GI data was retrieved from a digenic knockdown screen of 472x472 genes in Jurkat and K562 cells[34] (table S5 of reference 34). For both yeast and human data, unless otherwise indicated, analyses were performed using all GIs regardless of significance scores.

Yeast protein-protein interactions were taken from a near exhaustive whole-proteome affinity enrichment coupled to an affinity purification-mass spectrometry (AP-MS) screen[21] (Yeast_Interactome_Edges.csv from supplementary file 2 of the reference 28). Human protein-protein interactions and stoichiometries were taken from two quantitative AP-MS screens: one in which genes were tagged with GFP and expressed on bacterial artificial chromosomes in HeLa cells[19] (table S2 of the reference 19), and a second in which genes are endogenously tagged in HEK 293T cells with the small fragment of the split mNeonGreen$_2$ tag[20] (table S4 of the reference 20). Both human cell line interactomes were concatenated but common interactions between the two screens were kept as separate data points for all analyses.

Computing interaction stoichiometries and abundance stoichiometries in yeast

Interaction stoichiometries were calculated from data generated in an AP-MS screen[21] using a previously described strategy[19]. The starting preprocessed table containing normalized and imputed label-free quantification (LFQ) intensities from the AP-MS screen were provided by the authors. Briefly, computing interaction stoichiometries was done in three mains steps: 1) Subtracting the median intensity of a protein across all sample, 2) Dividing LFQ intensities by the number of theoretically observable peptides of lengths between 7 and 30 amino acids, 3) dividing

the values observed for the prey protein by that of the bait protein. Importantly, the number of
theoretically observed peptides were computed using sequences from the Uniprot database[64]
(https://www.uniprot.org/proteomes/UP000002311) using LysC only digestion (as opposed to
LysC and Tryptic digestion). Negative interaction stoichiometries (which arise because of
normalization in step 1) were dropped from the dataset. The final dataset includes interaction
stoichiometries for 7348 unique PPIs.

Protein abundances were taken from a dataset unifying measurements from various sources [65]
(table S4 of the referenced article). Abundance stoichiometries were determined by calculating the
ratio between the median copy numbers of the prey over that of the bait, as described previously[19].
Abundance measurements for both interactors were available for 7336 out of the 7348 PPIs for
which interaction stoichiometry values were calculated. The 22 interactions which had interaction
stoichiometry values, but not abundances stoichiometry values, were conserved in the dataset for
analyses in which abundance stoichiometry was not relevant.

Analysis of genetic interactions and protein-protein interaction stoichiometries

The relationship between GIs and stoichiometries was explored in yeast and human cell lines (see
first section for data collection). To merge GI and PPI stoichiometry datasets, tables were merged
such that if genes comprising a GI also code for proteins that interact, the GI score is assigned to
that PPI. Four stoichiometric regions were graphically defined within stoichiometry plots
according to previous criteria[19]. Hein et al. defined region 1 as the space enclosed by a circle
centered at (-0.5, 0) where the abscissa is interaction stoichiometry, the ordinate is abundance
stoichiometry, and coordinates are in base 10 logarithmic scale. Using these definitions as a
reference, region 3 was defined using circles of identical radii (0.75) centered at (-0.5, 1.5), region
2 was centered at (-2.5, 0) for yeast and (-3.5, 0) for human, and region 4 was centered at (-2.5, -
1.5) for yeast and (-3.5, -1.5) for human.

Estimating protein-protein interaction dissociation constants

591 To estimate dissociation constants from quantitative mass spectrometry data, we first assumed all
592 individual PPIs to be independent of all others (no competitive binding) and their binding
593 dynamics to be describable by a two-state system:

$$A + B \rightleftharpoons AB \tag{1}$$

$$K_d = \frac{[A][B]}{[AB]} \equiv \frac{[Bait][Prey]}{[Bait::Prey]} \tag{2}$$

Where K_d is the dissociation constant, $[A]$ the molar concentration of unbound protein A, $[B]$ the
concentration of unbound protein B, and $[AB]$ the concentration of the complex formed by proteins
A and B. In a given mass spectrometry experiment, one of the two proteins will be the bait (tagged
protein retained on column) and the other a prey.

Next, we sought to estimate the number of bait-prey complexes using interaction stoichiometries.
By definition, interaction stoichiometry is a proxy measurement of the ratio of prey protein bound
to bait protein over the total abundance of bait protein. Under the previously mentioned
assumptions, the amount of prey bound to bait is equal to the number of bait-prey complexes.

$$Interaction\ Stoichiometry = \frac{Prey_{LFQ_{Intensity}}}{Bait_{LFQ_{Intensity}}} \approx \frac{Prey_{Bound\ to\ Bait}}{Bait_{Total}} = \frac{Bait::Prey}{Bait_{Total}} \tag{3}$$

611 Therefore, we used the product of interaction stoichiometry and total abundance of bait (in copy
612 numbers) to estimate the number of bait-prey complexes in a given cell.

$$Bait::Prey = \frac{Bait::Prey}{Bait_{Total}} \cdot Bait_{Total} = (Interaction\ Stoichiometry) \cdot Bait_{Total} \tag{4}$$

For yeast PPIs, the cellular copy numbers of bait proteins were taken from the same source as that
described to calculate abundance stoichiometries[65]. For PPIs in human cell lines, abundances were
taken from whole proteome quantification experiments which were performed alongside the AP-
MS measurements of interaction stoichiometries[19,20]
(opencell.czbiohub.org/data/datasets/opencell-protein-abundance.csv [downloaded April 4th
2022]; Hein et al., table S3)[19].
622
623   Next, the number of unbound bait and prey proteins were estimated by subtracting the number of
624   bait-prey complexes[39,66]. In a few instances, the estimated number of bait-prey complexes was
625   higher than that of the total abundance of bait or prey, resulting in a negative number of unbound
626   proteins. This almost only arose when interaction stoichiometry is above 1. These negative values
627   were removed.
628

$$Bait_{Unbound} = Bait_{Total} - Bait::Prey \qquad (5)$$

$$Prey_{Unbound} = Prey_{Total} - Bait::Prey \qquad (6)$$

633   Copy numbers of bait proteins, prey proteins and bait-prey complexes were next converted to
634   molar concentrations (mol/L) using Avogadro's constant and cell volumes. Below is an example
635   formula for bait concentrations; concentrations prey and bait-prey complexes were calculated in
636   the same way. Dissociation constants were calculated using the resulting concentrations.
637

$$[Bait] = \frac{Bait_{unbound}/(6.022 \cdot 10^{23} \ mol^{-1})}{Cell \ Volume} \qquad (7)$$

$$K_d = \frac{[Bait][Prey]}{[Bait::Prey]} \qquad (8)$$

A cell volume of 4.2 $\times$ 10^{-14} Liters was used for yeast PPIs (which were measured in haploid
cells)[67]. A cell volume of 2.6 $\times$ 10^{-14} Liters was used for PPIs measured in HeLa cells by Hein et
al.[68] Cho et al. measured PPIs and stoichiometries in HEK293 cells, for which no experimental

measurement of volume was found. However, the diameter of HEK293 cells was measured to be

approximately 13 micrometers (Bionumbers ID: 108893). Therefore, an estimated cell volume of

1.15×10^{-14} Liters was calculated from this diameter by approximating the shape of a cell to that

of a sphere. The above-mentioned values were all found via the BioNumbers database[69].

Validating estimated dissociation constants using values from literature

Experimental dissociation constants from literature were taken from the protein-protein complex

subset of the PDBbind database (2020 version)[70,71]. Taxonomic identification numbers were

extracted from the PDB file of each database entry. Only interactions involving proteins from

budding yeast and human were retained. PDB files which listed more than two proteins were

removed, as it would not be possible to assign dissociation constants to the correct protein pair.

Systematic names or IDs for proteins and genes are absent from the PDBbind database. For this

reason, the PDB ID for each complex was used to access their structure's RCSB Protein Data Bank

web page and retrieve the Uniprot IDs of the proteins within the structure. Web scraping for the

data published in this article was performed on April 23rd 2024, using the *BeautifulSoup4 Python3*

package. For yeast PPIs, Uniprot IDs were then converted to systematic ORF IDs using the

org.Sc.sgd.db R package. For human PPIs, a Uniprot to HGNC symbol conversion table was

downloaded (on February 28th 2024) from the "Custom downloads" page of the HGNC website[72].

Information on the original source, which is often different from that of the structure, and details

on the experimental setup of each dissociation constant measurement is also absent from the

database. Although this limits quality control, it is known from curated sets for structure-based

modelling of PPI affinities that common methods to measure PPI binding affinities are best for

interactions in the micromolar to nanomolar range[41]. Therefore, measurements were limited to

dissociation constants no smaller than 1 nM and no greater than 1 mM ($1 \text{ nM} \leq K_d < 1 \text{ mM}$). The

final reference dataset contains 505 K_d measurements for human PPIs and 59 for yeast PPIs.

Estimated K_d values for PPIs were merged with experimental values for the reference dataset. For

PPIs in human cell lines, 70 estimated K_d values were mapped to values from literature. Only 4

estimated K_d values were mapped to values from literature for yeast PPI.

Comparison of mass spectrometry K_d estimates in yeast using PCA K_d estimates
678

Colony intensities and z-scores were collected from a genome-wide screen for protein-protein
interaction using a survival-selection protein-fragment complementation assay (PCA) [73](table S4
of the referenced article) and filtered to only contain the final 2770 protein-protein interactions
(table S3 of the referenced article) reported in the article. Next colony intensities were collected
from a proteome-wide abundance screen based on the GFP-PCA method[74] (Data Set S2 of the
reference article). GFP-PCA intensities were merged to cellular copy numbers from a compilation
of abundances in yeast[65] to generate a linear regression model relating PCA colony intensities to
copy numbers. Protein-protein interaction PCA intensities were then rescaled to a range similar to
that of the abundance GFP-PCA intensities. The linear regression model was then used to convert
protein-protein interaction intensities to numbers of complexes. With the number of copies of
complexes and individual protein copy numbers at hand, dissociation constants were estimated as
described in the "Estimating protein-protein interaction dissociation constants" section. In total,
1480 K_d estimates were calculated from PCA data, of which 48 also had K_d estimated from mass
spectrometry data.

Reconstructing a GI network matrix from a PPI binding network

Yeast GI and PPI dissociation constant (K_d) tables were merged such that if genes comprising a
GI also code for proteins that interact, the GI score is assigned to that PPI. The final table contains
3265 interactions, each with a GI score and a K_d value. A graph was then generated, using
individual proteins as nodes/vertices (V_{PPI}), PPIs as edges (E_{PPI}), and the reciprocal negative
logarithm of K_d values as weights (W_{PPI}):

702
$$G_{PPI} = \{V_{PPI}, E_{PPI}, W_{PPI}\} \tag{9}$$

704
$$w_{PPI}(\{u, v\} \in E_{PPI}) = \frac{1}{-log10(K_{d,u,v})} \tag{10}$$

Where $u, v \in V_{PPI}$, $K_{d,u,v}$ is the dissociation constant between the interacting proteins u and v, and
$w_{PPI}(e)$ is the weight assigned to the edge e (where $e = \{u, v\}$ connects u and v).

To establish an equation relating weights and GI scores, PPIs were first binned into quantiles

according to their weights. The mean weight and mean GI score within each quantile were then

calculated and an exponential equation was used to fit the data (See supplementary Figure 5a for

example with 25 quantiles). The root mean square error (RMSE) of the fit for different numbers

of quantiles were tested to find the most optimal (Supplementary Figure 5b). A dip of RMSE

values and narrowing of RMSE spread was observable at around 25 quantiles, which justified its

choice as the ideal number of quantiles for the final fitting (Supplementary Figure 5c and 5a).

To mimic the completeness of GI networks, the PPI graph was converted to an "association" matrix

similar to that described by Rives and Galitsky[75]. Briefly, the *shortest_path_length* function of the

Networkx python library was used on G_{PPI}, using W_{PPI} as the weight parameter, to generate a $|V| \times$

$|V|$ matrix (*d*) in which d[i,j] denotes the length of the shortest path between the nodes (proteins) i

and j. This all-pairs-shortest-path (APSP) matrix was used as a graph (G_{APSP}), such that:

722
$$G_{APSP} = \{V_{APSP}, E_{APSP}, W_{APSP}\} \tag{11}$$

724
$$w_{APSP}(\{x, y\}) = shortest\_path\_length(u, v) \tag{12}$$

Where $\{u, v\} \in E_{PPI}$, $\{x, y\} \in E_{APSP}$, and both edges represent the same protein interaction. The

exponential equation relating means of quantized weights and GI scores was then used on W_{APSP}

to generate a new graph (G_{GI_pred}) with weights W_{GI} containing estimated GI scores. The adjacency

matrix of G_{GI_pred} was clustered with average linkage hierarchical clustering using Pearson

correlation in Cluster 3.0[76].

Analysis of dissociation constants in yeast paralogs

Whole genome duplication (WGD) paralogs were collected from Kellis et al.[77] (Supplementary

file 8) and small-scale duplication (SSD) paralogs were collected form VanderSluis et al.[46]

Paralogs were separated into 3 categories as described in the main text. The rate of non-

synonymous mutations per non-synonymous site (dN) was used as a proxy for sequence

divergence. To calculate dN values, paralog pairs were aligned pairwise at the protein level with

ClustalW (version 2.1) using the PAM protein weight matrix. Using the *reverse.align* function of

the *seqinr* R package, protein-coding nucleic acid sequences were next aligned using protein

alignments as a guide. The reverse alignments of each paralog pair were then used to compute dN

using the *kaks* function of *seqinr*. Amino acid sequences and coding nucleic acid sequences were

downloaded from the SGD sequence archives (http://sgd-

archive.yeastgenome.org/sequence/S288C_reference/, March 2024) for the S288C reference

strain. A sequence divergence threshold of 0.2 was selected to categorize paralog pairs into "high"

and "low" sequence divergence (dN) (Supplementary Figure 7a). For protein-protein interactions

where both interactors have a paralog, the interaction was categorized into "high" when both

interactors had a dN value above 0.2.

Mapping GI Modules to PPI Modules

GI clusters were downloaded from an analysis of a near-complete double knockout study in yeast[1]

(Data File S6). Yeast protein complexes were downloaded from a species-specific dataset of the

ComplexPortal database[56]; protein Uniprot IDs were converted to systematic yeast ORFs using the

org.Sc.sgd.db R library. All GI clusters were compared to all PPI complexes by their common

ORFs, the degree of overlap was scored using the Jaccard index. For each GI cluster, the protein

complex which had the highest overlap (Jaccard Index) was retained.

If a GI cluster overlapped better with connector nodes (proteins outside complexes), it was dropped

from the dataset. To validate the significance of the overlap, GI clusters were randomized and

mapped to protein complexes. GI clusters were randomized by retaining the size of the clusters

and randomly redistributing ORFs. The randomization was performed for 10 000 iterations on a

general-purpose computer cluster. A p-value was derived from a z-score in which the mean Jaccard

score from the experimental dataset was compared to the distribution of mean Jaccard scores of

randomized clusters (Figure 4c).

Identifying Nodes Connecting GI and PPI Modules

The GI and PPI networks used for the analysis of "connector" nodes were taken from the near-

complete double knockout screen in yeast and whole-proteome affinity enrichment coupled to

mass spectrometry screen described in the *Collecting yeast and human genetic and protein-protein*

interaction data section. Only negative genetic interactions were used for the analysis in figure 4.

Because shortest paths are sensitive to noise, we used the moderately stringent cutoff of GI score

< -0.08 and p-value < 0.05 recommended by the authors[1]. Identifying nodes (gene/proteins)

connecting modules in GI and PPI networks was performed in 4 main steps. First, genetic modules

and protein modules are mapped 1-to-1 as described in the *Mapping GI Modules to PPI Modules*

section above.

The 1-to-1 mapping can be expressed mathematically, where M_{GI} and M_{PPI} are each a set

containing all modules in the GI and PPI network, respectively, and a module is itself a set of

genes/proteins. Connectors can be defined as the sets C_{GI} and C_{PPI}, which contain all the connector

genes/proteins in the GI and PPI networks respectively. Importantly, a gene/protein cannot be both

a connector and within a module. In other words:

785
$$C_{GI} \cap \bigcup M_{GI} = \emptyset \tag{13}$$

786
$$C_{PPI} \cap \bigcup M_{PPI} = \emptyset \tag{14}$$

787   We note that here the large union symbol denotes the union of all genes/proteins that are within

788   modules into a single large set. Next, a module $X \in M_{GI}$ is related (mapped) to a module $A \in M_{PPI}$

789   if and only if $J(X, A) > J(X, C_{PPI})$, and $J(X, A) > J(X, B)$ for all $B \in M_{PPI}$. This relation (mapping)

790   is denoted as R, where $(X, A) \in R$ for all modules X in M_{GI} and for all modules A in M_{PPI}. Note that

791   J denotes the Jaccard index:

792
$$J(X, A) = \frac{X \cap A}{X \cup A} \tag{15}$$

Second, for all combinations of modules within a network, the shortest path between all pairs of

nodes were computed. This was done independently for GI and PPI networks. In mathematical

terms, for a graph $G = \{V, E, W\}$ (representing the GI or PPI network)*,* let $\{a, b, c\}$ be a set of

genes within module A such that $A = \{a, b, c\}$ and $\{a, b, c\} \subset$ of V, and let $\{x, y, z\}$ be a set of

genes within module B such that $B = \{x, y, z\}$ and $\{x, y, z\} \subset V$. Importantly, $B \cap A = \emptyset$. For any

pair $(u, v) \in A \times B$, the shortest paths between nodes u and v were computed using Dijkstra's

algorithm (*all_short_paths* function of the *networkx* python package).

Third, connector nodes that bridge the same modules in both GI and PPI networks were

systematically identified. If the sequence of nodes (i, j, k) is the shortest path between module A

and module B in the PPI network (G_{PPI}) and the sequence (k, m, n) is the shortest path between

module X and module Y in the GI network (G_{GI}), then $(i, j, k) \cap (k, m, n)$ is the shortest path

common to both networks. Note that here modules A and X, as well as B and Y are mapped to each

other; $(A, X) \in R$ and $(B, Y) \in R$. Importantly, to avoid including nodes from the source and target

modules in the shortest path, nodes belonging to a module were removed from the shortest path.

Fourth, the overlapping features in both networks *i.e.,* modules and shortest paths connectors, are

combined to generate an integrated network. To determine if the extent of the overlap in connectors

is due to network topology as opposed to random chance, the approach described above was

performed on randomized networks. The networks were randomized by first conserving edges

(interactions) within modules and then performing degree preserved randomization[78] for all other

edges. Overlapping connectors were identified for 1 000 random GI and PPI networks on a general-

purpose computer cluster. A p-value was derived from a z-score in which the number of

overlapping connectors from the experimental dataset was compared to the distribution of

overlapping connectors from randomized networks (Figure 4d).

Code and Data Availability

All original scripts as well as information on how to properly access data and run the code is

provided in a GitHub repository dedicated to this article (https://github.com/Xavier-Castellanos-

Girouard/PPI_Binding_Epistasis).

1. Costanzo, M. *et al.* A global genetic interaction network maps a wiring diagram of cellular

function. *Science* **353**, aaf1420–aaf1420 (2016).

2. Costanzo, M. *et al.* Global Genetic Networks and the Genotype-to-Phenotype Relationship.

Cell **177**, 85–100 (2019).

3. Sharan, R., Ulitsky, I. & Shamir, R. Network-based prediction of protein function. *Molecular*

Systems Biology **3**, 88 (2007).

4. Beyer, A., Bandyopadhyay, S. & Ideker, T. Integrating physical and genetic maps: from

genomes to interaction networks. *Nat Rev Genet* **8**, 699–710 (2007).

5. Fang, G. *et al.* Discovering genetic interactions bridging pathways in genome-wide

association studies. *Nat Commun* **10**, 4274 (2019).

6. VanderSluis, B. *et al.* Integrating genetic and protein–protein interaction networks maps a

functional wiring diagram of a cell. *Current Opinion in Microbiology* **45**, 170–179 (2018).

7. Mani, R., St.Onge, R. P., Hartman, J. L., Giaever, G. & Roth, F. P. Defining genetic

interaction. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 3461–3466 (2008).

8. Phillips, P. C. Epistasis — the essential role of gene interactions in the structure and

evolution of genetic systems. *Nat Rev Genet* **9**, 855–867 (2008).

9. Zuk, O., Hechter, E., Sunyaev, S. R. & Lander, E. S. The mystery of missing heritability:

Genetic interactions create phantom heritability. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 1193–

1198 (2012).

10. Sackton, T. B. & Hartl, D. L. Genotypic Context and Epistasis in Individuals and

Populations. *Cell* **166**, 279–287 (2016).

11. Manolio, T. A. *et al.* Finding the missing heritability of complex diseases. *Nature* **461**, 747–

753 (2009).

12. Aguirre, L., Hendelman, A., Hutton, S. F., McCandlish, D. M. & Lippman, Z. B.

Idiosyncratic and dose-dependent epistasis drives variation in tomato fruit size. *Science* **382**,

315–320 (2023).

13. Mair, B., Moffat, J., Boone, C. & Andrews, B. J. Genetic interaction networks in cancer

cells. *Current Opinion in Genetics & Development* **54**, 64–72 (2019).

14. Shen, J. P. & Ideker, T. Synthetic Lethal Networks for Precision Oncology: Promises and

Pitfalls. *Journal of Molecular Biology* **430**, 2900–2912 (2018).

15. Costanzo, M. *et al.* The Genetic Landscape of a Cell. *Science* **327**, 425–431 (2010).

16. Oliver, S. Guilt-by-association goes global. *Nature* **403**, 601–602 (2000).

17. Sahni, N. *et al.* Widespread Macromolecular Interaction Perturbations in Human Genetic

Disorders. *Cell* **161**, 647–660 (2015).

18. Kelley, R. & Ideker, T. Systematic interpretation of genetic interactions using protein

networks. *Nat Biotechnol* **23**, 561–566 (2005).

19. Hein, M. Y. *et al.* A Human Interactome in Three Quantitative Dimensions Organized by

Stoichiometries and Abundances. *Cell* **163**, 712–723 (2015).

20. Cho, N. H. *et al.* OpenCell: Endogenous tagging for the cartography of human cellular

organization. *Science* **375**, eabi6983 (2022).

21. Michaelis, A. C. *et al.* The social and structural architecture of the yeast protein interactome.

Nature **624**, 192–200 (2023).

22. Bandyopadhyay, S., Kelley, R., Krogan, N. J. & Ideker, T. Functional Maps of Protein

Complexes from Quantitative Genetic Interaction Data. *PLoS Comput Biol* **4**, e1000065

(2008).

23. Collins, S. R. *et al.* Functional dissection of protein complexes involved in yeast

chromosome biology using a genetic interaction map. *Nature* **446**, 806–810 (2007).

24. Braberg, H. *et al.* From Structure to Systems: High-Resolution, Quantitative Genetic

Analysis of RNA Polymerase II. *Cell* **154**, 775–788 (2013).

25. Braberg, H. *et al.* Genetic interaction mapping informs integrative structure determination of

protein complexes. *Science* **370**, eaaz4910 (2020).

26. Halabi, N., Rivoire, O., Leibler, S. & Ranganathan, R. Protein Sectors: Evolutionary Units of

Three-Dimensional Structure. *Cell* **138**, 774–786 (2009).

27. Marks, D. S. *et al.* Protein 3D Structure Computed from Evolutionary Sequence Variation.

PLoS ONE **6**, e28766 (2011).

28. Faure, A. J. *et al.* Mapping the energetic and allosteric landscapes of protein binding

domains. *Nature* **604**, 175–183 (2022).

29. Ho, B., Baryshnikova, A. & Brown, G. W. Unification of Protein Abundance Datasets Yields

a Quantitative Saccharomyces cerevisiae Proteome. *Cell Systems* **6**, 192-205.e3 (2018).

30. Lynch, M. & Conery, J. S. The Origins of Genome Complexity. *Science* **302**, 1401–1404

(2003).

31. Lynch, M. & Force, A. The Probability of Duplicate Gene Preservation by

Subfunctionalization. *Genetics* **154**, 459–473 (2000).

32. Crow, K. D. What Is the Role of Genome Duplication in the Evolution of Complexity and

Diversity? *Molecular Biology and Evolution* **23**, 887–892 (2006).

33. Dolinski, K. & Botstein, D. Orthology and Functional Conservation in Eukaryotes. *Annu.*

Rev. Genet. **41**, 465–507 (2007).

34. Horlbeck, M. A. *et al.* Mapping the Genetic Landscape of Human Cells. *Cell* **174**, 953-

967.e22 (2018).

35. Van Leeuwen, J. *et al.* Exploring genetic suppression interactions on a global scale. *Science*

354, aag0839 (2016).

36. He, X., Qian, W., Wang, Z., Li, Y. & Zhang, J. Prevalent positive epistasis in Escherichia

coli and Saccharomyces cerevisiae metabolic networks. *Nat Genet* **42**, 272–276 (2010).

37. Segrè, D., DeLuna, A., Church, G. M. & Kishony, R. Modular epistasis in yeast metabolism.

Nat Genet **37**, 77–83 (2005).

38. Xie, X., Sun, X., Wang, Y., Lehner, B. & Li, X. Dominance vs epistasis: the biophysical

origins and plasticity of genetic interactions within and between alleles. *Nat Commun* **14**,

5551 (2023).

39. Diss, G. & Lehner, B. The genetic landscape of a physical interaction. *eLife* **7**, e32472

(2018).

40. Jarmoskaite, I., AlSadhan, I., Vaidyanathan, P. P. & Herschlag, D. How to measure and

evaluate binding affinities. *eLife* **9**, e57264 (2020).

41. Kastritis, P. L. *et al.* A structure-based benchmark for protein–protein binding affinity.

Protein Science **20**, 482–491 (2011).

42. Diss, G., Dubé, A. K., Boutin, J., Gagnon-Arsenault, I. & Landry, C. R. A Systematic

Approach for the Genetic Dissection of Protein Complexes in Living Cells. *Cell Reports* **3**,

2155–2167 (2013).

43. Wright, P. E. & Dyson, H. J. Linking folding and binding. *Current Opinion in Structural*

Biology **19**, 31–38 (2009).

44. Castellanos-Girouard, X. *GI_experimental_full.Png*. 11784216 Bytes (figshare, 2024).

doi:10.6084/M9.FIGSHARE.27020701.

45. Castellanos-Girouard, X. *GI_reconstructed_full.Png*. 6541289 Bytes (figshare, 2024).

doi:10.6084/M9.FIGSHARE.27020704.

46. VanderSluis, B. *et al.* Genetic interactions reveal the evolutionary trajectories of duplicate

genes. *Molecular Systems Biology* **6**, 429 (2010).

47. Kuzmin, E. *et al.* Exploring whole-genome duplicate gene retention with complex genetic

interaction analysis. *Science* **368**, eaaz5667 (2020).

48. Nowak, M. A., Boerlijst, M. C., Cooke, J. & Smith, J. M. Evolution of genetic redundancy.

Nature **388**, 167–171 (1997).

49. Vavouri, T., Semple, J. I., Garcia-Verdugo, R. & Lehner, B. Intrinsic Protein Disorder and

Interaction Promiscuity Are Widely Associated with Dosage Sensitivity. *Cell* **138**, 198–208

(2009).

50. Rice, A. M. & McLysaght, A. Dosage-sensitive genes in evolution and disease. *BMC Biol*

15, 78 (2017).

51. Papp, B., Pál, C. & Hurst, L. D. Dosage sensitivity and the evolution of gene families in

yeast. *Nature* **424**, 194–197 (2003).

52. Ascencio, D. *et al.* Expression attenuation as a mechanism of robustness against gene

duplication. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2014345118 (2021).

53. Qian, W., Liao, B.-Y., Chang, A. Y.-F. & Zhang, J. Maintenance of duplicate genes and their

functional redundancy by reduced expression. *Trends in Genetics* **26**, 425–430 (2010).

54. Michnick, S. W. & Levy, E. D. The modular cell gets connected. *Science* **375**, 1093–1094

(2022).

55. Pereira-Leal, J. B., Levy, E. D. & Teichmann, S. A. The origins and evolution of functional

modules: lessons from protein complexes. *Phil. Trans. R. Soc. B* **361**, 507–517 (2006).

56. Meldal, B. H. M. *et al.* Complex Portal 2018: extended content and enhanced visualization

tools for macromolecular complexes. *Nucleic Acids Res* **47**, D550–D558 (2019).

57. Tamanoi, F. Ras Signaling in Yeast. *Genes & Cancer* **2**, 210–215 (2011).

58. Cazzanelli, G. *et al.* The Yeast Saccharomyces cerevisiae as a Model for Understanding RAS

Proteins and their Role in Human Tumorigenesis. *Cells* **7**, 14 (2018).

59. Ho, J. & Bretscher, A. Ras Regulates the Polarity of the Yeast Actin Cytoskeleton through

the Stress Response Pathway. *MBoC* **12**, 1541–1555 (2001).

60. Xie, Y. & Miao, Y. Polarisome assembly mediates actin remodeling during polarized yeast

and fungal growth. *Journal of Cell Science* **134**, jcs247916 (2021).

61. Bonomelli, B., Busti, S., Martegani, E. & Colombo, S. Active Ras2 in mitochondria

promotes regulated cell death in a cAMP / PKA pathway-dependent manner in budding

yeast. *FEBS Letters* **597**, 298–308 (2023).

62. Han, J.-D. J. *et al.* Evidence for dynamically organized modularity in the yeast protein–

protein interaction network. *Nature* **430**, 88–93 (2004).

63. Usaj, M. *et al.* TheCellMap.org: A Web-Accessible Database for Visualizing and Mining the

Global Yeast Genetic Interaction Network. *G3 Genes|Genomes|Genetics* **7**, 1539–1549

(2017).

64. The UniProt Consortium *et al.* UniProt: the Universal Protein Knowledgebase in 2023.

Nucleic Acids Research **51**, D523–D531 (2023).

65. Ho, B., Baryshnikova, A. & Brown, G. W. Unification of Protein Abundance Datasets Yields

a Quantitative Saccharomyces cerevisiae Proteome. *Cell Systems* **6**, 192-205.e3 (2018).

66. Celaj, A. *et al.* Quantitative analysis of protein interaction network dynamics in yeast.
Molecular Systems Biology **13**, 934 (2017).

67. Jorgensen, P., Nishikawa, J. L., Breitkreutz, B.-J. & Tyers, M. Systematic Identification of
Pathways That Couple Cell Growth and Division in Yeast. *Science* **297**, 395–400 (2002).

68. Zhao, L. *et al.* Intracellular water-specific MR of microbead-adherent cells: the HeLa cell
intracellular water exchange lifetime. *NMR in Biomedicine* **21**, 159–164 (2008).

69. Milo, R., Jorgensen, P., Moran, U., Weber, G. & Springer, M. BioNumbers—the database of
key numbers in molecular and cell biology. *Nucleic Acids Research* **38**, D750–D753 (2010).

70. Liu, Z. *et al.* PDB-wide collection of binding data: current status of the PDBbind database.
Bioinformatics **31**, 405–412 (2015).

71. Wang, R., Fang, X., Lu, Y. & Wang, S. The PDBbind Database: Collection of Binding
Affinities for Protein−Ligand Complexes with Known Three-Dimensional Structures. *J.*
Med. Chem. **47**, 2977–2980 (2004).

72. Seal, R. L. *et al.* Genenames.org: the HGNC resources in 2023. *Nucleic Acids Research* **51**,
D1003–D1009 (2023).

73. Tarassov, K. *et al.* An in Vivo Map of the Yeast Protein Interactome. *Science* **320**, 1465–
1470 (2008).

74. Levy, E. D., Kowarzyk, J. & Michnick, S. W. High-Resolution Mapping of Protein
Concentration Reveals Principles of Proteome Architecture and Adaptation. *Cell Reports* **7**,
1333–1340 (2014).

75. Rives, A. W. & Galitski, T. Modular organization of cellular networks. *Proc. Natl. Acad. Sci.*
U.S.A. **100**, 1128–1133 (2003).

76. De Hoon, M. J. L., Imoto, S., Nolan, J. & Miyano, S. Open source clustering software.

Bioinformatics **20**, 1453–1454 (2004).

77. Kellis, M., Birren, B. W. & Lander, E. S. Proof and evolutionary analysis of ancient genome

duplication in the yeast Saccharomyces cerevisiae. *Nature* **428**, 617–624 (2004).

78. Maslov, S. & Sneppen, K. Specificity and Stability in Topology of Protein Networks.

Science **296**, 910–913 (2002).