Proteome-wide reverse molecular docking reveals folate receptor as a mediator of PFAS-induced neurodevelopmental toxicity

Ally Xinyi Kong[1*], Maja Johnson[1*], Aiden F Eno[1], Khoa Pham[1], Ping Zhang[1], Yijie Geng[1 2 #]

[1]Department of Environmental and Occupational Health Sciences, Seattle, WA 98105, USA.

[2]Department of Pharmacology, University of Washington, Seattle, WA 98105, USA.

*These authors contributed equally

[#]Corresponding author. Email: yjgeng@uw.edu

ABSTRACT

Per- and polyfluoroalkyl substances (PFAS) are a class of long-lasting chemicals with widespread use and environmental persistence that have been increasingly studied for their detrimental impacts to human and animal health. Several major PFAS species are linked to neurodevelopmental toxicity. For example, epidemiological studies have associated prenatal exposure to perfluorooctanoate (PFOA) and perfluorononanoate (PFNA) with autism risk. However, the neurodevelopmental toxicities of major PFAS species have not been systematically evaluated in an animal model, and the molecular mechanisms underlying these toxicities have remained elusive. Using a high-throughput zebrafish social behavioral model, we screened six major PFAS species currently under regulation by the Environmental Protection Agency (EPA), including PFOA, PFNA, perfluorooctane sulfonate (PFOS), perfluorohexanesulfonic acid (PFHxS), perfluorobutane sulfonate (PFBS), and hexafluoropropylene oxide dimer acid ammonium salt (GenX). We found that embryonic exposure to PFNA, PFOA, and PFOS induced social deficits in zebrafish, recapitulating one of the hallmark behavioral deficits in autistic individuals. To uncover protein targets of the six EPA-regulated PFAS, we screened a virtual library containing predicted binding pockets of over 80% of the 3D human proteome through reverse molecular docking. The screen predicts that folate receptor beta (FR-β, encoded by the gene *FOLR2*) interacts strongly with PFNA, PFOA, and PFOS but to a lesser degree with PFHxS, PFBS, and GenX, correlating positively with their *in vivo* toxicity. These predictions were validated through *in silico* molecular docking, *in vitro* protein binding analysis, and *in vivo* loss-of-function verification. Furthermore, embryonic co-exposure to folic acid effectively rescued social deficits induced by PFAS. The folate pathway has been implicated in autism, indicating a novel molecular mechanism for PFAS in autism etiology.

INTRODUCTION

Autism spectrum disorder (ASD) is a complex neurodevelopmental condition marked by social communication deficits. Over recent decades, the prevalence of autism has risen significantly, with reported cases in the United States increasing from 1 in 150 children diagnosed in 2000 to 1 in 36 in 2020, according to the Centers for Disease Control and Prevention (CDC). This increase is only partially accounted for by improved diagnostic criteria and awareness[1,2]. Meanwhile, the ever-growing chemical industry continues to introduce new chemicals into our environment, most of which with unknown toxicity to humans. An estimated 10 million new chemicals are synthesized annually, and within them, ~ 700 become mass produced for commercial use[3]. Emerging

research indicates that environmental factors contribute substantially to autism risk, with recent
estimates attributing up to 40% of ASD etiology to environmental influences[4,5]. Understanding
how environmental factors contribute to ASD is critical to mitigating the disease prevalence and
could inform targeted prevention and treatment strategies.

However, environmental contributors to ASD remain poorly understood, creating a gap in research
methods and assessment approaches to identify and evaluate these risk factors comprehensively.
Among environmental toxins with known associations with ASD, per- and polyfluoroalkyl
substances (PFAS) have become a particular focus of recent research[6,7]. PFAS is a class of
synthetic chemicals known for their persistence in the environment and biological systems. They
are widely present in nonstick cookware, stain-resistant textiles, and various industrial
applications. Their pervasive use has led to widespread human exposure, with the CDC reporting
PFAS detected in the blood of 97% of Americans. The health risks associated with PFAS exposure
are substantial, including links to kidney and liver disease, immune and thyroid dysfunction,
cancer, and, critically, developmental challenges[8]. Based on these pioneering studies, beginning
in April 2024, the U.S. Environmental Protection Agency (EPA) established legally enforceable
levels for six PFAS in drinking water, including perfluorooctanoic acid (PFOA), perfluorooctane
sulfonic acid (PFOS), perfluorononanoic acid (PFNA), perfluorohexanesulfonic acid (PFHxS),
perfluorobutane sulfonate (PFBS), and hexafluoropropylene oxide dimer acid ammonium salt
(GenX). Despite these progresses, however, with 15,538 PFAS compounds identified in the
Distributed Structure-Searchable Toxicity (DSSTox) database by the EPA[9] and over 7 million
PFAS species reported in PubChem to date[10], a comprehensive understanding of their biological
impact remains elusive, particularly regarding how prenatal exposures to various PFAS species
contribute to ASD risk. For example, although epidemiology studies linked PFNA and PFOA to
increasing ASD risk[6,7,11], whether the other four EPA-regulated PFAS contribute equally to ASD
remains unclear. Mechanisms underlying their neurodevelopmental toxicity are also largely
unknown.

To address these gaps, we developed a novel model and methodology using zebrafish. Zebrafish
offer a unique advantage as a model organism for studying social behavior[12]—a core behavioral
deficit in ASD. They share approximately 70% of their genome with humans, begin social
interactions at an early developmental stage, and are amenable to high-throughput chemical
screening and behavioral assays. In our previous work, we designed the Fishbook social
behavioral assay[13] to quantitatively assess social behavioral deficits in zebrafish, a behavior
analogous to social deficits in ASD. In this assay, a test fish is exposed to a social stimulus in a
controlled environment, allowing us to measure a social score based on proximity behaviors.
Notably, the Fishbook assay has observed social deficits in zebrafish following ASD-related gene
knockouts and exposure to ASD-related environmental toxicants[13-15], indicating its relevance for
examining environmental toxicants' influence on ASD. The Fishbook system enables us to screen
different PFAS compounds in parallel using the same assay platform for their impacts on sociality,
alongside genetic knockout models to elucidate their molecular mechanisms.

Moreover, to uncover potential molecular targets of PFAS, we conducted virtual screening through
reverse molecular docking using the HPRoteome-BSite database[16] to explore interactions
between PFAS and the predictable 3D human proteome. HProteome-BSite is a database of
predicted binding pockets covering over 80% of the UniProt entries in the AlphaFold human
proteome database[17,18]. The reverse molecular docking approach, unlike traditional docking
methods which typically screen a large number of chemicals against one or a few specific proteins,
allows us to comprehensively evaluate how different PFAS species interact with a range of
proteins involved in neurodevelopmental pathways.

In this study, we leverage our unique zebrafish model and assay with advanced virtual screening
techniques and validation experiments to evaluate the impacts of the six EPA-regulated PFAS on

social behavioral development and identified their molecular targets in the human proteome.
Among the molecular targets predicted by the virtual screen, folate receptors were found to
differentially bind to the six PFAS species, with the binding affinities positively correlating with their
toxicity on social development. *In silico*, *in vitro*, and *in vivo* experiments verified folate receptor
beta as a biological target of PFNA, PFOA, and PFOS. Co-exposure to folic acid rescued social
deficits caused by PFAS. Given the importance of folic acid in early neurodevelopment[19-21] and
its implications for autism[22], our findings strongly suggest that PFAS' interaction with folate
receptors likely contribute to their ASD-related neurodevelopmental toxicity and point to a
potential strategy for preventive interventions.

104   **RESULTS**

The six EPA-regulated PFAS showed differential impacts on social behavioral development
in zebrafish.

To compare the impacts of major PFAS species on social behavioral development, we tested the
six EPA-regulated PFAS, namely PFNA, PFOA, PFOS, PFHxS, PFBS, and GenX (Figure 1A),
using the Fishbook assay (Figure 1B). After an initial embryonic toxicity testing, a range of doses
were determined for each PFAS to assess their dose responses. The highest doses for the dose-
response analysis were set as 50% of the lowest doses that caused 100% mortality in the initial
toxicity test. The remaining doses for the dose-response analysis were set as 50% serial
reductions from the highest dose. Specifically, PFNA was evaluated at 0.78 μM to 200 μM, PFOA
at 3.125 μM to 1600 μM, PFOS at 0.25 μM to 12 μM, PFHxS at 3.125 μM to 400 μM, PFBS at
3.125 μM to 1600 μM, and GenX at 1.56 μM to 1600 μM. Embryos were exposed without
dechorionation, with no observed impacts on hatchability due to PFAS exposures. Embryos were
exposed to a single dose of PFAS at the concentrations listed above from 3 hours post fertilization
(hpf) to 3 days post fertilization (3 dpf).

Following the embryonic exposure, embryos were raised to the juvenile stage for Fishbook testing
at 23 dpf. The survival rates of each PFAS treatment condition at 23 dpf are shown in
Supplementary Figure 1A-1F. Note that there is a ~50% attrition during the nursery growth period
(5 dpf to 23 dpf) for even the control samples (Supplementary Figure 1A-1F). The range of survival
rates for all concentrations of PFNA (41.7%-58.3%; Supplementary Figure 1A), PFOA (26.7%-
66.7%; Supplementary Figure 1B), PFOS (53.3%-85%; Supplementary Figure 1C), GenX
(41.7%-68.3%; Supplementary Figure 1E), and PFBS (36.7%-73.3%; Supplementary Figure 1F)
exposures were each comparable to the survival rates of their respective vehicle control (45% for
PFNA, 38.3% for PFOA, 73.3% for PFOS, 50% for GenX, and 51.7% for PFBS). While the range
of survival rates for the lower doses of PHFxS (55%-75%; Supplementary Figure 1D) remained
comparable to the control (53.3%), the highest two doses of PHFxS (200 μM and 400 μM;
Supplementary Figure 1D) appeared to negatively impact survival (25% and 18.3%, respectively).
For Fishbook assessment of social behavioral deficits induced by the exposures, we observed a
clear dose-response for PFNA (Figure 1C; Supplementary Figure 1G), PFOA (Figure 1D;
Supplementary Figure 1H), and PFOS (Figure 1E; Supplementary Figure 1I) but not for PFHxS
(Figure 1F; Supplementary Figure 1J), PFBS (Figure 1G: Supplementary Figure 1K), and GenX
(Figure 1H) (Supplementary Figure 1L). The IC_{50} values were determined to be 23.85 μM (11.068
ppm) for PFNA (Figure 1C), 40.15 μM (16.63 ppm) for PFOA (Figure 1D), and 0.1892 μM (0.095
ppm) for PFOS (Figure 1E) by fitting the dose-response data to a four-parameter logistic sigmoidal
model using nonlinear least-squares regression. These results affirm PFNA and PFOA's
neurodevelopmental toxicity, which are aligned with previous epidemiology findings linking PFNA
and PFOA to ASD risk. Our results further demonstrate the toxicity of PFOS in social behavioral
development and suggest its potential implication for ASD risk.

Figure 1. Fishbook assay revealed differential neurodevelopmental toxicity of six EPA-regulated PFAS on social behavioral development in zebrafish. (**A**) Chemical structures of the six EPA-regulated PFAS. (**B**) A schematic of the dosing and testing workflow. (**C-H**) Dosage curves generated by plotting average social scores (y-axis) against PFAS concentrations (x-axis). PFAS tested include PFNA (C), PFOA (D), PFOS (E), PFHxS (F), GenX (G), and PFBS (H). IC_{50} values are shown for PFNA (C), PFOA (D), and PFOS (E). Sample sizes are as follows for the lowest to highest concentrations for each compound: n=27, 29, 29, 34, 25, 34, 35, 25, 32, 29, 31 for PFNA (C); n=23, 34, 37, 29, 40, 31, 16, 23, 30, 38, 16 for PFOA (D); n=44, 35, 44, 32, 51, 50, 47, 40, 38 for PFOS (C); n=32, 45, 38, 43, 40, 41, 33, 15, 11 for PFHxS (F); n=30, 28, 32, 29, 28, 30, 41, 31, 30, 34, 38, 25 for GenX (G); and n=31, 44, 26, 35, 31, 37, 22, 29, 38, 30, 35 for PFBS (H). Data were fitted to a sigmoidal curve using a four-parameter logistic (4PL) model, with X being concentration, through least squares fit.

Discovering protein binding targets of PFAS via reverse molecular docking against the human proteome.

To discover the protein targets of PFAS, we conducted a reverse molecular docking screen against predicted binding pockets[16] of the human 3D proteome based on the AlphaFold database[17,18] (Figure 2A). A single domain was used for pocket prediction for most of the proteins, while multiple domains were used for predictions for a subset of proteins. For each domain, pockets were predicted based on both sequence and 3D structure. The highest-ranking sequence-based and 3D-structure-based pocket predictions were selected for each prot ein domain for the screen. A total of 33446 predicted pockets were screened. Each pocket was screened once via molecular docking for each PFAS. The combined screening results are shown in Supplementary Table 1, screening results for each PFAS are separately shown in Supplementary Tables 2-7.

Figure 2. Reverse molecular docking discovered molecular targets of PFAS in the human proteome. (**A**) A schematic drawing demonstrating the reverse molecular docking approach. (**B**) A Venn diagram showing overlaps between the predicted protein targets of PFNA, PFOA, and PFOS. Hits were identified by selecting protein-PFAS interactions with predicted binding energy < -10 kcal/mol. The 7 proteins predicted to bind to all three PFAS are listed in the table pointed by the arrow. (**C-E**) Lists of the top 10 hits with predicted binding energy <-10 kcal/mol for PFNA (C), PFOA (D), and PFOS (E). (**F**) Summary of the reverse molecular docking screen. All molecular docking scores of six EPA-regulated PFAS compounds against predicted binding pockets in 80% of human proteome are shown as a bar chart at the top. The top 50 highest affinity binding interactions (the lowest scores) are zoomed in and shown as a bar chart at the bottom. Interactions between PFAS (PFNA, PFOS, and PFOA) and FOLR2 are highlighted in orange. Ranking is based on average binding scores for each PFAS-domain interaction, calculated from binding energies between each PFAS and binding pockets predicted by structural-based prediction and sequence-based prediction for each protein domain.

We set a cutoff at binding energy of -10 kcal/mol. Using this cutoff, no hit was identified for PFBS and GenX, and only 1 hit was identified for PFHxS (Supplementary Table 5). A number of hits were identified for PFNA, PFOA, and PFOS (Figure 2B & Supplementary Figure 2A). We noticed that folate receptor beta, or FOLR2, consistently appeared among the top hits for the three PFAS (Figure 2B), ranking #2 among the predicted targets of PFNA with a binding energy of -10.9 kcal/mol (Figure 2C), #3 for PFOA at -10.2 kcal/mol (Figure 2D), and #1 for PFOS at -10.73 kcal/mol (Figure 2E). The other two folate receptor subtypes, folate receptor alpha (FOLR1) and folate receptor gamma (FOLR3) were also targeted by PFAS, although at lower affinities with FOLR1 binding to PFNA at an energy of -8.6 kcal/mol, to PFOA at -8.9 kcal/mol, and to PFOS at -9.4 kcal/mol, and with FOLR3 binding to PFNA at an energy of -9.9 kcal/mol, to PFOA at -9.4 kcal/mol, and to PFOS at -10 kcal/mol (Supplementary Tables 2-4). Another two proteins involved in the folate metabolic pathway, folate hydrolase 1 (FOLH1) and methenyltetrahydrofolate

synthetase (MTHFS), were also predicted to be bound by PFAS (Figure 2B & Supplementary
Tables 2-4), with FOLH1 binding to PFNA at an energy of -10.1 kcal/mol, to PFOA at -10.1 kcal/mol,
and to PFOS at -10.1 kcal/mol, and with MTHFS binding to PFNA at -10 kcal/mol, to PFOA at -
9.5 kcal/mol, and to PFOS at -6.5 kcal/mol. The combined outcome of the screening is
summarized in Figure 2F and Supplementary Figure 2B.

Consistent with our observations, enrichment analyses identified the folate pathway in PFNA,
PFOA, and PFOS targeted proteins among the Gene Ontology (GO) pathways (Figure 3A &
Supplementary Figures 3A-3C) and the Kyoto Encyclopedia of Genes and Genomes (KEGG)
pathways (Figure 3B & Supplementary Figures 3D-3F). Protein-protein interaction (PPI) analysis
using stringDB for the top hits (with predicted binding energy < -10 kcal/mol) of PFNA (Figure 3C),
PFOA (Figure 3D), and PFOS (Figure 3E) consistently identified an interaction between FOLR2
and FOLH1. We did not find additional PPIs that consistently connect to FOLR2.

Figure 3. Enrichment analysis and protein-protein interaction analysis consistently identified the folate pathway as a target of PFAS. (**A-B**) Enrichment analyses for the GO (A) and KEGG (B) pathways consistently identified the folate pathway for PFNA, PFOA, and PFOS. Values inside each bar represents adjusted *p*-value. (**C-F**) stringDB analysis for the top hits of PFNA (C), PFOA (D), and PFOS (E) found no consistent PPI networks except a connection between FOLR2 and FOLH1.

***In silico*, *in vitro*, and *in vivo* validations confirm folate receptor as a biological target of**
PFAS.

To verify FOLR2 as a binding target for PFNA, PFOA, and PFOS, and compare their binding
affinities with the endogenous ligand of FOLR2, folic acid (or folate), in its natural binding pocket,
we conducted molecular docking using a 3D structure of FOLR2 from the Protein Data Bank (PDB)
in complex with folate (4KMZ). We found that the natural binding pocket of folate (Figure 4A) can

be effectively bound by PFNA (Figure 4B), PFOA (Figure 4C), and PFOS (Figures 4D). In fact, an
average of 5 random docking trials showed that PFNA, PFOA, and PFOS all bind to FOLR2 at a
higher potency (-11.04±0.2191 kcal/mol, -10.34±0.1949 kcal/mol, and -10.28±0.04472 kcal/mol,
respectively) compared to folate (-9.68±0.04472 kcal/mol), while PFHxS, PFBS, and GenX bind
to FOLR2 at a lower potency compared to folate (-9.3±0.0 kcal/mol, -8.82±0.2049 kcal/mol, and -
8.1±0.0 kcal/mol, respectively) (Figure 4E). We also conducted molecular docking with FOLR1
using a 3D structure of FOLR1 from the PDB in complex with folate (4LRH) (Supplementary
Figures 4A-4D) and found a similar pattern of differential binding affinity among the PFAS species:
PFOS and PFNA bind to FOLR1 at a potency higher than folate (PFOS: -10.08±0.08357 kcal/mol,
PFNA: -9.8±0.2236 kcal/mol, folate: -9.58±0.1095 kcal/mol), whereas PFOA, PFHxS, PFBS, and
GenX bind to FOLR1 at a lower potency compared to folate (-9.3±0.0 kcal/mol, -8.7±0.0 kcal/mol,
-8.4±0.0 kcal/mol, and -7.7±0.0 kcal/mol, respectively) (Supplementary Figure 4E). Consistent
with a previous report which associated PFAS exposure to lower folate levels in human blood[23],
our findings suggest that exposure to PFNA, PFOA, and PFOS may out-compete FOLR2's natural
ligand, folate, and lead to reduced folate uptake *in vivo*.

We then examined whether PFNA, PFOA, and PFOS can physically bind to the FR-β protein. We
applied an *in vitro* assay called proteome integral solubility alteration (PISA), which detects the
binding of a chemical to its target protein by comparing the soluble portion of the compound-
protein mixture with a control mixture following a temperature gradient treatment[24]. His-tagged
recombinant human and mouse FR-β proteins were incubated with PFOS, PFOA, PFNA, folic
acid, or vehicle control for 30 minutes at room temperature. The treated samples were subjected
to a thermal gradient (34–66°C) in a thermocycler, pooled, and centrifuged to isolate soluble
proteins. Supernatants were applied to a nitrocellulose membrane at n=10 replicates for dot blot
analysis. We found that PFOS (Figure 4F), PFOA, PFNA, and folic acid (Figure 4G) all
significantly altered the amount of protein remaining in the soluble fraction after thermal gradient
treatment, demonstrating their bindings with the human FR-β protein. Similar results were
observed for PFOS (Figure 4H), PFOA, and PFNA (Figure 4I) against the mouse FR-β protein.
These results demonstrate that PFOS, PFOA, and PFNA can indeed physically bind to the FR-β
protein.

Finally, we investigated whether folate deficiency contributes to social behavioral deficits in our
zebrafish social behavior model. The zebrafish only possess one gene encoding the folate
receptor, *folr*. To evaluate how *folr* loss-of-function affects sociality, we knocked out the zebrafish
folr using CRISPR/Cas9 by injecting fertilized eggs with a mixture of three *folr* gRNAs and Cas9
protein. The resulting F0 mutants show significant social deficits compared to wild-type controls
(*p*<0.0001, Cohen's *d*=0.70 indicating a medium effect) (Figure 4J), demonstrating that folate
pathway is required for social behavioral development in zebrafish. The F0 CRISPR/Cas9
knockout strategy has been shown to faithfully reproduce disease-relevant phenotypes without
the confounding influence of compensatory gene expression[25,26]: by simultaneously targeting
multiple loci within a single gene, this approach has been reported to consistently achieve over
90% biallelic knockout efficiency and enable rapid phenotypic assessment in F0 animals[26-28].
Indeed, PCR amplification of the gRNA-targeted genomic regions using genomic DNA extracted
from injected F0 embryos show that all three gRNAs ablated the WT amplicon, indicating that all
three gNRAs effectively mutated the targeted *folr* gene (Supplementary Figure 5).

Figure 4. *In silico*, *in vitro*, and *in vivo* assays validated folic acid receptor as the biological target of PFAS. (**A-D**) *In silico* validation via molecular docking. Showing docking images for FOLR2 against folic acid (FA) (A), PFNA (B), PFOA (C), and PFOS (D). FOLR2 structure was acquired using the PDB identifier 4KMZ. (**E**) Comparing the binding energies for folic acid and the six EPA-regulated PFAS against FOLR2. Binding energies are shown as average and standard deviation of 5 independent docking attempts. Significance values are calculated to compare PFAS data with folic acid. ***: $p < 0.001$, ****: $p < 0.0001$. (**F-G**) Dot blot analyses detected significantly different protein levels in the pooled soluble fractions of PISA samples between PFAS-treated samples and vehicle controls. PISA was conducted using recombinant human FR-β protein (hFR-β) against PFOS (10 μM) (F), PFOA (1600 μM) (G), and PFNA (400 μM) (G). Folic acid (12.5 μM) (FA) was used as a positive control (G). Datapoints represent raw signal intensities of the dot blot assays. Each sample were tested at n=10 replicates. (**H-I**) PISA and dot blot analyses revealed similar results using recombinant mouse FR-β protein (mFR-β) against PFOS (H), PFOA (I), and PFNA (I). Datapoints represent raw signal intensities of the dot blot assays. Each sample were tested at n=10 replicates. (**J**) CRISPR-Cas9 induced F0 knockout of the zebrafish folic acid receptor *folr* induced social deficits in zebrafish. Each datapoint represents the social score of an individual fish. A total of n=86 wild-type control fish (WT) and n=59 *folr* KO fish were tested. Significance was calculated by one-way ANOVA and Dunnett's multiple comparison test when comparing multiple groups (E, G, I) or two-tailed Student's *t* test when comparing two groups (F, H, J). *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$.

Folic acid rescued social deficits induced by PFAS through embryonic co-exposure.

Finally, to evaluate whether PFAS induces social deficit through inhibiting the folate pathway, we
attempted to rescue social deficits caused by PFAS exposure by co-exposing with folic acid
(Figure 5A). We hypothesized that certain doses of folic acid may outcompete PFAS binding,
restoring folate metabolism in the developing embryos, without causing developmental toxicity.
We co-exposed the three PFAS species including 400 μM PFNA, 1600 μM PFOA, and 10 μM
PFOS with a range of folic acid doses (7 doses ranging from 3.125 μM to 200 μM in 2-fold
increments) during the first 3 days of embryonic development. Folic acid co-exposure effectively
rescued social deficits in all three PFAS treatments. PFNA was rescued by folic acid at the lowest
applied doses of 3.125 μM (p=0.008, Cohen's d=0.74 indicating a medium effect) and 25 μM
(p=0.03, d=0.58 indicating a medium effect) (Figure 5B & Supplementary Figure 6A). Both PFOA
and PFOS showed a bell-shaped rescue curve when co-exposed to folic acid. PFOA was most
effectively rescued by folic acid at 6.25 μM (p=0.018, d=0.88 indicating a large effect) to 12.5 μM
(p=0.0095, d=0.95 indicating a large effect) (Figure 5C & Supplementary Figure 6B), whereas

PFOS was robustly rescued by 12.5 µM (p=0.0002, d=1.16 indicating a large effect), 25 µM (p
<0.0001, d=1.39 indicating a large effect), 50 µM (p=0.02, d=0.82 indicating a large effect), and
100 µM (p=0.017, d=0.77 indicating a medium effect) folic acid treatments (Figure 5D &
Supplementary Figure 6C). These results indicate that PFAS likely induces social deficit by
inhibiting the folate pathway, and this inhibition can be prevented by embryonic co-exposure to
folic acid.

Figure 5. Embryonic co-exposure to folic acid rescued social deficits induced by PFAS. (**A**) A schematic of the co-treatment and testing workflow. (**B-D**) Dosage curves generated by plotting average social scores (y-axis) against folic acid (FA) concentrations (x-axis). Showing both linear (top) and log2 (bottom) scales for the x-axis. All treatment groups in each experiment were exposed to 400 µM PFNA (B), 1600 µM PFOA (C), or 10 µM PFOS (D), respectively, plus a range of folic acid (FA) doses (3.125 µM to 200 µM) to assess its rescue effect. Red dots marks the no-FA control (0 µM FA) of each experiment in the linear plots. Sample sizes are as follows for the lowest to highest FA concentrations in each rescue experiment: n=43, 44, 42, 46, 42, 29, 45, 22 for PFNA+FA (B); n=28, 29, 29, 32, 29, 30, 27, 18 for PFOA+FA (C); and n=32, 23, 36, 37, 20, 25, 31, 40 for PFOS+FA (D). Data were fitted to a bell-shaped curve using a four-parameter logistic (4PL) model, with X being concentration, through least squares fit. Significance was calculated by one-way ANOVA and Dunnett's multiple comparison test. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$.

DISCUSSION

259 Our study provides critical insights into the neurodevelopmental toxicity of six EPA-regulated
260 PFAS compounds—PFNA, PFOA, PFOS, PFHxS, PFBS, and GenX—on social behavioral
261 development in zebrafish. Using the Fishbook assay, we observed distinct dose-dependent
262 responses for PFNA, PFOA, and PFOS, which displayed neurodevelopmental impacts on social
263 behavior in the exposed embryos, while PFHxS, PFBS, and GenX exhibited no apparent dose-

dependent effects on social behavior within the tested concentration range. IC_{50} values were established for PFNA, PFOA, and PFOS, reinforcing epidemiological links between PFNA, PFOA, and ASD risk, while suggesting a potential contribution of PFOS to ASD risk. This specificity in behavioral effects aligns with our molecular docking results, where PFNA, PFOA, and PFOS displayed stronger binding affinities to the folate receptor FOLR2, suggesting that these PFAS may disrupt folate metabolism and thereby influence neurodevelopmental pathways implicated in social behavioral development. This proposed mechanism is consistent with previous human biomonitoring evidence showing an inverse association between folic acid and PFAS levels in human red blood cells and serum samples[23], indicating that PFAS exposure in humans likely impedes folic acid uptake or bioavailability. Based on these convergent behavioral, molecular docking, and epidemiological observations, we hypothesize that a subset of PFAS with specific physicochemical properties disrupt neurodevelopment by interfering with folate receptor-mediated folate transport during critical developmental windows, thereby increasing susceptibility to ASD-relevant social behavioral deficits.

PFAS exposure is nearly ubiquitous for people living in the U.S., but the magnitude and composition of exposure vary substantially across different exposure scenarios such as the general population, communities at risk of exposure, and occupational exposure settings. Understanding these real-world exposure levels is essential for interpreting our results and evaluating the translational relevance of our findings. According to the National Health and Nutrition Examination Survey (NHANES), in the U.S. general population (GP), median serum concentrations of PFOS and PFOA in women of childbearing age have declined significantly over the past two decades due to regulatory actions and changes in industrial practices. Among the PFAS species examined in this study, PFOS decreased from approximately 24 ng/mL (48 nM) in 1999–2000 to 3 ng/mL (6 nM) in 2017–2018, and PFOA decreased from 5 ng/mL (12 nM) to 1 ng/mL (2.4 nM) over the same period. In contrast, PFNA exhibited an initial rise from 0.5 ng/mL in 1999–2000 to 1.0 ng/mL (2 nM) in 2009–2010 before declining to 0.3 ng/mL (0.5 nM) in 2017–2018. Despite these downward trends, PFAS remain detectable in the blood of more than 97% of Americans, reflecting their environmental persistence, bioaccumulation, and widespread distribution in food and drinking water. Exposed Communities (EC) near industrial plants, military installations, or municipal water supplies continue to experience disproportionately high PFAS exposure. For example, residents served by the Little Hocking Water Association in Ohio had a mean PFOA serum level of 227.6 ng/mL (0.55 µM), while exposed communities in Decatur, Alabama reported a PFOS concentration of 39.8 ng/mL (80 nM). These levels are tens to hundreds of times higher than those observed in the general population. Occupational exposure (OE) settings represent the highest end of the PFAS exposure spectrum. Historical data from PFAS manufacturing workers in Decatur, Alabama, showed mean serum concentrations approaching 899 ng/mL (2.2 µM) for PFOA and 941 ng/mL (1.9 µM) for PFOS[29], which are orders of magnitude above exposure levels of the general population (**Table 1**).

	General Population Low (GPL)	General Population High (GPH)	Exposed Communities (EC)	Occupational Exposure (OE)
PFOS	3 ng/mL (6 nM)	24 ng/mL (48 nM)	39.8 ng/mL (80 nM)	941 ng/mL (1.9 µM)
PFOA	1 ng/mL (2.4 nM)	5 ng/mL (12 nM)	227.6 ng/mL (0.55 µM)	899 ng/mL (2.2 µM)
PFNA	0.3 ng/mL (0.5 nM)	1.0 ng/mL (2 nM)	N.A.	N.A.

Table 1. Human exposure levels of PFOS, PFOA, and PFNA.

Our preliminary data revealed an IC_{50} value of 189.2 nM for PFOS (Figure 1E), indicating that
human PFOS exposures (GP: 6-48 nM, EC: 80 nM, OE: 1.9 µM) may approach an at-risk range.
Moreover, because PFOS, PFOA, and PFNA share a conserved molecular target, their combined
presence in humans may interfere with folate transport during neurodevelopment through additive
or synergistic effects, even though individual serum levels are lower than their respective toxic
doses in zebrafish. This possibility raises the need to consider cumulative PFAS burden rather
than focusing on single compounds. In short, among the six PFAS species regulated by the EPA,
PFOS remains the most relevant to ongoing human exposure, yet PFOA and PFNA cannot be
overlooked given their shared folate receptor-binding potential and likelihood of additive
neurotoxicity. Investigating these compounds and other environmental PFAS species with folate
receptor-binding activities is therefore crucial for exposure mitigation, mechanistic understanding,
and health-risk assessment in vulnerable populations.

The selective behavioral and molecular effects observed among the tested PFAS suggest a
potential structure-activity relationship. PFNA, PFOA, and PFOS share longer carbon chain
lengths and greater hydrophobicity compared to PFBS, PFHxS, and GenX, properties that may
enhance protein binding affinity, membrane interactions, and bioaccumulation. These
physicochemical features could facilitate stronger or more sustained interactions with folate
receptors or associated transport machinery, contributing to their observed neurodevelopmental
effects. However, additional factors such as functional head groups, branching, and differential
tissue distribution may also play important roles and warrant systematic investigation in the future.

Recent studies have revealed a potential role of folic acid in ASD[22]. Low maternal folate levels
have been associated with a higher incidence of ASD in offspring[30-32]. A number of studies have
linked maternal folic acid supplementation with reduced risk of ASD in offspring[32-38]. For example,
maternal folic acid supplementation at 600 µg has been associated with a significantly reduced
risk of ASD in offspring[33], suggesting that adequate folate levels during pregnancy may be
beneficial for disease prevention. Specifically to PFAS exposure, a recent epidemiology study
based on the Shanghai Birth Cohort detected an association between prenatal PFOA exposure
and increased risk of autistic traits in the offspring, and found that the PFOA-induced autistic risk
was mitigated by pre-pregnancy folic acid supplementation[11], directly supporting our findings.
However, the dosage of folic acid supplementation appears to affect its protective effects, with
higher or lower levels of supplementation associated with increased risk of ASD[35].
Supplementation with folic acid and/or folinic acid in children diagnosed with ASD improved their
neurological and behavioral symptoms[39-43]. Several studies have reported an elevated prevalence
of serum auto-antibodies against folate receptor alpha in children with ASD[44-46]. Finally,
polymorphisms in the methylenetetrahydrofolate reductase (MTHFR) gene have been associated
with ASD risk[47-53]. These findings align with our observations that PFAS-induced folate pathway
disruption can lead to social behavioral deficits.

The folate pathway may contribute to ASD through several potential mechanisms. Folic acid is
required for DNA methylation, a critical process during brain development: it is a source of the
one-carbon group used to methylate DNA and required for the production of S-
adenosylmethionine, the primary methyl group donor for DNA methylation. Folic acid is also
essential for neural tube formation. It is worth noting that the optimal protective effects of folic acid
against ASD was observed when folic acid was taken during preconception and the first trimester
of pregnancy[33,38], the latter of which overlaps with the period of neural tube formation. Both DNA
methylation and neural tube formation are critical processes for brain development, pointing to
potential mechanisms of action for folic acid's involvement in ASD. Additionally, folic acid is
essential for the production of neurotransmitters that affect mood, motivation, stress, and
cognitive performance, including serotonin, norepinephrine, and dopamine[54,55], influencing neural
signaling pathways that are often disrupted in ASD.

Some studies indicate a possible link between elevated maternal folate levels and an increased risk of ASD, although findings are inconsistent and complex due to variations in individual metabolism, timing, and dosage of supplementation[56]. These findings suggest that while folic acid is crucial for brain development[19-21], there is a need for more nuanced guidance on optimal intake levels to balance benefits with potential risks, especially for populations at risk for ASD. This is aligned with our finding that while moderate to low doses of folic acid exerted beneficial rescue effects on social behavior when co-exposed to PFAS, higher doses showed significantly reduced rescue potencies.

Although prior work has utilized reverse molecular docking on selected proteins to predict binding targets of the environmental toxicant toluene[57], to our knowledge, this work represents the first proteome-wide reverse molecular docking in toxicology to discover *in vivo* protein targets for environmental toxicants. In reverse molecular docking, a small molecule drug or toxicant is virtually docked into the potential binding pockets of a large set of proteins. The binding characteristics of these interactions are then analyzed to rank the protein targets based on their binding affinities to the small molecules. This approach is different from traditional molecular docking which typically screens a large number of small molecules against a predetermined protein target, with the goal of identifying a ligand with high binding affinity to this protein. The successful application of proteome-wide reverse molecular docking in toxicology demonstrates a rapid and systematic *in silico* approach to predict protein targets for key environmental toxicants, which is expected to significantly promote the advance of mechanistic toxicology.

While our findings support a folate receptor-centered mechanism, alternative and complementary pathways may also contribute to PFAS-induced neurodevelopmental toxicity. PFAS are known to interact with multiple biological targets, including nuclear receptors such as PPARs[58-61], lipid metabolism pathways[62-66], thyroid hormone signaling[67-69], oxidative stress responses[70-72], and calcium homeostasis[73-75], all of which can influence brain development and behavior. Disruption of these pathways could act independently of, or synergistically with, folate pathway perturbations to shape neurodevelopmental outcomes. In addition, several limitations of the present study should be acknowledged. Although zebrafish provides a powerful vertebrate model for high-throughput behavioral and mechanistic studies, differences in PFAS toxicokinetics, placental transfer, and folate transport mechanisms between zebrafish and humans may influence dose-response relationships and translational interpretation. Moreover, our molecular docking analyses predict binding affinity but do not directly measure *in vivo* receptor occupancy or functional inhibition of folate transport, which will require future biochemical and genetic validation.

In summary, this study provides evidence that specific PFAS, notably PFNA, PFOA, and PFOS, may contribute to social behavioral deficits through their interaction with folate receptors, an effect mitigated by co-exposure to folic acid. Our findings align with existing evidence on the association between folate pathway disruptions and ASD risk, contributing to a deeper understanding of the biochemical mechanisms by which environmental toxicants may influence neurodevelopmental and disease outcomes. Furthermore, the ability of folic acid to rescue PFAS-induced deficits highlights the potential of folate supplementation to counteract PFAS toxicity in developing embryos. Together, these results underscore the need for further research into folate-based interventions and the long-term implications of exposure to various species of PFAS on neurodevelopmental disorders, particularly ASD. In addition, by incorporating the innovative proteome-wide reverse molecular docking approach, we provide a comprehensive framework for advancing the field of mechanistic toxicology through the identification of novel toxicological targets.

MATERIALS AND METHODS

Zebrafish husbandry

Zebrafish were housed at 26°C–27°C on a 14-hour light, 10-hour dark light cycle. Wild-type AB strain was used for all experiments. Nursery fish were fed twice a day at roughly 9 AM to 12 PM and 2 PM to 5 PM during the weekdays and once during weekends at no set time. Nursery fish were fed only rotifers from 5 dpf to 10 dpf, and were fed a mixture of rotifers and artemia between 10 dpf to 23 dpf. All zebrafish experiments were approved by the Institutional Animal Care and Use Committee at the University of Washington.

Chemical exposure

Fertilized embryos were sorted and transferred into 60 mm diameter and 20 mm deep glass petri dishes at 30 embryos per dish. Each dish was filled with 10 mL of HEPES-buffered E3 medium. Compound stocks were prepared in their specified solvents to maintain stability and stored at -20°C. Each compound was added to duplicate dishes (30 embryos per dish, two dishes per condition) at various concentrations, with negative controls receiving an equal volume of the corresponding solvent. Embryos were exposed without dechorionation. To prevent media contamination, dead embryos were removed at 1 and 2 days post-fertilization (dpf). At 3 dpf, all viable larvae were rinsed with E3 medium and transferred into clean petri dishes containing fresh E3 medium. At 5 dpf, larvae from each petri dish were transferred to separate nursery tanks at a final density of no more than 50 fish per tank and raised to 23 dpf for Fishbook assays.

The following per- and polyfluoroalkyl substances (PFAS) were obtained from Cayman Chemical (Ann Arbor, MI, USA): Perfluorooctanesulfonic Acid (PFOS, CAS# 1763-23-1, Item no. 37233), Perfluorooctanoic Acid (PFOA, CAS# 335-67-1, Item no. 37232), Perfluorononanoic Acid (PFNA, CAS# 375-95-1, Item no. 37250), Perfluorohexanesulfonic Acid (PFHxS, CAS# 355-46-4, Item no. 37248), Perfluorobutanesulfonic Acid (PFBS, CAS# 375-73-5, Item no. 37243), and Folic Acid (CAS# 59-30-3, Item no. 20515). Ammonium perfluoro(2-methyl-3-oxahexanoate) (GenX, CAS# 62037-80-3, Item no. C23990) was obtained from Sigma-Aldrich. PFOS came as a solution in 100% ethanol. PFBS came in liquid form. To prepare stock solutions, PFOA, PFNA, and folic acid were dissolved in HEPES-buffered E3 media, GenX and PFHxS were dissolved in dimethyl sulfoxide (DMSO).

Fishbook assay

The Fishbook assay was run as described previously[13]. Briefly, the test arena consists of a total of 44 3D printed, 10-mm-deep, 8.5-mm-wide, and 80-mm-long rectangular chambers grouped together in parallel with each other. Each chamber is divided into three compartments by two transparent acrylic windows (1.5 mm thick): a 60-mm-long middle testing chamber to place the test subject and two 8.5-mm-long end chambers to place the social stimulus fish or remain empty, respectively. Test subjects (PFAS treated 23 dpf fish or controls) were each placed inside an individual test chamber using a plastic transfer pipette with its tip cut off to widen the opening. A 3D printed white comb-like structure was placed in front of the social stimulus compartment to block the test subject's visual access to social stimulus fish before testing begins. After test subjects were placed inside the chambers, the arena was placed inside the imaging station, and the combs were removed to visually expose the social stimulus fish to the test subjects. Following a brief acclimation period, a 10-min test session was video recorded.

Videos were streamed through the software Bonsai[76]. Videos were analyzed in real time during recording, and the frame-by-frame x and y coordinates of each fish relative to its own test compartment were exported as a CSV file. Data were analyzed using custom Python scripts to calculate social scores and generate tracking plots. Social score was defined as a fish's average y-axis position for all frames. The middle of each test chamber was designated as the origin of the y axis, with an assigned value of zero. A value of 1 was assigned to the end of the chamber

next to the social stimulus fish and a value of −1 to the other end of the chamber next to the empty
control compartment. In this coordinate system, all social scores have values between −1 and 1.
A higher social score demonstrates a shorter average distance between a test subject and a social
stimulus fish during a test, which suggests a stronger social preference.

CRISPR-Cas9 gene knockout

We designed 3 sets of sgRNAs targeting the zebrafish *folr* gene to maximize efficiency of F0
knockout[26-28]. CHOPCHOP[77] was used for the sgRNA design. The *folr* sgRNA sequences
designed are as follows: gRNA1
"ATTTAGGTGACACTATATGAGGGACAGTTATATCAGCGTTTTAGAGCTAGAAATAGCAAG",
gRNA2
"ATTTAGGTGACACTATATGTATCCAGGGCCCCAGGTGGTTTTAGAGCTAGAAATAGCAAG",
and gRNA3
"ATTTAGGTGACACTATAGTCCATGTTCTGCTCAGGTGGTTTTAGAGCTAGAAATAGCAAG".
sgRNAs targeting early coding exons were selected to introduce nonsense mutations as early as
possible and increase likelihood of loss-of-function mutations. The SP6 promoter sequence was
added before the target RNA sequence, followed by an overlap adapter, which is complementary
to the 5' end of an 80 bp constant oligo according to the published protocol[78]. Oligonucleotides
were synthesized by Eurofins Scientific. Double-stranded DNAs were generated by Phusion Hot-
Start Flex DNA Polymerase (New England Biolabs) using the gene specific oligos and the
constant oligo. The double-stranded DNA was cleaned up using the Zymo Clean and
Concentrator-5 kit (Zymo Research). *In vitro* sgRNA transcription was conducted using the
MEGAscript SP6 transcription kit (Thermo Fisher Scientific) and cleaned up using the Zymo RNA
Clean and Concentrator-5 kit (Zymo Research).

To perform zebrafish embryonic microinjections, pairs of male and female adult zebrafish were
kept in a mating cage overnight while separated by a divider. 1-cell-stage embryos were collected
the next day early in the morning immediately before injection by pulling out the divider to allow
breeding. Injection solution was prepared by combining the three sgRNAs together with 2 µM Spy
Cas9 NLS and 1X NEBuffer (New England Biolabs). All three gRNAs were injected together to
induce multiple point mutations in the same gene. Approximately 1 nL of sgRNA–Cas9 RNP
complex was injected into each 1-cell-stage embryo using a pneumatic microinjector (Parker
Instrumentation). Approximately 300 embryos were injected with an estimated 60% survival rate
24 hours post-injection. To verify successful knockout of *folr*, we extracted genomic DNA from
injected embryos and amplified the gRNA-targeted genomic regions using PCR. The following
forward and reverse primers were used: sgRNA1 forward primer:
GTGTGATGTGTCTGTGTTGTGC; sgRNA1 reverse primer: TGTGTGTTGCAGTGAAGTGTGT;
sgRNA2 forward primer: AAGAAGCGCATCAGGATAACTC; sgRNA2 reverse primer:
CTGATGTTTAAAACTGGTGGCA; sgRNA3 forward primer: CAACGTGATTGCCTCTATGAAA;
sgRNA3 reverse primer: AAACTCAGAAGCATGTGGTGTG. PCR was conducted in a Bio-Rad
thermal cycler using a 20µL reaction volume containing 10µL of Taq 2X Master Mix (NEB) and
0.4µL of each primer. The following program was used: a 3 minute initial denaturation at 95°C,
followed by 39 cycles of 15 sec at 95°C for denaturation, 30 sec at 54 °C for annealing, and 30
sec at 68 °C for extension, and a final extension step at 68 °C for 5 min. PCR products were
examined by electrophoresis at 100 V for 30 minutes in a 1.5% (w/v) agarose gel in 1 x TAE
buffer. The gel was imaged on a UV light box at 365 nm. Lack of amplification products or smeared
amplification products indicate successful knockout (Supplementary Figure 5).

Reverse molecular docking screen and molecular docking validation

All *in silico* calculations were performed using a Digital Storm workstation with AMD Ryzen
Threadripper PRO 5955WX (16-Core) processor (Up to 4.0 GHz), 128GB DDR4 memory, and

NVIDIA RTX A2000 6GB graphics card. Both reverse molecular docking screen and molecular
docking validation were conducted using Autodock Vina[79] and Open Babel[80]. Ligand structures
were downloaded from PubChem as sdf files and converted to pdbqt. Binding pocket predictions
were acquired from the HPRoteome-BSite database[16] which covers over 80% of the UniProt
entries in the AlphaFold human proteome database[17,18]. In the HPRoteome-BSite database, a
binding pocket was predicted for a single domain for most of the proteins, whereas a subset of
proteins had a binding pocket predicted for multiple domains. For each domain, pockets were
predicted based on either the peptide sequence or the 3D structure of the domain; both types of
predictions were used for the screen. The highest-ranking sequence-based predictions and the
highest-ranking 3D-structure-based pocket predictions were respectively selected for each
protein domain, resulting in a total of 33446 predicted pockets selected for the screen. The domain
structures were downloaded from the HPRoteome-BSite database and prepared for molecular
docking by removing all water molecules, ions, and other substructures. The docking coordinates
of each predicted binding pocket were provided by the HPRoteome-BSite database. Pocket size
of 60 × 60 × 60, energy range of 3, number of modes of 9, and an exhaustiveness of 8 were
employed for the molecular docking screen and validation. Each pocket was screened once for
each PFAS.

Protein-protein interaction analysis

514 Protein-protein interaction (PPI) analysis was conducted using STRING[81] (https://string-db.org/).

Gene enrichment analysis

Molecular targets with a binding energy of less than -10 kcal/mol from the reverse molecular
binding virtual screen were selected for gene enrichment analysis. Gene Ontology (GO), and
Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analysis were conducted using
the clusterProfiler package in R (version 4.12.5). The GO analysis included Biological Process
(BP), Cellular Component (CC), and Molecular Function (MF) categories, with the org.Hs.eg.db
database used for annotation (v3.19.1). Results were ranked based on RichFactor and adjusted
p-value. Adjusted p-value was calculated using the Benjamini-Hochberg (BH) method with a cutoff
of 0.05 for significance. RichFactor is defined as the ratio of input genes that are annotated under
a certain term to all genes annotated under this term. Results were visualized as bar plots using
the ggplot2 package (v3.5.1).

Proteome integral solubility alteration (PISA) assay

His-tagged recombinant human folate receptor beta (FR-β) protein (ProSpec, PRO-2673, amino
acids 22-230) and mouse FR-β protein (Antibodies-Online, ABIN7274670, amino acids 21-227)
were reconstituted in distilled water at a concentration of 500 µg/mL. The proteins were then
diluted to 50 µg/mL and incubated with 10 µM PFOS, 400 µM PFNA, 1600 µM PFOA, 12.5 µM
folic acid, or an equal volume of vehicle control (DMSO or ethanol) for 30 minutes at room
temperature on a rotator. All compounds were prepared as 100x stock solutions compared to their
final concentration. Each compound-treated sample was aliquoted into 9 PCR tubes and
subjected to a 5 min temperature gradient treatment (from 34°C to 66°C, in 4°C increments) using
a Bio-Rad Thermocycler. After temperature treatment, the samples were placed on ice for at least
5 minutes. Contents of the 9 PCR tubes were pooled into a new 1.5 mL tube and centrifuged at
15,000g for 30 minutes at 4°C. The supernatant was transferred to a new 1.5 mL tube. For dot
blot analysis, 2 µL of supernatant was applied onto a nitrocellulose membrane, and the procedure
was repeated 10 times for each sample (10 technical replicates). Each membrane was air-dried
and blocked for 30 minutes in Tris-buffered saline with Tween 20 (TBST) containing 2% bovine
serum albumin (BSA). The membrane was then incubated with an anti-His Tag Antibody
(SinoBiological, 105327-MM02T) diluted 1:1000 in TBST containing 0.1% BSA and 0.1% sodium
azide for 30 minutes. Following primary antibody incubation, the membrane was washed 3 × 5

min by TBST, and incubated with horseradish peroxidase (HRP)-conjugated secondary antibody (Jackson ImmunoResearch) diluted 1:10,000 in TBST containing 0.1% BSA for 30 minutes. After washing in TBST once for 15 minutes and twice for 5 minutes, the membrane was washed once in TBS for 5 minutes, and stored in TBS at 4°C if needed. For imaging, each membrane was exposed to ECL substrate (Bio-Rad) and scanned using the iBright Imaging System (Invitrogen) with chemiluminescence smart exposure. Quantification of the dot blot image was performed using FIJI[82].

Statistical analysis

Graphs were generated using GraphPad Prism. Two-tailed Student's *t* test was used to analyze data when there were two groups. For analysis of multiple groups, normal distribution of datasets was confirmed using Shapiro-Wilk test, and if more than half the data was normally distributed and standard deviations fell within the variance ratio, one-way analysis of variance (ANOVA) assuming gaussian distribution and equal standard deviation was performed, using Dunnett's multiple comparison test. *P* values less than 0.05 were considered significant.

ACKNOWLEDGEMENTS

We thank the University of Washington Office of Comparative Medicine for providing zebrafish husbandry support. This work was supported by the National Institute of Environmental Health Sciences (NIEHS) of the NIH under the award number R00ES031050. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

AUTHOR CONTRIBUTIONS

M.J. and A.X.K. designed and conducted the experiments and analyzed data. A.E. and K.P assisted in conducting experiments. P.Z. performed pathway enrichment analysis. Y.G. conceived the study and interpreted the data. M.J., A.X.K., and Y.G. wrote the manuscript. All authors contributed meaningful insights during discussions and reviewed and approved the final version of the manuscript.

COMPETING INTERESTS

The authors declare that they have no competing interests.

DATA AND MATERIALS AVAILABILITY

All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials.

REFERENCES

1   Hansen, S. N., Schendel, D. E. & Parner, E. T. Explaining the increase in the prevalence of autism spectrum disorders: the proportion attributable to changes in reporting practices. *JAMA Pediatr* **169**, 56–62 (2015).
2   King, M. & Bearman, P. Diagnostic change and the increased prevalence of autism. *Int J Epidemiol* **38**, 1224–1234 (2009). PMC2800781.

3   Muir, D. C. G., Getzinger, G. J., McBride, M. & Ferguson, P. L. How Many Chemicals in
Commerce Have Been Analyzed in Environmental Media? A 50 Year Bibliometric Analysis.
Environ Sci Technol **57**, 9119–9129 (2023). PMC10308830.

4   Gaugler, T. *et al.* Most genetic risk for autism resides with common variation. *Nat Genet*
46, 881–885 (2014). PMC4137411.

5   Sandin, S. *et al.* The familial risk of autism. *JAMA* **311**, 1770–1777 (2014). PMC4381277.

6   Ames, J. L. *et al.* Prenatal Exposure to Per- and Polyfluoroalkyl Substances and Childhood
Autism-related Outcomes. *Epidemiology* **34**, 450–459 (2023). PMC10074577.

7   Oh, J. *et al.* Prenatal exposure to per- and polyfluoroalkyl substances in association with
autism spectrum disorder in the MARBLES study. *Environ Int* **147**, 106328 (2021).
PMC7856021.

8   Fenton, S. E. *et al.* Per- and Polyfluoroalkyl Substance Toxicity and Human Health Review:
Current State of Knowledge and Strategies for Informing Future Research. *Environ Toxicol*
Chem **40**, 606–630 (2021). PMC7906952.

9   Patlewicz, G. *et al.* Development of chemical categories for per- and polyfluoroalkyl
substances (PFAS) and the proof-of-concept approach to the identification of potential
candidates for tiered toxicological testing and human health assessment. *Comput Toxicol*
31, 100327 (2024). PMC12181936.

10   Schymanski, E. L. *et al.* Per- and Polyfluoroalkyl Substances (PFAS) in PubChem: 7
Million and Growing. *Environ Sci Technol* **57**, 16918–16928 (2023). PMC10634333.

11   Huang, Y. *et al.* Prenatal exposure to per- and polyfluoroalkyl substances, genetic factors,
and autistic traits: Evidence from the Shanghai birth cohort. *J Hazard Mater* **480**, 135857
(2024).

12   Geng, Y. & Peterson, R. T. The zebrafish subcortical social brain as a model for studying
social behavior disorders. *Dis Model Mech* **12** (2019). PMC6737945.

13   Geng, Y. *et al.* Top2a promotes the development of social behavior via PRC2 and
H3K27me3. *Sci Adv* **8**, eabm7069 (2022). PMC9683714.

14   Geng, Y., Yates, C. & Peterson, R. T. Social behavioral profiling by unsupervised deep
learning reveals a stimulative effect of dopamine D3 agonists on zebrafish sociality. *Cell*
Rep Methods **3**, 100381 (2023). PMC9939379.

15   Diaz, L. *et al.* Butyrate rescues chlorpyrifos-induced social deficits through inhibition of
class I histone deacetylases. *bioRxiv* (2025).

16   Sim, J., Kwon, S. & Seok, C. HProteome-BSite: predicted binding sites and ligands in
human 3D proteome. *Nucleic Acids Res* **51**, D403–D408 (2023). PMC9825455.

17   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583–589 (2021). PMC8371605.

18   Varadi, M. *et al.* AlphaFold Protein Structure Database: massively expanding the structural
coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res* **50**,
D439–D444 (2022). PMC8728224.

19   Balashova, O. A., Visina, O. & Borodinsky, L. N. Folate action in nervous system
development and disease. *Dev Neurobiol* **78**, 391–402 (2018). PMC5867258.

20   Irvine, N., England-Mason, G., Field, C. J., Dewey, D. & Aghajafari, F. Prenatal Folate and
Choline Levels and Brain and Cognitive Development in Children: A Critical Narrative
Review. *Nutrients* **14** (2022). PMC8778665.

21   Naninck, E. F. G., Stijger, P. C. & Brouwer-Brolsma, E. M. The Importance of Maternal
Folate Status for Brain Development and Function of Offspring. *Adv Nutr* **10**, 502–519
(2019). PMC6520042.

22   Hoxha, B. *et al.* Folic Acid and Autism: A Systematic Review of the Current State of
Knowledge. *Cells* **10** (2021). PMC8394938.

23   Zhang, Y. *et al.* Folate concentrations and serum perfluoroalkyl and polyfluoroalkyl
substance concentrations in adolescents and adults in the USA (National Health and

Nutrition Examination Study 2003-16): an observational study. *Lancet Planet Health* **7**, e449–e458 (2023). PMC10901144.

Gaetani, M. *et al.* Proteome Integral Solubility Alteration: A High-Throughput Proteomics Assay for Target Deconvolution. *J Proteome Res* **18**, 4027–4037 (2019).

Buglo, E. *et al.* Genetic compensation in a stable slc25a46 mutant zebrafish: A case for using F0 CRISPR mutagenesis to study phenotypes caused by inherited disease. *PLoS One* **15**, e0230566 (2020). PMC7092968.

Kroll, F. *et al.* A simple and effective F0 knockout method for rapid screening of behaviour and other complex phenotypes. *Elife* **10** (2021). PMC7793621.

Kim, B. H. & Zhang, G. Generating Stable Knockout Zebrafish Lines by Deleting Large Chromosomal Fragments Using Multiple gRNAs. *G3 (Bethesda)* **10**, 1029–1037 (2020). PMC7056962.

Wu, R. S. *et al.* A Rapid Method for Directed Gene Knockout for Screening in G0 Zebrafish. *Dev Cell* **46**, 112–125 e114 (2018).

Olsen, G. W. *et al.* An occupational exposure assessment of a perfluorooctanesulfonyl fluoride production site: biomonitoring. *AIHA J (Fairfax, Va)* **64**, 651–659 (2003).

Gao, Y. *et al.* New Perspective on Impact of Folic Acid Supplementation during Pregnancy on Neurodevelopment/Autism in the Offspring Children - A Systematic Review. *PLoS One* **11**, e0165626 (2016). PMC5119728.

DeVilbiss, E. A., Gardner, R. M., Newschaffer, C. J. & Lee, B. K. Maternal folate status as a risk factor for autism spectrum disorders: a review of existing evidence. *Br J Nutr* **114**, 663–672 (2015).

Steenweg-de Graaff, J., Ghassabian, A., Jaddoe, V. W., Tiemeier, H. & Roza, S. J. Folate concentrations during pregnancy and autistic traits in the offspring. The Generation R Study. *Eur J Public Health* **25**, 431–433 (2015).

Schmidt, R. J., Iosif, A. M., Guerrero Angel, E. & Ozonoff, S. Association of Maternal Prenatal Vitamin Use With Risk for Autism Spectrum Disorder Recurrence in Young Siblings. *JAMA Psychiatry* **76**, 391–398 (2019). PMC6450282.

Levine, S. Z. *et al.* Association of Maternal Use of Folic Acid and Multivitamin Supplements in the Periods Before and During Pregnancy With the Risk of Autism Spectrum Disorder in Offspring. *JAMA Psychiatry* **75**, 176–184 (2018). PMC5838577.

Raghavan, R. *et al.* Maternal Multivitamin Intake, Plasma Folate and Vitamin B(12) Levels and Autism Spectrum Disorder Risk in Offspring. *Paediatr Perinat Epidemiol* **32**, 100–111 (2018). PMC5796848.

DeVilbiss, E. A. *et al.* Antenatal nutritional supplementation and autism spectrum disorders in the Stockholm youth cohort: population based cohort study. *BMJ* **359**, j4273 (2017). PMC6168830.

Nilsen, R. M. *et al.* Analysis of self-selection bias in a population-based cohort study of autism spectrum disorders. *Paediatr Perinat Epidemiol* **27**, 553–563 (2013). PMC3851582.

Suren, P. *et al.* Association between maternal use of folic acid supplements and risk of autism spectrum disorders in children. *JAMA* **309**, 570–577 (2013). PMC3908544.

Adams, J. B. *et al.* Comprehensive Nutritional and Dietary Intervention for Autism Spectrum Disorder-A Randomized, Controlled 12-Month Trial. *Nutrients* **10** (2018). PMC5872787.

James, S. J. *et al.* Efficacy of methylcobalamin and folinic acid treatment on glutathione redox status in children with autism. *Am J Clin Nutr* **89**, 425–430 (2009). PMC2647708.

Kaluzna-Czaplinska, J., Michalska, M. & Rynkowski, J. Vitamin supplementation reduces the level of homocysteine in the urine of autistic children. *Nutr Res* **31**, 318–321 (2011).

Moretti, P. *et al.* Cerebral folate deficiency with developmental delay, autism, and response to folinic acid. *Neurology* **64**, 1088–1090 (2005).

43   Sun, C., Zou, M., Zhao, D., Xia, W. & Wu, L. Efficacy of Folic Acid Supplementation in
Autistic Children Participating in Structured Teaching: An Open-Label Trial. *Nutrients* **8**
(2016). PMC4924178.
44   Zhou, J. *et al.* High prevalence of serum folate receptor autoantibodies in children with
autism spectrum disorders. *Biomarkers* **23**, 622–624 (2018).
45   Frye, R. E. *et al.* Folinic acid improves verbal communication in children with autism and
language impairment: a randomized double-blind placebo-controlled trial. *Mol Psychiatry*
23, 247–256 (2018). PMC5794882.
46   Ramaekers, V. T., Blau, N., Sequeira, J. M., Nassogne, M. C. & Quadros, E. V. Folate
receptor autoimmunity and cerebral folate deficiency in low-functioning autism with
neurological deficits. *Neuropediatrics* **38**, 276–281 (2007).
47   Ismail, S. *et al.* Study of C677T variant of methylene tetrahydrofolate reductase gene in
autistic spectrum disorder Egyptian children. *Am J Med Genet B Neuropsychiatr Genet*
180, 305–309 (2019).
48   El-Baz, F., El-Aal, M. A., Kamal, T. M., Sadek, A. A. & Othman, A. A. Study of the C677T
and 1298AC polymorphic genotypes of MTHFR Gene in autism spectrum disorder.
Electron Physician **9**, 5287–5293 (2017). PMC5633227.
49   Shaik Mohammad, N. *et al.* Clinical utility of folate pathway genetic polymorphisms in the
diagnosis of autism spectrum disorders. *Psychiatr Genet* **26**, 281–286 (2016).
50   Guo, T., Chen, H., Liu, B., Ji, W. & Yang, C. Methylenetetrahydrofolate reductase
polymorphisms C677T and risk of autism in the Chinese Han population. *Genet Test Mol*
Biomarkers **16**, 968–973 (2012).
51   James, S. J. *et al.* A functional polymorphism in the reduced folate carrier gene and DNA
hypomethylation in mothers of children with autism. *Am J Med Genet B Neuropsychiatr*
Genet **153B**, 1209–1220 (2010). PMC2943349.
52   Goin-Kochel, R. P. *et al.* The MTHFR 677C-->T polymorphism and behaviors in children
with autism: exploratory genotype-phenotype correlations. *Autism Res* **2**, 98–108 (2009).
53   Mohammad, N. S. *et al.* Aberrations in folate metabolic pathway and altered susceptibility
to autism. *Psychiatr Genet* **19**, 171–176 (2009).
54   Zhou, Y., Cong, Y. & Liu, H. Folic acid ameliorates depression-like behaviour in a rat model
of chronic unpredictable mild stress. *BMC Neurosci* **21**, 1 (2020). PMC6961331.
55   Miller, A. L. The methylation, neurotransmitter, and antioxidant connections between folate
and depression. *Altern Med Rev* **13**, 216–226 (2008).
56   Wiens, D. & DeSoto, M. C. Is High Folic Acid Intake a Risk Factor for Autism?-A Review.
Brain Sci **7** (2017). PMC5704156.
57   McCarthy, M. J., Chushak, Y. & Gearhart, J. M. Reverse molecular docking and deep-
learning to make predictions of receptor activity for neurotoxicology. *Comput Toxicol* **24**
(2022).
58   Evans, N. *et al.* In vitro activity of a panel of per- and polyfluoroalkyl substances (PFAS),
fatty acids, and pharmaceuticals in peroxisome proliferator-activated receptor (PPAR)
alpha, PPAR gamma, and estrogen receptor assays. *Toxicol Appl Pharmacol* **449**, 116136
(2022). PMC9341220.
59   Khazaee, M. *et al.* Perfluoroalkyl Acid Binding with Peroxisome Proliferator-Activated
Receptors alpha, gamma, and delta, and Fatty Acid Binding Proteins by Equilibrium
Dialysis with a Comparison of Methods. *Toxics* **9** (2021). PMC7996760.
60   Takacs, M. L. & Abbott, B. D. Activation of mouse and human peroxisome proliferator-
activated receptors (alpha, beta/delta, gamma) by perfluorooctanoic acid and
perfluorooctane sulfonate. *Toxicol Sci* **95**, 108–117 (2007).
61   Vanden Heuvel, J. P., Thompson, J. T., Frame, S. R. & Gillies, P. J. Differential activation
of nuclear receptors by perfluorinated fatty acid analogs and natural fatty acids: a
comparison of human, mouse, and rat peroxisome proliferator-activated receptor-alpha, -

beta, and -gamma, liver X receptor-beta, and retinoid X receptor-alpha. *Toxicol Sci* **92**,
476–489 (2006).

62   Birchfield, A. S., Musayev, F. N., Castillo, A. J., Zorn, G. & Fuglestad, B. Broad PFAS
binding with fatty acid binding protein 4 is enabled by variable binding modes. *bioRxiv*
(2025). PMC11974712.

63   Fan, X. *et al.* Binding of per- and polyfluoroalkyl substances with liver and serum proteins
in rats: implications for physiologically based pharmacokinetic modelling. *Environ Int* **201**,
109591 (2025).

64   Haug, M., Dunder, L., Lind, P. M., Lind, L. & Salihovic, S. Associations of perfluoroalkyl
substances (PFAS) with lipid and lipoprotein profiles. *J Expo Sci Environ Epidemiol* **33**,
757–765 (2023). PMC10541331.

65   Ji, C., Pavuk, M. & Ruiz, P. Global Serum Per- and Polyfluoroalkyl Substances Exposures
and Their Correlation with Lipids: A Systematic Review and Meta-Analysis. *Environ Sci*
Technol **59**, 19611–19629 (2025).

66   Kashobwe, L. *et al.* Legacy and alternative per- and polyfluoroalkyl substances (PFAS)
alter the lipid profile of HepaRG cells. *Toxicology* **506**, 153862 (2024).

67   Coperchini, F. *et al.* Thyroid Disrupting Effects of Old and New Generation PFAS. *Front*
Endocrinol (Lausanne) **11**, 612320 (2020). PMC7851056.

68   Du, X. *et al.* Association between PFAS exposure and thyroid health: A systematic review
and meta-analysis for adolescents, pregnant women, adults and toxicological evidence.
Sci Total Environ **953**, 175958 (2024).

69   Noyes, T. S. *et al.* Per and polyfluoroalkyl substances affect thyroid hormones for people
with a history of exposure from drinking water. *Sci Rep* **15**, 12502 (2025). PMC11992217.

70   Solan, M. E., Koperski, C. P., Senthilkumar, S. & Lavado, R. Short-chain per- and
polyfluoroalkyl substances (PFAS) effects on oxidative stress biomarkers in human liver,
kidney, muscle, and microglia cell lines. *Environ Res* **223**, 115424 (2023).

71   Taibl, K. R. *et al.* Associations of per- and polyfluoroalkyl substances (PFAS) and their
mixture with oxidative stress biomarkers during pregnancy. *Environ Int* **169**, 107541 (2022).
PMC9846434.

72   Wielsoe, M., Long, M., Ghisari, M. & Bonefeld-Jorgensen, E. C. Perfluoroalkylated
substances (PFAS) affect oxidative stress biomarkers in vitro. *Chemosphere* **129**, 239–
245 (2015).

73   Brown-Leung, J. M. & Cannon, J. R. Neurotransmission Targets of Per- and
Polyfluoroalkyl Substance Neurotoxicity: Mechanisms and Potential Implications for
Adverse Neurological Outcomes. *Chem Res Toxicol* **35**, 1312–1333 (2022).
PMC10446502.

74   Di Nisio, A. *et al.* Association of perfluoroalkyl substance (PFAS) on vitamin D biomarkers
in a highly exposed population of the Veneto Region in Italy. *Chemosphere* **374**, 144230
(2025).

75   Starnes, H. M., Rock, K. D., Jackson, T. W. & Belcher, S. M. A Critical Review and Meta-
Analysis of Impacts of Per- and Polyfluorinated Substances on the Brain and Behavior.
Front Toxicol **4**, 881584 (2022). PMC9035516.

76   Lopes, G. *et al.* Bonsai: an event-based framework for processing and controlling data
streams. *Front Neuroinform* **9**, 7 (2015). PMC4389726.

77   Montague, T. G., Cruz, J. M., Gagnon, J. A., Church, G. M. & Valen, E. CHOPCHOP: a
CRISPR/Cas9 and TALEN web tool for genome editing. *Nucleic Acids Res* **42**, W401–407
(2014). PMC4086086.

78   Gagnon, J. A. *et al.* Efficient mutagenesis by Cas9 protein-mediated oligonucleotide
insertion and large-scale assessment of single-guide RNAs. *PLoS One* **9**, e98186 (2014).
PMC4038517.

79   Trott, O. & Olson, A. J. AutoDock Vina: improving the speed and accuracy of docking with
a new scoring function, efficient optimization, and multithreading. *J Comput Chem* **31**,
455–461 (2010). PMC3041641.
80   O'Boyle, N. M. *et al.* Open Babel: An open chemical toolbox. *J Cheminform* **3**, 33 (2011).
PMC3198950.
81   Szklarczyk, D. *et al.* STRING v11: protein-protein association networks with increased
coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic*
Acids Res **47**, D607–D613 (2019). PMC6323986.
82   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**, 676–682 (2012). PMC3855844.

797