Structures of vertebrate R2 retrotransposon complexes during target-primed reverse transcription and after second strand nicking

Akanksha Thawani[1,2,†,*], Anthony Rodríguez-Vargas[2,†], Briana Van Treeck[2], Nozhat T Hassan[3], David L Adelson[3], Eva Nogales[1,2,4,5,*] and Kathleen Collins[1,2,*]

[1]California Institute for Quantitative Biosciences (QB3), Berkeley, CA, USA
[2]Department of Molecular and Cell Biology, University of California Berkeley, Berkeley, CA, USA
[3]School of Biological Sciences, University of Adelaide, Adelaide, Australia
[4]Howard Hughes Medical Institute, Chevy Chase, MD 20815, USA
[5]Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

[†]These authors contributed equally: Akanksha Thawani, Anthony Rodríguez-Vargas

[*]Corresponding authors. Email: athawani@berkeley.edu, enogales@lbl.gov, kcollins@berkeley.edu

Abstract

R2 retrotransposons are model site-specific eukaryotic non-LTR retrotransposons that copy-and-paste into gene loci encoding ribosomal RNAs. Recently we demonstrated that avian A-clade R2 proteins achieve efficient and precise insertion of transgenes into their native safe-harbor loci in human cells. The features of A-clade R2 proteins that support gene insertion are not characterized. Here, we report high resolution cryo-electron microscopy structures of two vertebrate A-clade R2 proteins, avian and testudine, at the initiation of target-primed reverse transcription and one structure after cDNA synthesis and second strand nicking. Using biochemical and cellular assays we discover the basis for high selectivity of template use and unique roles for each of the expanded A-clade zinc-finger domains in nucleic acid recognition. Reverse transcriptase active site architecture is reinforced by an unanticipated insertion motif in vertebrate A-clade R2 proteins. Our work brings first insights to A-clade R2 protein structure during gene insertion and enables further improvement and adaptation of R2-based systems for precise transgene insertion.

Introduction

Non-long terminal repeat (non-LTR) retrotransposons are mobile genetic elements that are widespread in eukaryotic species. Retrotransposon-derived DNA expression, mobilization, and rearrangement are recognized as major drivers of genome evolution and expansion (*1–3*). In mammals, retrotransposons have expanded via a copy-and-paste mechanism to compose a large portion of genomes. For example, nearly one-third of the human genome originated in the activity of the non-LTR retrotransposon Long Interspersed Element 1 (LINE-1), whose specialized insertion preference for DNA architecture is linked to replication fork progression with a degenerate DNA sequence recognition (*4–6*). The abundant cDNA-derived genome content shapes nuclear organization, chromatin landscape, and transcription of genes and regulatory RNAs (*3, 7–10*).

Other non-LTR retrotransposons are more target site selective (*11, 12*). R2 retrotransposons with sequence specificity for insertion to the tandemly repeated ribosomal RNA (rRNA) gene locus (the rDNA) are found within the genomes of multicellular animals including insects, crustaceans and non-mammalian vertebrates (*13, 14*). R2 protein (R2p) from a moth *Bombyx mori*, hereafter R2Bm, has long been the model system for biochemical characterization of target-primed reverse transcription (TPRT), where nicking of one of the two strands of the target site creates a primer for cDNA synthesis directly into the genome (*15, 16*) (Fig. 1a). R2p-mediated TPRT was recently re-purposed to insert transgenes into rDNA loci in cultured human cells (*17–20*). This technology, called precise RNA-mediated insertion of transgenes (PRINT), relies on an avian R2p translated from an engineered mRNA co-delivered with a second RNA that templates transgene synthesis (*17*).

The avian R2 retrotransposons belong to the A-clade, which among other clade-distinguishing differences has an expanded number of N-terminal zinc-finger domains (ZnFs) compared to D-clade R2Bm (*13*). Recent structural studies have revealed the architecture of R2Bm ribonucleoprotein (RNP) bound to duplex DNA or launched into TPRT (*21, 22*), but A-clade R2p remains under-characterized both biochemically and structurally. In particular, the role of the expanded array of ZnFs has not been elucidated, other than its significance for generating a more precise rDNA location of transgene 5′ junction formation with PRINT (*20*). Besides the N-terminal ZnFs, other distinct structural features are likely to exist due to the early divergence of A-clade and D-clade R2s (*13, 23*). Understanding the structural features and biochemical properties of A-clade vertebrate R2ps will enable rational engineering of these proteins for gene delivery and potential gene therapy applications. Further, while the initial TPRT stage has recently been characterized for R2Bm (*21*), subsequent stages, such as when cDNA synthesis for the first strand has completed and second strand nicking occurs, remain uncharted. In this work, using cryogenic electron microscopy (cryo-EM), we determine structures of A-clade avian (zebrafinch *Taeniopygia guttata*, R2Tg) and testudine (big-headed turtle *Platysternon megacephalum*, R2Pm) R2 RNPs with target site DNA. We also determine R2Pm protein domain configuration after completion of cDNA synthesis and second strand nicking, and we investigate the functional significance of A-clade-specific R2p structural features with biochemical and cellular assays.

Results

TPRT and PRINT activities of avian and testudine R2p

R2Tg and also R2p from the white-throated sparrow (*Zonotrichia albicollis*, R2Za) support PRINT (*17*). For comparison to avian R2p, we bioinformatically mined A-clade R2s from reptiles, which are the evolutionary predecessors of Aves (Fig. 1b). We found that testudine and avian R2 3′ untranslated regions (UTRs) have divergent primary sequence but share a possible pseudoknot-hinge-stem loop architecture at the 3′ end of their 3′UTR (Fig. S1a). We assayed the biochemical activities of bacterially expressed and purified R2Tg and R2Pm (Fig. S1b) in combination with 3 RNAs: the optimal avian R2 3′UTR (*17*, *19*) from the medium ground finch *Geospiza fortis* (292 nucleotide (nt) full-length Gf3 or the equally effective Gf98 containing the terminal 98 nt); R2Pm 3′UTR (210 nt full-length Pm3 or shortened Pm112), or R2Bm 3′UTR (248 nt full-length Bm3). Each 3′UTR sequence was followed by 5 nt of downstream rRNA (R5) that can base-pair with primer created by the first strand nick. R2Tg and R2Pm both efficiently used Gf98 and Pm112 RNA for TPRT *in vitro* (Fig. 1c). In competition assays using an RNA mixture for TPRT, both R2p had equal or greater preference for use of Gf98 (Fig. S1c). On the other hand, neither R2Tg nor R2Pm efficiently used Bm3 as a TPRT template (Fig. 1c), suggesting that like R2Tg (*14*), R2Pm has inherent RNA template recognition specificity.

To investigate R2Pm use of template RNA in cells, we tested PRINT efficiency with template RNAs that generate an autonomous GFP expression cassette, comprised of a modified CMV promoter, GFP ORF, and polyadenylation signal. Template RNAs have a 5′ module for RNA stability and a 3′ module with 3′UTR sequence followed by R4 and 22 adenosines (A22) for optimal PRINT (*17*, *19*, *24*). Template RNAs with either Gf3 or Pm3 in the 3′ module were delivered to human RPE-1 cells paired with an mRNA encoding R2Tg or R2Pm (Fig. 1d-e). Template RNA alone gave only background GFP signal (Fig. S1d). R2Tg paired with Gf3 template RNA generated 28% GFP-positive (GFP+) cells, whereas with Pm3 template RNA, only ~2% of cells were GFP+. R2Pm paired with Gf3 template RNA generated slightly less than 1% GFP+ cells, and as observed for R2Tg, the Pm3 template RNA was used with much lower efficiency (Fig. 1e, Fig. S1d). We conclude that although R2Tg has higher efficiency for transgene insertion than R2Pm, both proteins prefer PRINT template RNA with Gf3. We speculate that this reflects a more favorably homogeneous folding of Gf3 RNA, compared to Pm3 and the previously tested other avian R2 3′ UTRs that all share similar predicted secondary structure (*17*).

Structures of R2Tg and R2Pm during first strand synthesis

We sought to capture cryo-EM structures of A-clade R2p RNPs during TPRT. We used bacterially expressed and purified R2Pm and R2Tg and Gf3 or Gf98 RNA to assemble TPRT complexes by incubating the proteins with biotinylated rDNA target site duplex (Fig. S2a-b). We halted elongation after 1 nt of cDNA synthesis with dideoxythymidine triphosphate (ddTTP) and isolated complexes using a streptavidin-based pulldown strategy (Fig. S2b). All intended components of ternary complexes were present in the eluted samples, and both proteins had nicked the first strand and initiated cDNA synthesis (Fig. S2c-d). Cryo-EM structure determination for R2Pm with Gf3 in TPRT initiation stage reached an overall resolution of 3.2 Å (Fig. 1f, Fig. S2e, Fig. S3a and Fig. S4a-c). While initial attempts to determine high resolution cryo-EM structure of R2Tg with Gf3 RNA did not succeed due to low particle density, the particle density improved when we use the truncated Gf98 RNA (Fig. S2d, f), and we were able to obtain a structure of R2Tg RNP in the TPRT initiation stage at an overall resolution of 3.3 Å (Fig. 1g, Fig. S3b and Fig. S4a-c). The cryo-

EM density maps allowed us to model nearly the entire protein chain for R2Pm and R2Tg as well
as the upstream and downstream rDNA and an RNA pseudoknot-hinge-stem fold that forms an
extensive surface for protein interaction (Fig. 1h-i, Fig. 2a, Fig. S5a). We also resolved density for
ddTTP bound in the active site that is unable to join the cDNA 3′ end due to the incorporated
ddTTP (Fig. S5b).
The overall architectures of the A-clade R2p ternary complexes have both similarities and
differences with the D-clade R2Bm ternary complex captured at a similar stage of cDNA synthesis
(*21*). The shared R2p core domains include the reverse transcriptase (RT) fingers and palm motifs
(colored as RT domain) followed by the Thumb, a Linker, the C-terminal zinc-knuckle (ZnK), and
the restriction-like endonuclease domain (RLE) (Fig. 1f-i). As shown for R2Bm, the A-clade R2p
ZnK and RLE domains melt double-stranded DNA into single-stranded DNA across the first strand
nick site. Instead of the two NTEs (NTE 0 and -1) observed in the R2Bm structures (*21*, *22*), The
A-clade R2p RT core is preceded by three segments of N-terminal extension (NTE), two
previously recognized (NTE 0, -1) and a third (NTE -2) that was not described in the TPRT
initiation complex of R2Bm (*21*) or structures of bacterial retroelement proteins (*25*, *26*) (Fig.
S5c). NTE motifs are in turn preceded by an evolutionarily variable length of Spacer and the N-
terminal ZnF and Myb domains (Fig. 1f-i) that engage rDNA upstream of the first strand nick.
Large differences are present, however, in the architecture of A-clade versus D-clade R2p
interactions with RNA (see below) and in the shared and unique A-clade R2p ZnF contacts with
DNA and RNA that had not been predicted from previous biochemical assays (*20*, *27*). Overall,
our structures establish a divergence of A-clade and D-clade R2p nucleic acid interactions.
144
RNA recognition by ZnF3 and target site DNA
Of the 292 nt in Gf3 or 98 nt in Gf98, only the region within the 3′ 65 nt is visible in the cryo-EM
density map (Fig. 1f, h). The resolved regions of RNA correspond to a 5′ pseudoknot and a 3′ stem
connected by a 6 nt hinge (Fig. 2a, Fig. 1f-i). The 4 nt of single-stranded RNA between the 3′ stem
and the RNA paired to primer and ddTTP (Fig. S5b) were also resolved. We note that the fold and
topology of Gf3 engaged with the two A-clade R2p and of *B. mori* 3′UTR engaged with R2Bm
differ significantly, and there is more length of RNA density visible for Gf3 than was visible for
RNA bound to R2Bm, potentially due a more stabilized Gf3 3′ end RNA fold (Fig. S5d). The R2p
NTE -1, Linker and Thumb domains form a large surface for RNA recognition (Fig. 2a-b, Fig.
S5a, e). Key interactions include base-specific hydrogen bonds that Arg911 (R2Pm) or Arg960
(R2Tg) make with G-256, and Lys712 (R2Pm) or Lys763 (R2Tg) make with A-258 in the RNA
hinge (Fig. 2a-b, Fig. S5a, e). The sequence specific recognition of GGAAAAG motif in the hinge
and adjacent end of the pseudoknot is likely to contribute to the shared template selectivity of avian
and testudine R2p.
The A-clade R2p ZnF2 and ZnF3 fold together through a previously unanticipated
interaction of beta strands. This folding unit is sandwiched on target site DNA between ZnF1 and
RLE (described below) and bookends the RNA pseudoknot from the side opposite NTE -1 (Fig.
1h-i, Fig. 2a-b). ZnF3 contacts the pseudoknot with hydrogen bonding interactions to both
backbone and bases (Fig. 2c, d, Fig. S5c). Our structures also reveal that the rDNA target site itself
contributes to RNA recognition. We find that bases within the DNA region melted by R2p face
toward the pseudoknot. In both R2Pm and R2Tg structures, the base dA(-3) of the second strand
creates a sequence-specific hydrogen bond with the base of G-255 at the junction between the
pseudoknot and the hinge (Fig. 2e).
To assay the functional significance of the visualized RNA secondary structure and its

sequence, we made mutations in the pseudoknot and hinge regions and assessed change in PRINT
efficiency. Mutating the hinge base G-256 to A reduced PRINT efficiency and disrupting the
pseudoknot base-pairing via G-255 to A or C-254 to A mutation drastically reduced PRINT
efficiency (Fig. 2f). Further, restoring the pseudoknot base-pairing with compensatory mutations
(G-235>U, C-254>A) restored PRINT activity to a level comparable to the wild-type pseudoknot
sequence (Fig. 2f). Altogether, our structural and functional assays demonstrate that multiple
regions of the protein recognize and position template RNA, particularly the RNA pseudoknot and
the hinge sequence, during the initiation of TPRT.
177
Target site recognition by R2 N-terminal DNA binding domains
As also shown for R2Bm in a previous work (*21*), the A-clade R2p ZnK and RLE domains split
double-stranded DNA around the first strand nick site (Fig. 3a, Fig. S6a). The nicked first strand
upstream of the target site, including its 5′ end, remains buried within the ZnK and RLE domains
(Fig. 3a). As a second similarity with R2Bm, the R2Tg and R2Pm motif 6a within the RT domain
wedges into a distortion of the upstream target site DNA (Fig. 3b, Fig. 2a, Fig. S5a). Together, the
ZnF and Myb domains of A-clade R2p create an extended surface protecting the target site, using
the entirety of the 4 domains and also connecting amino acid segments between them (Fig. 3c). In
comparison, R2Bm ZnF and Myb domains occupy a much smaller surface of upstream target site,
even compared to the A-clade R2p ZnF1 and Myb domains alone (Fig. 3c). A-clade R2p ZnF2
and ZnF3 engage the target site close to the first strand nick site (Fig. 3c, Fig. 2a, Fig. S5a). ZnF2
makes sequence-specific contacts, whereas ZnF1 and ZnF3 predominantly make sequence non-
specific contacts with the phosphate backbone of the target DNA (Fig. 3d, Fig. S6b-c). In contrast,
R2Bm relies on the ZnF corresponding to the A-clade R2p ZnF1 for sequence-specific contacts
(*21*, *22*).
In previous work using R2Za (*20*), we found that deletion of ZnF2 and ZnF3 had minimal
impact on TPRT and reduced, but did not eliminate, PRINT (*20*), suggesting that the ZnF3-2
contacts to RNA and DNA can be lost without severe disruption of RNA and DNA binding
specificity. However, removal of ZnF3-2 strikingly decreased the positional precision of transgene
5′ junction formation from the rDNA side (*20*). Contacts between ZnF3 and the pseudoknot would
be removed by cDNA synthesis, but ZnF2 contacts to upstream target site could remain (explored
below). These contacts, potentially dynamic with continued cDNA synthesis, could influence
DNA positioning for second-strand nicking. In concurrence with this idea, a contribution of ZnF3-
2 to second strand nicking has been detected using purified proteins under some conditions (*20*).
However, future studies are necessary to explore the relationship between R2p's *in vitro* second
strand nicking and productive second strand nicking in cells.
A major difference between the R2Pm and R2Tg structures, in comparison with each other,
is the disposition of the Spacer, the region that connects the N-terminal DNA binding domains to
the NTE motifs (Fig. 1g-h). R2Tg has a Spacer of ~80 amino acids that could not be resolved in
our cryo-EM map, whereas R2Pm has a Spacer of only ~30 amino acids that we partially observe
in our structure as it makes contacts with the RT core (Fig. 3e). To investigate whether the
difference in Spacer length and/or the N-terminal DNA binding domains gives R2Pm its lower
PRINT efficiency than R2Tg, we used human cells to express chimeric R2Pm proteins with
segments swapped to have an avian R2p Spacer, ZnF3-2, or the entire N-terminal region before
the NTE motifs. Purified domain-chimera proteins had similar or slightly better TPRT activity
than wild-type R2Pm, but each of the domain-chimera proteins suffered a large loss of PRINT
efficiency (Fig. 3f-g). Curiously, R2Pm with the entire N-terminus of R2Tg had substantially

lower PRINT efficiency than R2Pm with the entire N-terminus of R2Za, which nonetheless
remained compromised for PRINT relative to wild-type R2Pm (Fig. 3g). Altogether, these results
demonstrate structural and functional divergence of the N-terminal nucleic acid binding domains
and Spacer within vertebrate R2 A-clade proteins to an extent that they are not exchangeable
modules of R2p domain architecture. This is suggestive of co-evolution of the catalytic domains
with the Spacers and with the DNA binding domains.
221
Vertebrate R2p expansion of the C-terminal Insertion
A structural feature specific to the two vertebrate A-clade R2p, relative to R2Bm, is a sequence
insertion (hereafter C-terminal insertion, CTI) that threads from after the Thumb to the RT fingers
and back (Fig. S7a, Fig. 4a-b). While this Linker sub-region in R2Bm has 11 amino acids
connecting two alpha helices, R2Tg and R2Pm have a much longer 44 or 46 amino acids,
respectively (Fig. S7a). The CTI anchors to the RT domain with an EWE amino acid triplet (Fig.
4a-b). The R2Pm CTI has a short α-helix that is not present in the R2Tg CTI (Fig. 4a-b). Further,
while the entire R2Pm CTI could be easily visualized in the cryo-EM density, the density for the
part of the R2Tg CTI that is not facing the RNA-cDNA duplex is only visible at low density
thresholds.
To investigate the functional significance of the longer CTI in A-clade R2p, we truncated
the CTI in R2Tg and R2Pm to match the length of this region in R2Bm (ΔCTI mutants) with the
goal of deleting the intramolecular EWE anchor without changing the fold of adjacent regions
(Fig. S7a). This design was guided by AlphaFold3 (*28*). Wild-type and ΔCTI versions of R2Tg
and R2Pm were purified after bacterial expression and assayed for TPRT using Gf68, with the
minimized 68 nt of pseudoknot-hinge-stem loop sequence. Due to CTI positioning, we reasoned
that it could underlie the previously described avian R2p requirement for base-pairing of primer
with the template 3′ tail (*17*). We tested TPRT with Gf68 RNAs harboring different lengths of
downstream rRNA (Fig. 4c). In agreement with our previous findings, a 3′ tail of R4 but not R0 or
R3 supported TPRT activity of wild-type R2Tg, and the same specificity was observed for wild-
type R2Pm (Fig. 4c, lanes 1-3). Additionally increasing the homology length to R5, R8, or R12
had little if any influence on first-strand nicking or cDNA synthesis (Fig. 4c, compare lanes 4-7;
note that the adenosine present at the 3′ end of R8 inhibits template jumping). Curiously, CTI
truncation did not alter TPRT dependence on R4, but it did decrease unproductive first-strand
nicking when the template RNA 3′ tail was too short to support productive TPRT (Fig. 4c, lanes
8-13).
In contrast to reconstituted TPRT, PRINT by both R2Tg and R2Pm was severely inhibited
by CTI truncation (Fig. 4d). The percentage of full-length transgene insertions was not
proportionally reduced comparing wild-type and ΔCTI R2Tg proteins (Fig. S7b), suggesting that
the PRINT deficit is not caused by a substantially lowered processivity of cDNA synthesis in cells.
Altogether, our findings lead to the hypothesis that CTI expansion stabilized the active RT fold in
a manner critical for PRINT but not limiting for TPRT activity in reactions with purified protein.
In a recent study (*18*), the R2Tg CTI was assigned to be a disordered loop and used as a location
for insertion of accessory protein modules to optimize transgene insertion. Results from our assays
of R2p structure and function above recommend against CTI disruption, which we find to decrease
rather than increase transgene insertion efficiency.
258
Structure of R2Pm after cDNA synthesis and second strand nicking
Structural insight into a stage of the retrotransposon insertion process following initiation of TPRT

is lacking for any clade of R2. We first assayed whether R2Tg and R2Pm proteins had second
strand nicking activity dependent on the catalytic activity of the endonuclease domain. Second
strand nicking has been demonstrated for R2Bm and recently for two A-clade R2p (R2Za and R2p
from flour beetle *Tribolium castaneum*) but is weak compared to first strand nicking (*16*, *20*). We
designed a first strand pre-nicked target site DNA with different dye labels at the top and bottom
strand 5′ ends. We purified wild-type R2Tg and R2Pm proteins as well as RT or RLE active-site
mutants (RTD and END, respectively). When combined with the target site DNA and Gf68 RNA,
wild-type and RTD proteins, but not END proteins, nicked the second strand (Fig. 5a). Second
strand DNA nicking improved when the wild-type protein was able to perform first strand
synthesis upon addition of dNTPs (Fig. 5a). Based on denaturing PAGE migration of the cleavage
products, the position of second strand nicking in vitro is similar to the 2 bp offset from the first
strand nick detected for all other R2p assayed to date (*16*, *20*)..
For structure determination, we assembled R2Pm with nucleic acid substrates that mimic the
completion of first strand synthesis. The first strand cDNA was annealed to Gf68 with an R5 3′
tail (Fig. 5b, Fig. S8a). The template RNA had a single-nt 5′ overhang that a functional R2p
complex would use to complete cDNA synthesis. We added dideoxycytidine (ddCTP) to allow
cDNA synthesis completion and then purified complexes and analyzed their composition by
denaturing PAGE (Fig. S8b). Some complexes included an intact second strand, but complexes
with the second strand nicked were also evident (Fig. S8b). The cryo-EM density reconstructed
was for a complex with the second strand nicked. The cryo-EM structure of R2Pm after cDNA
synthesis and second strand nicking had an overall resolution of 4.6 Å (Fig. S8c, Fig. S9 and Fig.
5b-c). While some of the 2D class averages visualize the long RNA:cDNA duplex emerging from
the protein density, the full length of this duplex was not resolved in 3D reconstructions due to
flexibility (Fig. 5b-c).
Our structure revealed a configuration of R2p with the ZnF and Myb domains still bound to
upstream target site, as they were at the launch of TPRT. However, considering the entire length
of double-stranded DNA, change in its overall positioning was evident comparing R2Pm structures
at the start of first strand synthesis and after second strand nicking (Fig. 5d). The single-stranded
region of the upstream second strand could be traced towards the first-strand nick site until base
dA(-3), consistent with second strand nicking 2 bp upstream from the first strand nick (Fig. 5e).
Of note, upon second strand nicking, the 3′ end of the second strand moves into a position occupied
by the template RNA pseudoknot at the initiation of first strand synthesis, closer to ZnF3-2 and
NTE -1 (Fig. 5d-e). We propose that this positioning would enable R2p to protect the nicked
second strand 3′ end from exposure to DNA repair machinery.
295
296
297 **Discussion**
298
Structural adaptations in R2 evolution
In this study, we investigated the structural basis for steps in the site-specific insertion mechanism
of A-clade R2 retrotransposons, which are in a different clade from the best studied D-clade R2Bm
system due to an expanded array of N-terminal ZnFs (*12*, *14*). Observations from our cryo-EM
along with biochemical and cellular assays demonstrate that each of the three A-clade R2p ZnFs
have entirely different nucleic acid recognition principles and roles during gene insertion. We find
that these ZnFs, when assayed together in full-length protein context, occupy distinct positions
along the upstream rDNA target site. While two ZnFs, together with the Myb domain, bind an

extensive length of double-stranded target site DNA, the most N-terminal ZnF, ZnF3, interacts
primarily with a newly determined pseudoknot of 3′UTR RNA. Although the ZnF of R2Bm R2p,
which corresponds to A-clade ZnF1, has sequence-specific contacts with DNA, it is ZnF2 that has
these specific contacts in vertebrate A-clade R2p. The A-clade is believed to be more ancestral
than the D-clade (29), suggesting that loss of the most N-terminal A-clade ZnFs was accompanied
by gain of sequence-specific interaction by the solo D-clade ZnF. Loss of ZnF3-2 may have
enabled the D-clade Myb-ZnF DNA binding domains to develop sequence-specific interaction
with both downstream and upstream target site sequences (20, 30).
Our work highlights structural differences among the R2p studied at the biochemical level
to date, with differences both across clades and also among vertebrate A-clade R2p. Included
among these differences is the variable disorder of the Spacer bridging the N-terminal DNA
binding domains with the RT-RLE. Unexpectedly, the Spacer and DNA binding domains do not
appear to function as a module separable from the RT-RLE. Another particularly divergent
structural feature is the CTI. It is of high interest to investigate CTI sequence and structure across
a wider diversity of R2p and link this diversity to functional differences at the biochemical and
cellular levels.
323
R2 retrotransposition and PRINT
Our cryo-EM structure of an R2p complex after second strand nicking reveals that an A-clade R2p
remains bound to the upstream target site even after first strand synthesis and second strand
nicking. This would ensure close proximity and protection of the upstream and downstream sides
of an R2 insertion site during cDNA synthesis. accomplished by a single R2p retained at the site
of its initial recruitment. Repositioning of the nicked second strand 3′ end does not place it near
the RT active site; instead, the second strand 3′ end is in a protected position that it can occupy
after TPRT removes the initially bound pseudoknot-hinge-stem loop RNA. While R2p can make
an appropriately positioned second strand nick *in vitro*, questions of whether R2p makes the second
strand nick in cells, and if so whether this is mediated by the initially recruited R2p or by a second
R2p acting in concert, remain unresolved. As a correlation, deletion of ZnF3-2 inhibits second
strand nicking under some conditions *in vitro* and strongly decreases the fidelity of 5' junction
formation for transgenes inserted by PRINT (20). However, loss of fidelity in 5' junction formation
could also result from increased R2p dissociation from the upstream target site during cDNA
synthesis. Future studies are necessary to explore the mechanisms of second-strand nicking and
synthesis in cells.
As a working model, we propose that the persistent upstream binding of A-clade R2p
protects otherwise free DNA strand ends but does not launch second strand synthesis. The
expanded A-clade R2p ZnF-array recognition and protection of upstream target site DNA could
contribute to the favorable function of avian A-clade R2p in transgene insertion by PRINT.
However, as A-clade R2Tg and R2Pm have similar RNA binding specificities and similar DNA
binding domain configurations on the target site, yet differ strikingly in their ability to support
PRINT, factors inherent to the RT-RLE core of R2p are also relevant for efficient PRINT. To
develop a site-programmable transgene insertion technology that exploits efficient R2p TPRT in
human cells, one possibility would be to replace or supplement the ZnF array with heterologous
sequence-specific DNA binding domains, adopting a design principle from zinc-finger nucleases
and transcription activator-like effector nucleases (31, 32). Yet, this is unlikely to be
straightforward given the lack of domain modularity evident in the deleterious Spacer and DNA
binding domain chimeras assayed to date. In combination with extensive target site DNA

recognition, the high specificity of vertebrate A-clade R2p for template use by copying the terminal
region of 3′UTR RNA would be beneficial for selective insertion of the intended transgene. Our
findings inform future improvements and possible reprogramming of R2p-based transgene
insertion to the human genome.
357
358

Acknowledgements
We thank members of the Collins and Nogales laboratories for discussions in this collaborative
project. We thank Isabella Bartmess for preliminary studies of pseudoknot significance. We thank
D. Toso and R. Thakkar at the Cal-Cryo EM facility at UC Berkeley for help with EM data
acquisition. **Funding:** Damon Runyon Postdoctoral Fellowship and Damon Runyon-Dale F. Frey
Award (A.T.), the National Institutes of Health grants F32 GM139306 and California Institute for
Regenerative Medicine training grant EDUC4-12790 (B.V.T.), Fulbright Future Scholarship (The
Kinghorn Foundation) (N.T.H.), the University of Adelaide (D.L.A.), the National Institutes of
Health grant R35-GM127018 (E.N.), and the National Institutes of Health grant DP1 HL156819
(K.C.). E.N. is a Howard Hughes Medical Institute Investigator.

Competing interests: K.C. is an equity holder and scientific advisor for Addition Therapeutics,
Inc., using a retrotransposon-based genome engineering technology. K.C. and B.V.T. are listed
inventors on patent applications filed by University of California, Berkeley related to the PRINT
platform.
374

375 **Author contributions**
376 Conceptualization: A.R.V., A.T., K.C.; Methodology: A.T., A.R.V., B.V.T. and N.T.H.;
377 Investigation: A.T., A.R.V., B.V.T. and N.T.H.; Visualization: A.T., A.R.V., B.V.T. and N.T.H.;
378 Supervision: K.C., E.N. and D.L.A.; Writing—original draft: A.T., A.R.V., and K.C.; Writing—
379 review & editing: all authors.
380

381 **Data Availability**
382 The cryo-EM maps reported in this work are deposited under EMD-XXXX, EMD-XXXX and
383 EMD-XXXX in the Electron Microscopy Data Bank and the corresponding atomic model under
384 PDB YYY, PDB YYY and PDB YYY on the Protein Data Bank. All other datasets generated and
385 analyzed during the current study are available from the corresponding authors on request.

References

1. J. S. Han, Non-long terminal repeat (non-LTR) retrotransposons: mechanisms, recent developments, and unanswered questions. *Mob DNA* **1**, 15 (2010).

2. L. M. Payer, K. H. Burns, Transposable elements in human genetic disease. *Nat Rev Genet* **20**, 760–772 (2019).

3. P. Mita, J. D. Boeke, How retrotransposons shape genome regulation. *Curr Opin Genet Dev* **37**, 90–100 (2016).

4. D. A. Flasch, Á. Macia, L. Sánchez, M. Ljungman, S. R. Heras, J. L. García-Pérez, T. E. Wilson, J. V. Moran, Genome-wide de novo L1 Retrotransposition Connects Endonuclease Activity with Replication. *Cell* **177**, 837-851.e28 (2019).

5. A. Thawani, A. J. F. Ariza, E. Nogales, K. Collins, Template and target-site recognition by human LINE-1 in retrotransposition. *Nature* **626**, 186–193 (2024).

6. P. Mita, A. Wudzinska, X. Sun, J. Andrade, S. Nayak, D. J. Kahler, S. Badri, J. LaCava, B. Ueberheide, C. Y. Yun, D. Fenyö, J. D. Boeke, LINE-1 protein localization and functional dynamics during the cell cycle. *Elife* **7**, e30058 (2018).

7. M. Percharde, C.-J. Lin, Y. Yin, J. Guan, G. A. Peixoto, A. Bulut-Karslioglu, S. Biechele, B. Huang, X. Shen, M. Ramalho-Santos, A LINE1-Nucleolin Partnership Regulates Early Development and ESC Identity. *Cell* **174**, 391-405.e19 (2018).

8. X. Li, L. Bie, Y. Wang, Y. Hong, Z. Zhou, Y. Fan, X. Yan, Y. Tao, C. Huang, Y. Zhang, X. Sun, J. X. H. Li, J. Zhang, Z. Chang, Q. Xi, A. Meng, X. Shen, W. Xie, N. Liu, LINE-1 transcription activates long-range gene expression. *Nat Genet* **56**, 1494–1502 (2024).

9. S. Li, X. Shen, Long interspersed nuclear element 1 and B1/Alu repeats blueprint genome compartmentalization. *Curr Opin Genet Dev* **80**, 102049 (2023).

10. J. Sharif, H. Koseki, N. F. Parrish, Bridging multiple dimensions: roles of transposable elements in higher-order genome regulation. *Curr Opin Genet Dev* **80**, 102035 (2023).

11. H. S. Malik, W. D. Burke, T. H. Eickbush, The age and evolution of non-LTR retrotransposable elements. *Mol Biol Evol* **16**, 793–805 (1999).

12. H. Fujiwara, Site-specific non-LTR retrotransposons. *Microbiol Spectr* **3**, MDNA3-0001–2014 (2015).

13. K. K. Kojima, Y. Seto, H. Fujiwara, The Wide Distribution and Change of Target Specificity of R2 Non-LTR Retrotransposons in Animals. *PLoS One* **11**, e0163496 (2016).

14. T. H. Eickbush, D. G. Eickbush, Integration, Regulation, and Long-Term Stability of R2 Retrotransposons. *Microbiol Spectr* **3**, MDNA3-0011–2014 (2015).

15. W. D. Burke, C. C. Calalang, T. H. Eickbush, The site-specific ribosomal insertion element
type II of Bombyx mori (R2Bm) contains the coding sequence for a reverse transcriptase-
like enzyme. *Mol Cell Biol* **7**, 2221–2230 (1987).

16. D. D. Luan, M. H. Korman, J. L. Jakubczak, T. H. Eickbush, Reverse transcription of R2Bm
RNA is primed by a nick at the chromosomal target site: a mechanism for non-LTR
retrotransposition. *Cell* **72**, 595–605 (1993).

17. X. Zhang, B. Van Treeck, C. A. Horton, J. J. R. McIntyre, S. M. Palm, J. L. Shumate, K.
Collins, Harnessing eukaryotic retroelement proteins for transgene insertion into human
safe-harbor loci. *Nat Biotechnol*, doi: 10.1038/s41587-024-02137-y (2024).

18. Y. Chen, S. Luo, Y. Hu, B. Mao, X. Wang, Z. Lu, Q. Shan, J. Zhang, S. Wang, G. Feng, C.
Wang, C. Liang, N. Tang, R. Niu, J. Wang, J. Han, N. Yang, H. Wang, Q. Zhou, W. Li,
All-RNA-mediated targeted gene integration in mammalian cells with rationally engineered
R2 retrotransposons. *Cell* **187**, 4674-4689.e18 (2024).

19. S. M. Palm, C. A. Horton, X. Zhang, K. Collins, Structure and sequence at an RNA template
5' end influence insertion of transgenes by an R2 retrotransposon protein. *RNA* **30**, 1227–
1245 (2024).

20. R. J. Lee, C. A. Horton, B. Van Treeck, J. J. R. McIntyre, K. Collins, Conserved and
divergent DNA recognition specificities and functions of R2 retrotransposon N-terminal
domains. *Cell Rep* **43**, 114239 (2024).

21. M. E. Wilkinson, C. J. Frangieh, R. K. Macrae, F. Zhang, Structure of the R2 non-LTR
retrotransposon initiating target-primed reverse transcription. *Science*, eadg7883 (2023).

22. P. Deng, S.-Q. Tan, Q.-Y. Yang, L. Fu, Y. Wu, H.-Z. Zhu, L. Sun, Z. Bao, Y. Lin, Q. C.
Zhang, H. Wang, J. Wang, J.-J. G. Liu, Structural RNA components supervise the
sequential DNA cleavage in R2 retrotransposon. *Cell* **186**, 2865-2879.e20 (2023).

23. W. Bao, K. K. Kojima, O. Kohany, Repbase Update, a database of repetitive elements in
eukaryotic genomes. *Mob DNA* **6**, 11 (2015).

24. S. M. Palm, B. Van Treeck, K. Collins, Experimental considerations for precise RNA-
mediated insertion of transgenes. *Methods Enzymol* **705**, 1–24 (2024).

25. J. L. Stamos, A. M. Lentzsch, A. M. Lambowitz, Structure of a Thermostable Group II
Intron Reverse Transcriptase with Template-Primer and Its Functional and Evolutionary
Implications. *Mol Cell* **68**, 926-939.e4 (2017).

26. D. B. Haack, X. Yan, C. Zhang, J. Hingey, D. Lyumkis, T. S. Baker, N. Toor, Cryo-EM
Structures of a Group II Intron Reverse Splicing into DNA. *Cell* **178**, 612-623.e12 (2019).

27. B. K. Thompson, S. M. Christensen, Independently derived targeting of 28S rDNA by A-
and D-clade R2 retrotransposons: Plasticity of integration mechanism. *Mob Genet Elements*
1, 29–37 (2011).

28. J. Abramson, J. Adler, J. Dunger, R. Evans, T. Green, A. Pritzel, O. Ronneberger, L.
Willmore, A. J. Ballard, J. Bambrick, S. W. Bodenstein, D. A. Evans, C.-C. Hung, M.
O'Neill, D. Reiman, K. Tunyasuvunakool, Z. Wu, A. Žemgulytė, E. Arvaniti, C. Beattie, O.
Bertolli, A. Bridgland, A. Cherepanov, M. Congreve, A. I. Cowen-Rivers, A. Cowie, M.
Figurnov, F. B. Fuchs, H. Gladman, R. Jain, Y. A. Khan, C. M. R. Low, K. Perlin, A.
Potapenko, P. Savy, S. Singh, A. Stecula, A. Thillaisundaram, C. Tong, S. Yakneen, E. D.
Zhong, M. Zielinski, A. Žídek, V. Bapst, P. Kohli, M. Jaderberg, D. Hassabis, J. M.
Jumper, Accurate structure prediction of biomolecular interactions with AlphaFold 3.
Nature **630**, 493–500 (2024).

29. A. Luchetti, B. Mantovani, Non-LTR R2 element evolutionary patterns: phylogenetic
incongruences, rapid radiation and the maintenance of multiple lineages. *PLoS One* **8**,
e57076 (2013).

30. S. M. Christensen, J. Ye, T. H. Eickbush, RNA from the 5' end of the R2 retrotransposon
controls R2 protein binding to and cleavage of its DNA target site. *Proc Natl Acad Sci U S*
A **103**, 17602–17607 (2006).

31. F. D. Urnov, E. J. Rebar, M. C. Holmes, H. S. Zhang, P. D. Gregory, Genome editing with
engineered zinc finger nucleases. *Nat Rev Genet* **11**, 636–646 (2010).

32. J. K. Joung, J. D. Sander, TALENs: a widely applicable technology for targeted genome
editing. *Nat Rev Mol Cell Biol* **14**, 49–55 (2013).

33. Z. Zhang, S. Schwartz, L. Wagner, W. Miller, A greedy algorithm for aligning DNA
sequences. *J Comput Biol* **7**, 203–214 (2000).

34. E. Gasteiger, A. Gattiker, C. Hoogland, I. Ivanyi, R. D. Appel, A. Bairoch, ExPASy: The
proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Res* **31**,
3784–3788 (2003).

35. A. Rodríguez-Vargas, K. Collins, Distinct and overlapping RNA determinants for binding
and target-primed reverse transcription by Bombyx mori R2 retrotransposon protein.
Nucleic Acids Res **52**, 6571–6585 (2024).

36. Messenger RNA encoding the full-length SARS-CoV-2 spike glycoprotein. (2020).
https://web.archive.org/web/20210105162941/https://mednet-
communities.net/inn/db/media/docs/11889.doc.

37. A. Patel, D. Toso, A. Litvak, E. Nogales, "Efficient graphene oxide coating improves cryo-
EM sample preparation and data collection from tilted grids" (preprint, Biophysics, 2021);
https://doi.org/10.1101/2021.03.08.434344.

38. S. Q. Zheng, E. Palovcak, J.-P. Armache, K. A. Verba, Y. Cheng, D. A. Agard, MotionCor2:
anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat*
Methods **14**, 331–332 (2017).

39. J. Zivanov, T. Nakane, B. O. Forsberg, D. Kimanius, W. J. Hagen, E. Lindahl, S. H. Scheres,
New tools for automated high-resolution cryo-EM structure determination in RELION-3.
Elife **7**, e42166 (2018).

40. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron
micrographs. *J Struct Biol* **192**, 216–221 (2015).

41. E. F. Pettersen, T. D. Goddard, C. C. Huang, E. C. Meng, G. S. Couch, T. I. Croll, J. H.
Morris, T. E. Ferrin, UCSF ChimeraX: Structure visualization for researchers, educators,
and developers. *Protein Sci* **30**, 70–82 (2021).

42. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta*
Crystallogr D Biol Crystallogr **66**, 486–501 (2010).

43. P. D. Adams, P. V. Afonine, G. Bunkóczi, V. B. Chen, I. W. Davis, N. Echols, J. J. Headd,
L.-W. Hung, G. J. Kapral, R. W. Grosse-Kunstleve, A. J. McCoy, N. W. Moriarty, R.
Oeffner, R. J. Read, D. C. Richardson, J. S. Richardson, T. C. Terwilliger, P. H. Zwart,
PHENIX: a comprehensive Python-based system for macromolecular structure solution.
Acta Crystallogr D Biol Crystallogr **66**, 213–221 (2010).

Figure 1

Fig. 1. TPRT and PRINT activities and cryo-EM structures of A-clade R2 RNPs initiating TPRT. (a) Schematic of biochemical steps during DNA insertion. (b) Phylogenetic analysis of R2p from the A-clade (birds, turtle, red flour beetle) and D-clade (silk moth and fruit fly) characterized in this and previous work (*17*, *20*). Tree branch length is indicative of substitutions per aligned site. (c) Denaturing PAGE of TPRT reaction products. Orange triangles indicate expected TPRT product lengths for copying a single full-length template (TPRT cDNA). Multiple templates may also be copied in series (template jumping products). R2Pm and R2Tg proteins were assayed with annealed rDNA target site oligonucleotides and different template RNAs, each with an R5 3′ tail: Gf98, Pm112, Bm3. (d) PRINT assay schematic. An mRNA encoding R2Pm or R2Tg protein is transfected with an engineered template RNA comprised of a 5′ module (5′M), modified CMV promoter (PRO), GFP ORF, polyadenylation signal (PA), and 3′ module (3′M) with a 3′tail containing rRNA and A22. (e) PRINT assays with 2-RNA transfection of the R2Pm or R2Tg mRNA and an engineered template RNA with either Gf3 or Pm3 followed by R4A22. Note the log-scale y-axis. (f-g) At top, domains of A-clade R2Pm and R2Tg are illustrated with amino acid numbering; abbreviations given in the text. Cryo-EM density of R2Pm (f) or R2Tg (g) first strand synthesis complex assembled with rDNA target site and either Gf3 full-length 3′UTR RNA (f) or Gf98 RNA (g) is shown and colored by domain. (h-i) Ribbon diagrams of R2Pm (h) or R2Tg (i) first strand synthesis complex structure colored by domains.

Figure 2

Fig. 2. Protein and DNA recognition of R2 3′UTR RNA. (a) Schematic of direct interactions
between R2Pm protein, rDNA target site, and 3′UTR RNA in a TPRT initiation complex. Color
scheme is consistent with Figure 1. Solid navy lines denote direct hydrogen bonds with the
nucleobases or ribonucleobases, while dashed navy lines represent hydrogen bonds with the
phosphate backbone or sugars. Solid mustard lines denote pi-stacking contacts with the
nucleobases or ribonucleobases. Black circles represent base-pairs in DNA duplex; RNA-DNA or
RNA-RNA base-pairing is indicated by apposition. DNA numbering (green and gray strands) is
negative upstream or positive downstream of the first strand nick. RNA numbering (red strand) is

from the start of Gf3. (b-c) Recognition of the 3′UTR RNA involves the NTE -1, Thumb, Linker
and ZnF3 domains. (b) Base-specific hydrogen bonds between bases G-256 and A-258 in the hinge
region of 3′UTR RNA and side chains within the Thumb and Linker domains in R2Pm. (c) ZnF3
domain from R2Pm and R2Tg contacts the pseudoknot of 3′UTR RNA. (d) Side chains in ZnF3
make base-specific hydrogen bonds: R2Pm with G-236. R2Pm ZnF3 also makes a contact with
the phosphate backbone of base G-237 at the junction of hinge and pseudoknot and R2Tg's ZnF3
with the phosphate backbone of base C-253. The helix segmentation is an artifact of automated
secondary structure assignment. Here and in subsequent figure panels, heteroatom representation
has oxygens in red and nitrogens in blue. (e) Base-specific hydrogen bonds between pseudoknot
bases and a bases in a single-stranded region of the second strand DNA. (f) PRINT assays using
mRNA encoding R2Tg and template RNA with 3′ module Gf98, or a variant Gf98, and R4A22 3′
tail. Base substitutions are numbered according to their position in Gf3, as annotated in (a), with
specific mutations described in the main text.
823
824

Figure 3

825
Fig. 3. Protein recognition of the target DNA and N-terminal domain requirements for TPRT
and PRINT. (a) RLE and ZnK domains surrounding the nicked first strand and single-stranded
second strand are illustrated for the R2Pm complex. (b) The motif 6a loop within the RT domain
is shown protruding into a distortion in target DNA. (c) Configuration on target DNA of the N-
terminal DNA binding domains: the three ZnF and the Myb domain for A-clade R2Pm and R2Tg
are compared with the single ZnF and Myb in D-clade R2Bm. (d) Base-reading hydrogen bonds
between ZnF2 and the target DNA proximal to the nick site. (e) The unstructured R2Pm Spacer
and its interaction with the RT and NTE 0 domains are depicted. (f) Denaturing PAGE of TPRT

reaction products with wild-type R2Tg, R2Za, R2Pm and chimeric proteins: R2Pm with the N-
terminus (Spacer, Myb, and three ZnFs) from R2Tg (NTg) or R2Za (NZa), R2Pm with ZnF3-2
domains from R2Tg (ZFTg), R2Pm with Spacer from R2Tg (spacTg). Gf68 RNA with R5 3′ tail
was used for all assays. Different regions of the same gel are shown, with first strand DNAs and
second strand DNAs imaged separately using different 5′ dye. (g) PRINT assays using mRNA
encoding R2Pm or the chimeras described in (f). The template RNA 3′ module was Gf3 followed
by R4A22.
841
842

Figure 4

843
Fig. 4. A C-terminal insertion in A-clade R2p. (a) The CTI is rendered in yellow against the RT
and Linker domains and RNA:cDNA duplex. The shorter loop present in R2Bm is shown for
comparison. (b) Side chains of the conserved EWE motif that anchors the CTI to the RT are
displayed for R2Pm. (c) Denaturing PAGE of TPRT reaction products with wild-type R2Tg, R2Tg
ΔCTI (CTI truncation) mutant, wild-type R2Pm and R2Pm ΔCTI mutant. Gf68 RNA was
synthesized with a variable length of the 3′ tail that base-pairs to target site primer: 0, 3, 4, 5, 8 and
12 nt. Different regions of the same gel are shown, with first strand DNAs and second strand DNAs
imaged separately using different 5′ dye. (d) PRINT assays were performed by 2-RNA transfection
of the indicated R2p mRNA and template RNA with Gf3 followed by R4A22.
853
854

Figure 5

Fig. 5. Biochemical activity and cryo-EM structure of A-clade R2 retrotransposon during second strand nicking. (a) Denaturing PAGE of target site nicking and TPRT reaction products from assays using wild-type R2Tg or its RTD and END variants. Gf68 RNA with R5 was used as template. Different regions of the same gel are shown, with first strand DNAs and second strand DNAs imaged separately using different 5′ dyes. Small triangle (mustard) indicates TPRT cDNA. (b) Nucleic acid substrate design to capture a post-TPRT structure for an R2Pm complex. 2D class averages from cryo-EM analysis are shown with inferred range of positions of RNA:cDNA duplex exiting the protein density. (c) Cryo-EM density and ribbon diagram of R2Pm second strand nicked complex assembled, colored by domains. (d) Comparison of upstream target site DNA position in the R2Pm first strand synthesis complex versus second strand nicked complex relative to the R2Pm (NTE to RLE) core (white) and bound 3′UTR RNA (red). After second strand nicking, the nicked single-stranded second strand DNA is displaced towards the RT core and the double-

868   strand DNA bend angle changes near the ZnF1 and Myb domains. (e) Nicked ends of upstream
869   target site DNA are illustrated with nearby R2Pm protein regions NTE -1 and ZnF3-2.