WMH-DualTasker: A Dual-Task Deep Learning Model with Self-supervised Consistency for Automated Segmentation and Visual Rating of White matter Hyperintensities - a Multicentre study

Yilei Wu[1,5] • Zijian Dong[1,5,6] • Hongwei Bran Li, PhD[2] • Yao Feng Chong, MD[1,5] • Fang Ji, PhD[1,5] • Joanna Su Xian Chong, PhD[1,5] • Nathanael Ren Jie Tang[1,5] • Saima Hilal, PhD[4] • Huazhu Fu, PhD[3] • Christopher Li-Hsian Chen, MD[4] • Juan Helen Zhou, PhD[1,5,6,7] • for the Alzheimer's Disease Neuroimaging Initiative*

[1]Centre for Sleep and Cognition & Centre for Translational Magnetic Resonance Research, National University of Singapore, Singapore; [2]Athinoula A. Martinos Centre for Biomedical Imaging, Massachusetts General Hospital, Harvard Medical School, USA; [3]Institute of High Performance Computing, A*STAR, Singapore; [4]Memory Aging & Cognition Centre, National University Health System, Singapore; [5]Human Potential Translational Research Program and Department of Medicine, [6]Yong Loo Lin School of Medicine, National University of Singapore, Singapore; Department of Electrical and Computer Engineering, National University of Singapore, Singapore; [7]Integrative Sciences and Engineering Programme (ISEP), NUS Graduate School, National University of Singapore, Singapore

*Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu) and the Australian Imaging Biomarkers and Lifestyle Study of Aging (AIBL) database (www.aibl.csiro.au). As such, the investigators within the ADNI and AIBL contributed to the design and implementation of ADNI and AIBL and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Address correspondence to:

Juan Helen Zhou

Centre for Sleep and Cognition & Centre for Translational Magnetic Resonance Research, National University of Singapore

Tel: 65-96501661

Email: helen.zhou@nus.edu.sg

Address: 12 Science Drive 2, Singapore 117549

Funding: This study was supported by the Singapore National Medical Research Council (NMRC/OFLCG19 May-0035, NMRC/CIRG/1485/2018, NMRC/CSA-SI/0007/2016, NMRC/MOH-00707-01, NMRC/CG/435 M009/2017-NUH/NUHS, CIRG21nov-0007 and HLCA23Feb-0004), RIE2020 AME Programmatic Fund from A*STAR, Singapore (No. A20G8b0102), Ministry of Education (MOE-T2EP40120-0007 & T2EP2-0223-0025, MOE-T2EP20220-0001), and Yong Loo Lin School of Medicine Research Core Funding, National University of Singapore, Singapore.

Competing interests: Associate Editor is co-author – <Juan Helen Zhou> is a handling editor of <Human Brain Mapping> and co-author is of this article. To minimize bias, they were excluded from all editorial

Acknowledgements

We extend our heartfelt gratitude to all the participants and the dedicated research teams at MACC and SINGER for their invaluable contributions. A special thanks goes to Dr. Hugo Kuijf for not only hosting the MICCAI-WMH segmentation challenge but also for making the dataset widely accessible to the public. Additionally, we are deeply appreciative of the Alzheimer's Disease Neuroimaging Initiative (ADNI) for generously making their dataset available to the research community. These collaborative efforts are fundamental to the advancement of our work.

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.

Keywords: White matter hyperintensity, Weakly supervised segmentation, Dementia, MRI

Abstract

Background: White matter hyperintensities (WMH) are neuroimaging markers linked to an elevated risk of cognitive decline. WMH severity is typically assessed via visual rating scales and through volumetric segmentation. While visual rating scales are commonly used in clinical practice, they offer limited descriptive power. In contrast, supervised volumetric segmentation requires manually annotated masks, which is labor-intensive and challenging to scale for large studies. Therefore, our goal was to develop an automated deep learning model that can provide accurate and holistic quantification of WMH severity with minimal supervision.

Methods: We developed WMH-DualTasker, a deep learning model that simultaneously performs voxel-wise segmentation and visual rating score prediction. The model leverages self-supervised, transformation-invariant consistency constraints, using WMH visual ratings from clinical settings as the sole supervisory signal. Additionally, we assessed its clinical utility by applying it to identify individuals with mild cognitive impairment (MCI) and to predict dementia conversion.

Findings: The volumetric quantification performance of WMH-DualTasker was either superior to or on par with existing supervised methods, as demonstrated on the MICCAI-WMH dataset (N=60, Dice=0.602) and the SINGER dataset (N=64, Dice=0.608). Furthermore, the model exhibited strong agreement with clinical visual rating scales on an external dataset (SINGER, MAE=1.880, K=0.77). Importantly, WMH severity metrics derived from WMH-DualTasker improved predictive performance beyond conventional clinical features for MCI classification (AUC=0.718, p<0.001), MCI conversion prediction (AUC=0.652, p<0.001) using the ADNI dataset.

Interpretations: WMH-DualTasker substantially reduces the reliance on labor-intensive manual annotations, facilitating more efficient and scalable quantification of WMH severity in large-scale population studies. This innovative approach has the potential to advance preventive and

precision medicine by enhancing the assessment and management of vascular cognitive impairment associated with WMH. Code and model weights are publicly available at https://github.com/hzlab/WMH-DualTasker.

1 Introduction

White matter hyperintensities (WMH) are neuroimaging markers that are usually indicative of cerebral small vascular disease — a condition characterized by damage to the small blood vessels in the brain. Using a T2-weighted fluid-attenuated inversion recovery (FLAIR) sequence on magnetic resonance imaging (MRI) [1,2], these hyperintensities manifest as areas of abnormal heightened signal intensity in the white matter. The presence of WMH carries significant clinical implications: studies have demonstrated their association with a myriad of neurological complications, including accelerated cognitive decline, a heightened risk of dementia, and an increased probability of stroke [3-5]. These links underscore WMH crucial influence on cognition and cerebrovascular disease.

There are two main ways to quantify WMH: voxel-wise segmentation and visual rating (Figure 1). Segmentation provides a volumetric assessment and an objective measure of WMH volume. This approach offers granularity and precision, capturing detailed information about the extent of WMH in the brain. However, the segmentation process might be time-consuming and technically challenging. On the other hand, visual rating is a method that assesses WMH based on standardized scales [2, 6-8]. Trained raters, such as radiologists and neurologists, evaluate and grade the severity and distribution of WMH based on their appearance in the scans. These visual ratings serve as a qualitative assessment of the WMH load. One scale commonly used for this purpose is the age-related white matter change (ARWMC) visual rating scale [7]. This scale assigns a global score for each brain, with values ranging from 0 to 30, where a higher score indicates increased severity. Notably, visual rating scales such as ARWMC are often more time-efficient than the manual creation of voxel-wise WMH masks for segmentation. However, the subjective nature of visual ratings might introduce variability among different raters. Considering the distinct advantages offered by comprehensive voxel-

wise segmentation and the efficiency and qualitative insights derived from visual ratings, both methodologies contribute valuable and complementary information in the evaluation of WMH. In the early era of automated WMH segmentation research, methodologies primarily relied on conventional machine learning techniques [9-11], including but not limited to random forest and logistic regression-based frameworks. However, these conventional techniques were limited by their reliance on manual feature extraction and lacked the ability to capture complex imaging patterns. With the advancement of deep learning techniques in the realm of medical image segmentation, approaches specific to WMH segmentation have naturally evolved to incorporate such techniques. Recent studies [12-14] highlight the preliminary successes of deep learning-based models for this task.

However, despite these advances, one outstanding challenge that remains is the dependence of these models on manually labelled WMH masks for effective training. The labelling process is labor-intensive; typically, a domain expert, such as a radiologist, dedicates between 0.5 to 2 hours to the meticulous labelling of a single MRI volume. This requirement for specialized expertise and the associated time commitment accentuates the difficulty in collecting large-scale manually labelled WMH datasets. Without a large number of such datasets, it is unlikely that the data will be sufficiently expansive and comprehensive to train a generalizable and performant segmentation model.

In this paper, we introduce WMH-DualTasker, a deep learning-based approach for simultaneous WMH segmentation and visual rating prediction. WMH-DualTasker predicts both the ARWMC score and the WMH mask using only visual rating scales for supervision, thus overcoming the challenge of acquiring manual WMH segmentation masks. Specifically, for weakly supervised segmentation, we leverage activation maps generated from the regression task as the initialization for WMH segmentation. We introduce an equivalent regularization loss to improve the localization capability of the activation map by ensuring it

learns transformation invariance. This enhancement increases its sensitivity to less distinguishable lesion regions while concurrently suppressing activations in non-lesion areas. Additionally, considering the tiny and hyperintense nature of WMH, we propose a hyperintense map to leverage intensity information, further improving the sensitivity of WMH segmentation. To the best of our knowledge, our work is the first to present a 3D weakly supervised segmentation framework guided by image-level weak supervision, distinct from methods guided by an object bounding box or partial annotations.

To investigate the effectiveness of WMH-DualTasker and understand how well it performs on independent test sets, we rigorously evaluated WMH-DualTasker on three distinct datasets. Remarkably, our framework not only demonstrates broad applicability but also exhibits performance metrics on par with supervised methods [9-11], underscoring its efficacy. WMH-DualTasker also demonstrates good to high agreement on ARWMC prediction with our radiologist and outperform on both internal and external validation. Additionally, to further substantiate the clinical utility of WMH-DualTasker for tasks such as dementia diagnosis, we apply it to the Alzheimer's Disease Neuroimaging Initiative (ADNI) dataset [15] to examine the predictive capability of the ARWMC rating and WMH segmentation mask generated by WMH-DualTasker. Our findings show that by combining the estimated volume of WMH and the ARWMC score with conventional clinical features, we can improve the accuracy in distinguishing between severe and mild stages of cognitive impairment. This underscores WMH-DualTasker's potential not only as a proficient tool for WMH identification but also as a clinically relevant aid in distinguishing between different stages of cognitive impairment.

2 Methods

2.1 Study design and datasets

Throughout our experiment, we use four datasets that span a range of ages from middle-aged to elderly, as well as multiple sites and ethnicities. Figure 2 illustrates our study design and

configuration among the four datasets and the type of annotation they have. We describe each dataset in the following part of this section and in Table 1, and more demographic information can be found in Tables S2-S5 in the supplementary material.

2.1.1 MACC dataset

A total of 660 T2-FLAIR MRI images from the Memory, Ageing and Cognition Centre (MACC) dataset are used. This dataset includes NCI (No Cognitive Impairment), CIND (Cognitive Impairment, No Dementia), AD (Alzheimer's Disease), and VaD (Vascular Dementia) participants whose ARWMC ratings have been determined by a radiologist with over ten years of experience. All images were quality-checked and were visually rated by clinical experts using the ARWMC scale. For WMH masks, we randomly select 40 images and manually annotate WMH masks by clinical experts as the reference standard for evaluating mentation performance (MACC-Test). Each image will be annotated by two radiologists with over ten years of experience independently and be reviewed by the third senior radiologist if the agreement is not achieved. Overall, we have three splits: MACC-Train (n=600), MACC-Val (n=20), and MACC-Test (n=40) (see Figure 2). To prevent information leakage, we use MACC-Val for selecting hyper-parameters, and MACC-Test was hidden from the model development. Additionally, our model was trained on MACC-Train with image-level weak signals **only**, and the best model was selected based on the lowest loss on the MACC-Val.

2.1.2 MICCAI-WMH and SINGER dataset

We utilize a total of 60 images from the MICCAI-WMH challenge [16] and 64 images from the Singapore Geriatric Intervention Study (SINGER) [17] to examine the performance of the WMH-DualTasker across different datasets. The MICCAI-WMH, a public dataset from an open challenge, provides MRI scans (from multiple sites) with WMH masks that are manually annotated and cross-verified by two independent annotations from radiologists with over ten years of experience. The SINGER study is designed to assess the efficacy and safety of

multidomain lifestyle interventions aimed at delaying cognitive decline in older adults at risk for dementia, and targets participants aged 60-77 with specific risk factors for dementia. For our analysis, we leverage 64 MRI scans from SINGER, accompanied by manually annotated WMH masks and visual ratings performed by radiologists using the ARWMC scale (similar to the MACC). We assess our model using the MICCAI-WMH and SINGER as external testing datasets to evaluate the generalizability of the WMH-DualTasker for both WMH segmentation and regression.

2.1.3 ADNI dataset

We employ the Alzheimer's Disease Neuroimaging Initiative (ADNI) dataset to assess the clinical significance of the WMH-DualTasker model. Data used in the preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). The ADNI was launched in 2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner, MD. The primary goal of ADNI has been to test whether serial magnetic resonance imaging (MRI), positron emission tomography (PET), other biological markers, and clinical and neuropsychological assessment can be combined to measure the progression of mild cognitive impairment (MCI) and early Alzheimer's disease (AD). For up-to-date information, see www.adni-info.org. We selected 394 ADNI subjects who had completed cognitive evaluations and MRI scans at their baseline visits (Figure 2). According to their baseline diagnoses, these subjects comprised 152 Cognitively Normal (CN), 149 Mild Cognitive Impairment (MCI), and 93 Alzheimer's Disease (AD) participants. Each subject's MRI scan included FLAIR images acquired in the axial plane.

2.1.4 Pre-processing

We pre-process all images using the same steps for all datasets. For a fair comparison, all images were resampled to the same resolution $\left(1.00 \times 1.00 \times 3.00(\text{mm}^3)\right)$ and center-

cropped to the size of $256 \times 256 \times 64$. We then normalized the intensities in each image using z-score transformation.

2.1.5 Ethics

The MACC study the ethics approval from NHG DOMAIN SPECIFIC REVIEW BOARD (DSRB) APPROVAL with a reference number of 2015/00406 (approval period: 01 November 2017 to 31 March 2020). The NHG DSRB operates in accordance to the ICH GCP and all applicable laws and regulations. Subjects were recruited into the study from the NUHS memory clinics and NUHS MACC research cohorts. Considering the retrospective nature of this study, the informed consent was waived.

The SINGER study obtained ethics approval from NUS-IRB REVIEW BOARD APPROVAL with a reference number of 2020/398 (approval period: 28 Aug 2020 to 31 May 2022). This is human biomedical research that is regulated by the Human Biomedical Research Act (HBRA) and researchers are required by law to comply with all the relevant regulatory requirements of the HBRA.

2.2 WMH-DualTasker - automatic segmentation and visual rating of WMH

This section introduces the formulation of WMH-DualTasker, which aims to provide a holistic solution for automated WMH analysis by simultaneously accomplishing two tasks: WMH segmentation and ARWMC prediction. It is trained in an end-to-end manner, using only the ARWMC score as the supervisory signal. We begin with the challenge of identifying White Matter Hyperintensities (WMH) using image-level weak labels. Then, we describe how self-supervised equivalent regularization loss enhances this process, and how integrating a hyperintense map further optimizes it by leveraging the prior characteristics of WMH. Finally, in Section 2.2.3 the inference strategy employed is introduced. WMH-DualTasker utilizes a 3D model and takes MRI volumes as its input (see Figure 3). More details about the model structure can be found in Figure S1 in the supplementary material.

2.2.1 Weakly supervised WMH segmentation through optimized CAM with self-supervised transformation equivalent regularization loss

Our approach to weakly supervised WMH segmentation centers on optimizing the Class Activation Map (CAM [19]) technique, a powerful tool for visualizing influential regions in image classification tasks. It works by generating a salient map that highlights the parts of an image most relevant to a particular class label. In this work, we extract a CAM that highlights WMH by training a 3D convolutional neural network to predict ARWMC given a brain MRI volume. We adopt the 3D light-weight structure (SFCN) [20] as the backbone network in our method (unless stated otherwise) due to its lower memory footprint.

Due to the weak supervisory signal in the classification task compared to the segmentation task, the visualization of CAM is usually blurry and thus cannot accurately localize white matter abnormalities. In addition, as WMH frequently appears in periventricular areas, we observe that the CAM simply highlights the centre of the brain. This motivates us to improve the localization capabilities of CAM from two perspectives. First, we refine its resolution by eliminating the max-pooling layers from standard 3D backbone networks, thereby obtaining feature maps with a higher spatial resolution. Specifically, our feature maps have a resolution of $\frac{H}{4} \times \frac{W}{4} \times \frac{D}{2}$ where H, W, D indicate the height, width, and depth of the input volume. Second, we introduce a regularization loss to address the limited localization capability of CAM from the regression task. Using CAM as an initial seed would result in poor WMH segmentation quality. To overcome this, we incorporate equivariance properties [21] into the learning objective. This integration promotes robust regularization, ensuring CAM consistency under perturbations (such as affine image transformations), thereby enhancing its spatial alignment with abnormal white matter (WMH) regions.

We implement the equivalent regularization loss for our volumetric application (see Figure 3). We denote the backbone network (without global average pooling (GAP) layers)as

$\theta(\cdot)$ and image-transform functions (e.g., scale, rotate, flip, translate) as $\pi(\cdot)$ respectively. For a brain MRI volume X, we have an activation map CAM and visual rating prediction $\hat{y}$ where $CAM = \theta(X)$ and $\hat{y} = GAP(CAM)$. Our equivalent regularization loss can be written as:

$$\mathcal{L}_{ER} = |\theta(X) - \pi^{-1}(\theta(\pi(X)))|_1 \quad (1)$$

The supervisory signal comes from the regression loss for predicting ARWMC visual ratings. We add a global average pooling (GAP) layer at the end of our model to aggregate the CAM of the input image. We adopt mean square error (MSE) loss for both original and transformed images. Our regression loss (MSE loss) can be written as follows:

$$\mathcal{L}_{Reg} = \frac{1}{2}\left(\mathcal{L}_2(\hat{y}^o, y) + \mathcal{L}_2(\hat{y}^t, y)\right) \quad (2)$$

Combining them, our overall training loss is:

$$\mathcal{L} = \mathcal{L}_{Reg} + \lambda\mathcal{L}_{ER} \quad (3)$$

where the λ is the hyper-parameter controls the strength of $\mathcal{L}_{ER}$.

2.2.2 Extraction of hyperintense map for enhanced WMH boundary delineation

WMH only comprises less than 1% of the volume of the whole brain [3] in healthy older subjects and it is dispersed throughout the white matter. Even with enhanced CAM, it can only reflect high-level semantics of the target with a smooth boundary, which makes it difficult to delineate a crisp boundary (see Figure 4). Moreover, we observe that directly using the enhanced CAM as the seed for segmentation, while helpful for large lesions, falls short of capturing the full extent of smaller WMH.

To address the challenge posed by the nature of WMH, we extract a hyperintense map from brain MRI volumes, using the intensity information as prior knowledge for segmentation mask generation (see Equation 4). The hyperintense map (HM) is defined as follows. The values of μ and σ stand for the mean and standard deviation of the grey matter (GM) intensity distribution, which were calculated from a GM mask generated with FSL-FAST [18]. α is a scaling parameter that adjusts the minimum signal intensity of outliers.

$$HM_i^{\text{FLAIR}} = \begin{cases} 1 & \text{if } X_i > \mu + \alpha\sigma \\ 0 & \text{otherwise} \end{cases} \forall i \in \Omega \ (4)$$

2.2.3 Inference of WMH-DualTasker

Lastly, the weighted sum of HM and CAM was post-processed by a 3D dense conditional random field (CRF) [22,23] to generate our final WMH segmentation mask output (see lower half of Figure 3).

$$WMH_{\text{pred}} = dCRF(\beta * HM + (1 - \beta) * CAM, \text{MRI}) \ (5)$$

The symbol $dCRF_{MRI}$ represents the 3D dense CRF function, and the coefficient β facilitates the linear combination of HM (Hyperintense Map) and CAM (Class Activation Map), while the MRI image acts as a reference within the CRF algorithm. This integration ensures that the segmentation of WMH is aligned with the specific attributes of each MRI scan. By incorporating the MRI data, the CRF framework effectively harnesses both spatial and intensity information intrinsic to the MRI images. This leads to a segmentation that is more contextually enriched.

2.2.4 Model Implementation and Training Details

We used SFCN [20] as our backbone architecture with the last two pooling layers removed for high-resolution CAM. To train the models, we use the AdamW optimizer [24] with an initial learning rate of 0.005 with linear learning rate decay to 0.00005 over 400 epochs. The weight for the equivalent regularization loss, lambda, is initially set to 0. This approach avoids enforcing the equivalent regularization loss at the start of training, which we empirically observe could degrade the quality of the CAM. Starting from the 200th epoch, lambda undergoes a stepwise increase, updating every epoch until it reaches 0.5. This gradual adjustment supports better training outcomes. We included scale and flip as image transform functions. We set the rescale ratio to 0.5-0.7 and flipped along any axis randomly for each iteration. Standard L2 and L1 loss were used to calculate our regression loss and equivalent regularization loss, respectively. Hyper-parameters alpha and beta were set to be 1.0 and 0.4,

respectively. More details about the parameter search range can be found in Table S1 in the supplmentary material.

2.3 Evaluation of WMH-DualTasker

We designed our experiments to rigorously evaluate WMH-DualTasker's performance, understand its components' contributions, and assess its potential clinical impact across different populations and cognitive states. Our experimental setup consisted of three main phases: performance evaluation, ablation studies, and downstream analysis (Figure 2).

2.3.1 Performance evaluation and ablation studies

We trained WMH-DualTasker on the MACC-Train dataset (n=600) using only visual ratings. To assess its accuracy in WMH segmentation and visual rating prediction, we tested it on MACC-Test (n=40), SINGER (n=64), and MICCAI-WMH (n=60) datasets (Section 3.2). We compared our model's performance against established supervised methods requiring manual annotation during training.

We conducted ablation studies on the MACC-Test dataset to examine each component's contribution to WMH-DualTasker (Section 3.3). These studies explored various configurations, including different backbone architectures, CAM methods, and our proposed enhancements such as high-resolution CAM and self-supervised regularization.

To evaluate the segmentation performance of WMH-DualTasker and other methods (Section 3.2), we adopted two primary metrics: Dice Score and Absolute Volume Difference (AVD). The Dice Score measures voxel-wise prediction accuracy for WMH compared to manual annotations, while AVD, a volume-based metric, provides information about a participant's overall WMH load. We chose to include AVD because WMH is frequently reported based on its volume in many research and clinical settings. This approach can differ from some other medical image segmentation applications (e.g., tumor, organ) where voxel-wise precision is often emphasized. As WMH-DualTasker can jointly predict the visual rating

score, we also evaluated its performance in this aspect. We rounded the predicted scores to the nearest integer and calculated the mean absolute error (MAE), Pearson correlation coefficient (R), and Kappa score to assess agreement with the visual rating scores manually annotated by radiologists. This comprehensive evaluation approach allows us to assess both the segmentation accuracy and the clinical relevance of the WMH-DualTasker's predictions.

$$\text{Dice} = \frac{2 \times (WMH_{\text{pred}} \cap WMH_{\text{mask}})}{WMH_{\text{pred}} + WMH_{\text{mask}}} \quad (6)$$

$$AVD = \frac{\left|V_{WMH_{\text{pred}}} - V_{WMH_{\text{mask}}}\right|}{V_{WMH_{\text{mask}}}} \quad (7)$$

$$\text{MAE} = \frac{\left|A\overline{RMW}C_{\text{pred}} - A\overline{RMW}C_{\text{manual}}\right|}{N} \quad (8)$$

2.3.2 Downstream validation analyses

For downstream validation analyses, we evaluated the utility of WMH volume and ARWMC derived from WMH-DualTasker in predicting cognitive performance across external datasets. We utilized two distinct cohorts: the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (n=394) as a disease cohort, and the SINGER dataset (n=470) as a community-based cohort. This dual-cohort approach allowed us to investigate the applications of these biomarkers in both clinical and general population settings.

In the ADNI dataset (http://adni.loni.usc.edu), we performed three types of analyses (Figure 5): classification between Cognitively Normal (CN) and Mild Cognitive Impairment (MCI) (155 CN & 146 MCI subjects; Figure 5a), prediction of MCI prognosis (40 MCI non-converters & 40 MCI converters; Figure 5b), and regression analyses of cognitive outcomes (394 subjects; Figure 5c). The regression analyses estimated scores for Mini-Mental State Examination (MMSE), Alzheimer's Disease Assessment Scale-Cognitive Subscale (ADAS-Cog), and Clinical Dementia Rating Scale sum of boxes (CDRSB).

We analyzed these tasks using four different combinations of input variables: clinical and demographic factors only, clinical factors plus WMH volume from nnU-Net, clinical factors plus WMH volume from WMH-DualTasker, and clinical factors plus both WMH volume and ARWMC rating from WMH-DualTasker. To ensure robust results, we employed ten-fold nested cross-validation, using logistic regression for classification tasks and kernel ridge regression for cognitive score prediction.

For the SINGER dataset, we explored the relationship between WMH burden and cognitive deterioration, specifically investigating the association between WMH volume predicted by WMH-DualTasker and the CAIDE Dementia Risk Score. The CAIDE score assesses the risk of developing dementia later in life, providing insights into the utility of our biomarkers in evaluating future dementia risk in a general population.

This comprehensive experimental setup allows us to thoroughly evaluate WMH-DualTasker's performance, understand its inner workings, and assess its potential clinical impact across various populations and cognitive states.

In the downstream analysis (Section 3.4), the area under the receiver operating characteristic (AUC) curves were conducted to evaluate the predictive capability of the input features. We applied ten-fold cross-validation for each feature combination on each task. A paired T-test was conducted across all methods, highlighting all pairs with significant differences. A statistically significant difference was set as $p < 0.05$. All statistical analyses were performed using Python version 3.7.11 with the Sklearn package and Statsmodels package.

3　Results

Our experiments demonstrate the effectiveness of WMH-DualTasker in WMH segmentation, ARWMC prediction, and clinical applications. Key findings include comparable or superior performance to supervised methods in segmentation tasks, significant improvements in ARWMC prediction, and added clinical value in cognitive impairment assessment and dementia risk prediction. We present detailed results comparing our model to established methods, followed by ablation studies and downstream clinical analyses.

3.2 Result for comparative analysis of WMH-DualTasker with segmentation and regression Baseline Models

Table 2 presents the performance of WMH-DualTasker compared to supervised segmentation methods across three datasets. On the MACC-Test dataset, WMH-DualTasker achieved a Dice score of 0.538±0.175, comparable to supervised methods like LST-LPA (0.551±0.230) [10] and superior to LST-LGA (0.403±0.253) [9]. Notably, WMH-DualTasker outperformed all methods in terms of Absolute Volume Difference (AVD), achieving 0.367±0.256 compared to the next best, Samseg [25,26], at 0.513±0.284.

On the SINGER dataset, WMH-DualTasker demonstrated the highest Dice score (0.608±0.146) and lowest AVD (0.365±0.277) among all methods, outperforming LST-LPA (Dice: 0.582±0.178, AVD: 0.431±0.344) and UBO-detector (Dice: 0.407±0.178, AVD: 0.416±0.226) [11]. Similarly, for the MICCAI-WMH dataset, our method achieved competitive Dice scores (0.602±0.222) and the best AVD (0.338±0.252), compared to LST-LPA (Dice: 0.632±0.188, AVD: 0.467±0.357).

Although nnU-Net [27] shows a higher Dice score than WMH-DualTasker on the MACC-Test (0.617±0.201 vs 0.538±0.175), WMH-DualTasker achieve better AVD evaluating WMH segmentation [28] (0.367±0.256 vs 0.461±0.302). The superior AVD

performance of WMH-DualTasker suggests its better capability in estimating overall WMH burden, which is crucial for clinical applications. Qualitatively, we also demonstrated that segmentation (Figure 4E) from WMH-DualTasker can accurately delineate the WMH from healthy tissue and is even very similar to our gold-standard reference (Figure 4F) from our radiologists.

For the ARWMC regression task (Table 3), WMH-DualTasker significantly outperformed standard 3D convolutional backbones [20,29]. On the MACC-Test dataset, our method achieved a mean absolute error (MAE) of 1.220±1.560, compared to 1.450±1.284 for the next best method SFCN. The Pearson's correlation (R) of 0.933±0.02 and Cohen's kappa (κ) of 0.94±0.04 for WMH-DualTasker were substantially higher than all other methods (next best: SFCN with R=0.854±0.03, κ=0.82±0.06), indicating excellent agreement with radiologists' visual ratings.

On the SINGER dataset, WMH-DualTasker maintained its superior performance with an MAE of 1.880±2.070, R of 0.612±0.08, and κ of 0.77±0.04, significantly outperforming other methods (next best: SFCN with MAE=3.274±2.368, R=0.420±0.19, κ=0.72±0.04). These results demonstrate the robust generalization of our method across different datasets and its potential as a reliable tool for automated ARWMC rating.

3.3 Result for ablation studies

To assess the effectiveness of each component in our proposed WMH-DualTasker approach, we conducted ablation studies, as shown in Table 4. The results revealed several important insights. First, using only the Class Activation Map (CAM) proved insufficient for satisfactory performance (Dice score: 0.133±0.109), underscoring the necessity of the hyperintense map for effective segmentation. Enhancing the resolution of CAM significantly improved segmentation by preserving more details, demonstrating its critical role (Dice score increased

to 0.312±0.180). Notably, different backbone networks provided comparable results under this enhancement (e.g., ResNet-10: 0.300±0.152, ResNet-18: 0.283±0.184). Second, the incorporation of self-supervised equivalent regularization loss led to notable improvements in both segmentation and regression performance (Dice: 0.474±0.233, MAE: 1.510±1.660 with scale transformation). This improvement was attributed to the enhanced quality of CAM, which more accurately reflected WMH locations, aiding in more precise Age-Related White Matter Changes (ARWMC) predictions. Specific affine transformations, such as rotation and flipping, further enhanced performance (Dice: 0.514±0.212, MAE: 1.220±1.560 with scale and flip), although combining all transformations indiscriminately did not yield additional benefits (Dice: 0.466±0.221, MAE: 1.467±1.872 with scale, flip, rotate, and translate). Finally, while Conditional Random Fields (CRF) provided incremental improvements (final Dice: 0.538±0.175), the use of more advanced CAM methods did not necessarily result in better performance (Grad-CAM: 0.503±0.196, Grad-CAM++: 0.527±0.189). Collectively, these findings highlight the critical components and configurations that contribute to the efficacy of the WMH-DualTasker model.

These findings highlight the critical components and configurations that contribute to the efficacy of the WMH-DualTasker model, with the combination of high-resolution CAM, hyperintense map, and self-supervised regularization providing the optimal performance.

3.4 Result for downstream validation analyses

3.4.1 Downstream analysis in AD cohort (ADNI dataset)

Our analysis revealed that incorporating WMH volume significantly enhanced the accuracy of cognitive impairment classification, prediction of disease progression, and estimation of cognitive performance scores (Figure 5).

The addition of WMH volume derived from WMH-DualTasker to clinical variables substantially improved the discrimination between Mild Cognitive Impairment (MCI) and Cognitively Normal (CN) subjects, increasing the Area Under the Curve (AUC) from 0.681 to 0.711 (p < 0.001). Combining both WMH volume and ARWMC scores further improved classification performance, achieving the highest AUC of 0.718 (Figure 5A).

WMH-DualTasker-derived biomarkers also enhanced the prediction of MCI conversion, with the AUC increasing from 0.618 to 0.652 (p < 0.001) for distinguishing between MCI converters and non-converters. This improvement underscores the potential of WMH features in identifying individuals at higher risk of progressing to dementia (Figure 5B).

Importantly, WMH volume predictions from WMH-DualTasker consistently outperformed those from nnUNet across both classification tasks, highlighting the robustness of our weakly supervised approach.

For continuous clinical outcomes (MMSE, CDRSB, and ADAS scores), incorporating WMH biomarkers consistently improved prediction correlations. The most notable improvements were observed for CDRSB and ADAS predictions, where the combination of clinical variables with WMH volume and ARWMC scores from WMH-DualTasker yielded the highest correlations (Figure 5C).

These comprehensive results demonstrate the added value of WMH biomarkers derived from WMH-DualTasker in assessing cognitive impairment and predicting disease progression. The consistent improvements across various cognitive measures and prediction tasks underscore the clinical utility of our approach, suggesting its potential as a valuable tool in both research and clinical settings for more accurate cognitive assessment and prognosis in the context of neurodegenerative diseases.

3.4.2 Downstream analysis in a community-based cohort (SINGER dataset)

To assess the utility of WMH-DualTasker in a general population setting, we analyzed the SINGER dataset, revealing a significant positive correlation between WMH volume predicted by WMH-DualTasker and the CAIDE Dementia Risk Score [30-32] (Spearman's rho = 0.19, p < 0.001). This finding supports the potential utility of WMH volume as a biomarker for future dementia risk in community-based populations.

The observed correlation, while modest, is clinically meaningful given the complex, multifactorial nature of dementia risk. It suggests that WMH-DualTasker can provide valuable information for assessing cognitive health in general populations, complementing traditional risk factors captured by the CAIDE score [33]. These findings, combined with the results from the ADNI cohort, demonstrate the versatility and clinical relevance of WMH-DualTasker across different population types and cognitive states.

3.5 Usage of WMH-DualTasker

The usage of WMH-DualTasker is relatively straightforward. Our GitHub repository provides examples to guide users for applying WMH-DualTasker. Our wrapper function will automatically perform the pre-processing for inputs with arbitrary resolution; training requires around 8GB of GPU memory on an RTX3090. Inference on a single subject is computationally efficient, requiring only a forward pass through the Deep Neural Network (DNN) followed by the application of Conditional Random Fields (CRF). This process is extremely fast, typically completed in a matter of seconds on standard GPU hardware and a few minutes on CPU.

4. Discussion

In this study, we present WMH-DualTasker, a deep learning-based algorithm that simultaneously performs white matter hyperintensity (WMH) segmentation and visual rating prediction. Our main contribution is the development of a model that requires only visual rating

scales for training, eliminating the need for labor-intensive, voxel-wise WMH annotations. This approach significantly reduces the time and expertise required for data preparation, making it particularly suitable for large-scale brain imaging studies. WMH-DualTasker employs a technique that leverages activation maps from the regression task to guide WMH segmentation. We enhance this process by incorporating self-supervised regularization and integrating a hyperintense map, improving lesion localization and sensitivity. By balancing accuracy, efficiency, and clinical relevance, our approach advances automated WMH analysis, offering a comprehensive tool for assessing cerebrovascular health in both research and clinical settings. The dual functionality of WMH-DualTasker in providing both segmentation and visual ratings makes it a versatile solution for quantifying WMH burden, potentially accelerating research in neurodegenerative and cerebrovascular diseases.

Existing WMH segmentation methods have limitations. Unsupervised methods [34-38] (LST-LGA) leverage the inherent MRI data characteristics, but often lack precision in identifying subtle WMH nuances. Supervised deep learning approaches [9, 11-13, 24, 39-42] (LST-LPA, UBO detector, and nnUNet), on the other hand, show promising results, however they rely on manually annotated WMH masks, which are time-consuming and expensive to produce, making these methods impractical for large-scale applications. Finally, weakly supervised methods leverage various forms of annotations, such as bounding boxes [45, 46] and partial annotations [47-53]. However, these approaches face challenges with WMH segmentation due to the dispersed nature of WMH. Moreover, bounding boxes and scribbles, in particular, are ill-suited for capturing the scattered distribution of WMH without extensive manual effort.

To address these limitations, WMH-DualTasker leveraged the Age-Related White Matter Changes (ARWMC) visual rating scale only as a supervisory signal. By utilizing activation maps generated from the regression task as initialization for WMH segmentation,

WMH-DualTasker can significantly reduce the time and expertise required for data preparation. To enhance the localization capability of the activation map, we introduce an equivalent regularization loss which ensures that it learns transformation invariance, increasing its sensitivity to less distinguishable lesion regions while suppressing activations in non-lesion areas.

We evaluated WMH-DualTasker on both internal and external datasets. Our results demonstrated that WMH-DualTasker achieves performance comparable or superior to the existing supervised methods in WMH segmentation on both internal (MACC) and external (MICCAI-WMH and SINGER) validations. This is particularly noteworthy given that WMH-DualTasker operates with only image-level labels, whereas the existing supervised methods require pixel-level annotations. Moreover, WMH-DualTasker can jointly predict visual rating scores. We observed good to high Kappa agreement concerning the radiologists' ratings on the external validation (SINGER). This indicates WMH-DualTasker's potential to serve as a valuable reference tool for clinical experts. Overall, our extensive evaluation across multiple datasets confirms that WMH-DualTasker consistently produces accurate WMH volume measurements and ARWMC scores, establishing it as a reliable and generalizable tool for assessing WMH across both Asian and Caucasian populations.

Moreover, we demonstrated the clinical utility of WMH-DualTasker's outputs in both AD-specific (ADNI) and community-based (SINGER) settings. We observed that incorporating both WMH volume and ARWMC scores significantly improved the accuracy of MCI/CN classification and MCI conversion prediction. In the community-cohort (SINGER), we found a correlation between WMH volume and the CAIDE Dementia Risk Score, indicating the relevance of WMH-DualTasker's outputs in assessing future dementia risk in a general population.

.

Given these compelling results, WMH-DualTasker shows great potential for integration into radiology and research workflows. Users can directly apply the pre-trained model (from our github repo) to their T2-FLAIR MRI images without fine-tuning, obtaining rapid and consistent WMH segmentations and ARWMC scores. This out-of-the-box functionality is particularly valuable for clinical settings or large-scale studies where manual annotations are impractical. For instance, a hospital could implement WMH-DualTasker to assist radiologists in assessing stroke risk or monitoring cerebrovascular disease progression across numerous patients. Researchers conducting population-wide studies could efficiently process thousands of MRI scans to investigate relationships between WMH burden and various neurological conditions or cognitive decline. If users wish to optimize performance for their specific dataset, they can fine-tune the model using only visual rating scores (ARWMC) for a small subset of their data, without the need for time-consuming manual WMH annotations. Our GitHub repository provides comprehensive documentation and scripts for both direct application and fine-tuning, ensuring WMH-DualTasker can be easily integrated into diverse clinical and research environments.

Our study has several limitations. While we utilized the ARWMC scale due to its availability in our datasets, we did not validate other clinically prevalent scales, such as the Fazekas scale [7]. This analysis focused on cross-sectional data, and future research should explore the algorithm's performance on longitudinal data to assess WMH progression over time. Additionally, further clinical validation through prospective studies in real-world settings is essential to evaluate the model's performance in routine clinical workflows and its impact on clinical decision-making. These steps will help position WMH-DualTasker as a reliable tool for assessing and managing cerebrovascular disorders associated with WMH, offering valuable assistance to healthcare professionals.

5. Conclusion

In conclusion, WMH-DualTasker, a novel dual-task deep learning model with self-supervised consistency, achieved accurate and generalizable performance for automatic White Matter Hyperintensity (WMH) segmentation and visual rating prediction using T2-FLAIR MRI. This weakly-supervised approach significantly reduces the annotation burden by utilizing only image-level visual ratings for training, while effectively performing both voxel-wise segmentation and global ARWMC score prediction. The model's robustness was demonstrated across multiple datasets, including both disease-specific and community-based cohorts. Considering its effectiveness in joint segmentation and regression tasks, and its proven clinical utility in cognitive impairment assessment and dementia risk prediction, WMH-DualTasker represents a valuable tool for both clinical practice and scientific research. Its ability to process large-scale datasets efficiently makes it particularly suited for population-wide studies, potentially advancing our understanding of cerebral small vessel disease and cognitive disorders.

Declarations

decision-making related to the acceptance of this article for publication.

Ethics approval and consent: The local Institutional Review Boards at each participating institution approved participant recruitment and data collection for the ADNI project. Informed written consent from participants was obtained at each participating site. For up-to-date information, see www.adni-info.org.

Consent for publication: All co-authors consent to publish the manuscript.

Reference

1. Wardlaw JM, Smith EE, Biessels GJ, et al. Neuroimaging standards for research into small vessel disease and its contribution to ageing and neurodegeneration. Lancet Neurol. 2013;12(8):822-38.

2. Duering M, Biessels GJ, Brodtmann A, et al. Neuroimaging standards for research into small vessel disease—advances since 2013. Lancet Neurology. 2023;22(7):602-18.

3. Debette S, Markus HS. The clinical importance of white matter hyperintensities on brain magnetic resonance imaging: systematic review and meta-analysis. BMJ. 2010;341:c3666.

4. Prins ND, Scheltens P. White matter hyperintensities, cognitive impairment and dementia: an update. Nat Rev Neurol. 2015;11(3):157-65.

5. Wardlaw JM, Valdés Hernández MC, Muñoz-Maniega S. What are white matter hyperintensities made of? Relevance to vascular cognitive impairment. J Am Heart Assoc. 2015;4(6):e001140.

6. Scheltens P, Erkinjuntti T, Leys D, et al. White matter changes on CT and MRI: an overview of visual rating scales. Eur Neurol. 1998;39(2):80-9.

7. Wahlund LO, Barkhof F, Fazekas F, et al. A new rating scale for age-related white matter changes applicable to MRI and CT. Stroke. 2001;32(6):1318-22.

8. Kapeller P, Barber R, Vermeulen RJ, et al. Visual rating of age-related white matter changes on magnetic resonance imaging: scale comparison, interrater agreement, and correlations with quantitative measurements. Stroke. 2003;34(2):441-45.

9. Schmidt P, Gaser C, Arsic M, et al. An automated tool for detection of FLAIR-hyperintense white-matter lesions in multiple sclerosis. Neuroimage. 2012;59(4):3774-83.

10. Schmidt P. Bayesian inference for structured additive regression models for large-scale problems with applications to medical imaging. LMU; 2017.

11. Jiang J, Liu T, Zhu W, et al. UBO Detector—A cluster-based, fully automated pipeline for extracting white matter hyperintensities. Neuroimage. 2018;174:539-49.

12. Li H, Jiang G, Zhang J, et al. Fully convolutional network ensembles for white matter hyperintensities segmentation in MR images. Neuroimage. 2018;183:650-65.

13. Sundaresan V, Zamboni G, Rothwell PM, et al. Triplanar ensemble U-Net model for white matter hyperintensities segmentation on MR images. Med Image Anal. 2021;73:102184.

14. Wu Y, Ji F, Chong YF, et al. Deeply supervised network for white matter hyperintensities segmentation with transfer learning. In: Medical Imaging with Deep Learning; 2022.

15. Jack CR, Bernstein MA, Fox NC, et al. The Alzheimer's disease neuroimaging initiative (ADNI): MRI methods. J Magn Reson Imaging. 2008;27(4):685-91.

16. Kuijf HJ, Biesbroek JM, De Bresser J, et al. Standardized Assessment of Automatic Segmentation of White Matter Hyperintensities and Results of the WMH Segmentation Challenge. IEEE Trans Med Imaging. 2019;38(11):2556-68.

17. Xu X, Chew KA, Wong ZX, et al. The Singapore geriatric intervention study to reduce cognitive decline and physical frailty (SINGER): study design and protocol. J Prev Alzheimers Dis. 2022;9(1):40-8.

18. Smith SM, Jenkinson M, Woolrich MW, et al. Advances in functional and structural MR image analysis and implementation as FSL. Neuroimage. 2004;23 Suppl 1:S208-S19.

19. Zhou B, Khosla A, Lapedriza A, et al. Learning deep features for discriminative localization. In: Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition; 2016. p. 2921-9.

20. Peng H, Gong W, Beckmann CF, et al. Accurate brain age prediction with lightweight deep neural networks. Med Image Anal. 2021;68:101871.

21. Wang Y, Zhang J, Kan M, et al. Self-Supervised Equivariant Attention Mechanism for Weakly Supervised Semantic Segmentation. In: Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR); 2020.

22. Kamnitsas K, Ledig C, Newcombe VF, et al. Efficient multi-scale 3D CNN with fully connected CRF for accurate brain lesion segmentation. Med Image Anal. 2017;36:61-78.

23. Krähenbühl P, Koltun V. Efficient Inference in Fully Connected CRFs with Gaussian Edge Potentials. In: Advances in Neural Information Processing Systems; 2011.

24. Loshchilov I, Hutter F. Decoupled Weight Decay Regularization. In: International Conference on Learning Representations; 2018.

25. Cerri S, Greve DN, Hoopes A, et al. An open-source tool for longitudinal whole-brain and white matter lesion segmentation. Neuroimage Clin. 2023;38:103354.

26. Cerri S, Puonti O, Meier DS, et al. A contrast-adaptive method for simultaneous whole-brain and lesion segmentation in multiple sclerosis. Neuroimage. 2021;225:117471.

27. Isensee F, Jaeger PF, Kohl SAA, et al. nnU-Net: a self-configuring method for deep learning-based biomedical image segmentation. Nat Methods. 2021;18(2):203-11.

28. Maier-Hein L, et al. Metrics reloaded: recommendations for image analysis validation. Nat Methods. 2024;21(2):195-212.

29. Hara K, Kataoka H, Satoh Y. Can Spatiotemporal 3D CNNs Retrace the History of 2D CNNs and ImageNet? In: Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition (CVPR); 2018. p. 6546-55.

30. Sindi S, Calov E, Fokkens J, et al. The CAIDE dementia risk score app: the development of an evidence-based mobile application to predict the risk of dementia. Alzheimers Dement Diagn Assess Dis Monit. 2015;1(3):328-33.

31. Keller JA, Kant IMJ, Slooter AJ, et al. Different cardiovascular risk factors are related to distinct white matter hyperintensity MRI phenotypes in older adults. Neuroimage Clin. 2022;35:103131.

32. McBride CA, Russom Z, Achenbach E, et al. Cardiovascular profiles associated with white matter hyperintensities in healthy young women. Front Physiol. 2023;13:979899.

33. Jiang J, Yao K, Huang X, et al. Longitudinal white matter hyperintensity changes and cognitive decline in patients with minor stroke. Aging Clin Exp Res. 2022;34(5):1047-54.

34. Damangir S, Westman E, Simmons A, et al. Reproducible segmentation of white matter hyperintensities using a new statistical definition. Magn Reson Mater Phy. 2017;30:227-37.

35. Zhan T, Zhan Y, Liu Z, et al. Automatic method for white matter lesion segmentation based on T1-fluid-attenuated inversion recovery images. IET Comput Vis. 2015;9(4):447-55.

36. Bowles C, Qin C, Guerrero R, et al. Brain lesion segmentation through image synthesis and outlier detection. Neuroimage Clin. 2017;16:643-58.

37. Wang R, Li C, Wang J, et al. Automatic segmentation and volumetric quantification of white matter hyperintensities on fluid-attenuated inversion recovery images using the extreme value distribution. Neuroradiology. 2015;57:307-20.

38. Iorio M, Spalletta G, Chiapponi C, et al. White matter hyperintensities segmentation: a new semi-automated method. Front Aging Neurosci. 2013;5:76.

39. Ronneberger O, Fischer P, Brox T. U-net: Convolutional networks for biomedical image segmentation. In: Medical Image Computing and Computer-Assisted Intervention—MICCAI 2015, Proceedings, Part III; 2015. p. 234-41.

40. Oktay O, Schlemper J, Folgoc LL, et al. Attention u-net: Learning where to look for the pancreas. arXiv preprint arXiv:1804.03999; 2018.

41. Zhou Z, Rahman Siddiquee MM, Tajbakhsh N, Liang J. Unet++: A nested u-net architecture for medical image segmentation. In: Medical Image Computing and Computer-Assisted Intervention—MICCAI 2018, Proceedings, Part III 18. Springer; 2018. p. 3-11.

42. Park G, Hong J, Duffy BA, et al. White matter hyperintensities segmentation using the ensemble U-Net with multi-scale highlighting foregrounds. Neuroimage. 2021;237:118140.

43. Wei Y, Liang X, Chen Y, et al. Stc: A simple to complex framework for weakly-supervised semantic segmentation. IEEE Trans Pattern Anal Mach Intell. 2016;39(11):2314-20.

44. Ahn J, Kwak S. Learning pixel-level semantic affinity with image-level supervision for weakly supervised semantic segmentation. In: Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition; 2018. p. 4981-90.

45. Kervadec H, Dolz J, Wang S, et al. Bounding boxes for weakly supervised segmentation: Global constraints get close to full supervision. In: Medical imaging with deep learning; 2020. p. 365-81.

46. Xie Y, Zhang Z, Chen S, Qiu C. Detect, Grow, Seg: A weakly supervision method for medical image segmentation based on bounding box. Biomed Signal Process Control. 2023;86:105158.

47. Kervadec H, Dolz J, Tang M, et al. Constrained-CNN losses for weakly supervised segmentation. Med Image Anal. 2019;54:88-99.

48. Chen Z, Tian Z, Zhu J, et al. C-cam: Causal cam for weakly supervised semantic segmentation on medical image. In: Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition; 2022. p. 11676-85.

49. Roth HR, Yang D, Xu Z, et al. Going to extremes: weakly supervised medical image segmentation. Mach Learn Knowl Extr. 2021;3(2):507-24.

50. Dubost F, Adams H, Bortsova G, et al. 3D regression neural network for the quantification of enlarged perivascular spaces in brain MRI. Med Image Anal. 2019;51:89-100.

51. Dubost F, Bortsova G, Adams H, et al. Gp-unet: Lesion detection from weak labels with a 3d regression network. In: Medical Image Computing and Computer Assisted Intervention—MICCAI 2017, Proceedings, Part III; 2017. p. 214-21.

52. Patel G, Dolz J. Weakly supervised segmentation with cross-modality equivariant constraints. Med Image Anal. 2022;77:102374.

53. Zhou Y, Chen H, Li Y, et al. Multi-task learning for segmentation and classification of tumors in 3D automated breast ultrasound images. Med Image Anal. 2021;70:101918.

54. Lei W, Su Q, Jiang T, et al. One-shot Weakly-Supervised Segmentation in 3D Medical Images. IEEE Trans Med Imaging. 2023. DOI: 10.1109/TMI.2023.3294975.

55. Selvaraju RR, Cogswell M, Das A, et al. Grad-cam: Visual explanations from deep networks via gradient-based localization. In: Proceedings of the IEEE International Conference on Computer Vision; 2017. p. 618-26.

56. Chattopadhay A, Sarkar A, Howlader P, Balasubramanian VN. Grad-cam++: Generalized gradient-based visual explanations for deep convolutional networks. In: 2018 IEEE Winter Conference on Applications of Computer Vision (WACV); 2018. p. 839-47.

57. Valverde S, Oliver A, Roura E, et al. Automated tissue segmentation of MR brain images in the presence of white matter lesions. Med Image Anal. 2017;35:446-57.

Table 1

The overview of the scanner and image details in this study.

Dataset	Usage	Site	Cohort	Scanner	Sequence	Voxel resolution (mm^3)	Number of subjects / cases with mask / cases with visual rating
MACC	Train / Validation	Singapore	SVD	3T Siemens Trio Tim	T2-FLAIR	1.0x1.0x3.0	660 / 60 / 660
SINGER	Validation	Singapore	SVD	3T Siemens Trio Tim	T2-FLAIR	1.0x1.0x1.0	60 / 60 / 60
SINGER*	Downstream Analysis	Singapore	SVD	3T Siemens Trio Tim	T2-FLAIR	1.0x1.0x1.0	470 / 0 / 0
MICCAI-UMC Utrecht	Validation	Singapore	SVD	3T Phillips Achieva	T2-FLAIR	1.0x1.0x3.0	20 / 20 / 20
MICCAI-NUHS	Validation	Netherlands	SVD	3T Siemens Trio Tim	T2-FLAIR	1.0x1.0x3.0	20 / 20 / 20
MICCAI-VU Amsterdam	Validation	Netherlands	SVD	3T GE Signa HDxT	T2-FLAIR	1.0x1.0x1.2	20 / 20 / 20
ADNI	Downstream Analysis	US	SVD, Control	3T Siemens Trio Tim 3T Siemens Skyra 3T Philips Ingenia 1.5T Siemens Avanto	T2-FLAIR	1.0x1.0x1.2	394 / 0 / 0

Note. SVD = Small Vessel Disease; T2-FLAIR = T2-weighted Fluid-Attenuated Inversion Recovery.

Table 2

Comparative evaluation of segmentation across all WMH segmentation tools.

Dataset		MACC-Test (40)		SINGER (64)		MICCAI-WMH (60)	
Methods	Mask	Dice ↑	AVD ↓	Dice ↑	AVD ↓	Dice ↑	AVD ↓
LST-LGA	Yes	$0.403_{\pm0.253}$	$0.580_{\pm0.304}$	$0.527_{\pm0.209}$	$0.401_{\pm0.260}$	$0.566_{\pm0.221}$	$0.378_{\pm0.266}$
LST-LPA		$0.551_{\pm0.230}$	$0.509_{\pm0.316}$	$0.582_{\pm0.178}$	$0.431_{\pm0.344}$	$\mathbf{0.632}_{\pm0.188}$	$0.467_{\pm0.357}$
UBO-detector		$0.340_{\pm0.221}$	$0.614_{\pm0.371}$	$0.407_{\pm0.178}$	$0.416_{\pm0.226}$	$0.572_{\pm0.183}$	$0.480_{\pm0.316}$
Samseg		$0.403_{\pm0.240}$	$0.513_{\pm0.284}$	$0.523_{\pm0.211}$	$0.407_{\pm0.228}$	$0.532_{\pm0.256}$	$0.428_{\pm0.270}$
nnU-Net*		$\mathbf{0.617}_{\pm0.201}$	$0.461_{\pm0.302}$	$0.605_{\pm0.201}$	$0.507_{\pm0.240}$	NA	
WMH-DualTasker	No	$0.538_{\pm0.175}$	$\mathbf{0.367}_{\pm0.256}$	$\mathbf{0.608}_{\pm0.146}$	$\mathbf{0.365}_{\pm0.277}$	$0.602_{\pm0.222}$	$\mathbf{0.338}_{\pm0.252}$

Note. WMH = White Matter Hyperintensities; Dice = Dice similarity coefficient; AVD = Absolute Volume Difference. For each metric, the method with the best performance is highlighted in bold. ↑ indicates higher values are better, ↓ indicates lower values are better. Mask:

- nnU-Net is trained with the MICCAI-WMH dataset. Therefore, we cannot report the evaluation result on the same dataset. The "Mask" column refers to whether the training requires WMH masks.

Table 3

Comparative Evaluation of Regression Across All Methods.

Dataset	MACC-Test (40)			SINGER (64)		
Method	MAE↓	R↑	κ↑	MAE↓	R↑	κ↑
ResNet-10	$1.550_{\pm 1.396}$	$0.806_{\pm 0.04}$	$0.55_{\pm 0.12}$	$3.340_{\pm 2.499}$	$0.257_{\pm 0.12}$	$0.42_{\pm 0.04}$
ResNet-18	$1.800_{\pm 1.249}$	$0.795_{\pm 0.03}$	$0.67_{\pm 0.08}$	$4.000_{\pm 2.987}$	$0.171_{\pm 0.14}$	$0.55_{\pm 0.04}$
ResNet-34	$1.550_{\pm 0.1322}$	$0.828_{\pm 0.02}$	$0.71_{\pm 0.06}$	$3.142_{\pm 2.720}$	$0.268_{\pm 0.15}$	$0.60_{\pm 0.04}$
SFCN	$1.450_{\pm 1.284}$	$0.854_{\pm 0.03}$	$0.82_{\pm 0.06}$	$3.274_{\pm 2.368}$	$0.420_{\pm 0.19}$	$0.72_{\pm 0.04}$
WMH-DualTasker	$\mathbf{1.220}_{\pm 1.560}$	$\mathbf{0.933}_{\pm 0.02}$	$\mathbf{0.94}_{\pm 0.04}$	$\mathbf{1.880}_{\pm 2.070}$	$\mathbf{0.612}_{\pm 0.08}$	$\mathbf{0.77}_{\pm 0.04}$

Note. MAE = Mean Absolute Error; R = Pearson's correlation; κ = Cohen's kappa. For each metric, the method with the best performance is highlighted in bold. ↑: Higher values indicate better performance. ↓:Lower values indicate better performance.

Table 4

Ablation studies results.

Backbone	CAM method	CAM resolution	Transform	Hyperintense Map	CRF	Dice↑	MAE↓
SFCN	CAM	$32 \times 32 \times 16$	-	-	-	$0.133_{\pm 0.109}$	$1.730_{\pm 0.970}$
SFCN	CAM	$32 \times 32 \times 16$	-	-	Yes	$0.154_{\pm 0.126}$	$1.730_{\pm 0.970}$
SFCN	CAM	$32 \times 32 \times 16$	-	Yes	-	$0.204_{\pm 0.092}$	$1.730_{\pm 0.970}$
SFCN	CAM	$64 \times 64 \times 32$	-	Yes	-	$0.312_{\pm 0.180}$	$1.700_{\pm 1.308}$
ResNet-10	CAM	$64 \times 64 \times 32$	-	Yes	-	$0.300_{\pm 0.152}$	$1.550_{\pm 1.532}$
ResNet-18	CAM	$64 \times 64 \times 32$	-	Yes	-	$0.283_{\pm 0.184}$	$1.750_{\pm 1.670}$
ResNet-34	CAM	$64 \times 64 \times 32$	-	Yes	-	OOM	OOM
SFCN	CAM	$64 \times 64 \times 32$	S	Yes	-	$0.474_{\pm 0.233}$	$1.510_{\pm 1.660}$
SFCN	CAM	$64 \times 64 \times 32$	S+F	Yes	-	$0.514_{\pm 0.212}$	$1.220_{\pm 1.560}$
SFCN	CAM	$64 \times 64 \times 32$	S+R	Yes	-	$0.489_{\pm 0.252}$	$1.430_{\pm 1.764}$
SFCN	CAM	$64 \times 64 \times 32$	S+T	Yes	-	$0.462_{\pm 0.247}$	$1.433_{\pm 1.775}$
SFCN	CAM	$64 \times 64 \times 32$	S+F+R+T	Yes	-	$0.466_{\pm 0.221}$	$1.467_{\pm 1.872}$
WMH-DualTasker	CAM	$64 \times 64 \times 32$	S+F	Yes	Yes	$0.538_{\pm 0.175}$	$1.220_{\pm 1.560}$
WMH-DualTasker	Grad-CAM	$64 \times 64 \times 32$	S+F	Yes	Yes	$0.503_{\pm 0.196}$	$1.220_{\pm 1.560}$
WMH-DualTasker	Grad-CAM++	$64 \times 64 \times 32$	S+F	Yes	Yes	$0.527_{\pm 0.189}$	$1.220_{\pm 1.560}$

Note. S: scale, F: flip, R: rotate, T: translate, HM: hyperintense map, CRF: conditional random field, OOM: out of memory. Dice: dice score for WMH segmentation; MAE: mean absolute error for ARWMC regression. ↑: Higher values indicate better performance. ↓:Lower values indicate better performance.

Figure legend

Figure 1: Illustration of WMH across varying loads. This figure presents a series of axial brain MRI scans from subjects, showcasing a gradient of WMH from minimal to pronounced, organized from left to right. In the top row, WMH is shown as high-intensity signals on T2-FLAIR MRI scans, contrasting sharply with the surrounding brain tissue and visually pinpointing the regions impacted by WMH. The bottom row complements this with WMH segmentation masks, rendered in striking red, to provide a precise visual quantification of the WMH load. We observe that WMH typically manifests as irregularly shaped, hyperintense lesions located in both periventricular and deep white matter regions on MRI scans. The final row features the ARWMC visual rating scale, introducing a standardized, qualitative approach to assess WMH severity. MRI: Magnetic Resonance Imaging. WMH: white matter hyperintensities. ARWMC: Age-Related White Matter Changes.

Figure 2. Comprehensive dataset utilization and downstream analysis framework for WMH-DualTasker. The schematic illustrates our multi-dataset approach and analytical pipeline. (A) For model development and evaluation, we utilized three primary datasets (MACC, SINGER, and MICCAI-WMH). The MACC dataset is divided into training, validation, and testing subsets, while SINGER and MICCAI-WMH are used as external test sets. Color-coding indicates the type of annotation available: yellow for visual ratings only, green for both visual ratings and manual segmentation masks, and blue for manual segmentation masks only. Subsequently, (B) We validate the utility of WMH-DualTask by performing our downstream analysis, demonstrating the application of WMH-DualTasker to two distinct cohorts: the ADNI dataset (a disease-specific cohort) and the SINGER dataset (a community-based cohort). These analyses encompass a range of clinically relevant tasks, including cognitive impairment classification, dementia conversion prediction, and assessment of dementia risk, showcasing the model's versatility and potential clinical utility across diverse population settings.

Figure 3. Schematic diagram of the WMH-DualTasker training and inference processes.
The top portion of the diagram depicts the training phase, supervised by WMH visual rating and self-supervised equivalent regularization loss. The bottom part illustrates the inference process, where the enhanced CAM is integrated with the Hyperintense Map (HM) to produce the final WMH segmentation. CAM: Class Activation Map.

(A) FLAIR　　　　(B) Baseline CAM　(C) Enhanced CAM　(D) Hyperintense Map　(E) **WMH prediction**　(F) Ground truth

Figure 4. WMH-DualTasker generates high-quality WMH masks by integrating enhanced CAM and Hyperintense Maps. Displayed across six rows for three subjects, this figure presents a side-by-side comparison of axial slices and their zoomed-in counterparts to enhance detail visualization. Progressing from left to right, the images showcase (A) FLAIR images, (B) baseline Class Activation Maps (CAM), (C) enhanced CAMs, (D) Hyperintense Maps, (E) WMH predictions from WMH-DualTasker, and finally, (F) the reference standard annotations. The baseline CAM identifies potential WMH regions but with blurry boundaries,

in contrast to the enhanced CAM, which delineates these regions with greater precision. Although Hyperintense Maps accurately outline WMH with clear boundaries, they are prone to including false positives. The integration of enhanced CAM with Hyperintense Maps by WMH-DualTasker culminates in the generation of high-quality WMH masks, achieving a similarity to manual annotations and substantially reducing false positives. WMH: White Matter Hyperintensities; CAM: Class Activation Map;

Figure 5. WMH biomarkers from WMH-DualTasker improve cognitive function prediction in Alzheimer's disease. (Panel A-B) Bar charts show the mean AUC across four models for (A) classification between CN and MCI subjects and (B) classification of MCI converters versus non-converters. Models compared are clinical factors only (pink), clinical + WMH volume from nnU-Net (green), clinical + WMH volume from WMH-DualTasker (blue), and clinical + WMH volume and ARWMC from WMH-DualTasker (purple). Error bars indicate standard deviation across 10-fold cross-validation. (Panel C) Box plots display correlation coefficients between predicted cognitive scores and actual scores for MMSE, ADAS-Cog, and CDRSB. Each box plot represents the distribution of correlation values obtained from 10-fold cross-validation for each model. Significant differences between models, identified through paired T-tests, are denoted by brackets. MMSE: Mini-Mental State

Examination; ADAS-Cog: Alzheimer's Disease Assessment Scale-Cognitive Subscale; CDRSB: Clinical Dementia Rating Scale Sum of Boxes; ARWMC: Age-Related White Matter Changes. CN: Cognitively Normal. MCI: Mild Cognitive Impairment. AUC: Area Under the Curve.

Figure 6. Higher WMH volumes are associated with increased dementia risk in a community-based cohort. Scatter plot illustrating the positive correlation between WMH volume predicted by WMH-DualTasker and CAIDE Risk Score in the SINGER cohort. This relationship suggests that increased WMH burden, as quantified by our model, is linked to higher risk of future dementia in a general population setting. CAIDE: Cardiovascular Risk Factors, Aging, and Incidence of Dementia.