Impacts of Structural Properties of Myosin II Filaments on Force Generation

Shihang Ding[1†], Pei-En Chou[2†], Shinji Deguchi[1*], Taeyoon Kim[3*]

[1]Graduate School of Engineering Science, Osaka University

Toyonaka, Osaka, Japan

[2] School of Mechanical Engineering, Purdue University

585 Purdue Mall, West Lafayette, IN 47907, USA

[3] Weldon School of Biomedical Engineering, Purdue University

206 Martin Jischke Drive, West Lafayette, IN 47907, USA

† equal contribution

* co-corresponding author

ABSTRACT

Cells need intracellular forces for their physiological functions, such as migration, cytokinesis, and morphogenesis. The actin cytoskeleton generates a large fraction of the forces via interactions between cytoskeletal components, such as actin filament (F-actin), myosin, and actin cross-linking proteins (ACPs). Myosin II plays the most important role in cellular force generation. Myosin II molecules self-assemble into filaments with different structures depending on myosin II isoforms and other conditions such as pH and ionic concentration. It has remained elusive how force generation in actomyosin structures is affected by the architecture of myosin II filaments. In this study, we employed an agent-based model to investigate the effects of the structural properties of myosin II filaments on force generation in disorganized actomyosin structures. We demonstrated that the magnitude of forces and the efficiency of force generation can vary over a wide range depending on the number and spatial distribution of myosin II filaments. Further, we showed that the number of myosin heads and the length of a bare zone at the center of myosin II filaments without heads highly affect the force generation process in bundles and networks. Our study provides insights into understanding the roles of the structural properties of myosin II filaments in actomyosin contractility.

INTRODUCTION

Cells require forces for a wide variety of physiological functions, including cytokinesis, migration, and morphogenesis [1]. It is well-known that mechanical forces are produced mainly by molecular interactions between filamentous actin (F-actin) and myosin II motor proteins in the actin cytoskeleton [2]. Myosin II motors walk toward the barbed end of F-actin using chemical energy stored in adenosine triphosphate (ATP). Myosin II molecules consist of two heads with a long tail, and they self-assemble into filamentous structures [3]. There are three isoforms of myosin II: non-muscle, smooth muscle, and skeletal muscle myosins [4]. These myosin isoforms form different filamentous structures whose length ranges from ~0.3 µm to ~1.5 µm with ~56 to ~800 heads [5-12] . Unlike smooth muscle myosin forming a side-polar filament, non-muscle and skeletal muscle myosins form a bipolar filament with two sets of myosin heads located at both ends of the filament and a bare zone at its center whose length is ~160 nm. The number of myosin II molecules in the myosin filament also varies depending on conditions, such as pH level and ionic concentration [13-16]. Individual myosin II heads have a relatively low duty ratio, meaning that they spend only a small fraction of their lifetime in the bound state unlike processive (i.e., high duty ratio) motors, such as myosin V or kinesin [17]. Thus, heads in a single myosin II molecule cannot walk along F-actin over a long distance by themselves. Myosin II circumvents this issue by forming the filamentous structure with multiple heads. If a myosin II filament has a sufficient number of heads, there are always a few myosin heads bound to F-actin, so the myosin II filament can remain proximal to the F-actin.

Heads with opposite polarities in the myosin II filament pull F-actins in opposite directions by walking toward the barbed ends of those F-actins, developing both tensile and compressive forces in disorganized actomyosin structures which are different from sarcomere found in muscle cells [18]. Theoretical and computational studies demonstrated that F-actin is easily buckled by compressive forces, leaving net tensile forces in disorganized bundles or networks that can mediate diverse contractile behaviors [19-21]. Various reconstituted actomyosin systems have been employed to illuminate how contraction and force generation emerge from interactions between F-actins, myosin II filaments, and actin cross-linking proteins (ACPs) [22-27]. Although they provided valuable insights, it has been understood poorly how various structural properties of myosin II filaments affect contraction or force generation in disorganized actomyosin structures.

Several computational models have been used to study the actomyosin contractility [28-34]. However, there were several limitations in the models. For example, some of the models treated myosin II filaments as either points [34, 35], rods only with two binding sites [31, 36, 37], or force dipoles [38]. Such drastically simplified structures significantly differ from the real structure of the myosin II filament. In this study, we used our well-established agent-based model with detailed descriptions of the structure of myosin bipolar filaments [20, 39-43], to illuminate how the number, length, bare zone size, and spatial distribution of myosin bipolar filaments influence force generation in disorganized actomyosin bundles and networks.

MATERIALS AND METHODS

Model overview

The detailed explanations about our model and all parameters used in the model are explained in Supplementary Text and Supplementary Tables. Our model consists of only three key cytoskeletal elements – F-actin, motor, and ACPs – among >100 proteins found in the actomyosin structures of cells [44]. They are simplified via cylindrical segments (Fig. 1A); F-actin is represented by serially connected segments with 140 nm in length. Each ACP comprises two arms with 23.5 nm in length connected to its center point. To mimic the structure of bipolar filaments, each motor has a backbone, consisting of serially linked segments, and two arms on each endpoint of the backbone segments that represent 8 myosin heads ($N_h = 8$). The reference length of backbone segments (L_{MB}) is 42 nm. The displacements of all the cylindrical segments at each time step are calculated by the Langevin equation and the forward Euler integration scheme. Deterministic forces in the Langevin equation include extensional and bending forces that maintain the equilibrium lengths of segments and equilibrium angles formed by segments, respectively, as well as a repulsive force exerted on overlapping pairs of actin segments for considering volume-exclusion effects.

Using the model, we simulate three types of systems: two filaments, bundles, and networks. In the two-filament simulations, a pair of anti-parallel F-actins with 19 μm in length are allocated in a rectangular computational domain (5×5×20 μm) with the periodic boundary condition (PBC) in x and y directions and the repulsive boundary condition in z direction. The barbed end of F-actins is clamped to two finite boundaries normal to the z direction. In the bundle simulations, we employ the same rectangular domain with the PBC in all directions (Fig. 1B). As in our previous study [42], we allocate F-actins in

specific x and y coordinates. The number of possible positions for the allocation in x or y direction is defined by N_F, and spacing between adjacent coordinates in each direction is set to 27 nm. In each set of x and y positions, we position two F-actins with 9 μm in length in random z position with random polarity. Thus, the total number of F-actins in the bundle is $2N_F^2$. In addition, the network simulations are conducted using a thin rectangular computational domain (20×20×0.1 μm) with the PBC in x and y directions and the repulsive boundary condition in z direction (Fig. 1C). In the domain, F-actins with 10 μm in length are allocated randomly in terms of positions and orientations. At the beginning of all simulations, while F-actins remain stationary, ACPs bind to F-actins to form permanent cross-linking points, and the arms of motors formed via the self-assembly of backbone segments bind to F-actins. After that, F-actins are allowed to move, and motor arms start walking and unbinding at force-dependent rates.

Variations in motor structures

In our study, we vary three structural properties of motors: the number of arms per motor (N_a), the length of the bare zone (L_{bz}), and spacing between motor arms (L_{sp}). We change the size of motors in three different ways. In the first way, N_a is increased by connecting more segments with arms for a backbone, whereas L_{bz} and L_{sp} are equal to L_{MB} (Fig. 1E, i). In the second way, L_{bz} is increased by adding more segments without arms to the bare zone, whereas N_a is unchanged, and L_{sp} is equal to L_{MB} (Fig. 1E, ii). In the third way, L_{sp} and L_{bz} are increased by making L_{MB} higher, whereas N_a is unchanged (Fig. 1E, iii).

Measurement of contractile forces

In the two-filament simulations and the bundle simulations, we measure tensile forces generated by the systems as follows. First, all segments crossing the cross-sections of the domain located every 200 nm in the z direction, which can be for either of F-actin, motor arm, motor backbone, or ACP arm, are identified. Then, the z component of spring forces acting on those segments is summed. Using these sums, four curves can be drawn as a function of z to show which segment supports more or less tensile loads in each z position. Note that at equilibrium, the sum of four forces is very similar regardless of z position, meaning that a large fraction of forces developed by motors act as spring forces rather than bending forces. The average of the sums calculated on all cross-sections at a steady state is considered the total force acting on the system, F_{tot}. In the network simulations, tensile forces are measured in a similar manner, using 20 cross-sections of the domain located evenly in the x and y directions. x or y component of tensile forces acting on all segments crossing each cross-section is summed at a steady state. The average of sums calculated on 40 cross-sections is considered F_{tot}.

RESULTS

The distribution of motors and ACPs plays a key role in force generation

A previous in vitro study reported that tension developed in a thin bundle was almost directly proportional to the number of myosin heads in each motor, not to the number of motors [4]. Although their experiments did not include any ACP, their

theoretical explanation was based on an assumption that the bundle consists of serially connected contractile units like sarcomeres in muscle cells [4]. They admitted that these contractile units do not have structural analogs in their disordered actomyosin bundles.

To verify their hypothesis, we first used a simple minimal model composed of two anti-parallel F-actins, two motors, and 16 ACPs (Figs. 1A and 2). The number of arms per motor was set to N_a = 24. We ran 20 simulations with random z positions of motors and ACPs and found that the total force generated by the system, F_{tot}, fell into the following range (Fig. 2A):

$$0.5 F_M^{max} \leq F_{tot} \leq F_M^{max}. \tag{1}$$

where F_M^{max} is the maximal force that all motors can generate in this system:

$$F_M^{max} = \frac{1}{2} \sum_i^{N_M} F_{M,z}^i \tag{2}$$

where $F_{M,z}^i$ is the z component of spring forces exerted by ith motor at a steady state, and N_M is the number of motors. In this example, N_M is 2. Note that only a quarter of motor arms (i.e., N_a / 4) can bind to one F-actin due to two constraints assumed for the binding of the motor arms: motor arms can bind to F-actin when the arms are properly aligned with the polarity of F-actin, and two arms connected to the same point on a backbone cannot bind to the same F-actin. With two antiparallel F-actins, up to half of the motor arms can stay in the bound state. Thus, $F_{M,z}^i$ of a motor bound to two anti-parallel F-actins at the steady state is close to $F_{st}N_hN_a/2$, where F_{st} is the stall force of one myosin head, and N_h indicates the number of myosin heads represented by each motor arm. We found that a difference in F_{tot} originated from the relative positions of motors and ACPs. When two motors were located in a very similar position with an almost full

overlap by chance, F_{tot} was close to F_M^{max}. When the two motors stayed apart without any overlap, F_{tot} was close to either $0.5F_M^{max}$ or F_M^{max} depending on whether or not there were ACPs between them. In the absence of ACPs between two motors, forces generated by the arms of two motors could add up to develop larger tensile forces on F-actins (Figs. 2B, E). However, when ACPs existed between the two motors, it resulted in the buckling of F-actin between the ACPs and motor arms on one side, and counterbalanced the tension generated by motor arms on the other side (Figs. 2C, F). As a result, F-actins ended up with feeling tension corresponding to a force generated by one motor. These ACPs between two motors divide the bundle into serially connected contractile units as the assumption of the theoretical explanation in the other study mentioned earlier [4, 45]. These ACPs were observed to experience larger tension than the rest of the ACPs since they directly counterbalanced large tension generated by motors (Fig. 2G). There were quite a few cases showing intermediate level of bundle tension between $0.5F_M^{max}$ and F_M^{max} (Fig. 2A). These cases had motors with a partial overlap and ACPs located within the z range spanned by one of these motors. Due to the partial overlap with ACPs, forces generated by two motors partially added up, resulting in the intermediate tension (Fig. 2D).

Force generation in disorganized bundles is also regulated by the same mechanism

To verify the importance of the spatial distributions of motors and ACPs for force generation in more physiologically relevant structures, we repeated simulations using disorganized bundles with various sizes between $N_F = 2$ (8 filaments) and $N_F = 7$ (98

filaments) (Figs. 1B and 3A). The densities of motors and ACPs were fixed at $R_{\text{ACP}} = 0.04$ and $R_{\text{M}} = 0.005$, resulting in N_{M} ranging between 4 and 52. In all cases, each motor still had 24 arms ($N_{\text{a}} = 24$). When the bundle became thicker by increasing N_{F}, a tensile force acting on the bundle (F_{tot}) also increased (Fig. 3B). Because the motor density was fixed, more motors were present in thicker bundles, generating larger F_{tot}. In case of the thickest bundle ($N_{\text{F}} = 7$), actin concentration was 12.25-fold higher (= 98 / 8) compared to the thinnest bundle ($N_{\text{F}} = 2$), so there were 13-fold more motors.

We defined the efficiency of force generation:

$$\eta = \frac{F_{\text{tot}}}{F_{\text{M}}^{\text{max}}} \tag{3}$$

In these bundles, $F_{\text{M}}^{\text{max}}$ is still defined by Eq. 2, but $F_{\text{M},z}^{i}$ was close to $F_{\text{st}}N_{\text{h}}N_{\text{a}}$ because there were more than one F-actin to bind for each polarity. Interestingly, η was quite similar in cases with $N_{\text{F}} = 4$, 6, and 7 (Fig. 3C). By contrast, cases with $N_{\text{F}} = 2$ exhibited much higher η than the other cases. We probed why motors could generate forces in the thinnest bundle in a more efficient manner. With $N_{\text{F}} = 2$, there were only 4 motors. The length of each motor with 24 arms is 462 nm (= 42 nm × 11). The sum of the length of all 4 motors is only ~9% of bundle length, 20 μm, so they were less likely to significantly overlap with each other. Instead, most of them stayed apart with various distances. Given high ACP density, there were more than one ACP between adjacent motors. Thus, these motors formed separate contractile units, and their forces could not add up. Thus, η for the thinnest bundle was roughly $1/N_{\text{M}}$ because F_{tot} was close to the force generated by a single motor. Indeed, η for the thinnest bundle was close to 0.25, confirming our rationale. If motors behave in the same manner in thicker bundles ($N_{\text{F}} > 2$), η will be smaller than 0.25 because N_{M} is proportional to N_{F}^{2} due to fixed motor density. Although η was

actually smaller than 0.25 in thicker bundles, it was ~0.12 in all cases with $N_F > 2$, which was larger than the prediction, $1/N_M$. As the bundle was thicker, there was a higher chance for motors to overlap with each other or stay closely due to higher N_M. Then, some of these proximal motors could add up their forces if there is no ACP between them as discussed earlier. If the bundle has any set of such "cooperative" motors, F_{tot} can become larger than a maximal force generated by a single motor since F_{tot} is determined by the largest force generated from one of the contractile units. As N_M increases due to high N_F, the number of the cooperative motors tends to increase, resulting in larger F_{tot}. Thus, as N_F increases, both F_{tot} and F_M^{max} increase, so η can be higher than $1/N_M$ and rather insensitive to a change in N_F when N_F is not small.

Bundles with more, localized motors generate larger forces

Based on prior observations, overlaps between motors highly affect the force generation process. The extent of overlaps is determined by N_M if the bundle length is fixed. To understand how force generation in the bundles depends on N_M more systematically, we ran simulations with a wide range of N_M, using the thickest bundle ($N_F = 7$) and the same $R_{ACP} = 0.04$ and $N_a = 24$. The bundle tension, F_{tot}, showed a tendency to increase with higher N_M, but η was inversely proportional to N_M (Fig. 4A). With low N_M, F_{tot} was less sensitive to an increase in N_M, whereas η was very sensitive to the increase in N_M. If there are a small number of motors in the bundle, adding a few more motors would not lead to a significant increase in F_{tot} because the new motors are not likely to be located in positions where they can add up forces with other motors (Fig. 4B). Instead, they will merely increase the number of contractile units, most of which have

only one motor. Thus, as explained in Eq. 3, η was close to $1/N_M$. With higher N_M, F_{tot} was sensitive to a change in N_M because adding more motors to the bundle contributes to an increase in the number of motors in each contractile unit (Fig. 4B). When N_M was very high, F_{tot} was almost directly proportional to N_M with slope ~ 1 since all parts of the bundle were already occupied by motors, so adding more motors caused a direct impact on the magnitude of F_{tot}. In the range of high N_M, η was much less sensitive to a change in N_M. For example, when N_M was varied from 10 to 1,000, η was reduced to approximately half. The insentitivity of η is attributed to an increase in both F_{tot} and F_M^{max} with higher N_M as explained earlier.

To verify this rationale, we developed a way to estimate the maximum number of overlapping motors, Ξ, using our simulation data:

$$\Xi = \max_{\substack{i=1...N_M \\ j=L \text{ or } R}} \left(1 + \sum_{k \neq i}^{N_M} \xi_{ik}^{j} \right) \tag{4}$$

where ξ_{ik}^{j} is:

$$\xi_{ik}^{j} = \begin{cases} 1 & \text{if } L_c \leq L_{ov} \leq L_M \\ L_{ov} / L_c & \text{if } 0 < L_{ov} < L_c \\ 0 & \text{if } L_{ov} = 0 \end{cases} \tag{5}$$

i and k are the indices of motors, j denotes the left or right side of a motor, and L_{ov} is an overlap distance between two motors. In addition, L_c is the critical length required for the cooperative overlap:

$$L_c = 2L_{sp} \left(\frac{N_a}{4} - 1 \right) \tag{6}$$

L_c is two-fold greater than the length occupied by motor arms on one side of a motor backbone. Two motors are considered a cooperative motor pair if L_{ov} is equal to or greater

than L_c (Fig. S1). With this cooperative overlap, ACPs cannot counterbalance forces exerted by motor arms. If L_{ov} is shorter than L_c and but greater than zero, two motors are considered as a partially cooperative motor pair. By comparing all the motor combinations, Ξ can be obtained. Ξ can range between 1 (no overlap between motors) and N_M (cooperative overlap between all motors). Using this Ξ, a force acting on the bundle can be estimated as F_{est}:

$$F_{est} = \frac{F_{st} N_h N_a \Xi}{2} \tag{7}$$

The sensitivity of F_{est} to an increase of N_M is high at large N_M with a slope close to 1 (Fig. 4C), which is consistent with our rationale described earlier. F_{est} was generally smaller than F_{tot} because this analysis does not account for actual bundle geometry consisting of multiple F-actins; if two motors are located far from each other in x or y direction, they may not counterbalance or add up forces. Nevertheless, we found that F_{est} captures the overall dependence of F_{tot} on parameters well.

So far, we have assumed that motors could be randomly located at any part of the bundle. If motors are confined within a smaller region, the extent of overlaps between motors (i.e., Ξ) could be increased with the same N_M, enhancing force generation. We tested the effects of motor distribution with $N_F = 7$, $R_{ACP} = 0.04$, $N_M = 52$, and $N_a = 24$. We allowed motors to be located only within a portion of the bundle defined by f between 0 and 1. $f = 1$ means that motors can be located anywhere. The center of the region for motor allocation is equal to the center of the bundle at $z = 10$ μm. It was observed that a bundle generated the largest F_{tot}, and η was the highest when motors were located near the center ($f = 0.06$) (Fig. 5A). Note that F_M^{max} in the denominator of Eq. 3 is identical in these cases, meaning that F_{tot} is directly proportional to η. Motors localized in relatively

the same position are more likely to overlap in the cooperative manner (i.e., $L_{ov} \geq L_c$).

Therefore, a large fraction of the motors were involved with the formation of a strong contractile unit (Fig. 5B). As f increased, motors were distributed on the bundle more sparsely. Then, many motors were partially overlapped or separated from each other. As a result, F_{tot} and η decreased (Fig. 5A). The effect of a variation in f on force generation was also reproduced well by the theoretical model explained earlier (Fig. 5C).

The structure of motors influences force generation in bundles

We have employed motors with 24 arms whose length is $L_M = 462$ nm with $L_{sp} = 42$ nm and $L_{bz} = 42$ nm. As mentioned earlier, the structure of myosin thick filaments can vary significantly, depending on myosin isoform and conditions. If motors are longer and have more arms, they may generate higher bundle forces. To verify this hypothesis, we tested cases with different N_a and the thickest bundle ($N_F = 7$), with the total number of arms in the system, $N_a N_M$, fixed. With higher N_a, L_M was increased, but N_M became lower (Fig. 1D, i). When L_M was increased by higher N_a, F_{tot} increased (Fig. 6A, red). Because these cases had the same total number of motor arms, F_M^{max} in Eq. 3 was identical in all the cases. Thus, η showed the same tendency as F_{tot} (Fig. S2A); with longer motors, η was higher. The same tendency was observed in F_{est} (Fig. 6B, red). As motors have motor arms, individual contractile units become stronger, so F_{tot} and η are directly proportional to L_M if there is no overlap between motors (Fig. 6C). However, these motors are hard to overlap in a fully cooperative manner because L_c is large (Fig. 6D, top and Eqs. 5 and 6). Thus, Ξ was actually smaller with higher L_M (Fig. S2B, red). This explains why the dependence of F_{tot} on L_M was weaker at high L_M than that at low L_M (Fig. 6A).

There are two additional ways to increase L_M without a change in N_a and N_M: increasing L_{bz} at the center of motors or increasing L_{sp} between motor arms uniformly (Fig. 1E, ii and iii). We ran simulations with a variation in either L_{sp} or L_{bz}, using the thickest bundle ($N_F = 7$) with $N_a = 16$. We compared results from these new simulations with those obtained with a variation in N_a shown earlier. Interestingly, when L_M was similar, F_{tot} acquired with a change in L_{bz} was similar to that with a variation in N_a despite different N_M (Fig. 6A, blue), whereas F_{tot} in cases with a change in L_{sp} was noticeably lower (Fig. 6A, green). η showed identical tendency as F_{tot} because F_M^{max} was the same in all the cases (Fig. S2A). F_{est} also reproduced a similar tendency (Fig. 6B). If L_M is fixed, a longer bare zone makes motors have their arms near the two ends of their backbone. Then, motors can have a higher possibility to add up their forces because L_c is smaller (Fig. 6D, bottom and Eqs. 5 and 6). Thus, Ξ was higher (Fig. S2B, blue), resulting in higher F_{tot} and η despite smaller N_a. By contrast, motors with uniform large spacing between arms need to have a more overlap to add up their forces (Fig. 6D, middle). Smaller Ξ and N_a led to lower F_{tot} and η (Fig. S2B, green).

Force generation in actin networks is regulated by the same mechanism

To check the generality of our findings, we performed simulations using a two-dimensional (2D) network where F-actins and motors are randomly oriented without any bias, which is different from those in bundles (Fig. 1C). We varied either of N_M, N_a, L_{bz}, or L_{sp} as done for bundles (Fig. 7A). F_{tot} was smaller than the values measured in bundles under the same conditions due to the random orientations of motors (Figs. 7B, C).

Interestingly, we found that F_{tot} is proportional to $N_M^{0.65}$, which is close to $\sqrt{N_M}$. Note that F_{tot} was directly proportional to N_M at large N_M in case of the bundles (Fig. 3A). Considering that motors are uniformly distributed on a 2D network, this weaker dependence on N_M is expected. η showed similar mganitudes to those measured in the bundles (Figs. 6B, S2, and S3). Note that η was still calculated using Eq. 3, but F_M^{max} was obtained by summing the x or y component of forces exerted by all motors network and then averaging them:

$$F_M^{max} = \frac{1}{4}\left(\sum_i^{N_M} F_{M,x}^i + \sum_i^{N_M} F_{M,y}^i\right) \qquad (8)$$

We found that the dependences of F_{tot} and η on these parameters are similar to those observed with bundles, meaning that force generation regulated by cooperative overlaps between motors takes place even in networks. Although forces exerted by motors are not oriented in the same direction in networks, these forces can be counterbalanced or add up depending on the relative positions of motors.

DISCUSSION

Actomyosin contractility is well-conserved machinery for generating mechanical forces in animal cells, facilitating cytokinesis, cell migration, and tissue morphogenesis [18]. Myosin II, which is the most important molecular motor for cellular force generation, exists as a highly organized form called thick filaments. Myosin thick filaments have been studied extensively during recent decades. However, despite various forms of thick filaments found in different types of cells, the effects of their structural properties on force

generation have not been understood well.

There was an in vitro study that employed thin disorganized actomyosin bundles consisting of a few F-actins with three myosin isoforms (skeletal muscle, smooth muscle, and non-muscle myosins) in order to investigate the effects of the size of myosin II filaments [4]. The study showed that a tensile force developed in bundles is directly proportional to the number of myosin heads in each myosin II filament if the total number of myosin heads is identical. They explained this direct proportionality via a theoretical bundle model consisting of serially-connected contractile units in which a single myosin II filament generates a tensile force. This is partially consistent with observations in our study. However, such explanation can be applied only to a bundle without any overlap between thick filaments, meaning that there are only a few thick filaments.

Although force generation by actomyosin contractility has been investigated in several theoretical and computational studies, most of the previous models used drastically simplified motors without consideration of thick filament structures [36, 37, 46]. Therefore, it was not feasible to investigate the importance of the structural properties of thick filaments in force generation in those studies. There were a few models accounting for thick filament structures. However, only one thick filament was simulated [32, 33] due to high computational cost, or force generation process was not probed [31].

In our study, using motors with the geometry of bipolar thick filaments, we systematically showed how the force generation process in disorganized actomyosin structures, including bundles and networks, is regulated by the number, spatial distribution, and structural properties of motors. First, using the simplest system with only two F-actins, we showed that motors with ACPs between them cannot add up their forces to generate a larger force (Fig. 2). Then, we probed the force generation process in

disorganized bundles with different thickness and found that the efficiency of force generation was lower in thicker bundles although the bundle-level force was higher (Fig. 3). This was attributed to an increase in the number of motors in thicker bundles. We then found that a larger force was developed in the same bundle in a less efficient way when there were a larger number of motors (Fig. 4). When motors were sparsely distributed without an overlap between them, the efficiency of force generation was inversely proportional to the number of motors (i.e., $\eta \propto 1/N_{\mathrm{M}}$). As the number of motors further increased, motors started cooperatively overlapping with each other, forming stronger contractile units. Thus, the efficiency became greater than $1/N_{\mathrm{M}}$. We also tested the effects of motor distribution on force generation to verify the importance of cooperative overlaps (Fig. 5). As motors were distributed in a more confined region, bundles generated larger forces because there could be more cooperative overlaps between more densely distributed motors. We also found that force generation was enhanced and more efficient when motors were longer with more arms because forces generated by the arms of one motor can simply add up (Fig. 6). In addition, longer motors with a long bare zone and a few arms could generate large forces in bundles than longer motors with a small bare zone and large spacing between arms since the former can achieve the cooperative overlap by a much smaller overlap (Fig. 6). Our results imply that the consideration of F-actin connectivity may not be enough to accurately predict force generation unlike predictions from a recent study [46].

Our findings can be applied to understand the structures of stress fibers that are known to mediate physiological processes, such as cell protrusion, cytokinesis, and cell shape maintenance [47, 48]. In stress fibers, non-muscle myosin II plays a main role in generating contractile forces. There are two types of contractile stress fibers: transverse

arcs and ventral stress fibers [49]. Ventral stress fibers, which are assembled from pre-existing stress fiber precursors, have a sarcomere-like structure with cascaded (i.e., serially connected) contractile units divided by α-actinin which is one type of ACPs. By contrast, transverse arcs are smaller without distinct repeated structures with much fewer ACPs [49]. ACPs in ventral stress fibers can counterbalance forces generated by motors as shown in Fig. 2. The structure with cascaded units and many ACPs prevents the bundle-generated force from increasing beyond a force generated by a single contractile unit. Therefore, it is expected that ventral stress fibers would generate smaller forces than transverse arcs if their thickness is similar. However, the bundle-generated force would be proportional to the thickness of bundles as shown in Fig. 3. Thus, ventral stress fibers, which are typically thicker, are known to generate larger tension than transverse arcs [50].

Although we focused on force generation, the contractile behaviors of actomyosin structures (i.e., a decrease in length) have also been of great interest. Our model can be used to study such contractile behaviors by deactivating the periodic boundary condition and removing connection between one end of bundle/network and a domain boundary as done previously [20]. To achieve higher contractile speed with the same total number of myosin heads, the existence of multiple contractile units would be better as suggested in a previous work [4]. This means that there is a trade-off between force generation and contractile speed. Previous studies also showed that the contractile speed of networks is proportional to motor density [18, 43, 51]. We may be able to use our model to systematically investigate how the contractile speed is regulated by parameters that we tested in this study, including the number, distribution, length, and structure of motors.

In conclusion, we probed how the force generation process in disorganized bundles and networks are regulated by the properties of myosin thick filaments. We found

that more motors enable bundles and networks to generate larger tensile forces, but the efficiency of force generation was lower. With the same number of motors, forces generated by bundles and networks can vary to a large extent, depending on i) whether they are sparsely or densely distributed, ii) how many myosin heads each thick filament has, and iii) how long the bare zone at the center is. Our results can be partly verified using different myosin isoforms under different conditions as done in the previous study [4]. In addition, synthetic myosin thick filaments (i.e., one made with DNA origami [52]) whose structure can be designed artificially could be used to verify our findings.

ACKNOWLEDGMENTS

We gratefully acknowledge the support from EMBRIO Institute, contract #2120200, a National Science Foundation (NSF) Biology Integration Institute and the support from JST SPRING (JPMJSP2138 to SDi), JSPS (Japan Society for the Promotion of Science) KAKENHI (21H03796 to SDe).

REFERENCES

[1] Lim, C. T., Zhou, E. H., & Quek, S. T. (2006). Mechanical models for living cells—a review. *Journal of biomechanics*, 39(2), 195-216.

[2] Salbreux, G., Charras, G., & Paluch, E. (2012). Actin cortex mechanics and cellular morphogenesis. *Trends in cell biology*, 22(10), 536-545.

[3] Craig, R., & Woodhead, J. L. (2006). Structure and function of myosin filaments. *Current opinion in structural biology*, 16(2), 204-212.

[4] Thoresen, T., Lenz, M., & Gardel, M. L. (2013). Thick filament length and isoform composition determine self-organized contractile units in actomyosin bundles. *Biophysical journal*, 104(3), 655-665.

[5] Trombitás, K., & Tigyi-Sebes, A. (1984). Cross-bridge interaction, with oppositely polarized actin filaments in double-overlap zones of insect flight muscle. *Nature*, 309(5964), 168-170.

[6] Dasbiswas, K., Hu, S., Schnorrer, F., Safran, S. A., & Bershadsky, A. D. (2018). Ordering of myosin II filaments driven by mechanical forces: experiments and theory. *Philosophical transactions of the royal society B: biological sciences*, 373(1747), 20170114.

[7] Oshima, K., Sugimoto, Y., Irving, T. C., & Wakabayashi, K. (2012). Head-head interactions of resting myosin crossbridges in intact frog skeletal muscles, revealed by synchrotron x-ray fiber diffraction. *PLoS One*, 7(12), e52421.

[8] Skubiszak, L., & Kowalczyk, L. (2002). Myosin molecule packing within the vertebrate skeletal muscle thick filaments. A complete bipolar model. *Acta biochimica polonica*, 49(4), 829-840.

[9] Niederman, R., & Pollard, T. D. (1975). Human platelet myosin. II. In vitro assembly and structure of myosin filaments. *The journal of cell biology*, 67(1), 72-92.

[10] Tyska, M. J., Dupuis, D. E., Guilford, W. H., Patlak, J. B., Waller, G. S., Trybus, K. M., ... & Lowey, S. (1999). Two heads of myosin are better than one for generating force and motion. *Proceedings of the national academy of sciences*, 96(8), 4402-4407.

[11] Saito, T., Huang, W., Matsui, T. S., Kuragano, M., Takahashi, M., & Deguchi, S. (2021). What factors determine the number of nonmuscle myosin II in the sarcomeric unit of stress fibers?. *Biomechanics and modeling in mechanobiology*, 20, 155-166.

[12] Huang, W., Matsui, T. S., Saito, T., Kuragano, M., Takahashi, M., Kawahara, T., ... & Deguchi, S. (2021). Mechanosensitive myosin II but not cofilin primarily contributes to cyclic cell stretch-induced selective disassembly of actin stress fibers. *American journal of physiology-cell physiology*, 320(6), C1153-C1163.

[13] Kaminer, B., & Bell, A. L. (1966). Myosin filamentogenesis: effects of pH and ionic concentration. *Journal of molecular biology*, 20(2), 391-401.

[14] Pollard, T. D. (1982). Structure and polymerization of Acanthamoeba myosin-II filaments. *The journal of cell biology*, 95(3), 816-825.

[15] Reisler, E., Smith, C., & Seegan, G. (1980). Myosin minifilaments. *Journal of molecular biology*, 143(1), 129-145.

[16] Josephs, R., & Harrington, W. F. (1966). Studies on the formation and physical chemical properties of synthetic myosin filaments. *Biochemistry*, 5(11), 3474-3487.

[17] Kee, Y. S., & Robinson, D. N. (2008). Motor proteins: myosin mechanosensors. *Current biology*, 18(18), R860-R862.

[18] Murrell, M., Oakes, P. W., Lenz, M., & Gardel, M. L. (2015). Forcing cells into shape: the mechanics of actomyosin contractility. *Nature reviews Molecular cell biology*, 16(8), 486-498.

[19] Lenz, M., Thoresen, T., Gardel, M. L., & Dinner, A. R. (2012). Contractile units in disordered actomyosin bundles arise from F-actin buckling. *Physical review letters*, 108(23), 238107.

[20] Li, J., Biel, T., Lomada, P., Yu, Q., & Kim, T. (2017). Buckling-induced F-actin fragmentation modulates the contraction of active cytoskeletal networks. *Soft matter*, 13(17), 3213-3220.

[21] Okamoto, T., Matsui, T. S., Ohishi, T., & Deguchi, S. (2020). Helical structure of actin stress fibers and its possible contribution to inducing their direction-selective disassembly upon cell shortening. *Biomechanics and modeling in mechanobiology*, 19, 543-555.

[22] Murrell, M. P., & Gardel, M. L. (2012). F-actin buckling coordinates contractility and severing in a biomimetic actomyosin cortex. *Proceedings of the national academy of sciences*, 109(51), 20820-20825.

[23] Bendix, P. M., Koenderink, G. H., Cuvelier, D., Dogic, Z., Koeleman, B. N., Brieher, W. M., ... & Weitz, D. A. (2008). A quantitative analysis of contractility in active cytoskeletal protein networks. *Biophysical journal*, 94(8), 3126-3136.

[24] Soares e Silva, M., Depken, M., Stuhrmann, B., Korsten, M., MacKintosh, F. C., & Koenderink, G. H. (2011). Active multistage coarsening of actin networks driven by myosin motors. *Proceedings of the national academy of sciences*, 108(23), 9408-9413.

[25] Mizuno, D., Tardin, C., Schmidt, C. F., & MacKintosh, F. C. (2007). Nonequilibrium mechanics of active cytoskeletal networks. *Science*, 315(5810), 370-373.

[26] Koenderink, G. H., Dogic, Z., Nakamura, F., Bendix, P. M., MacKintosh, F. C., Hartwig, J. H., ... & Weitz, D. A. (2009). An active biopolymer network controlled by molecular motors. *Proceedings of the national academy of sciences*, 106(36), 15192-15197.

[27] Linsmeier, I., Banerjee, S., Oakes, P. W., Jung, W., Kim, T., & Murrell, M. P. (2016). Disordered actomyosin networks are sufficient to produce cooperative and telescopic contractility. *Nature communications*, 7(1), 12615.

[28] Alvarado, J., Sheinman, M., Sharma, A., MacKintosh, F. C., & Koenderink, G. H. (2013). Molecular motors robustly drive active gels to a critically connected state. *Nature physics*, 9(9), 591-597.

[29] Alvarado, J., Sheinman, M., Sharma, A., MacKintosh, F. C., & Koenderink, G. H. (2017). Force percolation of contractile active gels. *Soft matter*, 13(34), 5624-5644.

[30] Åström, J. A., Kumar, P. S., & Karttunen, M. (2009). Aster formation and rupture transition in semi-flexible fiber networks with mobile cross-linkers. *Soft matter*, 5(15), 2869-2874.

[31] Cortes, D. B., Gordon, M., Nédélec, F., & Maddox, A. S. (2020). Bond type and discretization of nonmuscle myosin II are critical for simulated contractile dynamics. *Biophysical journal*, 118(11), 2703-2717.

[32] Stam, S., Alberts, J., Gardel, M. L., & Munro, E. (2015). Isoforms confer characteristic force generation and mechanosensation by myosin II filaments. *Biophysical journal*, 108(8), 1997-2006.

[33] Weirich, K. L., Stam, S., Munro, E., & Gardel, M. L. (2021). Actin bundle architecture and mechanics regulate myosin II force generation. *Biophysical journal*, 120(10), 1957-1970.

[34] Lenz, M. (2020). Reversal of contractility as a signature of self-organization in cytoskeletal bundles. *Elife*, 9, e51751.

[35] Grewe, J., & Schwarz, U. S. (2020). Mechanosensitive self-assembly of myosin II minifilaments. *Physical review E*, 101(2), 022402.

[36] Chandrasekaran, A., Upadhyaya, A., & Papoian, G. A. (2019). Remarkable structural transformations of actin bundles are driven by their initial polarity, motor activity, crosslinking, and filament treadmilling. *PLoS computational biology*, 15(7), e1007156.

[37] Freedman, S. L., Banerjee, S., Hocky, G. M., & Dinner, A. R. (2017). A versatile framework for simulating the dynamic mechanical structure of cytoskeletal networks. *Biophysical journal*, 113(2), 448-460.

[38] Ronceray, P., Broedersz, C. P., & Lenz, M. (2019). Fiber plucking by molecular motors yields large emergent contractility in stiff biopolymer networks. *Soft matter*, 15(7), 1481-1487.

[39] Bidone, T. C., Jung, W., Maruri, D., Borau, C., Kamm, R. D., & Kim, T. (2017). Morphological transformation and force generation of active cytoskeletal networks. *PLoS computational biology*, 13(1), e1005277.

[40] Mak, M., Zaman, M. H., Kamm, R. D., & Kim, T. (2016). Interplay of active processes modulates tension and drives phase transition in self-renewing, motor-driven cytoskeletal networks. *Nature communications*, 7(1), 10323.

[41] Jung, W., Murrell, M. P., & Kim, T. (2015). F-actin cross-linking enhances the stability of force generation in disordered actomyosin networks. *Computational particle mechanics*, 2, 317-327.

[42] Kim, T. (2015). Determinants of contractile forces generated in disorganized actomyosin bundles. *Biomechanics and modeling in mechanobiology*, 14, 345-355.

[43] Yu, Q., Li, J., Murrell, M. P., & Kim, T. (2018). Balance between force generation and relaxation leads to pulsed contraction of actomyosin networks. *Biophysical journal*, 115(10), 2003-2013.

[44] Liu, S., Matsui, T. S., Kang, N., & Deguchi, S. (2022). Analysis of senescence-responsive stress fiber proteome reveals reorganization of stress fibers mediated by elongation factor eEF2 in HFF-1 cells. *Molecular biology of the cell*, 33(1), ar10.

[45] Craig, R., & Megerman, J. (1977). Assembly of smooth muscle myosin into side-polar filaments. *The journal of cell biology*, 75(3), 990-996.

[46] Eliaz, Y., Nedelec, F., Morrison, G., Levine, H., & Cheung, M. S. (2020). Insights from graph theory on the morphologies of actomyosin networks with multilinkers. *Physical review E*, 102(6), 062420.

[47] Deguchi, S., Ohashi, T., & Sato, M. (2006). Tensile properties of single stress fibers isolated from cultured vascular smooth muscle cells. *Journal of biomechanics*, 39(14), 2603-2610.

[48] Kaunas, R., & Deguchi, S. (2011). Multiple roles for myosin II in tensional homeostasis under mechanical loading. *Cellular and molecular bioengineering*, 4, 182-191.

[49] Lehtimäki, J. I., Rajakylä, E. K., Tojkander, S., & Lappalainen, P. (2021). Generation of stress fibers through myosin-driven reorganization of the actin cortex. *Elife*, 10, e60710.

[50] Lee, S., Kassianidou, E., & Kumar, S. (2018). Actomyosin stress fiber subtypes have unique viscoelastic properties and roles in tension generation. *Molecular biology of the cell*, 29(16), 1992-2004.

[51] Malik-Garbi, M., Ierushalmi, N., Jansen, S., Abu-Shah, E., Goode, B. L., Mogilner, A., & Keren, K. (2019). Scaling behaviour in steady-state contracting actomyosin networks. *Nature physics*, 15(5), 509-516.

[52] Fujita, K., Ohmachi, M., Ikezaki, K., Yanagida, T., & Iwaki, M. (2019). Direct visualization of human myosin II force generation using DNA origami-based thick filaments. *Communications biology*, 2(1), 437.

[53] Underhill, P. T., & Doyle, P. S. (2004). On the coarse-graining of polymers into bead-spring chains. *Journal of non-newtonian fluid mechanics*, 122(1-3), 3-31.

[54] Clift, R., Grace, J. R., & Weber, M. E. (2005). Bubbles, drops, and particles.

[55] Isambert, H., Venier, P., Maggs, A. C., Fattoum, A., Kassab, R., Pantaloni, D., & Carlier, M. F. (1995). Flexibility of actin filaments derived from thermal fluctuations: effect of bound nucleotide, phalloidin, and muscle regulatory proteins. *Journal of biological chemistry*, 270(19), 11437-11444.

[56] Kim, T., Hwang, W., Lee, H., & Kamm, R. D. (2009). Computational analysis of viscoelastic properties of crosslinked actin networks. *PLoS computational biology*, 5(7), e1000439.

[57] Sheetz, M. P., & Spudich, J. A. (1983). Movement of myosin-coated fluorescent beads on actin cables in vitro. *Nature*, 303(5912), 31-35.

[58] Erdmann, T., Albert, P. J., & Schwarz, U. S. (2013). Stochastic dynamics of small ensembles of non-processive molecular motors: The parallel cluster model. *The journal of chemical physics*, 139(17), 175104-175127.

[59] Erdmann, T., & Schwarz, U. S. (2012). Stochastic force generation by small ensembles of myosin II motors. *Physical review letters*, 108(18), 188101.

Fig. 1 Modeling setups. (A) In the model, F-actin (blue), actin cross-linking protein (ACP, yellow), and motor (red) are simplified by cylindrical segments. F-actin has polarity defined by barbed and pointed ends. Motors consist of a backbone with motor arms that can bind to and walk along F-actin. ACPs comprise two segments connected at the center point. κ_s and κ_b represent extensional and bending stiffnesses, respectively. (B) Two-filament system consisting of two F-actins whose barbed ends are clamped to rigid boundaries (gray). (C) The disorganized bundle system with $2N_F^2$ F-actins randomly located and oriented in the presence of the periodic boundary condition (PBC) in the z direction, where N_F is a parameter defining bundle thickness. (D) The two-dimensional network system consisting of F-actins with random positions and orientations with the PBC in x and y directions. (E) A variation in the motor structure in three different ways: (i) increasing the number of motor arms ($N_a\uparrow$), (ii) increasing the bare zone length ($L_{bz}\uparrow$), and (iii) increasing a spacing between motor arms ($L_{sp}\uparrow$).

Fig. 2 Interactions between motors and ACPs regulate force generation. (A) Time evolution of the force generated by two motors in the two-filament system. The upper and lower dashed lines indicate ideal upper and lower limits of a force that two motors can generate, respectively. (B) Without any ACP between two motors, they can generate a force close to the upper limit which is two-fold larger than a force that one motor can generate ($= F_M^{max}$). (C) With ACP(s) between two motors, ACPs counterbalance a force generated by one of the motors. Thus, they can generate a force close to the lower limit which is equal to the force that one motor can generate ($= F_M^{max}/2$). Part of F-actins is buckled due to two forces with opposite directions. (D) If two motors are close to each other, ACP can counterbalance a fraction of the force generated by one motor. Then, two motors can generate a force between the upper and lower limits. (E, F) Initial and final configurations (E) without or (F) with ACPs between two motors. The vertical dimension is increased 10 times to show the configurations clearly. (G) Measurement of tensile forces acting on F-actins (green), ACPs (blue), motor arms (red), motor backbones (black), or all (magenta) in z direction.

Fig. 3 With fixed motor density, thicker bundles generate larger force in a less efficient manner. (A) An example of disorganized bundles with $N_F = 7$ visualized at the beginning of the simulation, where N_F is a parameter defining bundle thickness. F-actins are visualized as transparent elements to show the positions of motors. (B) Bundle-level force and (C) the efficiency of force generation measured at a steady state with different N_F. In thicker bundles ($N_F > 2$), larger forces were generated, but the efficiency was lower than that in the thinnest bundle ($N_F = 2$). *** represents $p \leq 0.001$, and n.s. represents $p > 0.05$.

Fig. 4 An increase in the number of motors (N_M) in disorganized bundles results in larger forces but smaller force generation efficiency. The thickest bundle ($N_F = 7$) was used for all cases. (A) Bundle-level force (blue circles) and the efficiency of force generation (red triangles) with a wide range of N_M. With more motors, a larger force (F_{tot}) was generated, but the efficiency (η) was lower. (B) Configuration of motors with different N_M. Ξ represents the estimated number of motors in the strongest contractile unit. With small N_M, F_{tot} and Ξ are unlikely to increase significantly until an entire bundle is occupied by motors, so η is roughly $1/N_M$. By contrast, with high N_M, an increase N_M directly enhances F_{tot} and Ξ, and η almost remains constant. (C) Prediction of a force using the positions of motors. To find the estimated force (F_{est}), Ξ is calculated first, and Eq. 7 is used.

Fig. 5 Motor distribution affects the force generation in disorganized bundles. We varied the relative size of a region where motors were initially located, f. $f = 1$ means that motors can be located at any part of the bundle. (A) Bundle-level force (F_{tot}) and efficiency (η) depending on f. As motors were localized more closely to the center (i.e., smaller f), the force and the efficiency were higher. (B) Configuration of motors with different f. Ξ indicates the estimated number of motors in the strongest contractile unit. Given the number of motors, smaller f results in more cooperative overlaps (i.e., higher Ξ) and thus leads to higher F_{tot} and η. (C) Prediction of the bundle-level force with different f.

Fig. 6 The architecture of motors impacts the force generation process in disorganized bundles. (A) Bundle-level force (F_{tot}) depending on L_M. The motor length (L_M) is varied by changing either of the number of motor arms (N_a, red circles), the bare zone length (L_{bz}, blue triangles), or a spacing between motor arms (L_{sp}, green squares). (B) Prediction of the bundle-level force using the positions of motors. (C) As each motor has more arms ($N_a\uparrow$), F_{tot} and the efficiency of force generation (η) become higher because forces generated by motor arms are counterbalanced to a lesser extent. (D) Possible overlaps between two motors with different structures. A dark gray color indicates a fully cooperative overlap, and light gray indicates a partially cooperative overlap. To have the fully cooperative overlap, motors with many arms need to located very closely, whereas motors with the long bare zone can overlap in the fully cooperative manner with a relatively long distance between them.

Fig. 7 Force generation in two-dimensional actomyosin networks is governed by similar mechanisms. (A) Examples of networks at the initial state under the reference condition, with a smaller or larger number of motors, and with longer motors. These snapshots show only a quarter of networks to better visualize individual motors. (B) Network-level tension (F_{tot}) and the efficiency of force generation (η) with a different number of motors. (C) F_{tot} depending on motor length varied by changing either of the number of motor arms (N_a, red circles), the bare zone length (L_{bz}, blue triangles), or a spacing between motor arms (L_{sp}, green squares).