Targeting SLC7A11-mediated cysteine metabolism for the treatment of trastuzumab

resistant HER2 positive breast cancer

Yijia Hua[1,2,#], Ningjun Duan[1,#,*], Chunxiao Sun[1], Fan Yang[1], Min Tian[1], Yanting Sun[1],

Shuhan Zhao[3], Jue Gong[1], Qian Liu[1], Xiang Huang[1], Yan Liang[1], Ziyi Fu[1], Wei Li[1],

Yongmei Yin[1,2,4,*]

1. Department of Oncology, The First Affiliated Hospital of Nanjing Medical

University, Nanjing, China

2. Gusu School, Suzhou Municipal Hospital, The Affiliated Suzhou Hospital of

Nanjing Medical University, Suzhou, China

3. Department of General Surgery, The First Affiliated Hospital of Nanjing Medical

University, Nanjing, China

4. Jiangsu Key Lab of Cancer Biomarkers, Prevention and Treatment, Collaborative

Innovation Center for Personalized Cancer Medicine, Nanjing Medical University,

Nanjing, China

# These authors contributed equally to this work.

***Correspondence to**: Yongmei Yin, Email: ymyin@njmu.edu.cn

Ningjun Duan, Email: ningjun_official@outlook.com

Keywords:

HER2 positive breast cancer, trastuzumab primary resistance, cysteine metabolism,

epigenetic modifications.

Abbreviations:

4-HNE, 4-hydroxynonenal; 5-mC, 5-methylcytosine; ADCC, antibody dependent

cellular cytotoxicity; DAMPs, damage-associated molecular patterns; DMR,

differentially methylated regions; GABA, 4-aminobutyric acid; GSH, glutathione;

GSR, glutathione reductase; GSS, glutathione synthetase; GSSG, glutathione disulfide;

HER2, human epidermal growth factor receptor 2; MDA, Malondialdehyde; pCR,

pathological complete response; PUFAs, polyunsaturated fatty acids; PUMA, p53

upregulated modulator of apoptosis; sgRNA, small guide RNA; siRNA, short

interfering RNA; SNP, single nucleotide polymorphism; SNV, single nucleotide

variation; TCA, tricarboxylic acid; TME, tumor microenvironment; TNBC, triple

negative breast cancer; WGBS, whole-genome bisulfite sequencing.

Author Contributions

Y.M.Y. and N.J.D. conceived the project. Y.J.H., N.J.D., C.X.S., F.Y., Y.T.S., S.H.Z.,

Y.L. and J.G. performed the experiments. M.T. and Q.L. helped collect clinical

samples. Y.J.H. and N.J.D. conducted data analysis and wrote the manuscript. N.J.D.,

X.H., Z.Y.F., W.L. and Y.M.Y. supervised the study. All authors reviewed the

manuscript.

Acknowledgements

We would like to thank the Core Facility of the First Affiliated Hospital of Nanjing

Medical University for its help in the detection of experimental samples.

Funding

This project was supported by National Natural Science Foundation of China

(81972484 and 82272667 to Yongmei Yin; 82203488 to Ningjun Duan), High-level

Innovation Team of Nanjing Medical University Program (JX102GSP201727), The

Collaborative Innovation Center for Tumor Individualization Program

(JX21817902/008) and Beijing Xisike Clinical Oncology Research Foundation

(Y-2019AZZD-00680).

Conflict of interest statement

The authors declare that there is no conflict of interest.

Data Availability

Further information and requests for resources could be directed to corresponding

authors.

Abstract

Trastuzumab resistance remains a challenge for HER2 positive breast cancer treatment. Targeting metabolic reprogramming would provide novel insights for therapeutic strategies. Here, we integrated metabolomics, transcriptomics and epigenomics data of trastuzumab sensitive and primary resistant HER2 positive breast cancer to identify metabolic alterations. Aberrant cysteine metabolism was discovered in trastuzumab primary resistant breast cancer at both circulating and intracellular levels. The inhibition of SLC7A11 and cysteine starvation could synergize with trastuzumab to induce ferroptosis. Mechanistically, increased H3K4me3 and decreased DNA methylation enhanced SLC7A11 transcription and cystine uptake in trastuzumab resistant breast cancer. The regulation of epigenetic modifications modulated cysteine metabolism and ferroptosis sensitivity. These results revealed an innovative approach for overcoming trastuzumab resistance by targeting specific amino acid metabolism.

Introduction

Accounting for 15%-20% of all breast cancer cases, human epidermal growth factor receptor 2 (HER2) positive breast cancer has posed great threats to the health of women worldwide(1, 2). The overexpression of HER2 results in aberrant breast cancer cell proliferation, invasion, and progression(3). As a monoclonal antibody targeting HER2, trastuzumab has been recognized as the foundational medication in HER2 positive breast cancer treatment and provided remarkable clinical benefits for numerous patients(4). Around 50% HER2 positive breast cancer patients would experience trastuzumab resistance and disease progression during or after treatment settings(5). Current theories maintained that trastuzumab resistance mainly resulted from HER2 mutation(6), antigenic epitope variation(7), compensatory pathway activation(8) and tumor heterogeneity(9). However, there still lacks a comprehensive theory to thoroughly explain the mechanisms behind trastuzumab resistance, and exploring effective targets to overcome resistance remains a major topic in the field of

breast cancer research(3).

Metabolic reprogramming has been acknowledged as one crucial cancer hallmark(10).

Traced back to the first study carried out by Otto Warburg, decades of studies have

indicated that glycolysis, tricarboxylic acid (TCA) cycle, lipid metabolism, amino

acid metabolism, as well as other processes collectively modulate the development

and function of cancer cells and tumor microenvironment (TME) components(11-13).

Several studies have revealed that different breast cancer subtypes featured diverse

metabolic patterns(14, 15). Targeting specific enzymes or modulating nutrients uptake

might normalize the TME and enhance breast cancer treatment efficacy(16, 17). Yet

previous studies often focused on the overall metabolic subtypes of breast cancer,

with little attention given to abnormal metabolic changes and underlying genomic

alterations induced by specific drug resistance. This might lead to a poor

understanding on the tumor heterogeneity in drug resistant breast cancer cells.

Epigenetic modifications play crucial roles in all stages of cancer development and

contribute to cancer cell survival selection(18). Alterations in chromatin structure,

DNA methylation and histone modifications provide tumor growth advantage and

increase phenotypic variability(19). During the formation of drug resistance,

epigenetic dysregulation modulates key genes transcription associated with treatment

tolerance, leading to resistant clonal derivatives and disease progression(20). Previous

studies suggested that COMPASS complex inactivation suppressed tumor suppressor

genes transcription and accelerated breast cancer tumorigenesis(21). Aberrant DNA

methylation resulted in abnormal oncogenic signaling activation and drug resistance

acquisition(22). Targeting epigenetic reprogramming might provide novel treatment

options for overcoming breast cancer drug resistance.

By integrating metabolomic, transcriptomic and epigenomic data, we identified

metabolic alternations, especially cysteine-associated pathways, accompanied with

the epigenetic changes during the development of primary trastuzumab resistance of

HER2 positive breast cancer (**Figure 1A**). Based on these discoveries, targeting

certain metabolism-related protein, epigenetic modification or nutrition could all be

possible ways to overcome trastuzumab resistance. In all, our study provided novel

insights and potential solutions on primary trastuzumab resistance of HER2 positive

breast cancer.

Methods

Patient cohort and blood samples collection

One breast cancer patient cohort from Jiangsu Province Hospital was embedded in

this study, which consists of 26 primary trastuzumab resistant patients defined as

disease recurrence during or after ($\leq$ 12 months) trastuzumab adjuvant treatment and

26 trastuzumab sensitive patients that benefited from trastuzumab adjuvant treatment

more than 12 months(23).

All blood samples were collected before the first trastuzumab treatment cycle.

Samples collection was under the approval of the Ethics Committee and Institutional

Review Board of Jiangsu Province Hospital. Each patient provided written informed

consent for sample and data use.

Cell culture

Human breast cancer cell lines JIMT1, SKBR3 and human breast epithelial cell line

MCF10A were obtained from American Type Culture Collection (ATCC). JIMT1 and

SKBR3 cell lines were cultured in high-glucose Dulbecco's modified eagle medium

(DMEM; Gibco, 11965092) with 10% fetal bovine serum (FBS; Gibco, 16140071)

and 1% penicillin streptomycin (Gibco, 15140122) at 37□ with 5% CO_2. MCF10A

was cultured in mammary epithelial cell growth medium (MEGM) supplemented with

bovine pituitary extract, human epidermal growth factor, hydrocortisone, gentamicin

sulfate-amphotericin, and insulin (Lonza, CC-3150) with 100 ng/ml cholera toxin

(Sigma, C8052) at 37□ with 5% CO_2. Cell culture with cysteine starvation was

performed by cystine/cysteine-deficient DMEM (BIOTREE) and 1% penicillin

streptomycin at 37℃ with 5% CO_2.

DNA constructs, plasmids, and transfection

Short interfering RNA (siRNA) targeting ASH2L, SLC7A11 and GPX4 were

presented in Supplementary Table S1. The design of small guide RNA (sgRNA) was

based on the relevant methylated region and conducted via the online tool CRISPick(24, 25). Sequences of sgRNA were indicated in Supplementary Table S2. dCas9-DNMT3A was obtained from Addgene (#100090)(26).

Transfection was conducted by using Lipofectamine 3000 (Invitrogen, L3000015) according to the manufacturer's protocol. In brief, cells were cultured in a six-well plate at 70%-90% confluency. For RNA interference, 2500 ng siRNA and 3.75 μl Lipofectamine 3000 reagents were mixed and then added in each well. For CRISPR-based epigenetic editing, dCas9-DNMT3A plasmid and sgRNA were first mixed in 1:2 ratio, then mixed with 3.75 μl Lipofectamine 3000 reagent and 5 μl P3000 reagent in each well.

Protein extraction and Western blots

Cells were collected and then lysed in SDS lysis buffer with phenylmethyl-sulfonyl fluoride (PMSF; Servicebio, G2008) and phosphatase Inhibitor cocktail (Servicebio, G2007) on ice. Proteins were seprated in SDS-PAGE and transferred to PVDF membranes (Millipore, IPVH00010). After blocked, membranes were incubated with specific primary and secondary antibodies, and then visualized with imaging system and ECL chemiluminescence kit (Vazyme, E422-01). Primary antibodies included SLC7A11 (CST, 12691S), GPX4 (CST, 52455S), glutathione synthetase (Abcam, ab124811), glutathione reductase (Abcam, ab124995), ASH2L (CST, 5019T), H3K4me3 (Abcam, ab213224), GAPDH (proteintech, HRP-60004), and H3 (Abcam, ab1791). The secondary antibody was horseradish peroxidase (HRP) conjugated goat anti-rabbit IgG(H+L) (ImmunoWay, RS0002).

GSH/GSSG ratio assay

GSH/GSSG ratio was measured with a GSH and GSSG assay kit (Beyotime, S0053). Cells were lysed with protein removing reagent solution and several freeze-thaw cycles. After centrifuged, supernatants were collected and incubated with DTNB and NADPH.

The absorbance at 412 nm was detected to measure the GSH/GSSG ratio with standard curve.

C11 BODIPY (581/591) assay

180 Cells were collected and incubated with 5 μM C11 BODIPY (581/591) (invitrogen,

181 D3861) for 30 minutes at 37 □. The oxidated and reduced state were detected by flow

182 cytometry, with excitation and emission wavelength at 488/510 and 581/591 nm,

183 respectively.

DCFH-DA assay

185 Cells were collected and incubated with 10 μM DCFH-DA (MCE, HY-D0940) for 30

186 minutes at 37 □. The oxidated state was detected with laser excitation wavelength at

187 488nm.

Cystine uptake assay

189 The cystine uptake ability was measured by using cystine uptake assay kit (DOJINDO,

190 UP05). Cells were incubated with cystine analog solution and fluorescent probe. The

191 fluorescence intensity was measured with excitation and emission wavelength at 490

192 and 535 nm.

Quantification of intracellular cysteine

194 The concentration of intracellular cysteine was detected by using a cysteine

195 quantification kit (Solarbio, BC0185). Cells were lysed and incubated with solution

196 containing phosphotungstic acid. The absorbance at 600nm was detected to measure

197 intracellular cysteine levels.

ChIP

Chromatin extraction and the following immune precipitation were performed with

chromatin extraction kit (Abcam, ab117152) and ChIP kit (Abcam, ab117138)

according to manufacturer's protocols. Cells were harvested and treated with 1%

formaldehyde first, then sonication was applied on cross-linked chromatin to generate

200-1000 bp fragments. Chromatin fractions enriched with certain histone

modifications were captured by antibodies (H3K27me3 (CST, C36B11), H3K4me3

(Abcam, ab213224) and IgG) and released by protease digestion.

RT-qPCR was applied to quantify the enrichment of specific type of histone

modification at certain chromatin regions with ChamQ SYBR qPCR Master Mix

(Vazyme, Q311). While VAHTS® Universal DNA Library Prep Kit for Illumina

(Vazyme, ND607) and Illumina Novaseq 6000 platform were applied for sequencing

library preparation and the following sequencing. All primers used for ChIP-qPCR

were presented in Supplementary Table S3.

MeDIP

Methylated DNA immune precipitation was performed with MeDIP kit (Abcam,

ab117135) according to manufacturer's protocols followed by chromatin extraction as

ChIP protocol. After cross-linking and sonication, methylated chromatin fractions

were enriched by anti-5-Methylcytosine (5-mC) antibody (CST, 28692). RT-qPCR

was applied to quantify the enrichment of 5-mC at certain chromatin regions. All

primers used for MeDIP-qPCR were presented in Supplementary Table S3.

Dot Blots

Total genomic DNA was extracted by using nucleic DNA isolation kit (Vazyme,

DC112). Denaturized DNA was spotted on a nitrocellulose membrane, and blocked

with 5% skim milk in TBST (20mM Tris, 150mM NaCl, and 0.1% Tween-20).

Anti-5-mC primary antibody and HRP-conjugated secondary antibody together with

ECL chemiluminescence kit were applied for total methylated cytosine visualization.

Mice xenograft model construction

Six-week-old female BALB/c nude mice were obtained from Beijing Vital River

Laboratory Animal Technology, housed in a 12-hour light-dark lighting condition with

20-22 ℃ temperature and 60 ± 10% humidity. The animal diet and drinking water

followed the national standard for experimental animal feed. The diet underwent

Cobalt-60 irradiation sterilization and the drinking water followed purification and

sterilization.

All animal experiments were conducted according to the review and approval of

Institutional Animal Care and Use Committee in Nanjing Medical University

(IACUC-2204057). 3×10^6 JIMT1 cells were injected subcutaneously into the

mammary fat pad region to generate xenograft models.

The mice were divided into 6 groups: (1) DMSO treatment; (2) trastuzumab (Roche)

treatment (10mg/kg, injected intraperitoneally twice a week); (3) erastin (HY-15763)

treatment (40mg/kg, injected intraperitoneally every other day); (4) cysteine

starvation; (5) trastuzumab treatment plus cysteine starvation; (6) trastuzumab plus

erastin treatment. Cysteine starvation was performed by utilizing cystine/cysteine

deficient diet (Xietong Bio). Treatment was initiated when tumor volume reached

50mm^3. Tumor sizes and mice weight were measured three times a week. Tumor

volumes were calculated as $0.5 \times L \times W^2$ (L: the longest dimension, W: the

perpendicular dimension). The tumor growth was compared by paired individual fold

change of tumor volumes.

Tumor and spleen cell composition analysis

Tumor samples were digested in serum-free RPMI with 10mg/ml collagenase I

(Sigma, C0130) and 1mg/mg DNase I (Roche, 10104159001) for 30-60 minutes at

37 □. Spleen samples were gently crushed. Both of them were filtered in 70μm

strainers, followed by red blood cell removal with RBC lysis buffer (eBioscience,

00-4333-57), and then resuspended in flow cytometry staining buffer. Fc receptor was

blocked with mouse FcR blocking reagent (Miltenyi, 130-092-575).

Fluorescent antibodies including CD45 (clone 30-F11, eBioscience, 48-0451-82),

CD3 (clone 145-2C11, BioLegend, 100353), CD49b (clone DX5, BioLegend,

108907), CD11b (clone M1/70, eBioscience, 56-0112-82), CD27 (clone LG.3A10,

BioLegend, 124216), CD335 (NKp46) (clone 29A1.4, eBioscience, 17-3351-82) and

granzyme B (clone NGZB, eBioscience, 11-8898-82) together with fixable viability

dye (eBioscience, 65-0865-14) were chosen for cell staining. CytoFLEX (Beckamn

Coulter) and FlowJo software were applied for fluorescence measurement and

analysis.

Immunohistochemical staining

After formalin fixation and paraffin embedding, tissue sections were cut and mounted

on slides followed by deparaffinizing, rehydrating and antigen retrieval. Samples were

then stained with antibodies including SLC7A11 (Abcam, ab307601), GPX4 (Abcam,

ab125066), 4-Hydroxynonenal (4-HNE) (Abcam, ab48506), Malondialdehyde (MDA)

(Abcam, ab243066), and caspase-3 (Abcam, ab32150).

Differential transcriptome analyses

The human hg38 genome was utilized for RNA sequencing reads alignment.

Differential expression analyses were measured by the DESeq2 package in R.

Significant differential expression was defined with a threshold of P adj ≤ 0.05 and

|log2 fold change| ≥ 0.5.

Differential metabolome analyses

273   Metabolomics data was analyzed by the MetaboAnalystR package in R(27). Pathway

274   enrichment analyses was performed according to KEGG databases. Significant

275   difference was defined with a threshold of P adj ≤ 0.05 and |log2 fold change| ≥ 0.5.

Differential epigenome analyses

277   Sequencing reads of ChIP and whole-genome bisulfite sequencing (WGBS) were

278   aligned to the human hg38 genome. Differential methylation analyses was performed

279   by deepTools (version 3.5.5)(28). Differentially methylated regions (DMR) were

280   identified by Bismark (version 0.24.2)(29). Significant difference was defined with a

281   threshold of P adj ≤ 0.05 and |log2 fold change| ≥ 0.5.

Genomics analyses

283   The human hg38 genome was utilized to align WGS sequencing reads. Germline

284   single nucleotide polymorphisms (SNP) and Indels were identified by GATK (v

285   4.5.0.0). The annotation of germline mutations was obtained by utilizing ANNOVAR.

Statistical analyses

All statistical analyses were performed by utilizing GraphPad Prism (version 10.1.2)

and R (version 4.3.3). All data were represented as the mean $\pm$ standard error (SE) or

the mean $\pm$ SEM from a minimum of three independent experiments. Two-tailed

Student's t-test was applied to determine statistical in comparisons between two

groups and ANOVA test in multiple groups. The threshold for statistical significance

is $P < 0.05$, and was marked as * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Results

Different amino acids metabolic patterns between trastuzumab sensitive and

primary resistant HER2 positive breast cancer patients

To investigate metabolic patterns of HER2 positive breast cancer patients with

different trastuzumab response, we obtained and analyzed the plasma metabolites data

of both trastuzumab sensitive and primary resistant patients (26/26) with good

interclass correlations (**Figure S1A-C**). Of them, 203 metabolites (130 upregulated

and 73 downregulated in primary resistant groups) were identified with different

abundances among two patient groups (**Figure 1B and S1D**). KEGG-based pathway

analyses suggested the major enrichment in amino acids metabolism. For instance,

glycine, serine and threonine metabolism, valine, leucine and isoleucine biosynthesis,

as well as cysteine and methionine metabolism were reduced in primary resistant

patients; while arginine, proline, alanine, aspartate and glutamate metabolism were

enhanced (**Figure 1C and 1D**).

By comparing the abundance of 20 common protein-construction amino acids, we

realized that cystine and aspartate were higher while valine, lysine and leucine were

lower in primary resistant groups. However, glutamate kept stable in both groups

(**Figure 1E and 1F**). Non-protein amino acids also displayed difference among these

two groups. 2-aminoisobutyric acid and 4-aminobutyric acid (GABA) were lower in

resistant groups, while citrulline, kynurenine, ornithine and taurine levels were similar

(**Figure S1E**).

We then analyzed transcriptomic data of pre-treatment breast cancer tumors in I-SPY2

trial(30) to further explore metabolic gene alterations, in which HER2 positive breast

cancer patients received trastuzumab plus paclitaxel, trastuzumab combined with

pertuzumab and paclitaxel or T-DM1 plus paclitaxel during neoadjuvant settings. By

comparing patients with different responses to neoadjuvant trastuzumab-based

regimens (pathological complete response [pCR] and non-pCR), we found 1542 genes

upregulated and 1350 genes downregulated in non-pCR groups (**Figure S1F and 1G**).

Based on KEGG datasets, amino acids metabolism was found dysregulated in patients

with poor responses to trastuzumab, including aberrant lysine degradation, cysteine

and methionine metabolism, beta-alanine metabolism and histidine metabolism

(**Figure 1H**).

As cystine was one of the most significant amino acids in both metabolomic and

transcriptomic analyses, we investigated relations between cystine/cysteine metabolic

key genes and trastuzumab treatment outcomes. Patients with non-pCR featured

higher expression of SLC7A11, while no differences were found in GPX4, GSS and

GSR expression (**Figure 1I**), indicating SLC7A11 might become a potential target for

trastuzumab primary resistant patients.

In general, trastuzumab primary resistant HER2 positive breast cancer patients

featured distinct amino acids metabolic patterns, especially increased plasma cystine

levels and higher expression of SLC7A11.

Distinct cysteine metabolism in trastuzumab primary resistant HER2 positive

breast cancer

To comprehensively compare the metabolic heterogeneity between trastuzumab

sensitive and primary resistant HER2 positive breast cancer, we obtained and

analyzed both metabolomic and transcriptomic data of trastuzumab sensitive SKBR3

and trastuzumab resistant JIMT1 cell lines.

Significant differences were observed in metabolic data with good quality (**Figure**

S2A and S2B). Specifically, 181 upregulated and 235 downregulated hydrophilic

metabolites accompany 546 upregulated and 412 downregulated lipophilic

metabolites were found in JIMT1 cells compared to SKBR3 cells (**Figure 2A and**

S2C). For hydrophilic metabolites, enrichment analyses revealed that glycine, serine

and threonine metabolism were upregulated in JIMT1, while cysteine and methionine

metabolism were downregulated (**Figure 2B and 2C**). Enrichment analyses based on

lipophilic metabolites showed that linoleic acid metabolism, fatty transportation and

glycerophospholipid catabolism were upregulated in JIMT1, but glycerolipid

metabolism, phospholipid biosynthesis and retinoid metabolism downregulated

(**Figure S3A and S3B**).

Differential gene expression could also be uncovered from transcriptomic data. 4235

downregulated genes and 4923 upregulated genes were identified in the comparison

between JIMT1 and SKBR3 cells (**Figure S4A**). Based on KEGG datasets, activities

of cysteine and glutathione metabolism pathways were found dysregulated in JIMT1

cells (**Figure 2D**). More importantly, the joint pathway analyses combining

metabolomic and transcriptomic data indicated that JIMT1 featured aberrant

glutathione metabolism (**Figure 2E**).

The extracellular cystine is mainly imported by SLC7All for subsequent protein

synthesis, glutathione (GSH) production and oxidative catabolism (**Figure 2F**). Both

transcriptomic and protein evidence revealed the upregulation of SLC7A11 and

glutathione synthetase (GSS) as well as the downregulation of GPX4 and glutathione

reductase (GSR) in JIMT1 cells (**Figure S4B and 2G**), together with the

characteristics that JIMT1 cells showed higher cysteine uptake ability and

cytoplasmic cysteine level but lower glutathione/glutathione disulfide (GSH/GSSG)

ratio than SKBR3 cells (**Figure 2H-J**), indicating the probably increased

transsulfuration activity and decreased γ-glutamyl-peptides synthesis activity in

JIMT1 cells (**Figure S4C**). These results suggested that HER2 positive breast cancer

with primary trastuzumab resistance might prefer to consume more cystine and

feature intracellular cysteine metabolic reprogramming, showing high consistency

with metabolic characteristics of relevant breast cancer patients and may associate

with trastuzumab treatment response.

Stimulate ferroptosis sensitivity of trastuzumab primary resistant HER2 positive

breast cancer

Since cysteine metabolism plays key role in maintaining cellular oxidative-reductive

balance, disruption of which may turn breast cancer cells into a more sensitive state to

oxidative damage, including ferroptosis(31, 32). We first downregulated the

expression of SLC7A11 and GPX4 in JIMT1 by siRNAs, and the lipid peroxidation

of ferroptosis was found increased by measuring BODIPY-C11 (**Figure 3A and 3B**).

Similar results were also detected in SKBR3(**Figure S5A and S5B**). We then detected

the sensitivity of JIMT1 and SKBR3 to SLC7A11 inhibitor (erastin) and GPX4

inhibitor (RSL3) and found JIMT1 was more sensitive to these two inhibitors (**Figure**

S5C and S5D). The deprivation of cysteine could suppress the growth of both JIMT1

and SKBR3, but JIMT1 showed more tolerance to this nutrition starvation (**Figure**

S5E).

12 hours treatment of SLC7A11 inhibitor (erastin), GPX4 inhibitor (RSL3) and even

cysteine starvation were all able to induce ferroptosis in both JIMT1 and SKBR3 cells

(**Figure 3C and 3D**). However, JIMT1 cells indicated more rapidly accumulation of

lipid peroxidation under the treatment of erastin, which could be detected by both

increased BODIPY-C11 and DCFH-DA (**Figure 3E and 3F**). Compared with SKBR3,

JIMT1 cells were also more sensitive to RSL3 but more durable to cysteine starvation

(**Figure 3G and 3H**), which might be explained by its stronger transsulfuration

activities and decreased γ-glutamyl-peptides synthesis (**Figure S5F**). The utilization

of ferroptosis inhibitor Fer-1 could suppress the lipid peroxidation resulted from

erastin, RSL3 and cysteine starvation in both JIMT1 and SKBR3, which also

indicated the role of cysteine metabolism in modulating HER2 positive breast cancer

ferroptosis (**Figure S5G and S5H**).

To understand whether trastuzumab would modulate the ferroptosis sensitivity, we

treated JIMT1 cells with trastuzumab for different durations. We found that with a

prolonged exposure of trastuzumab for 72 hours, the total amount of SLC7A11 in

JIMT1 was decreased (**Figure 3I**). Correspondingly, cystine uptake ability was

suppressed after trastuzumab treatment for 48 and 72 hours (**Figure 3J**). In addition,

cells underwent trastuzumab treatment combining with erastin or cysteine starvation

could demonstrate stronger lipid peroxidation and enhanced ferroptosis (**Figure 3K**).

Restraining cysteine metabolism inhibited tumor growth in vivo

To further investigate the effects of SLC7A11 inhibition or cysteine limitation on

trastuzumab primary resistant HER2 positive breast cancer, JIMT1 xenograft mouse

models were generated and then treated with different strategies including DMSO,

trastuzumab, erastin, cysteine starvation, trastuzumab plus erastin, and trastuzumab

plus cysteine starvation (**Figure 4A**).

Of them, trastuzumab plus erastin and trastuzumab plus cysteine starvation could

effectively inhibit tumor growth, while cysteine starvation and erastin monotherapy

only showed limited effects on tumor progression, demonstrating the better treatment

responses of combination strategy (**Figure 4B, 4C and S6A**). In addition, almost no

weight differences were observed among all groups during treatment, partially

indicating the safety of such cysteine limitation treatment (**Figure 4D**).

The immune cell compositions in tumor and spleen of each group were measured to

analyze the immune response induced by different treatment strategies (**Figure S6B**

and S6C). The infiltration of NK cells (CD3-CD45+CD49b+) in tumors was

increased only upon the combination of cysteine deprivation and trastuzumab (**Figure**

4E). Nearly no differences were found in tumor infiltrating NK cell activation

(NKp46), toxic function (Granzyme B) and maturation status (CD11b, CD27) among

all these groups (**Figure 4F-H**). While similar results were also observed in spleen

NK cells (**Figure S6D-G**).

These results suggested a better effect of cysteine starvation than erastin in NK cell

recruitment and activation, resulting a more powerful synergy with trastuzumab to

increase the abundance of tumor infiltrating NK cells and show better inhibition of

trastuzumab primary resistant breast cancer. As the anti-tumor effects of trastuzumab

mainly relies on antibody dependent cellular cytotoxicity (ADCC) effects mediated by

NK cells(33), this combination therapy might provide more effective approaches for

HER2 positive breast cancer treatment.

We then analyzed the biomarkers of cysteine metabolism and cell death pathways in

tumor samples of different treatment groups (**Figure 4I**). Erastin and cysteine

deprivation monotherapy, or combined with trastuzumab, could upregulate SLC7A11

and GPX4 expression, as well as the abundance of MDA and 4-HNE. Ferroptosis

might increase endoplasmic reticulum stress and activate p53 upregulated modulator

of apoptosis (PUMA)(34). The combination of trastuzumab and erastin or

trastuzumab and cysteine starvation might also modify apoptosis sensitivity.

Apoptosis biomarker caspase-3 was found increasing in these synergic treatment

groups (**Figure S6H**). These results suggested that targeting cysteine metabolism

could induce ferroptosis in trastuzumab primary resistant HER2 positive breast cancer

and might have a synergic effect with trastuzumab to inhibit breast cancer

progression.

Aberrant promoter H3K4me3 regulates SLC7A11 gene expression and cysteine

metabolism

Abnormal changes of histone modification play critical roles in tumor progression

including drug resistance(35). Of them, altered methylations on promoter histone

H3K4 and H3K27 show most important effects on gene expression(36).

By comparing the ChIP-seq data, we detected 3934 and 3925 H3K4me3 peaks were

enriched in JIMT1 and SKBR3 respectively. These altered peaks primarily located

among the promoter regions of genes (**Figure 5A and S7A**), which have close

relationships with biological processes such as nucleic acid transcription, protein

modification, mRNA processing and ribosome processing (**Figure S8A**)

We also confirmed that 7048 and 7001 H3K27me3 peaks were enriched in JIMT1 and

SKBR3 respectively, mainly enriching at not only promoter regions but also distal

intergenic regions, and exons (**Figure 5B and S7B**) of genes that involve in cell-cell

adhesions, axon guidance, postsynaptic transmission, and neuron differentiation

(**Figure S8B**).

Particularly, we uncovered the enrichment of H3K4me3 at the promoter regions of

key cysteine transporter gene SLC7A11 in JIMT1 cells compared with SKBR3 cells

while H3K27me3 signals were stable (**Figure 5C**), showing the consistency with its

high expression in JIMT1 cells.

By inhibiting the expression of ASH2L, one crucial subunit of COMPASS

complex(37), we downregulated not only the overall H3K4me3 level but also the

signal at SLC7A11 promoter in JIMT1 cells, and successfully decreased the

expression of SLC7A11 (**Figure 5D-F**). Meanwhile, diminished downstream cysteine

uptake leaded to increased lipid peroxidation, the key symbol of ferroptosis (**Figure**

5G and 5H).

These results suggested that H3K4me3 may play a more important role in modulating

SLC7A11 transcription and cysteine metabolism in JIMT1 than H3K27me3.

Targeting H3K4me3 could regulate SLC7A11 expression, enhancing lipid

peroxidation and inducing ferroptosis.

DNA methylation modifies SLC7A11 transcription and cysteine metabolism

Besides histone modification, DNA methylation is another crucial epigenetic

modification to regulate genomic imprinting, modify transposable elements and

modulate transcription, even in tumor progression(38).

In general, JIMT1 cells featured a lower abundance of total 5-mC than SKBR3 cells

(**Figure S9A**). By investigating DNA methylation landscape with WGBS, we

identified 329419 CG-DMRs enriched in JIMT1 and 84634 CG-DMRs enriched in

SKBR3 (**Figure 6A and S9B**). Among non-CG DMRs (CHG, CHH; in which H

denotes A, C or T), 1604 CHG-DMRs and 3727 CHH-DMRs were upregulated in

JIMT1, while 2187 CHG-DMRs and 5158 CHH-DMRs were upregulated in SKBR3

(**Figure S9B**).

Across whole genome, CG-DMRs mainly enriched in intron, intergenic regions, and

exons (**Figure S9C**). Differentially methylated genes were related to several

biological processes, such as neurotransmitter biosynthesis, postsynaptic transmission,

cytoplasmic translation, and ribosomal activities (**Figure S10A and S10B**).

SLC7A11 is located at chromosome 4q28.3 and we found that, in comparison with

SKBR3, the CpG methylation at SLC7A11 promoter region was decreased in JIMT1

(**Figure 6B**). To further verify the influence of altered promoter GC-DMR on

SLC7A11 gene transcription, CRISPR/Cas9-based genomic editing tool

dCas9-DNMT3A together with SLC7A11 specific promoter-targeting sgRNA sets

were utilized to successfully increase the 5-mC level at SLC7A11 promoter regions

(**Figure 6C**) in JIMT1 cells, which is confirmed by MeDIP-qPCR (**Figure 6D**). As a

result, the expression of SLC7A11 and cysteine uptake were reduced (**Figure 6E and**

6F), and the following cellular lipid peroxidation and ferroptosis were stimulated in

JIMT1 cells (**Figure 6G**).

These results revealed that increased DNA methylation at SLC7A11 promoter regions

in JIMT1 might contribute to SLC7A11 overexpression. Precise modulation of 5-mC

could suppress cysteine uptake and result in ferroptosis, which might provide novel

methods to overcome trastuzumab resistance.

Discussion

Metabolic heterogeneity has been widely accepted as important characteristics in many cancer types, which are closely associated with cancer progression, distant metastasis, and even drug resistance(39, 40), while targeting dysregulated metabolism could also be novel therapeutic strategies(41).

By comparing the metabolomic data of cell lines and blood samples from patients, we realized that widespread metabolic reprogramming particular the alternation of cysteine metabolism was an important feature of primary trastuzumab resistance, extending our understanding of anti-tumor drug resistance (**Figure 7**).

Cysteine is a crucial non-essential amino acid and mainly obtained from dietary, protein catabolism and methionine transsulfuration(42). Due to oxidative extracellular environment, cystine, instead of cysteine, is normally intake by the major transporter SLC7A11 (also known as xCT)(43), which also has important roles in protein synthesis, posttranslational modifications, and redox balance maintenance(42, 44).

Meanwhile, biosynthesis of GSH by GSS is one important outlet of cellular cysteine, which helps maintaining oxidation-reduction balance and protecting membrane from oxidative damage including ferroptosis with GPX4 and GSR(45, 46). The downregulation of both GPX4 and GSR and the upregulation of GSS, together with the higher intake of cysteine mediated by raised SLC7A11, could stimulate GSH consumption and GSSG accumulation in JIMT1 cells, resulting in a weaker reductive cytoplasmic environment and higher sensitivity to oxidative damage, including ferroptosis.

As an important rate-limiting metabolite of the GSH biosynthesis, cysteine modulates ferroptosis through the cyst(e)ine/GSH/GPX4 axis(47). SLC7A11 inhibition, cysteine deprivation or targeting GPX4 have been reported to effectively enhance lipid peroxidation and induce ferroptosis in several cancer types(48-51). In our study, JIMT1 was much more sensitive to SLC7A11 and GPX4 inhibition, but tolerable to cysteine starvation, and this might result from increased methionine transsulfuration activity and γ-glutamyl-peptides synthesis, which compensated the deficiency of extracellular cysteine(52, 53). Whether similar effects could be observed on GSS or

GSR also need to be studied.

The inhibition of HER2 could result in impaired redox homeostasis and increased

ROS(54). Tumor cells might adapt to this metabolic stress by activating antioxidant

signaling pathways and acquire recurrence and dormancy(54, 55). Although JIMT1 is

resistant to trastuzumab, the extended exposure of trastuzumab was found to decrease

SLC7A11 expression. SLC7A11 overexpression and altered cysteine and glutathione

metabolism might function as compensating ways for trastuzumab adaptation. The

dual inhibition of cysteine metabolism and HER2 also enhanced ferroptosis both in

vivo and in vitro. In a similar manner, the inhibition of glutamine metabolism and

antioxidant transcription factor NRF2 could also suppress the proliferation of residual

breast cancer after HER2 inhibition and restore intracellular GSH metabolism(54).

These results indicated that re-establishing redox homeostasis and modulating

antioxidant metabolism might provide novel treatment options for trastuzumab

primary resistant HER2 positive breast cancer.

Besides amino acids metabolism, lipid metabolism, especially polyunsaturated fatty

acids (PUFAs) and relevant phospholipid metabolism, which are also key drivers of

ferroptosis(56), were identified dysregulated in both primary and acquired

trastuzumab resistance of HER2 positive breast cancer(57), demonstrating the

possibility that targeting PUFA metabolism to induce ferroptosis might provide novel

treatment options for overcoming trastuzumab resistance.

Not limited to cancer cells, altered cellular metabolism and oxidation-reduction

balance can also affect activities of immune cells within TME(58-60). For example,

lipid peroxidation stress might impair NK functions and immune surveillance.

Metabolites, such as L-kynurenine, might enhance ferroptosis in NK cells through

aryl hydrocarbon receptor and mediate immunosuppressive TME in gastric cancer(61).

Activation of NRF2 related antioxidant pathways could restore NK cell functions and

trigger enhanced anti-tumor responses(62). And The deprivation of cysteine was

reported to promote the metabolic reprogramming of TME and influence T cell

immune responses(63). We observed that the combination of cysteine starvation and

trastuzumab can increase tumor infiltrating NK cells but show limited effects on NK

cell development and toxic functions, relevant mechanisms still need to be discussed.

In addition, metabolites, cytokines, damage-associated molecular patterns (DAMPs)

and other immunomodulatory signals released from cancer cells that undergo

ferroptosis could function on immune cells nearby, stimulating or suppressing

anti-tumor immune responses(64, 65).

The emergence of drug resistance in cancers mainly depends on two scenarios:

Darwinian selection and Lamarkian induction(66, 67). During the evolution of breast

cancer, different subtypes of breast cancer might originate from different development

status of mammary stem cells, bipotent progenitors and differentiated cells(68, 69).

HER2 positive breast cancer was reported to transform from luminal progenitor and

mature luminal cells(70-72). The tumorigenesis of trastuzumab primary resistant

HER2 positive breast cancer might favor the Darwinian selection, as a small

population of pre-existing trastuzumab tolerant cancer cells might be enriched upon

treatment.

To investigate genetic and epigenetic issues stimulating the evolution of primary

trastuzumab resistant and sensitive breast cancer, we compared genomic variation,

histone modification and DNA methylation status among JIMT1, SKBR3 and

MCF10A (normal mammary epithelial cell).

During carcinogenesis, missense mutations, splicing mutations and in frame deletions

were the most frequent events and MUC3A, MUC6, OR8U1 and ZNF717 were most

frequent mutated genes (**Figure S11A and S11B**). Meanwhile, single nucleotide

polymorphism (SNP) was the major variation type and the substitution between

cytosine and thymine was the most common signature (**Figure S12A-B and**

S13A-C).

Compared with MCF10A, JIMT1 instead of SKBR3 featured distinct single

nucleotide variation (SNV) at the exon region of SLC7A11 (gain at Chr4:138183261

and loss at Chr4: 138219340) (**Figure S14A-C**).

SNVs located in exons have been reported to modulate gene transcription and

function. For example, SNVs in BRCA1 exons encoding RING and BRCT domains

might disrupt BRCA1 expression and function, therefore increasing breast cancer

risks(73). While noncoding SNVs, such as rs9383590 located at ESR1 enhancer

region, could interact with regulatory elements and activate oncogene

transcription(74). The accumulation of SNVs was also associated with driver gene

mutations and might promote breast cancer evolution in a cumulative manner(75).

Whether different SNVs in SLC7A11 exon regions could contribute to SLC7A11

expression and breast cancer initiation still needs to be studied.

Epigenetic issues also demonstrated their impact on carcinogenesis. H3K4me3 instead

of H3K27me3 peaks were shown more enrichment at SLC7A11 promoter regions in

MCF10A cells (**Figure S15A and S15B**), while 5-mC levels at CpG islands among

SLC7A11 promoter regions were also increased in MCF10A (**Figure S15C**),

revealing the important proposal of histone modifications and DNA methylation on

metabolic reprogramming during the evolution of HER2 positive breast cancer.

Although several studies have reported that SLC7A11 expression was modulated by

NRF2, ATF4 and other transcription factors(43, 76, 77), it remains unclear whether its

expression is subject to a broader genome-wide or histone-level regulation(78). The

theory that epigenetic modifications contribute to specific transcription regulation,

cancer cell development and metabolic reprogramming, to modulate treatment

response and drug resistance, has been widely accepted and proved(79). Our study

tried to explore the relationship between the altered cysteine metabolism signature

and epigenetic modifications in trastuzumab resistant HER2 positive breast cancer.

Elucidating epigenetic mechanisms would help better understand trastuzumab

resistance formation and provide novel therapeutic opportunities.

Several studies have described the regulatory roles of histone modifications on

oncogene transcription(79, 80). MLL mutations were identified among various

cancers and widespread shortening H3K4me3 peaks at tumor suppressor genes might

contribute to tumor initiation(81, 82) and H3K27me3 could modulate chemotherapy

tolerance in triple negative breast cancer (TNBC)(83). Moreover, the combination of

chemotherapy and H3K27me3 demethylation inhibitor could prevent the proliferation

of persister cells(83). Targeting KDM6A could inhibit JIMT1 tumor growth via

decreasing tRNA transcription(84). As we confirmed, reducing H3K4me3 can

decrease ASH2L, SLC7A11 expression and cystine uptake, leading to enhanced lipid

peroxidation and ferroptosis, modulating histone modification can be a promising

strategy in cancer treatment.

Compared with normal tissues, almost all cancers feature abnormal DNA

hypomethylation and local hypermethylation at specific CpG islands(81, 85).

Dysregulated DNA methylation contributes to breast cancer tumorigenesis and drug

resistance formation(86, 87). For instance, trastuzumab resistant HER2 positive breast

cancer was found associated with downregulated TGFBI, CXCL2 and SLC38A1,

which was silenced by DNA hypermethylation(88). Aberrant DNMT1, DNMT3A and

DNMT3B expression were correlated with endocrine therapy resistance and predicted

worsen pathologic status and prognosis(89). By utilizing CRISPR interference, we

verified the effect of 5-mC at SLC7A11 promoter on gene expression and downstream

cysteine metabolism. What's more, since increased H3K4me3 might interact with

DNMT3A-DNMT3L(90), the crosstalk between histone modification and DNA

methylation may have collaborative effects on cysteine metabolism in trastuzumab

primary resistant breast cancer.

In conclusion, our study revealed that trastuzumab primary resistant HER2 positive

breast cancer featured distinct cysteine metabolism, which was driven by altered

H3K4me3 and DNA methylation. Targeting cysteine metabolism could provide novel

treatment concepts for overcoming trastuzumab resistance.

Supplemental information

Figures S1-S15 and Tables S1-S3

References

1.   Giaquinto AN, Sung H, Miller KD, et al. Breast Cancer Statistics, 2022. CA Cancer J Clin.

2022;72(6):524-41.

2.   Harbeck N, Penault-Llorca F, Cortes J, et al. Breast cancer. Nat Rev Dis Primers.

2019;5(1):66.

3.   Swain SM, Shastry M, Hamilton E. Targeting HER2-positive breast cancer: advances and

future directions. Nat Rev Drug Discov. 2023;22(2):101-26.

4.   Marra A, Chandarlapaty S, Modi S. Management of patients with advanced-stage

HER2-positive breast cancer: current evidence and future perspectives. Nat Rev Clin Oncol.

2024;21(3):185-202.

5.   Cossetti RJ, Tyldesley SK, Speers CH, et al. Comparison of breast cancer recurrence

and outcome patterns between patients treated from 1986 to 1992 and from 2004 to 2008. J

Clin Oncol. 2015;33(1):65-73.

6.   Xu X, De Angelis C, Burke KA, et al. HER2 Reactivation through Acquisition of the HER2

L755S Mutation as a Mechanism of Acquired Resistance to HER2-targeted Therapy in

HER2(+) Breast Cancer. Clin Cancer Res. 2017;23(17):5123-34.

7.   Arribas J, Baselga J, Pedersen K, et al. p95HER2 and breast cancer. Cancer Res.

2011;71(5):1515-9.

8.   Baselga J, Cortes J, Im SA, et al. Biomarker analyses in CLEOPATRA: a phase III,

placebo-controlled study of pertuzumab in human epidermal growth factor receptor 2-positive,

first-line metastatic breast cancer. J Clin Oncol. 2014;32(33):3753-61.

9.   Marchio C, Annaratone L, Marques A, et al. Evolving concepts in HER2 evaluation in

breast cancer: Heterogeneity, HER2-low carcinomas and beyond. Semin Cancer Biol.

2021;72:123-35.

10. Hanahan D. Hallmarks of Cancer: New Dimensions. Cancer Discov. 2022;12(1):31-46.

11. Martinez-Reyes I, Chandel NS. Cancer metabolism: looking forward. Nat Rev Cancer.

2021;21(10):669-80.

12. Vander Heiden MG, DeBerardinis RJ. Understanding the Intersections between

Metabolism and Cancer Biology. Cell. 2017;168(4):657-69.

13. Kao KC, Vilbois S, Tsai CH, et al. Metabolic communication in the tumour-immune

microenvironment. Nat Cell Biol. 2022;24(11):1574-83.

14. Gong Y, Ji P, Yang YS, et al. Metabolic-Pathway-Based Subtyping of Triple-Negative

Breast Cancer Reveals Potential Therapeutic Targets. Cell Metab. 2021;33(1):51-64 e9.

15. Xiao Y, Ma D, Yang YS, et al. Comprehensive metabolomics expands precision medicine

for triple-negative breast cancer. Cell Res. 2022;32(5):477-90.

16. Miller KD, O'Connor S, Pniewski KA, et al. Acetate acts as a metabolic immunomodulator

by bolstering T-cell effector function and potentiating antitumor immunity in breast cancer. Nat

Cancer. 2023;4(10):1491-507.

17. Xiao Y, Yu TJ, Xu Y, et al. Emerging therapies in cancer metabolism. Cell Metab.

2023;35(8):1283-303.

18. Timp W, Feinberg AP. Cancer as a dysregulated epigenome allowing cellular growth

advantage at the expense of the host. Nat Rev Cancer. 2013;13(7):497-510.

19. Feinberg AP, Koldobskiy MA, Gondor A. Epigenetic modulators, modifiers and mediators

in cancer aetiology and progression. Nat Rev Genet. 2016;17(5):284-99.

20. Brown R, Curry E, Magnani L, et al. Poised epigenetic states and acquired drug

resistance in cancer. Nat Rev Cancer. 2014;14(11):747-53.

21. Langille E, Al-Zahrani KN, Ma Z, et al. Loss of Epigenetic Regulation Disrupts Lineage

Integrity, Induces Aberrant Alveogenesis, and Promotes Breast Cancer. Cancer Discov.

2022;12(12):2930-53.

22. Garcia-Martinez L, Zhang Y, Nakata Y, et al. Epigenetic mechanisms in breast cancer

therapy and resistance. Nat Commun. 2021;12(1):1786.

23. Wong H, Leung R, Kwong A, et al. Integrating molecular mechanisms and clinical

evidence in the management of trastuzumab resistant or refractory HER-2(+) metastatic

breast cancer. Oncologist. 2011;16(11):1535-46.

24. Doench JG, Fusi N, Sullender M, et al. Optimized sgRNA design to maximize activity and

minimize off-target effects of CRISPR-Cas9. Nat Biotechnol. 2016;34(2):184-91.

25. Sanson KR, Hanna RE, Hegde M, et al. Optimized libraries for CRISPR-Cas9 genetic

screens with multiple modalities. Nat Commun. 2018;9(1):5416.

26. O'Geen H, Ren C, Nicolet CM, et al. dCas9-based epigenome editing suggests

acquisition of histone methylation is not sufficient for target gene repression. Nucleic Acids

Res. 2017;45(17):9901-16.

27. Ewald JD, Zhou G, Lu Y, et al. Web-based multi-omics integration using the Analyst

software suite. Nat Protoc. 2024.

28. Ramirez F, Ryan DP, Gruning B, et al. deepTools2: a next generation web server for

deep-sequencing data analysis. Nucleic Acids Res. 2016;44(W1):W160-5.

29. Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for

Bisulfite-Seq applications. Bioinformatics. 2011;27(11):1571-2.

30. Clark AS, Yau C, Wolf DM, et al. Neoadjuvant T-DM1/pertuzumab and

paclitaxel/trastuzumab/pertuzumab for HER2(+) breast cancer in the adaptively randomized

I-SPY2 trial. Nat Commun. 2021;12(1):6428.

31. Dixon SJ, Olzmann JA. The cell biology of ferroptosis. Nat Rev Mol Cell Biol. 2024.

32. Ye L, Wen X, Qin J, et al. Metabolism-regulated ferroptosis in cancer progression and

therapy. Cell Death Dis. 2024;15(3):196.

33. Musolino A, Gradishar WJ, Rugo HS, et al. Role of Fcgamma receptors in HER2-targeted

breast cancer therapy. J Immunother Cancer. 2022;10(1).

34. Lee YS, Kalimuthu K, Park YS, et al. BAX-dependent mitochondrial pathway mediates the

crosstalk between ferroptosis and apoptosis. Apoptosis. 2020;25(9-10):625-31.

35. Jin ML, Jeong KW. Histone modifications in drug-resistant cancers: From a cancer stem

cell and immune evasion perspective. Exp Mol Med. 2023;55(7):1333-47.

36. Millan-Zambrano G, Burton A, Bannister AJ, et al. Histone post-translational modifications

- cause and consequence of genome function. Nat Rev Genet. 2022;23(9):563-80.

37. Piunti A, Shilatifard A. Epigenetic balance of gene expression by Polycomb and

COMPASS families. Science. 2016;352(6290):aad9780.

38. Mattei AL, Bailly N, Meissner A. DNA methylation: a historical perspective. Trends Genet.

2022;38(7):676-707.

39. Faubert B, Solmonson A, DeBerardinis RJ. Metabolic reprogramming and cancer

progression. Science. 2020;368(6487).

40. Kim J, DeBerardinis RJ. Mechanisms and Implications of Metabolic Heterogeneity in

Cancer. Cell Metab. 2019;30(3):434-46.

41. Stine ZE, Schug ZT, Salvino JM, et al. Targeting cancer metabolism in the era of

precision oncology. Nat Rev Drug Discov. 2022;21(2):141-62.

42. Bonifacio VDB, Pereira SA, Serpa J, et al. Cysteine metabolic circuitries: druggable

targets in cancer. Br J Cancer. 2021;124(5):862-79.

43. Koppula P, Zhuang L, Gan B. Cystine transporter SLC7A11/xCT in cancer: ferroptosis,

nutrient dependency, and cancer therapy. Protein Cell. 2021;12(8):599-620.

44. Lennicke C, Cocheme HM. Redox metabolism: ROS as specific molecular regulators of

cell signaling and function. Mol Cell. 2021;81(18):3691-707.

45. Sies H, Mailloux RJ, Jakob U. Fundamentals of redox regulation in biology. Nat Rev Mol

Cell Biol. 2024.

46. Wu K, El Zowalaty AE, Sayin VI, et al. The pleiotropic functions of reactive oxygen

species in cancer. Nat Cancer. 2024;5(3):384-99.

47. Stockwell BR. Ferroptosis turns 10: Emerging mechanisms, physiological functions, and

therapeutic applications. Cell. 2022;185(14):2401-21.

48. Dixon SJ, Lemberg KM, Lamprecht MR, et al. Ferroptosis: an iron-dependent form of

nonapoptotic cell death. Cell. 2012;149(5):1060-72.

49. Badgley MA, Kremer DM, Maurer HC, et al. Cysteine depletion induces pancreatic tumor

ferroptosis in mice. Science. 2020;368(6486):85-9.

50. Shi Z, Naowarojna N, Pan Z, et al. Multifaceted mechanisms mediating cystine

starvation-induced ferroptosis. Nat Commun. 2021;12(1):4792.

51. Yang F, Xiao Y, Ding JH, et al. Ferroptosis heterogeneity in triple-negative breast cancer

reveals an innovative immunotherapy combination strategy. Cell Metab. 2023;35(1):84-100

e8.

52. Zhang HF, Klein Geltink RI, Parker SJ, et al. Transsulfuration, minor player or crucial for

cysteine homeostasis in cancer. Trends Cell Biol. 2022;32(9):800-14.

53. Kang YP, Mockabee-Macias A, Jiang C, et al. Non-canonical Glutamate-Cysteine Ligase

Activity Protects against Ferroptosis. Cell Metab. 2021;33(1):174-89 e7.

54. Fox DB, Garcia NMG, McKinney BJ, et al. NRF2 activation promotes the recurrence of

dormant tumour cells through regulation of redox and nucleotide metabolism. Nat Metab.

2020;2(4):318-34.

55. Parida PK, Marquez-Palencia M, Nair V, et al. Metabolic diversity within breast cancer

brain-tropic cells determines metastatic fitness. Cell Metab. 2022;34(1):90-105 e7.

56. Pope LE, Dixon SJ. Regulation of ferroptosis by lipid metabolism. Trends Cell Biol.

2023;33(12):1077-87.

57. Duan N, Hua Y, Yan X, et al. Unveiling Alterations of Epigenetic Modifications and

Chromatin Architecture Leading to Lipid Metabolic Reprogramming during the Evolutionary

Trastuzumab Adaptation of HER2-Positive Breast Cancer. Adv Sci (Weinh). 2024:e2309424.

58. De Martino M, Rathmell JC, Galluzzi L, et al. Cancer cell metabolism and antitumour

immunity. Nat Rev Immunol. 2024.

59. Leone RD, Powell JD. Metabolism of immune cells in cancer. Nat Rev Cancer.

2020;20(9):516-31.

60. Chen X, Kang R, Kroemer G, et al. Broadening horizons: the role of ferroptosis in cancer.

Nat Rev Clin Oncol. 2021;18(5):280-96.

61. Cui JX, Xu XH, He T, et al. L-kynurenine induces NK cell loss in gastric cancer

microenvironment via promoting ferroptosis. J Exp Clin Cancer Res. 2023;42(1):52.

62. Poznanski SM, Singh K, Ritchie TM, et al. Metabolic flexibility determines human NK cell

functional fate in the tumor microenvironment. Cell Metab. 2021;33(6):1205-20 e5.

63. Han C, Ge M, Xing P, et al. Cystine deprivation triggers CD36-mediated ferroptosis and

dysfunction of tumor infiltrating CD8(+) T cells. Cell Death Dis. 2024;15(2):145.

64. Dang Q, Sun Z, Wang Y, et al. Ferroptosis: a double-edged sword mediating immune

tolerance of cancer. Cell Death Dis. 2022;13(11):925.

65. Zhou Q, Meng Y, Li D, et al. Ferroptosis in cancer: From molecular mechanisms to

therapeutic strategies. Signal Transduct Target Ther. 2024;9(1):55.

66. Boumahdi S, de Sauvage FJ. The great escape: tumour cell plasticity in resistance to

targeted therapy. Nat Rev Drug Discov. 2020;19(1):39-56.

67. Gatenby RA, Brown JS. Integrating evolutionary dynamics into cancer therapy. Nat Rev

Clin Oncol. 2020;17(11):675-86.

68. Polyak K. Breast cancer: origins and evolution. J Clin Invest. 2007;117(11):3155-63.

69. Chaffer CL, Weinberg RA. Cancer cell of origin: spotlight on luminal progenitors. Cell

Stem Cell. 2010;7(3):271-2.

70. Lim E, Vaillant F, Wu D, et al. Aberrant luminal progenitors as the candidate target

population for basal tumor development in BRCA1 mutation carriers. Nat Med.

2009;15(8):907-13.

71. Prat A, Perou CM. Mammary development meets cancer genomics. Nat Med.

2009;15(8):842-4.

72. Tharmapalan P, Mahendralingam M, Berman HK, et al. Mammary stem cells and

progenitors: targeting the roots of breast cancer for prevention. EMBO J.

2019;38(14):e100852.

73. Findlay GM, Daza RM, Martin B, et al. Accurate classification of BRCA1 variants with

saturation genome editing. Nature. 2018;562(7726):217-22.

74. Bailey SD, Desai K, Kron KJ, et al. Noncoding somatic and inherited single-nucleotide

variants converge to promote ESR1 expression in breast cancer. Nat Genet.

2016;48(10):1260-6.

75. Nishimura T, Kakiuchi N, Yoshida K, et al. Evolutionary histories of breast cancer and

related clones. Nature. 2023;620(7974):607-14.

76. Anandhan A, Dodson M, Schmidlin CJ, et al. Breakdown of an Ironclad Defense System:

The Critical Role of NRF2 in Mediating Ferroptosis. Cell Chem Biol. 2020;27(4):436-47.

77. Bai X, Ni J, Beretov J, et al. Activation of the eIF2alpha/ATF4 axis drives triple-negative

breast cancer radioresistance by promoting glutathione biosynthesis. Redox Biol.

2021;43:101993.

78. Wang Y, Hu J, Wu S, et al. Targeting epigenetic and posttranslational modifications

regulating ferroptosis for the treatment of diseases. Signal Transduct Target Ther.

2023;8(1):449.

79. Wang N, Ma T, Yu B. Targeting epigenetic regulators to overcome drug resistance in

cancers. Signal Transduct Target Ther. 2023;8(1):69.

80. Sun L, Zhang H, Gao P. Metabolic reprogramming and epigenetic modifications on the

path to cancer. Protein Cell. 2022;13(12):877-919.

81. Zhao S, Allis CD, Wang GG. The language of chromatin modification in human cancers.

Nat Rev Cancer. 2021;21(7):413-30.

82. Chen K, Chen Z, Wu D, et al. Broad H3K4me3 is associated with increased transcription

elongation and enhancer activity at tumor-suppressor genes. Nat Genet.

2015;47(10):1149-57.

83. Marsolier J, Prompsy P, Durand A, et al. H3K27me3 conditions chemotolerance in

triple-negative breast cancer. Nat Genet. 2022;54(4):459-68.

84. Duan N, Hua Y, Yan X, et al. An Imbalance in Histone Modifiers Induces tRNA-Cys-GCA

Overexpression and tRF-27 Accumulation by Attenuating Promoter H3K27me3 in Primary

Trastuzumab-Resistant Breast Cancer. Cancers (Basel). 2024;16(6).

85. Nishiyama A, Nakanishi M. Navigating the DNA methylation landscape of cancer. Trends

Genet. 2021;37(11):1012-27.

86. Sher G, Salman NA, Khan AQ, et al. Epigenetic and breast cancer therapy: Promising

diagnostic and therapeutic applications. Semin Cancer Biol. 2022;83:152-65.

87. Sukocheva OA, Lukina E, Friedemann M, et al. The crucial role of epigenetic regulation in

breast cancer anti-estrogen resistance: Current findings and future perspectives. Semin

Cancer Biol. 2022;82:35-59.

88. Palomeras S, Diaz-Lagares A, Vinas G, et al. Epigenetic silencing of TGFBI confers

resistance to trastuzumab in human breast cancer. Breast Cancer Res. 2019;21(1):79.

89. Jahangiri R, Jamialahmadi K, Gharib M, et al. Expression and clinicopathological

significance of DNA methyltransferase 1, 3A and 3B in tamoxifen-treated breast cancer

patients. Gene. 2019;685:24-31.

90. Du J, Johnson LM, Jacobsen SE, et al. DNA methylation pathways and their crosstalk

with histone methylation. Nat Rev Mol Cell Biol. 2015;16(9):519-32.

Figure 1. Analyses of amino acids metabolic patterns in trastuzumab primary resistant and sensitive HER2 positive breast cancer patients. (A) Workflow of analyses performed in this study. (B) Volcano plot of different circulating metabolites in the comparison between trastuzumab primary resistant and sensitive patients. (C) Enrichment analysis of circulating metabolites downregulated in trastuzumab primary resistant patients. (D) Enrichment analysis of circulating metabolites upregulated in trastuzumab primary resistant patients. (E-F) Volcano plot (E) and violin plots (F) of different circulating protein-construction amino acids in the comparison between trastuzumab primary resistant and sensitive patients. (G) Volcano plot of different genes between non-pCR and pCR patients in trastuzumab-based neoadjuvant treatment (I-SPY2, GSE181574). (H) Amino acids metabolic pathway analysis of non-pCR and pCR patients. (I) Relations between cysteine metabolic genes and trastuzumab treatment outcomes. Significances were determined by two-tailed unpaired t-test (F, I). ns, P ≥0.05; *, P < 0.05; **, P < 0.01; ***, P < 0.001.

Figure 2. Heterogeneity of cysteine metabolism characteristics in HER2 positive breast cancer with different trastuzumab responses. (A) Volcano plot of hydrophilic metabolites in JIMT1 and SKBR3. (B) Upregulated metabolic processes in JIMT1 based on enrichment analysis of metabolites. (C) Downregulated metabolic processes in JIMT1 based on enrichment analysis of metabolites. (D) Different metabolic activities between JIMT1 and SKBR3 based on transcriptomics data. (E) Joint pathway analysis based on metabolomics and transcriptomics data. (F) Map of essential genes involved with cystine/cysteine metabolism. Upregulated genes in JIMT1 are highlighted in red and downregulated genes in JIMT1 are highlighted in blue. (G) Expression of essential proteins participating in cysteine and glutathione metabolism. (H) The comparison of cystine uptake ability between JIMT1 and SKBR3. (I) The comparison of intracellular cysteine abundance between JIMT1 and SKBR3. (J) The comparison of GSH:GSSG ratio between JIMT1 and SKBR3. Significances were determined by two-tailed unpaired t-test (H-J). ns, P ≥0.05; *, P < 0.05; **, P < 0.01; ***, P < 0.001.

Figure 3. Ferroptosis sensitivity of HER2 positive breast cancer with different trastuzumab response. (A-B) Knockdown of SLC7A11 (A) and GPX4 (B) induced increased lipid peroxidation (BODIPY-C11) in JIMT1. (C-D) Treatment with erastin (10µm), RSL3 (1µm) and cysteine starvation for 12 hours induced lipid peroxidation (BODIPY-C11) in JIMT1 (C) and SKBR3 (D). (E-F) JIMT1 and SKBR3 featured different lipid peroxidation sensitivity to erastin by measuring BODIPY-C11 (E) and DCFH-DA (F). (G-H) JIMT1 and SKBR3 featured different lipid peroxidation sensitivity to RSL3 (G) and cysteine starvation (H). (I) SLC7A11 expression under different trastuzumab treatment time. (J) Cystine uptake ability under different trastuzumab treatment time. (K) The combination of erastin or cysteine starvation and trastuzumab increased lipid peroxidation in JIMT1. H, trastuzumab; Cys, cysteine starvation. Significances were determined by one-way ANOVA (A-D, J, K) and two-tailed unpaired t-test (E, G, H). ns, $P \geq 0.05$; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Figure 4. Targeting cysteine metabolism synergizes with trastuzumab to induce ferroptosis. (A) Schematic outline of treatment on cysteine metabolism in trastuzumab resistant breast cancer. JIMT1 cells were injected into the mammary fat pads of BALB/c nude mice at day -7. Treatment with DMSO, erastin, cysteine starvation, trastuzumab combined with or without erastin and cysteine starvation started at day 0 for 21 days. Tumor volume and mice weight were measured. The tumor growth was compared by paired individual fold change of tumor volumes. (B) Tumor of different treatment groups. (C) Fold change of tumor volume in different treatment groups. (D) Mice weight of different treatment groups. (E-H) Abundance of CD49b+ (E), NKp46+ (F), Granzyme B+ (G) and different development status (H) tumor infiltrating NK cells. (I) Representing immunohistochemistry images of SLC7A11, GPX4, MDA and 4-HNE in different treatment groups. Scale bar, 50 μm. H, trastuzumab; Cys, cysteine starvation. Significances were determined by two-way ANOVA (C, D) and one-way ANOVA (E-H). ns, $P \geq 0.05$; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Figure 5. Increased H3K4me3 modifies SLC7A11 transcription and regulated cystine

uptake. (A-B) Alterations of H3K4me3 (A) and H3K27me3 (B) peaks between JIMT1

and SKBR3. (C) Different abundance of H3K4me3 and H3K27me3 at SLC7A11

promoter regions in JIMT1 and SKBR3. (D) Decreased ASH2L inhibited total

H3K4me3 expression. (E) Decreased ASH2L inhibited SLC7A11 expression. (F)

Decreased ASH2L suppressed H3K4me3 expression at SLC7A11 promoter regions.

(G) Decreased ASH2L suppressed cystine uptake ability. (H) Decreased ASH2L

induced lipid peroxidation and ferroptosis. Significances were determined by

two-tailed unpaired t-test (F-H). ns, P ≥0.05; *, P < 0.05; **, P < 0.01; ***, P <

0.001.

Figure 6. Decreased 5-mC modifications on SLC7A11 promoter relate with enhanced cystine uptake. (A) Alterations of 5-mC peaks between JIMT1 and SKBR3. (B) Different abundance of 5-mC at SLC7A11 promoter regions in JIMT1 and SKBR3. (C) Schematic outline of CRISPR-based epigenetic editing. dCas9-DNMT3A increased methylation of specific CpG islands in SLC7A11 promoter regions and inhibited gene transcription. (D) dCas9-DNMT3A increased 5-mC levels at SLC7A11 promoter regions. (E) dCas9-DNMT3A inhibited SLC7A11 expression. (F) dCas9-DNMT3A suppressed cystine uptake ability. (G) dCas9-DNMT3A induced lipid peroxidation and ferroptosis. Significances were determined by one-way ANOVA (D, F, G). ns, $P \geq 0.05$; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

953 **Figure 7.** Summary of targeting cysteine metabolism in trastuzumab resistant HER2

954 positive breast cancer.

Trastuzumab sensitive

Trastuzumab resistant

Cysteine starvation

Ferroptosis

Erastin

NK Cell

Epigenetic modification