Sparse matrix factorization of GWAS summary statistics robust to sample sharing improves detection and interpretation of factors with diverse genetic architectures

Ashton R. Omdahl[1], Joshua S. Weinstock[2], Rebecca Keener[1], Surya B. Chhetri[1], Marios Arvanitis[1,3], Alexis Battle[1,4-7]
1) Department of Biomedical Engineering, Johns Hopkins University; Baltimore, MD, 21218.
2) Department of Human Genetics, Emory University; Atlanta, GA, 30322.
3) Division of Cardiology, Department of Medicine, Johns Hopkins University, Baltimore, MD
4) Department of Computer Science, Johns Hopkins University; Baltimore, MD, 21218
5) Department of Genetic Medicine, Johns Hopkins University; Baltimore, MD, 21218.
6) Malone Center for Engineering in Healthcare, Johns Hopkins University, Baltimore, MD, 21218.
7) Data Science and AI Institute, Johns Hopkins University, Baltimore, MD, 21218.

Abstract

Complex trait-associated genetic variation is highly pleiotropic. This extensive pleiotropy implies that multi-phenotype analyses are informative for characterizing genetic variation, as they facilitate the discovery of trait-specific and trait-shared variants and pathways ("genetic factors"). Previous efforts have estimated genetic factors using matrix factorization (MF) applied to numerous GWAS. However, existing methods are susceptible to spurious factors arising from residual confounding due to sample-sharing in biobank GWAS. Furthermore, MF approaches have historically estimated dense factors, loaded on most traits and variants, that are challenging to map onto interpretable biological pathways. To address these shortcomings, we introduce "GWAS latent embeddings accounting for noise and regularization" (GLEANR), a MF method for detection of sparse genetic factors from summary statistics. GLEANR accounts for sample sharing between studies and uses regularization to estimate a data-driven number of interpretable factors. GLEANR is robust to confounding induced by shared samples and improves the replication of genetic factors derived from distinct biobanks. We used GLEANR to evaluate 137 diverse GWAS, identifying 58 factors that decompose the genetic architecture of input traits and have distinct signatures of negative selection and degrees of polygenicity. These sparse factors can be interpreted with respect to disease, cell-type, and pathway enrichment, such as three factors capturing platelet measure phenotypes enriched for disease-relevant markers corresponding to distinct stages of platelet differentiation. Overall, GLEANR is a powerful tool for discovering both trait-specific and shared pathways underlying complex traits from GWAS summary statistics.

Introduction

Genome-wide association studies (GWAS) have identified thousands of associations between common genetic variants and complex traits[1,2]. Efforts to understand genetic relationships between diseases often leverage pairwise comparison of GWAS[3,4] and have found evidence of widespread pleiotropy among genetic loci.[5,6] Studies of pleiotropy focus either on individual variants[4,5] or examine the genetic correlation[3] between GWAS, which detect covariance between the genetic effects of two phenotypes. Genetic correlation may arise from common biology among traits at multiple scales, such as shared variants, pathways, and cell types [7,8]. Pleiotropy can thus serve to support identification of important genetic pathways with multi-trait impact. Several studies[7-13] have shown that these relationships can be captured by latent genetic components (factors). These factors map genetically related traits to groups of variants belonging to shared pathways or cell types potentially missed by genome-wide analysis or narrow consideration of single variants.

Most study designs for estimating such genetic factors across many traits employ matrix factorization (MF) approaches [7,8,10,13]. MF explicitly decomposes an input matrix of GWAS effects for all traits into the product of two matrices; the first matrix maps SNPs to latent factors, and the second matrix maps latent factors to traits. While MF approaches are accessible and scalable, existing studies have a few notable limitations. First, standard approaches yield dense factors, where every trait is loaded on every factor, which in turn has non-zero effects for every SNP. This presents a challenge for interpretation, as the factors are less likely to represent distinct pathways, cell-types, or subsets of affected traits, and may increase the susceptibility to overfitting. Second, present studies have not consistently accounted for sources of potential confounding in GWAS data, including non-independence of target studies due to sharing of samples. Finally, downstream analysis of factors has focused primarily on characterization of the top variants and traits for each factor, while properties such as factor genetic architecture or distinct contributions of trait-relevant cell types remain unexplored.

To estimate interpretable genetic factors that are robust to differences in GWAS effect size uncertainty estimates and shared samples between studies, we developed GWAS latent embeddings accounting for noise and regularization (GLEANR). GLEANR uses dynamic model selection to yield sparse factors while statistically accounting for confounding covariance from sample sharing and variation in estimation error. In simulations, GLEANR outperforms other factorization methods when input studies share samples. By adjusting for the effects of sample sharing, our approach successfully estimates genetic components which are more replicable across distinct cohorts. In an evaluation of 137 GWAS traits across the phenome from the UK Biobank[14,15] (UKBB), GLEANR identifies genetic factors with distinct tissue enrichments, polygenic architectures, and signatures of selection. Among these, we identify three factors related to platelet traits which capture distinct stages of platelet formation related to inherited platelet disorders. GLEANR is a powerful tool for hypothesis-free exploration of genetic variants impacting shared pathways among component traits.

Results

Overview of GLEANR model and pipeline:
Complex traits are highly pleiotropic[3-6,16,17]. To capture common genetic pathways or components among multiple traits, we developed GLEANR, an MF approach tailored to GWAS summary statistics. GLEANR estimates sparse latent genetic components, decomposing the full set of genetic effects across traits into a lower dimensional representation with factors shared by

traits. Importantly, GLEANR accounts for both the uncertainty inherent in GWAS effect size estimates and the sharing of samples between studies.

GLEANR decomposes an input matrix of LD-pruned marginal effect sizes of N SNPs on M GWAS traits, $\mathbf{B} \in \mathbb{R}^{N \times M}$, as

$$\mathbf{B} = \boldsymbol{UV}^T + \boldsymbol{E}$$

where $\boldsymbol{V} \in \mathbb{R}^{M \times K}$ weights phenotypes on K genetic factors and $\boldsymbol{U} \in \mathbb{R}^{N \times K}$ reflects the SNP effects contributing to those factors. Both $\boldsymbol{U}$ and $\boldsymbol{V}$ are sparse, as not all phenotypes are genetically correlated, nor do all SNPs make pleiotropic contributions to these shared effects[4]. Hence, these matrices are estimated with sparsity-inducing L_1 regularization[18,19]. However, to account for pleiotropic genetic associations that impact many traits, the first (ubiquitous) factor is not subject to regularization in $\boldsymbol{V}$, and thus weighted on all traits. Each row of residual noise matrix $\boldsymbol{E}$ ($\boldsymbol{\epsilon_1}, \boldsymbol{\epsilon_2}, \dots \boldsymbol{\epsilon_N}$) corresponds to a single SNP across all studies and reflects both measurement uncertainty and non-genetic correlation between studies (e.g. sharing of samples). This is modelled as $\boldsymbol{\epsilon_n} \sim MVN_M\big(0, \mathrm{diag}(\boldsymbol{s_n}) \times \boldsymbol{C} \times \mathrm{diag}(\boldsymbol{s_n})\big)$, where $\mathrm{diag}(\boldsymbol{s_n}) \in \mathbb{R}_{>0}^{M \times M}$ is a diagonal matrix of GWAS standard errors for SNP n and $\boldsymbol{C}$ is a block matrix of non-genetic correlations between studies, which may be estimated directly from summary statistics using cross-trait LD-score regression[3] (XT-LDSR; Fig. 1A).

To estimate $\boldsymbol{U}$ and $\boldsymbol{V}$, GLEANR minimizes the bi-convex objective function in Fig. 1A through an iterative procedure of alternating generalized least squares regression. One of $\boldsymbol{U}$ or $\boldsymbol{V}$ is fixed to estimate the other via regression against $\mathbf{B}$, with matrix elements weighted by standard errors ($\boldsymbol{W_s}$) and transformed by a decorrelating matrix $\boldsymbol{W_c}$ derived from $\boldsymbol{C}$. This weighting and decorrelation help eliminate uninformative or spurious factors. GLEANR dynamically selects sparsity weights (α and λ) by minimizing the Bayesian Information Criteria (BIC) during alternating regression steps[20,21].

Factors can be directly interpreted by examining the non-zero entries in $\boldsymbol{U}$ and $\boldsymbol{V}$, along with their magnitude and direction, or may be used in downstream analysis as latent representations of genetic pathways shared by traits. Factors may capture narrow trait-specific contribution of variants, such as in singleton factors loaded on just one study, or may reflect broad variant effects, as in the unregularized ubiquitous factor (Fig. 1B). As representations of genetic pathways, factor SNP weights can capture distinct genetic architectures with tissue-marker and functional enrichments.

Figure 1: GLEANR decomposes GWAS summary statistics into sparse latent genetic factors. **A**) GLEANR maps GWAS SNPs (in U) and studies (in V) onto a latent K-dimensional space. Covariance of a single SNP n across studies is modelled as the matrix product of SNP standard errors ($diag(s_n)$) and a matrix of cohort-overlap induced correlations (C). GLEANR minimizes the indicated cost function, weighting terms by the inverse of standard errors (W_s) and the decorrelating transformation W_c derived from C. **B**) GWAS summary statistics provide estimates of SNP effect sizes ($\widehat{B}$) and standard errors ($\widehat{S}$) and are used to estimate correlation due to sample sharing ($\widehat{C}$). GLEANR performs model selection and fitting to select K factors loaded on traits and SNPs. A latent factor (e.g. F2) consists of traits (V_2) and relevant loaded SNPs (U_2). The first factor spans all traits (ubiquitous factor), and some factors may load on only a single study (singleton factors). We evaluate factors by considering their member traits, top SNPs, genes, tissue enrichments, and factor architecture.

Figure 2: GLEANR more accurately characterizes latent genetic factors in simulation than other tested methods and harmonizes genetic components across cohorts. A) Violin plot of the number of nominated factors from various factorization methods evaluating simulated GWAS with no genetic component at increasing levels of sample sharing between studies. Each point represents one of 840 simulations. B) Variance explained by estimated $\hat{U}$ across methods with respect to a known U, at varying sample sizes (on horizontal axis, labels at the top) and different covariance block structures (far right). Summary statistics were simulated from sparse U and V matrices. Larger R^2 indicates better reconstruction of U. C) Upper triangle heatmap of differences in variance-scaled estimates of inflation due to cohort overlap between FinnGen and UKBB for traits with non-zero differences, following GLEANR covariance shrinkage. D) Visual overview of analysis comparing UKBB versus FinnGen cohorts. Table reports global Pearson correlation between U from each cohort, estimated using GLEANR with (dark orange) or without (light orange) covariance adjustment. The GLEANR covariance adjustment procedure yields a more similar genetic basis across cohorts. (Trait abbreviations: BMI, body mass index; CAD, coronary artery disease; IBD: irritable bowel disease; T2D, type 2 diabetes; UC, ulcerative colitis)

Modest cohort overlap results in spurious factors for standard methods:

Overlap among samples between GWAS studies is commonplace, particularly as biobanks grow. Existing GWAS matrix factorization studies[10,22] commonly analyze phenotypes with substantial overlap, as reported on the UKBB Genetic Correlation Portal[3,23] (Web Resources; Fig. S1), including traits with both strong and weak genetic heritable components. To characterize the impact of sample sharing between GWAS on MF, we evaluated several factorization and model selection methods on GWAS derived from simulated, correlated phenotypes with no causal genetic component. By omitting a genetic signal, we isolated the impact of sample sharing since any discovered factors would be spurious. We evaluated methods including the singular value decomposition (SVD), testing both the elbow point of eigenvalues (SVD (elbow)) and factors corresponding to eigenvalues greater than the average eigenvalue[10] (SVD (avg)) to select the number of factors (K). We also added our own modification of SVD adjusted with a decorrelating transformation[4,24] to account for sample sharing (SVD-adj), as a baseline method that does adjust for sample sharing, and the empirical Bayesian method flash[25] (which automatically performs model selection). We compared these methods with GLEANR, both with covariance adjustment (standard GLEANR) and with no adjustment (unadjusted GLEANR or GLEANR-U). In these simulations, GLEANR and SVD-adj were provided with a cohort overlap correlation matrix estimated from phenotype correlations and the number of overlapping samples[24] (Methods).

We simulated genotypes based on minor allele frequencies (MAF > 0.01) reported for Europeans in the UKBB[14,26] among LD-pruned HapMap3[27] SNPs. For phenotypes, we randomly simulated values from a Gaussian distribution with mean and variance based on estimates from European UKBB[15] individuals for height. We simulated two additional related phenotypes roughly based on relationship between height, weight, and BMI in the UKBB, assuming weight is linearly related to height and BMI is based on their ratio, both with added Gaussian noise (Methods). Using linear regression with the simulated genotypes and phenotypes, we estimated summary statistics at increasing levels of sample sharing in three populations for a total of nine GWAS, each conducted across 30,000 simulated individuals. This approach ensured that correlations between simulated phenotypes were reasonably approximate to those in real traits.

Given the independent generation of genotypes and phenotypes, no true genetic factors influence traits in this simulated data. In this scenario, we found that sharing of less than 25% of study individuals was sufficient to cause baseline MF model selection procedures to nominate spurious components (Fig. 2A). By contrast, the covariance adjustment procedure used in GLEANR avoided detection of spurious factors even at high levels (>90%) of shared samples (Fig. 2A, dark orange, far right panel). While GLEANR-U also avoided detection of spurious factors better than most other methods (light orange) at low and moderate sample overlap, presumably due to enforced sparsity, at the highest levels of cohort overlap GLEANR-U also yielded spurious factors. Both sparsity-inducing penalty and direct adjustment contribute to GLEANR performance in the presence of cohort overlap. Thus, in the absence of genetic signal, sample sharing results in the detection of spurious factors unless appropriate statistical adjustment is made.

GLEANR recapitulates ground truth genetic factors even with cohort overlap:

To evaluate MF estimation of known genetic factors, we simulated GWAS summary statistics from prespecified U and V as $\mathbf{B} = UV^T + E$. Our simulated U and V consisted of five factors drawn from a sparse genetic architecture (Methods). We added to this noise (E) drawn from a multivariate normal distribution, with covariance reflecting simulated standard error across multiple sample sizes (from 5,000 to 200,000) and three cohort overlap structures (single

block, two block, and no overlap, far right of Fig. 2B). These overlap structures correspond to estimated sample sharing found in existing UKBB GWAS factorization studies[10,22] (Fig. S1A). GLEANR and SVD-adj were provided with the simulated cohort overlap correlation matrix. We evaluated flash with both the default and the "Kronecker" covariance setting, which can account for more complex residual covariance structures, and FactorGo[22], a variational model designed specifically for large-scale GWAS factorization. To assess fit, we first matched columns of $\widehat{U}$ and $\widehat{V}$ to those in U and V by maximizing the correlation between factor pairs, and then calculated the R^2 across all matched factors, scaled to unit-norm.

In simulations with no cohort overlap (bottom row, Fig. 2B), GLEANR performs comparably to other methods for samples between 10,000 and 200,000 individuals. In simulations with either a single-block and two-block sample overlap correlation, GLEANR reconstructions of U (Fig. 2B) had higher median R^2 than all tested factorization methods in sample sizes above 10,000, with similar performance for V (Fig. S2A). This is consistent with other metrics of fit, including Cohen's Kappa, which measured discrimination between signed and 0 elements (Fig. S2B and C). Simply decorrelating Z-scores (SVD-adj) did not consistently improve reconstruction of the latent space relative to unadjusted SVD. At sample sizes below 10,000, GLEANR was slightly outperformed by SVD-based approaches. This is due to sparsity constraints removing signals when effects are very small or noisy, suggesting that GLEANR may not perform better than baseline methods for GWAS with very small sample sizes. Overall, GLEANR is robust to simulated confounding due to sample sharing between studies and this adjustment improves reconstruction of genetic factors relative to other common methods.

Sample sharing adjustment improves concordance of factors between distinct cohorts:
To evaluate GLEANR 's covariance adjustment procedure on published GWAS, we assessed the replication of factors discovered from multiple traits in two distinct cohorts: UKBB[26] and FinnGen[28]. Specifically, the two large cohorts have no samples in common, but within each cohort multiple traits are measured over highly overlapping sets of individuals. We hypothesized that adjusting for sample sharing within each study during factorization would yield genetic factors that were more consistent between cohorts, reflecting more accurate factors due to elimination of confounding from sample overlap.

We selected 42 diverse phenotypes with summary statistics publicly available from both the UKBB[14,26,29] and FinnGen[28] to evaluate with GLEANR (Table S1). Traits were chosen to have strong genetic correlation ($r_g \geq 0.8$, with adjusted p < 0.01) between cohorts. Estimates of sample sharing were generated using XT-LDSR[3], and adjusted to account for uncertainty and trait heritability (see Methods). These estimates reflected distinct cohort-specific sample sharing structure (differences visualized in Fig 2C), especially among several cardiac phenotypes (i.e. angina, myocardial infarction, CAD). We applied GLEANR and GLEANR-U to GWAS across traits from FinnGen, and then separately to GWAS from the UKBB. To minimize differences related to model selection, we initialized GLEANR at $K_{init} = 41$ for both cohorts.

To determine how correcting for cohort-specific sample overlap affected detection of genetic factors, we compared the estimated factors ($\widehat{U}, \widehat{V}$, Tables S2-S9) and reconstructed effect sizes ($\widehat{B} = \widehat{U}\widehat{V}^T$) between UKBB and FinnGen. GLEANR nominated both fewer total and a more consistent number of final factors than GLEANR-U (13/12 vs 19/17) and yielded a 4% reduction in overall RRMSE between UKBB and FinnGen reconstructed effect sizes. Globally, GLEANR factors were more strongly correlated across cohorts than GLEANR-U factors (0.62 vs 0.59 for U, Fisher's Z-test p < 0.001, Fig 2D), scaled to unit length. This improvement in factor similarity is consistent with the estimated magnitude of structural differences in cohort overlap between UKBB and FinnGen (Fig. 2C,. Fig. S3). This suggests that GLEANR factors are more accurate and less reflective of confounding, non-genetic effects.

GLEANR removed multiple singleton factors from both UKBB and FinnGen (8 and 5, respectively) unadjusted analyses, including a factor loaded on lipo-protein metabolism disorders present in both cohorts (Fig. S4). Importantly, a key heart disease related factor connecting angina pectoris, heart attack, ischemic heart diseases and coronary atherosclerosis remained across cohorts even after covariance adjustment, despite also being influenced by sample sharing. This demonstrates that GLEANR is able to appropriately adjust for cohort-specific patterns of sample sharing while preserving pleiotropic signals between traits.

Figure 3: GLEANR analysis across 137 diverse European UKBB GWAS phenotypes yields 58 factors. A) Traits span 19 phenotype categories from across the human phenome. B) Scree plot of variance explained by 58 factors estimated by GLEANR, from F1 to F58. Factors are ordered in approximately decreasing order of data variance explained. C) Heatmap representation of selected factors in V and the traits with non-zero entries in those factors. Trait colors match the corresponding categories in panel A, and columns are scaled to have Euclidian norm of 1. Selected factors include the first seven factors explaining the largest proportion of data variance in addition to six other factors discussed in the text.

GLEANR across the human phenome identifies 57 sparse factors and one ubiquitous factor:

To explore distinct genetic factors spanning the human phenome, we evaluated 137 diverse GWAS conducted on Europeans from the UKBB[14,26], chosen so none had pairwise genetic

correlation r_g > 0.7 (see Methods). These traits span 19 trait categories[30] (Fig 4A), including metabolic, skeletal, behavioral, respiratory, and immunological phenotypes (see Table S10 for the full list). Across these, we evaluated HapMap3[27] SNPs with MAF > 0.01 that achieved a nominal $p < 1 \times 10^{-5}$ in at least one study, then LD clumped resulting in a total of 41,259 variants. Our LD clumping procedure prioritized inclusion of variants achieving nominal significance in the largest number of target studies. Overall, only 1,455 variants achieve this p-value threshold in 10 or more traits, with nearly half of SNPs (20,137) passing this threshold in just one trait (Fig. S11).

From the 137 input phenotypes, GLEANR identified 58 factors, 27 of which are loaded (e.g. had non-zero values) on only a single trait (singleton factors) and one (the uqibuitous factor, F1) loaded on all 137 traits (Tables S11-12). The remaining factors were loaded on 2 to 13 traits (mean 3.6). Each trait was weighted on between one and ten factors (median 2). On average, 12.7% of SNPs per loading had values of exactly zero. Collectively, these 58 factors explained 72.6% of variance in the covariance-adjusted summary statistics (Fig 4B; a selection of 13 factors are visualized as a heatmap in 4C).

Factor 1, which explains the largest proportion of variance from input summary statistics (5.5%), was loaded on all traits (ubiquitous factor). SNP loading scores (U_1) correlated strongly ($\rho = -0.75$) with the mean effect size per SNP across input GWAS, while trait scores (V_1) correlated with the average genetic correlation of each input GWAS relative to all others ($\rho = -0.80$; Fig. S5B). Of all traits we analyzed, 42% loaded only on the ubiquitous factor (values of exactly zero on all other factors). These traits had fewer nominally significant SNPs included in our analysis than traits loaded on multiple factors (median of 29 SNPs per study, 1,013 SNPs per study, respectively), and were predominantly categorical phenotypes (88%). GLEANR loads traits with few high-confidence genetic associations on the ubiquitous factor, which reflects average SNP effects across all input studies.

To test for tissue-specific regulatory effects captured by ubiquitous factor variants, we estimated heritability enrichments among chromatin tissue markers with stratified-LDSR[31,32] (S-LDSR). Factor 1 SNPs had heritability enriched in central-nervous system related tissues ($q = 1.74 \times 10^{-21}$), particularly in brain markers in the fetal brain, prefrontal cortex, and germinal matrix tissues (all $q < 1.0 \times 10^{-7}$; Fig. S5A). Our analysis included several brain-related phenotypes (16 psychiatric, 5 neurological, and 5 cognitive traits), ten of which are loaded only on F1. This enrichment may also reflect the influence of polygenic phenotypes correlated with BMI (including whole body fat mass and whole body impedance), which has reported connections to brain expression markers[32–34] and proteins[35.] Consistent with this, gene set enrichment of top F1 genes (see Methods) in DisGeNET[36] gene sets nominated BMI ($q = 2.1 \times 10^{-8}$), intelligence ($q = 2.88 \times 10^{-7}$), and cognition ($p = 1.45 \times 10^{-5}$) as top traits associated with F1 genes. Thus, the ubiquitous factor captures average effects across each variant and implicates brain-specific tissue markers contributing to a pathway shared broadly across the studies we evaluated.

Figure 4 : GLEANR factors reflect diverse genetic architectures and distinguish gene activity at specific stages of platelet formation. A) Relative polygenicity (x axis) as a proportion of the total number of SNPs, and S_k (y axis) per factor. Points are colored by the number of traits loaded on each factor, and factors referenced in the text are labelled. B) Relationship between SNP factor effect size and minor allele frequency on F1 (weak selection signature) and F32 (strong selection signature). U effects have been scaled to have unit variance for visualization. C) Schematic representing stages of megakaryopoiesis and platelet formation, with top factor genes associated with inherited platelet disorders listed below. Gene color indicates the factor in which it is a top factor gene, and size indicates the average frequency with which it is nominated as a top factor gene. **D)** Factor trait weight of Factors 4,11,

and 23 across mean platelet volume (MPV), platelet count (PCT), and platelet distribution width (PDW). E) Enrichment of top factor genes in inherited platelet disorder genes at stages of platelet formation. Bars represent 90% confidence intervals and color indicates significance in permutation enrichment test. F) Cell type enrichment from PanglaoDB cell-type gene signatures for each factor in hematopoietic stem cells (HSCs), megakaryocytes, and platelets.

GLEANR factors span diverse polygenic architectures and signatures of selection:
Genetic architectures vary widely across traits[30,37,38] and regions of the genome[38]. Studies modeling GWAS effects as mixtures of multiple distributions[39,40] have provided evidence that complex traits have contributing components with different genetic architectures. We hypothesized that factors decomposing GWAS may have underlying polygenic architectures and signatures of selection that could meaningfully differ from their composite traits.

To characterize these properties, we first estimated factor polygenicity using a previously reported measure of the effective number of associated SNPs[37]. Scores are reported as a proportion of all SNPs under evaluation, with larger scores indicating a larger number of SNPs contributing to the factor[37] (Fig. 4A, x axis). Reported estimates reflect our ascertainment of pleiotropic variants in heritable traits, so are not directly comparable to genome-wide scores. F1 is among the most polygenic factors, with a broad effect size distribution and fewer than 8% of SNPs with a weight of zero. Other top polygenic factors (F52, F57, and F58) are loaded on neurological and behavioral traits including difficulty sleeping, narcolepsy, fluid intelligence score, smoking, and chronic pain. This is consistent with reported findings that behavioral and cognitive phenotypes are highly polygenic[30,37]. Factors with lower estimated polygenicity include singleton factors loaded on low-density lipoprotein (LDL; F18), and bilirubin (F32).

Second, we quantified the relationship between factor effect sizes and SNP allele frequencies to examine signatures of negative selection[30,41,42]. In this model, the variance of factor SNP weights are a function of MAF (previously reported as the α-model[42]), which we report by the parameter S_k[30]. Large negative values of S_k correspond to large effect variants enriched at low allele frequencies, often interpreted as a signature of negative selection[30,42]. F1 had the second-smallest S_k among all factors $(S_1 = -0.76)$, while F32, which had minimal polygenicity, reflected the strongest selection signature $(S_{32} = -1.17;$ Fig 4B). We emphasize that large S_k magnitudes likely reflect the ascertainment of the included variants, which all have nominal genome-wide significance in at least one phenotype. Thus, we limit the interpretation of S_k to a relative scoring among factors. All S_k are negative, and strong selective effects tend to appear in less polygenic architectures (Fig 4A). Factors with S_k scores of greater magnitude were loaded on fewer traits, suggesting that latent traits under strong selection may be less likely to share latent genetic effects with other phenotypes. This demonstrates that GLEANR decomposes traits into factors spanning a range of polygenic architectures and signatures of selection.

The four factors with the largest S_k magnitudes were all singleton factors, loaded on total bilirubin (F32), skin color (F37), blonde hair (F43), or low-density lipoprotein (LDL, F18). Factors 32 and 18 are most prominently enriched for heritability in liver-specific chromatin markers (S-LDSR tissue-adjusted factor q = 0.00026 and 0.00092 respectively, see also Table S13), consistent with the role of liver in cholesterol metabolism. We tested top factor genes corresponding to lead factor variants (matched using the Open Targets platform[43,44]) for KEGG pathway enrichments[45] (see Methods). F18 genes were enriched in cholesterol metabolism, fat digestion and absorption, and glycerolipid metabolism, reflecting cholesterol production and recycling pathways. F32 top factor genes were enriched in bile secretion and glycerolipid metabolism pathways, with top SNPs near UGT genes involved in bilirubin metabolism[46], capturing biology related to cholesterol elimination through the bile. F18 and F13 are minimally correlated $(\rho = 0.04)$, corresponding to distinct components of lipid metabolism with evidence of

negative selection, consistent with previous reports of a strong selection signature for dislipidemia[30] in genomes of Euoropean ancestry.

We also observed large S_k estimates on F37 and F43 (singletons for skin color and blonde hair, respectively), consistent with prior positive selection scans on European ancestry genomes [47,48]. Pigmentation genes under selection[47,48], including OCA2, HERC2, SLC45A2, TYR, and TYRP1, were among the top factor genes All together, GLEANR identifies genetic factors with prominent allele frequency-effect size relationships that reflect genetic loci previously reported in selection scans.

Factors weighted on platelet traits distinguish stages of platelet formation:

GLEANR factors across UKBB GWAS identified instances of disease-relevant biology shared across traits. Our analysis included three platelet-related phenotypes (mean platelet volume (MPV), platelet count (PLT), and platelet distribution width (PDW)) which were decomposed into three distinct factors (F4, F11, F23), in addition to the ubiquitous factor (F1; see Fig. 4D). To characterize differences between F4, F11, and F23 (henceforth, "platelet factors"), we closely examined their SNP loadings, genetic architecture profiles, and top factor gene enrichments. As expected, top variants nominated by these factors correspond to GWAS associations[43] for these phenotypes with strong evidence for regulatory roles as expression quantitative trait loci (eQTLs) in whole blood (Table S14). For instance, rs2228367, the second lead variant in F4, is a replicated GWAS hit for all three phenotypes[23,49,50] and an eQTL in whole blood in multiple studies[51,52] for tropomyosin 4, an actin-binding protein with well-documented effects in platelet diseases.[53,54]

Among platelet factors, F4 had the strongest selection coefficient estimate ($\hat{S} = 0.95$; fifth largest magnitude in the entire analysis). Consistent with negative selection acting on large effect SNPs, top F4 SNPs have larger average effect size magnitudes across MPV, PCT and PDW than F11 (Wilcox test, p<2e-16) and F23 (p = 5.5e-6) SNPs (Fig. S6). Given this selection signature, we speculated that variants captured by F4 might adversely influence human fitness.

We also observed that PCT and MPV have opposite directions of effect in F4 (Fig 5D), suggesting that F4 SNPs drive platelet volume and platelet count in opposing directions. This is consistent with previously reported negative correlation between GWAS effects for these traits[55] and with negative correlation of the traits themselves across patients[56,57]. Published work has hypothesized that alleles contributing to both lower PLT and elevated MPV could explain hereditary cases of the blood condition macrothrombocytopenia[55]. Hereditary macrothrombocytopenia (HMTP) is characterized by elevated platelet size (>12 femtoliters) and low platelet counts (< 150,000/μL)[58] and is associated with many other blood-related diseases (e.g. Sitosterolemia, myopathy, Grey Platelet Syndrome, Bernard-Soulier syndrome, cancer risk)[55]. While previously thought to be a Mendelian condition, growing evidence from GWAS suggests polygenic contributions to both HMTP and other blood diseases [50,55].

To evaluate whether F4 captured genetic effects contributing to HMTP, we tested F4 top factor genes for enrichment in known HMTP -causing genes [55]. Given that for some genes multiple SNPs were mapped to the same gene in our analysis, we considered only unique genes in our enrichment tests and used random subsamples of our background gene set to generate an empirical null reflecting this imbalance (Methods). We found that F4 was strongly enriched for HMTP genes (OR = 6.5, empirical p = 1e-04), as were F11 and F23 (OR 6.1 and 6.3, respectively), suggesting a possible source of differential fitness that may impact F4's selection signature.

Next, we sought to explore the cell-type and context specificity of the platelet factors, noting that disease-relevant processes can occur in early and transient stages of cellular differentiation[59–61] in addition to terminal and adult cell types[51,62,63]. We tested each factor for enrichment in genes known to cause inherited platelet disorders at distinct stages of platelet

formation[64,55], specifically during early megakaryopoiesis, late megakaryopoiesis, proplatelet formation, and mature platelet function (Fig 4C). We observed that F23 was strongly enriched for inherited platelet disorder genes in early megakaryopoiesis (corrected permutation q<0.00008) and was the only factor to capture thrombopoietin (THPO), the critical growth factor promoting megakaryopoiesis[65], as a top gene. F11 related to late megakaryopoiesis (q = 0.02), and F4 reflected the later stages of proplatelet formation (q< 0.00008) and mature platelet function (q<0.00008; Fig 4E).

The correspondence between factors and different stages of platelet formation is further supported by complementary cell-type and GO-term enrichments. In an enrichment test of top factor genes in cell type gene markers from the PanglaoDB[66,67], the strongest detected enrichment for F23 was in hematopoietic stem cells, while F11 was most strongly enriched in megakaryocyte gene markers, and F4 in platelets (Fig 4F), though these enrichments are not significant after adjusting for multiple hypothesis testing across all cell types.

Consistent with extensive transcriptional regulation during early megakaryopoiesis, F23 had top GO molecular function enrichments in DNA binding (q=0.005) and transcription factor binding (q=0.04) and had GO biological process enrichments in myeloid cell development (q=0.03), myeloid cell differentiation (q=0.08), and hemopoiesis (q=0.03). Capturing the cytoskeletal dynamics involved in proplatelet formation, F4, had top GO molecular function enrichments including tubulin binding (q=0.03), cadherin binding (q=0.03), and microtubule binding (q=0.05, Fig. S7). Our observations across these platelet factors provide strong evidence that sparse factorization of GWAS by GLEANR can decompose trait effects into interpretable components that capture distinct disease-relevant cellular processes.

Methods

GLEANR Model:
GLEANR operates on marginal GWAS summary statistics, with effect size estimates in the matrix $\mathbf{B} \in \mathbb{R}^{N \times M}$ and standard error estimates $\mathbf{S} \in \mathbb{R}_{>0}^{N \times M}$. SNPs are indexed by $n \in [1\ldots N]$ and traits by $m \in [1\ldots M]$. Typically, substantial covariance is observed across both columns and rows due to sample sharing among traits and linkage disequilibrium between SNPs, respectively. To account for the impact of linkage disequilibrium, GLEANR requires SNPs under evaluation to be approximately uncorrelated, selected by an appropriate pruning or clumping procedure (e.g. $r^2 < 0.3$ across 250KB windows). We model the covariance associated with estimate error and sample sharing as follows:
Let $\boldsymbol{\beta_n}$ be a single row of $\mathbf{B}$ and $\boldsymbol{s_n}$ a single row of $\boldsymbol{S}$ along SNP n, where:

$$\boldsymbol{\beta_n} \sim MVN\left(\boldsymbol{Vu_n}, \mathrm{diag}(\boldsymbol{s_n}) \times \boldsymbol{C} \times \mathrm{diag}(\boldsymbol{s_n})\right) \tag{1}$$

$\boldsymbol{V} \in \mathbb{R}^{M \times K}$ maps M related traits onto K latent factors, so traits sharing a pleiotropic genetic factor have non-zero values along the same column of $\boldsymbol{V}$. $\boldsymbol{u_n} \in \mathbb{R}^{K \times 1}$ (a single row of the matrix $\boldsymbol{U} \in \mathbb{R}^{N \times K}$) is a sparse vector indicating the contribution of variant N to each of K latent factors. We allow $\boldsymbol{U}$ and $\boldsymbol{V}$ to be signed, as the parity with which a SNP contributes to a phenotype association (e.g. if it is a risk or protective SNP) is dictated by the arbitrary measure of traits. $\mathrm{diag}(\boldsymbol{s_n}) \in \mathbb{R}^{M \times M}$ contains standard errors for SNP n across all studies along the diagonal, and $\boldsymbol{C} \in \mathbb{R}^{M \times M}$ is a block correlation matrix of non-genetic confounding covariance structure between GWAS.

GLEANR imposes sparsity constraints on both $\boldsymbol{U}$ and $\boldsymbol{V}$ such that $\sum_{i=1,j=2}^{M,K}|V_{i,j}| < t_\lambda$ and $\sum_{i,j=1}^{N,K}|U_{i,j}| < t_\alpha$, where t is a threshold corresponding to the LASSO regularization weights for $\boldsymbol{V}$ and $\boldsymbol{U}$ (λ and α respectively, see below).

Model fitting procedure:

GLEANR model fitting begins by initializing V to the first $K_{init} - 1$ right singular vectors of B. An additional column is appended to reflect the dominant direction of correlation across all studies, with signs corresponding to the first eigenvector of $\text{cor}(\mathbf{B})$. We perform alternating iterations of generalized least squares with L1 regularization to update $\widehat{U}$ by regressing $\widetilde{B}$ against $\widetilde{V}$, where the tilde indicates elements of the matrix have been transformed to adjust for GWAS estimate uncertainty and confounding correlation (rows of $\widetilde{B}$ are $\widetilde{\boldsymbol{\beta}_n} = \mathbf{W}_c \text{diag}(s_n)^{-1}\boldsymbol{\beta}_n$ and $\widetilde{V} = \mathbf{W_c}\text{diag}(s_n)^{-1}V$). We then regress $\widetilde{B}$ against $\widetilde{U}$ to update $\widehat{V}$, where elements of $\widetilde{U}$ are $\widetilde{U}_{n,k} = W_{c(j,m)}S_{n,m}U_{n,k}$ expanded over all $j, m \in [1..M]$ (see Suppl. Note 1). We repeat alternating regressions until our objective function achieves convergence (Equation 2):

$$L(\mathbf{B}, \boldsymbol{W}, \boldsymbol{W_c}, \boldsymbol{U}, \boldsymbol{V}, \alpha, \lambda) = \min_{U,V} \frac{1}{2}\left\| \boldsymbol{W_c}\left(\boldsymbol{W_S} \odot \left(\mathbf{B} - \boldsymbol{UV}^T\right)\right)^T \right\|_F^2 + \alpha\|\boldsymbol{U}\|_{1,1} + \lambda\left\|\boldsymbol{V}_{\backslash\mathbf{1}}\right\|_{1,1} \qquad (2)$$

$\boldsymbol{W_S}$ is a matrix of weights with elements $W_{S(n,m)} = \frac{1}{\hat{s}_{n,m}}$ and $\boldsymbol{W_c} \in \mathbb{R}^{N \times M}$ is a decorrelating transformation derived from $\widehat{C}$ (see below). Here, $\|\boldsymbol{X}\|_{1,1} = \sum_{i,j=1}^{N,K}|X_{i,j}|$, is the entry-wise L1 matrix norm. The first column of V is unregularized to retain the dominant correlation structure across traits.

This objective monotonically decreases at each iteration because it is bi-convex (see Fig. S8) and achieves convergence when the change between iterations is smaller than a specified small threshold ε, with a default of < 0.1% of the previous iteration's value. This procedure is implemented using sparse matrix functions in *glmnet in R 4.2.*3 (Suppl. Note 1). This model fitting procedure is summarized below:

Algorithm 1: GLEANR model fitting

Input: $\mathbf{B} \in \mathbb{R}^{N \times M}, \boldsymbol{W}_c \in \mathbb{R}^{M \times M}, \boldsymbol{W_S} \in \mathbb{R}^{N \times M}, \widehat{\boldsymbol{V}}_{init} \in \mathbb{R}^{M \times K}, \alpha, \lambda, K_{init}$

Output: $\widehat{\boldsymbol{V}} \in \mathbb{R}^{M \times K}, \widehat{\boldsymbol{U}} \in \mathbb{R}^{N \times K}$

while $\%\Delta L\left(\mathbf{B}, \boldsymbol{W}, \boldsymbol{W}_c, \widehat{\boldsymbol{V}}, \widehat{\boldsymbol{U}}, \alpha, \lambda\right) > \varepsilon$ **do**

$\qquad \widehat{\boldsymbol{U}} \leftarrow \min_U \frac{1}{2}\left\| \boldsymbol{W_c}\left(\boldsymbol{W_S} \odot \left(\mathbf{B} - \boldsymbol{U}\widehat{\boldsymbol{V}}^T\right)\right)^T \right\|_F^2 + \alpha\|\boldsymbol{U}\|_{1,1}$

$\qquad \widehat{\boldsymbol{V}} \leftarrow \min_V \frac{1}{2}\left\| \boldsymbol{W_c}\left(\boldsymbol{W_S} \odot \left(\mathbf{B} - \widehat{\boldsymbol{U}}\boldsymbol{V}^T\right)\right)^T \right\|_F^2 + \lambda\left\|\boldsymbol{V}_{\backslash\mathbf{1}}\right\|_{1,1}$

end while

Estimation of C and decorrelating transformation ($\boldsymbol{W_c}$)

$\widehat{\boldsymbol{C}}$ may be estimated from the intercept term of XT-LDSC[3,4], from phenotype correlations and the number of overlapping samples[24] (when available), or from the empirical correlation of truncated z-statistics[68,69]. In this work we favor estimates from XT-LDSC.

Empirically, GLEANR is sensitive to large elements $\widehat{\boldsymbol{C}}$. To mitigate these effects and account for uncertainty in estimates of $\widehat{\boldsymbol{C}}$, we apply shrinkage to its elements in the following steps:

1. Require $\widehat{\boldsymbol{C}}$ to have a block structure, reflecting our expectation that $\boldsymbol{C}$ is sparse and contains clusters of correlated traits[24]. Block elements are identified through hierarchical clustering on the matrix of cohort overlap estimates, followed by pruning branches with a cutoff corresponding to pairwise correlations ≥ 0.2.

2. Apply Wolf-Ledoit shrinkage[70] to the block correlation matrix, adjusting for uncertainty in correlation estimates[71] by calculating a matrix $\widehat{\boldsymbol{C}}_S$:

$$\widehat{\boldsymbol{C}}_S = \gamma I + (1 - \gamma)\widehat{\boldsymbol{C}}, \text{ where } \gamma = \frac{\sum_{i \neq j} \widehat{Var}(\hat{c}_{i,j})}{\sum_{i \neq j} \hat{c}_{i,j}^2}$$

This procedure has the added benefit of yielding a theoretically positive definite matrix $\widehat{\boldsymbol{C}}_S$ as the linear combination of positive definite and semi-definite matrices[70].

When $\widehat{\boldsymbol{C}}$ is estimated using XT-LDSR,[3] GLEANR provides the option to scale $g_{cov-int}$ statistics by the product of the standard deviation of input traits, to adjust for the selection of heritable traits in analysis.

Model selection procedure:
GLEANR has three model hyper-parameters: α, λ (which correspond to the degree of sparsity in U and V, respectively), and the number of factors K_{init} at which to initialize GLEANR. Although multiple approaches for model selection exist in the context of matrix factorization[20,72–74], we employ the Bayesian Information Criteria (BIC)[20,72,75,76] due to its interpretability and computational efficiency. In high-dimensional problems, the BIC favors overly complex models[77,78], so we include an additional term[77] to adjust for models with many covariates receiving a higher likelihood under the standard prior. We calculate the BIC as:

$$BIC_\alpha = \frac{\left\| \mathbf{B}^* - \widehat{\boldsymbol{U}} \boldsymbol{V}^{*T} \right\|}{\sigma_{OLS}^2} + \log(MN) \left\| \widehat{\boldsymbol{U}} \right\|_{0,0} + 2\gamma \log \left(\frac{p}{\left\| \widehat{\boldsymbol{U}} \right\|_{0,0}} \right) \tag{5}$$

$$BIC_\lambda = \frac{\left\| \mathbf{B}^* - \boldsymbol{U}^* \widehat{\boldsymbol{V}}^{T} \right\|}{\sigma_{OLS}^2} + \log(MN) \left\| \widehat{\boldsymbol{V}} \right\|_{0,0} + 2\gamma \log \left(\frac{p}{\left\| \widehat{\boldsymbol{V}} \right\|_{0,0}} \right) \tag{6}$$

where $\left\| \widehat{X} \right\|_{0,0} = \sum_{n=1}^{N} \| x_n \|_0$ (e.g. the number of non-zero elements in the matrix), is an unbiased estimate for the degrees of freedom of the lasso[21], and $\gamma = 1 - \log(NM)/2\log(P)$ is scaling term that controls the contribution of the BIC extension[77] (last term). p is the total number of covariates under consideration in each model comparison (NK for $\boldsymbol{U}$ or MK for $\boldsymbol{V}$). This "extended" BIC yields more consistent results across GWAS cohorts than the traditional BIC in our tests (see Fig. S9).

Selection of α, λ, and K_{init}:
During GLEANR's initial iterations, we select either the λ (when fitting $\boldsymbol{V}$)or α (when fitting $\boldsymbol{U}$) which minimizes the BIC of the current step until $L_{BIC} = BIC_\alpha + BIC_\lambda$ converges, at which point α, λ are fixed. We fit U at 100 settings of α along the log-scale of possible values (from α_{max}, where all of $\widehat{U} = 0$, to $\alpha_{max} \times 10^{-4}$). GLEANR selects $\widehat{\boldsymbol{U}} = \min_\alpha(BIC_\alpha)$, then repeats this procedure with respect to $\boldsymbol{V}$ and λ. Alternating iterations continue until L_{BIC} is no longer declining (or for at least 5 iterations), at which point α and λ are fixed at $\alpha, \lambda = \underset{\alpha,\lambda}{\operatorname{argmin}} L_{BIC}$ and GLEANR proceeds with model fitting (see Algorithm 1). In practice, we observe that L_{BIC} typically converges in fewer than 10 iterations (Fig. S10).

To select K_{init}, GLEANR performs an adaptive grid search to select the K_{init} which minimizes L_{BIC} at convergence (see Suppl. Note 1). GLEANR first samples K_{init} from quantiles of the range $[1, M - 1]$, then tests additional settings in the neighborhood of $K_{min} = \underset{K_{init}}{\min}(L_{BIC})$, halfway between previously tested values. This procedure continues for a specified number of search points (default =10) or until K_{min} no longer changes.

Algorithm 2: GLEANR model selection

1 Input: $\mathbf{B} \in \mathbb{R}^{N \times M}, \boldsymbol{W}_c \in \mathbb{R}^{M \times M}, \boldsymbol{W}_S \in \mathbb{R}^{N \times M}, K_{init}$

2 Output: $\widehat{\boldsymbol{V}} \in \mathbb{R}^{M \times K}, \alpha, \lambda, K_{init}$

3 Initialize $\widehat{\boldsymbol{V}} = [\text{sign}(\text{SVD}(\text{cor}(\mathbf{B}), K = 1), \text{SVD}(\mathbf{B}, K = K_{init} - 1)]$

4 **while** $\Delta L_{BIC}^{(K_{init})} > 0$ **do**

5
$$\widehat{\boldsymbol{U}}, \alpha \leftarrow \min_{\alpha} \text{BIC}_\alpha (\min_{U} \frac{1}{2} \left\| \boldsymbol{W}_c \left(\boldsymbol{W}_S \odot (\mathbf{B} - \boldsymbol{U}\widehat{\boldsymbol{V}}^T) \right)^T \right\|_F^2 + \alpha \|\boldsymbol{U}\|_{1,1})$$

6
$$\widehat{\boldsymbol{V}}, \lambda \leftarrow \min_{\lambda} \text{BIC}_\lambda (\min_{V} \frac{1}{2} \left\| \boldsymbol{W}_c \left(\boldsymbol{W}_S \odot (\mathbf{B} - \widehat{\boldsymbol{U}}\boldsymbol{V}^T) \right)^T \right\|_F^2 + \lambda \|\boldsymbol{V}_{\backslash \mathbf{1}}\|_{1,1})$$

7 $\quad L_{BIC}^{(K_{init})} \leftarrow BIC_\alpha(\widehat{\boldsymbol{U}}) + BIC_\lambda(\widehat{\boldsymbol{V}})$

8 **end while**

9 $\quad \alpha, \lambda, K_{init}, \widehat{\boldsymbol{V}} \leftarrow \min_{\alpha, \lambda, K_{init}} (L_{BIC}^{(K_{init})})$

Simulations:

Our study included two simulation frameworks: 1) to isolate the impact of sample sharing on the detection of spurious genetic factors and 2) to test reconstruction of known genetic factors. In (1), we directly specified the degree of sample sharing when estimating summary statistics. In (2), we simulate a latent basis $(\boldsymbol{U}, \boldsymbol{V})$ and add noise corresponding to covariance structure induced by sample sharing between studies.

1) Direct simulation of GWAS summary statistics:

Three populations of 91,000 individuals were simulated, with genotypes for 1000 SNPs sampled from Bernoulli distributions for maternal and paternal alleles. Count probabilities came from minor allele frequencies of randomly selected pruned SNPs nominated by UKBB studies in our comparative UKBB-FinnGen analysis. Phenotypes were modeled based on the height, weight, and BMI of 10,000 UKBB individuals. Population parameters (height mean and variance and weight predicted from height) were used to generate 3 random phenotypes as; $p_1 \sim N(169.9, 86.5)$, $p_2 = 0.93p_1 - 80 + \epsilon_2, \epsilon_2 \sim N(0,182.3)$ and $p_3 = \frac{p_2}{p_1^2} + \epsilon_3, \epsilon_3 \sim N(0,1)$

We selected 30,000 individuals from each cohort per phenotype and estimated GWAS summary statistics with linear regression on standardized phenotypes, such that no individual was used more than once (for a total of 9 sets of summary statistics). We repeated this, reusing samples within a cohort at levels of 10,15,20,23,26, and 30-thousand repeated samples. We considered all unique pairwise combinations of overlap levels across the 3 cohorts for a total of $\binom{7}{3}_{w/rep.} = 84$ overlap settings, repeated 10 times each.

We performed matrix factorization and model selection on each of these 840 sets of summary statistics using the singular value decomposition (SVD), flash, and GLEANR, as described in Table 1. SVD was implemented using the BiocSingular $runPCA$ function. "flash,"[25,79] a robust and flexible method for variational Bayesian matrix factorization, was implemented using the "flashier"[79] package, using backfitting to update factors after initial fitting with all other settings as default.   GLEANR without covariance adjustment is equivalent to standard GLEANR with $\boldsymbol{W}_c = \boldsymbol{I_M}$, the $M \times M$ identity matrix. To estimate $\widehat{\boldsymbol{C}}$, we used the known simulation sample overlap and empirical phenotypic correlation, with diagonal elements given by $\hat{C}_{i,j} = \frac{N_o\hat{\rho}}{\sqrt{N_1 N_2}}$[3,24] and standard error estimates $\widehat{s_C} = \frac{N_o \hat{s}_\rho}{\sqrt{N_1 N_2}}$, with $\hat{\rho}$ and $\hat{s}_\rho$ from the R $stats::cor.test$ utility. Performance was assessed by counting the number of non-zero factors nominated by each model selection procedure.

MF method	Model selection method (to select K)	Input
SVD	Factors above the "elbow" of the scree plot	Z-scores ($\mathbf{Z} = \mathbf{B} \odot \mathbf{W_S}$)
SVD	Factors eigenvalues greater than the average eigenvalue: $\{k_\lambda: \lambda_k > \frac{1}{M}\sum_{k=1}^{M}\lambda_k\}$	Z-scores ($\mathbf{Z} = \mathbf{B} \odot \mathbf{W_S}$)
SVD-adj	Factors above the "elbow" of the scree plot	"Decorrelated" Z-scores $\mathbf{Z_d} = (\mathbf{W_c}(\mathbf{B} \odot \mathbf{W_S})^{\mathrm{T}})^{\mathrm{T}}$
SVD-adj	Factors eigenvalues greater than the average eigenvalue: $\{k_\lambda: \lambda_k > \frac{1}{M}\sum_{k=1}^{M}\lambda_k\}$	"Decorrelated" Z-scores $\mathbf{Z_d} = (\mathbf{W_c}(\mathbf{B} \odot \mathbf{W_S})^{\mathrm{T}})^{\mathrm{T}}$
flash	"Greedy" addition of non-zero factors, followed by "backfitting" procedure	$\mathbf{B}, \mathbf{S}$
GLEANR (no adjustment)	Grid search for K_{init} followed by regularized model fitting	$\mathbf{B}, \mathbf{W_S}$
GLEANR	Grid search for K_{init} followed by regularized model fitting	$\mathbf{B}, \mathbf{W_S}, \mathbf{W_c}$

Table 1: Summary of methods evaluated in direct GWAS-based simulations and their corresponding inputs. Matrix names ($\mathbf{B}, S, W_S, W_c$) match those in the main text.

2) Simulation of latent genetic basis:
To evaluate latent space estimation in the presence of confounding covariance, we simulated summary statistics from genetic factors and added noise reflecting sample sharing across studies. We simulated effect sizes ($\mathbf{B}$) for 100 SNPs across 10 traits as $\mathbf{B} = \boldsymbol{UV^T a}$, with prespecified $\boldsymbol{U}, \boldsymbol{V}$ and $\boldsymbol{a}$ a vector to scale causal effect sizes to match variance of previously reported continuous trait phenotypes[39]. We generated 3 each of $\mathbf{U}$ and $\mathbf{V}$ with K=5 for a total of nine distinct $\mathbf{B}$. Elements of $\boldsymbol{U}$ were sampled (at k factors, n SNPs) from

$$u_{n,k} \sim \theta_k^{(U)}\delta(u_{n,k}) + \left(1 - \theta_k^{(U)}\right)\mathcal{N}(0, \frac{0.05}{100K}),$$
$$\theta_k^{(U)} \sim Bernoulli\left(p_k^{(U)}\right),$$
$$p_k^{(U)} \sim Unif(0.1, 0.7)$$

Elements of $\boldsymbol{V}$ (at k factors, m traits) were drawn from:
$$v_{m,k} \sim \mathbf{1}_{k \neq 1}(k) \times \theta_k^{(V)}\delta(v_{m,k}) + \left(1 - \theta_k^{(V)}\right)\mathcal{N}(0,1)$$
$$\theta_k^{(V)} \sim Bernoulli\left(p_k^{(V)}\right)$$
$$p_k^{(V)} \sim Unif(0.5, 75)$$

All columns of $\boldsymbol{U}$ and $\boldsymbol{V}$ were required to have at least 1 non-zero element.
We then added noise $\mathbf{E}$, where a single row of $\mathbf{E}, \boldsymbol{\epsilon_i} \sim MVN\left(0, \mathrm{diag}(\boldsymbol{s_n}) \times \boldsymbol{C} \times \mathrm{diag}(\boldsymbol{s_n})\right)$.
Elements of the vector $\boldsymbol{s_n}$ are given by $s_{n,m} = \frac{1}{\sqrt{(N_m 2 f_n(1-f_n))}}$, where f_n (minor allele frequency of SNP n) was sampled from the European UKBB variants used in our analysis of 137 diverse phenotypes. N_m (sample size of study m) was simulated at levels of 5,10,50,100 and 200 thousand samples for each study, and in simulation with a "mixed" number of samples (10-200 thousand).
$\boldsymbol{C}$ was simulated with elements

$$c_{i,j} = \begin{cases} i = j: & 1 \\ i \neq j & \frac{n_o \rho_{i,j}}{\sqrt{N_i N_j}} \end{cases}$$

where N_i corresponds to the total sample size of study i. n_o and $\rho_{i,j}$ each correspond to one of three different cohort overlap structures tested. In the case of no overlap, $\rho_{i,j} = n_o = 0$. For block cohort overlap structures, we let $n_o = 0.9 \times \min(N_i, N_j)$ and $\rho_{i,j} = 1$ for all diagonal

entries. In the "single block" setting, $\rho_{i,j} \sim Unif(0.75,0.99)$ for traits 4-8, and 0 for all others. In the "double block" setting $\rho_{i,j} \sim Unif(0.4,0.7)$ for traits 1-5 or 6-10, 0 otherwise. These resulted in a total of 162 different evaluation settings, each of which had E resampled 10 times. Each method was specified to yield K=5 (or a maximum of K=5 in the case of GLEANR and flash). In addition to the methods above, we included flash with the Kronecker variance setting, which models a block matrix covariance structure, and the GWAS-specific variational method FactorGo[8].

Matching factors and evaluating factorization performance:
To determine how closely the $\widehat{U}, \widehat{V}$ predicted by factorization methods matched simulated matrices, we evaluate squared correlation (R^2) across all predicted factors. Matrix factorization solutions are statistically unidentifiable, so different solutions may reflect equivalent matrix structure but differ by an arbitrary matrix rotation and scaling. To address this, we find the factor order (I_o) and sign vector (μ) that maximizes the global correlation across both matrices by optimizing the following:

$$\max_{I_o, S}(\text{diag}[(\text{cor}(V, \mu I_o \widehat{V}))] + \text{diag}[\text{cor}(U_1, \mu I_o \widehat{U})])^2 \qquad (7)$$

Where I_o is an orthogonal rotation matrix with exactly one non-zero element per column of value 1, and μ is a vector of signs with elements $\mu_i \in [1, -1]$. When the dimension of predicted matrices was fewer than the true matrices, columns were padded with 0s. We then calculated R^2 across all factors, each scaled to have unit norm.

FinnGen and UKBB Cross-cohort analysis:
Study selection
We compared GLEANR factors across 42 phenotypes from GWAS evaluated in the FinnGen[28] and UKBB[14] cohorts (Table S1). The FinnGen study is a large-scale collaborative genomics initiative that has analyzed over 500,000 Finnish biobank sample and has published summary statistics for over 2000 phenotype endpoints. Similarly, the UKBB is a large-scale biomedical dabase with genetic and health information for over 500,000 UK volunteer participants aged 40-69 years. Summary statistics for 41 disease phenotypes in the UKBB cohort were accessed from the PheWeb[29] server, an online resource of GWAS in over 1400 binary phenotypes from white UKBB participants. For the remaining continuous phenotype, summary statistics came from the Neale Lab GWAS portal[23] round 2 results (Web Resources) calculated using Hail 0.2[80]. FinnGen and PheWeb GWAS were generated using SAIGE[81], and all studies were controlled for sex, age or birth year, and multiple genotype PCs. To select traits for evaluation, we first manually matched phenotypes available in both cohorts. These were filtered to GWAS with at least 25 minor alleles among controls for SNPs with MAF > 0.01, an effective sample size > 5000, and 1 reported GWAS hit in the FinnGen GWAS manifest. Subsetting to HapMap3[27] SNPs, we performed XT-LDSR[3] to estimate the genetic correlation between cohorts on the 77 remaining traits, keeping only those with $r_g > 0.8$ at a Bonferroni-adjusted $p < 0.01$. From the remaining 42 studies, we selected SNPs for evaluation by filtering out ambiguous and multi-allelic variants and those in the HLA region and keeping only those with $p < \times \ 10^{-5}$ in at least 1 GWAS. 14750 SNPs passed these filters in both the FinnGen and UKBB GWAS, which we then pruned using plink2[82] (*–indep-pairwise 250kb 0.2)* and a European 1000 Genomes[83] LD reference panel. The remaining 1663 SNPs were input to GLEANR analysis, evaluated at $K_{init} = 41$. We estimated cohort-specific sample sharing from XT-LDSR,[3] scaling reported $g_{cov-int}$ statistics to adjust for the selection of heritable traits.

To assess the concordance of factors between cohorts, we calculate R^2 across all unit-scaled factors, matching factors as described above (Equation 7). We compared reconstructed effect sizes, $\widehat{B}_{FG} = \widehat{U_{FG}}\widehat{V}_{FG}^T$ and $\widehat{B}_{UK} = \widehat{U_{UK}}\widehat{V}_{UK}^T$, via relative root mean squared error (RRMSE):

$$RRMSE = \sqrt{\frac{\left\|\widehat{\mathbf{B}}_{FG} - \widehat{\mathbf{B}}_{UK}\right\|_F^2}{\left\|\widehat{\mathbf{B}}_{FG}\right\|_F^2}} \tag{8}$$

UKBB analysis across 137 diverse GWAS:
Study selection:
      Traits were selected from 7228 phenotypes with GWAS published by the Pan-UKBB[26] project. Summary statistics were generated in a standardized pipeline across 6 continental ancestries (including up to 420531 European individuals; see Web Resources) using the SAIGE[81] software. We considered for inclusion those traits passing Pan-UKBB recommended filters (valid LDSR[84] $h^2 \in (0,1)$ with a significant Z-score, normal λ_{gc} across top 3 powered ancestries, an LDSR inflation ratio < 0.3), studies with effective sample size > 5000, and LDSR[84]-estimated $h^2 > 0.05$. We limited GWAS of continuous phenotypes to inverse rank normal transformed values and removed phenotypes in categories not of biological interest (including activities, environment, nutrition, social interaction, operation codes, parental age, prescriptions, and those related to hospital admission or treatment specialties). Category assignments were extended from previously published "domain" assignments[5] integrated with UKBB coding information. Of the remaining traits, we added studies to our set in order of decreasing heritability, requiring that each had a pairwise $r_g < 0.7$ (estimated by XT-LDSR[3]) with all other studies in our set. This yielded 137 traits in total (Table S10).
SNP selection:
      From ~29.0 million variants evaluated by the Pan-UKBB[26], we considered biallelic SNPs designated as "high quality" (e.g. PASS variants in gnomAD and consistent frequencies across gnomAD ancestries), with INFO > 0.9, MAF > 0.01 in Europeans. From these 7.9 M variants, we selected HapMap3[27] SNPs not in the HLA locus with GWAS $p < 1 \times 10^{-5}$ in at least 1 study, then LD-clumped these variants with plink,[85] using a 1000 Genomes[83] European genotype reference. During clumping, SNPs were prioritized according to the number of studies in which they achieved our nominal significant threshold, favoring the inclusion of more pleiotropic SNPs and yielding a total of 41260 SNPs.

UKBB GLEANR analysis
      We performed GLEANR on the selected GWAS summary statistics using an adaptive grid search at eight settings of K_{init} to select $K_{init} = 134$ as yielding the minimizing L_{BIC} score. GLEANR model fitting converged in 34 iterations, following 7 iterations of model selection. $\widehat{C}$ was estimated using XT-LDSC and scaled to adjust for the selection of heritable traits.
GLEANR factor evaluation
Factor tissue-marker enrichment: To test for heritability enrichment among tissue-specific chromatin markers in GLEANR factors, we performed stratified-LDSR[31] using 489 tissue-specific chromatin markers,[32] LD scores, and weights provided by the authors (see Web resources). GLEANR loading scores (columns of $\widehat{U}$) were scaled into Z-scores for input into S-LDSR, where SNP n of factor k has value

$$z_{n,k} = \left(\sum_m^M \frac{V_{m,k}^2 W_{S_{n,m}}}{\sum_m^M V_{m,k}^2} \right) U_{n,k} \tag{9}$$

Factor sample sizes N_K were given as the weighted average of the sample sizes of traits in a given factor $(\boldsymbol{v_k})$, with $N_k = \sum_m^M \frac{V_{m,k}^2 N_M}{\sum_m^M V_{m,k}^2}$. Because only 41260 SNPs were available for

regression, we increased the number of LDSR jackknife blocks to 5000 (default 200) to improve standard error estimation.

To correct for multiple hypothesis testing, we perform Benjamini-Hochberg adjustment across all enrichment tests per factor, and report BH-adjusted p-values as "q-values"[86]. To evaluate which tissue categories yielded the most prominent enrichment across all 489 tested chromatin markers per factor, we needed to further adjust for the unequal number of markers per tissue category. To do this, we performed Bonferroni adjustment of p-values for all factor tests withing a given tissue category, retained only the enrichment with the smallest p-value per category, and then performed BH-adjustment across these adjusted p-values.

Factor genetic architecture: We estimated the polygenicity and the selection signature parameter[30,42] for each factor. To estimate the polygenicity of factor k, we leveraged the kurtosis (scaled 4th moment) of SNP effect sizes[37], reporting $p_k = \frac{3}{\kappa(\boldsymbol{u_k})}, \in (0,1)$, where $\kappa(\boldsymbol{u_k})$ is the empirical kurtosis of elements of $\boldsymbol{u_k}$. Values close to 1 reflect larger tails[87] on the effect size distribution and indicate higher polygenicity.

To estimate selection coefficient S_k, we model non-zero factor effect sizes as $U_{i,k} \sim N(0, (2f_i(1-f_i))^{S_k}\sigma_k^2)$, where f_i is the MAF at SNP i. Using R's $optim$ function, we find the S_k and σ_k^2 maximizing the empirical log-likelihood of non-zero effects in factor $\boldsymbol{u_k}$.

Top SNP selection: We define "top factor SNPs" as those in the top tertile above the "elbow" of factor SNP effect magnitudes. For some factor loading $\boldsymbol{u_k}$, we ranked SNPs according to their magnitude in decreasing order, and then selected the SNP with effect size magnitude (i.e. $|U_{n,k}|$) at the graphical "elbow" of this distribution (e.g. achieving maximal distance from a line between the largest (rank 1) and smallest (rank N) SNP). We designate SNPs in the top third of effect size magnitudes above this elbow as top factor SNPs. The number of SNPs per factor chosen in this manner correlate strongly with estimates of pleiotropy ($\rho = 0.85$; Fig. S11).

SNP-gene mapping and gene set enrichment analysis: All SNPs were mapped to a putative candidate gene using the latest variant-to-gene mapping from the Open Targets[43,44] platform (download version 2022-10-11). If multiple genes were assigned equal scores by Open Targets or no gene assignment was available, we selected the gene with the closest protein-coding transcript start site to the SNP.

To perform gene set enrichment on factors, we designated the set of distinct genes mapped to by top factor SNPs as the set of top factor genes. All distinct genes mapped to by a SNP in the analysis constituted our background set, yielding 10,512 unique genes (from 41259 candidate SNPs) and 414.8 unique top factor genes per factor, on average.

Enrichment testing in canonical gene set libraries (including GO Biological Process, GO Cellular Component, GO Molecular Function, DisGeNET, KEGG pathways, GWAS Catalog, and Panglao DB) was performed using the *enrichr*[66,88,89] API. Reported q-values represent p-values adjusted using the Benjamini-Hochberg[86] procedure for the number of tests per gene set library.

For gene set enrichment tests on sets related to inherited platelet disorders, genes were collected from those in previously reported studies [55,64] and grouped by the stage of platelet development which they disrupt in disease. Top factor genes were tested for enrichment against background genes using a Fisher's exact test. To account for the small gene set size and the non-uniform inclusion of genes in our analysis (many genes are mapped to by multiple SNPs), we generated an empirical null distribution of enrichment statistics. For factor k with N_k TFSs, we randomly subsampled N_k SNPs and performed a Fisher's exact test on the distinct set of

corresponding genes. This procedure was repeated 50,000 times to generate an empirical p-value where $p = \frac{\sum_{i}^{50000} I[OR_i \geq OR_k]}{50000}$ (the number of tests with an odds ratio as or more extreme than the odds ratio from the true top factor genes), which we corrected using the Benjamini-Hochberg procedure[86] for the number of tests performed across factors and gene sets. A similar permutation procedure with 10,000 samplings was used to validate top PanglaoDB[67] cell-type enrichments reported in Figure 4.

Discussion

In this work we present GLEANR for the identification of genetic factors through sparse matrix factorization of GWAS summary statistics. GLEANR offers two key advantages over existing approaches. First, it enables hypothesis-free exploration over 100s of traits robust to the effects of shared samples among the input GWAS. Second, GLEANR encourages sparsity in the estimated factors specific to the target summary statistics, yielding latent factors that are more readily interpretable than those from dense MF approaches.

As revealed through extensive simulations, we observed that existing matrix factorization methods are susceptible to reporting spurious and confounded latent genetic factors. These spurious factors occur due to non-biological influences, particularly the effects of sample sharing among the input GWAS, which increases the covariance in GWAS effect sizes simply due to shared residual noise. These spurious factors capture non-genetic signal, complicate interpretability, and reduce reproducibility across populations and biobanks. As shown theoretically and empirically with published GWAS, GLEANR is more robust to identifying these spurious factors than existing methods. These findings emphasize that appropriately accounting for sample sharing, even if it only influences a subset of target studies, improves the detection of replicable components and is critical to preserve the interpretability of latent genetic factors.

Across 137 diverse GWAS in the UKBB, we estimated one ubiquitous and 57 sparse factors that reflect biological pathways underlying GWAS effects. These genetic factors are a resource for distinguishing pathways shared by multiple traits and those which are trait-specific (singleton factors). Enrichments among top factor genes and in tissue-specific chromatin markers can support hypothesis generation through exploration of previously unobserved relationships among traits, such as the enrichment we observe in brain-specific chromatin markers corresponding to Factor 1, which was loaded on all traits.

We observe that GLEANR factors decompose trait effects into components with distinct genetic architectures. Variation in factor polygenicity scores highlight that the effective number of variants contributing to factors differ. This also influences factor interpretation and informs how many factor SNPs to consider when performing factor enrichment tests. Components with strong MAF-effect size relationships, indicative of negative selection, are trait-specific and have lower polygenicity, a relationship observed in forward evolutionary simulations that model parameters of genetic architecture[30]. These results suggest that global evaluation of trait architecture may obscure the contributing sub-architecture of distinct genetic components.

Of the 58 factors nominated by GLEANR, we identify three factors (in addition to a ubiquitous factor) loaded on platelet phenotypes. Gene set enrichment tests provide evidence that these factors stratify variant contributions to inherited platelet disorders at distinct stages of platelet formation including hematopoiesis, megakaryopoiesis, and platelet formation. Published studies of platelet phenotypes report corresponding enrichments in chromatin accessibility profiles for these platelet-related progenitor cell types[50,90]. By partitioning platelet traits with GLEANR, we were able to stratify these cell-type effects into factors reflecting distinct stages of platelet development. This work demonstrates the ability of GLEANR factors to distinguish trait-relevant cell-type processes shared across traits.

GLEANR has a few limitations. For example, enforcing sparsity in genetic factors may remove small effect or low-confidence signals, limiting its applicability to very rare traits, studies

with small sample sizes, or those with a paucity of associations. In addition, our analysis does not account for every source of confounding between studies, such as assortative mating. However, if estimates of the impact of these confounders are available, GLEANR could be extended to adjust for those effects by integrating those estimates into $\widehat{C}$. Finally, estimates of polygenicity and selection signatures from GLEANR factors are not numerically comparable to those from full GWAS, as they are based on the subset of variants chosen for inclusion in the analysis due to trait-association. Interpretation its thus limited to comparison with other factors in the same analysis.

In conclusion, GLEANR is a powerful tool for evaluation of pleiotropic components underlying genetic correlations. It can be applied to diverse GWAS and is available as a free software package along with supporting tools for interpretation of latent factors. GLEANR enables characterization of trait genetic architecture and biology into distinct components that clarify the impact of pleiotropy in the human phenome.

Web resources

GLEANR package and pipeline code: https://github.com/aomdahl/gleanr
Pan-UKBB: https://pan.ukbb.broadinstitute.org
Finngen: https://finngen.gitbook.io/documentation/r10/releases
Genetic Correlation Portal: https://ukbb-rg.hail.is
PheWeb summary statistics: https://pheweb.org
Neale Lab GWAS: http://www.nealelab.is/uk-biobank/
Tissue-specific S-LDSC Resources: https://github.com/bulik/ldsc/wiki/Cell-type-specific-analyses

Declaration of interests

A.B. is a co-founder and equity holder of CellCipher, Inc, a stockholder in Alphabet, Inc, and has consulted for Third Rock Ventures. J.S.W. was a consultant to Spiral Genetics.
The other authors declare no competing interests.

Acknowledgements:

The authors would like to thank Battle lab members for helpful discussion throughout the course of this work. We especially thank Seraj Grimes and April Kim for manuscript feedback and code review, and Ben Strober for contributions to early ideation of project analyses. Alexis Battle is supported by R35GM139580. M.A. was supported by NHLBI grant K08HL166690.

Author contributions:

Ashton Omdahl: Conceptualization, Software, Methodology, Formal analysis, Data Curation, Writing- original draft, Writing – review & editing
Joshua Weinstock: Methodology, Writing – review & editing, Supervision
Rebecca Keener: Writing – review & editing, Project Administration
Surya Chhetri: Methodology
Marios Arvanitis: Conceptualization, Supervision
Alexis Battle: Conceptualization, Resources, Writing – review & editing, Supervision

References

1. Uffelmann, E. *et al.* Genome-wide association studies. *Nature Reviews Methods Primers 2021 1:1* 1, 1–21 (2021).
2. Sollis, E. *et al.* The NHGRI-EBI GWAS Catalog: knowledgebase and deposition resource. *Nucleic Acids Res* 51, D977–D985 (2023).
3. Bulik-Sullivan, B. *et al.* An atlas of genetic correlations across human diseases and traits. *Nature Publishing Group* 47, 1–3 (2015).
4. Qi, G. *et al.* Genome-Wide Large-Scale Multi-Trait Analysis Characterizes Global Patterns of Pleiotropy and Unique Trait-Specific Variants. doi:10.1101/2022.06.03.494686.
5. Watanabe, K. *et al.* A global overview of pleiotropy and genetic architecture in complex traits. *Nat Genet* 51, 1339–1348 (2019).
6. Jordan, D. M., Verbanck, M. & Do, R. HOPS: A quantitative score reveals pervasive horizontal pleiotropy in human genetic variation is driven by extreme polygenicity of human traits and diseases. *Genome Biol* 20, (2019).
7. Udler, M. S. *et al.* Type 2 diabetes genetic loci informed by multi-trait associations point to disease mechanisms and subtypes: A soft clustering analysis. *PLoS Med* 15, 1–23 (2018).
8. Zhang, Z. *et al.* A scalable approach to characterize pleiotropy across thousands of human diseases and complex traits using GWAS summary statistics. *Am J Hum Genet* 110, 1863–1874 (2023).
9. Grotzinger, A. D. *et al.* Genomic structural equation modelling provides insights into the multivariate genetic architecture of complex traits. *Nature Human Behaviour 2019 3:5* 3, 513–525 (2019).
10. Tanigawa, Y. *et al.* Components of genetic associations across 2,138 phenotypes in the UK Biobank highlight adipocyte biology. *Nat Commun* 10, (2019).
11. Ballard, J. L. & O'Connor, L. J. Shared components of heritability across genetically correlated traits. *Am J Hum Genet* 109, 989–1006 (2022).
12. Chang, D. & Keinan, A. Principal Component Analysis Characterizes Shared Pathogenetics from Genome-Wide Association Studies. *PLoS Comput Biol* 10, (2014).
13. Burren, O. S. *et al.* Genetic feature engineering enables characterisation of shared risk factors in immune-mediated diseases. *Genome Med* 12, 1–17 (2020).
14. Sudlow, C. *et al.* UK Biobank: An Open Access Resource for Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age. *PLoS Med* 12, e1001779 (2015).
15. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* 562, 203 (2018).
16. Solovieff, N., Cotsapas, C., Lee, P. H., Purcell, S. M. & Smoller, J. W. Pleiotropy in complex traits: challenges and strategies. *Nature Reviews Genetics 2013 14:7* 14, 483–495 (2013).
17. Gratten, J. & Visscher, P. M. Genetic pleiotropy in complex traits and diseases: Implications for genomic medicine. *Genome Med* 8, 1–3 (2016).
18. Friedman, J., Hastie, T. & Tibshirani, R. *Regularization Paths for Generalized Linear Models via Coordinate Descent*. *JSS Journal of Statistical Software* vol. 33 http://www.jstatsoft.org/ (2010).
19. Tibshirani, R. *Regression Shrinkage and Selection via the Lasso*. *Source: Journal of the Royal Statistical Society. Series B (Methodological)* vol. 58 (1996).
20. Lee, M., Shen, H., Huang, J. Z. & Marron, J. S. Biclustering via Sparse Singular Value Decomposition. *Biometrics* 66, 1087–1095 (2010).

21.	Zou, H., Hastie, T. & Tibshirani, R. On the 'degrees of freedom' of the lasso. *Ann Stat* 35, 2173–2192 (2007).
22.	Zhang, Z. *et al.* A scalable variational approach to characterize pleiotropic components across thousands of human diseases and complex traits using GWAS summary statistics. *medRxiv* (2023) doi:10.1101/2023.03.27.23287801.
23.	UK Biobank — Neale lab. http://www.nealelab.is/uk-biobank/.
24.	LeBlanc, M. *et al.* A correction for sample overlap in genome-wide association studies in a polygenic pleiotropy-informed framework. *BMC Genomics* 19, (2018).
25.	Wang, W. & Stephens, M. *Empirical Bayes Matrix Factorization. Journal of Machine Learning Research* vol. 22 http://jmlr.org/papers/v22/20-589.html. (2021).
26.	Karczewski, K. J. *et al.* Pan-UK Biobank GWAS improves discovery, analysis of genetic architecture, and resolution into ancestry-enriched effects. doi:10.1101/2024.03.13.24303864.
27.	Auton, A. *et al.* A global reference for human genetic variation. *Nature 2015 526:7571* 526, 68–74 (2015).
28.	Kurki, M. I. *et al.* FinnGen provides genetic insights from a well-phenotyped isolated population. *Nature* 613, 508–518 (2023).
29.	Gagliano Taliun, S. A. *et al.* Exploring and visualizing large-scale genetic associations by using PheWeb. *Nature Genetics* vol. 52 550–552 Preprint at https://doi.org/10.1038/s41588-020-0622-5 (2020).
30.	Zeng, J. *et al.* Widespread signatures of natural selection across human complex traits and functional genomic categories. *Nat Commun* 12, 1164 (2021).
31.	Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nat Genet* 47, 1228–1235 (2015).
32.	Finucane, H. K. *et al.* Heritability enrichment of specifically expressed genes identifies disease-relevant tissues and cell types. *Nat Genet* 50, 621–629 (2018).
33.	Locke, A. E. *et al.* Genetic studies of body mass index yield new insights for obesity biology. *Nature* 518, 197 (2015).
34.	Leyden, G. M. *et al.* Harnessing tissue-specific genetic variation to dissect putative causal pathways between body mass index and cardiometabolic phenotypes. *Am J Hum Genet* 109, 240–252 (2022).
35.	Gagnon, E. *et al.* Genetic control of body weight by the human brain proteome. *iScience* 26, 106376 (2023).
36.	Piñero, J. *et al.* DisGeNET: a comprehensive platform integrating information on human disease-associated genes and variants. *Nucleic Acids Res* 45, D833 (2017).
37.	O'connor, L. J. *et al.* Extreme Polygenicity of Complex Traits Is Explained by Negative Selection. *The American Journal of Human Genetics* 105, 456–476 (2019).
38.	Shi, H., Kichaev, G. & Pasaniuc, B. Contrasting the Genetic Architecture of 30 Complex Traits from Summary Association Data. *The American Journal of Human Genetics* 99, 139–153 (2016).
39.	Zhang, Y., Qi, G., Park, J. H. & Chatterjee, N. Estimation of complex effect-size distributions using summary-level statistics from genome-wide association studies across 32 complex traits. *Nat Genet* 50, 1318–1326 (2018).
40.	O'Connor, L. J. The distribution of common-variant effect sizes. *Nature Genetics 2021 53:8* 53, 1243–1249 (2021).
41.	Park, J. H. *et al.* Distribution of allele frequencies and effect sizes and their interrelationships for common genetic susceptibility variants. *Proc Natl Acad Sci U S A* 108, 18026–18031 (2011).
42.	Schoech, A. P. *et al.* Quantification of frequency-dependent genetic architectures in 25 UK Biobank traits reveals action of negative selection. *Nat Commun* 10, (2019).

43. Ghoussaini, M. *et al.* Open Targets Genetics: systematic identification of trait-associated genes using large-scale genetics and functional genomics. *Nucleic Acids Res* 49, D1311–D1320 (2021).

44. Mountjoy, E. *et al.* An open approach to systematically prioritize causal variants and genes at all published human GWAS trait-associated loci. *Nature Genetics 2021 53:11* 53, 1527–1533 (2021).

45. Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res* 45, D353 (2016).

46. Meech, R. *et al.* The UDP-Glycosyltransferase (UGT) superfamily: New members, new functions, and novel paradigms. *Physiol Rev* 99, 1153–1222 (2019).

47. Voight, B. F., Kudaravalli, S., Wen, X. & Pritchard, J. K. A Map of Recent Positive Selection in the Human Genome. *PLoS Biol* 4, e72 (2006).

48. Wilde, S. *et al.* Direct evidence for positive selection of skin, hair, and eye pigmentation in Europeans during the last 5,000 y. *Proc Natl Acad Sci U S A* 111, 4832–4837 (2014).

49. Chen, M. H. *et al.* Trans-ethnic and Ancestry-Specific Blood-Cell Genetics in 746,667 Individuals from 5 Global Populations. *Cell* 182, 1198-1213.e14 (2020).

50. Vuckovic, D. *et al.* The Polygenic and Monogenic Basis of Blood Traits and Diseases. *Cell* 182, 1214-1231.e11 (2020).

51. Võsa, U. *et al.* Large-scale cis- and trans-eQTL analyses identify thousands of genetic loci and polygenic scores that regulate blood gene expression. *Nat Genet* 53, 1300–1310 (2021).

52. Aguet, F. *et al.* Genetic effects on gene expression across human tissues. *Nature 2017 550:7675* 550, 204–213 (2017).

53. Pleines, I. *et al.* Mutations in tropomyosin 4 underlie a rare form of human macrothrombocytopenia. *J Clin Invest* 127, 814–829 (2017).

54. Marín-Quílez, A. *et al.* A novel nonsense variant in TPM4 caused dominant macrothrombocytopenia, mild bleeding tendency and disrupted cytoskeleton remodeling. *Journal of Thrombosis and Haemostasis* 20, 1248 (2022).

55. Collins, J., Astle, W. J., Megy, K., Mumford, A. D. & Vuckovic, D. Advances in understanding the pathogenesis of hereditary macrothrombocytopenia. *Br J Haematol* 195, 25–45 (2021).

56. Giles, C. The platelet count and mean platelet volume. *Br J Haematol* 48, 31–37 (1981).

57. Rafieian-Kopaie, M. & Nasri, H. Platelet counts and mean platelet volume in association with serum magnesium in maintenance hemodialysis patients. *J Renal Inj Prev* 1, 17 (2012).

58. Ghosh, K., Bhattacharya, M., Chowdhury, R., Mishra, K. & Ghosh, M. Inherited Macrothrombocytopenia: Correlating Morphology, Epidemiology, Molecular Pathology and Clinical Features. *Indian Journal of Hematology & Blood Transfusion* 34, 387 (2018).

59. Strober, B. J. *et al.* Dynamic genetic regulation of gene expression during cellular differentiation. *Science (1979)* (2019) doi:10.1126/science.aaw0040.

60. Cuomo, A. S. E. *et al.* Single-cell RNA-sequencing of differentiating iPS cells reveals dynamic genetic effects on gene expression. *Nat Commun* 11, (2020).

61. Elorbany, R. *et al.* Single-cell sequencing reveals lineage-specific dynamic genetic regulation of gene expression during human cardiomyocyte differentiation. *PLoS Genet* 18, (2022).

62. Aguet, F. *et al.* Genetic effects on gene expression across human tissues. *Nature* 550, 204–213 (2017).

63. Yazar, S. *et al.* Single-cell eQTL mapping identifies cell type-specific genetic control of autoimmune disease. *Science (1979)* 376, (2022).

64. Lentaigne, C. *et al.* Inherited platelet disorders: toward DNA-based diagnosis. *Blood* 127, 2814–2823 (2016).
65. Geddis, A. E. Megakaryopoiesis. *Semin Hematol* 47, 212 (2010).
66. Xie, Z. *et al.* Gene Set Knowledge Discovery with Enrichr. *Curr Protoc* 1, e90 (2021).
67. Franzén, O., Gan, L. M. & Björkegren, J. L. M. PanglaoDB: a web server for exploration of mouse and human single-cell RNA sequencing data. *Database* 2019, 46 (2019).
68. Sengupta, S. Improved Analysis of Large Genetic Association Studies Using Summary Statistics. (2018).
69. Urbut, S. M., Wang, G., Carbonetto, P. & Stephens, M. Flexible statistical methods for estimating and testing effects in genomic studies with multiple conditions. *Nature Genetics 2018 51:1* 51, 187–195 (2018).
70. Ledoit, O. & Wolf, M. The Power of (Non-)Linear Shrinking: A Review and Guide to Covariance Matrix Estimation. doi:10.1093/jjfinec/nbaa007.
71. Schäfer, J., Strimmer, K., Schmidt-Heck, W., Guthke, R. & Bayer, K. *Statistical Applications in Genetics and Molecular Biology A Shrinkage Approach to Large-Scale Covariance Matrix Estimation and Implications for Functional Genomics A Shrinkage Approach to Large-Scale Covariance Matrix Estimation and Implications for Functional Genomics *. http://www.bepress.com/sagmb.
72. Sill, M., Kaiser, S., Benner, A. & Kopp-Schneider, A. Robust biclustering by sparse singular value decomposition incorporating stability selection. 27, 2089–2097 (2011).
73. Zou, H., Hastie, T. & Tibshirani, R. Sparse Principal Component Analysis. *Journal of Computational and Graphical Statistics* 15, 265–286 (2006).
74. He, Y. *et al.* Sn-spMF: Matrix factorization informs tissue-specific genetic regulation of gene expression. *Genome Biol* 21, 1–25 (2020).
75. Schwarz, G. Estimating the Dimension of a Model. *https://doi.org/10.1214/aos/1176344136* 6, 461–464 (1978).
76. Neath, A. A. & Cavanaugh, J. E. The Bayesian information criterion: background, derivation, and applications. *Wiley Interdiscip Rev Comput Stat* 4, 199–203 (2012).
77. Chen, J. *Extended Bayesian Information Criteria for Model Selection with Large Model Spaces*.
78. Manichaikul, A., Moon, J. Y., Sen, Ś., Yandell, B. S. & Broman, K. W. A Model Selection Approach for the Identification of Quantitative Trait Loci in Experimental Crosses, Allowing Epistasis. *Genetics* 181, 1077–1086 (2009).
79. Willwerscheid, J., Carbonetto, P. & Stephens, M. ebnm: An R Package for Solving the Empirical Bayes Normal Means Problem Using a Variety of Prior Families. (2021).
80. Hail Team. Hail 0.2.
81. Zhou, W. *et al.* Efficiently controlling for case-control imbalance and sample relatedness in large-scale genetic association studies. *Nat Genet* 50, 1335–1341 (2018).
82. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* 4, (2015).
83. Fairley, S., Lowy-Gallego, E., Perry, E. & Flicek, P. The International Genome Sample Resource (IGSR) collection of open human genomic variation resources. *Nucleic Acids Res* 48, D941–D947 (2020).
84. Bulik-Sullivan, B. *et al.* LD score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet* 47, 291–295 (2015).
85. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *Am J Hum Genet* 81, 559 (2007).
86. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* 57, 289–300 (1995).
87. Westfall, P. H. Kurtosis as Peakedness, 1905 – 2014. R.I.P. *Am Stat* 68, 191 (2014).

88.   Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res* 44, W90–W97 (2016).

89.   Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics* 14, (2013).

90.   Ulirsch, J. C. *et al.* Interrogation of human hematopoiesis at single-cell and single-variant resolution. *Nat Genet* 51, 683 (2019).