MicroRNA-29 acutely regulates Memory Stability, Expression of Synaptic Genes, and DNA Methylation in the Mouse Adult Hippocampus

Aurelia Viglione[1,2], Chiara Giannuzzi[1], Elena Putignano[3], Raffaele Mario Mazziotti[3,4,5], Sara Bagnoli[1], Paola Tognini[6,7], Alessandro Cellerino[1,8] and Tommaso Pizzorusso[1,3*]

[1] BIO@SNS lab, Scuola Normale Superiore, Pisa, Italy;

[2] Center for Behavioral Sciences and Mental Health, Istituto Superiore di Sanità, Rome, Italy;

[3] Institute of Neuroscience, National Research Council, Pisa, Italy;

[4] Department of Developmental Neuroscience, IRCCS Stella Maris Foundation, Pisa, Italy;

[5] Department of Neuroscience, Psychology, Pharmacology, and Child Health, University of Florence, Florence, Italy;

[6]Health Science Interdisciplinary Center, Scuola Superiore Sant'Anna, Pisa, Italy

[7]Cardiology division, Fondazione Toscana Gabriele Monasterio, Pisa, Italy

[8] Leibniz Institute on Aging – Fritz Lipmann Institute (FLI) Jena Germany

* Corresponding author: Tommaso Pizzorusso, Istituto Neuroscienze CNR and BIO@SNS lab, Scuola Normale Superiore via G. Moruzzi 1, 56124 Pisa, Italy, tel +390503153167, Fax +390503153220, e-mail: tommaso.pizzorusso@sns.it

Summary

MicroRNAs are key regulators of brain gene expression, with miR-29 family notably upregulated from development to adulthood and in aging, and showing links to cognitive decline. However, the extent to which miR-29 levels influence learning and memory processes, and its molecular mediators, remains to be determined. Here, we down- and up-regulated miR-29 levels in the dorsal hippocampus of adult mice to reveal miR-29 role in memory. Inhibition of miR-29 enhanced trace fear memory stability, increased Dnmt3a levels, and promoted DNA methylation in a DNMT3a-dependent manner. In contrast, increasing miR-29 impaired memory performances and decreased Dnmt3a levels, suggesting a destabilization of memory processes. Proteomic and transcriptomic analysis demonstrated that miR-29 antagonism upregulated RNA-binding and synaptic proteins and downregulated inflammation and myelin associated proteins. These results underscore miR-29's pivotal role in memory persistence, plasticity, and cognitive aging, suggesting that miR-29 modulation could offer potential strategies for cognitive enhancement and age-related memory decline.

Keywords

fear conditioning; epigenetics; non-coding RNA; microRNA; hippocampus; neural plasticity; cognitive decline; aging

Introduction

MicroRNAs (miRNAs) are small non-coding RNA molecules that modulate gene expression by binding to complementary sequences in target mRNAs, leading to the repression of translation and transcript destabilization [1].Through their ability to regulate hundreds to thousands of targets simultaneously, miRNAs enable fine-tuned, reversible control of gene expression programs. This regulatory flexibility is particularly important in the nervous system, where dynamic and coordinated changes in gene expression underlie synaptic plasticity, learning, and memory formation [2, 3].

One particular miRNA family, miR-29, has emerged as a potential regulator at the intersection of aging and memory. MiR-29 is extensively involved in a wide range of biological processes relevant to brain function such as apoptosis [4], metabolism [5], neuronal maturation [4], immunity [6, 7], and the aging-associated accumulation of iron in the brain [4, 8]. Notably, miR-29 expression increases across the lifespan in multiple tissues [9], including the nervous system of multiple species [10–12], and altered miR-29 levels have been associated with cognitive impairment [13]. However, the functional implications of these expression changes for neuronal plasticity and memory remain incompletely understood. Previous studies have reported seemingly opposing functional roles for miR-29 in the brain. Partial reduction of miR-29 levels has been shown to reduce lifespan, induce neurodegeneration, and promote molecular hallmarks associated with brain dysfunction [4, 8, 14]. On the other hand, however, overexpression of miR-29 has even been shown to induce premature aging in mice, suggesting that miR-29 is a driver of the aging process [14].

The miR-29 family, which includes miR-29a, miR-29b, and miR-29c, is encoded by two genomic loci: miR-29a/b-1 and miR-29b-2/c. Among these, miR-29a-3p (hereafter miR-29a) is particularly noteworthy, as it is the most upregulated during postnatal development and maturation of the mouse visual cortex [15] and its downregulation has been shown to slow cognitive decline and reduce beta-amyloid deposition [16]. At the molecular level, miR-29a could influence memory by modulating genes related to extracellular matrix and transcription regulation [15]. In particular, miR-29a directly targets DNA methyltransferases (DNMTs) [17–19], enzymes responsible for establishing and maintaining DNA methylation patterns. DNA methylation is a critical epigenetic mechanism for regulating gene expression in the brain and has been implicated in synaptic plasticity, learning, and memory [20, 21]. Rapid and dynamic changes in DNA methylation have been observed in response to neuronal activity [22, 23], and studies have demonstrated that DNA methylation writers are necessary for learning and memory stability [24–26]. Moreover, within the brain-expressed miRNAs, miR-29a emerges as notably synaptically enriched [27], suggesting the involvement of miR-29a in the regulatory network underpinning synaptic plasticity and the formation of memories [28]. However, the precise mechanisms through which miR-29a influences memory performance during development and in the adult brain remain elusive.

In particular, whether the age-dependent increase in miR-29 expression is protective or
disruptive for cognitive function is still to be ascertained. Our study shows that inhibiting
miR-29a using Locked Nucleic Acid (LNA) antagomirs (anti-miR29a) resulted in increased
DNMT3a levels and improved memory retention, while enhancing miR-29a levels had the
opposite effect. These effects could also be attributed to miR-29c, as anti-miR-29a may
target both isoforms due to their sequences differing by only one nucleotide and sharing the
same seed region, thereby largely regulating the same targets. These findings suggest that
miR-29a not only plays a role in age-related cognitive decline but also exerts a bidirectional
control over memory possibly by altering the epigenetic landscape. Indeed, we observe
increased CpG methylation in promoter and exon regions associated with CpG islands and
modulation of the hippocampal transcriptome and proteome. These effects of miR-29a
antagonization on DNA methylation were blocked by down regulation of *Dnmt3a*. Our
findings provide compelling evidence that miR-29a orchestrates the molecular changes
necessary for the maintenance of emotional memories, highlighting its potential role in age-
related cognitive decline.

Results

Age-Dependent increase of miR-29a levels correlates with *Dnmt3a* reduction
in the dorsal hippocampus

100
101

To investigate the temporal expression pattern of miR-29a in the dorsal hippocampus, we
conducted a quantitative PCR (qPCR) analysis at different time points. Our results revealed
a significant 10-fold increase in miR-29a levels after postnatal day 10 (P10), reaching a
plateau around P60 (Figure 1A). Because expression levels remain stable after this peak,
we conducted the subsequent experiments at P90 to ensure that our analyses captured
robust and sustained miR-29a expression. Interestingly, the specific temporal window of
miR-29a increase aligns with critical periods characterized by heightened plasticity in
various brain regions, accompanied by an increased sensitivity of chromatin configuration to
environmental stimuli [29, 30]. Simultaneously, we examined the age-related modulation of
Dnmt3a, a well-validated target of miR-29 [18, 31]. With increasing age, *Dnmt3a* expression
showed a significant downregulation (Figure 1B), displaying a subject-by-subject negative
correlation with miR-29a levels (Figure 1C). In a previous study, a comparable expression
pattern was reported for the visual cortex, wherein the age-related elevation of miR-29a and
decrease in *Dnmt3a* expression levels were associated with a significant decrease in ocular
dominance plasticity and the expression of molecular critical period regulators [15]. Given
that age-related decline in *Dnmt3a* levels has been proposed as a potential mechanism
contributing to age-related cognitive impairments [32], we decided to explore the
involvement of miR-29a in modulating fear memories. To evaluate whether and how the
reduction in miR-29a expression affects fear memory formation and extinction, we injected a
LNA oligonucleotide, whose sequence is complementary to miR-29a (anti-miR29a), or

scrambled oligonucleotide (scr) in the dorsal hippocampus of adult mice, and then we tested

hippocampal memories through a hippocampus dependent trace fear conditioning protocol

(TFC) (Figure 1D). As previously observed using LNA oligonucleotides against miR-29a in

the visual cortex [15], anti-miR29a demonstrated a stable and specific binding affinity to their

complementary miRNA, effectively blocking its activity, and exhibiting resistance to

enzymatic degradation. These attributes render LNAs highly tolerable and validated, as

underscored by their approval for clinical use in human subjects [33–35]. As expected, the

qPCR revealed that anti-miR29a treatment caused a decrease in miR-29a (Figure 1E) and

an increase in the *Dnmt3a* (Figure 1F). We also found a sample-by-sample negative

correlation between *Dnmt3a* and miR-29a levels after LNA or scr injection (Supplementary

Figure 1A) confirming the tight regulation of *Dnmt3a* levels by miR-29a. To control for the

spatial spreading of our treatment, we quantified *Dnmt3a* levels also in the ventral

hippocampus. The results revealed no significant difference in *Dnmt3a* expression

(Supplementary Figure 1B), confirming the specificity of the injection effects within the

dorsal hippocampus. No effect was observed on other factors related to DNA methylation

such as MeCP2 and Tet3 (Supplementary Figure 2A-D). To assess specificity of the

treatment among miR-29 isoforms, we assessed the expression of miR-29b and miR-29c in

anti-miR29a treated animals. We found that miR-29c expression was strongly reduced,

while no significant alterations were detected in miR-29b expression (Supplementary Figure

3A-B). This result is consistent with the fact that miR-29a and miR-29c share an identical

seed sequence and differ by only one nucleotide located outside of the seed region and

regulate, regulating largely overlapping target transcripts.

Figure 1: Age-dependent increase of miR-29a correlates with *Dnmt3a* expression in the dorsal
hippocampus. **(A)** Dorsal hippocampus miR-29a expression level (normalized to P10) at different
ages (Kruskal-Wallis: H(5) = 16.22, P < 0.006; Dunn's multiple comparison vs P10: *P < 0.05 , n = 3/4
mice for each age). **(B)** Dorsal hippocampus *Dnmt3a* expression level (normalized to P10) at different
ages (Kruskal-Wallis: H(5) = 17.10, P < 0.004; Dunn's multiple comparison vs P10: **P < 0.01; n = 3/4
mice for each age). **(C)** Animal by animal correlation between normalized miR-29a and *Dnmt3a* levels
in the dorsal hippocampus at different ages (Spearman's correlation). **(D)** Experimental design for
LNA anti-miR29a treatment in the dorsal hippocampus. **(E)** Effects of anti-miR29a treatment on miR-
29a expression. Fold change values normalized to scr treated animals (scr: N = 10, anti-miR29a N =
11; Mann-Whitney U test: U = 1, p < 0.0001). **(F)** Effects of LNA-anti-miR29a treatment on *Dnmt3a*
expression. Fold change values normalized to scr treated animals (scr: N = 10, anti-miR29a N = 11;
Mann-Whitney test: U = 13, p < 0.01). PND = Postnatal Day.

MiR-29 downregulation promotes memory stability during fear extinction

Three days after anti-miR29a injection, we conditioned animals using a TFC protocol.
During TFC the neutral conditioned stimulus (CS, tone) and the aversive unconditioned
stimulus (US, shock) are separated in time by a trace interval (trace). The absence of
contiguity between the tone and the shock critically involves the hippocampus and this
protocol can be used to evaluate hippocampal dependent learning and memory [36] (Figure
2A). We observed that the treatment did not affect learning acquisition, since both anti-
miR29a and scr treated mice showed similar freezing during learning (Figure 2B and
Supplementary Figure 4A-C). When testing freezing responses 24h after training, the two
groups still did not differ in the percentage of time freezing (Figure 2D). However, on the first
day of the extinction protocol (day 3, Early Extinction) we observed that anti-miR29a treated
mice showed a higher fear retention, as compared to the control group (Figure 2E). This
observation suggests that miR-29a developmental upregulation affects strength and
persistence of memory, but it does not affect memory formation.

We also assessed freezing behavior induced by exposure to the conditioned context (Figure
2A). Remarkably, mice subjected to anti-miR29a treatment exhibited a pronounced
enhancement in their response to the contextual cues compared with the scr control group
(Figure 2C). Despite this general amplification in fear response, anti-miR29a treated mice
exhibited a typical extinction profile. Following the second extinction session (Late
Extinction), both anti-miR29a and scr treated mice demonstrated significantly reduced
freezing responses (Figure 2F). These findings offer insights into miR-29 role in modulating
the persistence of fear memories. This effect may be mediated through the regulation of
miR-29 target genes, particularly those involved in DNA methylation pathways and gene
expression control.

Figure 2: Reducing hippocampal miR-29a expression promotes memory stability. **(A)** Diagram showing the TFC paradigm and timeline. **(B)** Anti-miR29a and scr-treated animals showed no differences at baseline or in fear memory acquisition, as indicated by similar freezing responses during the final two-tone, trace trials and ITI of the TFC training session (Two-way ANOVA, test interval × treatment interaction $F(3, 76) = 0.39$, $p = 0.76$; main effect of test interval $F(3, 76) = 64.33$, $p < 0.0001$; main effect of treatment, $F(1, 76) = 0.93$, $p = 0.34$). **(C)** Downregulation of miR-29a in the dorsal hippocampus enhances contextual memory recall (Mann-Whitney U test: $U = 16$, $p < 0.01$, scr: $N = 10$, anti-miR29a $N = 11$). **(D)** No treatment differences were observed at baseline or during the final two-tone and trace trials of the TFC fear recall session (Two-way ANOVA, test interval × treatment interaction $F(2, 57) = 0.59$, $p = 0.56$; main effect of test interval $F(2, 57) = 41.74$, $p < 0.0001$; main effect of treatment, $F(1, 57) = 2.33$, $p = 0.13$). **(E)** Percentage of freezing during Early extinction. anti-miR29a mice showed a higher fear retention compared to the control group during both tone (Two-way RM ANOVA, main effect of treatment $F(1, 19) = 4.78$, $p < 0.05$; main effect of trials $F (2.98, 56.58) = 6.69$, $p < 0.0001$; trials × treatment interaction $F(5, 95) = 0.29$, $p = 0.92$) and trace intervals (Two-way RM ANOVA, main effect of treatment $F(1, 19) = 4.47$, $p < 0.05$; main effect of trials $F(3.26, 62.02) = 9.36$, $p < 0.0001$; trials × treatment interaction $F(5, 95) = 0.77$, $p = 0.57$). **(F)** During the Late extinction, anti-miR29a treated mice showed no significant differences in the percentage of freezing compared with controls (Tone: Two-way RM ANOVA, main effect of treatment $F(1, 18) = 0.19$, $p = 0.66$; main effect of trials $F (2.56, 46.16) = 4.81$, $p < 0.01$; trials × treatment interaction $F(5, 90) = 0.87$, $p = 0.50$; Trace: Two-way RM ANOVA, main effect of treatment $F(1, 18) = 0.24$, $p = 0.63$; main effect of trials $F(3.95, 71.15) = 6.26$, $p < 0.001$; trials × treatment interaction $F(5, 90) = 2.12$, $p = 0.07$). scr: $N = 10$, anti-miR29a $N = 11$; *indicates the main effect of treatment, #indicates the main effect of trials. CS= conditioned stimulus

MiR-29 inhibition alters DNA methylation patterns and influences memory-associated pathways at the transcript and protein level

We employed reduced representation bisulfite sequencing (RRBS) to investigate changes in DNA methylation patterns induced by anti-miR29a treatment in comparison to scr treated

hippocampal samples. RRBS allows for an exploration of methylation alterations in a CpG-
enriched subset of the genome, primarily focusing on promoters [22, 37]. We analyzed
differential methylation of the sequenced CpGs and selected differentially methylated CpGs
(DMCs, q<0.05) with a difference greater than 5% in absolute value. Our analysis identified
35,246 DMCs after anti-miR29a treatment, with a notable prevalence of hypermethylation
(32,878 CpGs) over hypomethylation (2,368 CpGs) (Figure 3A). To test whether these
methylation changes are mediated by DNMT3a, we analyzed hippocampal samples treated
with anti-*Dnmt3a* alone or in combination with anti-miR29a. Anti-*Dnmt3a* treatment resulted
in 1,742 hypermethylated and 2,082 hypomethylated CpGs, showing a bias toward
hypomethylation. The combined treatment produced 3,491 hypermethylated and 6,143
hypomethylated CpGs, displaying a similar hypomethylation-skewed profile (Figure 3A).
These results indicate that DNMT3a inhibition counteracts the hypermethylation bias
induced by anti-miR29a and suggest that a substantial fraction of miR-29a-dependent
methylation changes are mediated through DNMT3a. Partial least squares discriminant
analysis (PLS–DA) of DNA methylation profiles revealed clear separation of anti-miR29a
samples from controls as well as from all other groups. Anti-*Dnmt3a* and combined
treatment samples largely overlapped with each other but remained distinct from controls,
which formed a separate cluster (Figure 3B). These findings suggest that DNMT3a activity
contributes substantially to the methylation landscape induced by miR-29a inhibition.
Comparison of DMCpGs between anti-miR29a alone and combined treatment revealed a
highly non-random distribution of concordant and discordant CpGs (χ^2 = 473.11, p < 2.2 ×
10^{-1}). DMCpGs were classified based on their methylation changes in the two treatments
into concordant and discordant quadrants. Concordant hypermethylated CpGs, which were
hypermethylated in both anti-miR29a and combined treatment, were enriched for synaptic
processes, including vesicle-mediated transport in synapse (Figure 3C). In contrast,
discordant CpGs, which were hypermethylated in anti-miR29a but hypomethylated in the
combined treatment, were the majority and were enriched for regulation of membrane
potential, positive regulation of nervous system development, forebrain development, and
cell junction assembly (Figure 3C). These results indicate that miR-29 inhibition affects
multiple neuronal pathways: some epigenetic changes persist even when DNMT3a is
inhibited, while others are reversed, highlighting a partial DNMT3a dependence of miR-29-
mediated DNA methylation in hippocampal neurons.

Gene Ontology (GO) enrichment analysis of hypermethylated CpGs identified after anti-
miR29a treatment revealed enrichment for pathways related to synaptic signalling,
extracellular matrix organisation, and kinase activity. Mapping methylation changes onto the
KEGG glutamatergic synapse pathway (mmu04724) using Pathview showed widespread
hypermethylation across multiple synaptic genes, suggesting that miR-29 regulates
epigenetic programs involved in hippocampal-dependent plasticity and memory persistence
(Figure 3D). Finally, analysis of global CH methylation differences across treatment groups
revealed significant alterations, with all pairwise comparisons reaching statistical
significance (Figure 3E), further supporting a DNMT3a contribution to the broader

epigenomic effects induced by miR-29a inhibition. The enhanced DNA methylation in
response to anti-miR29a treatment was confirmed also in an independent cohort of mice
(Supplementary Figure 5).

To investigate the possible downstream consequences of epigenetic changes we performed
RNAseq and mass-spectrometry-based proteomics (Figure 4). The analysis of differentially
expressed proteins retrieved 420 significant protein groups out of 4878 detected protein
groups. RNAseq detected 946 differentially expressed genes (DEGs). The analysis of the
most overrepresented miRNA binding site in up-regulated DEGs revealed the expected
enrichment for miR29 binding sites (miTEA [40], FDR < 10^{-16}), confirming effective
antagonism of miR-29 family. We first sought to identify pathways that regulated in a
congruent direction both at the transcript and protein level, among these, glutamatergic
transmission (KEGG mmu04724) stood out as glutamate receptors and transporters were
upregulated often both at transcript and protein levels (Figure 4A). This pattern is clearly
consistent with enhancement of synaptic plasticity, which is further supported by a large
fraction of overrepresented GO terms related to synaptic function and plasticity, as well as
neural development, among up-regulated genes (Supplementary Figure 6A). The analysis of
downregulated genes revealed the strongest enrichment for genes expressed in microglia,
followed by cells of both the myeloid and lymphoid lineage (Figure 4B). In addition, a striking
down-regulation of genes coding for lysosomal protein (KEGG mmu04142) was observed
(Figure 4C). This is further supported by a large fraction of overrepresented GO terms
related to both adaptive and innate immunity among up-regulated genes (Supplementary
Figure 6B). Downregulated proteins, on the other hand, showed a significant enrichment for
genes expressed in oligodendrocytes only (Figure 4D) and coding for myelin sheath (GO
0043209).

To complement the analysis of GO term overrepresentation, we explored the interactome of
regulated proteins and transcripts by analyzing the physical interactions of the upregulated
differentially-expressed proteins (DEPs, FDR <0.1, Log2(fold change) > 0.1). We applied
STRING network analysis [41] and found DEPs to be strongly connected (Supplementary
Figure 7, expected number of edges: 295; number of edges: 938; PPI enrichment p-value:<
$1.0e^{-16}$). The protein network consisted almost entirely of two tightly clustered communities
of functionally-interacting proteins: one made of synaptic proteins and the second of RNA
binding proteins. STRING network analysis found DEGs (FDR <0.1, DESeq2) to be strongly
connected (Supplementary Figure 7, expected number of edges: 345; number of edges:
660; PPI enrichment p-value:< 1.0e-16). Similarly to DEP network, also the DEG network
was strongly enriched for synaptic terms and RNA binding activity, indicating that the above-
described regulation is transcriptionally-driven. STRING analysis of downregulated DEPs
(FDR <0.1, Log2(fold change) <- 0.1) revealed a significantly connected network (expected
number of edges: 142; number of edges: 260; PPI enrichment p-value:< 1.0e-16) with
overrepresentation of terms related to immunity, myelin, mitochondria and ribosome. The
analysis of downregulated DEGs (FDR <0.1, DESeq2) revealed a significantly connected
network (expected number of edges:320 ; number of edges: 1482; PPI enrichment p-value:<

1.0e^{-16}) with prominent overrepresentation of immunity-related terms and an interesting
overrepresentation of the specific WiKI pathway WP3625 Tyrobp causal network in
microglia. This network was identified by causal analysis of AD-driving mutations [42].
These data confirm the analysis of GO term overrepresentation and indicate that antagonism
of miR-29 family dampens the inflammatory reaction induced by the injection via a
transcriptional mechanism and reduces myelin, a major plasticity brake [43, 44], via
transcript-independent mechanisms.

Figure 3: Dnmt3a mediates the epigenomic effects of miR-29a in the mouse dorsal

hippocampus. (A) Differentially methylated CpGs (DMCpGs). Bar plots showing the number of

significantly hypermethylated (magenta) and hypomethylated (blue) CpGs identified in each treatment

compared to control (q ≤ 0.05; |Δmethylation| > 5%). Numbers above the bars indicate the absolute

number of DMCpGs in each category. Grey boxes report the total number and percentage of

DMCpGs relative to the total CpGs analyzed in each comparison. **(B)** Sample clustering based on

DNA methylation profiles. PLS–DA analysis of genome-wide DNA methylation across experimental

groups. Each dot represents one animal. The first two components are shown (comp1: 10% variance

explained; comp2: 7%). Ellipses indicate the 95% confidence interval for each group, highlighting

separation among treatments. **(C)** Differentially methylated CpGs and GO enrichment analysis across

treatment conditions. Central panel: Scatter plot of DMCpGs comparing methylation changes in the

combined treatment (anti-miR29a + anti-*Dnmt3a* vs control; x-axis) with those in anti-miR29a alone

(vs control; y-axis). Each dot represents one DMCpG. Dashed lines indicate no change relative to

control and define four quadrants. Quadrants 1 and 3 represent concordant CpGs (same direction of

change), whereas quadrants 2 and 4 represent discordant CpGs (opposite direction). The number of

CpGs per quadrant is shown in red. A χ^2 test revealed a significant enrichment of concordant CpGs

compared to a uniform distribution (χ^2 = 473.11, p < 2.2e-16). Left and right panels: GO Biological

Process enrichment analyses performed on CpGs from quadrant 2 and quadrant 1, respectively. Bars

show significantly enriched GO terms, plotted as −log10(FDR). **(D)** Pathway mapping of differentially

methylated genes. Schematic representation of the glutamatergic synapse pathway with genes

associated with DMCpGs overlaid onto the pathway map. Nodes are color-coded according to the

direction and magnitude of methylation change (green: hypomethylation; red: hypermethylation; color

scale shown above). This analysis highlights coordinated methylation changes affecting multiple

components of synaptic signaling and plasticity pathways. **(E)** Global CH methylation changes.

Boxplots showing the distribution of Δ methylation (treated − control) at CH sites for each treatment

group. Values represent per-animal differences between treated and contralateral control

hemispheres. Statistical comparisons were performed using Welch's ANOVA followed by Bonferroni-

corrected pairwise t-tests. Horizontal lines indicate medians; boxes represent the interquartile range

(IQR); whiskers extend to 1.5× IQR. All pairwise comparisons were significant (p < 0.001).

Figure 4. Integrated transcriptomic and proteomic analysis reveals synaptic potentiation and downregulation of immune and lysosomal programs. (A) Mapping of differentially expressed genes and proteins onto the KEGG glutamatergic synapse pathway (mmu04724). Ionotropic (e.g., NMDA, AMPA subunits), metabotropic receptors, and glutamate transporters are predominantly upregulated at the transcript and/or protein level, indicating enhanced synaptic plasticity. Colors indicate: transcript (blue), protein (red), both (solid red). **(B)** Cell-type enrichment analysis of downregulated genes shows strong overrepresentation of microglia-expressed genes, followed by

344 myeloid and lymphoid cells. **(C)** Mapping of downregulated genes and proteins onto the KEGG
345 lysosome pathway (mmu04142) reveals marked reduction of lysosomal hydrolases, membrane
346 proteins, and acidification regulators. **(D)** Cell-type enrichment analysis of downregulated proteins
347 highlights significant enrichment for oligodendrocyte proteins and myelin sheath components
348 (GO:0043209).

350 Elevated miR-29a levels in the hippocampus of adult mice result in impaired
351 memory retention

To reveal whether miR-29a levels are quantitatively related to memory retention, we asked
whether enhancing miR-29a levels in the dorsal hippocampus of adult mice could result in
impaired retention. Before conditioning, we injected animals with a synthetic miR-29a mimic
(mim29a) and then we tested defensive freezing responses using the same TFC protocol
employed in the anti-miR29a experiments (Figure 5A). Mim29a treatment resulted in an
upregulation of miR-29a expression and a concurrent downregulation of *Dnmt3a* levels in
the dorsal (Figure 5B-C) but not in the ventral hippocampus (Supplementary Figure 8)
confirming the spatial selectivity and biological activity of the treatment. At the behavioral
level, increasing miR-29a levels affected the trace stability but not the CS response. Indeed,
during the extinction protocol (Early Extinction) mim29a mice showed an impaired trace fear
retention compared to the control group (Figure 5F). A trend toward a reduced response to
the trace interval was also evident during both the training and recall phases (Figure 5D and
Supplementary Figure 9A-C). As observed also with anti-miR29a, mim29a treated mice
displayed a normal extinction profile (Figure 5F-G). The bidirectional impact of miR-29a on
memory strength during the extinction was also evident for contextual memories. In
particular, mice receiving miR-29a mimics displayed a strong deficit in the recall of
contextual fear memories (Figure 5E) as compared to controls. The involvement of miR-29a
in the maintenance of hippocampal fear memory stability was also confirmed by the
presence of an inverse correlation between miR29a expression levels and the percentage of
freezing behavior exhibited during early extinction and contextual fear recall (Figure 6A-E).
Overall, these findings support the hypothesis that heightened miR-29a levels contribute to
the compromised stability of hippocampal fear memories [16, 32, 45].

Figure 5: Hippocampal miR-29a upregulation impaired memory retention. **(A)** Diagram showing the TFC paradigm and timeline. To guarantee elevated miR-29a levels during all the sessions of the TFC test, mice were injected 24 hours before the start of the behavioral tests and again 24 hours before the start of the extinction protocol. **(B-C)** Effects of miR-29a mimic treatment on **(B)** miR-29a expression levels (fold change values normalized to scr treated animals; scr: N=8, mim29a N=8; Mann-Whitney U test: U = 0, $p < 0.001$) and **(C)** *Dnmt3a* expression levels (fold change values normalized to scr treated animals; scr: N = 7, mim29a N = 6; Mann-Whitney U test: U = 0, $p < 0.01$). The lower sample size for *Dnmt3a* reflects insufficient RNA in some samples; all available data were included. **(D)** No treatment differences were observed at baseline or during the final two-tone and trace trials of the TFC fear recall session (Two-way ANOVA, test interval × treatment interaction $F_{(1, 42)} = 3.52$, $p = 0.07$; main effect of test interval $F_{(2, 42)} = 23.89$, $p < 0.0001$; main effect of treatment, $F_{(2, 42)} = 1.30$, $p = 0.28$). **(E)** MiR-29a levels increase resulted in impaired contextual memory recall (Mann-Whitney U test: U = 12, $p < 0.05$, scr: N = 8, miR-29a mimic N = 8). **(F)** Percentage of freezing during Early extinction. Mim29a treated mice showed lower fear retention compared to the control group during the trace (Two-way RM ANOVA, main effect of treatment $F_{(1, 19)} = 4.475$, $p < 0.05$; main effect of trials $F_{(3.26, 62.02)} = 9.365$, $p < 0.0001$; trials × treatment interaction $F_{(5, 95)} = 0.77$, $p = 0.57$) but not during the tone interval (Two-way RM ANOVA, main effect of treatment $F_{(1, 14)} = 2.30$, $p = 0.15$; main effect of trials $F_{(3.21, 44.97)} = 3.80$, $p < 0.05$; trials × treatment interaction $F_{(5, 70)} = 0.65$, $p = 0.66$). **(G)** During the Late extinction, mim29a treated mice showed no significant differences in the percentage of freezing compared with controls (Tone: Two-way RM ANOVA, main effect of treatment $F_{(1, 14)} = 1.33$, $p = 0.27$; main effect of trials $F_{(3.63, 50.83)} = 6.41$, $p < 0.001$; trials × treatment interaction $F_{(5, 70)} = 1.46$, $p = 0.21$; Trace: Two-way RM ANOVA, main effect of treatment $F_{(1,14)} = 2.54$, $p = 0.13$; main effect of trials $F_{(3.35, 46.88)} = 9.09$, $p < 0.0001$; trials ×

399 treatment interaction F(5, 70) = 2.14, p = 0.07). scr: N = 8, mim29a N = 8; *indicates the main effect of
400 treatment, #indicates the main effect of trials.

Figure 6: Behavioral correlation with miR-29a levels post LNA and mimic injection. **(A)** During
early extinction miR-29a levels negatively correlate with freezing responses at the first two tone and
(B) trace presentations (Spearman correlation). **(C)** MiR-29a levels negatively correlate with freezing
responses during contextual fear recall (Spearman correlation). **(D)** During early extinction miR-29
levels negatively correlate with freezing responses to the tone across all trials **(E)** and with freezing
responses during the trace period across all trials (Spearman correlation).

Discussion

Regulation of the miR-29 family is characterized by a striking degree of evolutionary
conservation. Previous studies have demonstrated that miR-29 isoforms exhibit an age-
associated upregulation across diverse species and tissues [11, 12, 46]. Furthermore, miR-
29a plays a pivotal role in governing age-dependent processes such as neuronal maturation
and iron accumulation within the brain [4, 8]. In the visual cortex, miR-29a is the microRNA
with the most significant upregulation during the visual critical period [10, 15] [15]. Our data
confirmed this age-dependent upregulation of endogenous miR-29a expression in the dorsal
hippocampus of wild-type mice, revealing a tenfold increase between P10 and P60.
Importantly, a large-scale miRNA association study in human postmortem brain samples
identified miR-29a among the miRNAs most strongly associated with cognitive trajectories,

with higher miR-29a levels emerging as the second most significant correlate of accelerated
cognitive decline (Wingo et al. 2022). Together, these observations suggest that elevated
miR-29 levels may be linked to reduced plasticity and cognitive performance. To directly
investigate the functional role of miR-29a in memory, we bidirectionally modulated its levels
in the dorsal hippocampus of adult mice. Downregulation of miR-29a enhanced
hippocampus-dependent memory strength, whereas miR-29a overexpression impaired
memory performance, resembling aspects of age-associated cognitive decline. It is
important to note that the anti-miR-29a antagomir used in our experiments also targets miR-
29c. Given that miR-29a and miR-29c differ by only a single nucleotide and share the same
seed sequence, they are expected to regulate largely overlapping target genes; thus, the
observed effects may reflect combined modulation of both isoforms. The tight relationship
between miR-29 levels and memory is emphasized by the negative correlation between
miR-29a levels and the percentage of freezing observed during early extinction and
contextual fear recall. Conversely, levels of *Dnmt3a* exhibited a positive correlation with the
percentage of freezing exhibited in the initial stages of trace extinction. This finding lends
support to the specific regulatory role of miR-29 in memory persistence. Recent work shows
that miR-29a downregulation in the hippocampus of 5XFAD mice results in amelioration of
memory performances, reduction of beta-amyloid deposition, and lower microglia and
astrocytes activation [16]. Collectively, these studies support a model in which miR-29a acts
as a key regulator of age-associated molecular pathways that constrain plasticity and
memory maintenance.

miR- 29 sculpts the epigenetic landscape of adult hippocampus

Epigenetic mechanisms have been increasingly recognized as important regulators of
cognitive functions through the modulation of transcriptional responses [47, 48]. Since we
found a strong negative correlation between miR-29a and *Dnmt3a* levels, we decided to
explore the profile of methylation of anti-miR29a treated mice. DNA methylation is an
essential epigenetic mechanism for various cellular processes and has emerged as a crucial
regulator of memory consolidation [49, 50]. anti-miR29a treatment in the adult visual cortex
reactivates ocular dominance plasticity in adult mice and upregulates genes associated with
the extracellular matrix and epigenetic remodeling, however whether manipulations of miR-
29 levels could result in global changes of DNA methylation was utterly unexplored.
Contrary to what is observed for miR-29 levels, the degree of DNA methylation decreases
with age in many tissues, including the nervous system [51, 52]. In particular, aging has
been associated with a reduction in the expression of the DNA methyltransferase *Dnmt3a2*
in the hippocampus, and restoring *Dnmt3a2* levels has been shown to rescue cognitive
functions affected by aging [32]. It has also been reported that hippocampal *Dnmt3a2* levels
determine cognitive abilities in both young adult and aged mice [53–56]. Kupke et al,
however, showed that Dnmt3a1 regulates hippocampus-dependent memory via Nrp1 [57].

Furthermore, DNA methylation induced selectively within neuronal ensembles has been
proposed as a mechanism for stabilizing engrams during consolidation, supporting

successful memory retrieval [58]. Consistent with this framework, inhibition of miR-29a led to
a robust increase in CpG methylation across the hippocampal genome, in parallel with
increased *Dnmt3a* expression. Importantly, these methylation changes were substantially
blocked by *Dnmt3a* downregulation, demonstrating that a major component of the
epigenetic remodeling induced by miR-29a antagonism depends on DNMT3a activity. A
direct comparison of DMCpGs between anti-miR29a alone and the combined anti-miR29a +
anti-DNMT3a treatment revealed a highly non-random distribution of concordant and
discordant CpGs, supporting a structured and pathway-specific epigenetic response.
Concordant hypermethylated CpGs, those remaining hypermethylated even after DNMT3a
inhibition, were enriched for synaptic processes, including vesicle-mediated transport at the
synapse. In contrast, discordant CpGs, hypermethylated after anti-miR29a but
hypomethylated when DNMT3a was simultaneously inhibited, were enriched for processes
such as regulation of membrane potential, nervous system development, forebrain
development, and cell junction assembly. These findings indicate that miR-29a inhibition
affects multiple neuronal pathways, with a subset of methylation changes being DNMT3a-
dependent and others likely mediated through additional mechanisms. Global CH
methylation was also significantly altered across treatment groups, further supporting a
broad DNMT3a contribution to the epigenomic remodeling triggered by miR-29a
antagonism. Gene Ontology and KEGG pathway analyses reinforced the functional
relevance of these methylation changes. Hypermethylated CpGs following anti-miR29a
treatment were enriched in pathways related to synaptic signaling, extracellular matrix
organization, kinase activity, and notably the glutamatergic synapse pathway. Mapping
methylation changes onto the KEGG glutamatergic synapse pathway revealed widespread
hypermethylation across synaptic genes, suggesting that miR-29a inhibition engages
epigenetic programs closely linked to hippocampal-dependent plasticity and memory
persistence. While promoter methylation is often associated with transcriptional repression,
gene body methylation is positively correlated with gene expression [59, 60], indicating that
the observed hypermethylation may also facilitate transcriptional activation of plasticity-
related genes rather than suppress them. To explore the functional consequences of this
epigenetic remodeling, we performed RNA sequencing and mass-spectrometry–based
proteomics. Both datasets converged on a coherent molecular signature. Upregulated
transcripts and proteins were strongly enriched for synaptic and plasticity-related pathways,
with glutamatergic transmission (KEGG mmu04724) standing out prominently. Glutamate
receptors and transporters, including NMDA receptor subunits, were upregulated at both
transcript and protein levels. Network analysis revealed highly interconnected clusters
composed predominantly of synaptic proteins and RNA-binding proteins, indicating a
coordinated and transcriptionally driven enhancement of synaptic machinery. The
enrichment of miR-29 binding sites among upregulated genes confirmed effective
antagonism of the miR-29 family and supports the view that these transcriptional changes
are directly linked to miR-29 inhibition.

Conversely, downregulated transcripts were strongly enriched for genes expressed in
microglia and other immune-related cell types, including components of the Tyrobp causal
network implicated in Alzheimer's disease. Lysosomal pathways were also significantly
reduced. At the protein level, downregulated groups were enriched for oligodendrocyte-
specific and myelin-associated proteins. Together, these findings indicate that miR-29
antagonism dampens inflammatory and microglial activation through transcriptional
mechanisms while reducing myelin-related proteins through mechanisms that may be
partially transcription-independent. Given that myelin constitutes a well-established brake on
plasticity [43, 44], its reduction may further contribute to the enhanced plastic potential
observed after miR-29 inhibition. Overall, these results delineate a coordinated molecular
program in which miR-29a antagonism promotes DNMT3a-dependent epigenetic
remodeling, enhances glutamatergic synaptic function, suppresses inflammatory signaling,
and reduces structural constraints on plasticity. This integrated regulation could be further
complemented by studies aimed at identifying the specific cell types involved in miR-29a
action, thereby providing a more refined mechanistic framework linking miR-29a levels to
memory persistence. Future studies examining memory performance across developmental
stages characterized by distinct endogenous miR-29a levels will further clarify how this
regulatory axis contributes to age-dependent changes in cognitive function.

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to the lead contact, Tommaso Pizzorusso (tommaso.pizzorusso@sns.it).

Materials availability

This study did not generate new unique reagents.

Data and code availability

- All data reported in the main text or supplemental information are available from the lead contact upon request.
- All original code is also available from the lead contact upon request.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Acknowledgments

We gratefully acknowledge NVIDIA Corporation's support with the Jetson AGX Xavier Developer Kit donation for this research. European Union- Next Generation EU, Mission 4 Component 1 CUP E53C24001460006, project TNE- NEUROBRIDGE and PRIN2022 20228RMXBE to T.P.

Author contributions

Conceptualization: AV, AC, TP

Methodology: AV, EP, CG, RM, SB, PT, AC, TP

Formal analysis: AV, CG, SB, PT, AC,

Investigation: AV,CG,EP, SB, PT,

Resources: AC, TP, RM

Data curation: AV, CG, SB, AC

Writing original draft: AV, AC, TP

Visualization: AV, CG, SB, AC

Supervision: TP, AC

Project administration: TP, AC

Funding acquisition: TP, AC

554 Declaration of interests

555 The authors declare no competing interests.

Experimental Procedures

Animals

All experiments utilized the C57BL/6J strain. Animals were maintained at 22°C under a 12-h light–dark cycle (average illumination levels of 1.2 cd/m2) and housed in standard cages according to current regulations about animal welfare. Food (4RF25 GLP Certificate, Mucedola) and water were available ad libitum. All the experiments were carried out according to the directives of the European Community Council (2011/63/EU) and approved by the Italian Ministry of Health (621/2020-PR). All tissue explants were performed at the same time of the day (10–12 AM, Central European Time (CET, Italy)). Both male and female mice were included in the study, and no significant sex × treatment interactions were detected (Supplementary Table 1).

Stereotaxic intracranial injections

Mice were anesthetized with isoflurane (3% induction and 1.5% maintenance) and secured in a stereotaxic frame (Sutter Instrument, Novato, CA, USA) using ear bars. Prilocaine was used as a local anesthetic for the acoustic meatus. Body temperature was maintained at 37° C using a heating pad. The eyes were treated with a dexamethasone-based ophthalmic ointment (Tobradex, Alcon Novartis) to prevent cataract formation and keep the cornea moist. Respiration rate and pedal reflex were checked periodically to maintain an optimal level of anesthesia. Prior to scalp removal, a subcutaneous local injection of lidocaine (2%) was performed. After exposing the skull, small burr holes were made using a surgical drill. In separate experiments, mice received either LNA oligonucleotide complementary to miR-29a (miR-29a-3p miRCURY LNA, QIAGEN, Cat. No. 339121, 50 μM) or a synthetic miR-29a mimic (mirVana™ miRNA miR-29a-3p Mimic, Thermo Fisher Scientific, Cat. No. 4464066, 100 μM). The corresponding scrambled oligonucleotides were used as controls (Negative Control miRCURY LNA miRNA Mimic, QIAGEN, Cat. No. YM00479902, 50 μM; mirVana™ miRNA Mimic Negative Control #1, Thermo Fisher Scientific, Cat. No. 4464058, 100 μM). To assess the contribution of *Dnmt3a* to the effects mediated by miR-29a, a GapmeR antisense oligonucleotide targeting murine *Dnmt3a* was injected. In these experiments, the GapmeR anti-*Dnmt3a* in combination with the miR-29a LNA was delivered into one hippocampus, while the contralateral hippocampus received the corresponding control oligonucleotides. Additional control conditions included injections of GapmeR anti-*Dnmt3a* alone or the respective control oligonucleotides. Injection in the dorsal hippocampus was done using the following stereotaxic coordinates: –2 mm anteroposterior, ±1.5 mm mediolateral, –1.5, –1.7 mm dorsoventral from bregma. A total of 2 μl of solution per hemisphere was injected, 1μl for each coordinate using a Hamilton syringe. The injection needle was left in place for an additional 60 s to allow the fluid to diffuse. The skin was subsequently sutured, and a physiological solution was injected subcutaneously to prevent dehydration. At the end of the surgical procedure, mice were left in a heated cage for

recovery. Once fully awake, they were returned to their home cage. Paracetamol was
administered in water ad libitum for two days. Due to the inherent stability of LNA
oligonucleotides, a single injection was administered to mice before the beginning of the
behavioral test. In contrast, the miR-29a mimic exhibited faster degradation within a specific
three-day timeframe. To maintain the levels of miR-29a consistently elevated during all the
sessions of the behavioral test, mice received injections every two days, following the
protocol reported in Napoli et al., 2020.

Synthesis and selection of GapmeR Antisense Oligonucleotides

Three LNA GapmeR antisense oligonucleotides targeting the sequence
(AGGGTACTGGCCGCCTCTTCTTTGAGTTCTACCGCCTCCTGCATGATGCGCGGCCCAA
GGAGGGAGATGATCGCCCCTTCTTCTGGCTCTTTGAGAATGTGGTGGCCATGGGCGTT
AGTGACAAGAGGGACATCTCGCGATTTCTTGAGGTATAGACCGAGACCTTGGTTTGGC
CAGCTCACTAATGGCTTCTACCTGGGACTGCTGCTTCGTCCCTGTCTTGTCTGCATTGC
GGAGCTGGGGGATTGGAGCTGGGGACTGGTGGCTTCTCTTTGCAAGGGACAGCTTGA
GGAAGATTTTCCATGTAGAGAGGAAGCAGTGCTAAAGACCCACCTAGGAAGAAAGTTC
TCTTGTTCAAAGAGGTGTGGTGGTCATCATCAAACAGATGGACTGGGGCCAGCCCAGT
TTTTCTGTGGAGAACTCCAAAATCAGTTTTTAAAATCATTCTCTGACTAGAATGGCCTGT
GCTCACTCTCTGGCACCCTTGTGGGTGTTTTGTGATACCTGAGAGAAAATCATGGCTTA
TCTTTTTGCTTTCATTTCTTGTTATGTGAGCCCCAGTTTAGCACTGAGCCATGTACAAGC
TCAATCATAGGGGATTGCTGCTGCCCAAGGCATTCTTTTCTTTTCTTTTCTTTTCTTTTCT
TTTCTTTTCTTTTCTTTTCTTTCTCTCTCTCTCTCTTTCTTTCTTTCTTCCTTTCTTTCTTTC
TTTTTGATTTATTTATTATTATATATAAGTAGCTGACTTCAGACACACCAGAAGAGGGTGT
CAGATCTCATGGGTGGTTGTGAGCCACCATGTGGTTGCTGGGATTTGAACTCAGGACC
CTCGGAAGAGCAGTCAATGCCCTTAACTGCTGAGCCATCTCTCCAGCCCTGCCCAAGG
CATTCTTGTGGTAGGCTGTCAGCTTATAGTCCTGTCAGCCTACGCTCAATAATAACCTC
AGATTGTAAATGTGAGGACTTAGTACACAAACAGGCTTGCTCCGGTAAGGTCTTATGTG
GTCATCAGCCTTGTGGCTATTACAAGAATGAACCTCACTTTAAGCACACAAGAGTCATT
CATGGTACTACTGGAATAATAACAAAGTTCAGTTGGATTGTGCTGGCTGTCCAGAATGT
TCTAGCCAGCCGGGCGTGGTGGCTCACGCCTTTAATCCCAGCACTTGGGAGGCAGAG
GCAGGTGGATTTCTGAGTTCGAAGCCACCCGGTCTACAAAGTGAGTTCCAGGACAGCC
AGGGCTATACAGAGAAACCCTGTCTTGAAAAATAAAAATTAAAAAAAAAAAAAAAAGAATG
TTCTAGCCAGCACATGTGGGATGGATTTAAGTTTGGGTCTTCAGGTTATTCCGGTGTTT
GGCTTCCTACGGAGGAAGTTCCTCTGGGGAGTTAAGACTCTTGGCCAGCCCAGTTGCC
CAGCAGTGTTTAATTAATACTTCCTCCTTGGTCATCTTGAAACCATCTCCTATTTTACAG)
of the murine *Dnmt3a* transcript were designed and synthesized by QIAGEN (Germany). A

non-targeting control GapmeR was also designed and synthesized by the manufacturer. All

GapmeRs contained LNA modifications and a phosphorothioate backbone to enhance

nuclease resistance and promote RNase H–mediated degradation of the target RNA,

according to the manufacturer's specifications. To determine the most effective GapmeR in

our conditions, adult C57BL/6J mice were anesthetized with isoflurane and placed in a

stereotaxic frame. GapmeRs were diluted in sterile PBS and injected into the visual cortex at

a final dose of 100 pmol in a total volume of 1 μL per injection site using a Hamilton syringe.

The control GapmeR was injected into the contralateral visual cortex of the same animal.

Seven days after injection, mice were euthanized and the cortices were rapidly dissected on

ice. Tissue samples were snap-frozen in liquid nitrogen and stored at −80 °C until further

processing. RNA extraction, reverse transcription and Quantitative real-time PCR for murine

Dnmt3a were performed as described below (see section: RNA extraction). GapmeR

efficacy was determined by comparing *Dnmt3a* mRNA levels in treated cortices relative to

those injected with the control GapmeR. Among the three candidates tested, the GapmeR

producing the greatest reduction in *Dnmt3a* transcript levels was selected for subsequent

experiments (anti-*Dnmt3a*).

Trace fear conditioning and extinction protocol

Mice were subjected to a trace fear conditioning and extinction procedure using a custom-

made PVC/acrylic apparatus consisting of three radially arranged arms converging at a

central junction. Each arm measured 50 cm in length, 15 cm in width, and 21 cm in

height, and was covered by a clear plastic lid to prevent escape and allow behavioral

observation. The floors consisted of electrifiable grids of parallel stainless steel bars,

permitting delivery of aversive foot shock stimuli. A circular speaker of 5 cm diameter was

mounted centrally in the lid of each arm to deliver precisely controlled auditory cues

(conditioned stimuli). The arms differed in wall and floor textures to provide distinct

contextual cues, allowing for contextual discrimination between conditioning and extinction

environments.. During the test days, the mice were transported in their home cages to a

room adjacent to the testing room and left for 2 h before behavioral testing. We used two

different contexts: In context A, the walls had white plastic circles and the floor was

completely black; in context B, the walls had white vertical plastic strips and a white floor.

Both chambers were covered with transparent plexiglass lids with a loudspeaker in its center

point. The shock grid on the floor was made of stainless steel. Only the grid of context A

was electrified by a shock generator (World Precision Instruments, Sarasota, FL) and

guided by an Arduino Uno for CS and US parameter control and footshock delivery. Mice

behavior was recorded by a camera controlled by the EthoVision XT 8 software (Noldus

Information Technology, The Netherlands). The apparatus was cleaned before and after

each animal with 70% ethanol or 1% acetic acid for context A and context B respectively,
since the mice may associate the smell with the context. After each session, mice were
housed separately until the end of the test to avoid possible observational fear learning.
During the Training phase of the test (Day 1), mice were allowed to explore the conditioning
chamber (Context A) for 2 min (baseline) before receiving five conditioning trials. Each trial
consisted of a 20 s pure tone (80 dB, 2900 Hz) and a 2 s shock (0.6 mA) separated by a 20
s stimulus-free trace interval. The intertrial interval (ITI) was 120 s. Mice were removed from
the chamber 120 s after the last trial. Twenty-four hours later (Day 2), mice were placed
back in the conditioning chamber (Context A) for a 5 min context test. The same day (Day2)
after 2 h from the context test, mice were placed in a different context (Context B) for a
Recall test consisting of a 2 min baseline period followed by four 20 s tone presentations
separated by a 20 s stimulus-free trace interval with 120 s ITI. On day 3 (Early Extinction)
and day 4 (Late Extinction), conditioned mice were subjected to the extinction training in
context B during which they received twelve 20 s tone presentations separated by 20 s
stimulus-free trace interval presentations with 120 s ITI each day.

Analysis of Freezing Behavior

Recorded videos were manually scored for freezing behavior by two separate experimenters
blind to treatment conditions. Mice were considered to be freezing if no movement was
detected for 2 s (defined as the complete absence of movement except for respiratory
movements). Evoked freezing behavior was analyzed by calculating the percentage of time
an animal spent freezing during a given phase of the test. During fear extinction, averages
were calculated by pooling freezing across 2 CS presentations if not indicated otherwise.

RNA extraction and quantification (qPCR)

In all experiments, mice were euthanized and brains were rapidly removed and placed on

ice. The dorsal and ventral hippocampus were dissected as reported in Jaszczyk et al.

2022. Briefly, the two hemispheres were separated and the brain was rotated to expose the

medial surface. The hippocampus was carefully exposed by gentle mechanical dissection,

detached from the surrounding tissue, and cleaned of residual white matter. The isolated

hippocampus was then divided into three portions—dorsal, intermediate, and ventral—

based on anatomical landmarks and millimeter measurements. Tissue samples were snap-

frozen in liquid nitrogen and stored at −80 °C until further processing. Tissue samples were

homogenized in the cell disruption buffer (Ambion). RNA was extracted by the addiction of

Phenol/guanidine-based QIAzol Lysis Reagent (Qiagen, cat. no. 79306). Chloroform was

added, and the samples were shaken for 15 s. The samples were left at 20–24°C for 3 min
and then centrifuged (12,000 g, 20 min, 4°C). The upper phase aqueous solution, containing
RNA, was collected in a fresh tube, and the RNA was precipitated by the addition of
isopropanol. Samples were mixed by vortexing, left at 20–24°C for 15 min, and then
centrifuged (12,000 g, 20 min, 4°C). The supernatant was discarded, and the RNA pellet
was washed in 75% ethanol by centrifugation (7500 g, 10 min, 4°C). The supernatant was
discarded, and the pellet was left to dry for at least 15 min; then, it was resuspended in
RNAse-free water. Total RNA concentrations were determined by NanoDrop
Spectrophotometer (Thermo Scientific 2000 C). RNA quality was analyzed through a gel
running (1% agarose). Total RNA was reverse transcribed using the QuantiTeck Reverse
Transcription Kit (Qiagen, cat. no. 205311), and miRNAs were reverse transcribed using the
TaqMan MicroRNA reverse transcription kit (Thermo Fisher, cat. no. 4366596). Gene
expression was analyzed by real-time PCR (Step one, Applied Biosystems). TaqMan
inventoried assays were used for miR-29a-3p (assay ID: 002112), sno234 (assay
ID:001234), and *Dnmt3a* (assay ID: Mm00432881_m1). TaqMan assay was used for
glyceraldehyde 3-phosphate dehydrogenase (Gapdh), GAPDH probe
ATCCCAGAGCTGAACGG, GADPH forward CAAGGCTGTGGGCAAGGT, and GADPH
reverse GGCCATGCCAGTGAGCTT. Quantitative values for cDNA amplification were
calculated from the threshold cycle number (Ct) obtained during the exponential growth of
the PCR products. The threshold was set automatically by the Step One software.
Expression levels were normalized using *Gapdh or sno234* as housekeeping. Relative gene
expression was calculated using the ΔΔCt method.

DNA extraction and validation

For the methylation analysis, we administered LNA-anti-miR29a to one hemisphere of seven
mice, while the other hemisphere received an injection of the scrambled sequence, serving
as an internal control. To mitigate potential lateralization effects, we randomized the side of
LNA injection. Seven days after the injection, tissues were collected and samples were
stored in an ultralow freezer (-80°C) until DNA extraction. DNA was isolated from snap-
frozen hippocampi using the QIAamp DNA Micro Kit (Qiagen) following the manufacturer's
protocol. DNA concentration was measured with NanoDrop 2000 (Thermo Scientific), using
1 µl of input DNA. Then, dsDNA concentration was quantified using Quant-iT Picogreen
dsDNA Assay (Thermo Scientific), and the distribution of DNA fragments was assessed by
TapeStation 4200 (Agilent).

Reduced Representation Bisulfite Sequencing (RRBS)

RRBS libraries were produced by Ovation RRBS Methyl-Seq with the TrueMethyl oxBS kit
(Tecan, Redwood City, CA, USA) according to the manufacturer's instructions. Briefly, 100
ng of genomic DNA was digested for 1h at 37°C with the methylation-insensitive restriction

enzyme MpsI. Then, fragments were ligated to methylated adapters and treated with
bisulfite to convert unmethylated cytosine into uracil. PCR amplification was then performed
to obtain the final DNA library. The resulting libraries were sequenced on the Illumina
NovaSeq6000 with 100 bp single-end sequencing with an average of 60 million reads per
sample.

RRBS data analysis

The sequencing reads were trimmed to remove the adapter and low-quality bases with Trim
Galore! (www.bioinformatics.babraham.ac.uk). Then trimmed reads were aligned to the
mouse reference genome mm39 with BSMAPz (github.com/zyndagj). Using the Python
script methratio.py in BSMAPz, methylation ratios were extracted from the mapping output.
Methylation analysis was performed using the R package methylKit (version 1.20.0) (Akalin
et al., 2012). Only CpGs covered by at least 10 reads and present in every sample were
retained for downstream analysis. The methylation ratio of each site was calculated by
dividing the number of reads called "C" by the total number of reads called either "C" or "T"
at the specific site. The methylation score for each CpG site is represented as a β-value
which ranges between 0 (unmethylated) and 1 (fully methylated). Differentially methylated
CpGs (DMCs) were detected using a logistic regression model based on a Chi-square test.
A false discovery rate (FDR) q value threshold of < 0.05 and methylation difference of ±5%
between treated and control groups was used to identify significant DMCs. The promoter
was defined as the region ±1kb from the transcription start site. CpG sites located on the X
and Y chromosomes were excluded from the methylation analysis due to a strong sex-
associated effect, which represented the main source of variability in the dataset. All
downstream analyses were therefore restricted to autosomal CpGs. Differential methylation
analysis was performed separately for each treatment group (anti-miR29a, anti-DNMT3a
and combined anti-miR29a + anti-DNMT3a) compared to control samples using methylKit.
Multivariate analysis was performed using partial least squares discriminant analysis (PLS–
DA) in mixOmics R package to assess sample clustering based on methylation profiles; the
first two components were used for visualization. To evaluate concordance of methylation
changes between treatment conditions, DMCpGs identified in the combined treatment were
compared to those identified in the anti-miR29a treatment alone. CpGs were classified into
four quadrants based on the direction of methylation change. A χ^2 test was performed to
assess deviation from a uniform distribution of concordant versus discordant CpGs. Non-
CpG (CH) methylation levels were also quantified. Per-animal Δ methylation values (treated
minus control hemisphere) were calculated and compared across treatment groups using
Welch's ANOVA followed by Bonferroni-corrected pairwise t-tests. Pathway analysis was
performed using the web tools WebGestalt (www.webgestalt.org). Pathway visualization
was performed using the R package Pathview to map methylation changes onto KEGG
pathways. The KEGG pathway ID mmu04724 (Glutamatergic synapse) was specifically
examined to visualize gene-level methylation changes following anti-miR29a treatment.

Sample preparation for proteomics analysis

Before lysis, a lysis buffer was added to each sample to reach final concentrations of 4% SDS, 100 mM HEPES (pH 8.5), and 50 mM DTT. Samples were then boiled at 95°C for 7 min and sonicated using a tweeter. Reduction was followed by alkylation with 200 mM iodoacetamide (IAA, final concentration 15 mM) for 30 min at room temperature in the dark. Samples were acidified with phosphoric acid (final concentration 2.5%), and seven times the sample volume of S-trap binding buffer was added (100 mM TEAB, 90% methanol). Samples were bound on a 96-well S-trap micro plate (Protifi) and washed three times with a binding buffer. Trypsin in 50 mM TEAB pH 8.5 was added to the samples (1 μg per sample) and incubated for 1 h at 47°C. The samples were eluted in three steps with 50 mM TEAB pH 8.5, elution buffer 1 (0.2% formic acid in water) and elution buffer 2 (50% acetonitrile and 0.2% formic acid). The eluates were dried using a speed vacuum centrifuge (Eppendorf Concentrator Plus, Eppendorf AG, Germany) and stored at -20° C. Before analysis, samples were reconstituted in MS Buffer (5% acetonitrile, 95% Milli-Q water, with 0.1% formic acid) and spiked with iRT peptides (Biognosys, Switzerland).

LC-MS Data independent analysis (DIA)

Peptides were separated in trap/elute mode using the nanoAcquity MClass Ultra-High Performance Liquid Chromatography system (Waters, Waters Corporation, Milford, MA,USA) equipped with a trapping (Waters nanoEase M/Z Symmetry C18, 5μm, 180 μm x 20 mm) and an analytical column (Waters nanoEase M/Z Peptide C18, 1.7μm, 75μm x 250mm). Solvent A was water and 0.1% formic acid, and solvent B was acetonitrile and 0.1% formic acid. 1 μl of the sample (~1 μg on column) was loaded with a constant flow of solvent A at 5 μl/min onto the trapping column. Trapping time was 6 min. Peptides were eluted via the analytical column with a constant flow of 0.3 μl/min. During the elution, the percentage of solvent B increased in a nonlinear fashion from 0–40% in 120 min. Total run time was 145 min. including equilibration and conditioning. The LC was coupled to an Orbitrap Exploris 480 (Thermo Fisher Scientific, Bremen, Germany) using the Proxeon nanospray source. The peptides were introduced into the mass spectrometer via a Pico-Tip Emitter 360-μm outer diameter × 20-μm inner diameter, 10-μm tip (New Objective) heated at 300 °C, and a spray voltage of 2.2 kV was applied. The capillary temperature was set at 300°C. The radio frequency ion funnel was set to 30%. For DIA data acquisition, full scan

mass spectrometry (MS) spectra with mass range 350–1650 m/z were acquired in profile

mode in the Orbitrap with resolution of 120,000 FWHM. The default charge state was set to

3+. The filling time was set at a maximum of 60 ms with a limitation of 3 × 10 6 ions. DIA

scans were acquired with 40 mass window segments of differing widths across the MS1

mass range. Higher collisional dissociation fragmentation (stepped normalized collision

energy; 25, 27.5, and 30%) was applied and MS/MS spectra were acquired with a resolution

of 30,000 FWHM with a fixed first mass of 200 m/z after accumulation of 3 × 10 6 ions or

after filling time of 35 ms (whichever occurred first). Data were acquired in profile mode. For

data acquisition and processing of the raw data Xcalibur 4.3 (Thermo) and Tune version 2.0

were used.

Proteomic data processing

DIA raw data were analyzed using the directDIA pipeline in Spectronaut v.18 (Biognosys,
Switzerland) with BGS settings besides the following parameters: Protein LFQ method =
QUANT 2.0, Proteotypicity Filter = Only protein group specific, Major Group Quantity =
Median peptide quantity, Minor Group Quantity = Median precursor quantity, Data Filtering =
Qvalue, Normalizing strategy = Local Normalization. The data were searched against a
UniProt (Mus musculus, v. 160106, 16.748 entries) and a contaminants (247 entries)
database. The identifications were filtered to satisfy FDR of 1 % on peptide and protein
level. Relative protein quantification was performed in Spectronaut using a pairwise t-test
performed at the precursor level followed by multiple testing correction according to
Benjamini-Hochberg.

RNA-Seq library preparation

Sequencing of RNA samples was done using Illumina's next-generation sequencing
methodology . In detail, quality check and quantification of total RNA was done using the
Agilent Bioanalyzer 2100 in combination with the RNA 6000 pico kit (Agilent Technologies,
5067-1513). Total RNA library preparation was done by introducing 500 ng total RNA into
Illumina's NEBNext Ultra II directional mRNA (UMI) kit (NEB, E7760S), following the
manufacturer's instructions. The quality and quantity of all libraries were checked using
Agilent's Bioanalyzer 2100 and DNA 7500 kit (Agilent Technologies, 5067-1506).RNAseq
analysis. All libraries were sequenced on a NovaSeq6000 SP 100 cycles v1.5.
Corresponding UMI file was used to add UMI sequences to the reads names of each
sample R1 file using UMI-tools extract v1.1.1 [73] to enable duplicate removal in the later
steps. Trimmomatic v0.36 [74] was applied to the reads to remove adapters and low quality
sequences   using   custom   adapter   fasta   file   and   the   following   options

adapters_3tools.fa:2:30:10:6 SLIDINGWINDOW:3:25 MINLEN:25. Trimmed reads were
then mapped onto the *Mus musculus* GRCm39 genome primary assembly from Ensembl
release 111 [75] using STAR v2.7.10a [76] and -outSAMtype BAM SortedByCoordinate -
outSAMmultNmax 1 --outSJfilterReads Unique --outSAMstrandField intronMotif --
alignIntronMax 100000 --outFilterMismatchNoverLmax 0.04 options to generate sorted bam
files. The resulting bam file was indexed using samtools v1.19.2-23-g1a63877 [77] and
duplicates were removed from it using UMI-tools dedup v1.1.1 [78]. In order to get gene-
level counts for each sample featureCounts v2.0.3 [79] was used with mouse GRCm39 GTF
annotation from Ensembl release 111 [80] and -s 2 flag. Counts from individual samples
were then combined into a single counts matrix for use in downstream analysis tools.

Differential expression Analysis

Differential expression analysis was performed using DESeq2 [81]. Exploratory data
analysis was performed with PCA to identify possible outliers. Counts were normalized by
variance stabilizing transformation (vst) and adaptive fold-change shrinkage (apeglm).
Differential expression analysis was performed with default parameters. Since samples are
paired, the following design was implemented: design = ~ Subject + Cond where Cond has
two categorical values = "Scramble" and "anti-miR29".

Statistical analysis

All the statistical analyses were performed using GraphPad Prism 8. The Shapiro–Wilk test
was used to assess the normality of the data. For datasets that deviated significantly from
normality, appropriate non-parametric tests were applied. For the two-way mixed ANOVA,
this parametric approach was retained because it efficiently assesses main effects and
interactions for both within- and between-subject factors. Although the sample sizes were
slightly unequal between groups, ANOVA is generally robust to such minor imbalances,
particularly when variances are similar. Moreover, the Geisser–Greenhouse correction was
applied to account for violations of sphericity, ensuring valid F-tests. Analyses were followed
by appropriate post hoc tests. Significance was set at $P<0.05$ for all tests. Error bars
represent s.e.m. in all figures.

References

1.   Bartel DP (2018) Metazoan MicroRNAs. Cell 173:20–51

2.   Rajman M, Schratt G (2017) MicroRNAs in neural development: from master regulators
to fine-tuners. Development 144:2310–2322

3.   Lippi G, Fernandes CC, Ewell LA, et al (2016) MicroRNA-101 Regulates Multiple
Developmental Programs to Constrain Excitation in Adult Neural Networks. Neuron
92:1337–1351

4.   Kole AJ, Swahari V, Hammond SM, Deshmukh M (2011) miR-29b is activated during
neuronal maturation and targets BH3-only genes to restrict apoptosis. Genes Dev
25:125–130

5.   Dalgaard LT, Sørensen AE, Hardikar AA, Joglekar MV (2022) The microRNA-29 family:
role in metabolism and metabolic disease. Am J Physiol Cell Physiol 323:C367–C377

6.   Ma F, Xu S, Liu X, et al (2011) The microRNA miR-29 controls innate and adaptive
immune responses to intracellular bacterial infection by targeting interferon-γ. Nat
Immunol 12:861–869

7.   Yee Mon KJ, Zhu H, Daly CWP, et al (2021) MicroRNA-29 specifies age-related
differences in the CD8+ T cell immune response. Cell Rep 37:109969

8.   Ripa R, Dolfi L, Terrigno M, et al (2017) MicroRNA miR-29 controls a compensatory
response to limit neuronal iron accumulation during adult life and aging. BMC Biol 15:9

9.   Wagner V, Kern F, Hahn O, et al (2024) Characterizing expression changes in
noncoding RNAs during aging and heterochronic parabiosis across mouse tissues. Nat
Biotechnol 42:109–118

10.   Mazziotti R, Baroncelli L, Ceglia N, et al (2017) Mir-132/212 is required for maturation of
binocular matching of orientation preference and depth perception. Nat Commun
8:15488

11.   Somel M, Guo S, Fu N, et al (2010) MicroRNA, mRNA, and protein expression link
development and aging in human and macaque brain. Genome Res 20:1207–1218

12.   Baumgart M, Groth M, Priebe S, et al (2012) Age-dependent regulation of tumor-related
microRNAs in the brain of the annual fish Nothobranchius furzeri. Mech Ageing Dev
133:226–233

13.   Wingo AP, Wang M, Liu J, et al (2022) Brain microRNAs are associated with variation in
cognitive trajectory in advanced age. Transl Psychiatry 12:47

14.   Swahari V, Nakamura A, Hollville E, et al (2024) miR-29 is an important driver of aging-
related phenotypes. Commun Biol 7:1055

15.   Napoli D, Lupori L, Mazziotti R, et al (2020) MiR-29 coordinates age-dependent
plasticity brakes in the adult visual cortex. EMBO Rep 21:e50431

16.   Mei Z, Liu J, Schroeder JP, et al (2024) Lowering Hippocampal miR-29a Expression
Slows Cognitive Decline and Reduces Beta-Amyloid Deposition in 5×FAD Mice. Mol
Neurobiol 61:3343–3356

17. Swahari V, Nakamura A, Hollville E, et al (2021) MicroRNA-29 is an essential regulator
of brain maturation through regulation of CH methylation. Cell Rep 35:108946

18. Fabbri M, Garzon R, Cimmino A, et al (2007) MicroRNA-29 family reverts aberrant
methylation in lung cancer by targeting DNA methyltransferases 3A and 3B. Proc Natl
Acad Sci U S A 104:15805–15810

19. Benetti R, Gonzalo S, Jaco I, et al (2008) A mammalian microRNA cluster controls DNA
methylation and telomere recombination via Rbl2-dependent regulation of DNA
methyltransferases. Nat Struct Mol Biol 15:998

20. Liu L, van Groen T, Kadish I, Tollefsbol TO (2009) DNA methylation impacts on learning
and memory in aging. Neurobiol Aging 30:549–560

21. Ciccarone F, Tagliatesta S, Caiafa P, Zampieri M (2018) DNA methylation dynamics in
aging: how far are we from understanding the mechanisms? Mech Ageing Dev 174:3–
17

22. Gu H, Smith ZD, Bock C, et al (2011) Preparation of reduced representation bisulfite
sequencing libraries for genome-scale DNA methylation profiling. Nat Protoc 6:468–481

23. Tognini P, Napoli D, Tola J, et al (2015) Experience-dependent DNA methylation
regulates plasticity in the developing visual cortex. Nat Neurosci 18:956–958

24. Miller CA, Gavin CF, White JA, et al (2010) Cortical DNA methylation maintains remote
memory. Nat Neurosci 13:664–666

25. Lubin FD, Roth TL, Sweatt JD (2008) Epigenetic regulation of BDNF gene transcription
in the consolidation of fear memory. J Neurosci 28:10576–10586

26. Miller CA, Sweatt JD (2007) Covalent modification of DNA regulates memory formation.
Neuron 53:857–869

27. Lugli G, Torvik VI, Larson J, Smalheiser NR (2008) Expression of microRNAs and their
precursors in synaptic fractions of adult mouse forebrain. J Neurochem 106:650–661

28. Wang W, Kwon EJ, Tsai L-H (2012) MicroRNAs in learning, memory, and neurological
diseases. Learn Mem 19:359–368

29. Reh RK, Dias BG, Nelson CA 3rd, et al (2020) Critical period regulation across multiple
timescales. Proc Natl Acad Sci U S A 117:23242–23251

30. Putignano E, Lonetti G, Cancedda L, et al (2007) Developmental downregulation of
histone posttranslational modifications regulates visual cortical plasticity. Neuron
53:747–759

31. Morita S, Horii T, Kimura M, et al (2013) miR-29 represses the activities of DNA
methyltransferases and DNA demethylases. Int J Mol Sci 14:14647–14658

32. Oliveira AMM, Hemstedt TJ, Bading H (2012) Rescue of aging-associated decline in
Dnmt3a2 expression restores cognitive abilities. Nat Neurosci 15:1111–1113

33. Braasch DA, Corey DR (2001) Locked nucleic acid (LNA): fine-tuning the recognition of
DNA and RNA. Chem Biol 8:1–7

34. Smith CIE, Zain R (2019) Therapeutic Oligonucleotides: State of the Art. Annu Rev
Pharmacol Toxicol 59:605–630

35. Khvorova A, Watts JK (2017) The chemical evolution of oligonucleotide therapies of
clinical utility. Nat Biotechnol 35:238–248

36. Bangasser DA, Waxler DE, Santollo J, Shors TJ (2006) Trace conditioning and the
hippocampus: the importance of contiguity. J Neurosci 26:8702–8706

37. Meissner A, Gnirke A, Bell GW, et al (2005) Reduced representation bisulfite
sequencing for comparative high-resolution DNA methylation analysis. Nucleic Acids
Res 33:5868–5877

38. Rattiner LM, Davis M, French CT, Ressler KJ (2004) Brain-derived neurotrophic factor
and tyrosine kinase receptor B involvement in amygdala-dependent fear conditioning. J
Neurosci 24:4796–4806

39. Dias BG, Goodman JV, Ahluwalia R, et al (2014) Amygdala-dependent fear memory
consolidation via miR-34a and Notch signaling. Neuron 83:906–918

40. Steinfeld I, Navon R, Ach R, Yakhini Z (2013) miRNA target enrichment analysis reveals
directly active miRNAs in health and disease. Nucleic Acids Res 41:e45

41. Szklarczyk D, Kirsch R, Koutrouli M, et al (2023) The STRING database in 2023:
protein-protein association networks and functional enrichment analyses for any
sequenced genome of interest. Nucleic Acids Res 51:D638–D646

42. Zhang B, Gaiteri C, Bodea L-G, et al (2013) Integrated systems approach identifies
genetic nodes and networks in late-onset Alzheimer's disease. Cell 153:707–720

43. Xin W, Kaneko M, Roth RH, et al (2024) Oligodendrocytes and myelin limit neuronal
plasticity in visual cortex. Nature 633:856–863

44. McGee AW, Yang Y, Fischer QS, et al (2005) Experience-driven plasticity of visual
cortex limited by myelin and Nogo receptor. Science 309:2222–2226

45. Wingo TS, Yang J, Fan W, et al (2020) Brain microRNAs associated with late-life
depressive symptoms are also associated with cognitive trajectory and dementia. NPJ
Genom Med 5:6

46. Ugalde AP, Ramsay AJ, de la Rosa J, et al (2011) Aging and chronic DNA damage
response activate a regulatory pathway involving miR-29 and p53. EMBO J 30:2219–
2232

47. Barrett RM, Wood MA (2008) Beyond transcription factors: the role of chromatin
modifying enzymes in regulating transcription required for memory. Learn Mem 15:460–
467

48. Oliveira AMM, Hemstedt TJ, Freitag HE, Bading H (2016) Dnmt3a2: a hub for
enhancing cognitive functions. Mol Psychiatry 21:1130–1136

49. Day JJ, Sweatt JD (2010) DNA methylation and memory formation. Nat Neurosci
13:1319–1323

50. Ferry L, Fournier A, Tsusaka T, et al (2017) Methylation of DNA Ligase 1 by G9a/GLP
Recruits UHRF1 to Replicating DNA and Regulates DNA Methylation. Mol Cell 67:550–
565.e5

51. Day K, Waite LL, Thalacker-Mercer A, et al (2013) Differential DNA methylation with age
displays both common and dynamic features across human tissues that are influenced

by CpG landscape. Genome Biol 14:R102

52. Jung M, Pfeifer GP (2015) Aging and DNA methylation. BMC Biol 13:7

53. Oliveira AMM (2016) DNA methylation: a permissive mark in memory formation and
maintenance. Learn Mem 23:587–593

54. Morris MJ, Adachi M, Na ES, Monteggia LM (2014) Selective role for DNMT3a in
learning and memory. Neurobiology of learning and memory 0:30

55. Gong Z, Zhou Q (2018) Dnmt3a in the dorsal dentate gyrus is a key regulator of fear
renewal. Scientific Reports 8:5093

56. Brito DVC, Esteves F, Rajado AT, et al (2023) Assessing cognitive decline in the aging
brain: lessons from rodent and human studies. npj Aging 9:23

57. Kupke J, Klimmt J, Mudlaff F, et al (2024) Dnmt3a1 regulates hippocampus-dependent
memory via the downstream target Nrp1. Neuropsychopharmacology 49:1528–1539

58. Gulmez Karaca K, Kupke J, Brito DVC, et al (2020) Neuronal ensemble-specific DNA
methylation strengthens engram stability. Nat Commun 11:639

59. Jones PA (1999) The DNA methylation paradox. Trends Genet 15:34–37

60. Jones PA (2012) Functions of DNA methylation: islands, start sites, gene bodies and
beyond. Nat Rev Genet 13:484–492

61. Guruharsha KG, Kankel MW, Artavanis-Tsakonas S (2012) The Notch signalling
system: recent insights into the complexity of a conserved pathway. Nat Rev Genet
13:654–666

62. Artavanis-Tsakonas S, Rand MD, Lake RJ (1999) Notch signaling: cell fate control and
signal integration in development. Science 284:770–776

63. Alberi L, Hoey SE, Brai E, et al (2013) Notch signaling in the brain: in good and bad
times. Ageing Res Rev 12:801–814

64. Brai E, Alina Raio N, Alberi L (2016) Notch1 hallmarks fibrillary depositions in sporadic
Alzheimer's disease. Acta Neuropathol Commun 4:64

65. Yamada K, Mizuno M, Nabeshima T (2002) Role for brain-derived neurotrophic factor in
learning and memory. Life Sci 70:735–744

66. Tyler WJ, Alonso M, Bramham CR, Pozzo-Miller LD (2002) From acquisition to
consolidation: on the role of brain-derived neurotrophic factor signaling in hippocampal-
dependent learning. Learn Mem 9:224–237

67. Minichiello L, Korte M, Wolfer D, et al (1999) Essential role for TrkB receptors in
hippocampus-mediated learning. Neuron 24:401–414

68. Mu JS, Li WP, Yao ZB, Zhou XF (1999) Deprivation of endogenous brain-derived
neurotrophic factor results in impairment of spatial learning and memory in adult rats.
Brain Res 835:259–265

69. Alonso M, Vianna MRM, Depino AM, et al (2002) BDNF-triggered events in the rat
hippocampus are required for both short- and long-term memory formation.
Hippocampus 12:551–560

70. Steiner DF, Thomas MF, Hu JK, et al (2011) MicroRNA-29 regulates T-box transcription
factors and interferon-γ production in helper T cells. Immunity 35:169–181

71. Liston A, Papadopoulou AS, Danso-Abeam D, Dooley J (2012) MicroRNA-29 in the
adaptive immune system: setting the threshold. Cell Mol Life Sci 69:3533–3541

72. Kluever V, Russo B, Mandad S, et al (2022) Protein lifetimes in aged brains reveal a
proteostatic adaptation linking physiological aging to neurodegeneration. Sci Adv
8:eabn4437

73. Smith T, Heger A, Sudbery I (2017) UMI-tools: modeling sequencing errors in Unique
Molecular Identifiers to improve quantification accuracy. Genome Res 27:491–499

74. Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30:2114–2120

75. Harrison PW, Amode MR, Austine-Orimoloye O, et al (2024) Ensembl 2024. Nucleic
Acids Res 52:D891–D899

76. Dobin A, Davis CA, Schlesinger F, et al (2013) STAR: ultrafast universal RNA-seq
aligner. Bioinformatics 29:15–21

77. Li H, Handsaker B, Wysoker A, et al (2009) The Sequence Alignment/Map format and
SAMtools. Bioinformatics 25:2078–2079

78. Smith T, Heger A, Sudbery I (2017) UMI-tools: modeling sequencing errors in Unique
Molecular Identifiers to improve quantification accuracy. Genome Res 27:491–499

79. Liao Y, Smyth GK, Shi W (2014) featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. Bioinformatics 30:923–930

80. Harrison PW, Amode MR, Austine-Orimoloye O, et al (2024) Ensembl 2024. Nucleic
Acids Res 52:D891–D899

81. Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol 15:550