1   **SARS-CoV-2 ORF3a blocks lysosomal cholesterol egress by disrupting VPS39-regulated**

2   **NPC2 trafficking and BMP metabolism**

3   Baley A. Goodson[1], Valeria Montenegro Vazquez[1], Aliza Doyle[1], Oralia M. Kolaczkowski[1], Rui

4   Liu[2], Jingyue Jia[3], Morié Ishida[4], Hu Wang[5], Xianlin Han[5], Chunyan Ye[3], Alison M. Kell[1], Steven

5   B. Bradfute[3], Monica Rosas Lemus[1], Jing Pu[1],*

6

7   [1]Department of Molecular Genetics and Microbiology, University of New Mexico Health

8   Sciences Center, Albuquerque, NM, USA.

9   [2]Department of Pharmaceutical Sciences, College of Pharmacy, University of New Mexico

10   Health Sciences Center, Albuquerque, NM, USA.

11   [3]Department of Internal Medicine, University of New Mexico Health Sciences Center,

12   Albuquerque, NM, USA.

13   [4]Neurosciences and Cellular and Structural Biology Division, Eunice Kennedy Shriver National

14   Institute of Child Health and Human Development, National Institutes of Health, Bethesda,

15   MD, USA.

16   [5]Department of Medicine, UT Health San Antonio Long School of Medicine, San Antonio, TX,

17   USA.

18   *Correspondence: jpu@salud.unm.edu; T: 505-272-8890

19   Run title: ORF3a blocks lysosomal cholesterol via NPC2 and BMP pathways

Abstract

Cholesterol homeostasis relies on lysosomes, which release free cholesterol from degraded

lipids. We show that SARS-CoV-2 blocks lysosomal cholesterol export through its protein

ORF3a. ORF3a binds the HOPS subunit VPS39, and disrupting this interaction restores

cholesterol trafficking. Two mechanisms underlie this defect. First, ORF3a–VPS39 interaction

traps the sorting receptor CI-MPR and the retrieval complex retromer in endosomes/lysosomes,

impairing trafficking of the cholesterol transporter NPC2. Retromer deletion reproduced these

defects. Second, ORF3a reduces bis(monoacylglycerol)phosphates (BMPs), lysosomal lipids

required for cholesterol export. Lipidomics and proteomics revealed altered metabolism of BMP

precursors, mitochondrial phosphatidylglycerols (PGs), and reduced mitochondrial proteins at

lysosomes. ORF3a–VPS39 interaction decreased lysosome–mitochondrion membrane contact

sites (MCS), excluding autophagy or mitochondrion-derived vesicles as routes for PG transfer.

VPS39 deletion decreased the MCS and BMPs. These findings identify VPS39 as a regulator of

NPC2 trafficking and BMP biosynthesis and reveal that ORF3a disrupts both pathways to block

cholesterol egress.

INTRODUCTION

Cholesterol plays a vital role in maintaining cellular structure and functions. Thus, cells tightly regulate cholesterol levels and distribution through multiple pathways. Cholesterol pathways are not only disturbed in metabolic disorders but also hijacked in infectious diseases. COVID-19, caused by the SARS-CoV-2 virus, has been associated with significant disruptions in cholesterol metabolism[1–5]. In cases of long COVID, an increased risk of dyslipidemia and cardiovascular diseases is marked by elevated low-density lipoprotein (LDL) cholesterol and reduced high-density lipoprotein (HDL) cholesterol levels[4,5].

Lysosomes, the degradation organelles, receive and break down internalized lipoprotein particles and cellular macrolipids delivered via endocytosis and autophagy, respectively, releasing free cholesterol in the process[6]. The released cholesterol is then distributed to other cellular compartments to support structural and metabolic demands[7,8]. The cholesterol delivered to the endoplasmic reticulum (ER) serves as a signal to regulate the expression of genes that control cholesterol synthesis and uptake[9]. The cholesterol supplied to the plasma membrane is the major source for maintaining the local cholesterol pool that modulate plasma membrane fluidity and stability[10]. SARS-CoV-2 exploits plasma membrane cholesterol to enhance infectivity[11], as evidenced by a reduced binding between the viral spike protein and its host receptor after plasma membrane cholesterol depletion[12]. Therefore, the lysosomal cholesterol transport is central to cellular cholesterol homeostasis and may also play a role in SARS-CoV-2 infection.

Lysosomal cholesterol egress is the first step of cholesterol distribution. This process is primarily mediated by the lysosomal transmembrane cholesterol transporter Niemann-Pick C1 (NPC1) and the lumenal cholesterol-binding protein Niemann-Pick C2 (NPC2), which act cooperatively to transfer cholesterol out of lysosomes[8]. Additional lysosomal proteins, including the lumenal saposins and membrane proteins LAMP2[13,14], also contribute to cholesterol egress[8]. The proper trafficking and localization of newly synthesized lysosome transmembrane proteins depend on sorting signals and clathrin coats[15]. In contrast, the lysosome lumenal proteins are directed by sorting receptors, such as the mannose-6-phosphate receptors (M6PRs)[15]. These receptors deliver lysosome lumenal proteins from the trans-Golgi network (TGN) to endosomes and are subsequently recycled back to the TGN. This recycling requires retromer complexes at endosomes and lysosomes[16–19] and Golgi-associated retrograde

protein (GARP) complex at the TGN[20]. Disruption of these transport pathways by impairing

M6PR, HOPS or GARP functions compromises the delivery of newly synthesized proteins to

lysosomes and results in accumulation of cholesterol[18,21,22]. Such lysosomal cholesterol

sequestration profoundly disrupts lipid homeostasis and impairs cell functions, a defining

feature of Niemann-Pick disease type C, caused by mutations in NPC1 or NPC2[23,24]. Beyond

genetic disorders, certain viruses exploit lysosomal cholesterol transport pathways to complete

their life cycles. For instance, NPC1 is hijacked by Ebola virus for their infection[25].

In addition to protein-mediated cholesterol transport, the phospholipid

bis(monoacylglycero)phosphates (BMPs, also called lysobisphosphatidic acids or LBPAs), play

an important role in lysosomal cholesterol egress. BMPs are highly enriched in the late

endosome and lysosome membrane, where they contribute to lysosome sorting and

degradation functions[9,26–28]. BMP has also been implicated in viral infection by promoting viral

fusion with late endosome membranes[29–32]. Beyond these function, BMPs directly interact with

NPC2 to promote cholesterol egress in an NPC1-independent way, as supplementation with

BMPs rescues cholesterol sequestration in NPC1-defecient cells but has no effect in NPC2-

defecient cells[33,34]. The synthesis and turnover of BMPs depend on multiple lysosomal

enzymes that convert the mitochondrial phospholipids phosphatidylglycerols (PGs) into BMPs

and degrade BMPs through hydrolysis[35–39].

In this study, we show that SARS-CoV-2 infection drives cholesterol sequestration in lysosomes,

primarily through the viral protein ORF3a. ORF3a binds the HOPS subunit VPS39, disrupting

lysosomal lumenal protein trafficking by retaining the cation-independent mannose 6-

phosphate receptor (CI-MPR) and retromer complex at endosomes and lysosomes. In addition,

ORF3a–VPS39 interaction impairs lysosome-mitochondrion membrane contact sites, potentially

limiting lipid transport from mitochondria to lysosomes required for BMP synthesis. By

simultaneously disturbing these two VPS39-dependent pathways, ORF3a effectively sequesters

cholesterol within lysosomes, revealing a mechanistic link between SARS-CoV-2 infection and

lysosomal lipid dysregulation.

RESULTS

Increased lysosomal cholesterol in SARS-CoV-2-infected cells

To examine how SARS-CoV-2 infection alters host cholesterol at cellular levels, we infected

human lung carcinoma epithelial cells A549, stably expressing the viral entry receptor ACE2 of

human species (A549-hACE2), with the Washington strain of SARS-CoV-2. We confirmed

infection in approximately 80% of cells by immunostaining for double-stranded RNA (dsRNA)

(Figs. 1A & S1A). Using filipin staining to detect free cholesterol, we observed a marked

increase in filipin signal in infected cells, appearing as punctate structures, compared to mock-

treated cells (Fig. 1A). Similar observations were made in a different ACE2-overexpressing A549

cell line (Fig. S1B). Co-staining with an antibody against late endosomal and lysosomal

membrane protein LAMP1 revealed that the filipin puncta colocalized with LAMP1-positive

vesicles (Fig. 1A), indicating that cholesterol was sequestrated in the late endosomes and

lysosomes. Due to the similarity of these two organelles, we refer them to lysosomes hereafter.

Quantification of confocal images showed that lysosomal cholesterol levels significantly

increased at 24- and 48-hours post-infection (Fig. 1B). Although filipin puncta were also present

at 72 hours post-infection, mock cells showed elevated filipin signal, likely due to cell responses

associated with high cell density[40] (Fig. 1B). Further analysis of over 1,500 infected cells using a

high-content imaging system confirmed a consistent increase in filipin-positive puncta (Fig. 1C).

To validate this cholesterol alteration, we infected monkey kidney cells Vero E6 with SARS-

CoV-2, resulting approximately 80% infection rate (Fig. S1C). High-content imaging revealed

that the infected cells displayed a significant 50–60% increase in cholesterol relative to LAMP1,

compared with mock-treated cells, at 18 and 24 hours post-infection. (Figs. 1D, 1E, S1D),

confirming that SARS-CoV-2 infection induces an increase in lysosomal cholesterol. This change

persisted beyond the viral eclipse phase (~10 hours)[41]. Gas chromatography-mass

spectrometry analysis of total cholesterol levels in mock and infected A549 cells showed no

significant difference (Fig. 1F), suggesting that SARS-CoV-2 infection mainly alters cholesterol

distribution within cells, possibly due to impaired lysosomal cholesterol transport.

Viral protein ORF3a is responsible for increased lysosomal cholesterol

To identify the specific viral proteins responsible for lysosomal cholesterol sequestration, we

transfected A549-hACE2 cells with 28 plasmids individually, which encode SARS-CoV-2

proteins, and quantified filipin puncta 24 hours post-transfection. Most viral proteins (24 out of

28) did not show a significant effect on cholesterol distribution. NSP10 and NSP13 had modest

effects, ORF10 doubled the filipin signal, and ORF3a produced the strongest increase in filipin

puncta (Fig. 2A). ORF3a, expressed from a SARS-CoV-2 subgenomic RNA, is an accessory

protein. Unlike NSP10, NSP13, or ORF10, which were predominantly cytoplasmic (Fig. S2),

ORF3a displayed punctate or halo-like structures that partially co-localized with LAMP2 (Fig.

2B), consistent with previous reports of ORF3a localization to endosomes/lysosomes[42–44].

Remarkably, ORF3a-positive lysosomes appeared swollen and were filled with filipin (Fig. 1B,

arrow in enlarged frame 1), in contrast to ORF3a-negative lysosomes in the same cells, which

remained small and displayed minimal filipin signal (Fig. 1B, arrow in enlarged frame 2). This

suggests that ORF3a, particularly when localized on lysosomes, plays a critical role in

cholesterol sequestration.

To further validate this observation, we transfected HeLa cells with an ORF3a plasmid tagged

with a V5 epitope. ORF3a-V5 also formed punctate or halo structures, partially co-localized

with LAMP2, and significantly increased filipin signal (Fig. 2C,D), demonstrating that ORF3a

enhances lysosomal cholesterol accumulation across different cell types. Additionally, we

included ORF3a from SARS-CoV, which shares over 70% sequence identity with SARS-CoV-2

ORF3a, as a comparison. Interestingly, while SARS-CoV ORF3a also displayed a punctate

distribution, it did not increase filipin (Fig. 2C,D), suggesting a specificity in cholesterol

sequestration by SARS-CoV-2 ORF3a.

Together, these results demonstrate that SARS-CoV-2 infection sequesters host lysosomal

cholesterol primarily through the action of ORF3a. Expression of ORF3a alone is sufficient to

drive cholesterol accumulation in lysosomes, likely due to its lysosomal localization. The

differential effects on lysosomal cholesterol between ORF3a proteins from SARS-CoV and

SARS-CoV-2 may offer insights into the underlying mechanisms.

The Structural difference between SARS-CoV and SARS-CoV-2 ORF3a

Previous studies revealed that SARS-CoV-2 ORF3a, but not SARS-CoV ORF3a, interacts with

the host protein VPS39[42–45] (Figs. 3A & S3B), a subunit of the HOPS complex. Our recent work

showed that the HOPS complex is essential for lysosomal cholesterol egress, using VPS39

CRISPR-knockout (KO) cells[18]. SARS-CoV-2 infection did not downregulate the expression of

VPS39 or VPS11, another HOPS subunit (Fig. S3A). These results suggest that SARS-CoV-2

ORF3a disrupts lysosomal cholesterol egress by inhibiting VPS39 functions. To test this, we

aimed to disrupt the ORF3a-VPS39 interaction by introducing specific point mutations.

Miao et al. identified the residues S171 and W193 in SARS-CoV-2 ORF3a as crucial for its

interaction with VPS39, as mutating these residues to those in SARS-CoV ORF3a abolished the

interaction[43]. Structural analysis of SARS-CoV-2 ORF3a[46,47] revealed that W193, Y184, S171,

and H182 cluster near each other (Fig. 3B, upper), with W193 and Y184 forming a hydrophobic

surface likely involved in protein-protein interactions, while S171 and H182 are positioned

outside this hydrophobic surface (Fig. 3B, lower). In our alanine mutation screen, we found that

W193A and Y184A mutations weakened or disrupted the interaction, whereas S171A or H182A

mutations did not (Figs. 3C & S3C,D). This indicates that W193 and Y184, but not S171 or H182,

are essential for binding. Further structural analysis indicates that the hydrophobic surface

formed by W193 and Y184 (7.7 Å) in SARS-CoV-2 ORF3a is occluded in SARS-CoV ORF3a due

to a hydrogen bond between R193 and E171 (2.6 Å) and Y184 (2.8 Å), resulting a "closed

conformation" (Fig. 3B, upper). This model is supported by the observation that the S171E

mutation impairs the interaction[43], likely due to the large hydrophilic side chain of glutamic

acid obstructing the hydrophobic surface formed by W193 and Y184, thereby blocking the

interaction. Collectively, these findings suggest that W193 and Y184 form the primary binding

site, while S171 and H182 contribute to an optimal binding environment.

Given that both W193 and Y184 are aromatic amino acids, we hypothesized that their aromatic

rings play a role in the interaction. To test this, we mutated W193 to various amino acids and

evaluated the effects on binding. Replacing W193 with another hydrophobic aromatic amino

acid, such as phenylalanine, preserved the interaction. However, substituting W193 with a

charged aromatic amino acid, like histidine, or with a hydrophobic but non-aromatic amino

acid, like alanine, reduced binding. Furthermore, replacing W193 with a charged, non-aromatic

amino acid, such as lysine or aspartic acid, nearly abolished the interaction (Fig. 3D). These

results suggest that both the aromatic ring and hydrophobicity of W193 are critical for the

binding of SARS-CoV-2 ORF3a to VPS39.

ORF3a-VPS39 interaction interrupts NPC2 trafficking

With the understanding of structural basis of the interaction interface, we chose W193A

mutation to block the interaction between ORF3a and VPS39. To minimize potential artifacts

from overexpression, we individually introduced SARS-CoV ORF3a, SARS-CoV-2 ORF3a, the

SARS-CoV-2 ORF3a W193A mutant, and a linker peptide that serves as a control into HeLa-Flp-

In cells to generate tetracycline-inducible protein-expression cell lines. This system controls

protein expression levels via a pre-engineered recombination site in the cells[48,49]. Likely due to

low protein expression, we could not detect the ORF3a proteins by immunofluorescence (Fig.

S4A), but immunoblotting showed expected bands in response to induction by the tetracycline

analog doxycycline (Fig. 4A). Using immunoprecipitation, we confirmed that, in the Flp-In cells,

CoV-2 ORF3a interacted with endogenous VPS39, and the W193A mutation abolished this
interaction (Fig. 4B). High-content imaging quantification revealed that doxycycline-induce
expression of ORF3a increased filipin signaling by 50-60% compared to the control cells (Figs.
4C,D & S4B). Importantly, the W193A mutant significantly reduced ORF3a-induced cholesterol
accumulation (Figs. 4C,D & S4B), demonstrating that the ORF3a-VPS39 interaction is
responsible for the cholesterol defect.
212
Previous research has shown that ORF3a expression increases lysosome pH[42] and
compromises lysosomal membrane integrity[50], raising the question of whether lysosomal
membrane damage or dysfunction leads to cholesterol sequestration. To investigate this, we
permeabilized lysosomal membranes using the lysosomotropic agent Leu-Leu-O-Me (LLOMe)
and confirmed membrane damage with the marker galectin-3[51] (Fig. S4C). Elevated lysosomal
cholesterol was detected in the LLOMe treatment as well as in the treatment by lysosome
deacidifying reagents, such as the ionophore monensin or V-ATPase inhibitors bafilomycin and
chloroquine (Fig. S4D). These results showed that lysosome damage and pH increase could
sequestrate cholesterol. However, we did not detect galectin-3 puncta in the Flp-in cells (Fig.
S4E), suggesting that ORF3a does not markedly damage lysosomes in these cells, and that there
are other mechanisms that interrupt cholesterol independent of lysosome damage.
224
We next examined the lysosomal cholesterol transporters NPC proteins in the Flp-In cells.
Immunoblotting the endogenous proteins did not show an obvious difference in NPC protein
levels between the four Flp-in cell lines (Fig. S4F) or between mock and infected cells (Fig. S4G).
Immunofluorescence confocal microscopy revealed that NPC1 colocalized with LAMP2 (Fig.
4E), and no difference of Pearson correlation coefficient was detected by high-content imaging
quantification (Fig. 4F). In contrast, a mislocalization of NPC2 from the lysosomes was observed
in the CoV-2-ORF3a cells (Fig. 4G), reproduced in a quantitative measurement of Pearson
correlation coefficient between NPC2 and LAMP1 (Fig. 4H). This mislocalization was largely
rescued by the W193A mutant (Fig. 4G,H). Further analysis of protein secretion revealed an
increased level of NPC2 and another lysosomal lumenal protein, cathepsin D, in the culture
media of CoV-2-ORF3a cells (Fig. 4I,J). These results motivated us to test whether lysosomal
transmembrane proteins, such as LAMPs, can localize to the lysosomes. Distinct from lysosomal
transmembrane proteins that are transported from TGN to lysosomes through biosynthesis
pathway, Ragulator complex is associated with lysosome membrane through lipidation of its
subunit LAMTOR1, which scaffolds other subunits. By measuring the colocalization between
LAMP2 and the Ragulator subunit LAMTOR4, we did not see a difference between the Flp-In

cells (Fig. S4H,I). Together, these results indicated that ORF3a induces a trafficking defect of

lysosome lumenal proteins, likely due to a dysregulation of protein transport between TGN and

endosomes.

ORF3a disturbs the endosome-to-TGN transport

The NPC2 trafficking defect observed in CoV-2-ORF3a cells resembled what was observed in

VPS39-deleted cells in our previous study, in which the sorting receptor for lysosomal lumenal

proteins, CI-MPR, is abnormally degraded in lysosomes[18]. In CoV-2-ORF3a cells, CI-MPR

protein level did not reduce (Fig. S4F), but it dispersed from its usual location near the

microtubule-organizing center (MTOC) to the cell periphery (Fig. 5A). In addition, a more

severe dispersion of the TGN protein TGN46 was observed in CoV-2-ORF3a cells (Fig. 5A). The

positioning changes of CI-MPR and TGN46 was confirmed by high-content imaging analysis

(Fig. 5B,C), in which cellular area between MTOC and cell boundary was divided into "center",

"middle", and "periphery" to reflect protein cellular distribution[52]. We also examined Golgi

matrix protein 130 (GM130) but did not see a dispersion (Fig. S5A), indicating that the

dispersion of CI-MPR and TGN46 in the Flp-in cells was not due to Golgi fragmentation[53,54]}.

The dispersed CI-MPR and TGN46 were partially colocalized with LAMP2 (Fig. 5A, arrows),

and quantitative results showed an increased Pearson correlation coefficient between CI-MPR

and LAMP2 (Fig. 5D). Consistently, CoV-2-ORF3a cells had an increased localization between

TGN46 and Rab7 (Fig. 5SB,C). These results indicate that CI-MPR and TGN46 were retained

within the Rab7- or LAMP2-positive endosomes and lysosomes in the CoV-2-ORF3a cells,

suggesting that TGN-to-endosome transport was increased, and/or retrieval from endosomes

and lysosomes was decreased, resulting in protein prolonged retention within the endosome-

lysosome system.

Retrieval of endosomal proteins relies on the retromer[16,55] and retriever[56] complexes. KO of

their shared subunit VPS29 resembled the defects observed in CoV-2 ORF3a cells, including

increased NPC2 secretion (Fig. 5E,F), lysosomal cholesterol accumulation (Fig. 5G), and

enhanced colocalization of CI-MPR and TGN46 with LAMP1 (Fig. 5H,I). These results indicate

that disruption of endosomal protein retrieval impairs lysosome cholesterol egress. We next

examined the Flp-In cells and found that the retromer subunit VPS35, which mediates

endosome-to-Golgi transport, exhibited increased colocalization with LAMP2 in CoV-2-ORF3a

cells (Fig. 5J, arrows). High-content imaging quantification showed an elevated Pearson's

correlation coefficient between VPS35 and LAMP2, which was rescued in W193A cells (Fig. 5K).

Together, these findings indicate that the ORF3a-VPS39 interaction disrupts endosome-to-TGN

transport by retaining retromer in LAMP2-positive endosomes/lysosomes, resulting in reduced

recycling of CI-MPR and targeting of NPC2 to lysosomes.

ORF3a reduced BMPs

BMPs, localized exclusively on late endosome and lysosome membranes, physically interacts

with NPC2 to facilitate cholesterol egress[33,34]. Using an antibody against BMPs[26], we observed

a significant reduction in BMP levels in SARS-CoV-2-infected Vero E6 cells, starting 12 hours

post-infection (Fig. 6A,B). To test if this alteration was related to ORF3a, we examined the Flp-In

cells and found that the BMP level was reduced by 20% in CoV-2-ORF3a cells, compared to the

control cells. This reduction was partially rescued in W193A cells, while the CoV-ORF3a cells

exhibited no change (Fig. 6C,D). Shotgun lipidomics of total lipids extracted from the Flp-In

cells confirmed these results: the two most abundant BMP species, 18:1-18:1 and 18:1-22:6, were

reduced by approximately 20% in CoV-2-ORF3a cells and restored in W193A cells (Fig. 6E).

These results indicate that CoV-2-ORF3a alone can decrease BMP levels, which is dependent on

the ORF3a-VPS39 interaction, aligned with the observed cholesterol sequestration. Furthermore,

exogenous BMP treatment for 2 hours in serum-free media decreased the lysosomal cholesterol

levels by 25% in CoV-2-ORF3a cells (Fig. 5F). Together, these results suggest that reduced BMP

is an additional mechanism underlying SARS-CoV-2-induced cholesterol sequestration.

ORF3a interrupted lysosome-mitochondrion interaction

To investigate the mechanism underlying BMP reduction, we examined key enzymes involved

in BMP biosynthesis and turnover, including CLN5[37], PLD3, PLD4[38], ABHD6[35], and

PLA2G15[39]. With the exception of PLA2G15, which did not display a punctate pattern upon

transfection, we did not observe a reduced colocalization with lysosomes (Fig. S6A-H) or

reduced expression levels (Fig. S6I) of these proteins, suggesting these enzymes are not

responsible for the BMP reduction in CoV-2-ORF3a cells.

We then isolated the lysosomes by Lyso-IP[57] followed by mass spectrometry of lysosomal

proteomes from the control, CoV-2-ORF3a and W193A Flp-In cells (Tab. 1). Comparison

revealed 277 proteins uniquely present in CoV-2-ORF3a samples (gained, Tab. 2), and 348

proteins increased at least 2-fold relative to W193A cells (increased, Tab. 3). In contrast, 160

proteins were undetected (lost, Tab. 2), and 87 proteins decreased by ≥50% in CoV-2-ORF3a

samples (decreased, Tab. 3). Gene ontology analysis using DAVID Bioinformatics indicated that

the gained and increased proteins largely originated from cytoplasm (32%) and nuclei (25%)

(Fig. 7A), while the lost and decreased proteins were predominantly mitochondrial (45%) and

ER-derived (21%) (Fig. 7B,C,D). Among the top 20% of lost proteins, 43.7% proteins were

mitochondrial, including outer membrane, inner membrane, and matrix proteins (Fig. 7C).

Similarly, in the reduced proteins, the majority was from mitochondria, regardless their

mitochondrial localization (Fig. 7D). These results suggest a reduced lysosome-mitochondrion

interaction in CoV-2-ORF3a cells.

Confocal imaging showed frequent lysosome–mitochondria interactions in control cells (Fig. 7E,

upper, arrows), while CoV-2 ORF3a cells exhibited fewer physical contacts, though organelles

remained proximal (Fig. 7E, lower). To quantify the physical interactions, we defined the

cytoplasm area in each cell and excluded the perinuclear clouds to reduce the false overlap due

to a usual crowdedness of organelles at that area (Fig. 7F). Indicated by Pearson correlation

coefficient, the interactions between mitochondria and lysosomes was significantly decreased in

the CoV-2-ORF3a cells and was rescued in the W193A cells (Fig. 7G). High-content imaging of

TOMM20 revealed slightly increased mitochondrial abundance and similar morphology in both

CoV-2-ORF3a and W193A cells (Fig. S7A,B,C), confirming that the reduced lysosome-

mitochondrion interaction was not due to fewer mitochondria or altered mitochondrial

morphology.

We hypothesized that impaired mitochondrion-lysosome interaction may limit mitochondrial

transportation of PGs, the precursors for BMP biosynthesis. Lipidomics revealed unchanged PG

levels among the three cell types, but mitochondrial cardiolipin, a PG derivative, increased in

CoV-2 ORF3a cells (Fig. 7H), suggesting that interrupted PG exportation from mitochondria

resulted in an increased local PG conversion and decreased BMP synthesis in lysosomes.

ORF3a disturbed mitochondrion-lysosome membrane contact sites

The direct physical interactions between mitochondria and lysosomes can be achieved through

autophagy, mitochondrion-derived vesicles (MDVs), and mitochondrion-lysosome membrane

contact sites (MCS). It was shown that autophagy flux is interrupted by ORF3a-VPS39

interaction, which inhibits HOPS-mediated fusion between autophagosomes and lysosomes[32].

We observed the same defect shown as increased LC3-II levels in CoV-2-ORF3a cells (Fig. S8A).

However, inhibiting autophagy fusion by STX17 KO (Fig. 8A), or interrupting earlier autophagy

steps by knockdown (KD) of ATG5, ATG7, or ULK1 (Fig. 8C) did not alter BMP levels (Fig.

8B,D). These results excluded autophagy as a PG transport pathway.

We next tested MDVs, identified by a combined staining of the MDV marker TOMM20 and a

mitochondrial but not MDV protein, pyruvate dehydrogenase (PDH) (Fig. 8E, arrows). We also

included a lysosome marker to visualize the fusion between MDV and lysosomes (Fig. 8E,

circles). CoV-2-ORF3a cells exhibited more MDVs and MDV-lysosome fusion events (Fig. 8F,G),

which did not align with the BMP reduction (Fig. 6C). Furthermore, KD of MDV biogenesis

genes, DRP1 or MIRO1/2[58] (Fig. 8H), did not decrease BMP levels (Fig. 8I), indicating that

MDVs do not mediate PG delivery to lysosomes.

Last, we applied electron microscopy to assess mitochondrion-lysosome MCS. We randomly

chose cells at low magnification and image areas containing abundant mitochondria at higher

magnification. Lysosome-to-mitochondrion ratios were similar across cell types (Fig. S8B), but

MCS events, identified by intact membrane within 10 to 30 nm distance (Fig. 8J), decreased in

CoV-2-ORF3a cells and were rescued in the W193A cells, though it was not statistically

significant (Fig. 8K). VPS39 is known to mediate the MCS formation in a HOPS-independent

manner in yeast[59]. Consistently, VPS39 KO decreased TOMM20-LAMP2 overlap and BMP

level, whereas VPS29, VPS41, or Rab7 KO did not (Fig. 8L,M). There findings suggest that

ORF3a inhibited VPS39-mediated lysosome-mitochondrion MCS formation and impaired PG

transportation, consequently reducing BMP synthesis.

DISCUSSION

Our study identifies SARS-CoV-2 ORF3a as a potent disruptor of lysosomal cholesterol transport through dual interference with NPC2 trafficking and BMP biogenesis. By binding VPS39, ORF3a blocks retromer-mediated retrieval of CI-MPR and reduces mitochondrion-to-lysosome lipid transfer, thereby sequestering cholesterol within lysosomes. These findings expand our understanding of cellular cholesterol regulation and the functions of HOPS and VPS39, point to a regulatory pathway for BMP biosynthesis, and reveal a viral strategy that links lysosomal lipid metabolism to SARS-CoV-2 pathogenesis.

Cholesterol is not only a structural component of cellular membranes but also a dynamic signaling molecule whose intracellular distribution dictates multiple metabolic pathways. Our results demonstrate that SARS-CoV-2 impedes lysosomal cholesterol egress (Fig. 1), adding to the growing recognition that compartmentalized cholesterol pools, rather than bulk cholesterol, drive cellular physiology[60,61]. ORF3a's ability to retain cholesterol within lysosomes (Figs. 2&4C,D) reveals a new viral strategy to manipulate host lipid homeostasis and may explain systemic lipid alterations in COVID-19 patients[1–5]. More broadly, our findings place lysosomal cholesterol transport at the crossroads of host–pathogen interactions and suggest that this pathway may represent a therapeutic vulnerability.

BMPs play essential roles in lysosomal biology and have been implicated in a wide range of human diseases[27,28], such as infectious disease[30,32,62,63], and neurodegenerative disorders[64,65]. Here, we demonstrate that SARS-CoV-2 and its ORF3a protein reduce BMP levels (Fig. 6A-E), and that restoring BMP partially rescues cholesterol egress (Fig. 6F). This finding highlights BMP as a limiting factor for lysosomal cholesterol export, in agreement with prior studies[33,33,34]. Notably, ORF3a alters PG metabolism by decreasing its lysosomal products (BMPs) and increasing its mitochondrial products (cardiolipins) (Fig. 7H). This shift uncovers a previously unrecognized regulatory mechanism in BMP homeostasis, the supply of mitochondrial precursors. While BMP levels were thought to be controlled primarily by lysosomal synthetase and hydrolases[35,37–39], our data suggest that cross-organelle lipid transfer provides an additional layer of regulation. Importantly, by testing multiple possible routes, we found that mitochondrion-lysosome membrane contact sites likely serve as the pathway for PG transport, rather that autophagy or mitochondrial-derived vesicles (Fig. 8). These finding establish BMP regulation as a point of crosstalk between mitochondrial phospholipid metabolism and

397 lysosomal cholesterol transport, with broad implications for both cellular physiology and viral
398 pathogenesis.
399
The canonical view of HOPS has centered on its role as a tethering complex that mediates late
endosome–lysosome and autophagosome–lysosome fusion[17]. By leveraging the VPS39–ORF3a
interaction and the W193A mutant, our study expands this paradigm by showing that VPS39
also regulates two distinct aspects of lysosomal activities, regulation of retromer-dependent
trafficking[66] (Fig. 5) and formation of mitochondrion–lysosome membrane contact sites
necessary for BMP synthesis (Fig. 8J-M). These findings position VPS39 as a multifunctional
coordinator at the lysosome, integrating protein trafficking and lipid exchange. The dual
disruption of these VPS39-dependent functions by ORF3a reframes HOPS/VPS39 not solely as
a fusion mediator, but as a broader regulatory hub essential for lysosomal homeostasis.
409
Our study establishes ORF3a as a viral effector that directly manipulates host lipid metabolism.
Unlike SARS-CoV ORF3a[67], its SARS-CoV-2 counterpart acquired the ability to bind
VPS39[43,44,68], suggesting an evolutionary adaptation to exploit lysosomal cholesterol regulation.
Since plasma membrane cholesterol promotes SARS-CoV-2 entry[11,12], restricting cholesterol
export from lysosomes could reduce cholesterol at the plasma membrane and thereby limit the
viral entry. Consistent with this model, pharmacological blockade of NPC1 with U18666A
sequesters cholesterol in lysosomes and decreases SARS-CoV-2 infection (Fig. S8C–E). In
addition, BMPs are required for viral fusion within endosomes/lysosomes[63], and reduced BMP
levels may impair this step of the viral life cycle. Importantly, ORF3a is an accessory protein
that is not incorporated into virions but translated only after infection from viral subgenomic
RNAs[69]. Thus, its interference with cholesterol export and BMP synthesis is unlikely to affect
the current infection but may restrict secondary infection events. Furthermore, ORF3a's
previously reported activities, including increasing lysosomal pH and damaging lysosomal
membranes, may promote lysosome exocytosis, a route for viral egress[42,70]. Taken together,
these findings suggest that SARS-CoV-2 use ORF3a to modulate multiple lysosomal functions,
integrating cholesterol trafficking, lipid metabolism, and membrane dynamics to optimize
different stages of its life cycle.
427
Our study has several limitations. First, although we demonstrate ORF3a-induced cholesterol
sequestration and identify VPS39-dependent mechanisms, the downstream consequences for
viral replication, immune responses, and cell survival remain to be fully elucidated *in vivo*.
Second, while we observed reduced mitochondrion–lysosome contacts and altered BMP levels,

the molecular intermediates mediating phosphatidylglycerol transfer remain undefined.

Finally, the interplay between cholesterol sequestration and systemic lipid dysregulation

observed in COVID-19 patients requires further integration with clinical data. Notably, SARS-

CoV-2 infects hepatocytes[71], and given the conserved role of HOPS in lysosomal trafficking,

similar lipid defects may occur in the liver, a central regulator of cholesterol and lipoprotein

metabolism. Future work will therefore examine hepatocytes to determine whether ORF3a-

mediated lysosomal disruptions contribute to the lipoprotein abnormalities observed in

COVID-19 and long COVID.

In summary, this work uncovers cholesterol transport and BMP biogenesis as dual regulatory

nodes governed by VPS39 and reveals how SARS-CoV-2 ORF3a exploits these pathways to

rewire lysosomal lipid metabolism. These findings advance fundamental understanding of

lysosomal protein trafficking and lysosome–mitochondrion crosstalk, broaden the functional

repertoire of HOPS/VPS39, and provide new perspectives on viral pathogenesis and

cholesterol-related diseases.

MATERIALS AND METHODS

Cells and virus

A549 and HeLa cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% heat-inactivated fetal bovine serum (FBS), 25 mM HEPES, and MycoZap Plus-CL (Lonza, Basel, Switzerland) at 37°C with 5% CO_2. Vero E6 cells (ATCC, CRL-1586) were cultured in DMEM supplemented with 10% heat-inactivated FBS, 1% pen/strep, 2mM L-glut, 1% non-essential amino acids, 1% HEPES. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2, Isolate USA-WA1/2020) was acquired from BEI Resources (NR-52281) and propagated in Vero E6 cells for 3 days. Infectious virus was isolated by harvesting cellular supernatant and spinning at 1000 g for 10 minutes to remove cellular debris and stored at -80°C. SARS-CoV-2 infectious particles were quantified in media by standard plaque assay[72]

SARS-CoV-2 infections

Coverslips or optic 96-well plates (Agilent, Santa Clara, CA) were coated with collagen solution (0.03-0.04 mg/ml collagen in 0.02 N acetic acid) for > 30 min and then washed twice with PBS before receiving cells. A549-hACE2 cells were seeded on coverslips at 0.1×10^6 cells per well or in optical 96-well plates at 0.012×10^6 cells per well and infected with SARS-CoV-2 in DMEM (4% FBS) at MOI: 4 and incubated at 37°C for 1 hr on the next day. Cells were then washed twice with 1x PBS and normal culture media was added. At the indicated times post-infection, cells were fixed with 4% PFA. Vero E6 cells were seeded in optical 96-well plate around 0.025×10^6 cells per well and infected with SARS-CoV-2 at MOI: 0.1 in 2% FBS media. The control group of cells were incubated with the media only.

Plasmid and siRNA transfection

Cells were seeded a day before transfection and receive plasmids mixed with transfection reagent Lipofectamine 2000 (Thermo Fisher Scientific) or TransIT-LT1 (Mirus Bio, Madison, WI), following the manufactures' instructions. Cells were analyzed at 24- or 48-h post transfection. On-TARGETplus SMARTpool siRNAs were applied (Horizon Discovery, Cambridge, UK) for two-shot transfection at Day 1 and Day 3 after seeding. The transfection reagents Lipofectamine 2000 (Thermo Fisher Scientific) or TransIT-X2 (Mirus Bio) were used to deliver siRNA, and cells were analyzed 6-7 days after seeding.

CRISPR KO

Genes were edited using the CRISPR/Cas9 system. Guide RNAs (gRNAs) were designed using Benchling () Four to six 20-base pair (bp) gRNAs were selected and tested for the cleavage in

pilot experiments. gRNA was introduced into the px458 plasmid containing GFP sequence

(Addgene, Cambridge, MA). Cells were transfected with the plasmids, and GFP-positive cells

were collected by cell sorter 48 h after transfection. Transformants were kept in normal

medium for another 12 days to allow single colony formation. Genomic DNA was extracted

from individual colonies, and cleavage of the target sequence was tested by PCR followed by

Sanger sequencing and immunoblotting. Three to six single colonies were pooled for

experiment use.

Guide RNA target sequences used in this study:

Rab7 KO: TAGTTTGAAGGATGACCTCT and TTGCTGAAGGTTATCATCCT

VPS29 KO: GCAAACTGTTGCACCGGTGT and CATAACTCTCTTTGGTGCAA

Stx17 KO: TGGAGAAGACAGCTGTTACCAGGG and TTAAGATAGTAATCCCAACAGACC

Generation of flip-in cells

HeLa Flp-In host cells (Thermo Fisher Scientific, Waltham, MA) were transfected with the

plasmids encoding FLAG-tagged proteins or linker peptide and subjected to hygromycin

selection after 48 h post transfection in DMEM as described above but replacing regular FBS

with tetracycline-free FBS (Thermo Fisher Scientific) (tet-free media). Following 2-week

selection, transformants were kept in tet-free medium for another 12 days to allow single colony

formation. Protein expression was induced by addition of 1 μg/ml doxycycline for 16 hours.

The colonies had relatively low expression levels of FLAG were chosen and pooled for

experiments, determined by immunoblotting of FLAG.

Immunofluorescence, Filipin staining, and confocal microscopy

Fixed cells were washed 3 times with PBS, and primary antibodies were diluted in 1% BSA in

PBS supplemented with 0.2% saponin and applied to cells for 1 h at 37°C or overnight at 4°C.

Alexa-conjugated secondary antibodies (Thermo Fisher Scientific) were applied with the same

saponin-contained BSA-PBS solution for 30 min at 37°C. Coverslips were mounted with DAPI-

contained mounting reagents (Electron Microscopy Sciences, Hatfield, PA) and allowed to dry

at 37°C for at least 1 h. To stain free cholesterol, 25 μg/ml filipin complex (Millipore Sigma, St.

Louis, MO) in PBS was applied to fixed cells at 37°C for 30 min. When immunostaining was

performed with filipin, primary antibodies and Alexa-conjugated secondary antibodies along

with DRAQ7 (Novus Biologicals, Centennial, CO) were sequentially applied to filipin-stained

cells without detergent permeabilization. Then cells were mounted with mounting reagents

without DAPI (Electron Microscopy Sciences) at 37°C for at least 1 h. Images were acquired

using a Zeiss LSM800 confocal microscope equipped with the software ZEN (Zeiss,

Oberkochen, Germany).

High-content imaging and image analysis

Cells in optical 96-well plates were fixed and stained as described above and maintained in PBS for immediate imaging. Imaging and analysis were performed using the CellInsight Microscope platform (Thermo Fisher Scientific). DAPI- or DRAQ7-stained nuclei were scanned to determine focus through the programmed autofocus function. Cytoplasmic signals from HCS CellMask Near-IR Stain (Thermo Fisher Scientific) were applied to define individual cell boundaries. Cells were excluded based on criteria including incomplete cell boundaries at image edges, abnormal size or shape, and signs of cell death.

For cellular punctum analysis, the cell boundary-enclosed areas are defined as ROI_A. Filipin, LAMP, or BMP puncta were detected as spots (2 pixels in size) using the software "spot detection" function. Line-shaped filipin signals from the plasma membrane were excluded by setting the validation criteria for shape. Puncta that overlapped with ROI_A were quantified by their fluorescence intensity. To measure colocalized signals, the software function of "colocalization" was applied, and the output feature "ROI_A(B) Correlation Coefficient" was extracted for the quantification of Pearson Correlation Coefficient.

To analyze the infected cells, fixed cells were immunostained with an dsRNA antibody, and its signals were used to set the threshold for detection of infected cells, using the function "Reference" of the software.

Sample Processing for electron microscopy

Cells on glass coverslips were fixed in 0.1 M PIPES buffer containing 3% formaldehyde, 2% glutaraldehyde, and 1.5 mM CaCl2, then washed in several buffer changes before secondary fixation in 0.1 M PIPES containing: 1% osmium tetroxide for 15 min, 5 min PIPES rinse, 15 min 1% carbohydrazide in PIPES, 5 min PIPES rinse, and 15 min 1% osmium tetroxide (OCO). Samples were then washed with buffer, followed by ultrapure water, before being dehydrated through a graded series of ethanol to propylene oxide, infiltrated with Epon-Araldite resin, embedded, and heat-cured. The glass was removed from the cured resin blocks so the cells could be re-mounted for en face sectioning. Thin sections (60-80 nm) were mounted on Cu grids, post-stained with uranyl acetate and Reynold's lead citrate, and examined in a Hitachi HT7700 TEM operated at 80 kV. Images were captured with an AMT XR-81 CCD camera.

Cholesterol measurement by GC-MS

Cells were collected in PBS with a cell scraper after washing twice with PBS, and the total lipids

were extracted by adding lipid-extraction solvent (chloroform:methanol:acetic acid = 50:50:1) of

90% volume of PBS. The organic phase and aqueous phase were separated by 10 minute-

centrifugation at 21,000 g. Organic phase was collected and transferred into a new tube and

dried with vacuum. The extracted lipids were stored at -80°C for further analysis.

Cholesterol calibration standard solutions were prepared from the concentration of 50 μg/ml to

400 μg/ml along with 80 μg/ml 5a-cholestane as the internal standard. The linear calibration

curve was obtained with R2 =0.9884. The dried lipids were dissolved with hexane and mixed

with 4 μg 5a-cholestane. The GC-MS analysis was conducted on Agilent 6890/5973 with

selected ion monitoring (SIM) mode.

Shotgun Lipidomics

Lipid species were analyzed using a multidimensional mass spectrometry-based shotgun

lipidomics approach[73]. In brief, each cell sample homogenate containing 0.3 mg of protein

which was determined with a Pierce BCA assay was accurately transferred to a disposable glass

culture test tube. A premixture of lipid internal standards was added prior to conducting lipid

extraction for quantification of the targeted lipid classes. Lipid extraction was performed using

a modified Bligh and Dyer procedure (Wang and Han 2014), and each lipid extract was

reconstituted in chloroform:methanol (1:1, v:v) at a volume of 400 μL/mg protein.

Derivatization of lipid extracts for analysis of free fatty acid (FFA) and

bis(monoacylglycero)phosphate (BMP) was performed as described previously[74,75] before

lipidomics analysis. Lyso-phosphatidylglycerol (LPG) in the aqueous phase was enriched using

a HybridSPE cartridge. After washing with methanol, lysophospholipids were eluted with

methanol/ammonia hydroxide (9:1), dried and reconstituted in methanol for lipidomics

analysis [76]. For shotgun lipidomics, individual lipid samples prepared as aforementioned was

further diluted to a final concentration of ~500 fmol total lipids per μL. Mass spectrometric

analysis was performed on a triple quadrupole mass spectrometer (TSQ Altis, Thermo Fisher

Scientific, San Jose, CA) and a Q Exactive mass spectrometer (Thermo Scientific, San Jose, CA),

both of which were equipped with an automated nanospray device (TriVersa NanoMate,

Advion Bioscience Ltd., Ithaca, NY) as described[77]. Identification and quantification of lipid

species were performed using an automated software program[78,79]. Data processing (e.g., ion

peak selection, baseline correction, data transfer, peak intensity comparison, and quantitation)

was performed as described[79]. The results were normalized to the protein content (nmol

lipid/mg protein).

Analysis of intracellular and extracellular proteins by Western blotting

To detect secreted NPC2 and cathepsin D, cells were kept in a minimum volume of DMEM without serum for 6 h after three-time washing with the same medium. The media were collected and centrifuged at 1,000 g for 5 min to pellet any detached cells. The supernatants were mixed with SDS loading buffer and subjected to SDS-PAGE and immunoblotting. After collecting the media, cells were incubated in complete DMEM for 2 h. Cell lysates were obtained by using 2X SDS sample buffer.

Quantification and statistical analysis

Image J was used for quantifying fluorescence signals of randomly-taken images from at least three independent experiments. The signal intensity of vesicular filipin was measured using the function of "Analyze particles", and co-localization was analyzed with the plug-in "PSC Colocalization"[83]. High-content imaging results were obtained from more than 1,000 cells per well and at least 3 wells per experiment. The number of trials is specified in the legends where appropriate. The bar graphs show the pooled results displayed as mean ± SD. Colocalization results include frequency distribution. Statistical significance was determined by comparing two datasets using a t test with one-tailed distribution or more than two datasets using one-way ANOVA, using the software Prism 9 (GraphPad). Probability values and number of trials are given in the figure captions and the legends where appropriate. The statistical significance is generally denoted as follows: *, $p < 0.05$, **, $p < 0.01$, ***, $p < 0.001$, ****, $p < 0.0001$, and n.s., not significant.

AUTHOR CONTRIBUTIONS

ACKNOWLEDGMENTS

We thank A. Kajon for gift of A549-hACE2 cells, P. Lobel for mch-NPC2 construct and NPC2

antibody, J. Bonifacino for VPS29-KO HeLa cells, and J. Anderson for experimental assistance.

EM data were generated in the HSC-Electron Microscopy Facility, supported by University of

New Mexico Health Sciences Center (HSC). This research made use of the CellInsight

Microscope Platform and confocal microscope from Autophagy, Inflammation, and Metabolism

Center of Biomedical Research Excellence, funded by NIGMS, NIH (P20GM121176). Lipidomics

analysis was performed at Barshop Institute Functional Lipidomics Core funded by NIA, NIH

(P30AG013319). The funding support was from NIGMS, NIH (T32GM144834 and

R35GM147419).

SUPPLEMENTAL INFORMATION

633 Supplemental Information includes 8 figures and 3 tables.

REFERENCES

1. Li, Y., Zhang, Y., Lu, R., Dai, M., Shen, M., Zhang, J., Cui, Y., Liu, B., Lin, F., Chen, L., Han, D., Fan, Y., Zeng, Y., Li, W., Li, S., Chen, X., Li, H. & Pan, P., Lipid metabolism changes in patients with severe COVID-19. Clin Chim Acta 2021 517 33639119

2. Rezaei, A., Neshat, S. & Heshmat-Ghahdarijani, K., Alterations of Lipid Profile in COVID-19: A Narrative Review. Curr Probl Cardiol 2022 47(3) 34272088

3. Zickler, M., Stanelle-Bertram, S., Ehret, S., Heinrich, F., Lange, P., Schaumburg, B., Kouassi, N. M., Beck, S., Jaeckstein, M. Y., Mann, O., Krasemann, S., Schroeder, M., Jarczak, D., Nierhaus, A., Kluge, S., Peschka, M., Schlüter, H., Renné, T., Pueschel, K., Kloetgen, A., Scheja, L., Ondruschka, B., Heeren, J. & Gabriel, G., Replication of SARS-CoV-2 in adipose tissue determines organ and systemic lipid metabolism in hamsters and humans. Cell Metab 2022 34(1) 34895500

4. Xu, E., Xie, Y. & Al-Aly, Z., Risks and burdens of incident dyslipidaemia in long COVID: a cohort study. Lancet Diabetes Endocrinol 2023 11(2) 36623520

5. López-Hernández, Y., Oropeza-Valdez, J. J., García Lopez, D. A., Borrego, J. C., Murgu, M., Valdez, J., López, J. A. & Monárrez-Espino, J., Untargeted analysis in post-COVID-19 patients reveals dysregulated lipid pathways two years after recovery. Front Mol Biosci 2023 10 36936993

6. Luo, J., Yang, H. & Song, B. L., Mechanisms and regulation of cholesterol homeostasis. Nat Rev Mol Cell Biol 2020 21(4) 31848472

7. Luo, J., Jiang, L. Y., Yang, H. & Song, B. L., Intracellular Cholesterol Transport by Sterol Transfer Proteins at Membrane Contact Sites. Trends Biochem Sci 2019 44(3) 30415968

8. Meng, Y., Heybrock, S., Neculai, D. & Saftig, P., Cholesterol Handling in Lysosomes and Beyond. Trends Cell Biol 2020 30(6) 32413315

9. Matsuo, H., Chevallier, J., Mayran, N., Le Blanc, I., Ferguson, C., Fauré, J., Blanc, N. S., Matile, S., Dubochet, J., Sadoul, R., Parton, R. G., Vilbois, F. & Gruenberg, J., Role of LBPA and Alix in multivesicular liposome formation and endosome organization. Science 2004 303(5657) 14739459

10. Spector, A. A. & Yorek, M. A., Membrane lipid composition and cellular function. J Lipid Res 1985 26(9) 3906008

11. Sanders, D. W., Jumper, C. C., Ackerman, P. J., Bracha, D., Donlic, A., Kim, H., Kenney, D., Castello-Serrano, I., Suzuki, S., Tamura, T., Tavares, A. H., Saeed, M., Holehouse, A. S., Ploss, A., Levental, I., Douam, F., Padera, R. F., Levy, B. D. & Brangwynne, C. P., SARS-CoV-2 requires cholesterol for viral entry and pathological syncytia formation. Elife 2021 10 33890572

12. Wang, H., Yuan, Z., Pavel, M. A., Jablonski, S. M., Jablonski, J., Hobson, R., Valente, S., Reddy, C. B. & Hansen, S. B., The role of high cholesterol in SARS-CoV-2 infectivity. J Biol Chem 2023 299(6) 37119851

13. Eskelinen, E. L., Schmidt, C. K., Neu, S., Willenborg, M., Fuertes, G., Salvador, N., Tanaka, Y., Lüllmann-Rauch, R., Hartmann, D., Heeren, J., von Figura, K., Knecht, E. & Saftig, P., Disturbed cholesterol traffic but normal proteolytic function in LAMP-1/LAMP-2 double-deficient fibroblasts. Mol Biol Cell 2004 15(7) 15121881

14. Popovic, K., Holyoake, J., Pomès, R. & Privé, G. G., Structure of saposin A lipoprotein discs. Proc Natl Acad Sci U S A 2012 109(8) 22308394

15. Braulke, T. & Bonifacino, J. S., Sorting of lysosomal proteins. Biochim Biophys Acta 2009 1793(4) 19046998

16. Seaman, M. N., McCaffery, J. M. & Emr, S. D., A membrane coat complex essential for endosome-to-Golgi retrograde transport in yeast. J Cell Biol 1998 142(3) 9700157

17. Solinger, J. A. & Spang, A., Tethering complexes in the endocytic pathway: CORVET and HOPS. FEBS J 2013 280(12) 23351085

18. Anderson, J., Walker, G. & Pu, J., BORC-ARL8-HOPS ensemble is required for lysosomal cholesterol egress through NPC2. Mol Biol Cell 2022 33(9) 35653304

19.   Schleinitz, A., Pöttgen, L. A., Keren-Kaplan, T., Pu, J., Saftig, P., Bonifacino, J. S., Haas, A.
& Jeschke, A., Consecutive functions of small GTPases guide HOPS-mediated tethering of
late endosomes and lysosomes. Cell Rep 2023 42(1) 36640308
20.   Bonifacino, J. S. & Hierro, A., Transport according to GARP: receiving retrograde cargo at
the trans-Golgi network. Trends Cell Biol 2011 21(3) 21183348
21.   Willenborg, M., Schmidt, C. K., Braun, P., Landgrebe, J., von Figura, K., Saftig, P. &
Eskelinen, E. L., Mannose 6-phosphate receptors, Niemann-Pick C2 protein, and
lysosomal cholesterol accumulation. J Lipid Res 2005 46(12) 16177447
22.   Wei, J., Zhang, Y. Y., Luo, J., Wang, J. Q., Zhou, Y. X., Miao, H. H., Shi, X. J., Qu, Y. X., Xu,
J., Li, B. L. & Song, B. L., The GARP Complex Is Involved in Intracellular Cholesterol
Transport via Targeting NPC2 to Lysosomes. Cell Rep 2017 19(13) 28658628
23.   Carstea, E. D., Morris, J. A., Coleman, K. G., Loftus, S. K., Zhang, D., Cummings, C., Gu, J.,
Rosenfeld, M. A., Pavan, W. J., Krizman, D. B., Nagle, J., Polymeropoulos, M. H., Sturley,
S. L., Ioannou, Y. A., Higgins, M. E., Comly, M., Cooney, A., Brown, A., Kaneski, C. R.,
Blanchette-Mackie, E. J., Dwyer, N. K., Neufeld, E. B., Chang, T. Y., Liscum, L., Strauss, J.
F., Ohno, K., Zeigler, M., Carmi, R., Sokol, J., Markie, D., O'Neill, R. R., van Diggelen, O.
P., Elleder, M., Patterson, M. C., Brady, R. O., Vanier, M. T., Pentchev, P. G. & Tagle, D. A.,
Niemann-Pick C1 disease gene: homology to mediators of cholesterol homeostasis. Science
1997 277(5323) 9211849
24.   Naureckiene, S., Sleat, D. E., Lackland, H., Fensom, A., Vanier, M. T., Wattiaux, R., Jadot,
M. & Lobel, P., Identification of HE1 as the second gene of Niemann-Pick C disease.
Science 2000 290(5500) 11125141
25.   Carette, J. E., Raaben, M., Wong, A. C., Herbert, A. S., Obernosterer, G., Mulherkar, N.,
Kuehne, A. I., Kranzusch, P. J., Griffin, A. M., Ruthel, G., Dal Cin, P., Dye, J. M., Whelan, S.
P., Chandran, K. & Brummelkamp, T. R., Ebola virus entry requires the cholesterol
transporter Niemann-Pick C1. Nature 2011 477(7364) 21866103
26.   Kobayashi, T., Stang, E., Fang, K. S., de Moerloose, P., Parton, R. G. & Gruenberg, J., A
lipid associated with the antiphospholipid syndrome regulates endosome structure and
function. Nature 1998 392(6672) 9515966
27.   Showalter, M. R., Berg, A. L., Nagourney, A., Heil, H., Carraway, K. L. & Fiehn, O., The
Emerging and Diverse Roles of Bis(monoacylglycero) Phosphate Lipids in Cellular
Physiology and Disease. Int J Mol Sci 2020 21 33137979
28.   Medoh, U. N. & Abu-Remaileh, M., The Bis(monoacylglycero)-phosphate Hypothesis:
From Lysosomal Function to Therapeutic Avenues. Annu Rev Biochem 2024 93(1)
38603559
29.   Le Blanc, I., Luyet, P. P., Pons, V., Ferguson, C., Emans, N., Petiot, A., Mayran, N.,
Demaurex, N., Fauré, J., Sadoul, R., Parton, R. G. & Gruenberg, J., Endosome-to-cytosol
transport of viral nucleocapsids. Nat Cell Biol 2005 7 15951806
30.   Roth, S. L. & Whittaker, G. R., Promotion of vesicular stomatitis virus fusion by the
endosome-specific phospholipid bis(monoacylglycero)phosphate (BMP). FEBS Lett 2011
585(6) 21333650
31.   Bissig, C., Lenoir, M., Velluz, M. C., Kufareva, I., Abagyan, R., Overduin, M. & Gruenberg,
J., Viral infection controlled by a calcium-dependent lipid-binding module in ALIX. Dev
Cell 2013 25(4) 23664863
32.   Markosyan, R. M., Marin, M., Zhang, Y., Cohen, F. S. & Melikyan, G. B., The late
endosome-resident lipid bis(monoacylglycero)phosphate is a cofactor for Lassa virus
fusion. PLoS Pathog 2021 17(9) 34492091
33.   Chevallier, J., Chamoun, Z., Jiang, G., Prestwich, G., Sakai, N., Matile, S., Parton, R. G. &
Gruenberg, J., Lysobisphosphatidic acid controls endosomal cholesterol levels. J Biol
Chem 2008 283(41) 18644787
34.   McCauliff, L. A., Langan, A., Li, R., Ilnytska, O., Bose, D., Waghalter, M., Lai, K., Kahn, P.
C. & Storch, J., Intracellular cholesterol trafficking is dependent upon NPC2 interaction
with lysobisphosphatidic acid. Elife 2019 8 31580258

35. Pribasnig, M. A., Mrak, I., Grabner, G. F., Taschler, U., Knittelfelder, O., Scherz, B.,
Eichmann, T. O., Heier, C., Grumet, L., Kowaliuk, J., Romauch, M., Holler, S., Anderl, F.,
Wolinski, H., Lass, A., Breinbauer, R., Marsche, G., Brown, J. M. & Zimmermann, R., α/β
Hydrolase Domain-containing 6 (ABHD6) Degrades the Late Endosomal/Lysosomal
Lipid Bis(monoacylglycero)phosphate. J Biol Chem 2015 290(50) 26491015
36. Grabner, G. F., Fawzy, N., Pribasnig, M. A., Trieb, M., Taschler, U., Holzer, M., Schweiger,
M., Wolinski, H., Kolb, D., Horvath, A., Breinbauer, R., Rülicke, T., Rabl, R., Lass, A.,
Stadlbauer, V., Hutter-Paier, B., Stauber, R. E., Fickert, P., Zechner, R., Marsche, G.,
Eichmann, T. O. & Zimmermann, R., Metabolic disease and ABHD6 alter the circulating
bis(monoacylglycerol)phosphate profile in mice and humans. J Lipid Res 2019 60(5)
30894461
37. Medoh, U. N., Hims, A., Chen, J. Y., Ghoochani, A., Nyame, K., Dong, W. & Abu-
Remaileh, M., The Batten disease gene product CLN5 is the lysosomal
bis(monoacylglycero)phosphate synthase. Science 2023 381(6663) 37708259
38. Singh, S., Dransfeld, U. E., Ambaw, Y. A., Lopez-Scarim, J., Farese, R. V. & Walther, T. C.,
PLD3 and PLD4 synthesize S,S-BMP, a key phospholipid enabling lipid degradation in
lysosomes. Cell 2024 187(24) 39423811
39. Nyame, K., Xiong, J., Alsohybe, H. N., de Jong, A. P. H., Peña, I. V., de Miguel, R.,
Brummelkamp, T. R., Hartmann, G., Nijman, S. M. B., Raaben, M., Simcox, J. A., Blomen,
V. A. & Abu-Remaileh, M., PLA2G15 is a BMP hydrolase and its targeting ameliorates
lysosomal disease. Nature 2025 642(8067) 40335701
40. Cansell, M., Gouygou, J. P., Jozefonvicz, J. & Letourneur, D., Lipid composition of
cultured endothelial cells in relation to their growth. Lipids 1997 32(1) 9075191
41. Bar-On, Y. M., Flamholz, A., Phillips, R. & Milo, R., SARS-CoV-2 (COVID-19) by the
numbers. Elife 2020 9 32228860
42. Ghosh, S., Dellibovi-Ragheb, T. A., Kerviel, A., Pak, E., Qiu, Q., Fisher, M., Takvorian, P.
M., Bleck, C., Hsu, V. W., Fehr, A. R., Perlman, S., Achar, S. R., Straus, M. R., Whittaker, G.
R., de Haan, C. A. M., Kehrl, J., Altan-Bonnet, G. & Altan-Bonnet, N., β-Coronaviruses Use
Lysosomes for Egress Instead of the Biosynthetic Secretory Pathway. Cell 2020 183(6)
33157038
43. Miao, G., Zhao, H., Li, Y., Ji, M., Chen, Y., Shi, Y., Bi, Y., Wang, P. & Zhang, H., ORF3a of
the COVID-19 virus SARS-CoV-2 blocks HOPS complex-mediated assembly of the SNARE
complex required for autolysosome formation. Dev Cell 2021 56(4) 33422265
44. Zhang, Y., Sun, H., Pei, R., Mao, B., Zhao, Z., Li, H., Lin, Y. & Lu, K., The SARS-CoV-2
protein ORF3a inhibits fusion of autophagosomes with lysosomes. Cell Discov 2021 7(1)
33947832
45. Miller, A. N., Houlihan, P. R., Matamala, E., Cabezas-Bratesco, D., Lee, G. Y., Cristofori-
Armstrong, B., Dilan, T. L., Sanchez-Martinez, S., Matthies, D., Yan, R., Yu, Z., Ren, D.,
Brauchi, S. E. & Clapham, D. E., The SARS-CoV-2 accessory protein Orf3a is not an ion
channel, but does interact with trafficking proteins. Elife 2023 12 36695574
46. Kern, D. M., Sorum, B., Mali, S. S., Hoel, C. M., Sridharan, S., Remis, J. P., Toso, D. B.,
Kotecha, A., Bautista, D. M. & Brohawn, S. G., Cryo-EM structure of SARS-CoV-2 ORF3a
in lipid nanodiscs. Nat Struct Mol Biol 2021 28(7) 34158638
47. Dolan, K. A., Dutta, M., Kern, D. M., Kotecha, A., Voth, G. A. & Brohawn, S. G., Structure
of SARS-CoV-2 M protein in lipid nanodiscs. Elife 2022 11 36264056
48. Andrews, B. J., Proteau, G. A., Beatty, L. G. & Sadowski, P. D., The FLP recombinase of the
2 micron circle DNA of yeast: interaction with its target sequences. Cell 1985 40(4) 3879971
49. Argos, P., Landy, A., Abremski, K., Egan, J. B., Haggard-Ljungquist, E., Hoess, R. H.,
Kahn, M. L., Kalionis, B., Narayana, S. V. & Pierson, L. S., The integrase family of site-
specific recombinases: regional similarities and global diversity. EMBO J 1986 5(2) 3011407
50. Jia, J., Wang, F., Bhujabal, Z., Peters, R., Mudd, M., Duque, T., Allers, L., Javed, R., Salemi,
M., Behrends, C., Phinney, B., Johansen, T. & Deretic, V., Stress granules and mTOR are

791   regulated by membrane atg8ylation during lysosomal damage. J Cell Biol 2022 221(11)
792   36179369
793 51. Jia, J., Claude-Taupin, A., Gu, Y., Choi, S. W., Peters, R., Bissa, B., Mudd, M. H., Allers, L.,
794   Pallikkuth, S., Lidke, K. A., Salemi, M., Phinney, B., Mari, M., Reggiori, F. & Deretic, V.,
795   Galectin-3 Coordinates a Cellular System for Lysosomal Repair and Removal. Dev Cell
796   2020 52(1) 31813797
797 52. Kolaczkowski, O. M., Goodson, B. A., Vazquez, V. M., Jia, J., Bhat, A. Q., Kim, T. H. & Pu,
798   J., Synergistic Role of Amino Acids in Enhancing mTOR Activation Through Lysosome
799   Positioning. bioRxiv 2024 39416115
800 53. Tang, D., Mar, K., Warren, G. & Wang, Y., Molecular mechanism of mitotic Golgi
801   disassembly and reassembly revealed by a defined reconstitution assay. J Biol Chem 2008
802   283(10) 18156178
803 54. Wei, J. H. & Seemann, J., Golgi ribbon disassembly during mitosis, differentiation and
804   disease progression. Curr Opin Cell Biol 2017 47 28390244
805 55. Seaman, M. N., Marcusson, E. G., Cereghino, J. L. & Emr, S. D., Endosome to Golgi
806   retrieval of the vacuolar protein sorting receptor, Vps10p, requires the function of the
807   VPS29, VPS30, and VPS35 gene products. J Cell Biol 1997 137(1) 9105038
808 56. McNally, K. E., Faulkner, R., Steinberg, F., Gallon, M., Ghai, R., Pim, D., Langton, P.,
809   Pearson, N., Danson, C. M., Nägele, H., Morris, L. L., Singla, A., Overlee, B. L., Heesom, K.
810   J., Sessions, R., Banks, L., Collins, B. M., Berger, I., Billadeau, D. D., Burstein, E. & Cullen,
811   P. J., Retriever is a multiprotein complex for retromer-independent endosomal cargo
812   recycling. Nat Cell Biol 2017 19(10) 28892079
813 57. Abu-Remaileh, M., Wyant, G. A., Kim, C., Laqtom, N. N., Abbasi, M., Chan, S. H.,
814   Freinkman, E. & Sabatini, D. M., Lysosomal metabolomics reveals V-ATPase- and mTOR-
815   dependent regulation of amino acid efflux from lysosomes. Science 2017 358(6364)
816   29074583
817 58. König, T., Nolte, H., Aaltonen, M. J., Tatsuta, T., Krols, M., Stroh, T., Langer, T. & McBride,
818   H. M., MIROs and DRP1 drive mitochondrial-derived vesicle biogenesis and promote
819   quality control. Nat Cell Biol 2021 23(12) 34873283
820 59. González Montoro, A., Auffarth, K., Hönscher, C., Bohnert, M., Becker, T., Warscheid, B.,
821   Reggiori, F., van der Laan, M., Fröhlich, F. & Ungermann, C., Vps39 Interacts with Tom40
822   to Establish One of Two Functionally Distinct Vacuole-Mitochondria Contact Sites. Dev
823   Cell 2018 45(5) 29870720
824 60. Ikonen, E. & Zhou, X., Cholesterol transport between cellular membranes: A balancing act
825   between interconnected lipid fluxes. Dev Cell 2021 56(10) 34004151
826 61. Johnson, K. A. & Radhakrishnan, A., Accessibility of cholesterol at cell surfaces. J Lipid
827   Res 2020 61(10) 32327483
828 62. Zaitseva, E., Yang, S. T., Melikov, K., Pourmal, S. & Chernomordik, L. V., Dengue virus
829   ensures its fusion in late endosomes using compartment-specific lipids. PLoS Pathog 2010
830   6(10) 20949067
831 63. Birtles, D., Abbas, W. & Lee, J., Bis(Monoacylglycero)Phosphate Promotes Membrane
832   Fusion Facilitated by the SARS-CoV-2 Fusion Domain. J Phys Chem B 2024 128(11)
833   38466655
834 64. Maloney, M. T., Wang, X., Ghosh, R., Andrews, S. V., Maciuca, R., Masoud, S. T., Agam,
835   M., Caprioli, R. M., Astarita, G., Bondar, V. V., Chen, J., Chiu, C. L., Davis, S. S., Ho, A. C.,
836   Nguyen, H. N., Propson, N. E., Reyzer, M. L., Davis, O. B., Deen, M. C., Zhu, S., Di Paolo,
837   G., Vocadlo, D. J., Estrada, A. A., de Vicente, J., Lewcock, J. W., Arguello, A., Suh, J. H.,
838   Huntwork-Rodriguez, S. & Henry, A. G., LRRK2 kinase activity regulates Parkinson's
839   disease-relevant lipids at the lysosome. Mol Neurodegener 2025 20(1) 40770658
840 65. Qureshi, Y. H., Patel, V. M., Berman, D. E., Kothiya, M. J., Neufeld, J. L., Vardarajan, B.,
841   Tang, M., Reyes-Dumeyer, D., Lantigua, R., Medrano, M., Jiménez-Velázquez, I. J., Small,
842   S. A. & Reitz, C., An Alzheimer's Disease-Linked Loss-of-Function CLN5 Variant Impairs

Cathepsin D Maturation, Consistent with a Retromer Trafficking Defect. Mol Cell Biol
2018 38(20) 30037983

66. Walia, K., Sharma, A., Paul, S., Chouhan, P., Kumar, G., Ringe, R., Sharma, M. & Tuli, A.,
SARS-CoV-2 virulence factor ORF3a blocks lysosome function by modulating TBC1D5-
dependent Rab7 GTPase cycle. Nat Commun 2024 15(1) 38448435

67. Shen, S., Lin, P. S., Chao, Y. C., Zhang, A., Yang, X., Lim, S. G., Hong, W. & Tan, Y. J., The
severe acute respiratory syndrome coronavirus 3a is a novel structural protein. Biochem
Biophys Res Commun 2005 330(1) 15781262

68. Gordon, D. E., Jang, G. M., Bouhaddou, M., Xu, J., Obernier, K., White, K. M., O'Meara, M.
J., Rezelj, V. V., Guo, J. Z., Swaney, D. L., Tummino, T. A., Hüttenhain, R., Kaake, R. M.,
Richards, A. L., Tutuncuoglu, B., Foussard, H., Batra, J., Haas, K., Modak, M., Kim, M.,
Haas, P., Polacco, B. J., Braberg, H., Fabius, J. M., Eckhardt, M., Soucheray, M., Bennett, M.
J., Cakir, M., McGregor, M. J., Li, Q., Meyer, B., Roesch, F., Vallet, T., Mac Kain, A., Miorin,
L., Moreno, E., Naing, Z. Z. C., Zhou, Y., Peng, S., Shi, Y., Zhang, Z., Shen, W., Kirby, I. T.,
Melnyk, J. E., Chorba, J. S., Lou, K., Dai, S. A., Barrio-Hernandez, I., Memon, D.,
Hernandez-Armenta, C., Lyu, J., Mathy, C. J. P., Perica, T., Pilla, K. B., Ganesan, S. J.,
Saltzberg, D. J., Rakesh, R., Liu, X., Rosenthal, S. B., Calviello, L., Venkataramanan, S.,
Liboy-Lugo, J., Lin, Y., Huang, X. P., Liu, Y., Wankowicz, S. A., Bohn, M., Safari, M., Ugur,
F. S., Koh, C., Savar, N. S., Tran, Q. D., Shengjuler, D., Fletcher, S. J., O'Neal, M. C., Cai, Y.,
Chang, J. C. J., Broadhurst, D. J., Klippsten, S., Sharp, P. P., Wenzell, N. A., Kuzuoglu-
Ozturk, D., Wang, H. Y., Trenker, R., Young, J. M., Cavero, D. A., Hiatt, J., Roth, T. L.,
Rathore, U., Subramanian, A., Noack, J., Hubert, M., Stroud, R. M., Frankel, A. D.,
Rosenberg, O. S., Verba, K. A., Agard, D. A., Ott, M., Emerman, M., Jura, N., von Zastrow,
M., Verdin, E., Ashworth, A., Schwartz, O., d'Enfert, C., Mukherjee, S., Jacobson, M.,
Malik, H. S., Fujimori, D. G., Ideker, T., Craik, C. S., Floor, S. N., Fraser, J. S., Gross, J. D.,
Sali, A., Roth, B. L., Ruggero, D., Taunton, J., Kortemme, T., Beltrao, P., Vignuzzi, M.,
García-Sastre, A., Shokat, K. M., Shoichet, B. K. & Krogan, N. J., A SARS-CoV-2 protein
interaction map reveals targets for drug repurposing. Nature 2020 583(7816) 32353859

69. Murigneux, E., Softic, L., Aubé, C., Grandi, C., Judith, D., Bruce, J., Le Gall, M.,
Guillonneau, F., Schmitt, A., Parissi, V., Berlioz-Torrent, C., Meertens, L., Hansen, M. M.
K. & Gallois-Montbrun, S., Proteomic analysis of SARS-CoV-2 particles unveils a key role
of G3BP proteins in viral assembly. Nat Commun 2024 15(1) 38245532

70. Chen, D., Zheng, Q., Sun, L., Ji, M., Li, Y., Deng, H. & Zhang, H., ORF3a of SARS-CoV-2
promotes lysosomal exocytosis-mediated viral egress. Dev Cell 2021 56(23) 34706264

71. Wanner, N., Andrieux, G., Badia-I-Mompel, P., Edler, C., Pfefferle, S., Lindenmeyer, M. T.,
Schmidt-Lauber, C., Czogalla, J., Wong, M. N., Okabayashi, Y., Braun, F., Lütgehetmann,
M., Meister, E., Lu, S., Noriega, M. L. M., Günther, T., Grundhoff, A., Fischer, N.,
Bräuninger, H., Lindner, D., Westermann, D., Haas, F., Roedl, K., Kluge, S., Addo, M. M.,
Huber, S., Lohse, A. W., Reiser, J., Ondruschka, B., Sperhake, J. P., Saez-Rodriguez, J.,
Boerries, M., Hayek, S. S., Aepfelbacher, M., Scaturro, P., Puelles, V. G. & Huber, T. B.,
Molecular consequences of SARS-CoV-2 liver tropism. Nat Metab 2022 4(3) 35347318

72. Mendoza, E. J., Manguiat, K., Wood, H. & Drebot, M., Two Detailed Plaque Assay
Protocols for the Quantification of Infectious SARS-CoV-2. Curr Protoc Microbiol 2020
57(1) 32475066

73. Han, X. *Lipidomics: Comprehensive mass spectrometry of lipids* (John Wiley & Sons, 2016).

74. Cheng, H., Jiang, X. & Han, X., Alterations in lipid homeostasis of mouse dorsal root
ganglia induced by apolipoprotein E deficiency: a shotgun lipidomics study. J Neurochem
2007 101(1) 17241120

75. Han, X., Yang, K., Cheng, H., Fikes, K. N. & Gross, R. W., Shotgun lipidomics of
phosphoethanolamine-containing lipids in biological samples after one-step in situ
derivatization. J Lipid Res 2005 46(7) 15834120

76. Wang, M. & Han, X., Multidimensional mass spectrometry-based shotgun lipidomics.
Methods Mol Biol 2014 1198 25270931

77.   Han, X., Yang, K. & Gross, R. W., Microfluidics-based electrospray ionization enhances the
intrasource separation of lipid classes and extends identification of individual molecular
species through multi-dimensional mass spectrometry: development of an automated
high-throughput platform for shotgun lipidomics. Rapid Commun Mass Spectrom 2008
22(13) 18523984
78.   Wang, M., Wang, C., Han, R. H. & Han, X., Novel advances in shotgun lipidomics for
biology and medicine. Prog Lipid Res 2016 61 26703190
79.   Yang, K., Cheng, H., Gross, R. W. & Han, X., Automated lipid identification and
quantification by multidimensional mass spectrometry-based shotgun lipidomics. Anal
Chem 2009 81(11) 19408941
906

Figure 1

908
Figure 1 Characterization of cellular free cholesterol in SARS-CoV-2 infected cells
A-C. A549 cells stably expressing human ACE2 receptor (A549-hACE2) were infected with
SARS-CoV-2 and fixed at the indicated time post-infection (P.I.). Filipin was applied to visualize
free cholesterol. A dsRNA antibody was used to identify the infected cells. A LAMP1 antibody
was used to label lysosomes. Confocal microscopy was performed to collect fluorescence
images, and the images at 24 h P.I. were shown in **A**, with arrows pointing colocalized filipin
and LAMP1. The intensity of Filipin puncta was quantified with FIJI, and the mean values of
total filipin in each cell from three independent experiments were plotted, shown as in **B**.
Vesicular filipin was quantified at 24 h P.I. from at least 1,000 infected cells per well and three
replicate wells in an experiment by CellInsight microscope platform. Three independent
experiments were performed, and pooled results were shown in **C**. **D,E.** Similar experiments

were perform as above in Vero E6 cells, and filipin was quantified at the indicated time points.

F. Total lipids were extracted from mock and infected A549-hACE2 cells at 24 h P.I. and

subjected to free cholesterol measurement with GC-MS. Total protein levels were measure by

BCA assays. Two independent experiments with 3 replications were performed. Bar graphs are

presented as min-max (**B**) or mean ± SD (others). p values were determined using t' test or One-

way ANOVA test. **, $p<0.001$, ***, $p<0.001$, ****, $p<0.0001$, n.s., not significant. Scale bars, 5 µm.

Figure 2

Figure 2 Identification of SARS-CoV-2 proteins responsible for lysosome cholesterol

sequestration

A. Plasmids encoding SARS-CoV-2 proteins were transfected into A549-hACE2 cells

individually. Cells were fixed at 24 h post-transfection and co-stained with filipin to visualized

free cholesterol and antibodies against the epitope tags to identify transfected cells. Spike

protein has no tag and was identified with an antibody. Confocal imaging was performed to

randomly take images, and filipin intensity was quantified with FIJI in twenty to fifty cells from

two independent experiments. **B.** A549-hACE2 cells were transfected with ORF3a-GFP plasmid

as in **A** and co-stained with filipin and GFP and LAMP2 antibodies. **C,D**. HeLa cells were

transfected with plasmids encode V5 fused ORF3a proteins in SARS-CoV or SARS-CoV-2. V5

alone served as a control. Staining and quantification was performed as described above. Bar

graphs are presented as mean $\pm$ SD. p values were determined using t' test or One-way ANOVA

test. *, $p<0.05$, **** (vs. control) or #### (CoV vs CoV-2), $p<0.0001$, n.s., not significant. Scale

bars, 5 μm. Framed images in C were enlarged 9.36 folds.

Figure 3

942
Figure 3 Structural analysis of ORF3a-VPS39 interaction interface in ORF3a

A. HeLa cells were co-transfected with mCh-VPS39 and different GFP-ORF3a constructs as

indicated and lysed in a non-denature detergent-contained buffer for co-immunoprecipitation

of mCh-VPS39. Immunoblotting was performed to detect GFP signals. **B.** The structures of

ORF3a proteins from SARS-CoV-2 (beige) and SARS-CoV (green) were shown with the cartoon

representation of the indicated residues (upper panel). The side chains are depicted as sticks

with color code for carbon (beige or green), nitrogen (blue), and oxygen (red). The presence of

W193 in SARS-CoV-2 induces an open conformation of the divergent loop that is 7.7 Å away

from S171 (dotted line), while R193 in SARS-CoV forms hydrogen bonds with D171 or Y184

(dotted lines). The lower panels represent overlays between electrostatics surface and the

cartoon representations. Positively charged surface is depicted in blue, hydrophobic surface in

white, and negatively charged surface in red. All images and measurements were performed

using PyMol V 2.4.0. **C,D.** Transfection and co-immunoprecipitation was performed as

described in A to detect the interaction between mCh-VPS39 and GFP-tagged ORF3a and its

mutants.

Figure 4

960 **Figure 4 Characterization of NPC proteins in ORF3a-inducible expression cells**

961 **A.** HeLa-Flp-In cells were established as described in Methods and treated with or without 1

962 μg/ml doxycycline for 16 hours. Immunoblotting was performed with the indicated antibodies.

963 **B.** Co-immunoprecipitation was performed with FLAG antibody (M2)-coated magnetic beads.

Endogenous VPS39 was detected by immunoblotting (arrow). **C,D.** HeLa-Flp-In cells were fixed

and stained with filipin, CellMask, and DAPI for high-content imaging. Vesicular filipin was

quantified in at least 1,000 cells per well, 6 wells per experiments, 5 independent experiments.

E,F. Antibodies against NPC1 or LAMP2 were used for immunostaining in fixed cells. Confocal

microscopy (**E**) and high-content imaging (**F**) was performed to analyze colocalization between

NPC1 and LAMP2. **G,H.** Cells were transfected with NPC2-mCherry plasmid and

immunostained with antibodies of mCherry and LAMP1. Confocal microscopy was performed,

and FIJI was used to quantify colocalization between NPC2-mCherry and LAMP1 in the cells

from two independent experiments. **I,J.** Cell culture media was collected and subjected to

immunoblotting using the antibodies indicated. FIJI was used to quantify the bands from 3

replications in each of three independent experiments. Bar graphs are presented as mean $\pm$ SD. p

values were determined using t' test (D,H, CoV-2 vs W193A) or One-way ANOVA test. * or #,

$p<0.05$, ** or ##, $p<0.01$, ***, $p<0.001$, **** (vs. control) or #### (CoV-2 vs W193A), $p<0.0001$, n.s.,

not significant. Scale bars, 5 µm. Control (Ctrl), linker peptide, CoV, SARS-CoV ORF3a, CoV-2,

SARS-CoV-2 ORF3a, W193A, SARS-CoV-2 ORF3a-W193A mutant.

Figure 5

Figure 5 Decreased endosome-to-TGN trafficking

A-D, HeLa Flp-In cells were fixed after 16-h induction of protein expression and immunostained with the indicated antibodies. Confocal microscopy (images) and high-content imaging (bar charts) were performed to analyze protein cellular distribution and colocalization.

Periphery was defined as a ring area shrunk from cell boundary with a gap distance 15 pixels

(B,C). Center was defined as a dilated circle from nuclear boundary with a distance of 20 pixels

(B). Middle was defined as the area between periphery and center (B). **E,F**. Wildtype and

VPS29-KO HeLa cell were cultured in serum-free media for 6 hours, and the media were

collected before cells were lysed. The media and cell lysates were subjected to immunoblotting

with the indicated antibodies. **G**. Vesicular filipin was measured by high-content imaging and

normalized to lysosomes in the indicated cells. **H-K,** Cells were fixed and immunostained with

the indicated antibodies. Confocal microscopy (images) and high-content imaging (bar charts)

was performed to analyze protein colocalization. Bar graphs are presented as mean $\pm$ SD. p

values were determined using t' test or One-way ANOVA test *, $p<0.05$, **, $p<0.01$, ***, $p<0.001$,

**** (vs. control) or #### (C, CoV-2 vs W193A), $p<0.0001$, n.s., not significant. Scale bars, 5 µm.

Control (Ctrl), linker peptide. CoV, SARS-CoV ORF3a. CoV-2, SARS-CoV-2 ORF3a. W193A,

SARS-CoV-2 ORF3a-W193A mutant.

Figure 6 Analysis of BMP levels

A,B. Vero E6 cells were infected with SARS-CoV-2, fixed at the indicated time points post-infection, and immunostained with BMP and LAMP1 antibodies. LAMP1-positive BMP puncta were quantified with the high-content imaging system. Representative images were shown in B.

C,D. BMP in the HeLa Flp-In cells were measured as described in A. **E.** The indicated cells were harvested and lysed for total lipid extraction. Shotgun lipidomics was performed to identify and quantify bio(monoacylglycero)phosphate (BMP) species based on the fatty acid chains, labeled as carbon number : double-bond number. BMP was normalized to protein concentrations. **F.** Cells were incubated with serum-free media containing 1% BSA and 1 μM BMP for 2 hours, fixed and stained with filipin. filipin levels were measured with high-content imaging. Bar graphs are presented as mean ± SD. p values were determined using t' test or One-way ANOVA test *, $p<0.05$. **, ##, $p<0.01$. ***, $p<0.001$. ****, ####, $p<0.0001$, n.s., not significant.

1011 Scale bars, 5 μm. Control (Ctrl), linker peptide. CoV, SARS-CoV ORF3a. CoV-2, SARS-CoV-2

1012 ORF3a. W193A, SARS-CoV-2 ORF3a-W193A mutant.

1013

A. Gained and increased proteins from Lyso-IP (ORF3a-CoV-2 vs W193A)

B. Lost and reduced proteins from Lyso-IP (ORF3a-CoV-2 vs W193A)

C. Lost proteins from Lyso-IP (ORF3a-CoV-2 vs W193A, top 20% abundant)

D. Reduced proteins from Lyso-IP (ORF3a-CoV-2 vs W193A)

1014

Figure 7. Decreased lysosome-mitochondrion interaction

A-D. Lysosomes were isolated from ORF3a and W193A Flp-In cells, and the total protein were

extracted for mass spectrometry analysis. The proteins only presenting in ORF3a lysosomes

(gain) or increased more than 2-folds compared to W193A lysosomes (increased) are shown in

the pie chart based on their cellular localization (**A**). The proteins only presenting in W193A

lysosomes (lost) or decreased more than 50% compared to W193A (reduced) are shown in **B.**

Amount the lost proteins, as defined above, top 20% abundant proteins in W193A lysosomes

are shown in **C.** The reduced proteins, as defined above, were plotted in **D**. The color codes

reflect their cellular localization or functions. **E**. The control and CoV-2 ORF3a cells were fixed

and immunostained with the indicated antibodies. Confocal microscopy was applied for

imaging. Arrows indicate the lysosome-mitochondrion physical interactions. Scale bar, 5 μm.

F,G. The Flp-In cells were fixed and immunostained with the indicated antibodies accompanied

with CellMask and DAPI. High-content imaging system was used for image collection and

analysis. Cell boundary and perinuclear area were defined based on CellMask and DAPI,

respectively. Lysosome and mitochondrial masks were defined with LAMP2 and TOMM20

signals, respectively. The colocalization between lysosomes and mitochondria were quantified

and presented as Pearson correlation coefficient. **H.** Total lipids were extracted from the

indicated cells and subjected to shotgun lipidomics analysis. PG, BMP, and Cardiolipin with the

18:1 fatty acid chains are presented. PG, phosphatidylglycerol. BMP,

bis(monoacylglycerol)phosphate. Bar graphs are presented as mean ± SD. p values were

determined using One-way ANOVA test **, $p<0.01$. ***, $p<0.001$. n.s., not significant.

1036

1037

Figure 8. Involvement of impaired lysosome-mitochondrion membrane contact sites in BMP

reduction

A.B, Wildtype and STX17-knockout (KO) HeLa cells were lysed for immnoblotting with the

indicated antibodies (**A**) or fixed for immunostaining and hign-content imaging quantification

(**B**). **C.D,** Wildtype HeLa cells were transfected with siRNA targeting the indicated autophagy

genes. Non-targeting siRNA (N.T.) served as a control. Protein levels were examined by

immunoblotting (**C**), and the BMP levels were measured by immunostaining followed by high-

content imaging (**D**) at 2 days post-transfection. **E-G,** The Flp-In cells were fixed and

immunostained with a TOMM20 (marker of mitochondrion-derived vesicles, MDVs), PDH

(maker of mitochondria but not MDVs), and LAMP2 (lysosome marker) antibodies. Confocal

microscopy was performed to identify MDVs (arrows) and the MDVs fused with lysosomes

(circles) (**E**). The MDV numbers (**F**) and the number of lysosome-fused MDVs (**G**) were

quantified using FIJI. Scale bar, 5 μm. **H.I,** HeLa cells were transfected with the siRNAs of N.T.,

DRP1, or combined MIRO1 and MIRO2 and subjected to immunoblotting (**H**) or high-content

imaging quantification (**I**). **J.K,** The Flp-In cells were seeded on glass coverslips, fixed, and

subjected to electron microscopy. Randomized cells were imaged at mitochondrion-existing

areas. Lysosome-mitochondrion membrane contact sites were identified by their membrane

structures and the distance between membranes (**J**) and quantified from 20 cells of each type

(**K**). Scale bar, 0.5 μm. **L.M,** Wildtype and VPS39-knockout (KO) HeLa cells were

immunostained and subjected to high-content imaging quantification. Bar graphs are presented

as mean ± SD. *p* values were determined using *t′* test or One-way ANOVA test. *, $p<0.05$. **,

$p<0.01$. ***, $p<0.001$. ****, $p<0.0001$. n.s., not significant.