Epigenetic modulation of NLRP6 inflammasome sensor as a therapeutic modality

to reduce necroptosis-driven gastrointestinal mucosal dysfunction in HIV/SIV

infection

Lakmini S. Premadasa[1] and Marina McDew-White[1], Luis Romero[1], Beverly Gondo[1], Jade A.

Drawec[1], Binhua Ling[1], Chioma M. Okeoma[2,3], Mahesh Mohan[1*]

[1]Southwest National Primate Research Center, Texas Biomedical Research Institute, San

Antonio, Texas, USA.

[2]Department of Pathology, Microbiology, and Immunology, New York Medical College, Valhalla,

NY 10595-1524, USA

[3]Lovelace Biomedical Institute, Albuquerque, NM 87108-5127, USA.

*Address correspondence to Mahesh Mohan, D.V.M., M.S, Ph.D., Southwest National Primate

Research Center, Texas Biomedical Research Institute, 8715 West Military Drive, San Antonio,

Texas- 78227-5302.

E-mail: mmohan@txbiomed.org

Telephone: 210-258-9170

Conflict of Interest: The authors have declared that no conflict of interest exists

Abstract

The epigenetic mechanisms driving persistent gastrointestinal mucosal dysfunction in HIV/SIV infection is an understudied topic. Using reduced-representation bisulfite sequencing, we identified HIV/SIV infection in combination anti-retroviral therapy (cART)-naive rhesus macaques (RMs) to induce marked hypomethylation throughout promoter-associated CpG islands (paCGIs) in genes related to inflammatory response (*NLRP6, cGAS*), cellular adhesion and proliferation in colonic epithelial cells (CEs). Moreover, low-dose delta-9-tetrahydrocannabinol (THC) administration reduced NLRP6 protein expression in CE by hypermethylating the *NLRP6* paCGI and blocked polyI:C induced NLRP6 upregulation in vitro. In cART suppressed SIV-infected RMs, NLRP6 protein upregulation associated with significantly increased expression of necroptosis-driving proteins; phosphorylated-RIPK3(Ser199), phosphorylated-MLKL(Thr357/Ser358), and HMGB1. Most strikingly, supplementing cART with THC effectively reduced NLRP6 and necroptosis-driving protein expression to pre-infection levels. These findings for the first time demonstrate that NLRP6 upregulation and ensuing activation of necroptosis promote HIV/SIV-induced gastrointestinal mucosal dysfunction and that epigenetic modulation using phytocannabinoids represents a feasible therapeutic modality for alleviating HIV/SIV-induced gastrointestinal inflammation and associated comorbidities.

56
57
58 **Introduction**
59
60

The gastrointestinal (GI) tract, the predominant site of human/simian immunodeficiency virus

(HIV/SIV) replication, persistence, and dissemination, plays the triple role of nutrient, water, and

anti-retroviral drug absorption while simultaneously protecting the host from pathogenic bacteria

and their metabolites. Chronic inflammation of the colon, home to trillions of commensal

bacteria, can perturb the homeostatic balance between the host and gut microbiota, causing

dysbiosis and epithelial barrier dysfunction that can lead to microbial translocation, which can

perpetuate systemic inflammation [1]. Despite viral suppression by combination anti-retroviral

therapy (cART), HIV/SIV-associated enteropathy that persists in people living with HIV (PLWH)[2]

may result in systemic inflammation that can contribute to numerous other non-AIDS-associated

comorbidities, thereby decreasing their overall quality of life [1,3-5]. Persistent GI inflammation has

also been shown to occur in people living with inflammatory bowel disease (IBD), despite the

absence of overt GI symptoms [6]. More importantly, inflammation-driven chronic intestinal barrier

impairment has been linked to extra-intestinal comorbidities, such as nonalcoholic fatty liver

disease and type-2 diabetes, and is associated with neurocognitive and inflammatory diseases[1]

including Parkinson's disease, Alzheimer's disease, and multiple sclerosis [7]. We previously

showed that an impaired intestinal barrier caused by HIV/SIV-induced intestinal inflammation

can alter signaling along the microbiota-gut-brain-axis, where dysbiotic microbes and microbial-

derived by-products can increase intestinal inflammation, increase type-I interferon responses,

endoplasmic reticulum, and oxidative stress in the neurons of the basal ganglia, and could lead

to neuroinflammation and potentially cognitive decline in PLWH [8].

While the molecular mechanisms underlying GI epithelial dysfunction in PLWH are ill-

defined, emerging evidence in other GI inflammatory diseases, such as IBD and irritable bowel

syndrome (IBS), points to epigenetic origins. In this context, aberrant DNA methylation and non-

coding RNAs (microRNAs and long non-coding RNAs) have been shown to contribute to

epithelial dysfunction in IBD and irritable bowel syndrome (IBS) [9-31]. Moreover, therapeutic drugs

to treat intestinal epithelial barrier impairment, and restore barrier function are currently

unavailable, and therefore, this topic is of significant interest for future drug development [6,7].

Due to the heritable nature of epigenetic changes, the role of DNA methylation in chronic

intestinal inflammation needs to be carefully examined, and promising immune modulation

strategies should be explored to reduce inflammation and disease progression in not only

PLWH but also those who suffer from IBD, as only 40-60% of the population respond to current

therapies [32].

Here, we used reduced representation bisulfite sequencing (RRBS) to better understand

how epigenetic regulation, through DNA methylation of colonic epithelial genes, can impact pro-

inflammatory genes and lead to chronic intestinal inflammation and epithelial dysfunction in SIV-

infected rhesus macaques (RMs). While extremely challenging in PLWH, intestinal resections

(~5 cm) were collected from SIV-infected RMs without any adverse effects, providing a unique

avenue to identify epigenetically altered protein-coding genes involved in intestinal barrier

disruption [33-39]. We found marked alterations in DNA methylation, specifically hypomethylation

occurring in promoter-associated CpG islands (paCGI), in SIV-infected RMs. These genes were

linked to epithelial cell proliferation and adhesion, apoptosis, inflammatory response, dsDNA

damage response, and oxidative stress. More importantly, we showed that long-term, low-dose

phytocannabinoid administration to SIV-infected RMs successfully modulated methylation in

paCGIs, notably hypermethylation in the CpG island of the NOD-like receptor family pyrin

domain containing 6 (NLRP6), which markedly decreased protein expression of NLRP6, a

sensor component of the NLRP6 inflammasome known to maintain intestinal epithelial barrier

function and protect against microbial invasion, relative to control RMs. Persistent NLRP6

activation was associated with significantly high mRNA and protein expression of necroptosis-

promoting RIPK1-RIPK3-MLKL pathway. In in vitro cultured human small intestinal epithelial

cells, both THC and cannabidiol (CBD) successfully blocked NLRP6 protein upregulation in

response to polyI:C and lipoteichoic acid. Lastly, while long-term cART administration alone did

not fully reduce NLRP6 protein expression in the intestines of SIV-infected RMs, supplementing

long-term cART with low-dose THC successfully reduced NLRP6 protein compared to the

increase seen one month after infection.

115   Our findings provide novel and translational insights into the epigenetic regulation of

intestinal epithelial gene expression in HIV/SIV infection and phytocannabinoid-induced

alteration of DNA methylation as an important underreported molecular mechanism attributable

to its anti-inflammatory and cellular protective properties. Benefits could also expand to other

diseases associated with intestinal inflammation, such as IBD, where interventions to address

the cause and restoration of barrier integrity have not been successful. Additionally, we show

that chronic low-dose phytocannabinoid administration could be used as a safe [40] intervention

and restorative therapy for persistent intestinal inflammation, and its use alone or as an adjunct

to cART can be beneficial for PLWH[40] with no access to cART or to those who fail to fully

suppress HIV under cART.

Results

Plasma, colon, and jejunum viral loads and histopathology

cART-naïve RMs in groups 2 and 3 had substantial plasma (0.01×10^6 to 2.6×10^8) and colon

(1×10^6 to 9.36×10^9) viral loads at 6MPI. cART-treated RMs in groups 4 and 5 had

undetectable plasma and jejunum viral loads at 5MPI (Table 1). Histopathological analysis

revealed moderate colitis and cryptitis and lymphoid hyperplasia in four group 2 and two group 3

RMs (Table 1). SIV syncytial cells were detected in one group 3 macaque. No opportunistic

infections were detected in any animals.

Chronic HIV/SIV infection is characterized by global DNA hypomethylation

concentrated mainly in promoter-associated CpG islands in colonic epithelium

Based on findings from DNA methylation profiling of CE using RRBS, relative to controls, both

VEH/SIV and THC/SIV RMs, and VEH/SIV compared to THC/SIV RMs showed more changes

in CpG methylation ratios [Δβ-value] (a decrease indicates hypomethylation change, while an

increase indicates hypermethylation change); p-value <0.05, absolute Δβ-value >0.1) distributed

throughout all chromosomes, with the least amount of differentially methylated CpG sites

(DMSs) occurring in chromosome Y (Figure 1A). All data presented below is described in the

flow chart in Figure S1. As shown in Figure 1A, compared to controls, THC/SIV RMs had more

DMSs than VEH/SIV RMs. In addition, THC/SIV RMs showed more hypomethylation (values

below 0) changes in all comparisons than hypermethylation (values above 0) (Figure 1A). Only

DMSs within annotated genes (intergenic and unannotated regions excluded) were analyzed.

Compared to controls, VEH/SIV RMs had a total of 13,831 [out of 94,279 (14.67%)] DMSs

within annotated genes (Figure 1B). A similar number of DMSs were detected in THC/SIV

compared to control RMs; 13.59% of DMSs were within annotated genes (16,730 out of

123,135 DMSs; Figure 1C). Lastly, compared to THC/SIV, VEH/SIV RMs had a total of 6,796

[out of 64,069 (10.61%)] DMSs within annotated genes (Figure 1D).

We focused on methylation changes that occurred within the CpG-rich genomic regions

called CpG islands (CGIs) that also included the promoter, exon, and intron intragenic regions.

CGIs that spanned a promoter region (continuing into the exon and intron regions) were

classified as paCGIs. Nearly 70% of gene promoters reside within CGIs and are normally

unmethylated [41-43]. Interestingly, paCGIs are devoid of common promoter elements (TATA

boxes and core promoter elements), but contain binding sites for transcription factors, which

help their respective genes to maintain a transcriptionally permissive state. Out of all the DMSs

within annotated genes, VEH/SIV RMs had 43.79% located within CGIs (6,056 out of 13,831),

relative to control RMs. These DMSs were identified in 2,688 genes, of which 2,504 genes had

5,084 (83.95%) DMSs within paCGIs (Figure 1E and 1H). Relative to controls, THC/SIV RMs

showed 7,526 out of 16,730 (44.99%) DMSs in CGIs within annotated genes. There were 2,622

unique genes that had 80.85% (6,085 out of 7,526) of DMSs in paCGIs within annotated genes

(Figure 1F and 1H). Finally, VEH/SIV RMs had 1,686 (24.81%) DMSs within 540 CGIs of

annotated genes and contained 1,021 (60.56%) DMSs within the paCGIs of 343 genes, relative

to THC/SIV RMs (Figure 1G and 1H).

Intriguingly, untreated HIV/SIV infection was associated with more hypomethylation

changes in the paCGI [Figure 1B, 1C ($\Delta\beta$-values to the left of 0 on X-axis), 1H, and 1I ($\Delta\beta$-

values below 0)]. The largest $\Delta\beta$-value change detected was a decrease of 20% (Figure 1B and

1C), and both VEH/SIV and THC/SIV RMs had more promoters that were hypomethylated than

exons and introns in CGIs (Figure 1H and 1I). Relative to controls, THC/SIV RMs also showed

increased hypermethylation in the promoters, exons, and introns than VEH/SIV RMs (Figure

1H). Interestingly, all comparisons showed a large range of hypomethylation changes that

occurred in the paCGI (10-90% $\Delta\beta$-value decrease), but hypermethylation changes were

restricted to less than 50% (10-45% $\Delta\beta$-value increase, Figure 1I). There was also a significant

difference in the total number and hypomethylation changes detected in the paCGIs of VEH/SIV

and THC/SIV RMs compared to controls and in VEH/SIV relative to THC/SIV RMs (Figure 1I).

The same was true for the total number and hypermethylation changes that occurred in the

paCGIs in VEH/SIV and THC/SIV compared to controls, and in VEH/SIV vs. THC/SIV RMs,

relative to THC/SIV compared to controls (Figure 1I).

Figure S2A-C shows clustering heatmaps of the top 100 differentially methylated CpG

sites, including those located in intergenic regions, demonstrating major differences in

methylation among the three different groups. The top 100 DMSs are mostly hypomethylated in

VEH/SIV and THC/SIV relative to control RMs (Figures S2A and S2B). When compared to

THC/SIV, the top 100 DMSs in VEH/SIV RMs showed a range of methylation changes (Figure

S2C), with hypomethylation predominating. The Pearson's correlation coefficients for all CpG

sites between VEH/SIV vs. control, THC/SIV vs. control, and VEH/SIV vs. THC/SIV RMs were

0.9400, 0.9358, and 0.9359, respectively (Figure S2D-F). These values look very strong,

suggesting that each point where the methylation value is different indicates a major treatment

effect.

Hypomethylation of paCGI sites resulted in upregulation of their corresponding

gene expression, including the machinery responsible for demethylation and de

novo methylation

Next, we assessed the effect of the cumulative Δβ-value (average of all DMSs Δβ-values per

gene) of DMSs in individual paCGIs on gene expression. We compared the number of genes

that had a cumulative Δβ-value (hypo- or hypermethylation) in the paCGIs to the number of

genes that were differentially expressed in our recently published manuscript [44] that utilized a

human microarray platform designed to profile mRNAs and lncRNAs in the same RMs using CE

total RNA. Methylation changes in the paCGIs of genes showed an inverse association with

gene regulation (i.e., hypomethylated paCGIs and respective gene upregulation or

hypermethylation paCGIs and respective gene downregulation) in 11.5% [751 out of 6533

differentially expressed genes (DEGs)] and 11.4% (480 out of 4229 DEGs) of genes in VEH/SIV

and THC/SIV RMs, respectively, compared to controls (Figure 1J and 1K). Hypomethylation

was better associated with gene expression changes (increased expression) than

hypermethylation, as 99% (743/751 genes were hypomethylated and upregulated, compared to

8/751 genes that were hypermethylated and downregulated) (Figure 1J) and 97.5% (468/480

genes were hypomethylated and upregulated, compared to 12/480 genes that were

hypermethylated and downregulated) (Figure 1K) of all genes with DMSs in the paCGIs were

hypomethylated in VEH/SIV and THC/SIV RMs, respectively, compared to controls.

Surprisingly, methylation changes in the paCGIs had minimal influence on gene expression in

VEH/SIV when compared to THC/SIV RMs, as only 4 genes with methylation changes in the

paCGIs inversely impacted gene expression (Figure 1L). We also found that hypermethylation

in the paCGIs was associated with downregulation of gene expression, as well as an

association between hypomethylation and upregulation of gene expression. Consistent with

these findings, mRNA upregulation of the enzymes involved in demethylation (TET-2; 1.76-fold)

(Figure 1M) and *de novo* methylation [DNMT3a; 2.4-fold (Figure 1O) and DNMT3b; 2.06-fold)

(Figure 1P)] was detected in VEH/SIV (n=5) but not THC/SIV (n=6) compared to control RMs

(n=6). Although upregulated, TET-3 mRNA did not show statistical significance (Figure 1N).

Long-term, low-dose THC administration modulated gene expression associated

with intercellular signaling, communication, junction organization, and NLRP6

inflammasome assembly

Using Gene Ontology (GO) enrichment analysis, we investigated the effect of the cumulative

Δβ-value in the paCGIs of protein-coding genes on the biological processes in CE during

HIV/SIV infection. HIV/SIV infection-induced hypomethylation affected the expression of genes

linked to epithelial cell proliferation and its regulation, cell growth, cellular response to

transforming growth factor-β (TGF-β), regulation of and/or response to oxidative stress,

apoptotic signaling pathway, programmed cell death, canonical Wnt signaling pathway, cell

communication and signaling, and MAPK cascade (Figures 2A and 2B). Relative to controls,

phytocannabinoid administration had little impact on the number of genes hypomethylated in the

above biological processes, compared to VEH/SIV RMs (Figures 2A and 2B). When compared

to THC/SIV, VEH/SIV RMs had significantly more hypomethylated genes involved in positive

regulation of cell communication (Figure 2B). These animals also had hypermethylated genes

involved in the overall and negative regulation of cell communication, regulation of cell signaling,

and MAPK cascade (Figure 2B).

Not surprisingly, compared to controls, only VEH/SIV RMs showed hypomethylation

changes in genes involved in epithelial cell differentiation, positive regulation of cell population

proliferation, and regulation of epithelial cell migration, an indirect indicator of host response to

epithelial cell loss (Figure 2C). Notably, genes responsible for the regulation of oxidative stress-

induced cell death and autophagy were hypomethylated in THC/SIV relative to control RMs

(Figure 2C). Compared to THC/SIV, VEH/SIV RMs showed hypomethylation of genes involved

in cell junction organization and cell-to-cell signaling (Figure 2C). Most strikingly,

hypermethylation of genes responsible for the negative regulation of Wnt protein secretion,

positive regulation of MAPK cascade, and, more importantly, NLRP6 inflammasome complex

assembly (Figure 2D) were detected in THC/SIV relative to control RMs (Figure 3D).

Using a supervised analysis, we examined genes that showed differential methylation

throughout their respective paCGIs. Relative to controls, VEH/SIV RMs had DMSs (mostly

hypomethylation) in genes associated with inflammatory response [C-C motif chemokine ligand

12 (*CXCL12*), toll-interacting protein (*TOLLIP*), NLR family pyrin domain containing 6 (*NLRP6*;

Figure 2E), leptin (*LEP*; Figure 2F), Mab-21 Domain Containing 1 (*MB21D1*) also known as

cyclic GMP-AMP synthase (cGAS)], cellular adhesion [claudin 6 and 11, intercellular adhesion

molecule 1 (*ICAM1*), protein kinase C-beta (*PRKCB*), protocadherin 17 and 19 (*PCDH17* and

PCDH19; Figure 2G), adrenocorticotropic receptor beta 1 (*ADRB1*), cadherin 7 (*CDH7*; Figure

2H), genes associated with protection against oxidative stress [cytoglobin (*CYGB*), glutathione

peroxidase 3 (*GPX3*)], colonic epithelial cell proliferation [Wnt inhibitory factor 1 (*WIF1*), soluble

frizzled protein 1 (*SFRP1*; Figure 2I), telomerase reverse transcriptase (*TERT*; Figure 2J), heart

and neural crest derivatives expressed 2 (*HAND2*)], and apoptosis [death-associated protein

kinase 1 (*DAPK1*), BCL2-Like-1 (*BCL2L1*)]. Table S1 shows the respective number of DMSs for

each gene, as well as the average change in methylation due to these DMSs throughout the

CGI. The average methylation changes of 1-14 DMSs in the CGI for the genes above were

between -40.7% and 12.3% (columns 4, 6, and 8 of Table S1).

In contrast, relative to control RMs, although THC/SIV RMs showed DMSs in the genes

described above, key genes associated with the inflammatory response (Table S1) showed

more hypermethylated CpG sites (hyper-MSs), compared to VEH/SIV RMs, specifically *NLRP6*

(38 hyper-MSs and 6 hypo-MSs; Figure 2E), and *MB21D1* or *cGAS* (9 hyper-MSs and 1 hypo-

MS). The average methylation differences were 12.5% (*NLRP6*) and 24.2% (*MB21D1*).

Interestingly, while there were more hyper-MSs in *LEP* (Figure 2F), in THC/SIV than VEH/SIV

relative to control RMs, both had an average positive methylation change of 2.6% (VEH/SIV)

and 15.6% (THC/SIV). As with *LEP*, while *PCDH17* had more hyper-MSs (52; Figure 2G, Table

S1), there was a positive methylation change in both VEH/SIV and THC/SIV compared to

control RMs (3.7% and 24.1%, respectively). Genes associated with cellular adhesion (Table

S1) showed a similar pattern of methylation difference in THC/SIV and VEH/SIV compared to

control RMs, except for *CDH7* (Figure 2H), which had 20 hyper-MSs, one hypo-MS, and a

methylation change of 24.9% in THC/SIV compared to control RMs. Like VEH/SIV RMs, genes

associated with oxidative stress and apoptosis (Table S1) also showed similar methylation

changes in THC/SIV compared to controls. Finally, *SFRP1* (18 hyper-MSs and 3 hypo-MSs;

Figure 2I), *TERT* (10 hyper-MSs and one hypo-MS; Figure 2J), and *HAND2* (one hyper-MS),

associated with colonocyte proliferation, were hypermethylated in THC/SIV RMs compared to

controls, as opposed to VEH/SIV RMs.

Relative to THC/SIV, hypomethylation occurred in all the genes discussed above, with

an average methylation difference of between -28 and -11% in VEH/SIV RMs. Since our focus

was on methylation changes that occurred in paCGIs, we investigated the average methylation

changes that occurred in genes described above that had at least 10% of the CpG sites

differentially methylated, which included *NLRP6*, *LEP*, *CDH7*, and *SFRP1* (Figure 2E, 2F, 2H,

and 2I, Table S1). Unlike THC/SIV RMs, relative to control RMs, for the genes listed above,

VEH/SIV RMs displayed a negative (hypomethylation) methylation change in *TERT and SFRP1*

(Figure 2I, 2J and 2K), while *NLRP6* and *LEP* had a positive methylation change (Figure 2K).

Interestingly, while no DMSs were detected in the paCGIs of *CDH7* in VEH/SIV compared to

control RMs, several DMSs were detected in VEH/SIV relative to THC/SIV RMs.

Long-term, low-dose THC-induced hypermethylation of CpGs in the paCGI of

NLRP6 resulted in decreased NLRP6 protein expression in the colon of SIV-

infected RMs

The GO identification of a cluster of hypermethylated genes that were involved in NLRP6

inflammasome assembly (Figure 2D) prompted us to focus further on methylation changes in

NLRP6, the gene responsible for protecting the intestinal epithelial barrier from harmful gut

microbes and maintaining gut homeostasis, through inflammasome activation. However, in IBD,

although the host responds to microbial invasion initially by activating NLRP6, the persistence of

the inflammatory trigger drives NLRP6 hyperactivation, inflammasome activation, and

consequently damaging colonic inflammation and barrier disruption [45]. Within the *NLRP6* CGI,

VEH/SIV RMs showed a total of 14 DMSs (Figures 3A and 3B), relative to controls. These

included eight hypomethylated (boxed region in Figure 3B) and six hypermethylated sites.

Within the promoter region, two sites each were hypomethylated (green arrows in Figure 3B)

and hypermethylated (red arrows in Figure 3B). Relative to controls, THC/SIV RMs showed a

significantly higher number of DMSs in *NLRP6* CGI than seen in VEH/SIV RMs, with a total of

44 DMSs (Figure 3A and 3C). Out of these, 38 sites were hypermethylated (boxed region in

Figure 3C), and 6 were hypomethylated. Strikingly, except for one (hypomethylated) (green

arrow in Figure 3C), 26 sites were hypermethylated (red curly bracket in Figure 3C) in the

paCGI region. Lastly, when compared to THC/SIV, VEH/SIV RMs had 13 DMSs in the CGI of

NLRP6 (Figure 3A and 3D), out of which 12 were significantly hypomethylated (boxed region in

Figure 3D) and only one site located in the intron was hypermethylated (red arrow in Figure 3D).

The average Δβ-value in the paCGI of NLRP6 was not significantly different in VEH/SIV relative

to controls (Figure 3E) or THC/SIV RMs (Figure 3G). Inversely, there was a significant increase

in the Δβ-value of CpG sites of *NLRP6* paCGI in THC/SIV compared to control RMs (Figure 3F).

Since methylation changes alter gene expression, we next investigated NLRP6 protein

expression levels in the colon of all three groups. Since microarray data did not detect

significant changes in *NRLP6* mRNA expression, we focused on NLRP6 protein expression

because not all genes show similar changes at the mRNA and protein levels. NLRP6 protein

expression was detected in both the nucleus and cytoplasm, with a stronger signal in the

nucleus of CE and lamina propria mononuclear cells (LPMNCs). We found an inverse

relationship between hypermethylation changes in the paCGI of NLPR6 and protein expression,

although NLRP6 staining intensity was not significantly different in VEH/SIV RMs (n=6; Figure 3I

and 3K) or THC/SIV RMs (n=6; Figure 3J and 3K) compared to controls (n=5; Figure 3H and

3K). Nevertheless, NLRP6 staining intensity had a decreasing trend exclusively in CE of

THC/SIV RMs (Figure 3J and 3K) relative to controls and VEH/SIV RMs. Similar to the CE,

NLRP6 protein expression was reduced in LPMNCs (Figure 3L).

Acute HIV/SIV infection significantly increased NLRP6 protein expression, while

low-dose THC in combination with cART successfully decreased its expression in

the jejunum of SIV-infected RMs

Next, we looked at the effect of acute HIV/SIV infection [1 month post infection (MPI)] on NLRP6

protein expression and if cART alone or in combination with THC could decrease NLRP6

protein expression. Here, we focused on the jejunum, as it plays a critical role in both nutrient

and cART drug absorption, and its dysfunction can negatively impact both functions. Moreover,

high NLRP6 protein expression has been confirmed in the jejunum of mice [46] and humans [47,48].

Longitudinally, acute HIV/SIV infection (1MPI; n=8/group; Figure 3N, 3Q, and 3S) resulted in a

significant increase in NLRP6 protein expression compared to pre-infection (Figure 3M, 3P, and

3S). At 5MPI, NLRP6 expression in cART-suppressed (VEH/SIV/cART; n=8) RMs did not differ

from their 1MPI timepoint but significantly elevated from the pre-infection timepoint (Figure 3O,

3N, 3M, and 3S). Strikingly, THC/SIV/cART RMs (n=8) showed a significant decrease in NLRP6

protein expression from their respective 1MPI timepoints (Figure 3R, 3Q, and 3S) but showed

no significant decrease from VEH/SIV/cART RMs at 5MPI or the pre-infection timepoint (Figure

3O, 3P, and 3S). THC-induced reductions in NLRP6 protein expression in the jejunum LPMNCs

showed a similar trend (Figure 3T) to that observed in the epithelium (Figure 3S).

Phytocannabinoids blocked polyI:C and lipoteichoic acid-induced NLRP6 protein

upregulation in vitro and significantly increased mRNA expression of the NLRP6

deubiquitinating enzyme CYLD in the CE in vivo

To determine if phytocannabinoids directly blocked NLRP6 protein expression, we separately

added 3 µM THC and CBD to human small intestinal epithelial cultures after transfection with

2µg/mL of polyI:C followed by treatment with lipoteichoic acid (LTA) (10µg/mL) 4 h later. It is

clear from Figure 4B, 4C, and 4I that polyI:C alone or in combination with LTA significantly

increased NLRP6 protein expression in the cytoplasm (white arrows in Figure 4B and 4C)

compared to cells treated with DMSO (Figure 4A and 4I). Interestingly, pretreatment for 1 h with

THC (Figure 4D and 4I) or CBD (Figure 4G and 4I) or both (Figure 4H and 4I) successfully

blocked polyI:C and LTA-induced NLRP6 protein upregulation. Treatment of cells with an

antagonist of the cannabinoid receptor-1 (AM251) (Figure 4E and 4I) or cannabinoid receptor-2

(AM630) (Figure 4F and 4I) did not alter THC's ability to counter polyI:C and LTA-induced

NLRP6 protein upregulation. Therefore, future studies are needed to determine the role of other

receptors such as GPR55, PPARs and TRPV ion channels in cannabinoid blockade of NLRP6

protein induction.

Next, we examined if the gene expression levels of the *NLRP6* inflammasome

components and proinflammatory cytokines changed consequent to reduction of *NLRP6* protein

expression in response to hypermethylation of the paCGI. Intriguingly, mining our recently

published CE microarray data from the same RMs [44] for genes that showed differential

expression by at least 1.5-fold change and an adjusted p≤0.05 confirmed statistically significant

upregulation of *CYLD*, a deubiquitinase shown to remove K63-linked ubiquitin residues on

NLRP6 [45] and attenuate its activity. *CYLD* mRNA was significantly upregulated in CE of

THC/SIV (~2.7-fold) but not VEH/SIV RMs relative to controls (Figure 4J). Similarly, *DHX15*, an

RNA helicase and double-stranded viral RNA sensor shown to interact with NLRP6 and trigger

NLRP6 inflammasome assembly and activation [49], was significantly upregulated in CE of

VEH/SIV but not in THC/SIV RMs relative to controls (Figure 4K). In addition, while mRNA

expression of caspase-1 (*CASP1*, Figure 4L) and interleukin-18 (*IL-18*, Figure 4M) increased

significantly, the opposite trend (downregulated) was detected with interleukin-1β (*IL-1β*, Figure

4N) in CE of SIV-infected RMs irrespective of VEH or THC treatment relative to controls. Finally,

mRNA expression of the membrane pore-forming protein, Gasdermin D (*GSDMD*), the

downstream executor of pyroptosis, was significantly reduced in CE of both VEH/SIV and

THC/SIV relative to control RMs (Figure 4O).

THC-induced NLRP6 promoter hypermethylation associated with reduced necroptosis-

driving gene signature in the CE of SIV-infected RMs

Although *CASP1* and *IL18* mRNA expression was significantly upregulated in CE of both

VEH/SIV and THC/SIV RMs, the significantly decreased *GSDMD* mRNA expression in both

groups suggested that pyroptosis may not be the predominant pathway driving CE dysfunction.

Nevertheless, *NLRP6* has been shown to increase the expression of mixed lineage kinase

domain like pseudokinase protein (*MLKL*) and receptor-interacting serine/threonine-protein

kinase-3 (*RIPK3*) to induce necroptosis [50], a caspase-independent inflammatory mode of cell

death. Interestingly, again mining our recently published microarray data [44] for genes that

showed differential expression by at least 1.5-fold change and an adjusted p≤0.05 confirmed

statistically significant upregulation of *RIPK1*, *RIPK3*, and *MLKL* in CE of VEH/SIV RMs (Figure

4P-R). While expression of *RIPK1* mRNA showed statistically significant increase in both groups

(Figure 4P), mRNA expression of *RIPK3* and *MLKL* showed statistically significant increase only

in CE of VEH/SIV (Figure 4Q and 4R) compared to control RMs. Moreover, *RIPK3*,

phosphorylated on Ser227 and Ser199, plays a crucial role in the activation of *MLKL* by

phosphorylating it on Thr357 and Ser358 to enable its loading onto the necrosome [51]. In this

context, *PPM1B*, a phosphatase known to dephosphorylate *RIPK3* [51] was also significantly

upregulated only in CE of VEH/SIV RMs (Figure 4S), indirectly suggesting the presence of p-

RIPK3. Most strikingly, mRNA expression of *ADAM17* (Figure 5T), a positive regulator of

necroptosis [52], *TRPM7* (Figure 5U), a downstream target of *MLKL* that promotes its membrane

localization and Ca^{2+} ion influx for initiating necroptosis [53], and *HMGB1* (Figure 5V), a danger-

associated molecular pattern (DAMP) protein, known to propagate inflammation associated with

both pyroptotic and necroptotic cell death [54] was significantly increased only in CE of VEH/SIV

RMs.

We previously reported significantly reduced interferon (IFN)-stimulated gene expression

(ISG) in whole colon [39], CE [44] and basal ganglia (brain) [8] of THC/SIV RMs. Nevertheless, the

mechanism of THC action remained unknown. *MB21D1,* or *cGAS,* is an important DNA sensor

that drives type-I IFN responses [55]. In this context, within the *MB21D1* CGI (Figure S3A),

VEH/SIV RMs showed a total of 4 DMSs (Figure 4W1), relative to controls. These included

three hypomethylated (Figure 4W1) and one hypermethylated site. Within the paCGI, two sites

were hypomethylated (green arrows in Figure 4W1) and one hypermethylated (red arrow in

Figure 5W1). Relative to controls, THC/SIV RMs showed a significantly higher number of DMSs

in *MB21D1* CGI than seen in VEH/SIV RMs, with a total of 13 DMSs (Figure 4W2). Out of these,

nine sites, all in the paCGI region, were hypermethylated (boxed region and red curly bracket in

Figure 4W2), and four were hypomethylated (green curly bracket in Figure 4W2). Lastly, when

compared to THC/SIV, VEH/SIV RMs had six DMSs in the CGI of *MB21D1* (Figure 4W3), all of

which were in the paCGI region (boxed region in Figure 4W3). The average $\Delta\beta$-value in the

paCGI of *MB21D1* was not significantly different in VEH/SIV relative to controls (Figure 4X1). In

contrast, there was a significant increase in the $\Delta\beta$-value of CpG sites of *MB21D1* paCGI in

THC/SIV compared to control (Figure 4X2) and VEH/SIV (Figure 4X3) RMs. Further, the

expression of *TBK1*, a signaling adapter protein that transduces the signal emanating from

MB21D1 [56], was significantly increased in CE of VEH/SIV but not THC/SIV RMs (Figure 4Y).

Finally, *PQB1*, a proximal sensor of *MB21D1*-dependent increase in type-I IFN response to HIV-

1 infection [57], was also significantly increased in CE of VEH/SIV but not THC/SIV RMs (Figure

4Z).

Protein expression of necroptosis mediators p-RIPK3 (Ser-199), p-MLKL (Thr-357/Ser-358), and HMGB1 is significantly increased in the intestinal epithelium of SIV-infected RMs

Activated RIPK1 promotes RIPK3 phosphorylation, which then autophosphorylates itself on Ser199 (required for kinase function) and Ser227 (required for necroptosis), two sites crucial for the activation of MLKL. Interestingly, p-RIPK3(Ser199) showed significantly higher expression in CE of both ART-naïve VEH/SIV and THC/SIV relative to control RMs (Figure 5A to 5D) but not in ART-experienced RMs (Figure S3B-3E). In this context, protein expression of PPM1B, a phosphatase that inactivates RIPK3 by dephosphorylation, was significantly increased in CE of both ART-naïve VEH/SIV and THC/SIV relative to control RMs (Figure 5E to 5H) but not in ART-experienced RMs (Figure S3F-3I). The high expression of PPMB1 indirectly indicates the presence of activated p-RIPK3. Note that for both p-RIPK3(Ser199) and PPM1B, protein expression was slightly higher in VEH/SIV than THC/SIV relative to control RMs.

Consistent with high *MLKL* mRNA expression in CE of VEH/SIV RMs, protein expression of p-MLKL(Thr357), a marker of necroptosis, was significantly upregulated with intense cytoplasmic expression in CE of ART-naïve (Figures 5J and 5L) and ART-experienced VEH/SIV (Figure 5N and 5P) relative to control (Figures 5I and 5M) RMs. p-MLKL(Thr357) expression in CE of THC/SIV RMs was significantly higher than controls (Figures 5K and 5L). Most notably, p-MLKL(Thr357) protein expression was significantly reduced in CE of THC/SIV compared to VEH/SIV RMs in both ART-naïve (Figure 5K and 5L) and ART-experienced RMs (Figure 5O and 5P).

Likewise, p-MLKL(Ser358) protein expression was significantly increased in CE of both VEH/SIV and THC/SIV relative to control RMs (Figures 6A to 6D) in CE. Similar to p-MLKL(Thr357), protein expression of p-MLKL(Ser358) was more intense in the cytoplasm of VEH/SIV but not in THC/SIV compared to control RMs (Figure 6B and 6C). Notably, at 5MPI, p-

MLKL(Ser358) protein expression was significantly reduced in THC/SIV/ART compared to

VEH/SIV/ART RMs (Figure 6F, 6G, and 6H). While no significant differences in protein

expression of p-RIPK3(Ser199) were detected in the jejunum of SIV/ART RMs, it is possible that

p-RIPK3(Ser227) [51] and the Tam family of receptor tyrosine kinases [58] may be the predominant

kinases phosphorylating MLKL in the jejunum. Finally, expression of HMGB1, an important

DAMP released upon pyroptotic and necroptotic cell death and an initiator of secondary

inflammatory responses in neighboring cells[59], showed a marked but statistically nonsignificant

reduction in CE of ART-naïve THC/SIV RMs compared to VEH/SIV RMs (Figure 6I to 6L).

Nevertheless, unlike colon, protein expression of HMGB1 was significantly increased in the

jejunum epithelium of VEH/SIV/ART RMs compared to pre-infection and THC/SIV/ART RMs

(Figure 6M to 6P). Note the restoration of HMGB1 protein expression to pre-infection levels by

low-dose THC at 5MPI (Figure 6P). These findings demonstrate, for the first time, the activation

of necroptosis and its contribution to intestinal dysfunction in HIV/SIV infection under

suppressive cART and provide novel clinically relevant insights into how low-dose

phytocannabinoids, potentially through epigenetic modulation of NLRP6, can reduce

necroptosis signaling potentially via modulation of the RIPK3-MLKL pathway (Figure 7) that may

help reduce intestinal inflammation in PLWH and, by extension, IBD patients.

Discussion

481
482
Intestinal epithelial barrier disruption, triggered by mucosal immune dysfunction, is a common

sequela to HIV/SIV infection despite peripheral viral suppression by cART, which leads to

systemic inflammation, a major contributor to HIV-associated comorbidities [1]. Persistent

intestinal inflammation can trigger and sustain the accumulation of epigenetic modifications,

particularly alterations in levels of promoter DNA methylation in genes linked to epithelial

proliferation [27] leading to enterocyte senescence, epithelial barrier dysfunction, and dysbiosis [60].

Nevertheless, epigenetic control of epithelial barrier function in HIV/SIV infection remains an understudied topic. In this context, chronic intestinal inflammation and epigenetic alterations involving DNA methylation in gene promoters, together with transcriptome changes in intestinal epithelial cells, have been associated with IBD pathogenesis [61]. Moreover, since epigenetic signatures are cell type-specific, separating out enough purified epithelial cells from human colorectal pinch biopsies (maximum n=3 biopsies) is challenging. The feasibility of obtaining intact colon resections (~5 cm) without any adverse effects from SIV-infected RMs allows the isolation of ~85-95% pure epithelial cells and offers unique avenues to identify epigenetically altered genes involved in critical GI mucosal abnormalities. While we found significant global hypomethylation throughout the whole genome during chronic HIV/SIV infection, chronic low-dose phytocannabinoid (THC) administration caused significantly more hypermethylation and effectively increased methylation in paCGIs of genes associated with the inflammatory response (*NLRP6, MB21D1, LEP*), cellular adhesion (*CDH7*), and colonic epithelial cell proliferation (*SFRP1* and *TERT*).

HIV/SIV infection resulted in a significant decrease in methylation levels throughout the genome and specifically in paCGIs. Strikingly, chronic low-dose THC administration caused significantly more changes in methylation, both hypo- and hyper- methylation, than that observed in VEH/SIV RMs compared to controls. Remarkably, hypomethylation of just one DMS in the paCGI had an influence on gene expression (upregulation). This finding leads us to speculate that the traditional idea of methylation being inversely correlated to gene expression is gene location specific. While the paCGIs are regulatory regions, methylation changes in enhancer regions might have more influence on gene expression than paCGIs [62,63]. Interestingly, methylation changes that occurred outside of a gene region or mapped to an unannotated region were found to be in gene enhancers, which potentially could influence gene expression [63]. Since our analysis only looked for DMSs within a CGI, and if it was in a gene promoter, exon, or intron, we could not determine if the ~86% of unannotated sites excluded

from the analysis were a part of gene enhancers and the genes they regulated. Differential

methylation of enhancers has been implicated in osteoarthritis, Alzheimer's disease (AD), and

multiple sclerosis [64]. Therefore, further work to identify the unannotated DMSs that are an

integral part of gene enhancers with potential role(s) in HIV/SIV-induced intestinal dysfunction is

needed.

Significant hypomethylation detected in the paCGIs of genes associated with

inflammation, cellular adhesion, oxidative stress, cell proliferation, and apoptosis in CE of

VEH/SIV RMs suggests that alterations in the epigenetic landscape may facilitate their

increased expression, leading to the persistence of a proinflammatory environment. These

findings are consistent with increasing evidence pointing towards an association between DNA

hypomethylation and increased expression of proinflammatory genes (76-78), leading to

oxidative DNA damage, epithelial cellular senescence, barrier dysfunction, microbial

translocation, and systemic inflammation. Interestingly, compared to controls, THC/SIV RMs

had significantly more hypermethylation in the paCGI region of genes associated with

inflammatory responses, cellular adhesion, and colonic epithelial cell proliferation, specifically

NLRP6, *CDH7*, *SFRP1*, and *TERT*. These genes showed increased methylation levels in CE of

THC/SIV compared to control RMs that accounted for ~4-23% of CpG sites in their respective

CGIs, and the average methylation difference from all methylation changes in these genes was

~18%. More importantly, THC hypermethylated promotor CpG sites of *MB21D1*, also known as

cGAS, which interacts with *PQBP1* following its binding to retroviral reverse transcribed DNA in

the cytosol to activate the cGAS-STING pathway, leading to increased type-I interferon (IFN)

production. Interestingly, in addition to promoter CpG hypermethylation of *MB21D1*, *PQBP1*

gene expression was also significantly reduced in the CE of THE/SIV RMs, which may partly

explain the reduced type-I interferon gene expression in the CE of THC/SIV RMs we reported

previously [39,44], suggesting that phytocannabinoids are effective in inhibiting cGAS-STING

signaling, a pathway hyperactivated in ulcerative colitis leading to dysregulated intestinal

epithelial cell integrity, autophagy, and immune/inflammatory responses [55]. Interestingly, gene

ontology analysis identified significant hypomethylation of genes that regulate oxidative stress-

induced cell death and autophagy in CE of THC/SIV RMs. While cannabinoids are well-known

to induce autophagy pathways to reduce chronic inflammatory responses [65], our novel findings

demonstrate the involvement of epigenetic mechanisms underlying the enhancement of

autophagy (promoter DNA hypomethylation) via reduction of *MB21D1*-mediated type-I IFN

expression (promoter DNA hypermethylation) by cannabinoids. Thus, epigenetic modulation

represents a key mechanism by which phytocannabinoids alter inflammatory

signaling/response, epithelial cell adhesion and proliferation, and autophagy in HIV/SIV

infection.

When focusing on epigenetic changes in genes that regulate epithelial barrier

maintenance and homeostasis, inflammasome components and sensors, *NLRP6*, stood out

prominently. A critical function of *NLRP6* is to protect the intestinal epithelial barrier from

invading microbes or viruses, where it shows the highest expression [66,67]. *NLRP6* is thought to

regulate intestinal microbiome homeostasis through its ability to activate other proinflammatory

and antimicrobial genes [46,50,66,68]. In return, the gut microbiome modulates NLRP6

inflammasome through their microbial metabolites, prompting *NLRP6* to activate the production

of antimicrobial peptides by intestinal epithelial cells (paneth cells) or triggering goblet cells to

induce mucin exocytosis [66], [48]. While *NLRP6* has been shown to maintain intestinal

homeostasis, its perpetual activation due to inadequate regulation, as previously shown in

active ileal Crohn's disease (CD) (131-fold high) and colonic CD patients (3.9-fold high) [69], can

exacerbate inflammation, leading to gastrointestinal disease and even cancer. Quite

unexpectedly, we did not see a change in *NLRP6* mRNA expression in either SIV-infected RM

group compared to controls or to each other. Aberrant hypomethylation of *NLRP6* can expedite

inflammasome assembly, thereby activating inflammatory signaling pathways leading to CE

apoptosis, as seen in Kawasaki disease (KD). Increased *NLRP4*, *NLRP12*, and *IL-1β* gene

expression in KD patients was accompanied by hypomethylation of *NLRP4* and *NLRP12* [70].

Interestingly, treatment with intravenous immunoglobulin restored the gene expression and

methylation levels of *NLRP4*, *NLRP12*, and *IL-1β*. In the present study, we show that chronic

SIV-infection in cART-naïve RMs neither caused significant hypomethylation changes to *NLRP6*

paCGI (2 CpG sites compared to controls and 3 CpG sites compared to THC/SIV RMs) nor

increased protein expression significantly in CEs, compared to controls. Nevertheless,

hypomethylation of six other CpG sites in the exons and introns may impact *NLRP6* expression.

Contrariwise, chronic administration of low-dose THC to chronically SIV-infected RMs resulted

in a significant hypermethylation of the *NLRP6* paCGI compared to controls, which resulted in

markedly decreased *NLRP6* protein expression in the epithelium and lamina propria cells

compared to both control and VEH/SIV RMs.

The decrease in NLRP6 protein expression through hypermethylation was an interesting

finding that led us to question the effect of cART on *NLRP6* expression. During acute HIV/SIV

infection (1MPI), there was a significant increase in NLRP6 protein expression in the jejunum

epithelium, with cART initiated 2 weeks post infection. Noteworthy, at the chronic stage of

HIV/SIV infection (5MPI), after viral suppression for ~4.5 months, cART alone had no effect on

NLRP6 expression. However, and most importantly, phytocannabinoid administration, in

combination with cART, resulted in significantly decreased NLRP6 protein expression in

chronically infected RMs, bringing NLRP6 protein expression down to pre-infection levels. The

significantly high NLRP6 protein expression detected in the jejunum of THC/SIV/cART RMs at

1MPI despite initiating THC before SIV infection suggests that phytocannabinoids may produce

better anti-inflammatory effects when administered in combination with cART. The successful

lowering of NLRP6 protein expression at 5MPI also demonstrates that cART alone, regardless

of undetectable viral loads in plasma and tissues, is not sufficient to restore intestinal

homeostasis, but long-term, low-dose phytocannabinoids represent an excellent option to be

used as an adjunct to cART to reduce damaging intestinal inflammatory responses that persist

in PLWH. Folate produced by *Lactobacillus* and *Bifidobacteria*, also shown to be reduced in IBD

can affect host cellular DNA methylation, by regulating the methyl donor availability. The

increased methylation levels detected in THC/SIV RMs may be driven by the increased relative

abundance of *Lactobacillus* (n=28) and *Bifidobacterium* (n=18) species we recently

demonstrated [8], both of which are commensal microbes that synthesize folate for methyl group

donation [71]. As emphasized by Xu *et al.* [61] and Allen *et al.* [72], the interaction between gut

microbes and host epigenetics should not only be studied further but also therapeutically

targeted for HIV, IBD, and colorectal cancer therapy. Based on our findings, phytocannabinoids

may offer a safer and more effective therapeutic option to modulate microbe-host epigenetic

interactions in people with IBD and HIV.

Increased *NLRP6* expression in VEH/SIV RMs could be activated from the basolateral

side by dsRNA intermediates formed during the HIV replication cycle along with LTA from

dysbiotic gut bacteria. Such activation is strongly supported by the significantly high mRNA

expression of *DHX15*, an RNA helicase that cooperates with *NLRP6* to recognize dsRNA and

subsequently activate ISG expression through the mitochondrial antiviral signaling protein to

exert antiviral effects. Promoter DNA hypermethylation coupled with significantly high *CYLD*

mRNA expression in the CE of THC/SIV RMs suggests that phytocannabinoids can act

upstream at the level of the DNA to reduce *NLRP6* gene/protein expression and downstream by

upregulating *CYLD* to deubiquitylate and reduce NLRP6 protein activity. Further evidence

supporting THC's ability to inhibit NLRP6 activation comes from our recently published studies

[39,44] describing significantly reduced mRNA expression of several ISGs and defensins,

antimicrobial peptides produced by Paneth cells under the regulatory control of *NLRP6*.

Although NLRP6 has been described to activate both pyroptosis and necroptosis [50], the

significantly high mRNA expression of *RIPK1, RIPK3, MLKL, PPM1B, TRPM7, ADAM17,* and

HMGB1 suggested that necroptosis is likely the dominant cell death pathway potentially

activated by NLRP6 in the CE. Further, significantly high protein expression of p-RIPK3(Ser199)

and p-MLKL(Thr357/Ser358) together with increased protein expression of HMGB1, provided

strong evidence for the activation of necroptosis. More strikingly, excessive activation of

proteins associated with the necroptosis pathway was detected in the intestine even when SIV

replication was effectively suppressed by cART. As the terminal and crucial mediator of

necroptosis, the significantly high expression of p-MLKL(Thr357/Ser358) to our knowledge, for

the first time, identifies necroptosis as an important mechanism that could potentially drive

intestinal epithelial barrier disruption in PLWH on suppressive cART. Furthermore, the presence

of high HMGB1 protein expression means that necroptotic cell death could release HMGB1 into

the extracellular space thereby activating and further amplifying secondary inflammatory

responses leading to more widespread epithelial damage. Most importantly, the reduced mRNA

and protein expression of necroptosis-associated proteins in the intestinal epithelium of cART-

naïve and experienced THC/SIV RMs may be a direct consequence of downregulating NLRP6

(potentially other inflammasome sensors too) protein expression via promoter hypermethylation

(upstream) and its functional activity by promoting its deubiquitylation through upregulating

CYLD (downstream) (Figure 7). Given that the MB21D1 or cGAS-STING pathway also activates

necroptosis[56], THC-induced hypermethylation of MB21D1 provides an additional layer of

protection against necroptosis.

Despite these clinically relevant findings, our study does have limitations. First,

statistically significant NLRP6 upregulation was not detected in the colon of cART-naïve RMs.

This may be attributed to the variation associated with between-group cross-sectional sampling

performed on multiple animals at a single time point. However, in the cART-treated cohort, the

collection of surgical intestinal resections from the same animals before and after SIV infection

allowed the longitudinal detection of increased NLRP6 protein expression during acute infection

that continued to remain high into the chronic stage despite viral suppression by cART. Our data

also confirmed stronger NLRP6 expression in the jejunum than the colon and therefore should

receive more scrutiny in the future. From a functional perspective, persistent NLRP6 and

necroptosis activation in the jejunum could be more detrimental as it could interfere with nutrient

and cART drug absorption, leading to incomplete viral suppression, increased systemic

inflammation, and malnutrition. Moreover, the use of bulk CE made it impossible to tease apart

the specific enterocyte populations that were most impacted by the methylation changes. Our

future studies will utilize single-cell multiomics to investigate cell type-specific differences in

gene expression and chromatin accessibility changes within the same cell population.

In conclusion, our findings provide an unprecedented insight into the epigenetic

mechanisms regulating aberrant CE gene expression in chronic HIV/SIV infection, particularly in

the paCGI of CE genes during chronic HIV/SIV infection, that impact inflammatory responses,

cellular adhesions, and CE proliferation, including gene expression changes in the machinery

responsible for these methylation changes. Due to the heritable nature of epigenetic marks, we

postulate that these changes might partly explain a potential mechanism that drives chronic

intestinal inflammation and barrier dysfunction in PLWH. Given the absence of FDA-approved

inhibitors of NLRP6, we propose phytocannabinoids such as low-dose THC as a safe, feasible,

inexpensive, and effective strategy to modulate inflammasome formation via epigenetic

alteration of inflammasome sensors (i.e., *NLRP6*). Accordingly, modulation of inflammasome

activation and downstream inhibition of necroptosis in the intestine as a therapeutic modality will

benefit not only PLWH but also those suffering from other chronic inflammatory diseases like

IBD [73,74]. The important finding that CBD also blocked NLRP6 protein upregulation in vitro

indicates its potential use as an alternative to THC and presents the opportunity to be combined

with THC in appropriate ratios to reduce the psychotropic effects of THC while blocking NLRP6

hyperactivation. It is imperative that the hypermethylation of *NLRP6* and *MB21D1* not be

deemed too detrimental to intestinal homeostasis, as such a therapeutic approach may be

needed to suppress persistent intestinal inflammation in people with HIV and IBD. Nevertheless,

given the reversible nature of epigenetic changes, discontinuation of phytocannabinoids alone

after inflammation resolution should lead to restoration of normal methylation levels.

Materials and Methods

Animal care, ethics, and experimental procedures

All experiments using rhesus macaques were approved by the Tulane and LSUHSC Institutional Animal Care and Use Committee (Protocols 3574, 3581, and 3781). The Tulane National Primate Research Center (TNPRC) is an association for Assessment and Accreditation of Laboratory Animal Care International-accredited facility (AAALAC #000594). The NIH Office of Laboratory Animal Welfare assurance number for the TNPRC is A3071-01. All clinical procedures, including administration of anesthesia and analgesics, were carried out under the direction of a laboratory animal veterinarian. Animals were pre-anesthetized with ketamine hydrochloride, acepromazine, and glycopyrrolate, intubated and maintained on a mixture of isoflurane and oxygen. All possible measures were taken to minimize the discomfort of all the animals used in this study. Tulane University complies with NIH policy on animal welfare, the Animal Welfare Act, and all other applicable federal, state and local laws.

Animal model and experimental design

Fifty-eight age and weight-matched male Indian RMs were randomly distributed into six groups. Group 1 [uninfected controls (n = 16)] served as uninfected controls. Groups 2-5 (n = 42) were infected intravenously with 100 times the 50% tissue culture infective dose ($100TCID_{50}$) of SIVmac251. Groups 2 [VEH/SIV] (n=14) and 4 [VEH/SIV/ART] (n=8) received twice daily

injections of vehicle (VEH) (1:1:18 of emulphor:ethanol:saline), intramuscularly. Groups 3

[THC/SIV] (n=12) and 5 [THC/SIV/ART] (n=8) received twice daily injection of THC

intramuscularly four (THC/SIV) or two weeks (THC/SIV/ART) before SIV infection at 0.18mg/kg,

as used in previous studies [8]. This dose of THC was increased to 0.32mg/kg over a period of 2

weeks and maintained for the duration of the study. For animals in groups 4 and 5 (n = 16),

cART was given daily by subcutaneous injection and included PMPA (20mg/kg) (Gilead),

Emtricitabine (30mg/kg) (Gilead) and Dolutegravir (3mg/kg) (ViiV Healthcare) and was initiated

two weeks post-infection. As mentioned previously[44], four macaques in group 4 and four

macaques in group 5 received two injections of anti-α4β7 integrin (LN60, LC48, LM85, LJ21) or

control IgG (LM56, LH75, LH92, LI81) (50 mg/kg of anti- α4β7 or control IgG) beginning 4

months post infection (MPI) at three-week intervals as part of another study before the jejunum

resections were collected at 5 MPI.

Colon tissue (~5cm) was collected for epithelial cell isolation at necropsy from all

animals except animals from groups 4 and 5. For these animals, jejunum resections (5~cm)

were collected at pre-infection, 1 MPI and 5 MPI. For histopathological and

immunohistochemical evaluation, colon and jejunum tissues were fixed in Z-fix, embedded in

paraffin, sectioned at 5 μM for further analysis.

SIV levels in plasma, colon, and jejunum were quantified using the TaqMan One-Step

Real-time RT-qPCR assay that targeted the LTR gene [75]. SIV inoculum, infection duration,

plasma/colon/jejunum viral loads, and colon/jejunum histopathology can be found in Table 1.

Colonic epithelial cell isolation and DNA/RNA extraction

Colon epithelial cells were isolated [38] for Arraystar microarray [44] and RRBS. The purified

epithelial components were then collected in RNAlater (Thermo Fisher Scientific) for DNA

extraction (DNeasy Blood and Tissue Kit; Qiagen Inc, CA), or lyzed in Qiazol (Qiagen Inc, CA)

for total RNA extraction.

Reduced representation bisulfite sequencing (RRBS) library construction

Methyl-MiniSeq Library preparation was performed by Zymo Research. Libraries were prepared

from 200-500ng of genomic DNA digested with 60 units of Taqαl and 30 units of MspI (NEB)

sequentially and then extracted with DNA Clean & Concentrator^{TM}-5 (Zymo Research).

Fragments were ligated to pre-annealed adapters containing 5'-methyl-cytosine according to

Illumina's specified guidelines (www.illumina.com). Adapter-ligated fragments of 150-250bp and

250-300bp were recovered from a 2/5% NuSieve 1:1 agarose gel (Zymoclean Gel DNA

Recovery Kit, Zymo Research). The fragments were then bisulfite treated using EX DNA

Methylation-Lightning Kit (Zymo Research). Preparative-scale PCR was performed, and the

resulting products purified for sequencing on an Illumina HiSeq.

Sequence reads from bisulfite-treated EpiQuest libraries were identified using standard

Illumina based-calling software and then analyzed using Zymo Research propietrary analysis

pipeline,   which   is   written   in   Python   along   with   Bismark

(www.bioinfomatics.babraham.ac.uk/projects/bismark/) to perform the alignment. Index files

were constructed using the *bismark_genome_preparation*) command and the entire reference

genome. The *non-directional* parameter was applied while running Bismark. All other

parameters were set to default. Filled-in nucleotides were trimmed off while doing methylation

calling. The methylation level of each sampled cystosine (C) was estimated as the number of

reads reporting a C, divided by the total number of reads reporting a C or thymine. Fisher's

exact test or *t* test was performed for each CpG site which has at least five reads coverage, and

promoter, gene body, and CpG island annotations were added for each CpG included in the

comparison.

Global mRNA profiling

750 Sample preparation, microarray analysis, and statistics were performed by Arraystar, as

751 previously described [44].

GO enrichment analysis

755 To determine the biological functions of differentially methylated genes, Gene Ontology (GO)

756 enrichment analysis was carried out as previously decreased (52), but *Macaca mulatta* was the

757 selected organism.

Quantitative image analysis of colon and jejunum sections

Colon and jejunum sections from animals in Table 1 were stained with antibodies specific for

NLRP6 (Millipore Sigma: ABF29; colon 1:100 dilution), p-RIPK3(Ser199) (MyBiosource:

MBS1569089; 1:200 dilution), PPM1B (Proteintech: 67647-1-Ig; 1:500 dilution), p-

MLKL(Thr357) (Novus Biologicals: MAB91871, 1:200 dilution), p-MLKL(Ser358) (GeneTex;

GTX00973, 1:100 dilution), HMGB1 (D3E5) (Cell Signaling Technology: 6893; 1:50 dilution). At

least eight areas per tissues for each RM was scanned using a Zeiss LSM800 confocal

microscope (Carl ZEISS Microscopy, LLC) at 20X objective, and imported as digital images into

HALO software (Indica Labs) for image quantitation analysis. Since cells in the lamina propria

also stained positive for all protein markers, regions of interest (ROIs) were manually drawn

around the colon or jejunum epithelium using the HALO pen tool. The Indica Labs' Area

quantification module (FL v4.2.3) was then utilized to determine the average intensity of NLRP6

in each epithelial cell, in the ROIs. The output values (average positive area fluorescence) were

used to calculate the total fluorescence within the ROIs for colonic and jejunal sections.

Data availability

RRBS data will be submitted to Gene Expression Omnibus (GEO), and accession numbers will

be provided if we are approved for resubmission. mRNA profiling data was previously submitted

to GEO (accession no: GSE223482;

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE223482) [44].

Data analysis

Distribution of differentially methylated site in the paCGI region of protein-coding genes and the

average methylation change that occurred within NLRP6 paCGI, mRNA gene expression

changes, and NLRP6 and necroptosis-associated protein image quantitation in the colonic and

jejunum epithelium and lamina propria mononuclear cells were graphed and analyzed using

Prism v9 software (GraphPad).

After verifying data assumptions (normal distribution) using Anderson-Darling,

D'Agostino & Pearson, Shapiro-Wilk, and Kolmogorov-Smirnov tests, p-values were calculated

using either Kruskal-Wallis (differential methylation change in paCGIs and NLRP6 fluorescence

in colonic LPL) or Ordinary one-way ANOVA (Area fluorescence of NLRP6, p-RIPK3(Ser199),

PPM1B, p-MLKL(Thr357), p-MLKL(Ser358) and HMGB1 in colon and jejunum epithelium),

unpaired t test (VEH/SIV relative to control and THC/SIV RMs methylation ratio of DMSs within

NLRP6 and MB21D1 paCGI), Mann-Whitney U test (THC/SIV relative to control RMs

methylation ratio of DMSs within NLRP6 paCGI), or Mixed-effects analysis (NLRP6 area

fluorescence in jejunum epithelial cells. Heatmaps were made with TB Tools [76].

798
799
800
801   **Acknowledgements**
802
803

804   The authors would like to thank Ronald S. Veazey, Maurice Duplantis, Faith R. Schiro, Cecily C.

805   Midkiff, Coty Tatum (Tulane National Primate Research Center, Covington, Louisiana) and

806   Eunhee Lee (Texas Biomedical Research Institute, San Antonio, Texas) for their technical

807   assistance with nonhuman primate studies.

809
810   **Author Contributions**
811
812
813

LP and MMW equally contributed to the overall gene expression and DNA methylation data

processing, analysis, immunofluorescence studies, image analysis, cell culture and writing of

the manuscript. MM carried out the overall planning, direction, and design of the in vivo and in

vitro experiments. LP, MMW, LR, BG, JAD and MM carried out the day-day sampling

scheduling (animal experiments) and prepared the samples for RRBS and microarray gene

expression, and performed immunofluorescence/image quantification, and data analysis. BL

and CMO assisted with viral load assays, data analysis, and overall data interpretation and

conclusions. LP, MMW and MM wrote the manuscript with input from all authors. JAD, BL, and

CMO provided helpful suggestions and review of the manuscript. All authors read and approved

the final version of the manuscript.

Funding

Research reported in this publication was supported by the National Institutes of Health Award Numbers R01DA042524 and R01DA052845 to MM, R01DA050169 and R33DA053643 to CMO and MM, P30AI161943, P51OD011104 and P51OD111033. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. The funding agency (NIH) had played no role in the study design, data collection, data analyses, interpretation, or writing of the manuscript.

Figure 1. Hypomethylation changes in promoter associated CpG islands (paCGIs) associated with upregulation of their respective gene and methylation altering machinery gene expression. (**A**) Distribution of differentially methylated sites (DMSs) (p-value <0.05, absolute Δβ-value >0.1) in individual chromosomes of the rhesus macaque (RM) genome. After excluding intergenic DMSs, the bar graphs show the remaining 14.67% (**B**), 13.59% (**C**) and 10.61% (**D**) DMSs in annotated genes of VEH/SIV (**B**; red) and THC/SIV (**C**; blue) compared to control or VEH/SIV compared to THC/SIV (**D**; gold) RMs, respectively. Percentage of DMSs present in the promoter, exon and intronic regions contained within CpG islands (CGI) of annotated genes in VEH/SIV (**E**) and THC/SIV (**F**) compared to control or VEH/SIV (**G**)

compared to THC/SIV RMs. Number of genes that showed hypo or hypermethylation in their

paCGIs compared to their respective exons and introns in VEH/SIV (red) and THC/SIV (blue)

relative to control or VEH/SIV compared to THC/SIV RMs (gold) (**H**). Δβ-value of DMSs within

the paCGIs between VEH/SIV and THC/SIV compared to controls, and THC/SIV relative to

control vs VEH/SIV compared to THC/SIV RMs. Venn diagram showing relationship between

Δβ-values of DMSs in paCGIs and gene expression in VEH/SIV (**J**) and THC/SIV (**K**) relative to

control or VEH/SIV compared to THC/SIV RMs (**L**). Normalized signal intensity and fold change

of *TET2* (**M**), *TET3* (**N**), *DNMT3A* (**O**) and *DNMT3B* (**P**) mRNA in colonic epithelium of VEH/SIV

and THC/SIV compared to control RMs. Differences in methylation changes between groups

were analyzed using Kruskal-Wallis test (GraphPad Prism). A p-value of <0.05 was considered

significant. Data represented as mean $\pm$ SD.

Figure 2. Genes with methylation change throughout CGIs associated with inflammatory

response, cellular adhesion, and cell proliferation. Biological processes identified using

Gene ontology enrichment analysis of hypomethylated genes in VEH/SIV (red) and THC/SIV

(blue) compared to control or in VEH/SIV relative to THC/SIV RMs (gold) (**A-C**) or

hypermethylated genes in THC/SIV compared to control RMs (green) (**B&D**). Number of

differentially methylated CpGs in the CGI of *NLRP6* (**E**), *LEP* (**F**), *PCDH17* (**G**), *CDH7* (**H**),

SFRP1 (**I**), *TERT* (**J**), and the Δβ-values in promoter associated CGIs of *SFRP1*, *NLRP6*, *LEP*,

and *CDH7* (**K**). DMSs p-value <0.05, absolute Δβ-value >0.1.

Figure 3. Long-term low-dose THC administration significantly increased methylation in

the *NLRP6* promoter associated CGI (paCGI). UCSC Genome track for *NLRP6* showing the

location of the paCGI (black box), and DMSs in VEH/SIV (red) and THC/SIV (blue) relative to

control, and VEH/SIV compared to THC/SIV RMs (green) (**A**). Heat maps showing the DMSs in

NLRP6 CGI in VEH/SIV (**B**), and THC/SIV (**C**) compared to control or in VEH/SIV relative to

THC/SIV RMs (**C**). Cumulative β-values of NLRP6 paCGI in VEH/SIV (**E**) and THC/SIV (**F**) relative to control and in VEH/SIV compared to THC/SIV RMs (**G**). Immunofluorescence localization of NLRP6 protein expression in the colon of control (**H**), ART-naïve VEH/SIV (**I**), and THC/SIV RMs (**J**), and jejunum of cART-experienced RMs before SIV infection (**M&P**), at 1 (**N&Q**)- and 5 (**O&R**)-months post infection (MPI) administered Vehicle or THC. NLRP6 protein quantification in colon and jejunum epithelial (**K&S**) and lamina propria mononuclear cell (**L&T**) compartments.

Figure 4. Cannabinoids successfully blocked polyI:C and LTA-induced NLRP6 upregulation in vitro and reduced necroptosis-driven gene signature in colonic epithelium (CE) of SIV-infected rhesus macaques (RMs). Representative immunofluorescence images showing the expression of NLRP6 (green) protein in human small intestinal epithelial (HSIE) cells at 18 h following treatment with DMSO (**A**) or post transfection with 2 μg/mL of polyI:C alone (**B**) or also treated with lipoteichoic acid (LTA) (10 μg/mL) 4 h later (**C**). PolyI:C and LTA treated HSIE cells were preincubated for 1 h with THC (**D**), AM251 + THC (**E**), AM630 + THC (**F**), CBD (**G**) or THC and CBD (**H**), fixed after 18 h, stained and NLRP6 (green) protein expression quantitated (**I**). Experiments were performed in triplicate wells using 3 μM of THC and CBD, 10 μM of AM251/AM630, and repeated thrice. Microarray normalized signal intensity of *CYLD* (**J**), *DHX15* (**K**), *CASP1* (**L**), *IL-18* (**M**), *IL-1β* (**N**), *GSDMD* (**O**), *RIPK1* (**P**), *RIPK3* (**Q**), *MLKL* (**R**), *PPM1B* (**S**), *ADAM17* (**T**), *TRPM7* (**U**), *HMGB1* (**V**), *TBK1* (**Y**) and *PQBP1* (**Z**) in CE of SIV-infected RMs. Heat maps showing the DMSs in MB21D1 CGI in VEH/SIV (**W1**), and THC/SIV (**W2**) compared to control or in VEH/SIV relative to THC/SIV RMs (**W3**). Cumulative β-values of MB21D1 paCGI in VEH/SIV (**X1**) and THC/SIV (**X2**) relative to control and in VEH/SIV compared to THC/SIV RMs (**X3**). White arrows in panels A to G indicate epithelial regions. Immunofluorescence data were analyzed using one-way ANOVA followed by Tukey's multiple comparison post hoc test. A p-value of <0.05 was considered significant.

Figure 5. Chronic THC administration reduced HIV/SIV infection induced RIPK3(Ser199), PPM1B, p-MLKL(Thr357) protein expression in the intestine. p-RIPK3(Ser199) (red) (**A-C**), PPM1B (**E-G**) (green) and p-MLKL(Thr357) (green) (**I-K**) protein expression in colon tissues of uninfected control (**A, E&I**), VEH/SIV (**B, F&J**), and THC/SIV rhesus macaques (RMs) (**C, G&K**). p-MLKL(Thr357) (green) protein expression in jejunum tissues before SIV infection (**M**) and at 5 months post SIV infection in cART suppressed VEH/SIV (**N**) and THC/SIV (**O**) RMs. Representative immunofluorescence images were captured using a Zeiss confocal microscope at 20X magnification. Quantitation of p-RIPK3(Ser199) (**D**), PPM1B (**H**) and p-MLKL(Thr357) (**L&P**) average positive area fluorescence in colon and jejunum epithelium was performed using Halo software. Note the significantly reduced p-MLKL(Thr357) staining in the colon and jejunum epithelium of THC/SIV (**K, O, L&P**) compared to VEH/SIV (**J, N, L &P**) RMs. White arrows in panels **I** to **O** indicate epithelial regions. Immunofluorescence data were analyzed using one-way ANOVA followed by Tukey's multiple comparison post hoc test. A p-value of <0.05 was considered significant.

Figure 6. Chronic THC administered in combination with cART significantly reduced HIV/SIV infection induced p-MLKL(Ser358) and HMGB1 protein expression in the intestine. p-MLKL(Ser358) (green) (**A-C**), and HMGB1 (green) (**I-K**) protein expression in colon tissues of uninfected control (**A&I**), VEH/SIV (**B&J**), and THC/SIV (**C&K**) rhesus macaques (RMs). p- MLKL(Ser358) (green) (**E-G**) and HMGB1 (green) (**M-O**) expression in jejunum tissues before SIV infection (**E&M)**) and at 5 months post SIV infection in cART suppressed VEH/SIV (**F&N**) and THC/SIV (**G&O**) RMs. Representative immunofluorescence images were captured using a Zeiss confocal microscope at 20X magnification. Quantitation of MLKL(Ser358) (**D&H**), and HMGB1 (**L&P**) average positive area fluorescence in colon and jejunum epithelium was performed using Halo software. Note the significantly reduced p-

MLKL(Ser358) and HMGB1 staining in the jejunum epithelium of cART suppressed THC/SIV

(**G, O, H&P**) compared to VEH/SIV (**F, N, H&P**) RMs. White arrows in panels A to G indicate

epithelial regions. Immunofluorescence data were analyzed using one-way ANOVA followed by

Tukey's multiple comparison post hoc test. A p-value of <0.05 was considered significant.

Figure 7. HIV/SIV infection-induced DNA hypomethylation facilitates persistent activation of the

NLRP6 sensor and cGAS/MB21D1 protein expression in the intestinal epithelium, leading to

epithelial barrier breakdown and microbial translocation (**A**). Long-term low-dose THC

epigenetically modulates these changes by hypermethylating CpG sites in the NLRP6 and

cGAS promoters, thereby reducing gastrointestinal mucosal injury (**B**). NLRP6 inflammasome

assembly could potentially activate the RIPK1-RIPK3-MLKL pathway to induce necroptosis of

intestinal epithelium (**C**). TLR3 (Figure S3J) and NLRP6-DHX15 viral sensing complexes could

recognize dsRNA intermediates of HIV, leading to the phosphorylation of RIPK1, which in turn

activates RIPK3, culminating in the activation of the necroptosis executioner protein MLKL

through phosphorylation on Thr357 and Ser358. Activation of MLKL and concurrent activation of

the calcium ion channel TRPM7 enhances intracellular Ca++ influx, leading to necroptotic cell

death. Cell death facilitates the release and extracellular accumulation of the DAMP protein,

HMGB1, which can propagate secondary inflammatory responses by attracting neutrophils and

macrophages leading to more widespread epithelial damage (**D**). By hypermethylating the

NLRP6 promoter and increasing expression of the deubiquitinase CYLD, long-term low-dose

THC can block the downstream activation of the RIPK1-RIPK3-MLKL pathway and significantly

reduce HMGB1 protein expression to attenuate gastrointestinal inflammation and preserve

epithelial barrier integrity. Figures were generated using Biorender.

Supplemental Table S1. Number of CpG sites hypomethylated (in red) or hypermethylated (in

black) within CpG islands of unique genes in VEH/SIV and THC/SIV, compared to controls and

THC/SIV rhesus macaques.

Supplemental Figure S1. Flow chart showing stepwise processing of RRBS DNA methylation

data and identification of differentially methylated sites in paCGIs.

Supplemental Figure S2. Clustering heatmaps of the top 100 differentially methylated CpG

sites in VEH/SIV (**A**) and THC/SIV (**B**) relative to control rhesus macaques (RMs) and in

VEH/SIV (**C**) compared to THC/SIV RMs. Pearson's correlation coefficients for all CpG sites

between VEH/SIV vs control (**D**), THC/SIV vs control (**E**), and VEH/SIV vs THC/SIV (**F**) RMs

were 0.9400, 0.9358, and 0.9359, respectively.

Supplemental Figure S3. UCSC Genome track for *MB21D1* showing the location of the 108 bp

CGI (**A**). p-RIPK3(Ser199) (green) (**B-D**), and PPM1B (green) (**F-H**) protein expression in

jejunum tissues before SIV infection (**B&F)**) and at 5 months post SIV infection in cART

suppressed VEH/SIV (**C&G**) and THC/SIV (**D&H**) rhesus macaques. RIPK3(Ser199) (**E**) and

PPM1B (**I**) protein quantification in jejunum. Immunofluorescence data were analyzed using

one-way ANOVA followed by Tukey's multiple comparison post hoc test. A p-value of <0.05 was

considered significant. *TLR3* (**J**) gene expression in significantly upregulated in the colonic

epithelium of VEH/SIV but not in THC/SIV compared to control rhesus macaques.

References

1. Sim, J.H., Mukerji, S.S., Russo, S.C., and Lo, J. (2021). Gastrointestinal Dysfunction and HIV Comorbidities. Curr HIV/AIDS Rep *18*, 57-62. 10.1007/s11904-020-00537-8.

2. Somsouk, M., Estes, J.D., Deleage, C., Dunham, R.M., Albright, R., Inadomi, J.M., Martin, J.N., Deeks, S.G., McCune, J.M., and Hunt, P.W. (2015). Gut epithelial barrier and systemic inflammation during chronic HIV infection. Aids *29*, 43-51. 10.1097/qad.0000000000000511.

3. Hall, V.P. (2018). Common Gastrointestinal Complications Associated with Human Immunodeficiency Virus/AIDS: An Overview. Crit Care Nurs Clin North Am *30*, 101-107. 10.1016/j.cnc.2017.10.009.

4. Wang, H., and Kotler, D.P. (2014). HIV enteropathy and aging: gastrointestinal immunity, mucosal epithelial barrier, and microbial translocation. Curr Opin HIV AIDS *9*, 309-316. 10.1097/coh.0000000000000066.

5. Tincati, C., Merlini, E., Braidotti, P., Ancona, G., Savi, F., Tosi, D., Borghi, E., Callegari, M.L., Mangiavillano, B., Barassi, A., et al. (2016). Impaired gut junctional complexes feature late-treated individuals with suboptimal CD4+ T-cell recovery upon virologically suppressive combination antiretroviral therapy. Aids *30*, 991-1003. 10.1097/qad.0000000000001015.

6. Mehta, F. (2016). Report: economic implications of inflammatory bowel disease and its management. Am J Manag Care *22*, s51-60.

7. Muehler, A., Slizgi, J.R., Kohlhof, H., Groeppel, M., Peelen, E., and Vitt, D. (2020). Clinical relevance of intestinal barrier dysfunction in common gastrointestinal diseases. World J Gastrointest Pathophysiol *11*, 114-130. 10.4291/wjgp.v11.i6.114.

8.   McDew-White, M., Lee, E., Premadasa, L.S., Alvarez, X., Okeoma, C.M., and Mohan, M.

(2023). Cannabinoids modulate the microbiota-gut-brain axis in HIV/SIV infection by

reducing neuroinflammation and dysbiosis while concurrently elevating endocannabinoid

and indole-3-propionate levels. J Neuroinflammation *20*, 62. 10.1186/s12974-023-

02729-6.

9.   Legaki, E., and Gazouli, M. (2016). Influence of environmental factors in the

development of inflammatory bowel diseases. World J Gastrointest Pharmacol Ther *7*,

112-125. 10.4292/wjgpt.v7.i1.112.

10.   McDermott, E., Ryan, E.J., Tosetto, M., Gibson, D., Burrage, J., Keegan, D., Byrne, K.,

Crowe, E., Sexton, G., Malone, K., et al. (2016). DNA Methylation Profiling in

Inflammatory Bowel Disease Provides New Insights into Disease Pathogenesis. J

Crohns Colitis *10*, 77-86. 10.1093/ecco-jcc/jjv176.

11.   Kraiczy, J., Nayak, K., Ross, A., Raine, T., Mak, T.N., Gasparetto, M., Cario, E., Rakyan,

V., Heuschkel, R., and Zilbauer, M. (2016). Assessing DNA methylation in the

developing human intestinal epithelium: potential link to inflammatory bowel disease.

Mucosal Immunol *9*, 647-658. 10.1038/mi.2015.88.

12.   Gerecke, C., Scholtka, B., Löwenstein, Y., Fait, I., Gottschalk, U., Rogoll, D., Melcher,

R., and Kleuser, B. (2015). Hypermethylation of ITGA4, TFPI2 and VIMENTIN promoters

is increased in inflamed colon tissue: putative risk markers for colitis-associated cancer.

J Cancer Res Clin Oncol *141*, 2097-2107. 10.1007/s00432-015-1972-8.

13.   Marjoram, L., Alvers, A., Deerhake, M.E., Bagwell, J., Mankiewicz, J., Cocchiaro, J.L.,

Beerman, R.W., Willer, J., Sumigray, K.D., Katsanis, N., et al. (2015). Epigenetic control

of intestinal barrier function and inflammation in zebrafish. Proc Natl Acad Sci U S A

112, 2770-2775. 10.1073/pnas.1424089112.

14.   Gonsky, R., Deem, R.L., Landers, C.J., Haritunians, T., Yang, S., and Targan, S.R.

(2014). IFNG rs1861494 polymorphism is associated with IBD disease severity and

functional changes in both IFNG methylation and protein secretion. Inflamm Bowel Dis

20, 1794-1801. 10.1097/mib.0000000000000172.

15. Harris, R.A., Nagy-Szakal, D., Mir, S.A., Frank, E., Szigeti, R., Kaplan, J.L., Bronsky, J.,

Opekun, A., Ferry, G.D., Winter, H., and Kellermayer, R. (2014). DNA methylation-

associated colonic mucosal immune and defense responses in treatment-naïve pediatric

ulcerative colitis. Epigenetics *9*, 1131-1137. 10.4161/epi.29446.

16. Denizot, J., Desrichard, A., Agus, A., Uhrhammer, N., Dreux, N., Vouret-Craviari, V.,

Hofman, P., Darfeuille-Michaud, A., and Barnich, N. (2015). Diet-induced hypoxia

responsive element demethylation increases CEACAM6 expression, favouring Crohn's

disease-associated Escherichia coli colonisation. Gut *64*, 428-437. 10.1136/gutjnl-2014-

306944.

17. Karatzas, P.S., Gazouli, M., Safioleas, M., and Mantzaris, G.J. (2014). DNA methylation

changes in inflammatory bowel disease. Ann Gastroenterol *27*, 125-132.

18. Papadia, C., Louwagie, J., Del Rio, P., Grooteclaes, M., Coruzzi, A., Montana, C.,

Novelli, M., Bordi, C., de' Angelis, G.L., Bassett, P., et al. (2014). FOXE1 and SYNE1

genes hypermethylation panel as promising biomarker in colitis-associated colorectal

neoplasia. Inflamm Bowel Dis *20*, 271-277. 10.1097/01.MIB.0000435443.07237.ed.

19. Low, D., Mizoguchi, A., and Mizoguchi, E. (2013). DNA methylation in inflammatory

bowel disease and beyond. World J Gastroenterol *19*, 5238-5249.

10.3748/wjg.v19.i32.5238.

20. Koukos, G., Polytarchou, C., Kaplan, J.L., Morley-Fletcher, A., Gras-Miralles, B.,

Kokkotou, E., Baril-Dore, M., Pothoulakis, C., Winter, H.S., and Iliopoulos, D. (2013).

MicroRNA-124 regulates STAT3 expression and is down-regulated in colon tissues of

pediatric patients with ulcerative colitis. Gastroenterology *145*, 842-852.e842.

10.1053/j.gastro.2013.07.001.

21.   Ventham, N.T., Kennedy, N.A., Nimmo, E.R., and Satsangi, J. (2013). Beyond gene
1052   discovery in inflammatory bowel disease: the emerging role of epigenetics.
1053   Gastroenterology *145*, 293-308. 10.1053/j.gastro.2013.05.050.

1054   22.   Stylianou, E. (2013). Epigenetics: the fine-tuner in inflammatory bowel disease? Curr
1055   Opin Gastroenterol *29*, 370-377. 10.1097/MOG.0b013e328360bd12.

1056   23.   Cooke, J., Zhang, H., Greger, L., Silva, A.L., Massey, D., Dawson, C., Metz, A., Ibrahim,
1057   A., and Parkes, M. (2012). Mucosal genome-wide methylation changes in inflammatory
1058   bowel disease. Inflamm Bowel Dis *18*, 2128-2137. 10.1002/ibd.22942.

1059   24.   Vaiopoulou, A., Karamanolis, G., Psaltopoulou, T., Karatzias, G., and Gazouli, M.
1060   (2014). Molecular basis of the irritable bowel syndrome. World J Gastroenterol *20*, 376-
1061   383. 10.3748/wjg.v20.i2.376.

1062   25.   Dinan, T.G., Cryan, J., Shanahan, F., Keeling, P.W., and Quigley, E.M. (2010). IBS: An
1063   epigenetic   perspective.   Nat   Rev   Gastroenterol   Hepatol   *7*,   465-471.
1064   10.1038/nrgastro.2010.99.

1065   26.   Zilbauer, M., and Kraiczy, J. (2017). Epigenetics in Gastrointestinal Health and Disease:
1066   Spotlight on DNA Methylation in the Intestinal Epithelium. Nestle Nutr Inst Workshop Ser
1067   *88*, 35-44. 10.1159/000455211.

1068   27.   Howell, K.J., Kraiczy, J., Nayak, K.M., Gasparetto, M., Ross, A., Lee, C., Mak, T.N., Koo,
1069   B.K., Kumar, N., Lawley, T., et al. (2018). DNA Methylation and Transcription Patterns in
1070   Intestinal Epithelial Cells From Pediatric Patients With Inflammatory Bowel Diseases
1071   Differentiate Disease Subtypes and Associate With Outcome. Gastroenterology *154*,
1072   585-598. 10.1053/j.gastro.2017.10.007.

1073   28.   Kraiczy, J., Nayak, K.M., Howell, K.J., Ross, A., Forbester, J., Salvestrini, C., Mustata,
1074   R., Perkins, S., Andersson-Rolf, A., Leenen, E., et al. (2019). DNA methylation defines
1075   regional identity of human intestinal epithelial organoids and undergoes dynamic
1076   changes during development. Gut *68*, 49-61. 10.1136/gutjnl-2017-314817.

1077   29.   Chapman, C.G., and Pekow, J. (2015). The emerging role of miRNAs in inflammatory

1078   bowel   disease: a   review.   Therap   Adv   Gastroenterol   *8*,   4-22.

1079   10.1177/1756283x14547360.

1080   30.   Ye, D., Guo, S., Al-Sadi, R., and Ma, T.Y. (2011). MicroRNA regulation of intestinal

1081   epithelial   tight   junction   permeability.   Gastroenterology   *141*,   1323-1333.

1082   10.1053/j.gastro.2011.07.005.

1083   31.   Xu, X.M., and Zhang, H.J. (2016). miRNAs as new molecular insights into inflammatory

1084   bowel   disease: Crucial   regulators   in   autoimmunity   and   inflammation.   World   J

1085   Gastroenterol *22*, 2206-2218. 10.3748/wjg.v22.i7.2206.

1086   32.   Tartakover Matalon, S., Azar, S., Meiri, D., Hadar, R., Nemirovski, A., Abu Jabal, N.,

1087   Konikoff, F.M., Drucker, L., Tam, J., and Naftali, T. (2021). Endocannabinoid Levels in

1088   Ulcerative Colitis Patients Correlate With Clinical Parameters and Are Affected by

1089   Cannabis   Consumption.   Front   Endocrinol   (Lausanne)   *12*,   685289.

1090   10.3389/fendo.2021.685289.

1091   33.   Mohan, M., Chandra, L.C., Torben, W., Aye, P.P., Alvarez, X., and Lackner, A.A. (2014).

1092   miR-190b   is   markedly   upregulated   in   the   intestine   in   response   to   simian

1093   immunodeficiency virus replication and partly regulates myotubularin-related protein-6

1094   expression. J Immunol *193*, 1301-1313. 10.4049/jimmunol.1303479.

1095   34.   Mohan, M., Kumar, V., Lackner, A.A., and Alvarez, X. (2015). Dysregulated miR-34a-

1096   SIRT1-acetyl p65 axis is a potential mediator of immune activation in the colon during

1097   chronic simian immunodeficiency virus infection of rhesus macaques. J Immunol *194*,

1098   291-306. 10.4049/jimmunol.1401447.

1099   35.   Kumar, V., Torben, W., Kenway, C.S., Schiro, F.R., and Mohan, M. (2016). Longitudinal

1100   Examination   of   the   Intestinal   Lamina   Propria   Cellular   Compartment   of   Simian

1101   Immunodeficiency Virus-Infected Rhesus Macaques Provides Broader and Deeper

1102   Insights into the Link between Aberrant MicroRNA Expression and Persistent Immune

1103   Activation. J Virol *90*, 5003-5019. 10.1128/JVI.00189-16.

1104 36. Kumar, V., Mansfield, J., Fan, R., MacLean, A., Li, J., and Mohan, M. (2018). miR-130a

1105   and miR-212 Disrupt the Intestinal Epithelial Barrier through Modulation of PPARγ and

1106   Occludin Expression in Chronic Simian Immunodeficiency Virus-Infected Rhesus

1107   Macaques. J Immunol *200*, 2677-2689. 10.4049/jimmunol.1701148.

1108 37. Mohan, M., Kaushal, D., Aye, P.P., Alvarez, X., Veazey, R.S., and Lackner, A.A. (2012).

1109   Focused examination of the intestinal lamina propria yields greater molecular insight into

1110   mechanisms underlying SIV induced immune dysfunction. PLoS One *7*, e34561.

1111   10.1371/journal.pone.0034561.

1112 38. Mohan, M., Kaushal, D., Aye, P.P., Alvarez, X., Veazey, R.S., and Lackner, A.A. (2013).

1113   Focused examination of the intestinal epithelium reveals transcriptional signatures

1114   consistent with disturbances in enterocyte maturation and differentiation during the

1115   course of SIV infection. PLoS One *8*, e60122. 10.1371/journal.pone.0060122.

1116 39. Kumar, V., Torben, W., Mansfield, J., Alvarez, X., Vande Stouwe, C., Li, J., Byrareddy,

1117   S.N., Didier, P.J., Pahar, B., Molina, P.E., and Mohan, M. (2019). Cannabinoid

1118   Attenuation of Intestinal Inflammation in Chronic SIV-Infected Rhesus Macaques

1119   Involves T Cell Modulation and Differential Expression of Micro-RNAs and Pro-

1120   inflammatory Genes. Front Immunol *10*, 914. 10.3389/fimmu.2019.00914.

1121 40. Mboumba Bouassa, R.S., Needham, J., Nohynek, D., Singer, J., Lee, T., Bobeuf, F.,

1122   Samarani, S., Del Balso, L., Paisible, N., Vertzagias, C., et al. (2022). Safety and

1123   Tolerability of Oral Cannabinoids in People Living with HIV on Long-Term ART: A

1124   Randomized, Open-Label, Interventional Pilot Clinical Trial (CTNPT 028). Biomedicines

1125   *10*. 10.3390/biomedicines10123168.

1126 41. Deaton, A.M., and Bird, A. (2011). CpG islands and the regulation of transcription.

1127   Genes Dev *25*, 1010-1022. 10.1101/gad.2037511.

42. Vavouri, T., and Lehner, B. (2012). Human genes with CpG island promoters have a
distinct transcription-associated chromatin organization. Genome Biol *13*, R110.
10.1186/gb-2012-13-11-r110.

43. Moore, L.D., Le, T., and Fan, G. (2013). DNA methylation and its basic function.
Neuropsychopharmacology *38*, 23-38. 10.1038/npp.2012.112.

44. Premadasa, L.S., Lee, E., McDew-White, M., Alvarez, X., Jayakumar, S., Ling, B.,
Okeoma, C.M., Byrareddy, S.N., Kulkarni, S., and Mohan, M. (2023). Cannabinoid
enhancement of lncRNA MMP25-AS1/MMP25 interaction reduces neutrophil infiltration
and intestinal epithelial injury in HIV/SIV infection. JCI Insight *8*.
10.1172/jci.insight.167903.

45. Mukherjee, S., Kumar, R., Tsakem Lenou, E., Basrur, V., Kontoyiannis, D.L.,
Ioakeimidis, F., Mosialos, G., Theiss, A.L., Flavell, R.A., and Venuprasad, K. (2020).
Deubiquitination of NLRP6 inflammasome by Cyld critically regulates intestinal
inflammation. Nat Immunol *21*, 626-635. 10.1038/s41590-020-0681-x.

46. Elinav, E., Strowig, T., Kau, A.L., Henao-Mejia, J., Thaiss, C.A., Booth, C.J., Peaper,
D.R., Bertin, J., Eisenbarth, S.C., Gordon, J.I., and Flavell, R.A. (2011). NLRP6
inflammasome regulates colonic microbial ecology and risk for colitis. Cell *145*, 745-757.
10.1016/j.cell.2011.04.022.

47. Gremel, G., Wanders, A., Cedernaes, J., Fagerberg, L., Hallström, B., Edlund, K.,
Sjöstedt, E., Uhlén, M., and Pontén, F. (2015). The human gastrointestinal tract-specific
transcriptome and proteome as defined by RNA sequencing and antibody-based
profiling. J Gastroenterol *50*, 46-57. 10.1007/s00535-014-0958-7.

48. Li, R., and Zhu, S. (2020). NLRP6 inflammasome. Mol Aspects Med *76*, 100859.
10.1016/j.mam.2020.100859.

49.   Xing, J., Zhou, X., Fang, M., Zhang, E., Minze, L.J., and Zhang, Z. (2021). DHX15 is

1153   required to control RNA virus-induced intestinal inflammation. Cell Rep *35*, 109205.

1154   10.1016/j.celrep.2021.109205.

1155   50.   Ghimire, L., Paudel, S., Jin, L., and Jeyaseelan, S. (2020). The NLRP6 inflammasome in

1156   health and disease. Mucosal immunology *13*, 388-398. 10.1038/s41385-020-0256-z.

1157   51.   Seo, J., Nam, Y.W., Kim, S., Oh, D.B., and Song, J. (2021). Necroptosis molecular

1158   mechanisms: Recent findings regarding novel necroptosis regulators. Exp Mol Med *53*,

1159   1007-1017. 10.1038/s12276-021-00634-7.

1160   52.   Fuchslocher Chico, J., Falk-Paulsen, M., Luzius, A., Saggau, C., Ruder, B., Bolik, J.,

1161   Schmidt-Arras, D., Linkermann, A., Becker, C., Rosenstiel, P., et al. (2018). The

1162   enhanced susceptibility of ADAM-17 hypomorphic mice to DSS-induced colitis is not

1163   ameliorated by loss of RIPK3, revealing an unexpected function of ADAM-17 in

1164   necroptosis. Oncotarget *9*, 12941-12958. 10.18632/oncotarget.24410.

1165   53.   Cai, Z., Jitkaew, S., Zhao, J., Chiang, H.C., Choksi, S., Liu, J., Ward, Y., Wu, L.G., and

1166   Liu, Z.G. (2014). Plasma membrane translocation of trimerized MLKL protein is required

1167   for TNF-induced necroptosis. Nat Cell Biol *16*, 55-65. 10.1038/ncb2883.

1168   54.   Wen, S., Li, X., Ling, Y., Chen, S., Deng, Q., Yang, L., Li, Y., Shen, J., Qiu, Y., Zhan, Y.,

1169   et al. (2020). HMGB1-associated necroptosis and Kupffer cells M1 polarization underlies

1170   remote liver injury induced by intestinal ischemia/reperfusion in rats. Faseb j *34*, 4384-

1171   4402. 10.1096/fj.201900817R.

1172   55.   Ramos, A., Bizri, N., Novak, E., Mollen, K., and Khan, S. (2024). The role of cGAS in

1173   epithelial dysregulation in inflammatory bowel disease and gastrointestinal malignancies.

1174   Front Pharmacol *15*, 1409683. 10.3389/fphar.2024.1409683.

1175   56.   Decout, A., Katz, J.D., Venkatraman, S., and Ablasser, A. (2021). The cGAS-STING

1176   pathway as a therapeutic target in inflammatory diseases. Nat Rev Immunol *21*, 548-

1177   569. 10.1038/s41577-021-00524-z.

1178   57.   Yoh, S.M., Schneider, M., Seifried, J., Soonthornvacharin, S., Akleh, R.E., Olivieri, K.C.,

1179   De Jesus, P.D., Ruan, C., de Castro, E., Ruiz, P.A., et al. (2015). PQBP1 Is a Proximal

1180   Sensor of the cGAS-Dependent Innate Response to HIV-1. Cell *161*, 1293-1305.

1181   10.1016/j.cell.2015.04.050.

1182   58.   Najafov, A., Mookhtiar, A.K., Luu, H.S., Ordureau, A., Pan, H., Amin, P.P., Li, Y., Lu, Q.,

1183   and Yuan, J. (2019). TAM Kinases Promote Necroptosis by Regulating Oligomerization

1184   of MLKL. Mol Cell *75*, 457-468.e454. 10.1016/j.molcel.2019.05.022.

1185   59.   Frank, D., and Vince, J.E. (2019). Pyroptosis versus necroptosis: similarities,

1186   differences, and crosstalk. Cell Death Differ *26*, 99-114. 10.1038/s41418-018-0212-6.

1187   60.   Okumura, R., and Takeda, K. (2017). Roles of intestinal epithelial cells in the

1188   maintenance of gut homeostasis. Exp Mol Med *49*, e338. 10.1038/emm.2017.20.

1189   61.   Xu, J., Xu, H.M., Yang, M.F., Liang, Y.J., Peng, Q.Z., Zhang, Y., Tian, C.M., Wang, L.S.,

1190   Yao, J., Nie, Y.Q., and Li, D.F. (2022). New Insights Into the Epigenetic Regulation of

1191   Inflammatory Bowel Disease. Front Pharmacol *13*, 813659. 10.3389/fphar.2022.813659.

1192   62.   Del Real, A., Pérez-Campo, F.M., Fernández, A.F., Sañudo, C., Ibarbia, C.G., Pérez-

1193   Núñez, M.I., Criekinge, W.V., Braspenning, M., Alonso, M.A., Fraga, M.F., and Riancho,

1194   J.A. (2017). Differential analysis of genome-wide methylation and gene expression in

1195   mesenchymal stem cells of patients with fractures and osteoarthritis. Epigenetics *12*,

1196   113-122. 10.1080/15592294.2016.1271854.

1197   63.   Zhang, Y., Fukui, N., Yahata, M., Katsuragawa, Y., Tashiro, T., Ikegawa, S., and Lee,

1198   M.T. (2016). Genome-wide DNA methylation profile implicates potential cartilage

1199   regeneration at the late stage of knee osteoarthritis. Osteoarthritis Cartilage *24*, 835-843.

1200   10.1016/j.joca.2015.12.013.

1201   64.   Claringbould, A., and Zaugg, J.B. (2021). Enhancers in disease: molecular basis and

1202   emerging treatment strategies. Trends Mol Med *27*, 1060-1073.

1203   10.1016/j.molmed.2021.07.012.

65.   Fu, Z., Zhao, P.Y., Yang, X.P., Li, H., Hu, S.D., Xu, Y.X., and Du, X.H. (2023).

Cannabidiol regulates apoptosis and autophagy in inflammation and cancer: A review.

Front Pharmacol *14*, 1094020. 10.3389/fphar.2023.1094020.

66.   Zheng, D., Kern, L., and Elinav, E. (2021). The NLRP6 inflammasome. Immunology *162*,

281-289. 10.1111/imm.13293.

67.   Venuprasad, K., and Theiss, A.L. (2021). NLRP6 in host defense and intestinal

inflammation. Cell Rep *35*, 109043. 10.1016/j.celrep.2021.109043.

68.   Leng, F., Yin, H., Qin, S., Zhang, K., Guan, Y., Fang, R., Wang, H., Li, G., Jiang, Z.,

Sun, F., et al. (2020). NLRP6 self-assembles into a linear molecular platform following

LPS binding and ATP stimulation. Sci Rep *10*, 198-198. 10.1038/s41598-019-57043-0.

69.   Ranson, N., Veldhuis, M., Mitchell, B., Fanning, S., Cook, A.L., Kunde, D., and Eri, R.

(2018). Nod-Like Receptor Pyrin-Containing Protein 6 (NLRP6) Is Up-regulated in Ileal

Crohn's Disease and Differentially Expressed in Goblet Cells. Cell Mol Gastroenterol

Hepatol *6*, 110-112.e118. 10.1016/j.jcmgh.2018.03.001.

70.   Huang, Y.-H., Lo, M.-H., Cai, X.-Y., and Kuo, H.-C. (2018). Epigenetic hypomethylation

and upregulation of NLRC4 and NLRP12 in Kawasaki disease. Oncotarget *9*, 18939-

18948. 10.18632/oncotarget.24851.

71.   Wu, Y., Wang, C.Z., Wan, J.Y., Yao, H., and Yuan, C.S. (2021). Dissecting the Interplay

Mechanism between Epigenetics and Gut Microbiota: Health Maintenance and Disease

Prevention. Int J Mol Sci *22*. 10.3390/ijms22136933.

72.   Allen, J., and Sears, C.L. (2019). Impact of the gut microbiome on the genome and

epigenome of colon epithelial cells: contributions to colorectal cancer development.

Genome Med *11*, 11. 10.1186/s13073-019-0621-2.

73.   Suryavanshi, S.V., Kovalchuk, I., and Kovalchuk, O. (2020). Cannabinoids as Key

Regulators of Inflammasome Signaling: A Current Perspective. Front Immunol *11*,

613613. 10.3389/fimmu.2020.613613.

74.　Liu, C., Ma, H., Slitt, A.L., and Seeram, N.P. (2020). Inhibitory Effect of Cannabidiol on

the Activation of NLRP3 Inflammasome Is Associated with Its Modulation of the P2X7

Receptor　in　Human　Monocytes.　J　Nat　Prod　*83*,　2025-2029.

10.1021/acs.jnatprod.0c00138.

75.　McDew-White, M., Lee, E., Alvarez, X., Sestak, K., Ling, B.J., Byrareddy, S.N., Okeoma,

C.M., and Mohan, M. (2022). Cannabinoid control of gingival immune activation in

chronically SIV-infected rhesus macaques involves modulation of the indoleamine-2,3-

dioxygenase-1　pathway　and　salivary　microbiome.　EBioMedicine　*75*,　103769.

10.1016/j.ebiom.2021.103769.

76.　Chen, C., Wu, Y., Li, J., Wang, X., Zeng, Z., Xu, J., Liu, Y., Feng, J., Chen, H., He, Y.,

and Xia, R. (2023). TBtools-II: A "one for all, all for one" bioinformatics platform for

biological big-data mining. Mol Plant *16*, 1733-1742. 10.1016/j.molp.2023.09.010.

1242

1243

Table 1. Animal IDs, SIV infection (SIVmac251) duration time, cannabinoids and cART treatment, viral load for plasma, colon, and jejunum, and histopathology for colon and jejunum tissue.

Animal ID	Duration of Infection	Plasma viral loads 10^6/mL	Colon viral loads 10^6/mg RNA	Jejunum viral loads 10^6/mg RNA	Colon Histopathology	Jejunum Histopathology
Group 1: Uninfected Controls (Controls)						
CONT1, CONT2, CONT3, CONT4[a]	NA	NA	NA	NA	NA	NA
CONT5[a,b]	NA	NA	NA	NA	NA	NA
CONT6[b]	NA	NA	NA	NA	NA	NA
CONT7*[b]	NA	NA	NA	NA	NA	NA
CONT8*[b]	NA	NA	NA	NA	NA	NA
CONT9, CONT10, CONT11[c]	NA	NA	NA	NA	NA	NA
CONT12, CONT13, CONT14[c]	NA	NA	NA	NA	NA	NA
CONT15, CONT16[c]	NA	NA	NA	NA	NA	NA
Group 2: Chronic SIV-infected and Vehicle Treated (VEH/SIV)						
VEHSIV1[a,c]	150	4	147	NA	ND	NA
VEHSIV2[a]	180	0.02	2	NA	ND	NA
VEHSIV3[a]	180	2	300	NA	ND	NA
VEHSIV4[b,c]	180	0.1	786	NA	Lymphoid hyperplasia	NA
VEHSIV5[a,b,c]	180	9.4	29	NA	ND	NA
VEHSIV6[a,b,c]	180	0.38	320	NA	ND	NA
VEHSIV7[a,c]	180	0.04	4	NA	ND	NA
VEHSIV8[a,c]	180	0.5	20	NA	ND	NA
VEHSIV9*[b]	90	37	3000	NA	Moderate colitis/cryptitis	NA
VEHSIV10*[b,c]	180	9	59	NA	ND	NA
VEHSIV11[c]	180	30	3	NA	Lymphoid hyperplasia	NA
VEHSIV12[c]	180	3000	200	NA	Lymphoid hyperplasia	NA
VEHSIV13[c]	180	0.2	210	NA	ND	NA
VEHSIV14[b]	180	NA	11	NA	ND	NA
Group 3: Chronic SIV-infected and THC treated (THC/SIV)						
THCSIV1[b]	180	18.9	6726	NA	Mild hyperplasia/SIV syncytia	NA
THCSIV2[a]	150	260	9,360	NA	ND	NA
THCSIV3[a]	180	3	10	NA	ND	NA

THCSIV4[a,b,c]	180	0.01	3	NA	ND	NA
THCSIV5[b,c]	180	0.06	93.8	NA	ND	NA
THCSIV6[a,c]	180	0.02	10	NA	ND	NA
THCSIV7[a,c]	180	0.02	1	NA	ND	NA
THCSIV8[a,c]	180	1	300	NA	ND	NA
THCSIV9[b]	180	18.9	6,726	NA	ND	NA
THCSIV10[b,c]	180	1.5	1,261	NA	ND	NA
THCSIV11[b,c]	180	7.7	970	NA	ND	NA
THCSIV12[c]	150	0.66	35	NA	Mild colitis	NA
Group 4: Chronic SIV-infected, cART and Vehicle treated (VEH/SIV/cART)						
VEHSIVART1, VEHSIVART2, VEHSIVART3, VEHSIVART4, VEHSIVART5, VEHSIVART6, VEHSIVART7, VEHSIVART8*[c]	150	ND	NA	ND	NA	ND
Group 5: Chronic SIV-infected and THC treated and cART (THC/SIV/cART)						
THCSIVART1, THCSIVART2, THCSIVART3, THCSIVART4, THCSIVART5, THCSIVART6, THCSIVART7, THCSIVART8*[c]	150	ND	NA	ND	NA	ND

NA- not applicable, ND- none detected

* denotes animals pre-infection timepoint used as control

[a] denotes animals used for reduced representation bisulfite sequencing study

[b] denotes animals used for microarray gene expression study

[c] denotes animals used for NLRP6 and necroptosis immunofluorescence (confocal) studies

A VEH/SIV

HIV/SIV

Gene transcription

NLRP6 | MB21D1 (cGAS)

CpG island promoter

NLRP6 inflammasome activation

Dysbiotic microbes: *Gammaproteobacteria* and *Entrococcus faecalis*

Dysbiotic-microbe-derived metabolites

HIV/SIV

○ Decreased methylation at CpG Site

● Increased methylation at CpG site

B THC/SIV

HIV/SIV

Lactobacillus

Bifidobacteria

Methyl-donor via folate synthesis

Gene suppression

NLRP6 | MB21D1 (cGAS)

THC

NLRP6 inflammasome activation

Commensal microbes: *Clostridia, Lactobacillus,* and *Bifidobacteria*

Microbe-derived metabolites

C

HIV/SIV

LTA

Pathogenic bacteria

TLR2

HIV/SIV dsRNA

Poly I:C

HIV/SIV dsRNA

TLR3

LTA

NLRP6

NLRP6 | DHX15

CYLD

THC

P RIPK1

P RIPK1

RIPK3

P MLKL

Necrosome

Necroptosis

P RIPK3

P MLKL

TRPM7

Membrane disruption

Inflammation

HMGB1

THC

D

P MLKL | P MLKL | P MLKL

Ca²⁺

TRPM7

Cytoplasm

Ca²⁺

Extra Cellular Matrix

Ca²⁺

Plasma membrane disruption

Cytoplasm

Extra Cellular Matrix

Neutrophils

HMGB1

HMGB1

HMGB1

Macrophages

HMGB1

Inflammation ↑