Prey attracting but not avoiding predators suggests an asymmetric

investment in the predation sequence

Nicolas Ferry[1] – ferrynicolas@me.com

Christian Fiderer[1,2] – Christian.Fiderer@lazbw.bwl.de

Anne Peters[1,2] – Anne.Peters@npv-bw.bayern.de

Axel Ballmann[3] – Axel.Ballmann@npharz.de

Marco Heurich[1,2,4] – Marco.Heurich@npv-bw.bayern.de

[1] Department of National Park Monitoring and Animal Management, Bavarian Forest

National Park, Grafenau, Germany

[2] Chair of Wildlife Ecology and Wildlife Management, Faculty of Environment and Natural

Resources, University of Freiburg, Germany

[3] Harz National Park, Bad Harzburg, Germany

[4] Institute for Forest and Wildlife Management, Inland Norway University of Applied

Sciences, EvenstadNorway

Corresponding author

Nicolas Ferry – ferrynicolas@me.com

Keywords

behavioural ecology, predation, predation sequence, spatio-temporal interactions,

attraction/avoidance, camera-trap, recurrent events, large mammal, temperate forest.

Abstract

Understanding predator-prey interactions, particularly how species use space and time to influence encounter rates, is a central question in ecology. For large, free-ranging organisms, however, encounter rates are rarely quantified directly (e.g., via telemetry), due to both logistical and methodological challenges and are instead approximated through proxies such as spatial or temporal overlap. However, such proxies lack of precision regarding encounter rates estimation due to the exclusion of one of the dimensions (spatial or temporal). Camera traps, while not suited to directly measure encounters of large free-roaming species, can help overcome these limitations by estimating to what extent species tolerate or avoid one another's proximity. We analysed data from a one-year study across four German protected areas using 283 camera traps and applying recurrent event analysis to investigate interactions among three prey species (red deer *Cervus elaphus*, roe deer *Capreolus capreolus*, wild boar *Sus scrofa*) and two predators (grey wolf *Canis lupus*, Eurasian lynx *Lynx lynx*). Prey visitation rates were unaffected by predator presence, but wolves exhibited a strong attraction to prey, with visitation rates approximatively seven times higher right after prey detection. Limited data prevented robust conclusion regarding the Eurasian lynx. These findings point towards an asymmetry in predation sequence (i.e., spatio-temporal proximity, encounter, ignorance or avoidance post-encounter, capture or escape from attack): predators must succeed at every stage of the sequence to capture prey, while prey can "fail" at some steps (e.g. by decreasing spatio-temporal proximity, as in this study) but eventually escape predation by disrupting the process at any other stage.

Introduction

Understanding predator-prey interactions is fundamental to ecology, with implications for individual fitness, population dynamics, community structure, and ecosystem functions (Sih et al. 1985; Krebs et al. 1995). Holling (1959) as well as Lima & Dill (1990) decomposed the predation into a sequence of successive steps leading to the predatory event: (a) spatio-temporal proximity, (b) encounter, (c1) ignorance or (c2) avoidance post-encounter or eventually (c3) the attack leading either to (d1) the escape or (d2) the capture of the prey, i.e. the predation event. A key component is therefore the rate of encounters, i.e., when the detection range of the predator or prey exceeds the distance between them (*sensu* Lima & Dill, 1990). All other factors being equal, prey should seek to maximize use of safer areas and periods, following a dynamic landscape of fear (Messier & Barrette, 1985; Laundré et al.

2001, Palmer et al. 2017), i.e., the spatio-temporal variation of predation risk perceived by

prey (Gaynor et al. 2019), in order to minimize the encounter rate. On the other hand, optimal

foraging predicts that predators, seeking to maximize the encounter rate, should forage where

there is the higher chance to find and catch prey (Sih, 1980). These two opposite interests, i.e.

the "space race" (Sih, 1984), can lead to counter-intuitive patterns except when considering

both predator and prey simultaneously (e.g. Laundré et al. 2009, Kuijper et al. 2013). Further,

the physical landscape affects predator and prey spatial and temporal use, in part due to

energy expenditure (Dickinson et al., 2000) but also due to their perceived encountering

probability with their antagonist. Prey may choose habitats reducing encounter rates with

predators or lowering death probabilities upon encounters (Heithaus et al., 2009). However,

how the environment shapes the predation interaction is complex. For instance, vegetation

density can either provide safety or act as an obstacle, depending on context and species

(Hernandez & Laundré, 2005; Ordiz et al., 2011; Camp et al., 2013, Kuijper et al. 2013).

Estimating this encounter rate is challenging, especially for large free-ranging animals

(Hebblewhite et al., 2005, Hoffman et al., 2022). Traditionally, spatial overlap has been the

primary proxy used (e.g., occupancy, MacKenzie et al., 2004; resource selection probability,

Hebblewhite et al., 2005; habitat use similarity, Smith et al., 2019a). However, space overlap

solely is insufficient to approximate the encounter rate (Suraci et al. 2022), which depends on

predator and prey biology (e.g., population density, foraging tactic, locomotion, sensory

abilities) as well as environmental conditions (Sih, 1984; Lima & Dill, 1990). Even with

complete spatial overlap, low population densities (Travis & Palmer, 2005; Sims et al., 2006)

or slower movement speeds (Sih, 1984) can reduce encounter rates. Additionally, predators

and prey must co-occur in both space and time for encounters to happen (Sih, 1984; Minta,

1992; Caro, 2005; Sih, 2005; Suraci et al., 2022). Temporal overlap within cycles (e.g.,

circadian) can serve as another proxy for encounter rates, facilitated by analytical

advancements (e.g., diel activity pattern overlap, Ridout & Linkie, 2009) and technologies

(e.g., GPS collars, Eriksen et al., 2011; camera traps, Harmsen et al., 2009). However,

summarizing linear time in a circular pattern (yearly or daily) only provides a partial view.

High temporal and spatial overlap may occur, but individuals can avoid encounters by

consistently being out of sync by a day or more. Further, interpreting these overlaps can be

challenging in regard to the race between prey and predator without proper experimental

design (e.g., in presence/absence of the predator). Thus, considering spatial and temporal

overlaps separately is insufficient for understanding how predators and prey aim to win the

predation game. With large free-roaming species, telemetry is certainly the most accurate

method for estimating encounter rates and attraction/avoidance processes (e.g., Li et al., 2013,

Latombe et al., 2014). However, there are few empirical studies on this topic (e.g., Courbin et

al., 2013; Middleton et al., 2013; Isbell et al., 2018; Tallian et al., 2023; Elbroch et al., 2017;

Rafiq et al., 2020). This scarcity is likely due to the difficulty of simultaneously monitoring

two large free-roaming species with sufficient sample sizes and defining an adequate

detection radius to identify encounters (Lynch et al., 2015, Latombe et al., 2014). Stationary

monitoring devices, like camera traps, cannot accurately estimate encounter rates because

they rarely capture direct encounters (Triguero-Ocaña et al., 2020, Suraci et al., 2022), except

in specific situations such as predation on corralled livestock (e.g., Hoffman et al., 2022).

However, they can assess proximate co-occurrence, or the likelihood of one species being

detected after another at a location within a certain time frame, indicating whether species

tolerate or avoid each other (Schliep et al., 2018). Thus, camera traps can help bridge the gap

between spatial/temporal overlaps and encounter rates (Suraci et al., 2022).

Grey wolf (*Canis lupus*) and Eurasian lynx (*Lynx lynx*) populations are recovering in

western and central Europe after near-extinction (Chapron et al., 2014). Wolves are cursorial

predators, relying on long-distance movements (Mech, 1970; Peterson & Ciucci, 2003). Red

deer (*Cervus elaphus*), roe deer (*Capreolus capreolus*) and wild boar (*Sus scrofa*) are a

significant part of the wolves' diet (Newsome et al., 2016). Lynx are ambush predators,

mainly preying on roe deer in central Europe, occasionally on red deer and rarely on wild boar

(Khorozian & Heurich, 2023; Okarma et al., 1997; Molinari-Jobin et al., 2007; Krofel et al.,

2011; Heurich et al., 2016). Because of the recolonisation of European landscapes by wolf

and lynx, these three herbivores face an increasing risk of non-human predation, providing

new opportunities to examine predator-prey interactions in European ecosystems (see a

review from on wolf non-consumptive effects Gerber et al. 2024). From a prey perspective,

apart from human-induced mortality, wolf predation is the main cause of death for red deer

while lynx predation is less significant, with lynx preying mainly on juvenile and female red

deer (Kamler et al., 2007; Okarma et al., 1995; Belotti et al., 2015). Roe deer is similarly

predated by both lynxes and wolves, whereas wild boar is primarily predated by wolves and

almost never by lynx (Okarma et al., 1995). Using camera trap data from natural system in

which the three ungulate species and at least one of the large predators occur represents

therefore an ideal situation to assess how animals are using space and time according to

different level of predation pressure.

In this study, we used camera traps to investigate the proximate co-occurrence

between the two predators (wolf, lynx) and the three herbivores (red deer, roe deer and wild

boar). Based on prey prevalence in a predator's diet, predator impact on prey mortality, and

prey responses to predator hunting strategies (cursorial vs. ambush, Kuijper et al., 2014;

Sunde et al., 2022), we hypothesized various scenarios as follows (see Figure 1 for a visual

depiction of the hypotheses):

(H1). Red deer show avoidance towards wolves and, to a lesser extent, towards lynx.

(H2). Roe deer show avoidance towards both wolves and lynx.

(H3). Wild boars show avoidance towards wolves and remain neutral towards lynx.

(H4). Wolves are attracted to all three herbivores.

(H5). Lynx are attracted to roe deer, less attracted to red deer, and neutral towards

132   wild boars.

Further, information on previous species presence is mandatory for observing a reactive

avoidance or attraction effect. Olfactory clues like fur, skin, feces, or urine, are recognized as

important signals for heterospecific local presence (Kuijper et al., 2014; Parsons et al., 2018,

Lawson et al. 2019 and references therein) and are assumed to diminish over a few days or a

week (Lima, 1998; Parsons et al., 2018; Prat-Guitart et al., 2020). Consequently, we anticipate

that all visitation rates will return to baseline levels after a few days (Figure 1). Finally,

considering the influence of the landscape on predator-prey interactions, we hypothesize that:

(H6). Prey avoidance will be higher towards wolves in open areas and towards lynx in

denser areas.

(H7). Predator attraction will be higher towards prey in open areas for wolves and in

denser areas for lynx.

Figure 1: diagram of the expected visitation rate dynamics over time based on hypotheses H1-

H5. The pattern depends on the effects type (avoidance, neutral, attraction) and is relative to

the baseline visitation rate, i.e., under "usual" circumstances: 1) avoidance, starting lower than

baseline level and increasing until coming back to this last (H1-H3), 2) neutral, no change

from the baseline level (H3 and H5) and 3) attraction, starting higher than baseline level and

decreasing until coming back to this latter (H4-H5).

Methods

Study sites

Camera-trap data was collected from four large, protected areas in Germany: Bavarian

Forest National Park (BFNP), Müritz National Park (MNP), Harz National Park (HNP), and

Königsbrücker Heide wilderness area (KHWA, Figure 2). Due to the protection status of the

study areas, human activities are restricted to recreation, hunting, and forestry operations

necessary to meet area-specific management objectives. Across all areas, the primary goal of

hunting is to reduce browsing pressure and minimize conflicts arising from browsing and crop

damage outside the protected areas, thereby reducing conflicts with neighboring landowners,

such as farmers and foresters. In HNP, hunting of red deer, roe deer, and wild boar takes place

from August to December throughout the entire park, using both drive hunts and individual

hunting. In MNP, hunting is permitted in 35% of the area and is conducted through drive

hunts and individual hunting. Red deer and roe deer are hunted from August to January, with

male and yearling roe deer additionally hunted in May. The hunting season for fallow deer

extends from September to January, while wild boar are hunted year-round. In the BFNP,

hunting is limited to the management zone covering 25% of the park's border area. Here,

individual hunting and baiting are practiced. Red deer are hunted from June to January, and

wild boar are hunted year-round. In the KHWA, single hunt and drive hunts are permitted in

the bordering areas (ca. 19 percent of the total protected area). Hunting is mainly focused on

red deer, fallow deer (both from August to January), roe deer (males and young adults from

mid-April to January and females and fawns from August to January) and wild boar (year-

round). These areas were sampled in the project "Ungulate Monitoring in German National

Parks" from 2019 to 2020.

Lynx were reintroduced to the BFNP during the 1970s and 1980s (Wölfl et al., 2001).

In the HNP, reintroduction took place between 2000 and 2006, with animals originating from

German and Swedish zoos and wildlife parks (Mueller et al., 2020). Natural dispersal and the

reestablishment of lynx in currently unoccupied areas across Germany are hindered by habitat

fragmentation, large distances between populations, road mortality, and likely illegal killings

outside protected areas (Kramer-Schadt et al., 2004; Magg et al., 2016). As no reintroduction

efforts have been undertaken in MNP and KHWA, these areas are currently not inhabited by

lynx. Wolves naturally recolonized Germany since the early 2000s, with wolves migrating

from Eastern Europe, and were present in all areas during the time data was collected

(Jarausch   et   al.,   2021).

Table 1: study sites description. *m.a.s.l.: meter above sea level. **Definition of common mammal species was done based on the study camera-

trap monitoring.

Study area	Size (km²)	Hunting free area (%)	Elevation (m.a.s.l.*)	Overall habitat type	Main habitat	Common mammal species**
BFNP	243	75%	600 - 1,453	Low mountain range	Alluvial spruce-, mixed alpine-, mountain spruce forest (Kreft 2011)	Red deer, roe deer, wild boar, Eurasian lynx, red fox, badger, pine marten
HNP	247	0	200 - 1,141	Low mountain range	Mountain spruce-, mixed montane-, beech forests (Kreft 2011)	Red deer, roe deer, wild boar, Eurasian lynx, red fox, badger,
MNP	322	65%	62 - 143	lowland	Sub-alpine dwarf shrubland/moors/running water and rock biotopes (Kreft 2011)	Red deer, roe deer, wild boar, fallow deer, wolf, red fox, badger, racoon, hare
KHWA	69	81%	100 - 160	lowland	Riparian forest, pine bog forests, heaths, grassland, mires,	Red deer, roe deer, wild boar, wolf, red fox, badger, hare

<u>Camera-trap deployment</u>

A total of 285 camera traps were deployed between October 1, 2019, and October 1, 2020, resulting in 91,625 camera-trapping days (Table 2). When calculating trapping days, periods during which cameras were inactive or unable to record images—e.g., due to snow covering the lens—were excluded. In all study areas we used Cuddeback C2 cameras equipped with an infra-red flash. In MNP, Reconyx Hyperfire cameras were used at ten additional and randomly selected locations, while in BFNP, we used 50 Bushnell Trophy Cam cameras in the northern area and 50 Cuddeback C2 in the southern area. In July, all 100 cameras in the BFNP were replaced with Cuddeback G models.

Camera trap locations were selected using a systematic random sampling approach based on a randomly placed grid with a cell size of 1x1 km, where the center of each grid cell served as a potential camera location. Cameras were therefore not deliberately placed along game trails or other features that might increase the likelihood of wildlife detection (Wearn & Glover-Kapfer, 2017). The 1x1 km grid cell size was chosen to ensure spatial independence among camera trap locations (Wearn & Glover-Kapfer, 2017). Grid cells were excluded prior to final selection if (1) their centers did not overlap with the study area, or (2) if the grid cell center was located in an area unsuitable for camera deployment, such as water bodies, steep slopes, or fenced areas. From the remaining grid cells, 50–100 were randomly and evenly selected across each study area using ArcGIS software. In cases where the number of available cameras within a national park was insufficient to cover the entire area - such as in HNP and MNP - the park was divided into two sub-areas, and only one sub-area was surveyed. Consequently, in HNP, cameras were deployed only in the section located within the federal state of Lower Saxony (western part), while in MNP, only the northern section was surveyed (Fig. 1). Camera traps were installed within 25 m of the designated grid cell center, mounted either on trees or on wooden posts in open habitats. The lens height was set at 50 cm to ensure effective detection of medium- to large-sized mammals. Cameras were positioned to provide at least 8 m of unobstructed view along the line of sight and oriented northward ($\pm 10°$) to guarantee conscious orientation and minimize reflections from direct sunlight. All camera traps were configured to capture three consecutive still images with minimal delay upon activation. Video mode was not used.

218 Table 2: camera-trap deployment information per study site. Range of CT days represents the
219 minimum and maximum number of days a camera-trap was active. CT days is the
220 accumulated camera-trap days.

National Park	Cameras	Camera model	Range of CT days	CT days
BFNP	100	50 Cuddeback C2 + 50 Bushnell Trophy Cam / 100 Cuddeback G	29 - 367	30,576
HNP	60	Cuddeback C2	50 – 367	19,444
MNP	70	Cuddeback C2 + 10 Reconyx	7 - 367	22,377
KHWA	55	Cuddeback C2	283 - 367	19,228

221 Figure 2: Map of the four protected areas monitored in Germany from October 2019 to
222 October 2020. The number of camera traps deployed in each area is specified next to the
223 area's name (N). Black dots indicate camera trap locations. The number of independent wolf
224 and lynx events is provided for each area (N_{events}).

Vegetation density

In August 2019, vegetation density around each camera trap was measured. To assess
vegetation density, a stake was placed 8 m in front of the camera along its visual axis. Four
additional markers were placed at 4 m and 12 m along the camera's visual axis, and 4 m
west and east of the centre, forming a square around the central point with 90° angles.
Vegetation density was assessed at the four outer markers. A red towel was stretched from
each marker toward the centre and photographed at 100 cm lens height in landscape format,
using automatic mode without zoom or flash. The towel was held taut and parallel to the
ground, with edges aligned to the image frame; camera angles were adjusted on slopes as
needed. Inaccessible markers were visually estimated, and if obstacles prevented towel
placement, it was positioned at the nearest feasible location, with the adjusted distance
recorded (Abrams et al., 2018). Pictures were cropped to the extent of the red blanket and
converted to black and white using the image processing software Gimp2 (https://gimp.org).
Vegetation density was calculated as the ratio of black to white pixels, averaged across all
four photos (0 = completely open, 1 = completely closed). Analysis of the vegetation pictures
took place with the R-package *imageSeq* (Niedballa et al., 2022).

Camera trap data preparation and "survey" creation

All data preparations and analyses were done using R 4.3.0 open-source software (R Core
Team, 2023). Each event was a detection of either a single individual or a cluster of

individuals. Multiple detections of the same species and at the same location less than five

minutes apart were considered as a single independent event (Henrich et al., 2022). Following

Ferry et al. (2024) approach, when quantifying for instance the effect of wolf on red deer

visitation rate over time, wolf and red deer were designated as the *primary* (the one affecting)

and *secondary* (the one affected) species, respectively (*sensu* Ferry et al., 2024). Whenever a

wolf, the *primary*, was observed at a site, a new "post-primary survey" (*survey* hereafter)

commenced at that site and continued for a defined period (details on survey duration below).

Throughout this survey period, all observations of red deer, the *secondary* species, were

recorded and treated as *recurrent* events (Appendix 1 - Fig. S1). It was acknowledged that

additional species might also influence the secondary visitation rates, either through attraction

or avoidance. To mitigate potential confounding effects from these species when assessing the

impact of *primary* on *secondary* visitation rates, we established a *censoring* species pool

(Ferry et al. 2024). This censoring pool consisted of the primary species and *tertiary*

species—those other than the primary assumed to affect the secondary species through

avoidance or attraction. Therefore, if we assessed the effect of wolf on red deer visitation rate,

wolf was the *primary*, red deer was the *secondary* and lynx and human were considered as

tertiary potentially species affecting red deer visitation rate. The occurrence of any of these

three censoring species (wolf, lynx, human) ended the current survey (Fig. S1 A). If we

assessed the effect of wild boar (*primary*) on lynx (*secondary*), the censoring pool comprised

wild boar but also red deer, roe deer and human (*tertiaries*). Camera traps sometimes failed

due to issues such as empty batteries or snow obstruction. Extended inactivity periods

increase the chance of missing species observations, leading to an underestimation of the

event rate. Inactivity periods were defined as survey "cut-offs" based on i) deployment data

indicating camera trap daily activity and ii) the mean duration (in days) between two

independent animal observations at the camera trap site. If an inactivity period exceeded the

average duration between two events, it was considered a cut-off due to a high chance of

missing animal observations, concluding the survey (Appendix 1 - Fig. S1 B). Survey could

run up to 300 days long. However, considering olfactory cues diminish over time (Prat-

Guitart et al., 2020; Lima, 1998; Parsons et al., 2018), we set an arbitrary maximum survey

duration of 30 days. This timeframe balances capturing primary's effect on the secondary

visitation rate and ensuring model estimation is not solely influenced by unrelated factors

(e.g., resource availability phenology). As a summary, the survey ended with one of the

following censoring events: i) another species from the censoring pool observed (primary or

tertiary), ii) prolonged camera trap inactivity ("cut-off" period), iii) the survey reached a 30-
day duration threshold or iv) study's end.

The transformation of raw camera trap data into the recurrent event format was done
using the *recurrent* function from the *ct.recurrent* package (Ferry et al. 2024). For each
recurrent event, the time since the start of each survey (t.stop) and the time of the previous
recurrent event since the survey's start (t.start, equals 0 if it was the first recurrent event) were
determined. We recorded the event number and event status (0 = censoring event, 1 =
recurrent event). A second data transformation step was required, from the recurrent event
format, to obtain piece-wise exponential data using the *as_ped* function from the *pammtools*
package (Bender et al. 2018). Data has first to be discretized into J intervals with cutpoints 0
$= \tau_0 < \tau_1 < \cdots < \tau_J$. The j^{th} interval is defined as $(\tau_{J-1}, \tau_J]$ and τ_J as the maximum time set
arbitrarily, in this case, 30 days or less. Pseudo-observations were then created for each
interval during which the survey was at risk of an event (Bender et al. 2018; Ramjith et al.
2021, see an example with ecological data in Ferry et al. 2024).

<u>Analysis</u>

We then applied the Piece-wised Additive Mixed Model (PAMM, Bender & Scheipl,
2018; Ramjith et al. 2022) to estimate the temporal dynamics of the secondary species event
rate after the passage of the primary species and tested an effect of the vegetation density on
these dynamics. As part of the GAM family, this model estimates linear and non-linear
covariate effects using smoothing functions (see Ramjith et al., 2022 for a more detailed
mathematical description). It enables the non-parametric estimation of the effect of time since
the last primary event on the secondary's visitation rate. Separate models were conducted for
each prey-predator and predator-prey combination. We hypothesized different visitation rate
dynamics according to vegetation density and tested this by examining the interaction
between vegetation density and time after primary using bivariate penalized splines
represented by tensor products, when a large enough sample size allowed it (e.g. Ramjith et
al. 2022, Ferry et al. 2024). All models included a survey's random effect to address
dependency between recurrent events of the same survey (*survey_id*), as well as a random
effect of the National Park (*NP*). The effect of smooth functions is expressed in estimated
degrees of freedom (EDF), indicating the curvature's complexity. The p-values test whether
the smooth function significantly deviates from a flat line (Wood, 2017).

Results

Red deer were most frequently observed (n = 16276), followed by wild boar (n = 6980) and
roe deer (n = 3832, Fig. 3. Survey numbers were primarily constrained by predator events
(Figure 4). Wolves were the most prevalent predators in the four protected areas (n = 269),
while lynx were only observed in two (n = 41, Figure 2 & 4). This led to fewer post-lynx
surveys and limited lynx recurrent events (Figure 3 & 4). In contrast, there was ample
information on post-wolf surveys and wolf recurrent events. The limited data on combinations
involving roe deer, lynx, and wolves, or wild-boar and lynx, prevented testing the effect of
vegetation density (Table 1, Appendix 1).

Figure 3: Graph of prey recurrent events within post-predator survey, structured by prey
(rows) and predator species (columns, adapted from Chiou et al. 2021). Each grey line
represents a survey, sorted by decreasing duration (max. 30 days). Black dots represent
independent prey recurrent events within the surveys. The number of surveys and recurrent
events per combination is shown for each plot. Rows, from top to bottom: red deer, roe deer,
wild boar. Columns, from left to right: Eurasian lynx, wolf.

Figure 4: Graph of predator recurrent events within post-prey survey, structured by predator
(columns) and prey species (rows, adapted from Chiou et al. 2021). Each grey line represents
a survey, sorted by decreasing duration (max. 30 days). Black dots represent independent
predator recurrent events within the surveys. The number of surveys and recurrent events per
combination is shown for each plot. Rows, from top to bottom: Eurasian lynx, wolf. Columns,
from left to right: red deer, roe deer, wild boar.

The prey response to predators showed notable changes only for red deer following
lynx occurrences (Table 3). Red deer exhibited avoidance behavior, with a slight increase in
visitation rate after twenty days, though variability was substantial (Figure 5), and surveys
after twenty days were limited (Figure 3). No patterns were observed for wild boar or roe deer
(Table 3, Figure 5), and no interaction effect with vegetation density was found for the
combination wild boar following wolf (Table 3).

Lynx showed no clear response to prey, except for a slight decrease in visitation rate
over time after red deer occurrences, suggesting attraction (Table 3, Figure 6). Wolves
exhibited visitation rates ca. 6, 4.2, and 7 times higher immediately after encountering red
deer, roe deer, and wild boar, respectively, indicating strong initial attraction to all prey
(Table 1, values approximated from the difference of visitation between t = 0 day and either
the plateau or the end of the smooth function estimation around t = 10 days, dotted horizontal
lines Figure 6). The variability in wolf visitation rate after fifteen days, particularly following
red deer and roe deer encounters, was due to limited surveys and few wolves' recurrent events

(Figure 6). No wolf recurrent events occurred after fourteen days following wild boar

encounters, precluding further trend estimation (Figure 4 & 6). No interaction effect with

vegetation density was observed (Table 3).

<u>Table 3:</u> PAMM results for each primary-secondary combination include the estimated

degrees of freedom (EDF), indicating the polynomial degree required to describe curvature.

EDF is not indicative of effect size or direction (e.g., positive). The associated p-value tests

for non-linear effect of time, i.e. whether the smooth function significantly differs from a flat

line.

Primary	Secondary	Variable	EDF	Chi²	p-value
Lynx	Red Deer	Time	3.25	26.56	**<0.001**
		Time*Vegetation density	2.23	2.14	0.34
	Roe Deer	Time	1.15	0.06	0.86
	Wild Boar	Time	2.4	6.26	0.1
Wolf	Red Deer	Time	1.86	2.45	0.41
		Time*Vegetation density	1	0.01	0.91
	Roe Deer	Time	1	0.01	0.97
	Wild Boar	Time	2.92	8.56	0.05
		Time*Vegetation density	3.64	7.31	0.15
Red deer	Lynx	Time	1	2.07	0.15
	Wolf	Time	2.45	47.46	**<0.001**
		Time*Vegetation density	1	0.05	0.82
Roe Deer	Lynx	Time	1.01	0.75	0.39
	Wolf	Time	2.49	9.83	**0.02**
Wild Boar	Lynx	Time	1.75	2.06	0.392
	Wolf	Time	1	9.8	**0.001**
		Time*Vegetation density	2.77	6.48	**0.1**

<u>Figure 5:</u> Estimated prey visitation rate over 30 days after predator occurrence. Columns

represent prey species (red deer, roe deer, wild boar), and rows depict predators (Eurasian

lynx, wolf).

<u>Figure 6:</u> Estimated predator visitation rate over 30 days after prey occurrence. Columns

represent predator species (Eurasian lynx, wolf); rows depict prey species (red deer, roe deer,

wild boar). Horizontal dashed lines in the right column compare wolf visitation rates at t = 0

and t = 10 days. High variability after fifteen days in wolf estimates is due to limited surveys

(Fig. 4).

Discussion

Our findings did not support the hypotheses of avoidance effects by grey wolves on all

three prey species or by Eurasian lynx on roe deer and wild boar (H1, H2, H3). Only red deer

showed a slight, enduring avoidance towards lynx. Conversely, our results supported the

hypothesis that wolves respond to prey presence with a significant increase in visitation rates

(H4). Lynx showed only a slight attraction to red deer (H5) but not to roe deer or wild boar.

The limited number of post-roe deer surveys and lynx events may explain the lack of response

to roe deer. Placing camera traps on trails (Filla et al. 2017), where lynx are more active at

night (Heurich et al. 2014), could improve sample size. Consequently, our discussion will

focus on the wolves' response. Finally, no effect of vegetation density was observed for either

prey (H6) or predators (H7).

<u>The predation sequence</u>

The study aimed to assess proximate co-occurrence, with an emphasis on understanding

predator-prey interactions. While the focus was not on providing a complete approximation of

encounter rates, by considering also spatial and temporal overlaps, some interesting

conclusions can nonetheless be drawn. Wolves showed expected attraction patterns towards

prey, supporting our hypothesis H4. However, prey did not show the anticipated avoidance

patterns, as expected under the landscape of fear theory. We suggest three reasons behind this

asymmetry in reactive response.

First, predators and prey face different pressures all along the predation sequence.

Among all components (i.e. spatio-temporal proximity, encounter, ignorance or avoidance

post-encounter, capture or escape from attack), prey can optimize specific components of

their response to avoid predation, without necessarily addressing all the other components,

even completely ignoring some of them. For instance, some may specialize in avoiding risky

habitats, while others rely on early detection through vigilance, grouping, or alternative

defensive mechanisms (Gaynor et al., 2019, Gerber et al., 2024). Predators, while certainly

not always maximizing their predation rate (e.g. when resting, drinking, territory exploration

etc.), must however achieve a minimum success probability in each component when

attempting to complete the predation sequence. Thus, prey responses may seem

counterintuitive (e.g., Hebblewhite et al., 2005; Kauffman et al., 2007; Cuzack et al. 2017;

Prat-Guitart et al., 2020; Sunde et al., 2022) but would reflect this flexibility of managing

predation risks, showcasing the complexity of predator-prey interactions.

Second, this absence of response from prey might be related to the value of the

information on the current location of a predator. In their spatially explicit individual-based

model mimicking predator-prey space use and foraging, Fraker & Luttberg (2012) observed

that as movement and perception of predator increased relatively to those of prey, the latter

being more neutral toward predator location. Given a predator could be everywhere, the

information of where they were the step before had no value and prey were then selecting for
high quality patch rather than avoiding predators. In our study, given wolf are large-territory
animals (95% estimated AKDE Home Range = [50; 570] km², Vorel et al., 2024) compared to
their prey, and able to move fast, the cues of previous wolf presence might not be as
informative as they would from lynx, for instance.

Finally, in Ferry et al. (2024), the PAMM as well as the others tested approaches could
not efficiently detect (i.e., risk $\beta < 0.2$) weak or short-term effect (e.g. <12 hours) despite
hundreds of recurrent events available. Therefore, we could not completely rule out the
avoidance of prey towards predators, particularly regarding the response of roe deer towards
wolf, but this avoidance might be   too short or too weak to be detected solely based on
camera trap data and the current statistical approaches.

<u>Proximate co-occurrence on unmarked species</u>
Studies using unmarked species and camera traps face a limitation: the absence of individual-
level data. For example, when observing a flat visitation rate of red deer after wolf
occurrences, it's hard to distinguish between (i) the same individual repeatedly visiting or (ii)
several individuals appearing once and not returning. The first scenario suggests no response,
while the second indicates avoidance. The reality is likely a mix of but distinguishing them is
currently impossible. However, some signs can help infer the more likely scenario. Lower
population density and smaller home ranges increase the likelihood of observing the same
individual. For instance, given the red deer's population density (2 indiv/km², Tourani et al.,
2023), monthly home range (around 4.7 km²), and camera trap spacing (max. 1 trap/km²) in
the Bavarian Forest National Park, observing dozens of different individuals at a site during a
30-day survey is unlikely, suggesting the first scenario of absence of response. The low
density of predators compared to prey, especially for gregarious species like wolves (e.g.,
Mech & Peterson, 2003), makes it more likely to observe the same individual or group.

<u>The effect of the environment and human</u>
Dormann et al. (2018) stressed the importance of considering underlying processes and
confounding factors when interpreting species interactions from co-occurrence data. Multi-
Species Occupancy Models (MacKenzie et al. 2004) and joint Species Distribution Models
(Pollock et al. 2014) can help by accounting for covariate effects on species occurrence. In
spatio-temporal analyses, issues like a carcass causing apparent attraction between scavengers
or herbivores gathering at waterholes during the dry season (Ferry et al. 2016) can lead to
misinterpretations. Time-to-event analyses often struggle to incorporate covariates, but Prat-

Guitart et al. (2020) used a survival framework as an exception. PAMM, given enough data,
allowed us to examine vegetation density as a proxy for perceived predation risk and its
impact on species interaction. Our analysis found no influence of this parameter on
interactions. This may be due to the limited scope of our measurement, confined to a 4m
radius around camera traps, covering about 64 m². Additionally, camera trap placement often
avoids dense vegetation to maximize detection probability, which might not accurately reflect
the broader habitat context, hindering precise assessment of vegetation density effects.
Finally, the random placement of camera traps in the forest limited our ability to capture
human activities such as non-lethal recreation, which takes mostly place on trails, and
hunting. Both lethal and non-lethal activities can impact the spatio-temporal activity of both
predator and prey as well as their interactions (Peters et al., 2025; Courbin et al., 2022;
Bonnot et al., 2020; Gaynor et al., 2018). Often, detailed data on human activities is scarce,
limiting studies and methodology to address this topic. However, due to the marked and
widely recognised impact human activities have on wildlife behaviour, incorporating human
presence in data collection and analysis would be beneficial to understand predator-prey
interactions in Europe's human dominated landscape more holistically.

Conclusion
Our study delved into the spatio-temporal dynamics of interactions among three herbivore
species (red deer, roe deer, and wild boar) and two carnivores (wolf and lynx) using camera
trap data collected across four protected areas. Leveraging PAMM, we precisely estimated the
temporal patterns of visitation rates for each species following occurrence of another.
Contrary to our initial expectations, none of the herbivores exhibited discernible reactions to
the presence of predators. Conversely, we observed a notable attraction of wolves to all three
prey species, with lynxes showing a slight degree of attraction towards red deer. This stark
contrast suggests an inherent asymmetry in predator-prey interactions: while predators must
navigate through all stages of the predation sequence to successfully achieve a predation
event, prey species have the flexibility to evade predation by succeeding either in all or only a
portion of these stages. These intriguing findings highlight the potential existence of
asymmetry in predator-prey interactions, warranting further investigation.

Acknowledgements
The present publication is a partial result of the R&D project 'Ungulate monitoring in German
national parks' (FKZ: 3518 83 0200), which was supported by the Federal Agency for Nature
Conservation (BfN). We deeply thank Voler Spicher from Müritz National Park and Jürgen

Stein from Sachsenforst State Company, Nature Reserve Administration Königsbrücker
Heide / Gohrischheide Zeithain for providing data.

Statement of authorship
NF conceived this idea; NF led the writing of the manuscript; NF led analyses; CF, AP & AB
carried out the investigations, CF and MH acquired fundings and conceived the data
collection design. All authors contributed critically to the drafts and gave final approval for
publication.

Data accessibility statement
R   scripts   and   data   are   available   from   the   Zenodo   Digital   Repository
https://doi.org/10.5281/zenodo.13911909.

Conflict of interest disclosure
The authors declare that they comply with the PCI rule of having no financial conflicts of
interest in relation to the content of the article.

References
Abrams, J. F., Axtner, J., Bhagwat, T., Mohamed, A., Nguyen, A., Niedballa, J., Sollmann,
R., Tilker, A., Wilting, A. (2018): Studying terrestrial mammals in tropical rainforests.
A user guide for camera-trapping and environmental DNA. Leibniz-IZW, Berlin.
Bender, A. & Scheipl, F. (2018). pammtools: Piece-wise exponential additive mixed
modeling tools. arXiv:1806.01042 [stat]
Camp, M. J., Rachlow, J. L., Woods, B. A., Johnson, T. R., & Shipley, L. A. (2013).
Examining functional components of cover: the relationship between concealment and
visibility in shrub□steppe habitat. *Ecosphere*, *4*(2), 1-14.
Caro, T. M. (2005). *Antipredator defenses in birds and mammals*. University of Chicago
Press.

Chapron, G., Kaczensky, P., Linnell, J. D., Von Arx, M., Huber, D., Andrén, H., ... & Boitani,

L. (2014). Recovery of large carnivores in Europe's modern human-dominated

landscapes. *science*, *346*(6216), 1517-1519.

Chiou, S. H., Xu, G., Yan, J., & Huang, C. Y. (2021). Regression Modeling for Recurrent

Events Using R Package reReg. *arXiv preprint arXiv:2104.11708*.

Courbin, N., Fortin, D., Dussault, C., Fargeot, V., & Courtois, R. (2013). Multi□trophic

resource selection function enlightens the behavioural game between wolves and their

prey. *Journal of Animal Ecology*, *82*(5), 1062-1071.

Courbin, N., Garel, M., Marchand, P., Duparc, A., Debeffe, L., Börger, L., & Loison, A.

(2022). Interacting lethal and nonlethal human activities shape complex risk tolerance

behaviors in a mountain herbivore. Ecological Applications, 32(7), e2640.

https://doi.org/10.1002/eap.2640

Cusack, J. J., Dickman, A. J., Kalyahe, M., Rowcliffe, J. M., Carbone, C., MacDonald, D. W.,

& Coulson, T. (2017). Revealing kleptoparasitic and predatory tendencies in an

African mammal community using camera traps: a comparison of spatiotemporal

approaches. *Oikos*, *126*(6), 812-822.

Dickinson, M. H., Farley, C. T., Full, R. J., Koehl, M. A. R., Kram, R., & Lehman, S. (2000).

How animals move: an integrative view. *science*, *288*(5463), 100-106.

Dormann, C. F., Bobrowski, M., Dehling, D. M., Harris, D. J., Hartig, F., Lischke, H., ... &

Kraan, C. (2018). Biotic interactions in species distribution modelling: 10 questions to

guide interpretation and avoid false conclusions. Global ecology and biogeography,

27(9), 1004-1016.

Elbroch, L. M., López-González, C., Fitzgerald, J., Kusler, A., & Quigley, H. (2017).

Attraction-repulsion among top predators following reintroduction efforts.

Mammalian Biology, *86*, 66-69.

Eriksen, A., Wabakken, P., Zimmermann, B., Andreassen, H. P., Arnemo, J. M., Gundersen, H., ... & Storaas, T. (2011). Activity patterns of predator and prey: a simultaneous study of GPS-collared wolves and moose. *Animal Behaviour*, *81*(2), 423-431.

Ferry, N., Dupont, P., Bender, A., & Heurich, M. (2024). Introducing recurrent event analyses to assess species interactions based on camera-trap data: A comparison with time-to-first-event approaches. Methods in Ecology and Evolution, 15, 1233–1246. https://doi.org/10.1111/2041-210X.14338

Filla, M., Premier, J., Magg, N., Dupke, C., Khorozyan, I., Waltert, M., ... & Heurich, M. (2017). Habitat selection by Eurasian lynx (Lynx lynx) is primarily driven by avoidance of human activity during day and prey availability during night. *Ecology and Evolution*, *7*(16), 6367-6381.

Gaynor, K. M., Hojnowski, C. E., Carter, N. H., & Brashares, J. S. (2018). The influence of human disturbance on wildlife nocturnality. Science, 360(6394), 1232–1235. https://doi.org/10.1126/science.aar7121

Gaynor, K. M., Brown, J. S., Middleton, A. D., Power, M. E., & Brashares, J. S. (2019). Landscapes of fear: spatial patterns of risk perception and response. *Trends in ecology & evolution*, *34*(4), 355-368.

Gerber, N., Riesch, F., Bojarska, K., Zetsche, M., Rohwer, N. K., Signer, J., ... & Balkenhol, N. (2024). Do recolonising wolves trigger non□consumptive effects in European ecosystems? A review of evidence. *Wildlife Biology*, *2024*(6), e01229.

Hebblewhite, M., Merrill, E. H., & McDonald, T. L. (2005). Spatial decomposition of predation risk using resource selection functions: an example in a wolf–elk predator–prey system. *Oikos*, *111*(1), 101-111.

Heithaus, M. R., Frid, A., Wirsing, A. J., & Worm, B. (2008). Predicting ecological
consequences of marine top predator declines. *Trends in ecology & evolution*, *23*(4),
202-210.

Heurich, M., Hilger, A., Küchenhoff, H., Andrén, H., Bufka, L., Krofel, M., ... & Linnell, J.
D. (2014). Activity patterns of Eurasian lynx are modulated by light regime and
individual traits over a wide latitudinal range. *PloS one*, *9*(12), e114143.

Heurich, M., Zeis, K., Küchenhoff, H., Müller, J., Belotti, E., Bufka, L., & Woelfing, B.
(2016). Selective predation of a stalking predator on ungulate prey. PloS one, 11(8),
e0158449.

Henrich, M., Hartig, F., Dormann, C. F., Kühl, H. S., Peters, W., Franke, F., ... & Heurich, M.
(2022). Deer behavior affects density estimates with camera traps, but is outweighed
by spatial variability. *Frontiers in Ecology and Evolution*, *10*, 881502.

Hernández, L., & Laundré, J. W. (2005). Foraging in the 'landscape of fear'and its
implications for habitat use and diet quality of elk Cervus elaphus and bison Bison
bison. *Wildlife Biology*, *11*(3), 215-220.

Hoffmann, C. F., Pilfold, N. W., Ruppert, K. A., Letoluai, A., Lenguya, L., Limo, I., &
Montgomery, R. A. (2022). The Integral Nature of Encounter Rate in Predicting
Livestock Depredation Risk. *Frontiers in Conservation Science*, *3*, 808043.

Holling, C. S. (1959). The components of predation as revealed by a study of small-mammal
predation of the European Pine Sawfly1. *The canadian entomologist*, *91*(5), 293-320.

Isbell, L. A., Bidner, L. R., Van Cleave, E. K., Matsumoto-Oda, A., & Crofoot, M. C. (2018).
GPS-identified vulnerabilities of savannah-woodland primates to leopard predation
and their implications for early hominins. *Journal of Human Evolution*, *118*, 1-13.

Jarausch, A., Harms, V., Kluth, G., Reinhardt, I., & Nowak, C. (2021). How the west was
won: genetic reconstruction of rapid wolf recolonization into Germany's
anthropogenic landscapes. *Heredity*, *127*(1), 92-106.

Kamler, J. F., Jędrzejewski, W., & Jędrzejewska, B. (2007). Survival and Cause☐specific
Mortality of Red Deer Cervus Elaphus in Białowieża National Park, Poland. *Wildlife*
Biology, *13*(1), 48-52.

Kauffman, M. J., Varley, N., Smith, D. W., Stahler, D. R., MacNulty, D. R., & Boyce, M. S.
(2007). Landscape heterogeneity shapes predation in a newly restored predator–prey
system. *Ecology letters*, *10*(8), 690-700.

Khorozyan, I., & Heurich, M. (2023). Patterns of predation by the Eurasian lynx Lynx lynx
throughout its range: ecological and conservation implications. *Mammal Review*,
53(3), 177-188.

Kramer-Schadt, S., Revilla, E., Wiegand, T., & Breitenmoser, U. R. S. (2004). Fragmented
landscapes, road mortality and patch connectivity: modelling influences on the
dispersal of Eurasian lynx. *Journal of Applied Ecology*, *41*(4), 711-723.

Krebs, C. J., Boutin, S., Boonstra, R., Sinclair, A. R. E., Smith, J. N. M., Dale, M. R., ... &
Turkington, R. (1995). Impact of food and predation on the snowshoe hare cycle.
Science, *269*(5227), 1112-1115.

Kreft, V. S. (Ed.). (2011). Nationalparks in Deutschland: wild und schön.

Krofel, M., Huber, D., & Kos, I. (2011). Diet of Eurasian lynx Lynx lynx in the northern
Dinaric Mountains (Slovenia and Croatia) importance of edible dormouse Glis glis as
alternative prey. *Acta Theriologica*, *56*, 315-322.

Kuijper, D. P., De Kleine, C., Churski, M., Van Hooft, P., Bubnicki, J., & Jędrzejewska, B.
(2013). Landscape of fear in Europe: wolves affect spatial patterns of ungulate
browsing in Białowieża Primeval Forest, Poland. *Ecography*, *36*(12), 1263-1275.

Kuijper, D. P., Verwijmeren, M., Churski, M., Zbyryt, A., Schmidt, K., Jędrzejewska, B., &

Smit, C. (2014). What cues do ungulates use to assess predation risk in dense

temperate forests?. *PLoS One*, *9*(1), e84607.

Laundré, J.W. et al. (2001) Wolves, elk, and bison: reestablishing the "landscape of fear" in

Yellowstone National Park, U.S.A. Can J Zool 79, 1401–1409

Latombe, G., Fortin, D., & Parrott, L. (2014). Spatio□temporal dynamics in the response of

woodland caribou and moose to the passage of grey wolf. *Journal of Animal Ecology*,

83(1), 185-198.

Lawson, R. R., Fogarty, D. T., & Loss, S. R. (2019). Use of visual and olfactory sensory cues

by an apex predator in deciduous forests. *Canadian Journal of Zoology*, *97*(5), 488-

494.

Li, Z., Ding, B., Wu, F., Lei, T. K. H., Kays, R., & Crofoot, M. C. (2013). Attraction and

avoidance detection from movements. *Proceedings of the VLDB Endowment*, *7*(3),

157-168.

Lima, S. L., & Dill, L. M. (1990). Behavioral decisions made under the risk of predation: a

review and prospectus. *Canadian journal of zoology*, *68*(4), 619-640.

Lima, S. L. (1998). Nonlethal effects in the ecology of predator-prey interactions. *Bioscience*,

48(1), 25-34.

Lone, K., Loe, L. E., Gobakken, T., Linnell, J. D., Odden, J., Remmen, J., & Mysterud, A.

(2014). Living and dying in a multi□predator landscape of fear: roe deer are squeezed

by contrasting pattern of predation risk imposed by lynx and humans. *Oikos*, *123*(6),

641-651.

Lynch, E., Northrup, J. M., McKenna, M. F., Anderson Jr, C. R., Angeloni, L., & Wittemyer,

G. (2015). Landscape and anthropogenic features influence the use of auditory

vigilance by mule deer. *Behavioral Ecology*, *26*(1), 75-82.

MacKenzie, D. I., Bailey, L. L., & Nichols, J. D. (2004). Investigating species co‐occurrence

patterns when species are detected imperfectly. *Journal of Animal Ecology*, *73*(3),

546-555.

Magg, N., Müller, J., Heibl, C., Hackländer, K., Wölfl, S., Wölfl, M., ... & Heurich, M.

(2016). Habitat availability is not limiting the distribution of the Bohemian–Bavarian

lynx Lynx lynx population. *Oryx*, *50*(4), 742-752.

Marion, S., Demšar, U., Davies, A. L., Stephens, P. A., Irvine, R. J., & Long, J. A. (2021).

Red deer exhibit spatial and temporal responses to hiking activity. Wildlife Biology,

2021(3). https://doi.org/10.2981/wlb.00853

Mech, D.L. 1970. The wolf: the ecology and behaviour of an endangered species. Natural

History Press for the American Museum of Natural History, New York.

Mech, L. D., & Peterson, R. O. (2003). Wolf-prey relationships. In L. D. Mech, & L. Boitani

(Eds.), Wolves: Behavior, ecology, and conservation (pp. 131–161). Chicago, IL:

University of Chicago Press.

Messier, F., & Barrette, C. (1985). The efficiency of yarding behaviour by white-tailed deer

as an antipredator strategy. *Canadian Journal of Zoology*, *63*(4), 785-789.

Middleton, A. D., Kauffman, M. J., McWhirter, D. E., Jimenez, M. D., Cook, R. C., Cook, J.

G., ... & White, P. J. (2013). Linking anti‐predator behaviour to prey demography

reveals limited risk effects of an actively hunting large carnivore. *Ecology letters*,

16(8), 1023-1030.

Minta, S. C. (1992). Tests of spatial and temporal interaction among animals. *Ecological*

applications, *2*(2), 178-188.

Molinari‐Jobin, A., Zimmermann, F., Ryser, A., Breitenmoser‐Würsten, C., Capt, S.,

Breitenmoser, U., ... & Eyholzer, R. (2007). Variation in diet, prey selectivity and

home⊡range size of Eurasian lynx Lynx lynx in Switzerland. *Wildlife Biology*, *13*(4),

393-405.

Mueller, S. A., Reiners, T. E., Middelhoff, T. L., Anders, O., Kasperkiewicz, A., & Nowak,

C. (2020). The rise of a large carnivore population in Central Europe: genetic

evaluation of lynx reintroduction in the Harz Mountains. *Conservation Genetics*,

21(3), 577-587.

Newsome, T. M., Boitani, L., Chapron, G., Ciucci, P., Dickman, C. R., Dellinger, J. A., ... &

Ripple, W. J. (2016). Food habits of the world's grey wolves. *Mammal Review*, *46*(4),

255-269.

Niedballa, J., Axtner, J., Döbert, T. F., Tilker, A., Nguyen, A., Wong, S. T., Fiderer, C.,

Heurich, M., Wilting, A. (2022): imageseg: An R package for deep learning-based

image segmenta-tion. Methods in Ecology and Evolution 13(11), 2363–2371.

Okarma, H., Jędrzejewska, B., Jędrzejewski, W., Krasiński, Z. A., & Miłkowski, L. (1995).

The roles of predation, snow cover, acorn crop, and man-related factors on ungulate

mortality in Białowieża Primeval Forest, Poland. *Acta Theriologica*, *40*(2), 197-217.

Okarma, H., Jedrzejewski, W., Schmidt, K., KowALCZYK, R., & Jedrzejewska, B. (1997).

Predation of Eurasian lynx on roe deer and red deer in Bialowieza Primeral Forest,

Poland. *Acta Theriologica*, *42*(2), 203-224.

Ordiz, A., Støen, O. G., Delibes, M., & Swenson, J. E. (2011). Predators or prey? Spatio-

temporal discrimination of human-derived risk by brown bears. *Oecologia*, *166*, 59-

67.

Palmer, M. S., Fieberg, J., Swanson, A., Kosmala, M., & Packer, C. (2017). A

'dynamic'landscape of fear: prey responses to spatiotemporal variations in predation

risk across the lunar cycle. *Ecology letters*, *20*(11), 1364-1373.

Parsons, M. H., Apfelbach, R., Banks, P. B., Cameron, E. Z., Dickman, C. R., Frank, A. S., ...

& Blumstein, D. T. (2018). Biologically meaningful scents: a framework for

understanding predator–prey research across disciplines. *Biological Reviews*, *93*(1),

98-114.

Peters, A., Ferry, N., Dormann, C. F., & Heurich, M. (2025). Hunting shapes wildlife

behavioural responses to non-lethal recreational activities in a protected area. *Global*

Ecology and Conservation, e03994.

Peterson, R. O., and P. Ciucci. 2003. The wolf as a carnivore. Pages 104–130 in L. D. Mech

and L. Boitani editors. Wolves: behavior, ecology, and conservation. University of

Chicago Press, Chicago, Illinois, USA.

Pollock, L. J., Tingley, R., Morris, W. K., Golding, N., O'Hara, R. B., Parris, K. M., . . .

McCarthy, M. A. (2014). Understanding co-occurrence by modelling species

simultaneously with a joint species distribution model (JSDM). Methods in Ecology

and Evolution, 5, 397–406.

Prat-Guitart, M., Onorato, D. P., Hines, J. E., & Oli, M. K. (2020). Spatiotemporal pattern of

interactions between an apex predator and sympatric species. *Journal of Mammalogy*,

101(5), 1279-1288.

R Core Team (2023). R: A Language and Environment for Statistical Computing. R for

Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

Rafiq, K., Jordan, N. R., Wilson, A. M., McNutt, J. W., Hayward, M. W., Meloro, C., ... &

Golabek, K. A. (2020). Spatio⬜temporal factors impacting encounter occurrences

between leopards and other large African predators. *Journal of Zoology*, *310*(3), 191-

200.

Ramjith J, Bender A, Roes KCB, Jonker MA. Recurrent events analysis with piece-wise

exponential additive mixed models. 2022. Statistical Modelling., 2022

Ridout, M. S., & Linkie, M. (2009). Estimating overlap of daily activity patterns from camera

trap data. *Journal of Agricultural, Biological, and Environmental Statistics*, *14*(3),

322–337. https://doi.org/10.1198/jabes.2009.08038

Sih, A. (1984). The behavioral response race between predator and prey. The American

Naturalist, 123(1), 143-150.

Sih, A., Crowley, P., McPeek, M., Petranka, J., & Strohmeier, K. (1985). Predation,

competition, and prey communities: a review of field experiments. *Annual review of*

ecology and systematics, *16*(1), 269-311.

Sih, A. (2005). Predator-prey space use as an emergent outcome of a behavioral response

race. *Ecology of predator-prey interactions*, *256*, 78.

Sims, D. W., Witt, M. J., Richardson, A. J., Southall, E. J., & Metcalfe, J. D. (2006).

Encounter success of free-ranging marine predator movements across a dynamic prey

landscape. *Proceedings of the Royal Society B: Biological Sciences*, *273*(1591), 1195-

1201.

Smith, J. A., Donadio, E., Pauli, J. N., Sheriff, M. J., Bidder, O. R., & Middleton, A. D.

(2019). Habitat complexity mediates the predator–prey space race. *Ecology*, *100*(7),

e02724.

Sunde, P., Böcker, F., Rauset, G. R., Kjellander, P., Chrenkova, M., Skovdal, T. M., ... &

Heurich, M. (2022). Mammal responses to predator scents across multiple study areas.

Ecosphere, 13(8), e4215.

Suraci, J. P., Smith, J. A., Chamaillé□Jammes, S., Gaynor, K. M., Jones, M., Luttbeg, B., ...

& Sih, A. (2022). Beyond spatial overlap: harnessing new technologies to resolve the

complexities of predator–prey interactions. *Oikos*, *2022*(8), e09004.

Tallian, A., Mattisson, J., Stenbacka, F., Neumann, W., Johansson, A., Støen, O. G., &

Kindberg, J. (2023). Proximity□sensors on GPS collars reveal fine□scale predator–

prey behavior during a predation event: A case study from Scandinavia. *Ecology and*

Evolution, *13*(12), e10750.

Tourani, M., Franke, F., Heurich, M., Henrich, M., Peterka, T., Ebert, C., ... & Peters, W.

(2023). Spatial variation in red deer density in a transboundary forest ecosystem.

Scientific Reports, *13*(1), 4561.

Travis, J. M. J., & Palmer, S. C. F. (2005). Spatial processes can determine the relationship

between prey encounter rate and prey density. *Biology Letters*, *1*(2), 136-138.

Triguero-Ocaña, R., Vicente, J., Palencia, P., Laguna, E., & Acevedo, P. (2020). Quantifying

wildlife-livestock interactions and their spatio-temporal patterns: is regular grid

camera trapping a suitable approach?. *Ecological Indicators*, *117*, 106565.

Vorel, A., Kadlec, I., Toulec, T., Selimovic, A., Horníček, J., Vojtěch, O., ... & Barták, V.

(2024). Home range and habitat selection of wolves recolonising central European

human dominated landscapes. *Wildlife Biology*, *2024*(6), e01245.

Wearn, O. R., & Glover-Kapfer, P. (2017). Camera-trapping for conservation: a guide to best-

practices. *WWF conservation technology series*, *1*(1), 181.

Wikenros, C., Aronsson, M., Liberg, O., Jarnemo, A., Hansson, J., Wallgren, M., ... &

Bergström, R. (2017). Fear or food–abundance of red fox in relation to occurrence of

lynx and wolf. *Scientific reports*, *7*(1), 9059.

Wölfl, M., Heurich, M., Habel, H., Huber, T., & Poost, W. (2001). Distribution and status of

lynx in the border region between Czech Republic, Germany and Austria. Acta

Theriologica, 14

Wood, S. N. (2017). Generalized additive models: an introduction with R. chapman and

hall/CRC.

Germany

Harz National Park - N = 60

N events = 7
N events = 9

0 2 4 6 8 10
Kilometer

Müritz National Park - N=70

N events = 70

0 2 3 5 6 8
Kilometer

Königsbrücker-Heide Wilderness Area - N=55

N events = 181

0 1 2 3 4 5
Kilometer

Legend

• Locations of camera-traps

Bavarian Forest National Park - N=100

N events = 8
N events = 32

0 4 8 12
Kilometer

0 100 200 Kilometer

Surveys = 41
Events = 118

Surveys = 269
Events = 1549

Surveys = 41
Events = 19

Surveys = 269
Events = 199

Surveys = 41
Events = 60

Surveys = 269
Events = 685

Time after predator (days)

Time after predator (days)

Surveys = 5182
Events = 21

Surveys = 920
Events = 7

Surveys = 1481
Events = 6

Surveys = 16276
Events = 152

Surveys = 3832
Events = 25

Surveys = 6980
Events = 78

Time after prey (days)

Time after prey (days)

Time after prey (days)