Prey attracting but not repelled by predators suggests an asymmetric investment in the encounter-avoid-escape sequence.

Nicolas Ferry[1] – ferrynicolas@me.com

Christian Fiderer[1,2] – Christian.Fiderer@lazbw.bwl.de

Anne Peters[1,2] – Anne.Peters@npv-bw.bayern.de

Axel Ballmann[3] – Axel.Ballmann@npharz.de

Marco Heurich[1,2,4] – Marco.Heurich@npv-bw.bayern.de

[1] Department of National Park Monitoring and Animal Management, Bavarian Forest National Park, Grafenau, Germany

[2] Chair of Wildlife Ecology and Wildlife Management, Faculty of Environment and Natural Resources, University of Freiburg, Germany

[3] Harz National Park, Bad Harzburg, Germany

[4] Institute for Forest and Wildlife Management, Inland Norway University of Applied Sciences, EvenstadNorway

Corresponding author

Nicolas Ferry – ferrynicolas@me.com

Keywords

behavioural ecology, predation, encounter sequence, spatio-temporal interactions, attraction/avoidance, camera-trap, recurrent events, large mammal, temperate forest.

Abstract

Understanding predator-prey interactions, particularly how species use space and time to

influence encounter rates, is crucial in ecology. Camera traps, while not being able to directly

measure encounters of large free-roaming species, can help estimating how species tolerate or

avoid proximity with eachother. We used data from a one-year study in four German protected

areas used 283 camera traps and applied recurrent event analysis to explore interactions among

three prey species (red deer, roe deer, wild boar) and two predators (grey wolf, Eurasian lynx).

Prey visitation rates were unaffected by predators, but wolves showed a strong attraction to

prey, with visitation rates approximatively seven times higher right after prey occurrence.

Insufficient data hindered robust pattern estimation regarding the Eurasian lynx. This suggests

an asymmetry where predators need to succeed at every stage of the encounter sequence, while

prey can disrupt the process at any stage to avoid being caught.

Introduction

Understanding predator-prey interactions is fundamental to ecology, with implications

for individual fitness, population dynamics, community structure, and ecosystem functions (Sih

et al. 1985; Krebs et al. 1995). Predation is one possible outcome of the encounter sequence

(*sensu* Lima & Dill, 1990), with mutual ignorance, prey avoidance or escape (Lima & Dill,

1990). Encounters occur at a certain rate (Holling, 1959) when the predator or prey detection

radius exceeds the distance between them (Lima & Dill, 1990). Predators aim to maximize this

rate, while prey aim to minimize it. Estimating this encounter rate is challenging, especially for

large free-ranging animals (Hebblewhite et al., 2005). Estimating this encounter rate, depicting

the predator-prey "space race" (Sih, 1984) is challenging, especially for large free-ranging

animals (Hebblewhite et al., 2005).

Traditionally, spatial overlap has been the primary metric used (e.g., occupancy,

MacKenzie et al., 2004; resource selection probability, Hebblewhite et al., 2005; habitat use

similarity, Smith et al., 2019a). However, space overlap solely is insufficient to estimate the

encounter rate (Suraci et al. 2022), which depends on predator and prey biology (e.g.,

population density, foraging tactic, locomotion, sensory abilities) as well as environmental

conditions (Sih, 1984; Lima & Dill, 1990). Even with complete spatial overlap, low population

densities (Travis & Palmer, 2005; Sims et al., 2006) or slower movement speeds (Sih, 1984)

can reduce encounter rates. Additionally, predators and prey must co-occur in both space and

time for encounters to happen (Sih, 1984; Minta, 1992; Caro, 2005; Sih, 2005; Suraci et al.,

2022). Temporal overlap within cycles (e.g., circadian) can serve as a proxy for encounter rates,

facilitated by analytical advancements (e.g., diel activity pattern overlap, Ridout & Linkie,

2009) and technologies (e.g., GPS collars, Eriksen et al., 2011; camera traps, Harmsen et al.,

2009). However, summarizing linear time in a circular pattern (yearly or daily) only provides a

partial view. High temporal and spatial overlap may occur, but individuals can avoid encounters

by consistently being out of sync by a day or more. Thus, considering spatial and temporal

overlaps separately is insufficient for understanding how predators and prey aim to maximize

and minimize encounter rates.

Telemetry is the most accurate method for estimating encounter rates and

attraction/repulsion processes (e.g., Li et al., 2013). However, there are few empirical studies

on this topic (e.g., Courbin et al., 2013; Middleton et al., 2013; Isbell et al., 2018; Tallian et al.,

2023; Elbroch et al., 2017; Rafiq et al., 2020). This scarcity is likely due to the difficulty of

simultaneously monitoring two large free-roaming species with sufficient sample sizes and

defining an adequate detection radius to identify encounters (Lynch et al., 2015). Stationary

monitoring devices, like camera traps, cannot accurately estimate encounter rates because they

rarely capture direct encounters (Suraci et al., 2022). However, they can assess proximate co-

occurrence, or the likelihood of one species being detected after another at a location within a

certain time frame, indicating whether species tolerate or avoid each other (Schliep et al., 2018).

Thus, camera traps can help bridge the gap between spatial/temporal overlaps and encounter

rates (Suraci et al., 2022).

Grey wolf (*Canis lupus*) and Eurasian lynx (*Lynx lynx*) populations are recovering in

western and central Europe after near-extinction (Chapron et al., 2014). As a consequence,

larger herbivores, including red deer (*Cervus elaphus*), roe deer (*Capreolus capreolus*) and wild

boar (*Sus scrofa*), face an increasing risk of non-human predation, providing opportunities to

examine predator-prey interactions in European ecosystems. Apart from human-induced

mortality, wolf predation is the main cause of death for red deer while lynx predation is less

significant, and only for calves and hinds (Kamler et al., 2007; Okarma et al., 1995; Belotti et

al., 2015). Both lynx and wolves threaten roe deer, whereas wild boar is primarily predated by

wolves and almost never by lynx (Okarma et al., 1995). Wolves are cursorial predators, relying

on long-distance movements (Mech, 1970; Peterson & Ciucci, 2003), for which these three

herbivores are a significant part of its diet (Newsome et al., 2016). Lynx are ambush predators,

mainly preying on roe deer, occasionally on red deer and rarely on wild boar (Khorozian &

Heurich, 2023; Okarma et al., 1997; Molinari-Jobin et al., 2007; Krofel et al., 2011; Heurich et

al., 2016).

In this study, we used camera traps to investigate the proximate co-occurrence between

these two predators and three herbivores. Following Ferry et al. (2024), three different patterns

of visitation rate over time of a species after the occurrence of another scenarios could be

distinguished: i) a flat rate (neutral), ii) first low and increasing until reaching a plateau

(avoidance) or iii) first high and decreasing until reaching a plateau (attraction). Based on prey

prevalence, predator impact on prey mortality, and prey responses to hunting strategies

(cursorial vs. ambush, Kuijper et al., 2014; Sunde et al., 2022), we hypothesized various

scenarios:

(H1). Red deer show avoidance towards wolves and, to a lesser extent, towards lynx.

(H2). Roe deer show avoidance towards both wolves or lynx.

(H3). Wild boars show avoidance towards wolves and remain neutral towards lynx.

(H4). Wolves are attracted to all three herbivores.

(H5). Lynx are attracted to roe deer, somewhat attracted to red deer, and neutral

102   towards wild boars.

The physical landscape affects predator and prey spatial and temporal use (Lone et al.,

2014). Prey may choose habitats to reduce encounter rates with predators or lower death

probabilities upon encounters (Heithaus et al., 2009). Cover can either provide safety or act as

an obstacle, depending on context and species (Hernandez & Laundré, 2005; Ordiz et al., 2011;

Camp et al., 2013). We hypothesize that:

(H6). Prey avoidance will be higher towards wolves in open areas and towards lynx in

109   closed areas.

110   (H7). Predator attraction will be higher towards prey in open areas for wolves and in

111   closed areas for lynx.

Information on previous species presence is mandatory for observing a repulsion or

attraction effect. Olfactory clues like fur, skin, feces, or urine, are recognized as important

signals for heterospecific local presence (Kuijper et al., 2014; Parsons et al., 2018) and are

assumed to diminish over a few days or a week (Lima, 1998; Parsons et al., 2018; Prat-Guitart

et al., 2020). Consequently, we anticipate that all visitation rates will return to baseline levels

after a few days (H8).

Methods

<u>Study sites</u>

Camera-trap data was collected from four large protected areas in Germany: Bavarian

Forest National Park, Müritz National Park, Harz National Park, and Königsbrücker Heide

wilderness area. These areas were sampled in the "Ungulate Monitoring in German National

Parks" project from 2019 to 2020. Eurasian lynx were present only in Bavarian Forest and Harz

National Parks, while grey wolves were found in all four areas. Camera traps were set up within

a grid of 1 km² cells, covering at least 50 percent of the cells and a minimum of 55 cells (Fig.

1). Grid cells for camera traps were selected randomly, with traps placed within 25 meters of

the cell center, facing north. Due to the large size of Harz and Müritz National Parks, only sub-

areas were sampled: the northern part of Müritz and the Lower Saxony part of Harz (Fig. 1).

<u>Figure 1:</u> Map of the four protected areas monitored in Germany from October 2019 to October

2020. The number of camera traps deployed in each area is specified next to the area's name

(N). Black dots indicate camera trap locations. The number of independent wolf and lynx events

is provided for each area (N_{events}).

<u>Vegetation density</u>

In August 2019, vegetation density around each camera trap was measured. Four photos

were taken facing north, west, south, and east from the camera trap's center, each facing a red

blanket 4 meters away. Pictures were cropped to the extend of the red blanket and converted to

black and white using the image processing software Gimp2. cropped to the blanket's extent

and converted to black and white using Gimp2. Vegetation density was calculated as the ratio

of black to white pixels, averaged across all four photos (0 = completely open, 1 = completely

closed).

<u>Camera trap data preparation and analyses</u>

All data preparations and analyses were done using R 4.3.0 open-source software (R

Core Team, 2023). Each event was a detection of either a single individual or a cluster of

individuals. Multiple detections of the same species and at the same location less than five

minutes apart were considered as a single independent event (Heinrich et al., 2022). Following

Ferry et al. (2024) approach, we tested the effect of a primary species on a secondary species,

e.g. wolf on red deer, while considering the confounding effect of tertiary species, e.g. lynx and

human. We created surveys starting after a primary event, considering all following secondary

events, and ending when: i) another primary or tertiary species was observed, ii) prolonged

camera trap inactivity occured, iii) the survey reached a 30-day duration threshold or iv) study's

ended (see Supplementary Material 1 for survey definition and data formatting details). We

then applied the Piece-wised Additive Mixed Model (PAMM, Bender & Scheipl, 2018; Ramjith

et al. 2022) to estimate the temporal dynamics of the secondary species event rate after the

passage of the primary species and tested an effect of the vegetation density on this dynamics

(see Supplementary Material 1 for models details).

Results

Red deer were most frequently observed, followed by wild boar and roe deer. Survey

numbers were primarily constrained by predator events (Supplementary Material 2). Wolves

were the most prevalent predators in the four protected areas, while lynx were only observed in

two (Figure 1). This led to fewer post-lynx surveys and limited lynx recurrent events

(Supplementary Material 2). In contrast, there was ample information on post-wolf surveys and

wolf recurrent events. The limited data on combinations involving roe deer, lynx, and wolves,

and wild-boar/lynx, prevented testing the effect of vegetation density (Table 1, Supplementary

Material 3).

The prey response to predators showed notable changes only for red deer following lynx

occurrences (Table 1). Red deer exhibited avoidance behavior, with a slight increase in

visitation rate after twenty days, though variability was substantial (Figure 2), and surveys after

twenty days were limited (Supplementary Material 2). No patterns were observed for wild boar

or roe deer (Table 1, Figure 2), and no interaction effect with vegetation density was found

(Table 1).

Lynx showed no clear response to prey, except for a slight decrease in visitation rate

over time after red deer occurrences, suggesting attraction (Table 1, Figure 3). Wolves exhibited

visitation rates 6.6, 7, and 7 times higher immediately after encountering red deer, roe deer, and

wild boar, respectively, indicating strong initial attraction to all prey (Table 1, Figure 3). The

variability in wolf visitation rate after fifteen days, particularly following red deer and roe deer

encounters, was due to limited surveys and few wolf recurrent events (Supplementary Material

2). No wolf recurrent events occurred after fourteen days following wild boar encounters,

precluding further trend estimation (Figure 3). No interaction effect with vegetation density was

observed (Table 1).

Table 1: PAMM results for each primary-secondary combination include the estimated degrees

of freedom (EDF), indicating the polynomial degree required to describe curvature. EDF is not

indicative of effect size or direction (e.g., positive). The associated p-value tests for non-linear

effect of time, i.e. whether the smooth function significantly differs from a flat line.

Primary	Secondary	Variable	EDF	Chi²	p-value
Lynx	Red Deer	Time	3.25	26.56	**<0.001**

		Time*Vegetation density	2.23	2.14	0.34
	Roe Deer	Time	1.15	0.06	0.86
	Wild Boar	Time	2.4	6.26	0.1
Wolf	Red Deer	Time	1.86	2.45	0.41
		Time*Vegetation density	1	0.01	0.91
	Roe Deer	Time	1	0.01	0.97
	Wild Boar	Time	2.92	8.56	0.05
		Time*Vegetation density	3.64	7.31	0.15
Red deer	Lynx	Time	1	2.07	0.15
	Wolf	Time	2.45	47.46	**<0.001**
		Time*Vegetation density	1	0.05	0.82
Roe Deer	Lynx	Time	1.01	0.75	0.39
	Wolf	Time	2.49	9.83	**0.02**
Wild Boar	Lynx	Time	1.75	2.06	0.392
	Wolf	Time	1	9.8	**0.001**
		Time*Vegetation density	2.77	6.48	**0.1**

Figure 2: Estimated prey visitation rate over 30 days after predator occurrence. Columns represent prey species (red deer, roe deer, wild boar), and rows depict predators (Eurasian lynx, wolf).

Figure 3: Estimated predator visitation rate over 30 days after prey occurrence. Columns represent predator species (Eurasian lynx, wolf); rows depict prey species (red deer, roe deer, wild boar). Horizontal dashed lines in the right column compare wolf visitation rates at t = 0 and t = 10 days. High variability after fifteen days in wolf estimates is due to limited surveys (Supplementary Material 2).

Discussion

Our findings did not support the hypotheses of deterrence effects by grey wolves on all three prey species or by Eurasian lynx on roe deer and wild boar (H1, H2, H3). Only red deer showed a slight, enduring avoidance. Conversely, our results supported the hypothesis that wolves respond to prey presence with a significant increase in visitation rates (H4). Lynx showed only a slight attraction to red deer (H5) but not to roe deer or wild boar. The limited number of post-roe deer surveys and lynx events may explain the lack of response to roe deer. Placing camera traps on trails (Filla et al. 2017), where lynx are more active at night (Heurich et al. 2014), could improve sample size. Consequently, our discussion will focus on the wolves' response. Finally, no effect of vegetation density was observed for either prey (H6) or predators (H7).

The encounter sequence

The study aimed to assess proximate co-occurrence through a novel statistical approach
using camera trap data, with an emphasis on understanding predator-prey interactions. While
the focus was not on providing a complete approximation of encounter rates, by considering
also spatial and temporal overlaps, some interesting conclusions can nonetheless already be
drawn. Wolves showed expected attraction patterns towards prey, supporting the hypothesis of
encounter rate maximization. However, prey did not show the anticipated avoidance patterns.
This aligns with Sunde et al. (2022), who found no change in prey visitation after predator urine
was deposited. This asymmetry in predation interactions may be due to the different pressures
faced by predators and prey. Among all components of the encounter sequence (spatial and
temporal overlap, proximity, avoidance post-encounter, capture or escape from attack), prey
can optimize specific components of their response to avoid predation, such as reducing capture
probability through behaviors like mobbing (Crofoot, 2013), without necessarily addressing all
the other components to avoid predation. Predators, however, must achieve a minimul success
probability in each component to complete the encounter sequence. Thus, prey responses may
seem counterintuitive (e.g., Hebblewhite et al., 2005; Kauffman et al., 2007; Cuzack et al. 2017;
Prat-Guitart et al., 2020; Sunde et al., 2022) but would reflect this flexibility of managing
predation risks, showcasing the complexity of predator-prey interactions.

Proximate co-occurrence on unmarked species

Studies using unmarked species and camera traps face a limitation: the absence of
individual-level data. For example, when observing a flat visitation rate of red deer after wolf
occurrences, it's hard to distinguish between (i) the same individual repeatedly visiting or (ii)
several individuals appearing once and not returning. The first scenario suggests no response,
while the second indicates repulsion. The reality is likely a mix of both, but distinguishing them
is currently impossible. However, some signs can help infer the more likely scenario. Lower
population density and smaller home ranges increase the likelihood of observing the same
individual. For instance, given the red deer's population density (2 indiv.km², Tourani et al.,
2023), monthly home range (around 4.7 km²), and camera trap spacing (max. 1 trap.km²) in the
Bavarian Forest National Park, observing dozens of different individuals at a site during a 30-
day survey is unlikely, suggesting the first scenario of absence of response. The low density of
predators compared to prey, especially for gregarious species like wolves (e.g., Mech &
Peterson, 2003), makes it more likely to observe the same individual or group.

The effect of the environment

Dormann et al. (2018) stressed the importance of considering underlying processes and
confounding factors when interpreting species interactions from co-occurrence data. Multi-
Species Occupancy Models (MacKenzie et al. 2004) and joint Species Distribution Models
(Pollock et al. 2014) can help by accounting for covariate effects on species occurrence. In
spatio-temporal analyses, issues like a carcass causing apparent attraction between scavengers
or herbivores gathering at waterholes during the dry season (Ferry et al. 2016) can lead to
misinterpretations. Time-to-event analyses often struggle to incorporate covariates, but Prat-
Guitart et al. (2020) used a survival framework as an exception. PAMM, given enough data,
allowed us to examine vegetation density as a proxy for perceived predation risk and its impact
on species interaction. Unfortunately, our analysis found no influence of this parameter on
interactions. This may be due to the limited scope of our measurement, confined to a 4m radius
around camera traps, covering about 64 m². Additionally, camera trap placement often avoids
dense vegetation to maximize detection probability, which might not accurately reflect the
broader habitat context, hindering precise assessment of vegetation density effects.

Conclusion
Our study delved into the spatio-temporal dynamics of interactions among three
herbivore species (red deer, roe deer, and wild boar) and two carnivores (wolf and lynx) using
camera trap data collected across four protected areas. Leveraging PAMM, we precisely
estimated the temporal patterns of visitation rates for each species following occurence of
another. Contrary to our initial expectations, none of the herbivores exhibited discernible
reactions to the presence of predators. Conversely, we observed a notable attraction of wolves
to all three prey species, with lynxes showing a slight degree of attraction towards red deer.
This stark contrast suggests an inherent asymmetry in predator-prey interactions: while
predators must navigate through all stages of the encounter sequence to successfully achieve a
predation event, prey species have the flexibility to evade predation by succeeding either in all
or only a portion of these stages. These intriguing findings highlight the potential existence of
asymmetry in predator-prey interactions, warranting further investigation.

Acknowledgements
The present publication is a partial result of the R&D project 'Ungulate monitoring in German
national parks' (FKZ: 3518 83 0200), which was supported by the Federal Agency for Nature
Conservation (BfN). We deeply thank Voler Spicher from Müritz National Park and Jürgen
Stein from Sachsenforst State Company, Nature Reserve Administration Königsbrücker Heide
/ Gohrischheide Zeithain for providing data.

Statement of authorship

NF conceived this idea; NF led the writing of the manuscript; NF led analyses; CF, AB carried

out the investigations, CF and MH acquired fundings and conceived the data collection design.

All authors contributed critically to the drafts and gave final approval for publication.

Data accessibility statement

R scripts and data are available from the Zenodo Digital Repository

https://doi.org/10.5281/zenodo.13911909.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of

interest in relation to the content of the article.

References

Bender, A. & Scheipl, F. (2018). pammtools: Piece-wise exponential additive mixed modeling

tools. arXiv:1806.01042 [stat]

Camp, M. J., Rachlow, J. L., Woods, B. A., Johnson, T. R., & Shipley, L. A. (2013). Examining

functional components of cover: the relationship between concealment and visibility in

shrub-steppe habitat. *Ecosphere*, *4*(2), 1-14.

Caro, T. M. (2005). *Antipredator defenses in birds and mammals*. University of Chicago Press.

Chapron, G., Kaczensky, P., Linnell, J. D., Von Arx, M., Huber, D., Andrén, H., ... & Boitani,

L. (2014). Recovery of large carnivores in Europe's modern human-dominated

landscapes. *science*, *346*(6216), 1517-1519.

Chiou, S. H., Xu, G., Yan, J., & Huang, C. Y. (2021). Regression Modeling for Recurrent

Events Using R Package reReg. *arXiv preprint arXiv:2104.11708*.

Courbin, N., Fortin, D., Dussault, C., Fargeot, V., & Courtois, R. (2013). Multi-trophic resource

selection function enlightens the behavioural game between wolves and their prey.

Journal of Animal Ecology, *82*(5), 1062-1071.

Cusack, J. J., Dickman, A. J., Kalyahe, M., Rowcliffe, J. M., Carbone, C., MacDonald, D. W.,

& Coulson, T. (2017). Revealing kleptoparasitic and predatory tendencies in an African

mammal community using camera traps: a comparison of spatiotemporal approaches.

Oikos, *126*(6), 812-822.

Dormann, C. F., Bobrowski, M., Dehling, D. M., Harris, D. J., Hartig, F., Lischke, H., ... &

Kraan, C. (2018). Biotic interactions in species distribution modelling: 10 questions to

guide interpretation and avoid false conclusions. Global ecology and biogeography,

27(9), 1004-1016.

Elbroch, L. M., López-González, C., Fitzgerald, J., Kusler, A., & Quigley, H. (2017).

Attraction-repulsion among top predators following reintroduction efforts. *Mammalian*

Biology, *86*, 66-69.

Eriksen, A., Wabakken, P., Zimmermann, B., Andreassen, H. P., Arnemo, J. M., Gundersen,

H., ... & Storaas, T. (2011). Activity patterns of predator and prey: a simultaneous study

of GPS-collared wolves and moose. *Animal Behaviour*, *81*(2), 423-431.

Ferry, N., Dupont, P., Bender, A., & Heurich, M. (2024). Introducing recurrent event analyses

to assess species interactions based on camera-trap data: A comparison with time-to-

first-event approaches. Methods in Ecology and Evolution, 15, 1233–1246.

https://doi.org/10.1111/2041-210X.14338

Filla, M., Premier, J., Magg, N., Dupke, C., Khorozyan, I., Waltert, M., ... & Heurich, M.

(2017). Habitat selection by Eurasian lynx (Lynx lynx) is primarily driven by avoidance

of human activity during day and prey availability during night. *Ecology and Evolution*,

7(16), 6367-6381.

Hebblewhite, M., Merrill, E. H., & McDonald, T. L. (2005). Spatial decomposition of predation

risk using resource selection functions: an example in a wolf–elk predator–prey system.

Oikos, *111*(1), 101-111.

Heithaus, M. R., Frid, A., Wirsing, A. J., & Worm, B. (2008). Predicting ecological
consequences of marine top predator declines. *Trends in ecology & evolution*, *23*(4),
202-210.

Heurich, M., Hilger, A., Küchenhoff, H., Andrén, H., Bufka, L., Krofel, M., ... & Linnell, J. D.
(2014). Activity patterns of Eurasian lynx are modulated by light regime and individual
traits over a wide latitudinal range. *PloS one*, *9*(12), e114143.

Heurich, M., Zeis, K., Küchenhoff, H., Müller, J., Belotti, E., Bufka, L., & Woelfing, B. (2016).
Selective predation of a stalking predator on ungulate prey. PloS one, 11(8), e0158449.

Henrich, M., Hartig, F., Dormann, C. F., Kühl, H. S., Peters, W., Franke, F., ... & Heurich, M.
(2022). Deer behavior affects density estimates with camera traps, but is outweighed by
spatial variability. *Frontiers in Ecology and Evolution*, *10*, 881502.

Hernández, L., & Laundré, J. W. (2005). Foraging in the 'landscape of fear' and its implications
for habitat use and diet quality of elk Cervus elaphus and bison Bison bison. *Wildlife
Biology*, *11*(3), 215-220.

Holling, C. S. (1959). The components of predation as revealed by a study of small-mammal
predation of the European Pine Sawfly1. *The canadian entomologist*, *91*(5), 293-320.

Isbell, L. A., Bidner, L. R., Van Cleave, E. K., Matsumoto-Oda, A., & Crofoot, M. C. (2018).
GPS-identified vulnerabilities of savannah-woodland primates to leopard predation and
their implications for early hominins. *Journal of Human Evolution*, *118*, 1-13.

Kamler, J. F., Jędrzejewski, W., & Jędrzejewska, B. (2007). Survival and Cause-specific
Mortality of Red Deer Cervus Elaphus in Białowieża National Park, Poland. *Wildlife
Biology*, *13*(1), 48-52.

Kauffman, M. J., Varley, N., Smith, D. W., Stahler, D. R., MacNulty, D. R., & Boyce, M. S.
(2007). Landscape heterogeneity shapes predation in a newly restored predator–prey
system. *Ecology letters*, *10*(8), 690-700.

Khorozyan, I., & Heurich, M. (2023). Patterns of predation by the Eurasian lynx Lynx lynx

throughout its range: ecological and conservation implications. *Mammal Review*, *53*(3),

177-188.

Krebs, C. J., Boutin, S., Boonstra, R., Sinclair, A. R. E., Smith, J. N. M., Dale, M. R., ... &

Turkington, R. (1995). Impact of food and predation on the snowshoe hare cycle.

Science, *269*(5227), 1112-1115.

Krofel, M., Huber, D., & Kos, I. (2011). Diet of Eurasian lynx Lynx lynx in the northern Dinaric

Mountains (Slovenia and Croatia) importance of edible dormouse Glis glis as alternative

prey. *Acta Theriologica*, *56*, 315-322.

Kuijper, D. P., Verwijmeren, M., Churski, M., Zbyryt, A., Schmidt, K., Jędrzejewska, B., &

Smit, C. (2014). What cues do ungulates use to assess predation risk in dense temperate

forests?. *PLoS One*, *9*(1), e84607.

Li, Z., Ding, B., Wu, F., Lei, T. K. H., Kays, R., & Crofoot, M. C. (2013). Attraction and

avoidance detection from movements. *Proceedings of the VLDB Endowment*, *7*(3), 157-

168.

Lima, S. L., & Dill, L. M. (1990). Behavioral decisions made under the risk of predation: a

review and prospectus. *Canadian journal of zoology*, *68*(4), 619-640.

Lima, S. L. (1998). Nonlethal effects in the ecology of predator-prey interactions. *Bioscience*,

48(1), 25-34.

Lone, K., Loe, L. E., Gobakken, T., Linnell, J. D., Odden, J., Remmen, J., & Mysterud, A.

(2014). Living and dying in a multi-predator landscape of fear: roe deer are squeezed

by contrasting pattern of predation risk imposed by lynx and humans. *Oikos*, *123*(6),

641-651.

Lynch, E., Northrup, J. M., McKenna, M. F., Anderson Jr, C. R., Angeloni, L., & Wittemyer,

G. (2015). Landscape and anthropogenic features influence the use of auditory vigilance

by mule deer. *Behavioral Ecology*, *26*(1), 75-82.

MacKenzie, D. I., Bailey, L. L., & Nichols, J. D. (2004). Investigating species co-occurrence

patterns when species are detected imperfectly. *Journal of Animal Ecology*, *73*(3), 546-

555.

Marion, S., Demšar, U., Davies, A. L., Stephens, P. A., Irvine, R. J., & Long, J. A. (2021). Red

deer exhibit spatial and temporal responses to hiking activity. Wildlife Biology,

2021(3). https://doi.org/10.2981/wlb.00853

Mech, D.L. 1970. The wolf: the ecology and behaviour of an endangered species. Natural

History Press for the American Museum of Natural History, New York.

Mech, L. D., & Peterson, R. O. (2003). Wolf-prey relationships. In L. D. Mech, & L. Boitani

(Eds.), Wolves: Behavior, ecology, and conservation (pp. 131–161). Chicago, IL:

University of Chicago Press.

Middleton, A. D., Kauffman, M. J., McWhirter, D. E., Jimenez, M. D., Cook, R. C., Cook, J.

G., ... & White, P. J. (2013). Linking anti-predator behaviour to prey demography

reveals limited risk effects of an actively hunting large carnivore. *Ecology letters*, *16*(8),

1023-1030.

Minta, S. C. (1992). Tests of spatial and temporal interaction among animals. *Ecological*

applications, *2*(2), 178-188.

Molinari-Jobin, A., Zimmermann, F., Ryser, A., Breitenmoser-Würsten, C., Capt, S.,

Breitenmoser, U., ... & Eyholzer, R. (2007). Variation in diet, prey selectivity and home-

range size of Eurasian lynx Lynx lynx in Switzerland. *Wildlife Biology*, *13*(4), 393-405.

Newsome, T. M., Boitani, L., Chapron, G., Ciucci, P., Dickman, C. R., Dellinger, J. A., ... &

Ripple, W. J. (2016). Food habits of the world's grey wolves. *Mammal Review*, *46*(4),

255-269.

Okarma, H., Jędrzejewska, B., Jędrzejewski, W., Krasiński, Z. A., & Miłkowski, L. (1995). The

roles of predation, snow cover, acorn crop, and man-related factors on ungulate

mortality in Białowieża Primeval Forest, Poland. *Acta Theriologica*, *40*(2), 197-217.

Okarma, H., Jedrzejewski, W., Schmidt, K., KowALCZYK, R., & Jedrzejewska, B. (1997).

Predation of Eurasian lynx on roe deer and red deer in Bialowieza Primeral Forest,

Poland. *Acta Theriologica*, *42*(2), 203-224.

Ordiz, A., Støen, O. G., Delibes, M., & Swenson, J. E. (2011). Predators or prey? Spatio-

temporal discrimination of human-derived risk by brown bears. *Oecologia*, *166*, 59-67.

Parsons, M. H., Apfelbach, R., Banks, P. B., Cameron, E. Z., Dickman, C. R., Frank, A. S., ...

& Blumstein, D. T. (2018). Biologically meaningful scents: a framework for

understanding predator–prey research across disciplines. *Biological Reviews*, *93*(1), 98-

114.

Peterson, R. O., and P. Ciucci. 2003. The wolf as a carnivore. Pages 104–130 in L. D. Mech

and L. Boitani editors. Wolves: behavior, ecology, and conservation. University of

Chicago Press, Chicago, Illinois, USA.

Pollock, L. J., Tingley, R., Morris, W. K., Golding, N., O'Hara, R. B., Parris, K. M., . . .

McCarthy, M. A. (2014). Understanding co-occurrence by modelling species

simultaneously with a joint species distribution model (JSDM). Methods in Ecology and

Evolution, 5, 397–406.

Prat-Guitart, M., Onorato, D. P., Hines, J. E., & Oli, M. K. (2020). Spatiotemporal pattern of

interactions between an apex predator and sympatric species. *Journal of Mammalogy*,

101(5), 1279-1288.

R Core Team (2023). R: A Language and Environment for Statistical Computing. R for

Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

Rafiq, K., Jordan, N. R., Wilson, A. M., McNutt, J. W., Hayward, M. W., Meloro, C., ... &

Golabek, K. A. (2020). Spatio-temporal factors impacting encounter occurrences

between leopards and other large African predators. *Journal of Zoology*, *310*(3), 191-

200.

Ramjith J, Bender A, Roes KCB, Jonker MA. Recurrent events analysis with piece-wise

exponential additive mixed models. 2022. Statistical Modelling., 2022

Ridout, M. S., & Linkie, M. (2009). Estimating overlap of daily activity patterns from camera

trap data. *Journal of Agricultural, Biological, and Environmental Statistics*, *14*(3), 322–

337. https://doi.org/10.1198/jabes.2009.08038

Sih, A. (1984). The behavioral response race between predator and prey. The American

Naturalist, 123(1), 143-150.

Sih, A., Crowley, P., McPeek, M., Petranka, J., & Strohmeier, K. (1985). Predation,

competition, and prey communities: a review of field experiments. *Annual review of*

ecology and systematics, *16*(1), 269-311.

Sih, A. (2005). Predator-prey space use as an emergent outcome of a behavioral response race.

Ecology of predator-prey interactions, *256*, 78.

Sims, D. W., Witt, M. J., Richardson, A. J., Southall, E. J., & Metcalfe, J. D. (2006). Encounter

success of free-ranging marine predator movements across a dynamic prey landscape.

Proceedings of the Royal Society B: Biological Sciences, *273*(1591), 1195-1201.

Smith, J. A., Donadio, E., Pauli, J. N., Sheriff, M. J., Bidder, O. R., & Middleton, A. D. (2019).

Habitat complexity mediates the predator–prey space race. *Ecology*, *100*(7), e02724.

Sunde, P., Böcker, F., Rauset, G. R., Kjellander, P., Chrenkova, M., Skovdal, T. M., ... &

Heurich, M. (2022). Mammal responses to predator scents across multiple study areas.

Ecosphere, 13(8), e4215.

Suraci, J. P., Smith, J. A., Chamaillé-Jammes, S., Gaynor, K. M., Jones, M., Luttbeg, B., ... &

Sih, A. (2022). Beyond spatial overlap: harnessing new technologies to resolve the

complexities of predator–prey interactions. *Oikos*, *2022*(8), e09004.

Tallian, A., Mattisson, J., Stenbacka, F., Neumann, W., Johansson, A., Støen, O. G., &

Kindberg, J. (2023). Proximity-sensors on GPS collars reveal fine-scale predator–prey

behavior during a predation event: A case study from Scandinavia. *Ecology and*

Evolution, *13*(12), e10750.

Tourani, M., Franke, F., Heurich, M., Henrich, M., Peterka, T., Ebert, C., ... & Peters, W.

(2023). Spatial variation in red deer density in a transboundary forest ecosystem.

Scientific Reports, *13*(1), 4561.

Travis, J. M. J., & Palmer, S. C. F. (2005). Spatial processes can determine the relationship

between prey encounter rate and prey density. *Biology Letters*, *1*(2), 136-138.

Wikenros, C., Aronsson, M., Liberg, O., Jarnemo, A., Hansson, J., Wallgren, M., ... &

Bergström, R. (2017). Fear or food–abundance of red fox in relation to occurrence of

lynx and wolf. *Scientific reports*, *7*(1), 9059.

Wood, S. N. (2017). Generalized additive models: an introduction with R. chapman and

hall/CRC.

Harz National Park -N = 60
- N_{events} = 7
- N_{events} = 9

Kilometer
0 2 4 6 8 10

Müritz National Park - N=70
- N_{events} = 70

Kilometer
0 2 3 5 6 8

Königsbrücker-Heide Wildnerness Area - N=55
- N_{events} = 181

Kilometer
0 1 2 3 4 5

Bavarian Forest National Park - N=100
- N_{events} = 8
- N_{events} = 32

Kilometer
0 4 8 12

Legend
- Locations of camera-traps

0 50 100 200 Kilometer