A Chaperonin Complex Regulates Organelle Proteostasis in Malaria Parasites

Amanda Tissawak[1,2], Yarden Rosin[1,2], Shirly Katz Galay[1,2], Alia Qasem[1,2], Michal

Shahar[1,2], Nirit Trabelsi[1], Ora Furman-Schueler[1], Steven M Johnson[3], Anat

Florentin[1,2][*].

[1] Department of Microbiology and Molecular Genetics, Faculty of Medicine, The

Hebrew University of Jerusalem, Jerusalem, Israel

[2] The Kuvin Center for the Study of Infectious and Tropical Diseases, Faculty of

Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel

[3] Department of Biochemistry and Molecular Biology, Indiana University School of

Medicine, Indianapolis, IN, USA

[*]**Corresponding Author:** Anat Florentin, +972-2-6757976 (phone), Email:

anat.florentin@mail.huji.ac.il

Preprint Server: BioRxiv

Keywords: malaria*, Plasmodium falciparum*, parasitology, apicoplast, proteostasis,

chaperonin, CPN, HSP60, Clp system.

Abstract

The apicoplast of *Plasmodium* parasites serves as a metabolic hub that synthesize essential biomolecules. Like other endosymbiotic organelles, 90% of the apicoplast proteome is encoded by the cell nucleus and transported to the organelle. Evidence suggests that the apicoplast has minimal control over the synthesis of its proteome and therefore it is unclear how organelle proteostasis is regulated. Here, we identified and investigated a large and conserved chaperonin (CPN) complex with a previously unknown function. Using genetic tools, we demonstrated that ablation of the apicoplast CPN60 subunit leads to parasite death due to organellar damage, immediately within its first replication cycle, deviating from the delayed death phenotype commonly observed for apicoplast translation inhibitors. Unlike its close orthologues in other prokaryotic and eukaryotic cells, CPN60 is not upregulated during heat shock (HS) and does not affect HS response in the parasite. Instead, we found that it is directly involved in proteostasis through interaction with the Clp (*caseinolytic protease*) proteolytic complex. We showed that CPN60 physically binds both the active and inactive forms of the Clp complex, and manipulates its stability. A computational structural model of a possible interaction between these two large complexes suggests a stable interface. Finally, we screened a panel of inhibitors for the bacterial CPN60 orthologue GroEL, to test the potential of chaperonin inhibition as antimalarial. These inhibitors demonstrated an anti-*Plasmodium* activity that was not restricted to apicoplast function, with additional targets outsides of this organelle. Taken together, this work reveals how balanced activities of proteolysis and refolding safeguard the apicoplast proteome, and is essential for organelle biogenesis.

Author Summary The cell of the human malaria parasite *Plasmodium falciparum* has a unique organelle called the apicoplast that produces essential metabolites, but it is unclear how it maintains a stable proteome. Here, we address the question of organelle proteostasis by investigating the function of a large chaperonin complex and its main subunit CPN60. We show that CPN60 mutants die due to organellar damage immediately within the first replication cycle, avoiding the typical apicoplast-delayed cell death. We demonstrate that it binds and stabilizes another large proteolytic complex and use computational predicting tools to demonstrate how a stable interface is attained. We use bacterial inhibitors to explore their potential as an antimalarial drug target. This study reveals how balanced refolding and proteolysis

safeguard the apicoplast proteome and opens a new avenue for antimalarial drug
discovery.

Introduction

The malaria parasite, *Plasmodium falciparum*, remains one of the deadliest human
pathogens, causing 600,000 deaths yearly worldwide[1]. Limited vaccines efficacies
and spreading of drug resistance are major global concerns. Despite years of
research, many aspects of the parasite's basic cell biology remain obscure,
hindering rational drug design and improved therapeutic interventions. Like most
other related parasites from the phylum of *Apicomplexa*, this single-celled eukaryotic
organism contains a non-photosynthetic plastid called apicoplast[2,3]. This unique
organelle evolved via a two-step endosymbiosis; in the primary endosymbiotic event,
a cyanobacterium was incorporated into a eukaryotic cell to form the modern
chloroplast. During the secondary endosymbiotic event, a photosynthetic red alga
was further taken up by a protist and led to the formation of a secondary plastid[4].
Although not photosynthetic, the apicoplast harbors essential prokaryotic metabolic
pathways that are essential to the parasite throughout its complex life cycle,
including fatty acid, Fe-S clusters, and isoprenoids biosynthesis[5]. The *Plasmodium*
apicoplast shares molecular features with prokaryotes, plants and parasites, and
therefore encompasses multiple parasite-specific drug targets. Indeed, drugs that
target apicoplast biology are in clinical use[6]. Most of those drugs (e.g., doxycycline,
clindamycin) target the prokaryotic protein synthesis machinery in the apicoplast[6,7].
However, less than 10% of the apicoplast proteome is encoded by its own genome.
The vast majority of the hundreds of apicoplast proteins are encoded in the nuclear
genome and are transported to the organelle via the secretory pathway[8,9]. Available
transcriptomic data suggest that the apicoplast does not control the expression of
these nuclear-encoded proteins[10]. In fact, parasites without an apicoplast continue to
express and accumulate apicoplast proteins in vesicle-like structures in the
cytoplasm[11-14]. Due to the inability of the apicoplast to control its own protein
synthesis, it is unclear how the apicoplast maintains a stable proteome.

Previously, we suggested that protein degradation plays a central role in organelle
proteostasis and showed that apicoplast-targeted Clp (*caseinolytic protease*)
proteins form an essential proteolytic complex in the organelle[14,15]. The conserved
Clp family consists of proteases and chaperones that form multi-subunit proteolytic

complexes, with complex composition varying widely between different species and
organelles[16,17]. In bacteria and plant chloroplasts, the ClpP proteases typically
associate with Clp ATPase chaperones that unfold and feed substrates into the ClpP
barrel-like cavity for degradation[18–20]. However, the putative *Plasmodium* Clp
proteins differ significantly from their bacterial orthologs and their organellar,
molecular and enzymatic functions are unclear[21,22]. Moreover, despite detailed
genetic and molecular data illuminating complex composition, no regulator of the
apicoplast Clp complex has been identified.

Besides the Clp chaperones, the *Plasmodium* apicoplast is predicted to host other
molecular chaperones, among those, chaperonin 60. This 60 kDa Heat Shock
protein, often termed HSP60 or CPN60, is highly conserved and found in nearly all
organisms[23]. Its bacterial orthologue, GroEL, is involved in maintaining proteostasis
under different stressors[24], and in most eukaryotic cells, *Plasmodium* included, a
mitochondrial orthologue is required for organellar protein import[25]. In apicomplexan
parasites, a CPN60 homolog was reported to localize to the apicoplast[15,26], although
its functions are unknown.

Here, we report the molecular investigation of a non-mitochondrial CPN60
orthologue in the malaria parasite *P. falciparum*. We show that it localizes to the
apicoplast and is primarily expressed during the late stages of the parasite's life
cycle, when organelle biogenesis occurs. We found that CPN60 function is essential
for parasites viability, but unlike its orthologs in most organisms, it is not involved in
heat shock response. Instead, it is required for apicoplast functions and interacts
with the proteolytic Clp complex. We further demonstrate that it binds the proteolytic
Clp subunit and this interaction is required for Clp complex stability. Lastly, we
screened a panel of inhibitors of the bacterial HSP60 orthologue GroEL in order to
test the potential of chaperonin inhibition as antimalarial. Taken together, our studies
shed light on the activity of the apicoplast CPN60 and its central role in maintaining
organelle proteostasis and parasite's viability.

Results

***Plasmodium falciparum* chaperonin 60 is an apicoplast-localized protein**
Our previous work suggested that the apicoplast employs caseinolytic protease
(Clp)-mediated degradation to maintain organellar functions[14,15]. In order to further

investigate proteostasis regulation in the apicoplast, we performed an affinity screen,

immunoprecipitating several different Clp subunits coupled with mass spectrometry

analysis[15]. Notably, we found a conserved apicoplast-resident heat shock protein

(HSP60) homolog, termed chaperonin 60 (CPN60, PF3D7_1232100). The putative

apicoplast CPN60 in *P. falciparum* is 718 amino acids in length, while the

mitochondrial HSP60 (PF3D7_1015600) is 580 amino acids long, and the two

isoforms share 30% sequence identity (46% similarity) (Sup. Fig. 1). To test whether

and how CPN60 is involved in proteostasis and other apicoplast functions, we

employed a genetic approach using the tetR-Dozi system, a *Plasmodium*-specific

method[27] (Fig. 1A). We tagged the gene with a V5 tag followed by the tetR-Dozi

cassette. Using CRISPR/Cas9, we incorporated the regulatory cassette at the 3′ end

of the CPN60 gene, creating the CPN60$^{V5\text{-}apt}$ parasite line. Genotyping of

independently isolated clones was performed by PCR analysis, and showed correct

integration into the *cpn60* locus (Fig. 1B). Western blot analysis revealed expression

of endogenous V5-tagged CPN60 at the predicted 85 kDa size (Fig. 1C). The

presence of a double band suggested that CPN60 contains an apicoplast-targeting

transit peptide that is cleaved upon its organellar localization. To verify this, we

treated CPN60$^{V5\text{-}apt}$ parasites with the antibiotic chloramphenicol (CHL), which

specifically inhibits apicoplast protein translation and leads to organelle loss[6]. 96-h

post CHL treatment, we observed accumulation of CPN60 largest band and

disappearance of its smaller form (Fig. 1C). This indicates that CPN60 has a transit

peptide that cannot be cleaved off when organelle functions are compromised.

Homologs of the HSP60 family are often considered house keeping genes, retaining

constant expression levels throughout the cell cycle[23]. However, protein extraction

from synchronized parasites revealed that CPN60 levels were relatively low at the

first half of the 48-h asexual cycle, and increased only at the later trophozoite stage,

36 hours post-invasion (Fig. 1D, see protein levels and corresponding blood

smears). This was also manifested with increased CPN60 mRNA levels, as

measured by quantitative real-time PCR (Fig. 1E). Quantification of CPN60 protein

levels shows that expression peaks around the time that schizogony (nuclear

replication), and apicoplast development begin[28] (Fig. 1F).

We further monitored CPN60 expression and localization in different developmental

stages using immunofluorescence assay (IFA). Recently, we showed that apicoplast

development is tightly coordinated with nuclear replication and the cell cycle[29]. In

accordance with these observations, we saw CPN60 at an early stage in a dot-like

structure, in a typical elongated apicoplast morphology during the late trophozoite

stage, a branched structure during schizogony and divided daughter organelles post

segmentation, in the last hour before egress (Fig. 2A). We took advantage of

ultrastructure expansion microscopy to verify the details of these structures and

observed CPN60 in a consistent and uniform organelle localization (Fig. 2B). Lastly,

we combined this IFA with a mitotracker staining to demonstrate that the intricate

structure is distinct from the parasite's mitochondrion and therefore marks the

apicoplast (Fig. 2C).

CPN60 is essential for parasite's survival due to its function in apicoplast

biogenesis

To look into CPN60 cellular function, we activated the tet-aptamer system, to induce

translational repression of CPN60 (Fig. 3A). We induced protein knockdown by

removal of the stabilizing ligand anhydrotetracycline (aTC), effectively reducing

CPN60 protein levels within 24 hours, dropping below detection after 72 hours (Fig.

3B). The consequences of CPN60 knockdown were parasite's death, indicating that

its activity is essential for parasite's viability (Fig. 3C). It was shown that apicoplast-

related cytotoxicity can be reversed by addition of isopentenyl pyrophosphate (IPP),

an essential apicoplast-producing metabolite[11]. We observed that media

supplementation with IPP completely restored mutants' growth, indicating that

CPN60 essential activity is restricted to apicoplast function (Fig. 3D). Delayed cell

death due to apicoplast-associated damage is observed in parasites treated with

doxycycline or chloramphenicol. Such parasites complete their first 48 hours cell

cycle, reinvade new cells and die only at the end of the 2^{nd} cycle after 96 hours of

treatment[30]. We therefore synchronized the cultures and removed aTC from early

rings stage CPN60^{V5-apt} parasites and observed their intraerythrocytic development

(Fig. 3E). During the first cycle, knockdown parasites developed like +aTC control

until the trophozoite stage, but then could not process into normal schizogony (Fig.

3F). The few persisting parasites in the culture could not egress or reinvade and

failed to form new ring-stage parasites in the 2^{nd} cycle (Fig. 3G). For comparison, we

induced knockdown in a different apicoplast-resident protein, PfClpS, which we

previously showed to be required for organelle biogenesis[15]. Induction of PfClpS

knockdown using the same tet-aptamer system under the same experimental
settings resulted in a delayed cell death, with parasites death observed 5 days post
aTC removal (Sup. Fig. 2). This reinforces the observation that, unlike many other
apicoplast proteins, CPN60 ablation results in an apicoplast-immediate cell death.

We then tested whether essentiality of CPN60 is limited to supporting an apicoplast
metabolic pathway, or it is also required for the physical biogenesis of the organelle.
If the former is correct, then IPP supplementation during CPN60 knockdown should
be sufficient to retain an intact organelle, albeit dysfunctional. However, if the latter is
true and CPN60 is required for apicoplast biogenesis, its knockdown will result in
organelle loss, regardless of IPP supplementation. To distinguish between these two
possibilities, we grew CPN60$^{V5\text{-apt}}$ parasites for 17 days without aTC while
supplementing the media with IPP, and then added back aTC to restore CPN60
expression (Fig. 3H). We observed that this treatment resulted in parasites growth
that is completely dependent on IPP supplementation, regardless of aTC presence
(i.e. CPN60 expression) (Fig. 3H). Moreover, IFA showed that CPN60 in treated
parasites lost its distinct apicoplast elongated structure and appeared instead in
vesicle-like cytoplasmic puncta (Fig. 3I). Moreover, processing of CPN60 was lost
and it migrated on the gel at the size of its cytoplasmic form (Fig. 3J). Collectively,
these results demonstrate that CPN60 activity is required for apicoplast biogenesis,
and during knockdown the organelle is lost and cannot be retrieved even when
CPN60 expression is restored.

***Plasmodium* CPN60 is not involved in heat shock response**
In most organisms, HSP60 family members are involved in stress pathways,
particularly heat shock (HS) response. During infection, the malaria parasite faces
periodic fevers, making HS one of its main stressors. To investigate the function of
apicoplast CPN60 during HS, we measured its expression levels following 24 hours
incubation at 40^{o}C. *Wildtype* parasites subjected to HS, grew slower and exhibited
morphological defects (Fig. 4A). However, this response was not accompanied by
elevated CPN60 protein levels (Fig. 4B). We also followed potential changes in
transcript levels following HS, and as a positive control used leucine-rich repeat
protein (LRR5), a *Plasmodium* gene that is nonessential under ideal growing
conditions, however was found to be associated with HS response in the

parasite[31,32]. Here too, we found that while LRR5 transcripts were elevated by two-
and three-folds post HS, there was no effect on CPN60 transcription (Fig. 4C).

Next, we sought to analyze the effect of CPN60 depletion on parasite's survival
following HS. Because complete knockdown of CPN60 leads to parasite's death, we
were interested in decreasing CPN60 levels to the point where subtle effects could
be tested. To do that, we performed a half-maximal effective concentration (EC50)
assay for aTC, and found that 4.5 nM support 50% survival under normal
temperature as well as following incubation at 40°C (Fig. 4D). Next, we measured
CPN60 protein levels while growing the parasite's with 2X EC50 dose (8 nM aTC)
compared to its levels in non-induced parasites which are cultured at 100X EC50
(500 nM aTC, Fig. 4E). Quantification revealed that at this aTC concentration,
CPN60 cellular levels were decreased by 50% (Fig. 4E). Next, we measured
parasite's growth at different aTC concentrations following 24 hours incubation at
40°C. While *wildtype* parasites recover after heat exposure, their growth is slightly
attenuated (Fig. 4F, blue curve). Importantly, parasites with reduced CPN60 levels
(16, 8, 4 nM which correspond to purple, green and red curves, respectively in Fig.
4F) exhibited compromised growth, in an aTC dose-dependent manner. However,
exposure to heat did not alter their relative growth rate compared to the effect that it
had on *wildtype* parasites (Fig. 4F and Sup. Fig. 2B). These data suggest that unlike
canonical CPN60 orthologues, the *Plasmodium* CPN60 is probably not involved in
HS response and possesses an apicoplast-specific function.

Co-expression and co-localization of CPN60 and PfClpP, the active protease of
the Clp complex
We previously identified CPN60 in a mass spectrometry screen as a potential
interactor of the Clp complex[15]. To look deeper into this potential interaction, we
expressed a tagged copy of the Clp protease subunit, PfClpP. PfClpP is a highly
potent protease and in order to avoid adverse effects, we used a transgenic PfClpP
with a point mutation in its proteolytic active site (S264A), rendering it catalytically
inactive[15]. Furthermore, we designed integration into the parasite's genome under
the endogenous *hsp110* promoter, which enables constant but moderate levels of
expression[15]. This construct was transfected on the background of the CPN60
conditional mutants, generating the CPN60[V5-apt]; ClpP[DEAD-Ty] parasite line (Fig. 5A).
Clones were isolated and integration at the *hsp110* locus was verified using

diagnostic PCR (Sup. Fig.3A). Western blot demonstrated the co-expression of the

endogenous CPN60^{V5} at 85 KDa, as well as the transgenic ClpP$^{DEAD-Ty}$ (Fig. 5B).

The expected processing pattern of PfClpP is consistent with a model of a full-length

PfClpP, presumably in transit through the secretory pathway (fraction III), an

apicoplast-localized PfClpP (fraction II), and a mature PfClpP after proteolytic

removal of an inhibitory pro-domain (fraction I) (Fig. 5B). We verified that expression

of ClpP$^{DEAD-Ty}$ had no effect on parasite's growth or CPN60 function, as can be seen

by a consistent EC50 at 37^{o}C or following HS (Fig. 5C). Immunofluorescent

microscopy, revealed that CPN60 co-localizes with PfClpP in the apicoplast during

all cell cycle stages (Fig. 5D). Therefore, we concluded that ectopic expression of

ClpP$^{DEAD-Ty}$ does not affect its processing or overall apicoplast morphology.

Furthermore, it colocalizes with CPN60 without affecting its activity, indicating that

the co-expression could be used to evaluate interaction between the two.

CPN60 binds and stabilizes the Clp complex

To test for physical interactions between CPN60 and PfClpP, we performed

immunoprecipitation using the double-mutant parasites. The pulldown of ClpP$^{DEAD-Ty}$

resulted in the co-IP of CPN60, in the form of a weak band (Fig. 5E, right panel). The

reciprocal IP of CPN60 revealed that it interacts with the mature ClpP protease (I),

as well as with the zymogen form (II) but not with the full length PfClpP fraction (III)

(Fig. 5E, left panel). Importantly, pulldown of CPN60^{V5} was specific to PfClpP as it

failed to co-IP unrelated highly-expressed proteins (Sup. Fig. 3B), suggesting a

specific interaction with ClpP however weak or transient. This indicates that CPN60

and PfClpP interact only upon co-localization to the apicoplast and that the

chaperone binds the Clp complex both before and after protease activation. This

might be a direct interaction, or indirect through a third unknown partner.

Typically, Clp proteolytic subunits oligomerize into heptameric rings, consisting of the

core of the Clp complex[17]. Likewise, CPN60 orthologues are part of large heptameric

or tetradecameric chaperonin complexes[23]. Therefore, the interaction observed here

is unlikely to occur between individual monomers, rather at an interface between

large complexes. To better understand this interaction, we utilized previously solved

structures of CPN60 (PDB ID 7K3Z)[33] and PfClpP (PDB ID 2F6I)[22] and set to predict

how these two large complexes might interact in their 3D conformations (Sup. Fig.

3C&D). We used AlphaFold3 (AF3)[34], a state-of-the-art model with a diffusion-based

architecture capable of handling large protein complexes, offering improved

accuracy and speed compared to AlphaFold2[35]. The AF3 predictions provided a

model in which the two rings are stacked on each other, whereby the individual

monomers tilt in opposite directions. Each monomer of the PfClpP heptameric ring

makes contacts with two monomers in CPN60, and each CPN60 monomer contacts

two monomers in PfClpP (Sup. Fig. 3E). In this configuration it is possible for both

CPN60 and ClpP complexes to form tetradecameric barrels, which interact via a

stable interface.

We next examined the significance of this interaction, and tested possible effects on

PfClpP activity. We previously showed that PfClpP accumulates in the apicoplast as

an inactive zymogen, and upon assembly of the PfClpP oligomer, it is auto-cleaved

to remove the inhibitory pro-domain and becomes activated[15]. Thus, the ratio

between the inactive (band II) and active (band I) PfClpP forms is indicative of its

activation state. To test the processing pattern of PfClpP upon CPN60 knockdown,

we removed aTC from non-synchronous parasites and observed PfClpP processing

24 hours later, before parasites begin to die. We detected a reduction in the levels of

PfClpP active band (I) and accumulation of the inactive zymogen (II), once CPN60

levels drop (Fig. 5F). This shift in the ratio of active: inactive PfClpP could be

quantified (Fig. 5G), suggesting that the interaction between CPN60 and PfClpP is

essential for PfClpP activation and complex stability.

Exploiting the apicoplast chaperonin system as an antimalarial target

Inhibition of protein synthesis in the apicoplast is a clinically approved drug target, as

indicated by the antimalarial effect of doxycycline. Nevertheless, such treatment

directly affects only a fraction of the total apicoplast proteome. Although largely

unexplored, post-translational regulation in the organelle is no less critical and

likewise can prove druggable. In this regard, the essentiality of CPN60 described

here supports its viability as a potential antimalarial target.

To test this, we took advantage of a series of bacterial inhibitors designed to target the

prokaryotic CPN60 orthologue, GroEL. As far as we are aware, chaperonin inhibitors

have never been explored against *Plasmodium*. Thus, we conducted a pilot study to

test the efficacy of a panel of phenylbenzoxazole (PBZ) analogues. PBZ compounds

were previously shown to bind and inhibit the activity of GroEL in *E.* Coli, inducing a

separation and a twist of the two rings that decoupled their inter-ring allosteric

signaling, thereby blocking the folding cycle[36]. Detailed cryoEM structures showed that

the PBZ analogue PBZ1587 binds at the GroEL ring-ring interface[37]. Moreover, it was

shown that mutations of key residues within the interface binding site render GroEL

resistance both *in vitro* and in *E. coli*, demonstrating on-target antibacterial effects and

a validated tool for other studies. We therefore hypothesized that PBZ analogues have

the potential to bind the *Plasmodium* chaperonin systems, either the apicoplast

CPN60, the mitochondrial HSP60, or both. Sequence alignments indicated that *E. coli*

GroEL, human mtHSP60, and *P. falciparum* mitochondrial HSP60 shared high

conservation of their overall sequences and the PBZ1587 ring-ring interface binding

sites (Sup Fig. 1 and Sup Fig. 4), but apicoplast CPN60 was much less conserved.

However, upon generating homology models of apicoplast CPN60 and mitochondrial

HSP60 bound with PBZ1587 at the ring-ring interface as it does with GroEL, we were

encouraged to see that, despite the differences in binding site residues, many key

ligand-binding site interactions were predicted (Sup Fig. 5 and Sup Fig. 6), suggesting

the inhibitors could still be effective in *P. falciparum*. Furthermore, the apicoplast

CPN60 isoform contains an extended alpha-helix at the ring-ring interface near the

PBZ1587 binding site, a feature we hoped might provide selective targeting as the site

is more well defined compared with mitochondrial HSP60.

To test the antimalarial potential of the PBZ inhibitors, we incubated parasites with

serial dilutions of each (50 μM to 50 nM) and monitored growth after 72 or 96 hours

to calculate the inhibitor EC50 (Fig. 6A and Table 1). Although some inhibitors

showed moderate activity against *P. falciparum* growth (1134, 1135, 1259), others

produced an EC50 at a single-digit micromolar range (1001, 1064, 1133, 1207,

1221) or lower (1587), demonstrating a potent anti-*Plasmodium* activity (Fig. 6A and

Table 1). We then incubated synchronized ring-stage parasites with a selected panel

of the more potent inhibitors and followed their intraerythrocytic replication. During

the first cycle, treated parasites developed normally like untreated control until the

trophozoite stage, but then could not proceed to normal schizogony, and failed to

complete their cycle (Fig. 6B). The kinetics of this developmental failure was

comparable to the one observed upon CPN60 knockdown (Fig. 3G). We therefore

tested whether inhibitors cytotoxicity is associated with impaired apicoplast

development. We incubated non-synched CPN60^{V5-apt} parasites with PBZ inhibitors,

and fixed the cells after 24 hours, before parasites begin to die. Strikingly, IFA

demonstrated that PBZ treatments led to aberrant apicoplast morphologies, including

fragmented, shrunk, or undeveloped organelles (Fig. 6C). These results suggest at

least one apicoplast-associated target, with CPN60 being a likely candidate.

We tried to test this directly, and performed EC50 assays for the PBZ compounds

during CPN60 partial knockdown. This, however, did not lead to a significant EC50

shift for the compounds, hindering definitive conclusions (Sup. Fig. 7). We therefore

assessed specificity towards broader apicoplast function, by coincubation of PBZ

compounds with IPP. We incubated the parasites with a fixed inhibitor concentration

of 12.5 µM (roughly 2-3X of most EC50s), and monitored their growth over 96 hours.

While IPP did not affect growth of untreated parasites, it fully restores viability of

chloramphenicol-treated parasites, which inhibit apicoplast translation (Fig. 7A).

However, the toxic effect of the PBZ inhibitors could not be restored by IPP,

suggesting that (1) the 12.5 µM test concentration may have been too high above

inhibitor EC50 values, impeding rescue by IPP supplementation (i.e. PBZ1587), or

(2) inhibitors could be functioning against additional targets, the most likely candidate

being mitochondrial CPN60. However, the moderately potent inhibitor PBZ1214

(EC50 = 9.6-12.5 µM) presented an intriguing result as IPP treatment sensitized

parasites to the inhibitor (Fig. 7B & C). The reason(s) for IPP sensitization of

PBZ1214 is unclear at present and will require further investigation.

Discussion

Chaperonins (CPN60/ HSP60) function biochemically as ATP-dependent molecular

chaperones, working in concert with other chaperones and co-chaperones to

facilitate protein folding and maintain proteostasis. Chaperonins are originally

prokaryotic proteins and are essential in almost all bacteria, where they are known

as GroEL, as well as in all endosymbiotic organelles[23]. Indeed, an HSP60 homolog

is found in the mitochondria of almost all eukaryotes, including human and

Plasmodium, although the latter has not been well characterized. Another GroEL

homolog is found in chloroplasts[38] (an essential Rubisco binding subunit) as well as

in related non-photosynthetic plastids, including the apicoplast organelle of

Apicomplexan parasites. As we show here, the apicoplast resident CPN60, is quite

distinct from the *Plasmodium* mitochondrial Hsp60 which in turn shares a higher

degree of conservation with human mtHSP60 and *E. Coli* GroEL. Prior works in

Plasmodium and in *Toxoplasma gondii* reported on the expression of an apicoplast

CPN60[26,39]. However, so far, the cellular functions of the apicoplast CPN60 have not

been resolved.

In this work, we found that CPN60 expression profile correlates with the timing of

apicoplast elongation, branching and division[29,40], which supports its newly-observed

function in organelle biogenesis. Interestingly, CPN60 knockdown induced

immediate cell-death within the first replication cycle. This phenotype is different and

faster than the more typical apicoplast-delayed cell death, observed in parasites

treated with apicoplast-targeting antibiotics[30]. This kinetics is more similar to the

effect of drugs blocking isoprenoids synthesis, which also target the apicoplast but

kill within one replication cycle[11,41]. Thus, CPN60 knockdown adds to growing

evidence of immediate cell death resulting from inhibition of apicoplast biogenesis[42−

45], underlying its potential as a drug target.

In bacteria and plant chloroplasts, CPN60 is critical for heat shock (HS) response. In

the case of *Plasmodium*, the parasite copes and responds to periodic fever episodes

during infection of the human host[46,47]. Since the ER has a limited repertoire of

molecular chaperones[48], it would have made sense if the apicoplast chaperonin

complex complements the cellular stress response. Indeed, a recent large-scale

forward-genetic screen suggested a link between the apicoplast and fever-survival[31].

However, we found that during recovery after HS, CPN60 levels remained constant,

and that perturbating CPN60 levels did not affect HS response, most likely

precluding a role for CPN60 in this stress.

In the apicoplast of *P. falciparum*, Clp proteins form an organellar degradation

machinery that is independent of the cytoplasmic proteasome[15,22]. In previous

studies[14,15], we showed that a tightly synchronized sequence of events determines

the activity of the Clp complex. In the current work, we found that CPN60 interacts

with both the inactive and active PfClpP protease, affecting its activation turnover.

While these co-IP experiments demonstrate interaction in the native cellular

environment, they cannot reveal whether the binding is direct or rather mediated

through a third partner. Similar to *Plasmodium*, the chloroplast Clp system is the

primary machinery for protein degradation in the plastid stroma[19]. A recent cryo-EM

study describing the structure of the chloroplast Clp complex, found that CPN20 co-

chaperonin, a co-factor of CPN60, forms a cap on the top of ClpP and is shown to

repress ClpP proteolytic activity in vitro[49]. In *Plasmodium*, there are three potential

CPN20-like genes that are predicted to localize to the apicoplast. However, none of

the structural predictions that we tried could fit any of them into a stable interaction

model with either PfClpP or CPN60. Thus, future experimental work will be needed

to understand the nature of the Clp-CPN complex interaction and whether it is

mediated by co-chaperonins or other factors.

In the absence of such information, we used our IP data and previous structural

works to try and predict how a direct interaction between CPN and Clp complexes

might look. A previous study solved the crystal structure of *Plasmodium* CPN60[33].

Although in this study it was annotated as the *Plasmodium* mitochondrial HSP60,

careful examination of the sequence used indicates that it is rather the structure of

the apicoplast CPN60 (PF3D7_1232100). As seen in its solved structure, the basic

oligomeric state of CPN60 is a single heptameric ring, though a double-ring

tetradecameric structure is also possible (PDB ID 7K3Z, Sup. Fig. 3C). Similarly, a

solved X-ray structure of PfClpP demonstrates that it forms a tetradecamer,

consisting of two heptameric rings (PDB ID 2F6I, Sup. Fig. 3D)[22]. Our AF3

predictions provided a model in which the PfClpP heptameric ring tilts one way,

whereas the CPN60 ring tilts the other way (Sup. Fig. 3E). In this configuration it is

possible for both CPN60 and ClpP complexes to form tetradecameric barrels, which

interact via a stable interface (See our working model in Fig. 8A). The stability of this

interaction can be assessed by computational alanine scanning that identifies

potential interaction hotspots. This sort of analysis reveals significant hotspots in key

residues in both complexes; The ClpP predicted hotspots residue are Y41, I45 and

Y73 and the CPN60 hotspots residues are K243, F320 and N323 (Fig. 8B).

Additional experimental validation is required to confirm these interaction models and

to refine our understanding of the molecular mechanisms driving the CPN60-ClpP

complex formation. Interestingly, we discovered that upon CPN60 removal, active

PfClpP levels decline, while PfClpP zymogen levels increase (Fig. 8C). Until this

point, we operated under the hypothesis that, similar to chaperonin orthologs in other

organisms, the apicoplast CPN60 acts as a "conformational editor" to assist in

organellar folding processes. These unexpected results showed that at least part of

its activity is to stabilize and activate the Clp complex, thus supporting organellar

degradation processes. Based on these results, we propose a working model in

which the interdependence between two complexes with opposing functions,

facilitates a balanced proteostasis in the apicoplast (Fig. 8).

Over years of research, chaperonins have been identified as potential targets for the

development of novel agents affecting their activity. Many of those have implications

for new-generation antibiotics, cancer and neurodegeneration treatment, as well as

the development of diagnostic biomarkers[23,50–52]. Nevertheless, the potential of

mitochondrial and/or apicoplast CPN60 as viable targets for antimalarial

development have not been investigated. Here we tested the antimalarial activity of a

small panel of phenylbenzoxazole (PBZ) compounds that were originally identified as

inhibitors of the *E. coli* GroEL/ES chaperonin system[36]. These inhibitors exhibited

antibiotic effects from the low-μM to mid-nM range against a panel of Gram-positive

and Gram-negative bacteria[50], mycobacteria[53] and even *Trypanosoma* parasites[54],

highlighting the versatility of targeting chaperonins. Recently, Godek *et al.*

demonstrated PBZ analogues that functioned on-target against GroEL in *E. coli*[37].

Within the group of PBZ anti-GroEL inhibitors tested here, we found compounds with

antimalarial effect at the low and sub-micromolar range which affect apicoplast

biogenesis. However, their cytotoxic effects were not reversed with IPP

supplementation, suggesting additional targets beyond the apicoplast CPN60. This

lack of specificity could be explained by the presence of two other chaperonin

systems in the parasite in addition to the one in the apicoplast. One of them is the

mitochondrial HSP60, that like the apicoplast CPN60 belongs to the Type I

chaperonins which are common to all bacteria and endosymbiotic organelles in

humans, plants and parasites. Another is the type II chaperonin, which is found in

archaebacterial cytoplasm (thermosome) and eukaryotic cytosol (CCT/TRiC)[55].

Based on recognized and separated evolutionary lines, the TriC system is generally

poorly characterized comparing to the type I HSP60. Uncharacteristically, the

CCT/TRiC system in *Plasmodium* has been investigated more than its type I variant,

and shown to be a valid drug target[56–58]. Whether it is also targeted by PBZ

inhibitors, or can explain the IPP toxicity that is associated with some of them,

remains to be investigated.

To conclude, by investigating the relationship between the *Plasmodium* Chaperonin

and Clp complexes, we revealed two opposing mechanisms that interact to balance

proteostasis in the apicoplast. These detailed cellular, biochemical and bioinformatic

analyses shed light on how excessive degradation is avoided when folding is

compromised, to regain steady state. We further provide a foundation for systematic

evaluation of these mechanisms and the potential of targeting CPN60 as an

antimalarial strategy. Future studies focusing on their bacterial evolutionary origins

could use existing antibacterial compounds to streamline drug discovery.

Materials & Methods

Plasmids construction

Genomic DNA was isolated from *P. falciparum* using NucleoSpin blood kit (Machery-

Nagel). For the generation of the CPN60^{V5-apt}; conditional mutant, 500 bps from the

C-term and 3'UTR from the of *cpn60* gene (Pf3D7_1232100) were amplified from

genomic DNA using primers P1+P2 and P3+P4, respectively. The two products were

conjugated together using overlap extension PCR and inserted as one piece into

pMG74-tetR-Dozi vector[27]. For expression of a C-term CPN60 guide RNA, oligos

P8+P9 were annealed and inserted into pUF1-Cas9-guide. Primer P7 was used in

combination with primer P6 or P4 to test for accurate integration in the parasite

genome following transfections. Evaluating aptamer repeats number was done using

primers P5+P6.

For the generation of ClpP^{DEAD} parasites, we transfected parasites with the puc-

HSP110-ClpP$^{DEAD-Ty}$ plasmid which we used previously to study ClpP processing

and activation[15]. It encodes the entire Open Reading Frame of ClpP

(PF3D7_0307400), mutated in three residues: Ser264Ala, Glu308Arg and

Arg285Glu, following by a triple Ty tag. Expression is driven under the *Pfhsp110c*

gene (PF3D7_0708800). Briefly, the HSP110 expression system comprises of a

repair plasmid that contains homology sequences from the *Pfhsp110c* gene. The

repair sequences include the last 429 bp (not including the stop codon) from the

Pfhsp110c gene, followed by a 2A skip peptide, ClpP$^{DEAD-Ty}$ sequence and the first

400 bps from *Pfhsp110c* 3'UTR. For expression of a *Pfhsp110c* guide RNA, oligos

P10+P11 are expressed from pUF1-Cas9-guide. Primers P12 and P13 were used to

test for accurate integration in the parasite following transfections.

All constructs utilized in this study were confirmed by Sanger sequencing. PCR

products were inserted into the respective plasmids using In-Fusion Snap assembly

Master mix (Takara). All restriction enzymes used in this study were purchased from

New England Biolabs. Oligonucleotides used in this study are summarized in Table

2.

Cell culture and transfections

Parasites were cultured in RPMI medium supplemented with Albumax I (Gibco) and

transfected as described earlier[59–61]. Synchronization of cultures were achieved by

intermittent sorbitol and Percoll treatments. To generate CPN60^{V5-apt} parasites, a mix

of two plasmids was transfected into NF54 *wildtype* parasites in the presence of 0.5

mM anhydrotetracycline (aTC, Cayman Chemicals). The plasmid mix contained 40

µg of pUF1-Cas9-guide expressing the CPN60 guide, and 20 µg of pMG74-based

donor plasmid with CPN60 homology regions, linearized by EcoRV. Drug pressure

was applied 48 hours post transfection using 2.5 µg/ml Blasticidin (BSD, Sigma). For

each transfection, at least 2 clones (A4 and F11) were isolated via limiting dilutions

and used in all subsequent experiments.

To generate CPN60$^{V5\text{-}apt}$; ClpP^{DEAD} parasites, a mix of two plasmids was transfected

into CPN60$^{V5\text{-}apt}$ parasites. The mix contained 2 plasmids; 40 µg of pUF1-Cas9-

Pfhsp110c-guide and 40 µg of the marker-less repair plasmid puc57-HSP110-

ClpP^{DEAD}. Drug pressure was applied 48 hours post transfection, using 1 µM DSM1

(BEI Resources), selecting for Cas9 expression. DSM1 was removed from the

culturing media once parasites clones were isolated by limiting dilution. For each

transfection, at least 2 clones (D12 and H10) were isolated via limiting dilutions and

used in all subsequent experiments.

Growth assays

For CPN60 knockdown assays, asynchronous parasite cultures were washed 8

times and incubated without aTC. Throughout the course of the experiment,

parasites were sub-cultured to maintain parasitemia between 1-5% and parasitemia

was monitored every 24 hours via flow cytometry. Growth assay during PfClpS

knockdown was performed following the same steps. The parasitemia levels were

determined in comparison to negative control (uninfected RBCs). Cumulative

parasitemia at each time point was back calculated based on actual parasitemia

multiplied by the relevant dilution factors. Parasitemia in the presence of aTC at the

end of each experiment was set as the highest relative parasitemia and was used to

normalize parasites growth. Normalized data from technical replicates were analyzed

and fit to an exponential growth curve using Prism (GraphPad Software, Inc.) All

experiments shown are representative of at least three biological replicates. For IPP

rescue, media was supplemented with 200 µM IPP (Isoprenoids LC). Controls for

apicoplast ablation were done using 30 µM chloramphenicol. For apicoplast ablation

during CPN60 knockdown, parasites were incubated without aTC for 17 days while

being supplemented with IPP. After these 17 days, this culture was divided into two

groups, one with aTC (to allow CPN60 re-expression and accumulation) and one

without aTC. Both groups were supplemented with IPP. After additional 16 days,

each group was further divided into two cultures, with or without IPP. The growth of

the four treated groups were then measured by flow cytometry, as well as taken for

IFA and WB analysis.

Flow cytometry

551   Aliquots of parasite cultures (5 μl) were stained with 8 μM Hoechst (ThermoFisher

552   Scientific) in PBS. The fluorescence profiles of infected erythrocytes were measured

553   and analyzed by flow cytometry on a CytoFlex S (Beckman Coulter). The

554   parasitemia data were analyzed and presented using Prism (GraphPad Software,

555   Inc.).

Recovery after heat shock

To induce heat shock response, parasites were incubated at 40^{o}C for 24 hours and

then allowed to recover at 37^{o}C to mimic the characteristic malarial fever episodes.

Growth of recovered parasites was monitored over the next following days using flow

cytometry and protein samples were isolated as described below. When CPN60

levels were to be manipulated, aTC was first washed away, and parasites were

subjected to heat shock 24 hours later to allow protein levels to drop.

Half-maximal effective concentration (EC50)

To generate an EC50 curve for aTC, asynchronous cultures of CPN60$^{V5\text{-}apt}$ or

CPN60$^{V5\text{-}apt}$; ClpP$^{DEAD\text{-}Ty}$ parasites were washed 8 times and then seeded at 0.5%

parasitemia in a 96 well plate with varying concentrations of aTC. To measure aTC

EC50 after recovery from heat shock, parasites were similarly treated, incubated for

24 hours at 40^{o}C, and then returned to 37^{o}C for the rest of the experiment. To

generate EC50 curves for HSP60 inhibitors, NF54 *wildtype* parasites were seeded at

0.5% parasitemia in a 96 well plate with varying concentrations of each compound.

In all cases parasitemia was measured after 72- and 96-hours using flow cytometry.

Data were fit to a dose-response equation using Prism (GraphPad Software, Inc.).

Inhouse anti-Ty tag monoclonal antibody purification

The hybridoma BB2 clone cell line (a kind gift from Boris Striepen) was adapted to

serum-free medium. 500 ml spent medium of the hybridoma clone containing mAbs

was collected, concentrated and purified by HiTrap Protein G HP antibody

purification 1 ml column (Cytiva) operated by peristaltic pump, as per manufacturer

protocol. The bound antibody was eluted using 10 ml 0.1 M Glycine-HCl buffer, pH

2.7 for ten minutes, and then neutralized with 1M Tris-HCl, pH 9.2 buffer. The protein

content was determined and adjusted to 1 mg/ml.

Immunoprecipitation

Co-Immunoprecipitation (Co-IP) protocols were performed using anti-V5 antibody

(CPN60$^{V5\text{-apt}}$) or anti-Ty antibody (ClpP$^{DEAD\text{-Ty}}$). Pellets from 4×10^8 parasites were

isolated using cold saponin and were lysed and sonicated in Extraction Buffer (40

mM Tris HCl pH 7.6, 75 mM KCl, and 1 mM EDTA, 15% glycerol) supplemented with

HALT protease inhibitor (Thermo). 10% of the sample was kept for later analysis

(input sample). Rabbit anti-V5 (Cell Signaling, D3H8Q) or anti-Ty, BB2 (inhouse

purification) antibodies were crosslinked to Dynabeads protein G beads (Invitrogen)

by incubating with 5 mM BS3 crosslinker (CovaChem) for 30 minutes. Quenching of

crosslinker was performed using 1 M Tris HCl (pH 7.5) for 30 minutes and then

washing with 1.2M Glycine HCl (pH 2.5) to remove excess unbound antibody.

Antibody-conjugated beads were then washed 3 times with PBS and incubated with

the supernatant at 4°C. Washes were performed using a magnetic rack (Life

Technologies). Samples were run on SDS-page and blotted with anti-V5 or anti-Ty

antibodies.

Western blot

Western blots were performed as described previously[15]. Briefly, parasites were

collected and host red blood cells were permeabilized selectively by treatment with

ice-cold 0.04% saponin in PBS for 15 min, followed by a wash in ice-cold PBS. Cells

were lysed using RIPA buffer, sonicated, and cleared by centrifugation at 4oC. The

antibodies used in this study were mouse anti-Ty, BB2 (inhouse purification), rabbit

anti-V5, D3H8Q (Cell Signaling, 1:3000), mouse anti-V5, E9H80 (Cell Signaling,

1:3000), rabbit anti-aldolase ab207494 (abcam, 1:3000). The secondary antibodies

that were used are IRDye 680CW goat anti-rabbit IgG and IRDye 800CW goat anti-

mouse IgG (LICOR Biosciences, 1:20,000). The Western blot images and

quantifications were processed and analyzed using the Odyssey infrared imaging

system software (LICOR Biosciences). Three biological replicates were used for

quantification of protein expression levels using Prism (GraphPad Software, Inc.).

Microscopy and image processing

For IFA, cells were fixed using a mix of 4% paraformaldehyde and 0.015%
glutaraldehyde and permeabilized using 0.1% Triton-X100. Primary antibodies used
are mouse anti-Ty1, BB2 (Invitrogen, 1:100), and rabbit anti-V5, D3H8Q (Cell
Signaling, 1:100). Secondary antibodies used are Alexa Fluor 488 and Alexa Fluor
546 (Life Technologies, 1:100). For mitochondrial staining, live cells were incubated
with 300 nM Mito-tracker Deep Red (Invitrogen) for 15 minutes, and then fixed and
stained as described above. Cells were mounted on Fluoroshield with DAPI (Sigma).
The imaging was performed using a Nikon Spinning Disk confocal fluorescence
microscope equipped with a 1005 x/1.4NA objective. Images were collected as Z-
stack, and displayed as maximum intensity projection. Image processing, analysis
and display were performed using Zeiss ZEN 3.7 Software. Adjustments of
brightness and contrast were made for display purposes. Blood smears were imaged
using an upright Eclipse E200 Microscope (Nikon), equipped with x100 oil objective
Type NVH, and captured using a digital Sight 1000 microscope camera and NIS-
Elements Software (Nikon). Figures for this manuscript were generated using
BioRender.com.

Ultrastructure Expansion Microscopy

Ultrastructure expansion microscopy was performed as previously described[62] with
minor modifications. Briefly, parasite cells were adhered to poly-d-lysine coated
coverslips, washed and fixated in PFA, and then incubated on gels, followed by
transfer to denaturation buffer. Gels were then incubated at 95°C, and then
expanded in doubled-distilled water. The gels were then measured for expansion
factor (5X in this case). The gel is then blocked (3% BSA) and incubated with
antibodies. antibodies used in this study: rabbit anti-V5, D3H8Q (Cell Signaling,
1:100), Alexa Fluor 488 goat anti-rabbit (1:200, Cell Signaling), NHS-ester Alexa
Fluor 405 (1:250, Thermo Fisher Scientific) and SYTOX Deep Red (1:1000, Thermo
Fisher Scientific). The imaging was performed using Airyscan2 LSM980 confocal
microscope equipped with 63x/1.4NA objective.

RNA extraction and Quantitative Real Time PCR (qRT-PCR)

Total RNA was extracted using the Macherey-Nagel NucleoSpin RNA Prep Kit. RNA
concentration and purity were assessed using a NanoDrop spectrophotometer. For
reverse transcription, 10 µL of the total RNA (from a 40 µL eluate) were used to

synthesize complementary DNA (cDNA) using the QuantaBio qScript cDNA
Synthesis Kit.

Quantitative PCR was performed using the Luna Universal qPCR Master Mix (New
England Biolabs) on a QuantStudio 5 Real-Time PCR System. Gene-specific
primers were used to amplify CPN60 (P14+P15) , LRR5 (PF3D7_1432400,
P20+P21), and the two housekeeping genes aldolase (P16+P17) and arg-tRNA
synthetase (P18+P19) for normalization. Each reaction was performed in triplicate.
Relative expression levels were calculated using the ΔΔCt method, normalizing to
the geometric mean of the housekeeping genes and comparing expression levels in
heat-shocked samples to the non-treated control. Statistical analyses were
performed using GraphPad Prism.

AF3 structure prediction and Alanine scanning

The inputs are protein and peptide sequence. All different versions were run with
random seed 2 or 1. Runs was done on alphafold server from GitHub
(https://alphafoldserver.com/). Structural Visualization was done using the PyMOL
Molecular Graphics System (Version 2.4.0) and UCSF ChimeraX (Version 1.4).
Robetta computational alanine scanning was applied to define the critical interface
residues, as implemented in http:://robetta.org. Detailed protocol was described by
Kortemme *et al*[59].

Generating the apicoplast and mitochondrial PfCPN60 models with PBZ1587
bound at the ring-ring interface

CryoEM structures were recently reported indicating PBZ1587 binds with 7-fold
symmetry between adjacent subunits across the ring-ring interface of *E. coli*
GroEL[37]. To support whether PBZ inhibitors would similarly bind to and inhibit the
apicoplast and mitochondrial PfCPN60 isoforms, we generated homology models for
the two PfCPN60 isoforms based on the GroEL-1587 cryoEM structure (9C0B) using
Schodinger Maestro. We generated two models for each isoform, one with the apical
domains in a conformation where the substrate binding sites are oriented inwards
towards the central cavity (like in the GroEL-1587 9C0B structure), and the other
with the apical domains in an extended conformation and oriented upwards from the
central cavity (like in the apicoplast PfCPN60 structure), described as follows.

1. Apicoplast PfCPN60-1587 model with apical domains positioned inwards: We first
employed the build homology model, multiple templates (homomultimer) function
using default parameters and aligning the apicoplast PfCPN60 sequence onto the
GroEL-1587 tetradecamer structure (9C0B). This provided a reasonable initial model
for further refinements of the ligand binding sites. As the alpha-helices spanning
residues 496-541 present in the apicoplast PfCPN60 structure were not represented
as such in the homology model, we then spliced these secondary structures into the
model. Next, we independently refined each of the 7 binding sites using the Refine
Protein-Ligand function using default parameters, with the exception of refining
atoms within 10 Å of the ligands. We then conducted Induced Fit Docking of the
PBZ1587 inhibitor independently at each site using default parameters for Extended
Sampling, with the exception of allowing refining of residues within 8.0 Å of ligand
poses. The docked structure was then subjected to final refinement through the
Protein Preparation Workflow using default parameters.

2. Mitochondrial PfCPN60-1587 model with apical domains positioned inwards: We
first employed the build homology model, multiple templates (homomultimer) function
using default parameters and aligning the mitochondrial PfCPN60 sequence onto the
GroEL-1587 tetradecamer structure (9C0B). Next, we independently refined each of
the 7 binding sites using the Refine Protein-Ligand function using default
parameters, with the exception of refining atoms within 10 Å of the ligands. We then
conducted Induced Fit Docking of the PBZ1587 inhibitor independently at each site
using default parameters for Extended Sampling, with the exception of allowing
refining of residues within 8.0 Å of ligand poses. The docked structure was then
subjected to final refinement through the Protein Preparation Workflow using default
parameters.

3. Apicoplast PfCPN60-1587 model with apical domains extended upwards: We first
aligned the individual apicoplast PfCPN60 rings on those of the GroEL-1587
tetradecamer structure (9C0B) so as to retain the apical domains in the extended,
upwards conformation as in the reported PfCPN60 structure. Next, we independently
refined each of the 7 binding sites using the Refine Protein-Ligand function using
default parameters, with the exception of refining atoms within 10 Å of the ligands.
We then conducted Induced Fit Docking of the PBZ1587 inhibitor independently at
each site using default parameters for Extended Sampling, with the exception of

allowing refining of residues within 8.0 Å of ligand poses. The docked structure was
then subjected to final refinement through the Protein Preparation Workflow using
default parameters.

4. Mitochondrial PfCPN60-1587 model with apical domains extended upwards: We
first employed the build homology model, multiple templates (homomultimer) function
using default parameters and aligning the mitochondrial PfCPN60 sequence onto the
apicoplast-1587 homology model generated above, with the apical domains
extended upwards (model 3). Next, we independently refined each of the 7 binding
sites using the Refine Protein-Ligand function using default parameters, with the
exception of refining atoms within 10 Å of the ligands. We then conducted Induced
Fit Docking of the PBZ1587 inhibitor independently at each site using default
parameters for Extended Sampling, with the exception of allowing refining of
residues within 8.0 Å of ligand poses. The docked structure was then subjected to
final refinement through the Protein Preparation Workflow using default parameters.

Data Availability
All of the data described are included in the manuscript and Figures.

Acknowledgements

We thank Boris Striepen for the BB2 hybridoma cell line; Florentin lab members for
critical reading and comments on the manuscript; Yael Feinstein-Rotkopf at the
Imaging Core Facility at the Faculty of Medicine, En Kerem; VEuPathDB and
PlasmoDB for scientific, academic and community support. This work was supported
by grants from the Israel Science Foundation (400/22 and 2786/22 to AF, and
301/21 to OFS) and HUJI start-up funds to AF. AT and SKG are supported by PhD
scholarships from Science Training Encouraging Peace (STEP), and AT is also
supported by Isadore Sharp Einstein scholarship. AQ is supported by Carole and
Andrew Harper Diversity PhD Program. AF is supported by the Abisch-Frenkel
Faculty Development Lectureship. AF, AT, MS, AQ, SKG and YR acknowledge
support by The Kuvin Center for the Study of Infectious and Tropical Diseases.

Author Contributions:
Conceptualization: AF, AT
Formal analysis: AF, AT, SMJ
Funding Acquisition: AF

Investigation: AF, AT, YR, MS, SKG, AQ

Methodology: NT, OFS, SMJ, AF, AT, YR, MS, SKG, AQ

Project Administration: AF

Resources: NT, OFS, SMJ

Supervision: AF

Validation: AF, AT

Writing – Original Draft Preparation: AF

Writing – Review & Editing: AF, AT, SMJ, MS, NT, OFS

Declaration of Interests: S.M.J. is a founder of BioEL Inc, which was formed to

commercialize GroEL inhibitors for antibacterial applications. The remaining authors

declare no competing interests.

Supplemental information

Document Sup Figures: Figures S1–S7

Preprint Server: BioRxiv

Figures

Figure 1

Figure 1. Generation and characterization of transgenic CPN60^{V5-apt} parasites.

A. Schematic representation of CRISPR-mediated homologues recombination through CPN60 homology regions enabling the integration of the V5 tag and the tetR-Dozi system. **B.** PCR amplification of genomic DNA isolated from CPN60 transfected parasites. Primers used are depicted in A and in Table 2. Primers are integration specific and thus can amplify 1.6 KB (P7+P6) and 11 KB (P7+P4) products only if integration occurs at the desired genomic locus. **C.** Western blot depicting two clones of CPN60^{V5-apt} parasites (A4 and F11) following 96-hour incubation with or without 30 µM chloramphenicol (CHL) using an anti-V5 antibody and anti-aldolase as a loading control. **D.** CPN60^{V5-apt} parasites were synchronized by incubation with 5% sorbitol followed by Percoll-mediated schizont-enrichment the following day. Western blot depicting expression profile of tagged CPN60 throughout parasite life cycle using an anti-V5 antibody. Right: Blood smears demonstrate parasites' developmental stage at the time of protein extraction. **E.** CPN60^{V5-apt} parasite clones (A4 and F11) were synchronized using 5% sorbitol, and 8ml of

culture were collected at the following time points: early ring stage (~12 h post-

invasion), early trophozoite stage (~28 hpi), late trophozoite stage (~36 hpi), and

schizont stage (~44 hpi). Parasites were liberated from host red blood cells using

0.1% saponin, and total mRNA was extracted and reverse-transcribed to cDNA.

Quantitative real-time PCR (qRT-PCR) was performed to quantify transcript levels of

the apicoplast-encoded chaperonin CPN60. Primers P14+P15 were used to amplify

CPN60 and two housekeeping genes (aldolase P16+P17 and arg-tRNA synthetase

P18+P19) for normalization. Gene expression was normalized to the mean ΔCt of

both housekeeping genes (aldolase and arg-tRNA synthetase), and expression

levels were calculated relative to the early ring stage using the ΔΔCt method. Data

are presented as fold change ($2^{-\Delta\Delta Ct}$). Values represent the mean expression

from two parasite lines, and error bars indicate the standard deviation. Statistical

analyses were performed using GraphPad Prism. **F.** Densitometric analysis of

protein bands from Western blot in 1D. V5 signal was normalized to aldolase.

Figure 2

A.   Figure 2. Organellar Localization and expression of CPN60. A.
Immunofluorescence microscopy of CPN60$^{V5\text{-}apt}$ parasites in different developmental
stages. Z stack images processed as Maximum Intensity show from left to right:
merged DIC contrast, anti-V5 (CPN60, red), DAPI (parasite nucleus, blue), and
merge of fluorescentt channels. The imaging was performed using a Nikon Spinning

Disk confocal fluorescence microscope equipped with a 1005 x/1.4NA objective.

Scale bar is 2.5 µM. **B.** A representative image of an expanded parasite

demonstrates uniform apicoplast localization of CPN60 in a well-developed organelle

during mid-schizogony (eight nuclei). Images from left: NHS-Esther (total protein,

grey), merge, DNA (red), V5 (CPN60, green), and a 3D visualization. Images were

captured as Z-stacks (20 slices of 0.1 µm each) at 63x using Airyscan microscopy.

The images are displayed as maximum intensity projections. Expansion factor is 5X.

Scale bar is 5 µm. **C.** Immunofluorescence microscopy of CPN60[V5-apt] parasites

processed as Maximum Intensity show from left to right: merged DIC contrast, DAPI

(parasite nucleus, blue), mito-tracker (mitochondrion, magenta), anti-V5 (CPN60,

green), and merge of fluorescentt channels. The imaging was performed using a

Nikon Spinning Disk confocal fluorescence microscope equipped with a 1005

x/1.4NA objective. Scale bar is 2.5 µM.

Figure 3

Figure 3. CPN60 is essential for parasite's viability due to its apicoplast function. A. Schematic representation of the tetR-Dozi system, which enables protein knockdown upon removal of the tetracycline analogue aTC. **B.** Western blots depicting the expression of the tagged CPN60 using an anti-V5 antibody following knockdown induction. A significant reduction in protein levels is already observed after 24 hours, and falls below detection 72 hours after aTC removal. **C.** Growth curve of CPN60^{V5-apt} parasites. Independent CPN60^{V5-apt} clones A4 and F11 were grown with or without 0.5 mM aTC, and parasitemia was monitored every 24 h over two cycles via flow cytometry. The 100% of growth represents the highest value of calculated parasitemia (final parasitemia in the presence of aTC). Normalized data are represented as mean ± SEM for three technical replicates. **D.** Growth curve of CPN60^{V5-apt} parasites, clone A4, supplemented with 200 μM IPP. The 100% of growth represents the highest value of calculated parasitemia (final parasitemia in the presence of aTC). Normalized data are represented as mean ± SEM for three technical replicates. **E.** CPN60^{V5-apt} parasites were synchronized by incubation with 5% sorbitol followed by Percoll-mediated schizont-enrichment the following day. Synched parasites were washed 8 times and then incubated with or without aTC. Giemsa-stained blood smears were imaged using an upright Eclipse E200 Microscope. **F.** Growth rates of synchronous CPN60^{V5-apt} parasites. Clones A4 and F11 were grown with or without 0.5 mM aTC, and parasitemia was monitored every 24 h over two cycles via flow cytometry. Data are represented in columns as mean ± SEM for three technical replicates. **G.** Blood smears of synched CPN60^{V5-apt} parasites with or without aTC were analyzed for specific parasites developmental stages. Minus aTC parasites demonstrate a delayed trophozoite stage already on the first replication cycle, fail to egress, and die during schizogony. Dead schizonts are defined as morphologically aberrant mutant cells with more than one nucleus that linger for more than 48 hours but did not complete their first replication cycle. **H.** Parasites were cultured without aTC and supplemented with IPP for 17 days. Parasites culture was then divided into four groups treated as following: (A) +aTC, +IPP; (B) +aTC, -IPP; (C) -aTC, +IPP; (D) -aTC, -IPP. Parasitemia of these four groups was monitored every 24 hours over two cycles via flow cytometry. Normalized data are represented as mean ± SEM for three technical replicates. **I.** Immunofluorescence microscopy of untreated and group (A) CPN60^{V5-apt} parasites. Z

stack images processed as Maximum Intensity show from left to right: merged DIC

contrast, anti-V5 (CPN60, red), DAPI (parasite nucleus, blue), and merge of

fluorescentt channels. The imaging was performed using a Nikon Spinning Disk

confocal fluorescence microscope equipped with a 1005 x/1.4NA objective. Scale

bar is 2.5 μM. **J.** Western blot depicting expression of untreated and group (A)

CPN60$^{V5\text{-}apt}$ parasites using an anti-V5 antibody and anti-aldolase as a loading

control.

Figure 4

Figure 4: **CPN60 is not involved in Heat Shock response**. **A.** Workflow of HS
experiments and representative blood smears. Parasites are subjected to HS at
40°C for 24 hours and then allowed to recover at normal temperature (37°C). **B.**
Parasites were subjected to HS and recovery as depicted in (A). Left: Cellular
lysates were extracted every 24 hours over a span of four days and subjected to
Western blot analysis using an anti-V5 antibody to follow tagged CPN60 levels.

Right: Quantification of CPN60 levels post-heat shock demonstrate no significant

increase in protein levels. **C**. *P. falciparum* culture was subjected to heat shock at

40°C for 24 hours, followed by return to standard conditions (37°C). Twelve milliliters

of parasite culture were collected at 24, 48, and 72 hours post-heat shock for RNA

extraction and cDNA synthesis. Quantitative real-time PCR (qRT-PCR) was

performed to assess the expression levels of CPN60 (P14+P15), and the heat

shock-responsive gene LRR5 (P20+P21). Gene expression was normalized to the

mean ΔCt of two housekeeping genes; aldolase (P16+P17) and arg-tRNA

synthetase (P18+P19), and values were calculated relative to the non-heat-shocked

control using the ΔΔCt method. The graph displays –ΔΔCt values. Error bars

represent the standard deviation from three biological replicates. While LRR5

expression increased post-heat shock, CPN60 levels remained unchanged,

suggesting that CPN60 is not transcriptionally regulated in response to heat stress.

D. To calculate EC50 of aTC, CPN60$^{V5\text{-}apt}$ parasites were thoroughly washed to

remove aTC, and then incubated in serial aTC dilutions in a 96-well plate.

Parasitemia was measured after 4 days using flow cytometry, showing an EC50 of

4.3 nM at 37^{o}c or 4.5 nM following 24 h incubation at 40^{o}c. Data are fit to a dose-

response equation and are represented as mean ± SEM. One representative

experiment out of three is shown. **E**. Parasites were thoroughly washed and then

incubated with 8 nM aTC (2X EC50). Left: Parasites lysates were obtained every 24

hours and subjected to western blot analyses by being probed with antibodies

against V5 (green) and aldolase (loading control, red). The protein marker sizes that

co-migrated with the probed protein are shown on the left. Right: Densitometric

analysis of protein bands on the left indicating 50% drop in CPN60 expression. **F.**

CPN60$^{V5\text{-}apt}$ parasites were washed and incubated with different aTC concentrations

(8 nM, 4 nM, 16 nM and 500 nM), subjected to HS and then allowed to grow at 37^{o}C

for three days, while being measured daily by flow cytometry. Data are fit to an

exponential growth curve and are represented as mean ± SEM for three technical

replicates.

Figure 5

Figure 5: CPN60 co-localizes, binds and stabilizes PfClpP. A. A diagram
depicting integration of a transgenic inactive PfClpP with a point mutation in the
proteolytic active site (S264A) and a C-terminal Ty tag. CRISPR-mediated
integration into the parasite's genome under the endogenous *hsp110* promoter
enables constant and moderate expression. **B.** Western blot analysis of lysates from
CPN60^{V5-apt}; ClpP$^{DEAD-Ty}$ parasite line, probed with antibodies against V5 (red)

against TY (green) showing the co-expression of the two proteins. Bottom: Schematic representation of the PfClpP processing: Full length cytoplasmic PfClpP with the transit peptide (TP, III), apicoplast-localized fraction after TP removal (II) and mature PfClpP after proteolytic cleavage of the pro-domain (I). **C.** EC50 of aTC in CPN60$^{V5\text{-apt}}$; ClpP$^{DEAD\text{-Ty}}$ parasite in 37^{o}C (blue) and following 24 hours incubation at 40^{o}C (green). Parasites were washed to remove aTC, and then incubated in serial aTC dilutions in a 96-well plate. Parasitemia was measured after 4 days using flow cytometry, showing comparable EC50s of 4.5 nM and 3.9 nM. Data are fit to a dose-response equation and are represented as mean ± SEM. One representative experiment out of three is shown. **D.** Immunofluorescence microscopy of CPN60$^{V5\text{-apt}}$; ClpP$^{DEAD\text{-Ty}}$ in different developmental stages. Z stack images processed as Maximum Intensity show from left to right: merged fluorescentt channels and DIC contrast, anti-V5 (CPN60, green), anti-ty (ClpP$^{DEAD\text{-Ty}}$, red) and DAPI (parasite nucleus, blue), along with merged fluorescentt channels. The imaging was performed using a Nikon Spinning Disk confocal fluorescence microscope equipped with a 1005 x/1.4NA objective. Scale bar is 2.5 µM. Right: Cytofluorograms depicting the fluorescence of each pixel in green channel (x-axis) against the red channel (y-axis) of maximum projection confocal images on the left. Pearson's coefficients (r) above 0.5 indicate significant co-localization between two fluorescent channels. **E.** Co-IP of CPN60$^{V5\text{-apt}}$ and ClpP$^{DEAD\text{-Ty}}$. Parasites were isolated and sonicated, and extracts were incubated with either anti-V5 antibody-conjugated beads (for CPN60 pulldown, left) or anti-Ty antibody-conjugated beads (for PfClpP pulldown, right). Input and IP samples were loaded on SDS-page and blotted with anti-Ty and anti-V5 antibodies. **F.** CPN60$^{V5\text{-apt}}$; ClpP$^{DEAD\text{-Ty}}$ parasites were incubated without aTC for 24 h, and lysates were extracted and subjected to Western blot analysis using antibodies against V5 (CPN60, red), Ty (ClpP$^{DEAD\text{-Ty}}$ parasite, green) and aldolase (red). CPN60 knockdown results in the disappearance of the processed form of PfClpP (I) and accumulation of the inactive zymogen (II). Gel image is representative of three biological replicates. **G.** Densitometric analysis of PfClpP protein bands from Western blot in (F). Each PfClpP processed form (I, II, III) was calculated as a fraction of total PfClpP expression. Results indicate a significant reduction in PfClpP processing upon CPN60 knockdown, suggesting that the interaction with CPN60 is essential for PfClp complex activity.

Fig. 6

Figure 6. Testing phenylbenzoxazole (PBZ) GroEL inhibitors for anti-

***Plasmodium* activity. A.** *Wildtype* NF54 parasites were seeded at 0.5% parasitemia

and incubated in serial dilutions of PBZ compounds, ranging from 50 µM to 50 nM.

Parasitemia was measured after 72 or 96 hours to calculate the drugs' half-maximal

effective concentrations (EC50). Data are fit to a dose-response equation and are
represented as mean ± SEM. **B.** CPN60^{V5-apt} parasites were synchronized and
treated with 12.5 µM of different PBZ inhibitors. Giemsa-stained blood smears were
imaged using an upright Eclipse E200 Microscope. Blood smears of CPN60^{V5-apt}
parasites with or without PBZ treatment were analyzed for specific parasites
developmental stages. PBZ-treated parasites demonstrate a delayed trophozoite
stage already on the first replication cycle, fail to egress, and die during schizogony.
C. Immunofluorescence microscopy of CPN60^{V5-apt} parasites following 24-hour
incubation with 12.5 µM of different PBZ inhibitors. Z stack images processed as
Maximum Intensity show from left to right: merged DIC contrast, anti-V5 (CPN60,
red), DAPI (parasite nucleus, blue), and merge of fluorescent channels. The imaging
was performed using a Nikon Spinning Disk confocal fluorescence microscope
equipped with a 1005 x/1.4NA objective. Scale bar is 2.5 µM.

Figure 7

Figure 7. Testing specificity of PBZ compounds against apicoplast activity. A.

Wildtype NF54 parasites were incubated with either DMSO (blue, control), 30μM

Chloramphenicol (green, CHL) or 12.5μM of different PBZ compounds (shades of

red), with (dashed) or without (solid) 200μM IPP. Parasitemia was monitored every

24 h over at least two cycles via flow cytometry. 100% of growth represents the

highest value of calculated parasitemia (final parasitemia with DMSO). Normalized

data are represented as mean ± SEM for three technical replicates. **B.** *Wildtype*

NF54 parasites were incubated with DMSO (blue, control) or 12.5μM of PBZ1214

(red), with (dashed) or without (solid) 200μM IPP. Parasitemia was monitored every

24 h over two cycles via flow cytometry. Data are represented as mean ± SEM for

three technical replicates. **C.** *Wildtype* NF54 parasites were seeded at 0.5%

parasitemia and incubated in serial dilutions of PBZ1214 with (red) or without (blue)

200μM IPP. Parasitemia was measured after 72 hours to calculate the drugs' half-

maximal effective concentrations (EC50). Data are fit to a dose-response equation

and are represented as mean ± SEM.

Figure 8

960
Figure 8: A model for regulation of proteostasis in the apicoplast by CPN and

Clp complexes. A. Two opposing mechanisms interact and balance proteostasis in

the apicoplast; The CPN complex consists of the ATPase chaperonin CPN60 which

is arranged in two heptameric rings assisting in refolding organellar proteins. The Clp

proteolytic complex in its core contains the PfClpP protease which is, likewise,

arranged in two heptameric rings that degrade organellar proteins. The model shows

a complete representation of the interaction between the two tetradecamers of

CPN60 and PfClpP as predicted by AF3. PfClpP is depicted in blue and teal shades,

while CPN60 is colored in red, brown, and beige. A hypothetical model of the full

tetradecamers, constructed based on the solved structures, is shown in grey tones.

PfClpP rings can be found in two forms; one as an inactive zymogen which contains

an inhibitory pro-domain (C) and a processed form representing the proteolytically

active ring (shown in this model). Both forms bind the CPN complex, and the switch

between the two forms enable a balance between degradation and folding and

determine organellar proteostasis state. **B.** Alanine scanning predicting hotspot

residues in the interaction between PfClpP and CPN60. CPN60 is shown in salmon,

and PfClpP in light blue. Key interaction residues ("hotspots") are highlighted in dark

blue for PfClpP and dark red for CPN60. **C.** Structure prediction of the oligomerized

PfClpP zymogen coloured by confidence levels according to pLDDT scores. High-

confidence regions are shown in blue, while low-confidence regions are depicted in

red, indicating the varying reliability of the predicted structure. The upper panel

shows schematic representation of PfClpP posttranslational processing.

Test PBZ #	Structure	EC$_{50}$ Results (µM)	
		72 h	96 h
1587		0.91	0.94
1038		9.41	7.27
1001		3.90	3.68
1064		5.27	4.65
1133		4.54	3.76
1135		21.15	18.78
1134		25.18	25.15
1221		6.90	7.03
1039		8.56	10.60
1200		7.03	6.29
1207		5.42	4.98
1214		14.02	10.95

985   Table 2

Table 2: Primers and Oligos used in this study		
Primers	**Primer name**	**Primer sequence**
P1	Apt-CPN60-Cterm-EcoRV-F	TTTTTTTTGGATATCGAATCTGAAATGGAATTACAAAAAATGGGGGCTAATATAG
P2	CPN60 cterm apt R	TTGGTTTACCCTCGAGTTCATCATAATTGTATCCGTCGTTCATGCTATCTTCATCATC
P3	Apt-AflII-CPN60-3UTR-F	CTTTCCGGGCGCGCCTTAAGCACACATATATATATATATATATATATATAAATTG
P4	CPN60-3UTR-EcoRV-Apt-R	TTCAGATTCGATATCCAAAAAAAACAAAAATTTTTATGTATTTATCCATATAC
P5	aptamer Forward	CTTATGACGTACCTGATTATGCAC
P6	aptamer Reverse	GTAGACCCCATTGTGAGTACATAAATATATTATATAAACTAGACTAGG
P7	CPN60 integr specific F	GTT CAA ATA ATG ACG AAA AGA AAT ATC TTG AAC TAA TAG G
P8	U6 CPN60 guide F	TAAGTATATAATATTGATGAAGATTCTATGAATGAGTTTTAGAGCTAGAA
P9	U6 CPN60 guide R	TTCTAGCTCTAAAACTCATTCATAGAATCTTCATCAATATTATATACTTA
P10	U6 HSP110 guide F	TAAGTATATAATATTGACAAACTTGGATGCAAATGGTTTTAGAGCTAGAA
P11	U6 HSP110 guide R	TTCTAGCTCTAAAACCATTTGCATCCAAGTTTGTCAATATTATATACTTA
P12	HSP110-Cterm pcr test F	GAGATCGTATTTTACTTTCCTTAGATGATTATGAAAATTGG
P13	HSP110-UTR pcr test R	GCATCAATGCACAAAATAAAAATCATACAATTTGGGG
P14	CPN60 qPCR F	GTCATTTTCATCTTTACATAATTTTCTTTTC
P15	CPN60 qPCR R	TTTGTTGCTTGTAATATTTTTGTTATGTTTGCGG
P16	Aldolase qPCR F	GCAAAGAGTATTATAAAGCTGGTGCAAGGTTTGC
P17	Aldolase qPCR R	GCATATCTAGCCAATCCCCATGCAGTTTCGTG
P18	arginine-tRNA ligase qPCR F	AAGAGATGCATGTTGGTC
P19	arginine-tRNA ligase qPCR R	GTACCCCAATCACCTACA
P20	LRR5 qPCR F	AATGTTTTGGACACGTTAATAGAAGAACAG
P21	LRR5 qPCR R	CTACAGTATATTCCTTCTCATCAAAATCG

References

1. Venkatesan, P. The 2023 WHO World malaria report. *Lancet Microbe* **5**, e214 (2024).

2. McFadden, G. I., Reith, M. E., Munhalland, J. & Lan-Unnasch, N. Plastid in human parasites.
Nature **381**, 482–482 (1996).

3. Köhler, S. *et al.* A plastid of probable green algal origin in Apicomplexan parasites. *Science* **275**,
1485–1489 (1997).

4. Dooren, G. G. van & Striepen, B. The algal past and parasite present of the apicoplast. *Annual*
Review of Microbiology **67**, 271–289 (2013).

5. Ralph, S. A. *et al.* Metabolic maps and functions of the Plasmodium falciparum apicoplast. *Nat Rev*
Microbiol **2**, 203–216 (2004).

6. Fichera, M. E. & Roos, D. S. A plastid organelle as a drug target in apicomplexan parasites. *Nature*
390, 407–409 (1997).

7. Biddau, M. & Sheiner, L. Targeting the apicoplast in malaria. *Biochem. Soc. Trans.* **47**, 973–983
(2019).

8. Waller, R. F., Reed, M. B., Cowman, A. F. & McFadden, G. I. Protein trafficking to the plastid of
Plasmodium falciparum is via the secretory pathway. *Embo J* **19**, 1794–1802 (2000).

9. Foth, B. J. *et al.* Dissecting Apicoplast Targeting in the Malaria Parasite Plasmodium falciparum.
Science **299**, 705–708 (2003).

10. Swift, R. P. *et al.* A mevalonate bypass system facilitates elucidation of plastid biology in malaria
parasites. *PLoS Pathog.* **16**, e1008316 (2020).

11. Yeh, E. & DeRisi, J. L. Chemical Rescue of Malaria Parasites Lacking an Apicoplast Defines
Organelle Function in Blood-Stage Plasmodium falciparum. *PLoS Biol.* **9**, e1001138 (2011).

12. Walczak, M., Ganesan, S. M., Niles, J. C. & Yeh, E. ATG8 Is Essential Specifically for an
Autophagy-Independent Function in Apicoplast Biogenesis in Blood-Stage Malaria Parasites. *Mbio* **9**,
e02021-17 (2018).

13. Swift, R. P., Rajaram, K., Liu, H. B. & Prigge, S. T. Dephospho☐CoA kinase, a nuclear☐encoded
apicoplast protein, remains active and essential after Plasmodium falciparum apicoplast disruption.
Embo J e107247 (2021) doi:10.15252/embj.2020107247.

14. Florentin, A. *et al.* PfClpC Is an Essential Clp Chaperone Required for Plastid Integrity and Clp
Protease Stability in Plasmodium falciparum. *Cell reports* **21**, 1746–1756 (2017).

15. Florentin, A., Stephens, D. R., Brooks, C. F., Baptista, R. P. & Muralidharan, V. Plastid
biogenesis in malaria parasites requires the interactions and catalytic activity of the Clp proteolytic
system. *Proceedings of the National Academy of Sciences of the United States of America* **3**,
201919501 (2020).

16. Schirmer, E. C., Glover, J. R., Singer, M. A. & Lindquist, S. HSP100/Clp proteins: a common
mechanism explains diverse functions. *Trends in biochemical sciences* **21**, 289–296 (1996).

17. Nishimura, K. & Wijk, K. J. van. Organization, function and substrates of the essential Clp
protease system in plastids. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1847**, 915–930
(2015).

18. Bouchnak, I. & Wijk, K. J. van. Structure, Function and Substrates of Clp AAA+ protease systems
in cyanobacteria, plastids and apicoplasts; a comparative analysis. *J Biol Chem* **296**, 100338 (2021).

19. Olinares, P. D. B., Kim, J., Davis, J. I. & Wijk, K. J. van. Subunit Stoichiometry, Evolution, and
Functional Implications of an Asymmetric Plant Plastid ClpP/R Protease Complex in Arabidopsis.
Plant Cell **23**, 2348–2361 (2011).

20. Kim, J. *et al.* Subunits of the Plastid ClpPR Protease Complex Have Differential Contributions to
Embryogenesis, Plastid Biogenesis, and Plant Development in Arabidopsis. *The Plant Cell* **21**, 1669–
1692 (2009).

21. Bakkouri, M. E. *et al.* Structural insights into the inactive subunit of the apicoplast-localized
caseinolytic protease complex of Plasmodium falciparum. *Journal of Biological Chemistry* **288**,
1022–1031 (2013).

22. Bakkouri, M. E. *et al.* The Clp Chaperones and Proteases of the Human Malaria Parasite
Plasmodium falciparum. *J. Mol. Biol.* **404**, 456–477 (2010).

23. Singh, M. K. *et al.* Molecular Chaperonin HSP60: Current Understanding and Future Prospects.
Int. J. Mol. Sci. **25**, 5483 (2024).

24. Dupuy, E. & Collet, J.-F. Entering deeper into the mysteries of the GroEL–GroES nanomachine.
Curr. Opin. Microbiol. **79**, 102480 (2024).

25. Perez, R. F., Mochi, G., Khan, A. & Woodford, M. Mitochondrial Chaperone Code: Just warming
up. *Cell Stress Chaperones* **29**, 483–496 (2024).

26. Fellows, J. D., Cipriano, M. J., Agrawal, S. & Striepen, B. A Plastid Protein That Evolved from
Ubiquitin and Is Required for Apicoplast Protein Import in Toxoplasma gondii. *mBio* **8**,
10.1128/mbio.00950-17 (2017).

27. Ganesan, S. M., Falla, A., Goldfless, S. J., Nasamu, A. S. & Niles, J. C. Synthetic RNA-protein
modules integrated with native translation mechanisms to control gene expression in malaria
parasites. *Nature Communications* **7**, 10727 (2016).

28. Elaagip, A., Absalon, S. & Florentin, A. Apicoplast Dynamics During Plasmodium Cell Cycle.
Front. Cell. Infect. Microbiol. **12**, 864819 (2022).

29. Michal, S. *et al.* Organelle Development and Inheritance are Driven by Independent Nuclear and
Organellar Mechanisms in Malaria Parasites. *bioRxiv* 2025.03.06.641809 (2025)
doi:10.1101/2025.03.06.641809.

30. Dahl, E. L. & Rosenthal, P. J. Multiple Antibiotics Exert Delayed Effects against the Plasmodium
falciparum Apicoplast. *Antimicrob. Agents Chemother.* **51**, 3485–3490 (2007).

31. Zhang, M. *et al.* The apicoplast link to fever-survival and artemisinin-resistance in the malaria
parasite. *Nat. Commun.* **12**, 4563 (2021).

32. Thomas, P. *et al.* Phenotypic Screens Identify Parasite Genetic Factors Associated with Malarial
Fever Response in Plasmodium falciparum piggyBac Mutants. *mSphere* **1**, e00273-16 (2016).

33. Nguyen, B., Ma, R., Tang, W. K., Shi, D. & Tolia, N. H. Crystal structure of P. falciparum Cpn60
bound to ATP reveals an open dynamic conformation before substrate binding. *Sci. Rep.* **11**, 5930
(2021).

34. Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with AlphaFold 3.
Nature **630**, 493–500 (2024).

35. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–
589 (2021).

36. Johnson, S. M. *et al.* A biochemical screen for GroEL/GroES inhibitors. *Bioorg. Med. Chem. Lett.*
24, 786–789 (2014).

37. Godek, J. *et al.* Bis-sulfonamido-2-phenylbenzoxazoles Validate the GroES/EL Chaperone
System as a Viable Antibiotic Target. *J. Am. Chem. Soc.* **146**, 20845–20856 (2024).

38. Gruber, A. V. & Feiz, L. Rubisco Assembly in the Chloroplast. *Front. Mol. Biosci.* **5**, 24 (2018).

39. Sheiner, L. *et al.* A Systematic Screen to Discover and Analyze Apicoplast Proteins Identifies a
Conserved and Essential Protein Import Factor. *PLoS Pathog.* **7**, e1002392 (2011).

40. Elaagip, A., Absalon, S. & Florentin, A. Apicoplast dynamics during Plasmodium cell cycle.
Front. Cell. Infect. Microbiol. Parasite and Host (2022).

41. Bowman, J. D. *et al.* Antiapicoplast and Gametocytocidal Screening To Identify the Mechanisms
of Action of Compounds within the Malaria Box. *Antimicrob. Agents Chemother.* **58**, 811–819
(2014).

42. Boucher, M. J. & Yeh, E. Disruption of Apicoplast Biogenesis by Chemical Stabilization of an
Imported Protein Evades the Delayed-Death Phenotype in Malaria Parasites. *mSphere* **4**,
10.1128/msphere.00710-18 (2019).

43. Amberg-Johnson, K. *et al.* Small molecule inhibition of apicomplexan FtsH1 disrupts plastid
biogenesis in human pathogens. *eLife* **6**, e29865 (2017).

44. Uddin, T., McFadden, G. I. & Goodman, C. D. Validation of Putative Apicoplast-Targeting Drugs
Using a Chemical Supplementation Assay in Cultured Human Malaria Parasites. *Antimicrob. Agents*
Chemother. **62**, 10.1128/aac.01161-17 (2018).

45. Okada, M., Guo, P., Nalder, S. & Sigala, P. A. Doxycycline has distinct apicoplast-specific
mechanisms of antimalarial activity. *eLife* **9**, e60246 (2020).

46. Pires, C. V., Chawla, J., Simmons, C., Gibbons, J. & Adams, J. H. Heat-shock responses: systemic
and essential ways of malaria parasite survival. *Curr. Opin. Microbiol.* **73**, 102322 (2023).

47. Tintó-Font, E. & Cortés, A. Malaria parasites do respond to heat. *Trends Parasitol.* **38**, 435–449
(2022).

48. Harbut, M. B. *et al.* Targeting the ERAD pathway via inhibition of signal peptide peptidase for
antiparasitic therapeutic design. *Proc. Natl. Acad. Sci.* **109**, 21486–21491 (2012).

49. Wang, N. *et al.* The cryo-EM structure of the chloroplast ClpP complex. *Nat Plants* **7**, 1505–1515
(2021).

50. Abdeen, S. *et al.* GroEL/ES inhibitors as potential antibiotics. *Bioorg. Med. Chem. Lett.* **26**, 3127–
3134 (2016).

51. Zimbone, S., Rosa, M. C. D., Chiechio, S. & Giuffrida, M. L. Exploring the Role of Hsp60 in
Alzheimer's Disease and Type 2 Diabetes: Suggestion for Common Drug Targeting. *Int. J. Mol. Sci.*
24, 12456 (2023).

52. Bavisotto, C. C. *et al.* Hsp60 Post-translational Modifications: Functional and Pathological
Consequences. *Front. Mol. Biosci.* **7**, 95 (2020).

53. Washburn, A. *et al.* Dual-targeting GroEL/ES chaperonin and protein tyrosine phosphatase B
(PtpB) inhibitors: A polypharmacology strategy for treating Mycobacterium tuberculosis infections.
Bioorg. Med. Chem. Lett. **29**, 1665–1672 (2019).

54. Abdeen, S. *et al.* Targeting the HSP60/10 chaperonin systems of Trypanosoma brucei as a strategy
for treating African sleeping sickness. *Bioorg. Med. Chem. Lett.* **26**, 5247–5253 (2016).

55. Leitner, A. *et al.* The Molecular Architecture of the Eukaryotic Chaperonin TRiC/CCT. *Structure*
20, 814–825 (2012).

56. Lu, K.-Y. *et al.* Plasmodium chaperonin TRiC/CCT identified as a target of the antihistamine
clemastine using parallel chemoproteomic strategy. *Proc. Natl. Acad. Sci.* **117**, 5810–5817 (2020).

57. Spillman, N. J., Beck, J. R., Ganesan, S. M., Niles, J. C. & Goldberg, D. E. The chaperonin TRiC
forms an oligomeric complex in the malaria parasite cytosol. *Cell. Microbiol.* **19**, e12719 (2017).

58. Mbengue, A. *et al.* New Export Pathway in Plasmodium falciparum Infected Erythrocytes: Role
of the Parasite Group II Chaperonin, PfTRiC. *Traffic* **16**, 461–475 (2015).

59. Muralidharan, V., Oksman, A., Iwamoto, M., Wandless, T. J. & Goldberg, D. E. Asparagine
repeat function in a Plasmodium falciparum protein assessed via a regulatable fluorescent affinity tag.
Proceedings of the National Academy of Sciences **108**, 4411–4416 (2011).

60. Kudyba, H. M., Cobb, D. W., Florentin, A., Krakowiak, M. & Muralidharan, V. CRISPR/Cas9
Gene Editing to Make Conditional Mutants of Human Malaria Parasite P. falciparum. *Journal of*
visualized experiments : JoVE (2018) doi:10.3791/57747.

61. Muralidharan, V., Oksman, A., Pal, P., Lindquist, S. & Goldberg, D. E. Plasmodium falciparum
heat shock protein 110 stabilizes the asparagine repeat-rich parasite proteome during malarial fevers. -
PubMed - NCBI. *Nature Communications* **3**, 1310 (2012).

62. Liffner, B. *et al.* Atlas of Plasmodium falciparum intraerythrocytic development using expansion
microscopy. *eLife* **12**, RP88088 (2023).

A

1.2 Kb

P7 → CPN60 | 3UTR ← P4

P7 → CPN60 | C-term | V5 | aptamer x10 ← P6 | 3UTR ← P4

1.8 Kb

10 Kb

B

P7 + P4   P7 + P6

CPN60^{V5-apt}   CPN60^{V5-apt}

Size (KB)   *wt*   A4   F11   *wt*   A4   F11

10 -
3 -
2 -
1.5 -
1 -

C

CPN60^{V5-apt}:   clone A4   clone F11

(KDa)   +CHL   +CHL

100 —   α V5

50 —   α aldolase
35 —

D

Hours Post Invasion

(KDa)   14   22   36   44

100 —   α V5

50 —   α aldolase
35 —

14h   22h   36h   44h

E

relative mRNA expression vs. hours post invasion (hpi)

12   28   36   44

F

Relative Protein Expression vs. hours post invasion (hpi)

14   22   36   44

A

Early-stage

Trophozoite

Schizont

Late Schizont (segmented)

Merge | α V5 (CPN60) | DAPI

B

NHS-Esther | Merge | Nuclei | α V5 (CPN60) | 3D

C

Merge | DAPI | mito-tracker | α V5 (CPN60)