Biophysical network modeling of temporal and stereotyped sequence propagation of neural activity in the premotor nucleus HVC

Zeina Bou Diab[1], Marc Chammas[1], Arij Daou[1*]

1. Biomedical Engineering Program, American University of Beirut, Lebanon

* Corresponding author: arij.daou@aub.edu.lb

- Number of pages: 65
- Number of figures: 13
- Number of words of abstract: 244
- Number of words of introduction: 702
- Number of words of discussion: 1642

- Conflict of interest: The authors declare no competing financial interests.
- Acknowledgements: Daniel Margoliash engaged in valuable conversations as these results developed. Z.B.D., M.C., and A.D. conceptualized, design the research and constructed the network; Z.B.D performed simulations and prepared figures, Z.B.D and A.D. interpreted results of simulations and drafted manuscript. This work was supported in part by grants provided to A.D. by the University Research Board and Farouk Jabr Foundation at the American University of Beirut.

Abstract

Stereotyped neural sequences are often exhibited in the brain, yet the neurophysiological mechanisms underlying their generation are not fully understood. Birdsong is a prominent model to study such behavior particularly because juvenile songbirds progressively learn from their tutors and by adulthood are able to sing stereotyped song patterns. The songbird premotor nucleus HVC coordinate motor and auditory activity responsible for learned vocalizations. The HVC comprises three neural populations that has distinct *in vitro* and *in vivo* electrophysiological responses. Typically, models that explain HVC's network either rely on intrinsic HVC circuitry to propagate sequential activity, rely on extrinsic feedback to advance the sequence or rely on both. Here, we developed a physiologically realistic neural network model incorporating the three classes of HVC neurons based on the ion channels and the synaptic currents that had been pharmacologically identified. Our model is based on a feedforward chain of microcircuits that encode for the different sub-syllabic segments (SSSs) and that interact with each other through structured feedback inhibition. The network reproduced the *in vivo* activity patterns of each class of HVC neurons, and unveiled key intrinsic and synaptic mechanisms that govern the sequential propagation of neural activity by highlighting important roles for the T-type Ca^{2+} current, Ca^{2+}- dependent K^+ current, A-type K^+ current, hyperpolarization activated inward current, as well as excitatory and inhibitory synaptic currents. The result is a biophysically realistic model that suggests an improved characterization of the HVC network responsible for song production in the songbird.

Significance Statement

Learned motor sequences acquired through repetitive practice undergo stabilization and

are integrated into cortical motor circuits through a process of consolidation. The process

of mastering a complicated motor sequence requires thorough motor exploration to

achieve closer alignment with the desired outcome, yet the neural mechanisms

underlying sequence generation remain largely unexplored. In this study, we investigate

the neural circuitry in a premotor region of the songbird brain (known as HVC) that is well-

known for generating extremely precise learned sequences. We develop a biophysically

realistic network model that incorporate pharmacologically identified intrinsic and synaptic

currents for the three different classes of HVC neurons. The network highlights

fundamental intrinsic and synaptic mechanisms regulating the sequential propagation of

neural activity.

 Introduction

Learned temporal sequences are expressed in many brain regions and play a critical role in temporal information encoding, such as navigation (Skaggs et al., 1996), the timing of motor actions (Wang et al., 2018), time generation (MacDonald et al., 2011; Pastalkova et al., 2008), decision making (Harvey et al., 2012; Schmitt et al., 2017) and skilled movement (Peters et al., 2014). Zebra finches sing remarkably stereotyped songs rendering them as an excellent model for studying the mechanisms underlying neural sequences (Hahnloser et al., 2002). The anatomical basis for song production and learning is a highly developed neural network known as the song system (Fig. 1A), with nucleus HVC exhibiting a rhythmic pattern-generating role encoding for the syllable order and the overall temporal structure of the birdsong (Fee & Goldberg, 2011; Fee et al., 2004; Fee & Scharff, 2010; Mooney, 2009; Yu & Margoliash, 1996).

There are three main neuronal populations in the HVC exhibiting different functional, cellular and pharmacological properties (Daou et al., 2013; Kubota & Taniguchi, 1998; Mooney, 2000; Mooney & Prather, 2005): neurons that project to the robust nucleus of arcopallium (RA; HVC_{RA}), neurons that project to Area X (HVC_X), and interneurons (HVC_{INT}, Fig. 1B). HVC_{Av} neurons (projecting to nucleus Avalanche) exists and play a role in song learning, yet their intrinsic and synaptic properties remains unknown (Roberts et al., 2017). Intracellular recordings showed that HVC_{RA}, HVC_X, and HVC_{INT} neurons have distinctive *in vivo* and *in vitro* electrophysiological properties (Daou et al., 2013; Dutar et al., 1998; Kubota & Taniguchi, 1998; Mooney et al., 2001; Shea et al., 2010), which are orchestrated via a family of ion channels (Daou et al., 2013).

91 ***Figure 1***. *Song system overview and general intrinsic and network properties of HVC neurons in vivo and in vitro.*

92 ***A***. *Schematic diagram showing a sagittal view of the male zebra finch song system. The vocal motor pathway (VMP,*

93 *red color) contains circuits that directly pattern song output. The anterior forebrain loop (AFP, blue color) pathway*

94 *contains circuits that are important for song learning and plasticity.* ***B***. *HVC includes multiple classes of neurons;*

95 *HVC$_X$ neurons that project to area X (blue), HVC$_{RA}$ neurons that project to nucleus RA (red) and HVC interneurons*

During singing, HVC_{RA} neurons produce a single burst of spikes (~10 ms) that are tightly

locked to the song (Fig. 1C, (Hahnloser et al., 2002; Kozhevnikov & Fee, 2007). HVC_X

neurons in their turn elicit 1-4 bursts that are also time-locked to vocalizations (Fig. 1D,

(Kozhevnikov & Fee, 2007), while HVC_{INT} neurons exhibit tonic activation (Fig. 1C), with

bursting and suppression at different locations throughout the song (Amador et al., 2013;

Cannon et al., 2015; Hahnloser et al., 2002; Kosche et al., 2015; Kozhevnikov & Fee,

2007; Long et al., 2010; Markowitz et al., 2015).

Several models of how sequence is generated within HVC have been proposed (Cannon

et al., 2015; Drew & Abbott, 2003; Egger et al., 2020; Elmaleh et al., 2021; Galvis et al.,

2018; Gibb et al., 2009a, 2009b; Hamaguchi et al., 2016; Jin, 2009; Long & Fee, 2008;

Markowitz et al., 2015). These models either rely on intrinsic HVC circuitry to propagate

sequential activity, rely on extrinsic feedback to advance the sequence or rely on both.

The proposed models do not capture the complex details of spike morphology, do not

include the right ionic currents, do not incorporate all classes of HVC neurons, or do not

generate realistic firing patterns as seen *in vivo*. In this work, we ask a simple but powerful

question: Can the biologically realistic intrinsic properties of HVC neurons and the local

synaptic connections among them produce the precise sequence propagation seen in

birdsong? We are particularly interested in understanding what core biophysical

ingredients (ionic and synaptic currents) are truly necessary to generate sparse

sequences while preserving each HVC neuron activity in vivo. How do features like

bursting behavior of these neurons or patterns of inhibition help shape the flow of activity

throughout network?" To address these questions, we developed a physiologically

realistic network model incorporating the three classes of HVC neurons based on the ion

channels and the synaptic currents that had been pharmacologically identified (Daou et

al., 2013; Kosche et al., 2015; Mooney & Prather, 2005). Our model is based on a

feedforward chain of microcircuits that encode for the different sub-syllabic segments

(SSSs) and that interact with each other through structured feedback inhibition. The

network developed unveiled key intrinsic and synaptic mechanisms that govern the

sequential propagation of neural activity by highlighting important roles for the T-type Ca^{2+}

current, Ca^{2+}- dependent K^+ current, A-type K^+ current, hyperpolarization activated inward

current, as well as excitatory and inhibitory synaptic currents. Our model provides a new

way of thinking about sequence generation during birdsong vocalizations and in network

architectures more generally.

Methods

Single-compartment conductance-based Hodgkin-Huxley-type (HH) biophysical models

of cells from the HVC were developed and connected together via biologically realistic

synaptic currents. Simulations of these model neurons and of the model network

composed of synaptically coupled HVC_{RA}, HVC_X, and HVC_{INT} neurons were performed

using the ode45 numerical integrator in MATLAB (MathWorks). Source codes for each

network will be made available online at our lab's website as well as on ModelDB.

HVC model cells that are used to connect the networks exhibited ionic and synaptic

currents that had been shown to be expressed pharmacologically (Daou et al., 2013;

Kosche et al., 2015; Long et al., 2010; Mooney & Prather, 2005). The functional forms of

activation/inactivation functions and time constants were based on previous published

mathematical neural models (Daou et al., 2013; Destexhe & Babloyantz, 1993; Dunmyre

et al., 2011; Hodgkin & Huxley, 1952; Terman et al., 2002; Wang et al., 2003), and the

parameters that were varied were merely the maximal conductances of some ionic

currents that vary among the various neuronal subtypes (Daou et al., 2013), as well as

the synaptic conductances. Every model neuron is represented by ordinary differential

equations for the different state variables as illustrated below.

Ion channels of model HVC neurons

We added a hyperpolarization-activated inward current conductance (I_H) to HVC_X and

HVC_{INT} because it is responsible for the sag seen in these neurons (Daou et al., 2013;

Dutar et al., 1998; Kubota & Saito, 1991; Kubota & Taniguchi, 1998), and we added a

low-threshold T-type Ca^{2+} current (I_{CaT})conductance responsible for the post-inhibitory

rebound firing seen in HVC$_X$ and HVC$_{INT}$ neurons (Daou et al., 2013). A small-

conductance Ca^{2+}-activated K$^+$ current (I_{SK}) was added for HVC$_{RA}$ and HVC$_X$ neurons as

it is responsible for the spike frequency adaptation feature that these two classes exhibit

(Daou et al., 2013). For interneurons, we integrated a large magnitude of the delayed

rectifier K$^+$ current conductance allowing these neurons to undershoot the resting

membrane potential as seen experimentally (Daou et al., 2013; Dutar et al., 1998; Kubota

& Saito, 1991; Kubota & Taniguchi, 1998). For HVC$_{RA}$ neurons, we added an A-type K+

current that supports the delay to spiking seen in response to depolarizing current pulses

(Daou et al., 2013; Kubota & Taniguchi, 1998; Mooney & Prather, 2005). High-threshold

Ca^{2+} conductance was added to all classes of HVC neurons (Daou et al., 2013; Kubota

& Saito, 1991; Long et al., 2010). In total, the model was designed to include spike-

producing currents (I_K and I_{Na}), a high-threshold L-type Ca^{2+} current (I_{CaL}), a low-

threshold T-type Ca^{2+} current (I_{CaT}), a small-conductance Ca^{2+} - activated K$^+$ current (I_{SK}),

an A-type K$^+$ current (I_A), a hyperpolarization-activated current (I_H), and a leak current

(I_L). The membrane potential of each HVC neuron obeys the following equations:

$$C_m \frac{dV_{RA}}{dt} = -I_L - I_K - I_{Na} - I_{CaL} - I_A - I_{SK} \qquad (1)$$

$$C_m \frac{dV_X}{dt} = -I_L - I_K - I_{Na} - I_{CaL} - I_{CaT} - I_{SK} - I_H \qquad (2)$$

$$C_m \frac{dV_{INT}}{dt} = -I_L - I_K - I_{Na} - I_{CaL} - I_{CaT} - I_H \qquad (3)$$

where C_m is the membrane capacitance. The associated equations and parameters for

each of the activation/inactivation gating variables for each ionic current are given in

(Daou et al., 2013) and shown below. In total, every single model HVC$_{RA}$, HVC$_X$, and

HVC_{INT} neuron had a total of 6, 8 and 7 ODEs, respectively, that govern their intrinsic

dynamics. Every synaptic current that was integrated to any model neuron added a new

ODE to the set of ODEs governing the membrane potential of the corresponding model

neuron. See APPENDIX 1 for detailed equations.

Synaptic currents

In addition to the ionic currents above that orchestrate the internal dynamics of each HVC

neuron, we integrated synaptic currents in order to reproduce the biological features of

the voltage traces observed *in vivo*. Excitatory (AMPA) and inhibitory ($GABA_A$) synaptic

currents were used to connect neurons inside each architecture based on the

pharmacological dual synaptic connections as described by Mooney and Prather (2005).

Each synaptic current represents the synaptic input(s) from the presynaptic cell(s) to the

particular HVC model neuron, and is modeled as $I_{syn} = \sum_X I_{X \to Y}$ where $I_{X \to Y} =$

$g_{X \to Y} s_{X \to Y} (V - V_{X \to Y})$. Here the summation is taken over the presynaptic HVC neurons

where X represents a presynaptic cell, Y represents a postsynaptic cell, $V_{X \to Y}$ is the

reversal potential for the synapse in the postsynaptic cell with $V_{X \to Y} = V_{AMPA}$ for excitatory

input and $V_{X \to Y} = V_{GABA-A}$ for inhibitory input.

The model equations for the synaptic currents are detailed in APPENDIX 1 , taken after

(Destexhe et al., 1994; Varela et al., 1997). We limited our synaptic currents' choices in

all networks to AMPA (excitatory) and $GABA_A$ (inhibitory) without integrating NMDA and

$GABA_B$ for the following reasons: 1) both AMPA and $GABA_A$ currents are voltage-

dependent with simple activation kinetics that do not depend on further parameters that

are very hard to tune or calibrate (for example, NMDA current relies on Mg^{2+} concentration

and $GABA_B$ on G-proteins dynamics, both which require additional ODEs and series of

parameters that we don't know about in the HVC), 2) adding the additional synaptic

currents do not have a significant contribution to the network dynamics we're building

because the emphasis is on excitation and inhibition and we could convey the

mechanisms we envision that orchestrate each network with these two currents solely.

Therefore, model HVC_{RA} neurons send their excitatory afferents to other HVC_{RA} neurons

as well as to HVC_{INT} neurons via AMPA currents. Model HVC_{INT} neurons send their

inhibitory afferents to both HVC_{RA} and HVC_X neurons via $GABA_A$ currents. And lastly,

model HVC_X neurons excite HVC_{INT} neurons via AMPA currents.

Desired network activity

Our aim here is to generate the optimal and desired network activity that's generated by

the three classes of HVC neurons during singing. Therefore, we focused on model

parameters and their underlying mechanisms that play key roles in 1) reproducing the

patterns without breaking the sequence of activity propagation and 2) generate

biologically realistic traces for each class as shown using intracellular recordings *in vivo*

in a way to maintain spike shapes, burst patterns, rebound firing/bursting, subthreshold

oscillations, etc … In a nutshell, the behavior of the network was considered desired and

"good" if the model voltage traces for the total populations in each of the three classes of

HVC neurons in the network matched the following: 1) the time-locked and characteristic

bursting of HVC_{RA} neurons (3-6 spikes for a ~10 ms duration), with spikes riding on a

plateau, 2) HVC_X neurons eliciting 1-4 bursts (4-9 spikes per burst) that are also time-

locked and that are mostly rebound bursts from inhibition (Lewicki, 1996; Mooney, 2000),

3) HVC$_{INT}$ neurons exhibiting tonic activation with spiking and bursting throughout the

song, and 4) spike frequency, spike amplitude, sags and/or rebound upon inhibition,

resting membrane potential and other known features of the intrinsic properties of the

three classes of HVC neurons exhibited for each of the classes that exhibit them (Daou

et al., 2013).

Maximal and synaptic conductances variations

Automated adjustment of model parameters was performed to qualitatively reproduce

desired membrane potential trajectories, as described next. Fixed parameter values for

HVC neurons used in the simulations are given in Table 1 in APPENDIX 2. Parameters

that vary between the different model neurons are shown in Figure 2-A.

Some maximal conductances were fixed while others were allowed to vary. We fixed g_{Na}

and g_K for each class of the HVC neurons to values that had been shown earlier to

accurately fit the spike morphologies (upstrokes and downstrokes of action potentials,

plateaus, etc…) in response to applied current given *in vitro* (Daou et al., 2013). For

example, HVC$_{INT}$ neurons' spikes exhibit a relatively large undershoot of the resting

membrane potential, while HVC$_X$ and HVC$_{RA}$ spikes ride on a plateau with characteristic

properties (Daou et al., 2013). We also fixed the value of g_{CaL} because we could achieve

the same accuracy of fitting by varying g_{SK}, and so could not distinguish between the two.

The four key conductances in our model that played crucial roles in controlling not only

the intrinsic properties of the HVC neurons they're expressed in but also in shaping overall

network activity and sequence propagation are g_{SK}, g_h, g_A and g_{CaT}. As a recap, g_{SK} is

expressed in HVC$_X$ and HVC$_{RA}$, g_H and g_{CaT} in HVC$_X$ and HVC$_{INT}$, and g_A in HVC$_{RA}$ only

(Daou et al., 2013). Random variations in these four parameters were performed to

qualitatively reproduce membrane potential trajectories of the three classes of HVC

neurons as seen firing when the bird is singing.

In a previous study, Daou and Margoliash (2020) showed that intracellular recordings

from X-projecting neurons in adult zebra finch brain slices share similar spike waveform

morphologies, with modeling indicating similar magnitudes of their principal ion currents.

To that end, we fixed in our network the intrinsic properties of the population of HVC_X

neurons to the same values. Therefore, the automated variations in the conductances

held for HVC_X neurons were done at the population level by varying a corresponding ionic

conductance value (say, g_{SK}) and setting its value to all of the HVC_X population. We also

checked the effects of removing this constraint in Figure 13, that is, allowing the intrinsic

properties of HVC_X neurons to vary like other parameters.

We first manually selected the four key conductances to default values that generate the

desired behavior of the network as described in the previous section. Network robustness

to varied maximal conductances as well as the legitimate ranges for each maximal

conductance was determined by simulating the network many times, each time with a

random variation in the maximal conductance about its default value. The network

response was considered accurate if the network generated sequential bursting and all

of the desired features described earlier. The range of variation of the randomly-varying

parameter was increased or decreased randomly up until the point that the network

ceased to be accurate.

The variation in parameters was done at the population level (setting the maximal

conductance for all neurons of the same class to a single value and randomly changing

that value for all), well as on the individual neuronal level (varying the maximal

conductance for one neuron at a time of the same class while fixing the others to their

default values). This is different for the HVC_X parameters, where the corresponding

conductances at the population and the neuronal levels are considered the same since

we assumed same intrinsic properties as described earlier. If any of the simulations

generated networks where any of the HVC_{RA} neurons elicit bursts exhibiting spikes

outside the 3-6 number of spikes range or duration of any burst longer than 10 ms, then

we ignore that parameter value. Similarly, if any of the simulations generate networks

where any of the HVC_X neurons eliciting more than 4 bursts, individual bursts exhibiting

spikes outside the 4-9 spikes/burst range, then the corresponding parameters are

ignored. Moreover, we also ignore parameters that generate unrealistic intrinsic

properties of individual HVC neurons; for instance, if the resting membrane potentials of

individual HVC neurons were outside the reported ranges (Daou et al., 2013), if HVC_X

and HVC_{INT} neurons failed to generate sags and rebounds (depolarization or bursting) in

response to inhibition, or if spikes' amplitudes were not realistic (HVC_X and HVC_{RA} spikes

not riding on plateaus or HVC_{INT} spikes not undershooting the RMP). We realize this might

be inducing tough constraints on the selection of the parameters limiting the space for

which the conductances are allowed to vary, but we opted for this method in order to

generate the optimal ranges in which intrinsic and synaptic conductances are able to

reproduce the biophysically realistic firing patterns seen during singing. We therefore

ended up with lists of ranges for each conductance that was varied and in each class of

HVC neuron, such that the desired network activity is maintained. Figure 2-A shows the

ranges for each maximal conductance that had been allowed to vary for the three classes

of HVC neurons, while maintaining robust network propagation and biologically realistic

in vivo behavior.

Figure 2. *Box plots showing the ranges of ionic and synaptic currents that were allowed to vary while maintaining*

*robust network propagation and biologically realistic in vivo behavior of all HVC neuronal classes. **A.** The ionic*

conductances that were varied are g_A of HVC_{RA}, g_{SK} of HVC_{RA} and HVC_X, g_h and g_{CaT} of HVC_{INT} and HVC_X. The shown

ranges reflect values whereby each neuron class is able to maintain realism in terms of electrophysiological behavior

*and network properties. **B.** Ranges of values of synaptic conductances that connect two classes of HVC neurons while*

conserving sequential activity propagation and the general network activity.

Similar to what was done with the maximal ionic conductances, we conducted automated

and random variations for all synaptic conductances in the model (no synaptic

conductance was fixed). Figure 2-B reports the ranges of the synaptic conductances that

were able to maintain the robustness of network propagation and the general *in vivo*-like

desired behavior of all neuronal classes. All synaptic conductances showed considerable

ranges during which sequential activity is propagated and the overall desired network

activity is maintained, with the exception of the GABA conductance from HVC_{INT} to HVC_{RA}

($g_{GABA_{INT \to RA}}$). Increasing $g_{GABA_{INT \to RA}}$ to larger magnitudes would induce an inhibition in

the HVC_{RA} pushing its voltage below its resting membrane potential (due to the dense

bursting and firing in HVC_{INT}), and this is not realistic because it's been shown that during

singing HVC_{RA} neurons ride on a depolarizing plateau throughout the song (Long et al.,

2010). Moreover, while the intrinsic properties of HVC_X neurons were set to the same

values, the synaptic parameters associated with each HVC_X neuron (afferent and

efferent) were allowed to vary from one neuron to another.

Moreover, to account for synaptic variability, we introduced a stochastic input current of

the form $I_{noise}(t) = \sigma.\xi(t)$ where $\xi(t)$ is a Gaussian white noise with zero mean and unit

variance, and σ is the noise amplitude. This stochastic drive was introduced to every

model neuron and it mimics the fluctuations in synaptic input arising from random

presynaptic activity and background noise. For values of σ within 1-5% of the mean

synaptic conductance, the stochastic current has no effect on network propagation. For

larger values of σ, the desired network activity was disrupted or halted.

<p style="text-align: center;">**Results**</p>

Adult zebra finches generate intricate songs composed of sequences of distinct song elements, each characterized by a stereotypical acoustic pattern across every rendition of song. The neural circuitry that governs this behavior consists of HVC_{RA} neuronal population where each emits a single and stereotyped 6-10 ms burst during each rendition of song, HVC_X neurons eliciting 1-4 bursts that are similarly time locked to vocalizations, and HVC_{INT} neurons that tend to burst densely throughout song. Intrinsic and synaptic mechanisms that orchestrate these neurons' behaviors are well known (Daou et al., 2013; Kornfeld et al., 2017; Mooney & Prather, 2005). We next describe the steps of building our biophysical network model to describe this ongoing behavior in the following order: 1) tuning the synaptic parameters to fit the dual-intracellular recording traces collected experimentally by Mooney and Prather (2005) as well as Kosche et al (2015), and then 2) describing the network components that are essential for the patterned output of the system, as well as the internal dynamics of the network that govern the strength and duration of individual bursts, the duration of silent gaps between bursts, sparseness versus tonicity, the interplay between excitation and inhibition, role of intrinsic properties and so on that explain how the firing activity of the three classes of HVC neurons propagate through the network in sequential manner.

Tuning synaptic parameters:

We initiated our HVC network modeling study calibrating the synaptic parameters (excitatory and inhibitory currents' activation/inactivation constants, etc..) by reproducing the voltage traces elicited by the dual intracellular recordings from identified pairs of HVC neurons in brain slices conducted by Mooney and Prather (2005). While we are using off-

the-shelf synaptic currents from the literature (Destexhe et al., 1994; Varela et al., 1997),

we needed to make sure that the synaptic parameters used can replicate the dual

synaptic connectivity patterns (strengths of excitation/inhibition, magnitudes of voltage

deflections and other trace morphologies). Mooney and Prather's findings revealed robust

disynaptic feedforward inhibition from HVC_{RA} to HVC_X neurons (mediated by HVC_{INT}

neurons), potent monosynaptic excitation from HVC_{RA} and HVC_X to HVC_{INT} neurons (via

NMDA and AMPA currents), and substantial monosynaptic inhibition from HVC_{INT}

neurons to HVC_{RA} and HVC_X (via GABA currents).

Figure 3 displays the dual intracellular recordings conducted by Mooney and Prather (left

column) as well as the mathematical model replications (right column) after the synaptic

parameters' calibration. DC-evoked action potentials in HVC_{RA} neurons trigger inhibitory

postsynaptic potentials (iPSPs) in HVC_X neurons (Fig. 3-A1), as well as fast depolarizing

postsynaptic potentials (dPSP) in HVC_{INT} neurons (Fig. 3-B1). To replicate the effects of

stimulating HVC_{RA} neurons onto the other two classes of HVC neurons, we connected

one HVC_{RA} neuron to excite one HVC_{INT} neuron via an AMPA current. The HVC_{INT} neuron

in its turn was connected with one HVC_X neuron via a GABA current, thereby making a

di-synaptic pathway from HVC_{RA} to HVC_X. DC-evoked action potentials in the model

HVC_{RA} neuron (brief ~10 ms depolarizing current pulses (0.5 nA, similar to what Mooney

and Prather applied to HVC_{RA} neurons) evoked a fast-depolarizing postsynaptic potential

(dPSP) in the corresponding model HVC_{INT} neuron (Fig. 3-B2) as well as inhibitory

postsynaptic potential (iPSP) in the corresponding model HVC_X neuron (Fig. 3-A2)

mediated via HVC_{INT}. Similarly, DC-evoked action potentials in HVC_{INT} neurons generate

fast iPSPs in HVC_X neurons (Figures 3-C1, D1). The unequal magnitude of the four sags

elicited in the HVC_X neuron (due to the four brief stimuli) and the large sag after the stimuli

ends (Fig. 3-C1), as well as the jagged long sag in response to the repetitive action

potentials elicited by the HVC_{INT} neuron (Fig. 3-D1) is probably due to the fact that the

corresponding HVC_X neurons are receiving multiple synaptic inputs from neurons other

than the neurons being stimulated by Mooney and Prather, which adds to the underlying

nonlinearities of the responses being recorded. Model HVC_{INT} and HVC_X neurons were

connected and synaptic parameters were calibrated to generate similar waveforms by

giving brief ~10 ms depolarizing current pulses of 0.5 nA to HVC_{INT} (Fig. 3-C2), or giving

a DC-pulse of 0.5 nA for 500 ms (Fig. 3-D2). In the model, the sag seen in the HVC_X

neuron response is due to the build-up of the H-current as the model HVC_{INT} neurons

continues firing, exerting its inhibition onto the HVC_X neuron. Finally, model HVC_X-HVC_{INT}

monosynaptic connectivity (Fig. 3-E2) was calibrated to match the experimental findings

(Fig. 3-E1).

Mooney & Prather 2005   Model Simulations

A1

A2

B1

B2

C1

C2

D1

D2

E1

E2

*Figure 3. Model output compared to experimental results obtained by Mooney and Prather (2005). DC-evoked action potentials in HVC_{RA} neurons trigger iPSPs in HVC_X neurons (**A1**) as well as fast dPSPs in HVC_{INT} neurons (**B1**). Brief (~10 ms) depolarizing current pulses (0.5 nA) applied to model HVC_{RA} neuron (same values used as by Mooney and Prather (2005)) evokes similar responses in the corresponding model HVC_X (**A2**) and model HVC_{INT} (**B2**) neurons. DC-evoked action potentials in HVC_{INT} neurons generate fast iPSPs in HVC_X neurons (**C1, D1**). Similar responses were elicited in model HVC_X neurons when model HVC_{INT} neuron was stimulated by brief (10 ms) depolarizing pulses (0.5nA) (**C2**) or when it was given a DC-pulse of 0.5 nA for 500 ms (**D2**). Finally, HVC_{INT} neurons elicit fast dPSPs when HVC_X neurons are injected with 10 ms pulses of 0.5 nA current (**E1**), which was simulated in the model (**E2**). In this and subsequent figures (unless otherwise specified), HVC_X neurons' traces are represented in blue, HVC_{RA} neurons in red, and HVC_{INT} neurons in black. Panels in the left column adopted from Mooney and Prather (2005).*

Network Architecture: Non-sequential random sampling in HVC

We next describe the activity patterns generated by our network model, and explain how the firing activity of the three classes of HVC neurons propagate through the network in sequential bursts of activity. The network developed is composed of chains of HVC subnetworks or microcircuits, each with its own intrinsic dynamics. The microcircuit represents a basic architectural unit that encodes for a syllable or a sub-syllabic segment (SSS) in the motif (Fig. 4). Each neuron in a microcircuit is representative of a neural population. In other words, a model neuron (belonging to any class) firing is representative of a population of that neuronal class firing, which could be many neurons of the same class exhibiting very similar intrinsic and synaptic properties leading to their firing at the same time. We refer here to "microcircuits" in a more functional sense, rather than rigid, isolated spatial divisions (Cannon et al. 2015). A microcircuit in our model reflects the local rules that govern the interaction between all HVC neuron classes within the broader network, and that are essential for proper activity propagation. The number of microcircuits in the chain determines the number of SSSs in the motif. We envision the HVC to be composed of many copies of such microcircuit chains that are associated with

SSSs with roughly synchronized activity. The duration of the sub-syllabic segments need

not be the same; therefore, the number of neurons that each microcircuit encompasses

need not be equal as we will see next. Moreover, while silent gaps are integral to the

overall process of song production, we have not elaborated on them in this model due to

the lack of a clear, biophysically grounded representation for the gaps themselves at the

level of HVC. Our primary focus here is on modeling the active, syllable-producing phases

of the song, where the HVC network's sequential dynamics are critical for song. However,

one can think the encoding of silent gaps via similar mechanisms that encode SSSs,

where each gap is encoded by similar microcircuits comprised of the three classes of

HVC neurons (let's call them GAP, rather than SSS) that are active only during the silent

gaps. In this case, the propagation of sequential activity is carried throughout the GAPs

from the last SSS of the previous syllable to the first SSS of the subsequent syllable.

The network is comprised of a total pool of HVC_{RA} neurons (120 neurons, red circles), a

pool of interneurons (50 neurons, black circles) and a pool of HVC_X neurons (50 neurons,

blue circles), thereby maintaining a 2:1:1 proportionality factor across the populations of

HVC_{RA}: HVC_{INT}: HVC_X as reported earlier (Kornfeld et al., 2017; Wild et al., 2005). The

total number of neurons in the pool is arbitrary and can be made larger, but we limited it

to these values given the huge number of ODEs that are being simulated (~2000 ODEs).

Figure 4-A shows the network diagram illustrating three sample microcircuits encoding

for three sub-syllabic segments (SSS_{i-1}, SSS_i, and SSS_{i+1}) in sequence. Each microcircuit

(enclosed by a gray oval) is made up of a number of HVC_{RA} neurons (red circles) and a

number of HVC_{INT} neurons (black circles), all selected randomly from their corresponding

pools, as we will describe next. In our model, we limited the number of microcircuits to

twenty (i.e., the motif is encoded by twenty sub-syllabic segments), but this number is

also arbitrary and can be made larger or smaller.

***Figure 4.** Cartoon diagram illustrating the network architecture configuration. **A.** Each grey oval represents a*
microcircuit encoding for a sub-syllabic segment (SSS). The number of microcircuits is envisioned to be equal to the
number of SSSs representing the song. Each microcircuit contains a number of HVC_{RA}, HVC_X and HVC_{INT} neurons
*selected randomly from the total pool of neurons (see text). **B.** Within each microcircuit, HVC_{RA} neurons are connected*
to each other in a chain-like mode and they send excitatory afferents to HVC_{INT} neurons in the same and other
microcircuits, selected randomly. HVC_{INT} neurons send GABAergic synapses onto HVC_X neurons in the same
microcircuit only as well as to HVC_{RA} neurons in any other random microcircuit except the microcircuit they belong
to. Finally, HVC_X neurons send excitatory afferents to HVC_{INT} neurons in the same microcircuit. Activity starts by a

 small DC pulse to the first HVC$_{RA}$ neuron in the first microcircuit. Activity propagates from one microcircuit to

another by excitatory coupling between the last HVC$_{RA}$ neuron in microcircuit i and the first HVC$_{RA}$ neuron in

*microcircuit i+1. **C.** During singing, the propagation of activity unfolds across the chain of microcircuits, such that*

neurons belonging to microcircuit x gets activated and encode for SSS$_x$.

Network organization: The total pool of HVC$_{RA}$ neurons is comprised of smaller groups

of HVC$_{RA}$ neurons, the number in each group of which is chosen randomly from the pool

(red background circles in each gray oval). Each group of HVC$_{RA}$ neurons belongs to a

unique microcircuit and no HVC$_{RA}$ neuron is allowed to be part of more than one

microcircuit for reasons described next. Since we set the motif to be represented by

twenty microcircuits in our illustration, HVC$_{RA}$ neurons were recruited to their

corresponding microcircuits randomly, with each microcircuit allowing a random number

(3 – 10) neurons of the RA-projectors to belong to it. In addition to that, the numbers of

HVC$_{INT}$ neurons (black background circles) that a microcircuit exhibit is a random number

between 1 and 4, as well as the number of HVC$_X$ neurons (blue background circles) in

each microcircuit is random between 1 and 4, where the individual neurons are selected

arbitrarily one neuron at a time from their corresponding pools. Each HVC$_{INT}$ neuron can

belong to a single microcircuit and similarly each HVC$_X$ neuron can belong to a single

microcircuit for reasons described next.

Synaptic Connectivity: Within each microcircuit, HVC$_{RA}$ neurons are selected randomly,

one after the other, to send AMPA excitatory synapses to each other in a chain-like mode.

Specifically, if there are m HVC$_{RA}$ neurons recruited to belong to microcircuit i (where m

is the random number generated between 3 and 10 in this case), a neuron from the m is

first selected randomly and designated as the first neuron in the chain ($HVC_{RA_1}^i$). After

that, a second neuron ($HVC^i_{RA_2}$) from the remaining $m-1$ is selected randomly and an

AMPA synapse is connected from $HVC^i_{RA_1}$ to $HVC^i_{RA_2}$. Similarly, a third neuron ($HVC^i_{RA_3}$)

is selected randomly from the remaining $m-2$ neurons and an AMPA synapse is

connected from $HVC^i_{RA_2}$ to $HVC^i_{RA_3}$, and so on, until the HVC$_{RA}$ neurons in every

microcircuit are connected together (Fig. 4A, small red circles in each microcircuit).

Each HVC$_{INT}$ in a microcircuit was assigned a random number (between 3 and 8) of

excitatory AMPA connections from the HVC$_{RA}$ neurons in the same microcircuit it belongs

to, as well as from HVC$_{RA}$ neurons in the other microcircuits (Fig. 4B). In their turn, each

HVC$_{INT}$ neuron sends a random number (between 2 and 4) of GABAergic inputs to HVC$_{RA}$

neurons, chosen arbitrarily from any microcircuit in the chain except the microcircuit that

the HVC$_{INT}$ neuron belongs to, due to the following reason: if HVC$_{INT}$ inhibits HVC$_{RA}$ in

the same microcircuit, some of the HVC$_{RA}$ bursts in the microcircuit might be silenced by

the dense and strong HVC$_{INT}$ inhibition breaking the chain of activity. However, if HVC$_{INT}$

inhibits HVC$_{RA}$ in any other microcircuit, activity is ensured to propagate because the

HVC$_{INT}$ inhibition of the corresponding HVC$_{RA}$ would arrive at times that are not the

"assigned" times of the HVC$_{RA}$ to elicit their ultra-sparse code (Hahnloser et al., 2002).

Unlike HVC$_{RA}$, HVC$_{INT}$ neurons belonging to a particular microcircuit can burst at times

other than the moments when the corresponding encoded SSS is being "sung"; however,

we chose to house interneurons within microcircuits for the mere fact that any given

interneuron cannot inhibit any given HVC$_{RA}$ neuron, rather there are some "rules of

engagement" where we ensure that no inhibition arrives to any HVC$_{RA}$ neuron while it's

eliciting its burst of activity (Fig. 4B). In other words, what makes a particular interneuron

belong to this microcircuit or the other is merely the fact that it cannot inhibit HVC_{RA}

neurons that are housed in the microcircuit it belongs to for the reasons described. In this

regard, this arrangement is similar to the Cannon et al (2015) model in the context of

structured inhibition amid the ongoing feedforward excitation.

At the HVC_X side, each X-projecting neuron excites via AMPA currents a random

selection (1 - 3) of HVC_{INT} neurons that belong to the same microcircuit, and in their turn,

each HVC_{INT} neuron inhibits via GABA synapses a random selection (1 - 2) of HVC_X

neurons in the same microcircuit (Fig. 4B). These numbers are again arbitrary, but we

limited the number of connections from HVC_{INT} to HVC_X due to that fact that X-projecting

neurons in our model fire upon rebound from inhibition, and the more inhibitory inputs

they receive, the more rebound bursts they elicit, which is not realistic since HVC_X

neurons are known to elicit 1 - 4 bursts during singing (Fujimoto et al., 2011; Kozhevnikov

& Fee, 2007), which is what we achieved with this number of synapses. HVC_X neurons

were selected to be housed within microcircuits and their synapses connecting to

interneurons within the same microcircuit due to the following reason: if an HVC_X neuron

belonging to microcircuit i sends excitatory input to an HVC_{INT} neuron in microcircuit j,

and that interneuron happens to select an HVC_{RA} neuron from microcircuit i as its afferent

inhibitory connection (via random sampling), then the propagation of sequential activity

will halt, and we'll be in a scenario similar to what was described earlier for HVC_{INT}

neurons inhibiting HVC_{RA} neurons in the same microcircuit. Similarly, if an HVC_{INT} neuron

in microcircuit i inhibits an HVC_X neuron in another microcircuit j, and that HVC_X neuron

excites an interneuron that synapses onto an HVC$_{RA}$ from microcircuit i, then sequential

activity might be disrupted.

While HVC$_{RA}$ neurons are connected to each other in each microcircuit in a chain-like

mode as described earlier, the microcircuits interact with each other via the projections

from the last HVC$_{RA}$ in a microcircuit i to the first HVC$_{RA}$ in a following microcircuit $i + 1$.

The network is kick-started by a stochastic DC input to HVC_{RA}^1, that is, only HVC_{RA}^1 receives

input from outside HVC. The propagation of sequential activity along with the realistic

firing of the three classes of HVC neurons is maintained and orchestrated by HVC's

intrinsic and synaptic processes without relying on extrinsic inputs as shown next

(Figures. 5-11).

Sequential propagation of HVC$_{RA}$ activity

The activity patterns that RA-projecting neurons of this network display are illustrated in

Figure 5. HVC$_{RA}$ neurons burst extremely sparsely generating at most a single burst per

simulation (song motif) and with different HVC$_{RA}$ neurons bursting at different times in the

song. HVC$_{RA}$ bursts had a duration of 8.12 ± 0.89 ms, and comprised of 4.76 ± 0.48

spikes. The inset of Fig. 5 shows the delay from the onset of the first spike in an HVC$_{RA}$

neuron to the onset of the first spike in the next HVC$_{RA}$ it synapses to. This duration is

controlled by two factors: 1) the magnitude of the AMPA conductance, with larger

magnitudes corresponding to shorter delays and vice versa (Fig. 6A). In short, the faster

the AMPA current is (modeled as larger magnitude in the g_{AMPA} parameter that connects

two HVC$_{RA}$ neurons), the shorter the delay between the successive HVC$_{RA}$ bursts. 2) the

magnitude of the A-type K$^+$ current conductance (g_A) with larger magnitudes

corresponding to longer delays (Fig. 6B). This conductance increases rapidly on

depolarization due to I_A's fast activation. The rapid increase in g_A halts after a few

milliseconds and switches to a slow decrease that is due to the slow inactivation that this

current exhibit. The slow decrease is reflected in the voltage trace as a slow

depolarization in the membrane potential (encoding for the delay to spiking), and this

allows the model HVC$_{RA}$ neuron to eventually escape the inhibition produced by I_A and

fire its delayed burst.

Figure 5. Spiking patterns of 120 HVC$_{RA}$ neurons (labeled with numbers) showing the propagation of sequential activity. The neural traces are aligned by the acoustic elements of a spectrogram from an exemplar bird's song illustrating the firing of HVC$_{RA}$ neurons with respect to ongoing part of a song. The inset shows a zoomed version of two subsequent HVC$_{RA}$ neurons firing patterns illustrating the delay between their individual bursts.

The number of spikes in HVC$_{RA}$ neurons and the strength of the burst is controlled by three factors: 1) the strength of the AMPA conductance itself where stronger excitatory coupling corresponds to larger number of spikes (Fig. 6A), 2) the A-type K$^+$ conductance where larger magnitudes of the conductance reduce the general excitability of the HVC$_{RA}$ neuron and limit its number of spikes (Fig. 6B), and 3) the interplay between the L-type

Ca²⁺ (g_{CaL}) and the Ca²⁺-dependent K⁺ (g_{SK}) conductances that control the strong

adaptation which these neurons exhibit (Daou et al., 2013). In short, intracellular Ca²⁺

(due to I_{CaL}) accumulates during the HVC$_{RA}$ burst. This result in a buildup of Ca²⁺-

activated K⁺ current (I_{SK}) that terminates the HVC$_{RA}$ burst, which in turn terminates any

burst in a post-synaptic neuron since HVC$_{RA}$ can no longer provide excitation.

Figure 6. *Effects of the AMPA synaptic conductance and A-type K⁺ conductance on the delay between two successive*

HVC$_{RA}$ bursts. **A.** *Presynaptic model HVC$_{RA}$ neuron (top, black) is connected to a postsynaptic model HVC$_{RA}$ neuron*

and the corresponding AMPA excitatory conductance (g_{AMPA}) was increased from 10 nS (bottom, red) to 12 nS

(bottom, blue), while keeping all other parameters fixed. Increasing g_{AMPA} reduce the delay between the peaks of the

pre- and post- HVC$_{RA}$'s first spikes and increase the number of spikes in the postsynaptic neuron. **B.** *Larger*

magnitudes of the A-type K⁺ conductance (g_A) leads to longer delays to spiking. While keeping all intrinsic and

synaptic parameters fixed ($g_{AMPA} = 10$), increasing g_A from 10 nS (bottom, red) to 13 nS (bottom, blue) delayed the

onset to spiking and reduced the number of spikes. Bars on the top shows the duration in ms between the peak of first

action potential in the presynaptic neuron to the peak of the first action potential in the postsynaptic neuron.

The propagation of sequential bursting in HVC$_{RA}$ neurons halts and the chain of HVC$_{RA}$

activity is broken if any of the following is satisfied: 1) if an AMPA conductance for any of

the HVC$_{RA}$'s that connects it to the next HVC$_{RA}$ in its chain is small enough such that it's

not able to elicit sufficient excitability on the postsynaptic side, 2) if a g_A conductance (Fig.

7A) or a g_{SK} conductance (Fig. 7B) in any of the HVC$_{RA}$'s is large enough (mimicking an

up-regulation in the corresponding channel) to eliminate the corresponding HVC$_{RA}$'s

burst. Therefore, in order to generate accurate HVC$_{RA}$ bursting patterns that maintain the

sequential propagation of neural activity, accurate number of spikes and delays between

spikes across the population of HVC$_{RA}$'s in all microcircuits, as well as the general intrinsic

properties of the HVC$_{RA}$ neurons' spike morphologies (see Methods), both intrinsic and

synaptic constraints are needed with key roles in this process for the A-type K$^+$ and the

Ca^{2+}-dependent K$^+$ currents that HVC$_{RA}$ model neurons exhibit, as well as the AMPA

currents connecting the HVC$_{RA}$ population together within and across microcircuits.

Recall that we envision each HVC neuron of any class in our model as a representative

of a neural population of the same class that exhibits the same intrinsic as well as afferent

and efferent synaptic connectivity. Therefore, in Fig. 7 and the subsequent figures (10,12

and 13) where we show disruption of sequential activity due to changes in synaptic or

intrinsic properties of HVC neurons, the modeled synaptic/intrinsic changes to the

selected neurons shown are envisioned to be changes applied to the whole neural

population encoded by our model neuron. In other words, disrupting the properties of a

single neuron within that neural population will not cause harm to the propagation of

activity due to what could be thought of as homeostatic mechanisms of the network

(Golowasch et al., 1999; Marder & Goaillard, 2006; Williams et al., 2013). This

redundancy within the population allows the propagation of activity to be maintained. It is

an important feature of our model and is consistent with biological observations where

neural populations exhibit robust collective behavior and the loss of a single neuron does

not result in a major disruption of network activity.

Figure 7. Intrinsic changes in HVC_{RA} halts the propagation of sequential activity. Up-regulating the A-type K^+ current

(A) or the Ca^{2+} - dependent K^+ current (B) in exemplar neurons (HVC_{RA}^{60}, A) or (HVC_{RA}^{80}, B), by increasing g_A 10-fold

(A), or g_{SK} from 15-fold nS (B), reduces the excitability of corresponding HVC_{RA} neuron markedly, eliminating its

corresponding burst and breaking the sequence.

Bursting patterns of HVC$_{INT}$ and HVC$_X$ neurons

The activity patterns that HVC$_{INT}$ neurons exhibit in our network architecture are illustrated

in Figure 8A (shown here for 10 HVC$_{INT}$ neurons). In contrast to the sparse activity in RA-

projecting neurons, HVC interneurons generate multiple bursts and spike densely as

reported during the song (Hahnloser et al., 2002). The number of bursts each HVC_{INT}

neuron exhibits as well as the strength of each burst is controlled by network and synaptic

mechanisms as described next (Figures. 9, 10). HVC_X neurons in their turn generate 1 -

4 bursts per song motif (Fig. 8B, shown here for 10 HVC_X neurons) similar to experimental

results (Fujimoto et al., 2011; Hahnloser et al., 2002). Similarly, the number of bursts and

strength of each HVC_X burst depends on a set of synaptic and intrinsic mechanisms,

illustrated in Figures 11 and 12. HVC_X neurons differed in the number of bursts and the

number of spikes per burst rendering the results more realistic.

Figure 8. *Activity patterns for 10 HVC$_{INT}$ and 10 HVC$_X$ neurons are illustrated. **A**. HVC interneurons displays dense spiking and bursting throughout the song, due to the dense HVC$_{RA}$ – HVC$_{INT}$ and HVC$_X$ – HVC$_{INT}$ excitatory coupling (Fig. 4). **B**. HVC$_X$ neurons display 2-4 rebound bursts that vary in their strength and duration due to HVC$_{INT}$ – HVC$_X$ inhibitory coupling as well as intrinsic properties (Fig. 9).*

Dense bursting in HVC$_{INT}$

The random connections from HVC$_{RA}$ and HVC$_X$ neurons to HVC$_{INT}$ as well as the multiple

bursts HVC$_{INT}$ and HVC$_X$ exhibit in this network are illustrated in Figure 9 by highlighting

a sample interneuron (HVC$_{INT}^{10}$). Here the firing patterns of HVC$_{INT}^{10}$ in addition to all HVC$_{RA}$

and HVC$_X$ neurons that connect to it are shown. HVC$_{INT}^{10}$ received random synaptic inputs

from HVC$_{RA}^{11}$, HVC$_{RA}^{32}$, HVC$_{RA}^{83}$ and HVC$_{RA}^{120}$ neurons, as well as from X-projecting neurons

HVC$_X^7$, HVC$_X^8$ and HVC$_X^9$ in the same microcircuit it belongs to. Notice that each HVC$_{RA}$

neuron (which happen to belong to separate microcircuits in this case), bursts only once

as reported experimentally (Hahnloser et al., 2002). HVC$_{INT}^{10}$ generates multiple bursts as

well as spike sparsely. Each of the HVC$_{INT}^{10}$ bursts are aligned with one of the bursts in

HVC$_{RA}$ and/or HVC$_X$ neurons due to excitatory coupling.

The number of bursts an HVC$_{INT}$ neuron exhibits is largely determined by the number of

bursts that each HVC$_{RA}$ and each HVC$_X$ neuron that synapses onto that interneuron

exhibits. In other words, the larger the number of bursts that a population of HVC$_{RA}$ and

HVC$_X$ that connects to an interneuron exhibit, the larger the number of bursts elicited in

the interneuron. The strength of the bursts in HVC$_{INT}$ neurons is determined by two

factors: 1) the strength of the excitatory synaptic conductance from HVC$_{RA}$ and/or HVC$_X$

to HVC$_{INT}$, with stronger bursts in HVC$_{INT}$ corresponding to larger magnitudes of g_{AMPA}

from HVC$_{RA}$ and/or HVC$_X$ onto the interneuron, 2) the number of spikes/bursts in HVC$_{RA}$

and HVC$_X$ that are aligned and occur simultaneously at the HVC$_{INT}$ synapses. For

example, the asterisk shown in Figure 9 under the HVC$_{INT}^{10}$ spike train highlights a stronger

burst (compared to the rest of the spiking pattern) because HVC$_{RA}^{83}$, HVC$_X^7$, HVC$_X^8$ and HVC$_X^9$

neurons tended to fire/burst at the same moment (or within a close interval of each other),

thereby generating a stronger response in HVC_{INT}^{10}. Therefore, the characteristic tonic

activity that HVC interneurons exhibit with bursting and suppression at different locations

during singing is explained in our model by the excitatory and inhibitory coupling between

the interneurons and both classes of projecting neurons.

Figure 9. *Patterned activity of HVC interneurons illustrated for an exemplar interneuron (HVC_{INT}^{10}). For this neuron,*

HVC_{RA}^{11}, HVC_{RA}^{32}, HVC_{RA}^{83}, HVC_{RA}^{120}, HVC_{X}^{7}, HVC_{X}^{8} and HVC_{X}^{9} were selected randomly from the pool of HVC_{RA}'s and

HVC_{X}'s to form excitatory coupling. The number of bursts in HVC_{INT}^{10} is controlled by the number of bursts that each

of the HVC_{RA} and HVC_{X} neurons that connect to it exhibit. The strength of each of the HVC_{INT}^{10} bursts depend on the

magnitude of g_{AMPA} from the corresponding neuron(s) they cause it as well as the simultaneous bursting of any of the

projecting neurons. For example, the asterisk (*) shows a region of dense firing in HVC_{INT}^{10} because HVC_{RA}^{83}, HVC_{X}^{7},

HVC_{X}^{8} and HVC_{X}^{9} neurons elicit their spikes at similar times causing a potentiated response in HVC_{INT}^{10}. HVC_{X} neurons

exhibit multiple sags and rebounds because they're receiving inhibition from several interneurons (not shown here).

The inhibitory effect of HVC$_{INT}$ neurons, their interplay with the excitatory projection

neurons, and their intrinsic properties plays a key role in the modulation of sequence

propagation and overall desired network behavior. In particular, if the excitation from the

projection neurons onto the interneurons was very large (>> g_{AMPA}), then HVC$_{INT}$ neurons

enter regimes of very dense and continuous bursting/spiking (above their natural and

basic, yet already enhanced potentiation), thereby leading to the inhibition of the HVC$_{RA}$'s,

halting the sequence. Similarly, if the GABAergic conductances from HVC$_{INT}$ to HVC$_{RA}$

was relatively strong, outside their ideal ranges (Figure 2-B), then HVC$_{RA}$ neurons won't

be able to elicit their bursts.

Besides the synaptic modulations of HVC$_{INT}$ neurons on network activity, intrinsic

mechanisms orchestrated primarily by the T-type Ca^{2+} current and the hyperpolarization-

activated inward current need to be tightly regulated to ensure desired network activity.

The T-type calcium channel opens near resting membrane potential and markedly

influence neuronal excitability (Huguenard, 1996; Jagodic et al., 2008). In our model, if

the T-type Ca^{2+} current in an HVC$_{INT}$ neuron is up-regulated (due to a depolarizing shift

in its voltage-dependent inactivation or simply setting g_{CaT} to a relatively large value), then

the interneuron will fire with much larger firing frequency, silencing the corresponding

HVC$_{RA}$ and HVC$_X$ neurons that it connects to and breaking the sequence (Fig. 10A).

Similarly, the H-channels regulate the resting membrane potentials of the neurons they're

expressed in and play a key role in regulating the spontaneous firing activity (Datunashvili

et al., 2018; Funahashi et al., 2003; Yao et al., 2003). In our model, if the H conductance

of an HVC$_{INT}$ was upregulated (by increasing the magnitude of g_h), then the neuron

switches to continuous firing, silencing the HVC$_{RA}$'s that is connects to (Fig. 10B).

Therefore, both the interplay between excitation and inhibition between HVC$_{INT}$ and HVC

projection neurons as well as the intrinsic properties of HVC$_{INT}$ neurons (I_{CaT} and I_H) are

necessary elements to ensure an accurate propagation of sequential activity.

Figure 10. *Intrinsic changes in HVC$_{INT}$ halts the propagation of sequential activity. Up-regulating the T-type Ca^{2+}*

*current conductance (**A**) or the hyperpolarization-activated inward current conductance (**B**) in exemplar HVC$_{INT}$*

neurons eliminates sequence propagation. Increasing g_{CaT} in HVC$_{INT}^{23}$ 10-fold results in dense bursting and firing in

HVC$_{INT}^{23}$, which in its turn blocks the bursting of HVC$_{RA}^{45}$ due to the inhibitory GABA coupling between HVC$_{INT}^{23}$ and

*HVC$_{RA}^{45}$ (**A**). Similarly, increasing g_H in HVC$_{INT}^{41}$ 10-fold results in increased firing in HVC$_{INT}^{41}$, which in its turn blocks*

*the bursting of HVC$_{RA}^{76}$ due to the inhibitory GABA coupling between HVC$_{INT}^{41}$ and HVC$_{RA}^{76}$ (**B**). Sequence of HVC$_{RA}$*

bursts truncated at the level of HVC$_{RA}^{80}$ for better visualization purposes.

Rebound bursting in HVC$_X$

The activity patterns that HVC$_X$ neurons exhibit in the network are illustrated in Figure 8B (shown here for 10 HVC$_X$ neurons). The X-projecting neuronal firing is characterized by regions of inhibition throughout singing, interrupted by occasional rebound bursting. The strength of each burst and the number of total bursts in an HVC$_X$ depend on synaptic and intrinsic mechanisms summarized briefly as such: 1) the degree of inhibition from HVC$_{INT}$ neurons, 2) the number and timing of bursts of HVC$_{INT}$ neurons and 3) the intrinsic properties and magnitudes of the T - type Ca^{2+} and H – currents that the HVC$_X$ neurons exhibit. In short, the stronger the GABAergic maximal conductance(s) from the HVC$_{INT}$ neurons that inhibit a given HVC$_X$ neuron, the stronger the HVC$_X$ neuron's rebound burst (Figures 8-11). Similarly, if multiple HVC$_{INT}$ neurons' bursts that inhibit an HVC$_X$ neuron aligned simultaneously, then the rebound in HVC$_X$ is potentiated due to the stronger inhibition. This is primarily due to the T-type Ca^{2+} current and the H-current that HVC$_X$ neurons exhibit, facilitating rapid calcium influx into the neurons when they rebound from hyperpolarization. The calcium influx is correlated to the degree and the duration of inhibition that the neuron receives, and can trigger a potentiated burst of action potentials leading to more robust rebound responses. In other words, the stronger the inhibition of HVC$_X$, the stronger the activation of I_{CaT} and I_h , leading to stronger rebounds. Moreover, the number of spikes in each HVC$_X$ burst is controlled by several factors including 1) the degree of inhibition from HVC$_{INT}$ and its effect on I_{CaT} and I_h as described earlier, and 2) the strength of the Ca^{2+}-dependent K$^+$ conductance (g_{SK}) with stronger magnitudes of this conductance dampening the excitability of these neurons and reducing the number of spikes.

The interplay between HVC_{INT} and HVC_X in shaping the characteristic HVC_X responses

is illustrated in Figure 11 from exemplar neurons in the network. Two interneurons, HVC_{INT}^5

(black) and HVC_{INT}^6 (green) were selected randomly to inhibit HVC_X^3 (blue trace, Fig. 11A).

There are several regions in the HVC_X^3 membrane potential trajectory where the neuron is

inhibited (sags/sinks in the voltage trace). These regions correspond to the moments in

time when either HVC_{INT}^5 or HVC_{INT}^6 neurons, or both simultaneously, are bursting, thereby

silencing HVC_X^3. Eventually, HVC_X^3 is able to escape inhibition at a few instances in time

and generate multiple post inhibitory rebound bursts mediated by I_{CaT} and I_h . Figure 11B

shows a zoomed version of panel A illustrating the three rebound bursts in HVC_X^1 as a

result of escape from inhibition. HVC_{INT}^5 and HVC_{INT}^6 neurons generated multiple

successive bursts of firing, during which HVC_X^3 cannot escape the inhibition. There are

only a few intervals where HVC_X^3 is able to elicit rebound bursts; the first opportunity was

at the beginning of the trace (Fig. 11A) where the sag generated in HVC_X^3 as a result of

inhibition from HVC_{INT}^5 is clear, but the inhibition from HVC_{INT}^5 was not strong enough to

elicit a rebound burst in HVC_X^3, rather eliciting only a rebound depolarization. The

subsequent three opportunities where HVC_X^3 is able to escape the inhibition all elicited

rebound bursts due to the strong inhibitory input arriving from both interneurons.

All rebound bursts in HVC_X are orchestrated by the hyperpolarization-activated inward

current (I_H) and the T-type Ca^{2+} current (I_{CaT}). The longer the duration of the inhibition

prior to a rebound, the longer the time I_H and I_{CaT} take to fully activate and generate the

rebound burst. The shorter the inhibitory silent interval (that is, the interval in which HVC_X

is receiving no inhibition), the shorter the rebound and the fewer the number of spikes in

the rebound burst (Fig. 11A, first rebound burst). Moreover, HVC interneurons inhibit

multiple HVC$_X$ neurons in our network. Figure 11C shows an exemplar interneuron

(HVC_{INT}^5, black) that happens to inhibit two X-projecting neurons (HVC_X^3, blue and HVC_X^5,

orange). Each burst in HVC_{INT}^5 elicits an iPSP in HVC_X^3 and HVC_X^5, and can contribute to

subsequent rebound bursts in them unless silenced by other interneurons that connect to

HVC_X^3 or HVC_X^5 (Fig. 11D). All HVC$_X$ neurons differed in their corresponding number of

rebound bursts as well as in the number of spikes per burst rendering the results

biologically accurate.

Figure 11. *Activity patterns illustrating the interplay between HVC interneurons and X-projecting neurons. HVC_X^3*

(blue) is an exemplar projecting neuron that receives inhibition from HVC_{INT}^5 (black) and HVC_{INT}^6 (green) due to the

*random inhibitory coupling (**A**). HVC_X^3 is inhibited whenever HVC_{INT}^5 and HVC_{INT}^6 are firing, eventually escaping*

inhibition at some intervals and eliciting rebound bursts due to the activation of I_H and I_{CaT} . HVC interneurons can

*inhibit multiple HVC_X neurons. HVC_{INT}^5 is an exemplar from the network that inhibits HVC_X^3 and HVC_X^5 (orange) (**C**).*

Bursts in HVC_{INT}^5 elicit subsequent bursts in HVC_X^3 and HVC_X^5 unless silenced by other HVC_{INT} neurons that connect

*to them. Zoomed versions of (**A**) and (**C**) are show in (**B**) and (**D**).*

As mentioned earlier, the T-type Ca^{2+} and the H- conductances play a significant role in

modulating the rebound bursting in HVC_X neurons. Due to this interplay, we do not need

significant inhibition to generate rebound bursts, because the T-type Ca^{2+} current's

conductance can be stronger than the inhibitory conductance, leading to robust rebound

bursting, even when the degree of inhibition is not very strong. As a matter of fact, these

two conductances, along with the Ca^{2+} - dependent K^+ conductance can halt the

sequential propagation of activity and mess up the overall desired network behavior as

described next. Ramping up g_H in HVC_X neurons to values outside the allowed ranges

(Figure 2-A) will break the sequence of propagation and generate nonrealistic firing

patterns. The g_H parameter was increased in Figure 12A in an exemplar HVC_X neuron

(HVC_X^{36}) to large values, driving HVC_X^{36} into regimes of runaway excitation due to the up-

regulation of I_H and generating a non-realistic firing pattern for a typical X-projecting

neuron during singing. The reason the sequential propagation of HVC_{RA} is halted is

because HVC_X^{36} happens to have an excitatory coupling with HVC_{INT}^{34} (along with other

interneurons, but we illustrate it here for HVC_{INT}^{34}), and as a result of HVC_X^{36}'s increased

firing, HVC_{INT}^{34} generates a mostly-continuous firing trace of bursting and spiking, which

silences the HVC_{RA} neuron that it happens to inhibit (in this case, HVC_{RA}^{70}, Fig. 12A).

*Figure 12. Intrinsic changes in HVC_X halts the propagation of sequential activity. **A**. Up-regulating the*

hyperpolarization-activated inward current conductance in a sample HVC_X neuron (HVC_x^{36}, by increasing its g_H 10-

fold) leads to increased firing in all HVC_{INT} neurons it connects to (for example, HVC_{INT}^{34}), which in its turn inhibits

all HVC_{RA} neurons it connects to (for example, HVC_{RA}^{70}, being first in the pool that it inhibits) breaking the sequence

*at the level of HVC_{RA}^{70}. **B**. Up-regulating the T-type Ca^{2+} current conductance in a sample HVC_X neuron (HVC_x^{12}, by*

increasing its g_{CaT} 15-fold) leads to stronger rebound bursts in HVC_x^{12} which leads to increased firing in all HVC_{INT}

neurons it connects to (for example, HVC_{INT}^{16}), which in its turn inhibits all HVC_{RA} neurons it connects to (for example,

*HVC_{RA}^{80}, being first in the pool that it inhibits) breaking the sequence at the level of HVC_{RA}^{80}. **C**. Finally, down-regulating*

the Ca^{2+} - dependent K^+ current conductance in a sample HVC_X neuron (HVC_x^{27}, by setting its g_{SK} to zero) leads to

stronger rebound bursts in HVC_x^{27} which leads to increased firing in all HVC_{INT} neurons it connects to (for example,

HVC_{INT}^{30}), which in its turn inhibits all HVC_{RA} neurons it connects to (for example, HVC_{RA}^{15}, being first in the pool that

it inhibits) breaking the sequence at the level of HVC_{RA}^{15}. Sequence of HVC_{RA} bursts truncated at the level of HVC_{RA}^{85}

for better visualization purposes.

Similarly, altering g_{CaT} in HVC$_X$ neurons can have similar effects on network activity.

Figure 12B shows an exemplar HVC$_X$ neuron (HVC_X^{12}) where g_{CaT} was ramped up to large

values and as a result, stronger rebound bursts in HVC_X^{12} were elicited as well as larger

number of rebound spikes in each burst. This is primarily due to the up-regulation of the

T-type Ca^{2+} current that was induced, markedly influencing the neuronal excitability. The

stronger rebounding in HVC_X^{12} generated stronger excitation in their postsynaptic

interneurons' counterparts that they excite (for example, HVC_{INT}^{16}), which in their turn

silenced the HVC$_{RA}$ neurons that they inhibit (for example, HVC_{RA}^{80} is the first neuron in the

chain that HVC_{INT}^{16} inhibits, breaking the sequence at HVC_{RA}^{80}). And finally, the Ca^{2+} -

dependent K^+ conductance, which plays a key role in governing HVC$_X$ neurons'

excitability and their characteristic spike frequency adaptation (Daou et al., 2013) can

have similar effects if the channel was down-regulated (rather than up-regulated as in

I_{CaT} and I). Figure 12C shows the effects of blocking the g_{SK} conductance in an exemplar

HVC$_X$ neuron (HVC_X^{27}) leading to its increased firing, which in turn leads to stronger

bursting and spiking in the HVC$_{INT}$ neurons it excites (for example, HVC_{INT}^{30}). HVC_{INT}^{30} in its

turn generates stronger and wider-range inhibition onto the HVC$_{RA}$ neurons it sends its

axons to (for example, HVC_{RA}^{15} is the first neuron in the chain HVC_{INT}^{30} inhibits, silencing it

and breaking the sequence at HVC_{RA}^{15}).

Finally, we checked the consequences of altering the intrinsic properties of HVC$_X$ neurons

on the network's desired behavior. To do so, we varied the maximal conductances of the

three principal ionic currents of the X-projecting neurons (I_{CaT} , I_{SK}, I_H) across all neurons

of the population, while keeping this variation within the reported ranges shown in Figure

2A (because certainly going outside these ranges will disrupt network activity for other

reasons as reported in Figures 10 and 12). Varying those three key paramters across the

HVCx population had different results. In 100 different simulations that generated random

maximal conductances for I_{CaT}, I_{SK} and I_H, 41% of the simulations did not have any

considerable effect on the desired network activity, whereas 59% resulted in disrupted

network activity, and sometimes detrimental. For example, Figure 13 shows an example

where the sequential propagation of activity was halted and the firing patterns of some

interneurons and X-projecting neurons were rendered non-biophysically realistic.

Particularly, some interneurons switched to continuous spiking or phasic bursting with

little episodic bursting modes, while some HVC_X neurons generated longer rebounds,

fewer number of bursts or fewer number of spikes per burst (Fig. 13 B-C). Hence, changes

in the intrinsic properties of X-projecting neurons can disrupt activity propagation

necessary for song production, and produce biologically unrealistic bursting patterns in

HVC neurons. This can wreak havoc on our network model hinting to the finding that

biophysical parameters are distinct and consistent for an individual bird and this unique

combination is needed for song (Daou & Margoliash, 2020). The homogeneity in the

intrinsic properties of X-projectors might be a strategy allowing it to adapt or respond to

changes in the network.

Figure 13. *Altering the intrinsic properties of HVCx neurons disrupt network activity in 59% of the cases (out of 100 simulations) where the maximal conductances (gCaT, gSK and gH) of HVCx neurons are randomly varied within their allowed ranges (Figure 2A). As a result, some HVC interneurons (B) and X-projecting neurons (C) generated non-biological realistic firing patterns, halting the propagation of sequential activity in RA-projectors (A).*

In conclusion, we developed a detailed and biophysically realistic neural network model for sequence propagation in the HVC of the zebra finch. Our model consisted of chains of microcircuits, each comprised of a selection of HVC_{RA}, HVC_{INT} and HVC_X model neurons selected randomly from a total pool of neurons. The maximal conductances of the four key and principle ionic currents for each model neuron, the number of neurons of each class in any microcircuit, and the excitatory and inhibitory connections between the different classes within a microcircuit and across microcircuits are all selected randomly. This activity propagates throughout the chain of microcircuits causing a sequence of HVC_{RA} bursts while leaving behind realistic bursting patterns for all classes

of HVC neurons as seen during singing. The model incorporates all known ionic and

synaptic currents for each HVC neuron. The network architecture we developed was able

to replicate the *in vivo* biologically realistic firing behavior for each class by including

sparse timely-locked bursting in the RA-projecting neurons (with accurate intrinsic

properties for each burst in terms of number of spikes, duration and spike morphology),

multiple bursting in the X-projecting neurons that are also sparse and time-locked, and

dense bursting/spiking in the interneurons with few intermittent quiescence. The ability of

our network to reproduce the sequential propagation of activity in the presence of

excitatory and inhibitory connections involving all neuronal subclasses as well as over a

range of values for each synaptic and ionic maximal conductances is an indication of its

robustness. Our network unveiled key intrinsic and synaptic mechanisms that modulate

the sequential propagation of neural activity by highlighting important roles for the T-type

Ca^{2+} current and hyperpolarization activated (H) inward current in HVC_X and HVC_{INT}

neurons, Ca^{2+}-dependent K^+ current in HVC_X and HVC_{RA}, A-type K^+ current in HVC_{RA}, as

well as GABAergic and glutamatergic synaptic currents that connects all neuronal

subclasses together. The result is an improved characterization of the HVC network

responsible for song production in the zebra finch. Beyond replicating established HVC

firing patterns, our model provides testable hypotheses that intrinsic membrane

properties, particularly inhibitory timing and rebound bursts maintain robust sequential

propagation. This context generates clear experimental predictions: for example,

modulating I_H or I_{CaT} in HVC_X or HVC_{INT} neurons directly affects rebound spiking and

sequence propagation; or altering I_{SK} or I_A conductances disrupts the ability of HVC_{RA}

neurons to burst at the right time. The model also suggests that structured inhibition can

act as a temporal scaffold for burst timing guiding experiments that manipulate

interneuron dynamics or the overall network inhibition through optogenetics or

pharmacology. Hence, a significant strength of the proposed model is that it outs forward

suggestions for experimental manipulations in the form of targeted experiments whether

using optogenetics or pharmacology that would help validate the model's mechanisms

and further clarify the specific roles different HVC components play in driving sequential

activity.

Discussion

In this study, we developed a biophysically realistic neural network model to explore how intrinsic neuronal properties and local connectivity within the songbird nucleus HVC may support the generation of temporally precise activity sequences associated with zebra finch song. The biophysically realistic network architecture that we designed combine both classes of HVC projection neurons with local inhibitory interneurons. A fundamental goal that we have achieved in our design is a successful replication of the *in vivo* firing behaviors of all the HVC neuronal classes: single sparse timely-precise bursting (3-6 spikes for ~10 ms) in the RA-projecting neurons, multiple bursting (1-4 bursts with 4-9 spikes/burst) in the X-projecting neurons, dense and frequent bursting in the interneurons, as well as the general intrinsic properties that each class of HVC neurons exhibit (Daou et al., 2013; Lewicki, 1996; Long et al., 2010). The patterning activity in HVC is largely shaped in our model by the intrinsic properties of the individual neurons as well as the synaptic properties where excitation and inhibition play a major role in enabling neurons to generate their characteristic bursts during singing.

The three classes of model neurons incorporated to our network as well as the synaptic currents that connect them are based on Hodgkin-Huxley formalisms that contain ion channels and synaptic currents which had been pharmacologically identified (Daou et al., 2013; Kosche et al., 2015; Mooney & Prather, 2005). Our network showed that sequence propagation can be broken if several intrinsic mechanisms are perturbed. In particular, if I_{CaT} or I_H are upregulated in HVC_X or HVC_{INT}, if I_{SK} is downregulated in HVC_X or if I_{SK} is

upregulated in HVC$_{RA}$, then the corresponding chain of activity stops and the rhythmic

activity of the network is disrupted (Figures 7, 10 and 12). Synaptically, perhaps the most

critical role in our network design is played by interneurons which orchestrate the activity

of the two projection neurons in a structured manner. Interneurons adjust the timing of

HVC projection neurons' bursts (Amador et al., 2013; Kosche et al., 2015), and

developmental learning regulates inhibition onto HVC$_{RA}$ (Vallentin et al., 2016). While

findings of Kosche et al. (2015) emphasize the robustness of the HVC timing circuit to

inhibition, our model is more sensitive to inhibition, highlighting that HVC likely operates

with several, redundant mechanisms that overall ensure temporal precision. HVC$_{RA}$

neurons interact with HVC$_X$ through local interneurons, a disynaptic inhibitory pathway

that conveys information to HVC$_X$ neurons (Prather et al., 2008). Focal application of the

GABA$_A$ receptor antagonist, gabazine, restricted inhibitory impact in HVC leading to

stronger and faster responses relative to call onset (Benichov & Vallentin, 2020), showing

that local HVC interneurons form an inhibitory mask that can greatly constrain the spiking

activity of projecting neurons (Kornfeld et al., 2017; Kosche et al., 2015; Mooney &

Prather, 2005) suggesting that HVC model networks that lack inhibitory neurons are

inadequate for explaining sequential propagation of neural activity.

Various models of how the song is encoded within HVC have been proposed. Some

groups suggested that bursting activity propagates through a chain of synaptically

connected HVC$_{RA}$ neurons either as single neurons (Fee et al., 2004; Hahnloser et al.,

2002; Long et al., 2010) or as pools of HVC$_{RA}$ neurons, each group driving a distinct

ensemble of RA neurons (Jin et al., 2007; Leonardo & Fee, 2005; Li & Greenside, 2006).

These models assume that HVC_{RA} neurons generate a continuous, feed-forward

sequence of activity over time, with little or no role played by X-projecting HVC neurons

and interneurons. Other models have incorporated alternative temporal encoding

mechanisms by necessitating synaptic integration at the levels of HVC_{RA} and HVC_{INT}

populations (Drew & Abbott, 2003; Gibb et al., 2009a; Jin, 2009; Weber & Hahnloser,

2007) while yet other approaches gave emphasis to brainstem feedback processes by

incorporating inter-hemispheric coordination to activate sequences of syllable-specific

HVC_{RA} and HVC_{INT} neurons (Galvis et al., 2018; Gibb et al., 2009b). A prominent model

used spatially recurrent excitatory chains and local feedback inhibition to show how the

HVC network stabilize synchrony while propagating sequential activity (Cannon et al.,

2015; Markowitz et al., 2015).

All existing models that describe premotor sequence generation in the HVC either

assume a distributed model (Elmaleh et al., 2021) that dictates that local HVC circuitry is

not sufficient to advance the sequence but rather depends upon moment-to-moment

feedback through Uva (Hamaguchi et al., 2016), or assume models that rely on intrinsic

connections within HVC to propagate sequential activity. In the latter case, some models

assume that HVC is composed of multiple discrete subnetworks that encode individual

song elements (Glaze & Troyer, 2013; Long & Fee, 2008; Wang et al., 2008), but lacks

the local connectivity to link the subnetworks, while other models assume that HVC may

have sufficient information in its intrinsic connections to form a single continuous network

sequence (Long et al., 2010).

The network architecture we developed here exhibits overlap with the various models

presented. First, in agreement with the continuous model, our network architecture

displays a feed-forward mechanism regulating the circuit dynamics e.g., (Gibb et al.,

2009a; Jin, 2009; Jin et al., 2007; Li & Greenside, 2006; Long et al., 2010). Nonetheless,

diverging from a linear progression of HVC neurons directing the song, the network's

structure comprises sequences of microcircuits incorporating all classes of HVC neurons,

where sequential activity transmits from one microcircuit to the next, as opposed to

transitioning directly between individual neurons. Second, in agreement with the

subnetwork models, our model envisions HVC as comprised of multiple discrete

subcircuits (SSSs) where each microcircuit incorporates its own pool of neurons;

however, in our model HVC's connectivity is sufficient to link the microcircuits together

and extrinsic influences are not needed. Moreover, our network is in agreement with the

(Cannon et al., 2015) model where structured inhibition is needed to propagate sequential

activity, synchronize the firing of pools of neurons and stabilize spike timing along the

chain. The pivotal element in advancing sequential activity through time is the inhibition

exerted by HVC_{INT} onto HVC_X and HVC_{RA} neurons, facilitated in the case of HVC_X through

rebound firing, and all orchestrated by intrinsic mechanisms.

A potential drawback of our model is that it does not incorporate brainstem feedback

processes or address inter-hemispheric coordination as proposed by others (Galvis et al.,

2018; Gibb et al., 2009b). Another drawback is its sole focus on local excitatory

connectivity within the HVC (Kornfeld et al., 2017; Long et al., 2010). Moreover, HVC

neurons receive afferent excitatory connections (Akutagawa & Konishi, 2010; Nottebohm

et al., 1982) that plays significant roles in their local dynamics. For example, the excitatory

inputs that HVC neurons receive from Uvaeformis may be crucial in initiating (Andalman

et al., 2011; Danish et al., 2017; Galvis et al., 2018) or sustaining (Hamaguchi et al., 2016)

the sequential activity. In addition, while our simplified, somatically driven architecture

enables better exploration of mechanisms for sequence propagation, future extensions

of the model will incorporate dendritic compartments to more accurately reflect the

intrinsic bursting mechanisms observed in HVC_{RA} neurons. Moreover, our model was run

at a fixed physiological temperature, but it's well known going all the way back to Hodgkin

and Huxley that both ion channel activity and synaptic dynamics can change with

temperature. In future work, adding temperature scaling (like Q10 factors) could help us

explore how burst timing and sequence speed change with temperature changes, and

how neural activity in HVC would/would not preserve its precision under different

physiological conditions.

1015

The role of ion channels in controlling network activity

Our model highlights the role of principal ion channels (I_H, I_{CaT}, I_{SK} and I_A) in controlling

HVC's network dynamics and progressing its neural sequence. Hyperpolarization-

activated ionic conductances had been widely observed across various electrically

excitable cells (Pape, 1996) and play significant roles in rhythmogenesis (Budde et al.,

1997; Golowasch et al., 1992; Golowasch & Marder, 1992). In our network, model HVC_X

neurons are not able to elicit their rebound bursting without I_H and sequence is halted if

this conductance is upregulated in either of HVC_X or HVC_{INT} (Figures 8-10, 11 and 12).

Similarly, the T-type Ca^2+ current is recognized as crucial in various systems as an ionic

contributor to burst generation (Deschenes et al., 1982; Fraser & MacVicar, 1991;

Huguenard, 1996; Llinás & Yarom, 1981). Lewicki (1996) observed a significant

hyperpolarization in some HVC neurons *in vivo* before they emit their corresponding

bursts, and the intensity of the burst correlates with the degree of hyperpolarization. In

this study, we have illustrated its pivotal role in rebound spiking where up-regulating this

conductance in HVC_X or HVC_{INT} halts sequence propagation (Figures 8-10, 11 and 12).

Moreover, the A-type K^+ current is involved in several rhythmogenic activities controlling

membrane excitability (Coetzee et al., 1999; Ellis et al., 2007; Gross et al., 2016) and in

our network, upregulating I_A suppress bursting in model HVC_{RA} and breaks sequence

propagation (Fig. 7). Finally, the small conductance Ca^{2+}-activated potassium current

(I_{SK}) plays important roles in the regulation of excitable cells controlling network rhythmic

activity (Benítez et al., 2011; Chen et al., 2014; Pedarzani et al., 2005) and in our network

I_{SK} plays a significant role since its upregulation in HVC_{RA} or its downregulation in HVC_X

eliminates sequence propagation (Fig.7 and Fig. 12 respectively).

In conclusion, the network model developed provide a large step forward in describing

the biophysics of HVC circuitry, and may throw a new light on certain dynamics in the

mammalian brain, particularly the motor cortex (Shmiel et al., 2006) and the hippocampus

regions (Lee & Wilson, 2002) where precisely-timed sequential activity is crucial. We

suggest that temporally-precise sequential activity may be a manifestation of neural

networks comprised of chain of microcircuits, each containing pools of excitatory and

inhibitory neurons, with local interplay among neurons of the same microcircuit and global

interplays across the various microcircuits, and with structured inhibition and intrinsic

properties synchronizing the neuronal pools and stabilizing timing within the ongoing

sequence.

References

Akutagawa, E., & Konishi, M. (2010). New brain pathways found in the vocal control system of a songbird. *Journal of Comparative Neurology*, *518*(15), 3086-3100.

Amador, A., Perl, Y. S., Mindlin, G. B., & Margoliash, D. (2013). Elemental gesture dynamics are encoded by song premotor cortical neurons. *Nature*, *495*(7439), 59-64.

Andalman, A. S., Foerster, J. N., & Fee, M. S. (2011). Control of vocal and respiratory patterns in birdsong: dissection of forebrain and brainstem mechanisms using temperature. *PLoS One*, *6*(9), e25461.

Benichov, J. I., & Vallentin, D. (2020). Inhibition within a premotor circuit controls the timing of vocal turn-taking in zebra finches. *Nature communications*, *11*(1), 221.

Benítez, B. A., Belálcazar, H. M., Anastasía, A., Mamah, D. T., Zorumski, C. F., Mascó, D. H., Herrera, D. G., & De Erausquin, G. A. (2011). Functional reduction of SK3-mediated currents precedes AMPA-receptor-mediated excitotoxicity in dopaminergic neurons. *Neuropharmacology*, *60*(7-8), 1176-1186.

Budde, T., Biella, G., Munsch, T., & Pape, H. C. (1997). Lack of Regulation by Intracellular Ca2+ of the Hyper Polarization-Activated Cation Current in Rat Thalamic Neurones. *The Journal of physiology*, *503*(1), 79-85.

Cannon, J., Kopell, N., Gardner, T., & Markowitz, J. (2015). Neural sequence generation using spatiotemporal patterns of inhibition. *PLoS computational biology*, *11*(11), e1004581.

Chen, L., Deltheil, T., Turle-Lorenzo, N., Liberge, M., Rosier, C., Watabe, I., Sreng, L., Amalric, M., & Mourre, C. (2014). SK channel blockade reverses cognitive and motor deficits induced by nigrostriatal dopamine lesions in rats. *International Journal of Neuropsychopharmacology*, *17*(8), 1295-1306.

Coetzee, W. A., Amarillo, Y., Chiu, J., Chow, A., Lau, D., McCormack, T., Morena, H., Nadal, M. S., Ozaita, A., & Pountney, D. (1999). Molecular diversity of K+ channels. *Annals of the New York Academy of Sciences*, *868*(1), 233-255.

Danish, H. H., Aronov, D., & Fee, M. S. (2017). Rhythmic syllable-related activity in a songbird motor thalamic nucleus necessary for learned vocalizations. *PLoS One*, *12*(6), e0169568.

Daou, A., & Margoliash, D. (2020). Intrinsic neuronal properties represent song and error in zebra finch vocal learning. *Nature communications*, *11*(1), 952.

Daou, A., Ross, M. T., Johnson, F., Hyson, R. L., & Bertram, R. (2013). Electrophysiological characterization and computational models of HVC neurons in the zebra finch. *Journal of neurophysiology, 110*(5), 1227-1245.

Datunashvili, M., Chaudhary, R., Zobeiri, M., Lüttjohann, A., Mergia, E., Baumann, A., Balfanz, S., Budde, B., Van Luijtelaar, G., & Pape, H.-C. (2018). Modulation of hyperpolarization-activated inward current and thalamic activity modes by different cyclic nucleotides. *Frontiers in cellular neuroscience*, *12*, 369.

Deschenes, M., Roy, J., & Steriade, M. (1982). Thalamic bursting mechanism: an inward slow current revealed by membrane hyperpolarization. *Brain research*, *239*(1), 289-293.

Destexhe, A., & Babloyantz, A. (1993). A model of the inward current Ih and its possible role in thalamocortical oscillations. *Neuroreport*, *4*(2), 223-226.

Destexhe, A., Mainen, Z. F., & Sejnowski, T. J. (1994). Synthesis of models for excitable membranes, synaptic transmission and neuromodulation using a common kinetic formalism. *Journal of computational neuroscience*, *1*, 195-230.

Drew, P. J., & Abbott, L. (2003). Model of song selectivity and sequence generation in area HVc of the songbird. *Journal of neurophysiology*, *89*(5), 2697-2706.

Dunmyre, J. R., Del Negro, C. A., & Rubin, J. E. (2011). Interactions of persistent sodium and calcium-activated nonspecific cationic currents yield dynamically distinct bursting regimes in a model of respiratory neurons. *Journal of computational neuroscience*, *31*, 305-328.

Dutar, P., Vu, H. M., & Perkel, D. J. (1998). Multiple cell types distinguished by physiological, pharmacological, and anatomic properties in nucleus HVc of the adult zebra finch. *Journal of neurophysiology*, *80*(4), 1828-1838.

Egger, R., Tupikov, Y., Elmaleh, M., Katlowitz, K. A., Benezra, S. E., Picardo, M. A., Moll, F., Kornfeld, J., Jin, D. Z., & Long, M. A. (2020). Local axonal conduction shapes the spatiotemporal properties of neural sequences. *Cell*, *183*(2), 537-548. e512.

Ellis, L. D., Krahe, R. d., Bourque, C. W., Dunn, R. J., & Chacron, M. J. (2007). Muscarinic receptors control frequency tuning through the downregulation of an A-type potassium current. *Journal of neurophysiology*, *98*(3), 1526-1537.

Elmaleh, M., Kranz, D., Asensio, A. C., Moll, F. W., & Long, M. A. (2021). Sleep replay reveals premotor circuit structure for a skilled behavior. *Neuron*, *109*(23), 3851-3861. e3854.

Fee, M. S., & Goldberg, J. H. (2011). A hypothesis for basal ganglia-dependent reinforcement learning in the songbird. *Neuroscience*, *198*, 152-170.

Fee, M. S., Kozhevnikov, A. A., & Hahnloser, R. H. (2004). Neural mechanisms of vocal sequence generation in the songbird. *Annals of the New York Academy of Sciences*, *1016*(1), 153-170.

Fee, M. S., & Scharff, C. (2010). The songbird as a model for the generation and learning of complex sequential behaviors. *ILAR journal*, *51*(4), 362-377.

Fraser, D. D., & MacVicar, B. A. (1991). Low-threshold transient calcium current in rat hippocampal lacunosum-moleculare interneurons: kinetics and modulation by neurotransmitters. *Journal of Neuroscience*, *11*(9), 2812-2820.

Fujimoto, H., Hasegawa, T., & Watanabe, D. (2011). Neural coding of syntactic structure in learned vocalizations in the songbird. *Journal of Neuroscience*, *31*(27), 10023-10033.

Funahashi, M., Mitoh, Y., Kohjitani, A., & Matsuo, R. (2003). Role of the hyperpolarization-activated cation current (Ih) in pacemaker activity in area postrema neurons of rat brain slices. *The Journal of physiology*, *552*(1), 135-148.

Galvis, D., Wu, W., Hyson, R. L., Johnson, F., & Bertram, R. (2018). Interhemispheric dominance switching in a neural network model for birdsong. *Journal of neurophysiology*, *120*(3), 1186-1197.

Gibb, L., Gentner, T. Q., & Abarbanel, H. D. (2009a). Brain stem feedback in a computational model of birdsong sequencing. *Journal of neurophysiology*, *102*(3), 1763-1778.

Gibb, L., Gentner, T. Q., & Abarbanel, H. D. (2009b). Inhibition and recurrent excitation
in a computational model of sparse bursting in song nucleus HVC. *Journal of
neurophysiology*, *102*(3), 1748-1762.
Glaze, C. M., & Troyer, T. W. (2013). Development of temporal structure in zebra finch
song. *Journal of neurophysiology*, *109*(4), 1025-1035.
Golowasch, J., Buchholtz, F., Epstein, I. R., & Marder, E. (1992). Contribution of individual
ionic currents to activity of a model stomatogastric ganglion neuron. *Journal of
neurophysiology*, *67*(2), 341-349.
Golowasch, J., Casey, M., Abbott, L., & Marder, E. (1999). Network stability from activity-
dependent regulation of neuronal conductances. *Neural computation*, *11*(5), 1079-
1096.
Golowasch, J., & Marder, E. (1992). Ionic currents of the lateral pyloric neuron of the
stomatogastric ganglion of the crab. *Journal of neurophysiology*, *67*(2), 318-331.
Gross, C., Yao, X., Engel, T., Tiwari, D., Xing, L., Rowley, S., Danielson, S. W., Thomas,
K. T., Jimenez-Mateos, E. M., & Schroeder, L. M. (2016). MicroRNA-mediated
downregulation of the potassium channel Kv4. 2 contributes to seizure onset. *Cell
reports*, *17*(1), 37-45.
Hahnloser, R. H., Kozhevnikov, A. A., & Fee, M. S. (2002). An ultra-sparse code
underliesthe generation of neural sequences in a songbird. *Nature*, *419*(6902), 65-
70.
Hamaguchi, K., Tanaka, M., & Mooney, R. (2016). A distributed recurrent network
contributes to temporally precise vocalizations. *Neuron*, *91*(3), 680-693.
Harvey, C. D., Coen, P., & Tank, D. W. (2012). Choice-specific sequences in parietal
cortex during a virtual-navigation decision task. *Nature*, *484*(7392), 62-68.
Hodgkin, A. L., & Huxley, A. F. (1952). A quantitative description of membrane current
and its application to conduction and excitation in nerve. *The Journal of physiology*,
117(4), 500.
Huguenard, J. (1996). Low-threshold calcium currents in central nervous system neurons.
Annual review of physiology, *58*(1), 329-348.
Jagodic, M. M., Pathirathna, S., Joksovic, P. M., Lee, W., Nelson, M. T., Naik, A. K., Su,
P., Jevtovic-Todorovic, V., & Todorovic, S. M. (2008). Upregulation of the T-type
calcium current in small rat sensory neurons after chronic constrictive injury of the
sciatic nerve. *Journal of neurophysiology*, *99*(6), 3151-3156.
Jin, D. Z. (2009). Generating variable birdsong syllable sequences with branching chain
networks in avian premotor nucleus HVC. *Physical Review E*, *80*(5), 051902.
Jin, D. Z., Ramazanoğlu, F. M., & Seung, H. S. (2007). Intrinsic bursting enhances the
robustness of a neural network model of sequence generation by avian brain area
HVC. *Journal of computational neuroscience*, *23*, 283-299.
Kornfeld, J., Benezra, S. E., Narayanan, R. T., Svara, F., Egger, R., Oberlaender, M.,
Denk, W., & Long, M. A. (2017). EM connectomics reveals axonal target variation
in a sequence-generating network. *Elife*, *6*, e24364.
Kosche, G., Vallentin, D., & Long, M. A. (2015). Interplay of inhibition and excitation
shapes a premotor neural sequence. *Journal of Neuroscience*, *35*(3), 1217-1227.
Kozhevnikov, A. A., & Fee, M. S. (2007). Singing-related activity of identified HVC
neurons in the zebra finch. *Journal of neurophysiology*, *97*(6), 4271-4283.

Kubota, M., & Saito, N. (1991). Sodium-and calcium-dependent conductances of
neurones in the zebra finch hyperstriatum ventrale pars caudale in vitro. *The*
Journal of physiology, *440*(1), 131-142.
Kubota, M., & Taniguchi, I. (1998). Electrophysiological characteristics of classes of
neuron in the HVc of the zebra finch. *Journal of neurophysiology*, *80*(2), 914-923.
Lee, A. K., & Wilson, M. A. (2002). Memory of sequential experience in the hippocampus
during slow wave sleep. *Neuron*, *36*(6), 1183-1194.
Leonardo, A., & Fee, M. S. (2005). Ensemble coding of vocal control in birdsong. *Journal*
of Neuroscience, *25*(3), 652-661.
Lewicki, M. S. (1996). Intracellular characterization of song-specific neurons in the zebra
finch auditory forebrain. *Journal of Neuroscience*, *16*(18), 5854-5863.
Li, M., & Greenside, H. (2006). Stable propagation of a burst through a one-dimensional
homogeneous excitatory chain model of songbird nucleus HVC. *Physical Review*
E, *74*(1), 011918.
Llinás, R., & Yarom, Y. (1981). Properties and distribution of ionic conductances
generating electroresponsiveness of mammalian inferior olivary neurones in vitro.
The Journal of physiology, *315*(1), 569-584.
Long, M. A., & Fee, M. S. (2008). Using temperature to analyse temporal dynamics in the
songbird motor pathway. *Nature*, *456*(7219), 189-194.
Long, M. A., Jin, D. Z., & Fee, M. S. (2010). Support for a synaptic chain model of neuronal
sequence generation. *Nature*, *468*(7322), 394-399.
MacDonald, C. J., Lepage, K. Q., Eden, U. T., & Eichenbaum, H. (2011). Hippocampal
"time cells" bridge the gap in memory for discontiguous events. *Neuron*, *71*(4), 737-
749.
Marder, E., & Goaillard, J.-M. (2006). Variability, compensation and homeostasis in
neuron and network function. *Nature Reviews Neuroscience*, *7*(7), 563-574.
Markowitz, J. E., Liberti III, W. A., Guitchounts, G., Velho, T., Lois, C., & Gardner, T. J.
(2015). Mesoscopic patterns of neural activity support songbird cortical
sequences. *PLoS biology*, *13*(6), e1002158.
Mooney, R. (2000). Different subthreshold mechanisms underlie song selectivity in
identified HVc neurons of the zebra finch. *Journal of Neuroscience, 20*(14), 5420-
5436.
Mooney, R. (2009). Neurobiology of song learning. *Current opinion in neurobiology*, *19*(6),
654-660.
Mooney, R., Hoese, W., & Nowicki, S. (2001). Auditory representation of the vocal
repertoire in a songbird with multiple song types. *Proceedings of the national*
academy of sciences, *98*(22), 12778-12783.
Mooney, R., & Prather, J. F. (2005). The HVC microcircuit: the synaptic basis for
interactions between song motor and vocal plasticity pathways. *Journal of*
Neuroscience, 25(8), 1952-1964.
Nottebohm, F., Paton, J. A., & Kelley, D. B. (1982). Connections of vocal control nuclei in
the canary telencephalon. *Journal of Comparative Neurology*, *207*(4), 344-357.
Pape, H.-C. (1996). Queer current and pacemaker: the hyperpolarization-activated cation
current in neurons. *Annual review of physiology*, *58*(1), 299-327.
Pastalkova, E., Itskov, V., Amarasingham, A., & Buzsaki, G. (2008). Internally generated
cell assembly sequences in the rat hippocampus. *Science*, *321*(5894), 1322-1327.

Pedarzani, P., McCutcheon, J. E., Rogge, G., Jensen, B. S., Christophersen, P., Hougaard, C., Strøbæk, D., & Stocker, M. (2005). Specific enhancement of SK channel activity selectively potentiates the afterhyperpolarizing current IAHP and modulates the firing properties of hippocampal pyramidal neurons. *Journal of Biological Chemistry*, *280*(50), 41404-41411.

Peters, A. J., Chen, S. X., & Komiyama, T. (2014). Emergence of reproducible spatiotemporal activity during motor learning. *Nature*, *510*(7504), 263-267.

Prather, J. F., Peters, S., Nowicki, S., & Mooney, R. (2008). Precise auditory–vocal mirroring in neurons for learned vocal communication. *Nature*, *451*(7176), 305-310.

Roberts, T. F., Hisey, E., Tanaka, M., Kearney, M. G., Chattree, G., Yang, C. F., Shah, N. M., & Mooney, R. (2017). Identification of a motor-to-auditory pathway important for vocal learning. *Nature neuroscience*, *20*(7), 978-986.

Schmitt, L. I., Wimmer, R. D., Nakajima, M., Happ, M., Mofakham, S., & Halassa, M. M. (2017). Thalamic amplification of cortical connectivity sustains attentional control. *Nature*, *545*(7653), 219-223.

Shea, S. D., Koch, H., Baleckaitis, D., Ramirez, J.-M., & Margoliash, D. (2010). Neuron-specific cholinergic modulation of a forebrain song control nucleus. *Journal of neurophysiology*, *103*(2), 733-745.

Shmiel, T., Drori, R., Shmiel, O., Ben-Shaul, Y., Nadasdy, Z., Shemesh, M., Teicher, M., & Abeles, M. (2006). Temporally precise cortical firing patterns are associated with distinct action segments. *Journal of neurophysiology*, *96*(5), 2645-2652.

Skaggs, W. E., McNaughton, B. L., Wilson, M. A., & Barnes, C. A. (1996). Theta phase precession in hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus*, *6*(2), 149-172.

Terman, D., Rubin, J. E., Yew, A., & Wilson, C. (2002). Activity patterns in a model for the subthalamopallidal network of the basal ganglia. *Journal of Neuroscience*, *22*(7), 2963-2976.

Vallentin, D., Kosche, G., Lipkind, D., & Long, M. A. (2016). Inhibition protects acquired song segments during vocal learning in zebra finches. *Science*, *351*(6270), 267-271.

Varela, J. A., Sen, K., Gibson, J., Fost, J., Abbott, L., & Nelson, S. B. (1997). A quantitative description of short-term plasticity at excitatory synapses in layer 2/3 of rat primary visual cortex. *Journal of Neuroscience*, *17*(20), 7926-7940.

Wang, C. Z. H., Herbst, J. A., Keller, G. B., & Hahnloser, R. H. R. (2008). Rapid interhemispheric switching during vocal production in a songbird. *PLoS biology*, *6*(10), e250.

Wang, P., Liu, H., Wang, L., & Gao, R. X. (2018). Deep learning-based human motion recognition for predictive context-aware human-robot collaboration. *CIRP annals*, *67*(1), 17-20.

Wang, X.-J., Liu, Y., Sanchez-Vives, M. V., & McCormick, D. A. (2003). Adaptation and temporal decorrelation by single neurons in the primary visual cortex. *Journal of neurophysiology*, *89*(6), 3279-3293.

Weber, A. P., & Hahnloser, R. H. R. (2007). Spike correlations in a songbird agree with a simple Markov population model. *PLoS computational biology*, *3*(12), e249.

Wild, J. M., Williams, M. N., Howie, G. J., & Mooney, R. (2005). Calcium-binding proteins define interneurons in HVC of the zebra finch (Taeniopygia guttata). *Journal of Comparative Neurology*, *483*(1), 76-90.

Williams, A. H., O'Leary, T., & Marder, E. (2013). Homeostatic regulation of neuronal excitability. *Scholarpedia*, *8*(1), 1656.

Yao, H., Donnelly, D. F., Ma, C., & LaMotte, R. H. (2003). Upregulation of the hyperpolarization-activated cation current after chronic compression of the dorsal root ganglion. *Journal of Neuroscience*, *23*(6), 2069-2074.

Yu, A. C., & Margoliash, D. (1996). Temporal hierarchical control of singing in birds. *Science*, *273*(5283), 1871-1875.

1298 APPENDIX 1

1299 ***Voltage-gated ionic currents***
1300

The constant-conductance leak current is $I_L = g_L(V - V_L)$. The remaining voltage-gated

ionic currents have non-constant dependent currents with activation/inactivation kinetics:

$$I_K = g_K n^4 (V - V_K) \qquad (4)$$

$$I_{Na} = g_{Na} m_\infty^3(V) h(V - V_{Na}) \qquad (5)$$

$$I_A = g_A a_\infty(V) e(V - V_K) \qquad (6)$$

$$I_{CaL} = g_{Ca} s_\infty^2(V) \frac{Ca_{ex}}{1 - e^{\frac{2FV}{RT}}} \qquad (7)$$

where $x_\infty(V) = \dfrac{1}{1 + e^{\frac{V - \theta_x}{\sigma_x}}}$, $\qquad x = m, a \; or \; s \qquad (8)$

and $\dfrac{dx}{dt} = \dfrac{x_\infty(V) - V}{\tau_x}$, $x = n, h \; or \; e \quad (9)$

where $x_\infty(V)$ for n and e is given by (8) and for h_∞ as follows

$$h_\infty(V) = \frac{\alpha_h(V)}{\alpha_h(V) + \beta_h(V)} \qquad (10)$$

where $\quad \alpha_h(V) = 0.128 e^{\frac{V+15}{-18}} \qquad (11)$

and $\quad \beta_h(V) = \dfrac{4}{1 + e^{\frac{V+27}{-5}}} \qquad (12)$

Low-voltage activated T-type calcium current
1314

The low-voltage activated T-type Ca2+ current is described by the Goldman-Hodgkin-

Katz formula:

$$I_{CaT} = g_{CaT} (a_T)^3_\infty (V)(b_T)^2_\infty (r_T) \frac{Ca_{ex}}{1-e^{\frac{2FV}{RT}}} \quad (13)$$

where $a_{T_\infty}(V) = \dfrac{1}{1 + e^{\frac{V - \theta_{a_T}}{\sigma_{a_T}}}}$ (14)

and $b_{T_\infty}(r_T) = \dfrac{1}{1 + e^{\frac{r_T - \theta_b}{\sigma_b}}} - \dfrac{1}{1 + e^{\frac{-\theta_b}{\sigma_b}}}$ (15)

with $\dfrac{dr_T}{dt} = \dfrac{r_{T_\infty}(V) - r_T}{\tau_{r_T}(V)}$ (16)

and $r_{T_\infty}(V) = \dfrac{1}{1 + e^{\frac{V - \theta_{a_T}}{\sigma_{a_T}}}}$ (17)

and $\tau_{r_T}(V) = \tau_{r_0} + \dfrac{\tau_{r_1}}{1 + e^{\frac{V - \theta_{r r_T}}{\sigma_{r r_T}}}}$ (18)

Calcium-dependent Potassium current
1324

1325 The small conductance potassium current (I_{SK}) is modeled as

1326 $I_{SK} = g_{SK} k_\infty ([Ca^{2+}]_i)(V - V_K)$ (19)

1327 where $k_\infty ([Ca^{2+}]_i) = \dfrac{[Ca^{2+}]_i^2}{[Ca^{2+}]_i^2 + k_s^2}$ (20)

1328 and $\dfrac{d[Ca^{2+}]_i}{dt} = -f \, \varepsilon(I_{CaL} + I_{CaT}) + k_{Ca}([Ca^{2+}]_i - b_{Ca})$ (21)

1329

1330 ***Hyperpolarization-activated inward current***
1331

The hyperpolarization activated inward current's activation is modeled as in Destexhe and

Babloyantz (1993) using a fast component (r_f) and a slow component (r_s) as follows:

$I_H = g_H [k_r r_f + (1 - k_r) r_s](V - V_h)$ (22)

The fast activation component is given by:

$\dfrac{dr_f}{dt} = \dfrac{r_{f_\infty}(V) - r_f}{\tau_{r_f}(V)}$ (23)

where $r_{f_\infty}(V) = \dfrac{1}{1+e^{\frac{V-\theta r_f}{\sigma r_f}}}$ (24)

with its time constant $\tau_{r_f}(V) = \dfrac{p_{r_f}}{\frac{-7.4(V+70)}{e^{(\frac{V+70}{-0.8})}-1}+65\,e^{(\frac{V+56}{-23})}}$ (25)

The slow activation component obeys:

$\dfrac{dr_s}{ds} = \dfrac{r_{s\infty}(V)-r_s}{\tau_{r_s}}$ (26)

where $r_{s\infty}(V) = \dfrac{1}{1+e^{\frac{-(V-\theta r_s)}{\sigma r_s}}}$ (27)

1342

1343 Synaptic currents equations

1344

$I_{AMPA} = \overline{g_{AMPA}}s_{AMPA}(V - V_{AMPA})$

$I_{GABA-A} = \overline{g_{GABA-A}}s_{GABAA}(V - V_{GABA-A})$

where s_{AMPA} and s_{GABAA} are given by

$[T]_{pre} = \dfrac{T_{max}}{1+\exp(\frac{V_{pre}-V_T}{Kp})}$ and $\dfrac{ds}{dt} = a_r\,[T](1-s) - a_d s$

with $T_{max} = 1, K_p = 5, V_T = 2$. For GABA_A, $a_r = 5$ and $a_d = 0.18$, while for AMPA, $a_r =$

1.1 and $a_d = 0.19$.

1351

1352

1353

1354

1355

1357

Parameter	Value	Parameter	Value
V_L	$-70\ mV$	θ_{a_T}	$-65\ mV$
V_K	$-90\ mV$	θ_b	$0.4\ mV$
V_{Na}	$50\ mV$	θ_{r_T}	$-67\ mV$
V_{Ca}	$85\ mV$	$\theta_{r_{r_T}}$	$68\ mV$
V_H	$-30\ mV$	σ_m	$-5\ mV$
g_L	$2\ nS$	σ_n	$-5\ mV$
g_{Ca}	$19\ nS$	σ_s	$-0.05\ mV$
τ_n	$10\ msec$	σ_a	$-10\ mV$
τ_{hp}	$1000\ msec$	σ_e	$5\ mV$
τ_e	$20\ msec$	σ_{r_f}	$5\ mV$
τ_h	$1\ msec$	σ_{r_s}	$25\ mV$
τ_{r_s}	$1500\ msec$	σ_{a_T}	$-7.8\ mV$
τ_{r_0}	$200\ msec$	σ_b	$-0.1\ mV$
τ_{r_1}	$87.5\ msec$	σ_{r_T}	$2\ mV$
θ_m	$-35\ mV$	$\sigma_{r_{r_T}}$	$2.2\ mV$
θ_n	$-30\ mV$	f	0.1
θ_s	$-20\ mV$	ε	$0.0015\ pA^{-1}\ \text{microM}\ msec^{-1}$
θ_a	$-20\ mV$	k_{Ca}	$0.3\ msec^{-1}$
θ_e	$-60\ mV$	b_{Ca}	$0.1\ micro\ M$

θ_{r_f}	$-105\ mV$	k_s	$0.5\ micro\ M$
θ_{r_s}	$-105\ mV$	p_{r_f}	100
$g_{Na_{HVC_{RA}}}$	$300\ nS$	$g_{K_{HVC_{RA}}}$	$150\ nS$
$g_{Na_{HVC_X}}$	$450\ nS$	$g_{K_{HVC_X}}$	$80\ nS$
$g_{Na_{HVC_{INT}}}$	$800\ nS$	$g_{K_{HVC_{INT}}}$	$1200\ nS$

1358

1359   **Table 1.** Fixed parameter values used in all simulations.

1360