Lipid Nanoparticles from *L. meyenii* Walp Mitigate Sepsis through Multimodal Protein Corona Formation

Junsik J. Sung[1], Jacob R. Shaw[2], Josie D. Rezende[1], Shruti Dharmaraj[1], Andrea L. Cottingham[1], Mehari M. Weldemariam[1], Jace W. Jones[1], Maureen A. Kane[1], Ryan M. Pearson[1,2,3*]

1. Department of Pharmaceutical Sciences, University of Maryland School of Pharmacy, 20 N. Pine Street, Baltimore, MD 21201, USA

2. Department of Microbiology and Immunology, University of Maryland School of Medicine, 685 W. Baltimore Street, Baltimore, MD 21201, USA

3. Marlene and Stewart Greenebaum Comprehensive Cancer Center, University of Maryland School of Medicine, 22 S. Greene Street, Baltimore, MD 21201, USA

[*]Address correspondence to:
Ryan M. Pearson, Ph.D.
Associate Professor, Department of Pharmaceutical Sciences
Director, Bio- and Nano-Technology Center
University of Maryland School of Pharmacy
20 N. Pine Street
N525 Pharmacy Hall
Baltimore, MD 21201, US
Phone: 410-706-3257
Email: rpearson@rx.umaryland.edu

Short title: Plant-derived Nanoparticles Mitigate Sepsis

Abstract

Background: Plant-derived nanoparticles (PDNP) are nano-sized particles isolated from various edible plants that contain bioactive components involved in regulating cellular immune responses against pathogenic intrusion and inflammation.

Purpose: This study describes a novel PDNP derived from *Lepidium meyenii* Walp (maca) that efficiently captures pro-inflammatory cytokines and acute phase proteins in its protein corona to enhance survival in two representative lethal models of sepsis.

Methods: Lipid nanoparticles were isolated from maca (MDNP) and triacylglycerols and phytoceramides were identified as major constituents using lipidomics. The physicochemical properties of MDNPs were determined, anti-inflammatory effects of MDNP were evaluated using *in vitro* models and *in vivo* using endotoxemia and cecal ligation and puncture (CLP) polymicrobial sepsis models. Proteomic analysis of MDNP in healthy or LPS-induced inflammatory plasma was used to determine the composition and inflammatory pathways modulated due to the MDNP protein corona.

Results: *In vitro* studies showed that MDNP were non-toxic, reduced macrophage activation, and effectively sequestered pro-inflammatory cytokines to mitigate NF-κB activity under lipopolysaccharide (LPS) stimulation. In a pre-established LPS-induced endotoxemia model, MDNP-treated mice showed significantly reduced systemic pro-inflammatory cytokines and enhanced survival. Untargeted proteomics and pathway analysis of the MDNP protein corona identified an enrichment in acute phase proteins in MDNP-LPS plasma coronas. MDNP treatment also significantly improved survival in the CLP sepsis model in the absence of antibiotics.

Conclusion: This work identified MDNP as an efficient, plant-derived lipid NP that broadly sequesters and neutralizes a compilation of inflammatory mediators in their coronas, offering multimodal therapeutic potential for treating inflammatory diseases.

Key words

Plant-derived lipid nanoparticles, inflammation, cytokines, acute phase response, sepsis, protein corona

Abbreviations

AP-1: activator protein-1, APCS: Serum Amyloid P component, APP: acute phase proteins, APR: acute phase response, AST/SGOT: aspartate aminotransferase, ATL/SGPT: alanine aminotransferase, BMDM: bone marrow derived macrophages, BUN: blood urea nitrogen, Cer(Phyto): phytoceramide , CLP: cecal ligation and puncture, CREA: creatine, DAMP: damage-associated molecular pattern, DAPI: 4',6-diamidino-2-phenylindole, DGDG: digalactosyldiacylglycerol, DLS: dynamic light scattering, DSC: differential scanning calorimetry, FABP4: Fatty Acid Binding Protein 4, FFA: free fatty acids, FITC: fluorescein isothiocyanate, H&E: hematoxylin & eosin, HexCer(phyto): phytohexosylceramide, Hp: haptoglobin, HSP90AA: Heat Shock Protein 90 alpha, IFN-γ: interferon- gamma, IL-1β: interleukin-1beta, IL-6: interleukin-6, IP: intraperitoneal, IV: intravenous, LCN2: lipocalin-2, LPC: lysophosphatidylcholine, LPS: lipopolysaccharide, MDNP: maca-derived lipid nanoparticles, NF-κB: nuclear factor kappa-light-chain-enhancer of activated B cells, NGP4: neutrophilic granule protein, PAMP: pathogen-associated molecular pattern, PC: phosphatidylcholine, PDNP: plant-derived lipid nanoparticle, PEG: poly(ethylene glycol), PI: phosphatidylinositol, PLGA: poly(lactic-co-glycolic acid), Saa1: serum amyloid A1, SEAP: secreted alkaline phosphatase, SerpinA3N: serine protease inhibitor A3N, TG: triglyceride, TLR-4: toll-like receptor-4, TNF-α: tumor necrosis factor-alpha, TP: total protein

Introduction

Edible plant-derived nanoparticles (PDNP) are nanostructured and membrane-enveloped vesicles secreted by plant cells that serve as carriers of various endogenous bioactive substances. Although previously perceived as cellular debris, PDNP are recognized as crucial entities that regulate cell-cell communication and immune responses against pathogens (Shao et al., 2018). The evolving understanding of the biogenesis of PDNP has been described in previous literature (Chafe, 1969; Robarbs and Kidwai, 1969). Briefly, the process initiates with the formation of a trans-Golgi network or early endosome. Plant cells then generate the matured multivesicular endosome that integrate and fuse with the plasmalemma, releasing PDNP through the inward budding of multivesicular endosomes (Li et al., 2018). There are several studies pointing to the potential therapeutic benefits of PDNP, which in some cases demonstrate anti-inflammatory (Yang et al., 2021), antioxidant (Perut et al., 2021), and anti-cancer activity (Wang et al., 2013). This emerging class of naturally derived nanovesicles offers highly advantageous features for use as nanomedicines, exhibiting negligible toxicity or immunogenicity, efficient cellular uptake, and having capacity to deliver a variety of therapeutic agents through re-engineering (Fang et al., 2024).

Cytokine secretions and the acute phase response (APR) are early, systemic immune reactions that develop within minutes to hours as part of the body's early defense mechanism to injury, infection, or immune challenge (Chen et al., 2018; Markanday, 2015a). The APR is triggered by the release of pro-inflammatory cytokines such as interleukin-6 (IL-6), tumor necrosis factor-alpha (TNF-α), interleukin-1b (IL-1β), and others by macrophages, neutrophils, and various immune cells at the site of inflammation (Tanaka et al., 2014). This response triggers the production of acute phase proteins (APP), which are typically undetectable under healthy conditions but increased 10- to 1000-fold during inflammation (Ansar and Ghosh, 2016). These proteins play roles in modulating inflammation, enhancing pathogen clearance, and promoting tissue repair (Ehlting et al., 2021). The APR is generally recognized as beneficial for restoring homeostasis disturbed by such injuries, however, APP have been described to elicit a range of functions to induce pro- or anti-inflammatory responses. During severe inflammatory responses like sepsis, a dysregulated APR can potentiate the inflammatory response leading to coagulopathy, organ failure, and even death (Castellheim et al., 2009; Markanday, 2015b; Zhang and Ning, 2021).

Several groups have developed strategies to mitigate the inflammatory response by sequestering pro-inflammatory cytokines (Lasola et al., 2020). Multiple studies highlight that the use of membrane-coated nanoparticles that mimic biological cell membranes to trap and neutralize these inflammatory cytokines (Khatoon et al., 2022). By acting as macrophage decoys, these nanoparticles can be developed to bind and neutralize endotoxins, pro-inflammatory cytokines, and even venoms that would contribute to negatively contribute to survival outcomes (Thamphiwatana et al., 2017).

Similarly, neutrophil membrane-coated nanoparticles are produced using neutrophils
isolated from mouse bone marrow after LPS stimulation (Zhang et al., 2024). The
process involves membrane extraction, incubation, and extrusion or sonication with
polymeric NP cores. These membrane-camouflaged nanoparticles reduce inflammation
by binding and neutralizing endotoxins and cytokines such as IL-6, and TNF-α (Shen et
al., 2019; Thamphiwatana et al., 2017). A study showed that IL-6 can be effectively
captured and neutralized using chitosan/hyaluronic acid polymeric nanoparticle that
have antibodies immobilized on the surface (Lima et al., 2018). Moreover,
telodendrimers were developed to bind LPS and pro-inflammatory cytokines through
controlling hydrophobic and charge-based interactions (Shi et al., 2020a), and
poly(ethylene glycol) (PEG) hydrogels were developed to capture and neutralize
histones to improve sepsis survival (Koide et al., 2021a). Despite these advancements,
the complex synthesis and formulation required for these designs to sequester pro-
inflammatory mediators presents a significant challenge to clinical translation. As a
result, there is critical need to develop simpler anti-inflammatory strategies that can
sequester multiple inflammatory mediators to mitigate pro-inflammatory responses to
augment sepsis survival.
Lepidium meyenii Walp, also referred to as maca, is a biennial root plant
indigenous to the Peruvian Andes. Maca possesses high contents of fiber, amino acids,
fatty acids, and other essential nutrients, while also containing various bioactive
compounds that are involved in integral cellular activities within plants (Wang and Zhu,
2019). Several studies have shown that maca alleviates fatigue (Zhu et al., 2022),
oxidative stress (Fei et al., 2022), tumor formation (Cao et al., 2023), and inflammation
(Yang et al., 2023). However, studies have focused on the raw material or crude extract,
which contains mixture of different active ingredients, to evaluate in disease models (Fei
et al., 2020; Tenci et al., 2017). Despite these known advantages, maca remains
relatively underexplored.
This investigation demonstrates the isolation, characterization, and therapeutic
development of a PDNP isolated from maca root, termed maca-derived lipid
nanoparticle (MDNP), for the treatment of severe inflammation as exemplified using *in*
vitro and *in vivo* models of lipopolysaccharide (LPS)-induced endotoxemia and
polymicrobial sepsis. MDNP were isolated and characterized for physicochemical
properties, as well as lipid composition using lipidomic analysis. The toxicity profile of
MDNP was then established prior to determining their uptake profile and anti-
inflammatory properties. *In vitro*, MDNP efficiently sequestered multiple pro-
inflammatory cytokines, which led to comprehensive *in vivo* assessment of their
biodistribution and therapeutic activity. Therapeutic administration of MDNP to LPS-
challenged mice led to significant reductions in plasma pro-inflammatory cytokines,
reductions in inflammation-induced organ damage, and improved survival. When
nanoparticles are introduced into a biological fluid, they are rapidly covered by a layer of

biomolecules known as the biomolecule corona (or protein corona) (Pearson et al., 2014; Shaw and Pearson, 2022). To identify the mechanism by which MDNP elicited its anti-inflammatory effects, we performed untargeted proteomics analysis of the MDNP protein corona to uncover the inflammatory mediators sequestered by MDNP and corresponding pathways and upstream regulators modulated. MDNP was found to sequester and neutralize a variety of APP, which promote the propagation of pro-inflammatory immune responses, in addition to pro-inflammatory cytokines. Lastly, the efficacy of MDNP was assessed using a clinically relevant mouse model of polymicrobial sepsis and found to significantly increase survival. These results demonstrate the potential of MDNP as an abundant, cost effective, naturally derived therapeutic agent for use as a multimodal intervention for a variety of inflammatory diseases due to their ability to sequester a broad spectrum of inflammatory mediators.

Results

Isolation and characterization of maca-derived lipid nanoparticles (MDNP)

PDNPs were isolated from maca juice using differential ultracentrifugation and density gradient sucrose gradient centrifugation (**Figure 1**) (Sung et al., 2019). MDNP accumulated at the interface of 20/35% sucrose gradient, and the recovery was approximately 4 mg per 30 g of maca powder (**Figure 1B,C**). TEM revealed that MDNP displays a spherical morphology, and their core structure was consistent with that of solid lipid nanoparticles (**Figure 1D**). Size and zeta potential of MDNP measured an average of 174.5 nm and -9.6 mV, respectively (**Figure 1E**). We next assessed the stability of MDNP at multiple temperatures. MDNP could tolerate a freeze and thaw cycle and were stable at 4°C for up to 6 days, the maximum length of time tested. All different temperature storage conditions kept MDNP stable except for room temperature (RT), where after 3 days the size was significantly increased (**Figure 1F**). The sizes and zeta potential of MDNP were also stable upon lyophilization (**Supplementary Figure S1**). Differential scanning calorimetry (DSC) analysis determined the summit of melting peak of MDNP, which was measured as 74.3°C (**Figure 1G**).

To determine the lipid composition of MDNP, untargeted high throughput lipidomics was performed. A bar graph (**Figure 1H**) and a pie/vertical slice chart (**Figure 1I**) shows the total lipid abundance and ratio in percentage of each component in MDNP. The lipidomics identified a total of eight individual lipids in four categories including sphingolipids, galactolipids, neutral lipids, and phospholipids (**Supplementary Figure S2**). The most prevalent species within the MDNP was the triglyceride (TG) subclass with 51 identifications (77.45%), followed by phytoceramide (Cer (Phyto); 14.77%), phytohexosylceramide (HexCer (Phyto)) (2.7%), free fatty acids (FFAs) (1.95%), digalactosyldiacylglycerol (DGDG) (0.72%), phosphatidylcholine (PC) (0.96%), phosphatidylinositol (PI) (0.85%), and lysophosphatidylcholine (LPC) (0.13%).

The complete protein compositional analysis of MDNP using UPLC-MS/MS
revealed undetectable levels of proteins that could be linked to plant databases from
Uniprot and the Plant Proteome Database. This aligns with our lipidomics results
indicating that MDNP is primarily composed of lipids, with minimal or no associated
protein content. Overall, these data demonstrate the successful isolation, purification,
and characterization of MDNP from bulk maca extract.

Figure 1. Isolation and characterization of MDNP. (A) Schematic representation of the isolation process for maca-derived lipid nanoparticles (MDNP). (B) Before and after images of MDNP isolation. MDNP were isolated and purified from maca extract by sucrose density gradient purification. (C) MDNP recovery from pure maca powder. (D) A representative TEM image of MDNP. (E) Room temperature measurement of size (174.5 nm) and zeta potential (-9.6 mV) of freshly isolated MDNP. (F) Storage stability test of MDNP under various storage conditions up to day 6. (G) Heat flow curve of MDNP measured by differential scanning calorimetry showing the summit of melting peak at 74.3 ºC. (H) Lipid extraction and component analysis by liquid chromatography coupled to high resolution LC-MS/MS revealed eight different lipids (TG, Cer, HexCer, DGDG, LPC, PC, PI and FFA). Bar graph shows abundance of lipids. (I) Pie graph shows the whole ratio of each component and the vertical slice magnifies the smaller portion of composition excluding TG. N=3/group.

232 ***In vitro* cytotoxicity, internalization, and therapeutic effect of MDNP**
233　　　　To analyze potential toxicity, BMDM were treated with MDNP (1 mg/mL) for 8
234 hours prior to flow cytometry analysis to measure viability and apoptosis (**Figure 2A**).
235 MDNP treatment did not affect the cell viability or induce apoptosis. Furthermore, we
236 observed a time-dependent cellular uptake of Cy5.5-labeled MDNP using BMDM
237 (**Figure 2B**). The quantitative measure of mean fluorescence intensity (MFI) from the
238 confocal microscopy images can be found in **Supplementary Figure S3**. These
239 findings demonstrate the MDNP are non-toxic and efficiently internalized by BMDM.
240　　　　BMDM were treated with LPS followed by MDNP to assess their ability to
241 mitigate immune activation. The observed decrease in CD80 (reduced but not
242 significant), CD86, and MHCII expression following MDNP treatment suggests that
243 MDNP reduced macrophage activation, contributing to a decreased pro-inflammatory
244 response (**Figure 2C-E**). Modulation of macrophage polarization is one of the strategies
245 to modulate inflammation, support tissue regeneration and reestablish disrupted
246 homeostasis(Parisi et al., 2018). We also measured a reduction in CD206 as a marker
247 of anti-inflammatory M2-like macrophages, suggesting MDNP induced a shift in
248 macrophage phenotype towards a more undifferentiated or M0-like state (**Figure 2F**).
249 These results indicate that MDNP can impact the activation status of macrophages
250 suggesting that they aid in shifting their profile towards a more balanced and neutral
251 immune response.
252　　　　To assess the anti-inflammatory effects of MDNP, we investigated its prophylactic
253 and therapeutic activity by treating MDNP on LPS-challenged BMDM. In the
254 prophylactic setting, MDNP was treated for 8 hr prior to 4 hr LPS exposure (**Figure 2G**).
255 However, it did not significantly attenuate the release of pro-inflammatory cytokines (IL-
256 6 and TNF-α) (**Figure 2H,I**). In the therapeutic setting, BMDM were challenged with
257 LPS for 4 hrs prior to the treatment of MDNP for 8 hr (**Figure 2J**). The result shows that
258 pro-inflammatory cytokines including IL-6, and TNF-α induced by LPS were significantly
259 reduced, whereas IL-1β, and IFN-γ were only slightly reduced (**Supplementary Figure
260 S4**). Furthermore, additional testing was conducted on the crude extract after MDNP
261 isolation to determine if it possesses any anti-inflammatory efficacy; however, no
262 cytokine reduction was detected (**Supplementary Figure S5**). These results suggested
263 that MDNP can reduce pro-inflammatory responses, and the anti-inflammatory
264 component of maca is the purified MDNP.
265

Figure 2. *In vitro* cytotoxicity, cellular internalization, immunomodulation and therapeutic effects against LPS-induced inflammation. (A) PI/Annexin V assay by flow cytometry displayed that treatment with higher concentration (1 mg/mL) of MDNP did not induce any significant toxicity in BMDM. Quantitative analysis of live, apoptotic, and dead cells after MDNP treatment is accompanied. (B) Cy5.5 labeled MDNP were gradually internalized by BMDM after 4 and 8 hrs. Cells were stained with DAPI and FITC-PI to visualize the nucleus and F-actin, respectively. (C-F) Immunomodulatory activity of MDNP in LPS inflamed BMDM showed reduction in M1 (CD80, CD86, MHCII) and M2 (CD206) markers. (G) A timeline for prophylactic treatment. Prophylactic approach did not make significant change in levels of (H) IL-6 and (I) TNF-⍺. (J) A timeline for therapeutic treatment. Therapeutic treatment displayed significant reduction of (K) IL-6 and (L) TNF-⍺. Data are expressed as mean ± SD (n=5). Data are expressed as mean ± SD (n=3). *P<0.05, **p<0.05 versus LPS control group.

Sequestration of cytokines and mitigation of NF-κB activation

Interleukin-6 (IL-6) and tumor necrosis factor-α (TNF-α) are well known to be
involved in pathogenesis of chronic inflammation, autoimmune diseases, and cancer
(Hirano, 2021; Popa et al., 2007). Utilizing data from the *in vitro* anti-inflammatory assay,
an investigation was conducted to confirm whether MDNP possess the capability to
sequester pro-inflammatory cytokines in the absence of cells. In a time-dependent
experiment, BMDM were treated with 100 ng/mL LPS for 4 hrs, then the supernatant
rich in inflammatory cytokines was incubated with 100 µg/mL MDNP for up to 24 hrs
(**Figure 3A**). ELISA analysis of these supernatants indicated a gradual decrease in IL-6
levels, with a maximum of 48% reduction after 8 hrs (**Figure 3B**). A similar trend was
observed for TNF-α levels, with a 28% reduction over the same time period (**Figure 3C**).
Optimization of MDNP concentration revealed concentration-dependent cytokine
sequestration properties of MDNP and 100 µg/mL MDNP exhibited the most significant
sequestration of IL-6 and TNF-α, whereas PLGA nanoparticle (as control) did not
display any cytokine sequestration properties (**Supplementary Figure S6**). We also
confirmed the ability of MDNP to sequester recombinant IL-6 to exclude the possibility

Figure 3. *In vitro* sequestration of pro-inflammatory cytokines from LPS-induced bone marrow derived macrophages. (A) Time course reduction of LPS-stimulated pro-inflammatory cytokine by treatment of MDNP. 8 hr time point exhibited the most significant reduction for both (B) IL-6 and (C) TNF-α. (D) Recombinant IL-6 was sequestered through incubation with MDNP ex vivo without cellular interaction. (E) 100 µg/mL MDNP was the most effective concentration to remove IL-6 protein. (F) Expression of NF-κB in in RAW-Blue 264.7 cell is suppressed by the treatment of MDNP. (G) NF-κB activity level is downregulated after treatment of MDNP. PLGA was used as a control to show different effect on NF-κB. All data are expressed as means ± SD (n=5). **p<0.05, ***p<0.005, and ****p<0.0005 versus LPS or NT control group.

of other proteins contributing to cytokine removal *in vitro* (**Figure 3D,E**). These findings
confirm that MDNP can sequester pro-inflammatory cytokines to mitigate inflammatory
responses.
The innate immune responses that are triggered by LPS are mediated by Toll-like
receptor (TLR)-4 and subsequent activation of the transcription factors nuclear factor
kappa B (NF-κB) and activator protein 1 (AP-1) (Nyati et al., 2017). Upon encountering
various PAMPs and DAMPs, macrophages undergo rapid activation and secrete a
diverse range of cytokines and chemokines including IL-6 and TNF-α (Liu et al., 2017).
In this experiment, RAW-Blue cells, a NF-κB macrophage reporter cell line that releases
secreted alkaline phosphatase (SEAP) after activation, were used to gauge MDNP-
mediated effects following LPS stimulation (**Figure 3F**). The expression of IL-6 is tightly
regulated at multiple levels, including gene transcription, mRNA translation and mRNA
degradation levels. Furthermore, NF-κB activation was examined by exposing LPS-
activated RAW-Blue cells to 100 µg/mL MDNP for 1 hr (**Figure 3G**). PLGA was used as
a control and were unable to reduce NF-κB activation. The results demonstrate a
significant decrease in NF-κB activity, suggesting that MDNP possess the capacity to
mitigate pro-inflammatory cell signaling and reduce macrophage activation.
300
***In vivo* biodistribution and anti-inflammatory effects of MDNP**
For the *in vivo* biodistribution study, MDNP were labeled with Cy5.5 and
administered intraperitoneally (IP) or intravenously (IV) into LPS-challenged mice (5
mg/kg LPS dose) twice at an MDNP dose of 2 mg/injection at 0.5 and 2 hr (**Figure 4A**).
At 4 hr, mice were euthanized to image the biodistribution of Cy5.5-MDNP in different
organs including liver, spleen, heart, kidney, and lung with an *in vivo* imaging system
(IVIS). Most of the Cy5.5-labeled MDNP were delivered to the liver and kidney by both
IP and IV injection, and very low amounts were detected in the spleen, heart, and lung
(**Figure 4B, C**).
We also investigated how the route of administration (IP versus IV) affected the
in vivo therapeutic effects in mice. Mice were challenged with 5 mg/kg LPS IP followed
by treatment with 2 mg MDNP at 0.5 hr and 2 hr post-LPS treatment. After 4 hrs, mice
were euthanized to collect organs samples and plasma (**Figure 4A**). Plasma samples
were processed using a MAGPIX Luminex multiplexing immunoassay system. This
system was employed to analyze the expression level of a 7-plex panel of pro-
inflammatory cytokines which included IL-6, TNF-α, MCP-1, IL-10, IL-1β, IFN-β, and
GROα. The heatmap demonstrates overall reduction of pro-inflammatory cytokines for
both routes (**Figure 4D**). The analysis revealed a significant reduction in IL-6 level
following both administration routes (**Figure 4E**), and a notable reduction in TNF-
α, MCP-1, and GROα for IP administered MDNP (**Figure 4F-K**). IV administration of
MDNP also reduced systemic cytokine levels, however it was slightly less effective. We
further evaluated blood biochemistry of albumin, alanine transaminase, creatine,

aspartate transferase, globulin, total protein, and blood urea nitrogen. Although minor
reductions were observed for MDNP-treated mice, there was no significant alteration in
levels of blood biochemistry parameters compared to LPS-treated controls
(**Supplementary Figure S7**). The non-significant reductions in blood biochemistry
following MDNP-treatment could be due to the pre-established severe inflammatory
response, that could not be completely reversed by the endpoint of the study.

Figure 4. Route of administration biodistribution and anti-inflammatory effects of MDNP (A) Timeline of MDNP treatments following LPS challenge. The schematic illustrates the IP and IV injection performed on randomly selected C57BL/6J mice groups (LPS only, LPS+IP injection, and LPS+IV injection, n=3 per group). (B) Biodistribution of fluorescently labeled MDNP in mouse organs (liver, kidney, spleen, lung, and heart). (C) Fluorescence intensity measurement of MDNP biodistribution. (D) Heatmap of pro-inflammatory cytokine expression including: (E) IL-6, (F) TNF-α, (G) MCP-1, (H) IL-10, (I) IL-1β, (J) IFN-β, (K) GROα from mice plasma samples. Data are expressed as mean ± SD (n=3). *P<0.05, **p<0.05, ***p<0.005, and ****p<0.0005 versus PBS control group.

Treatment of MDNP accelerated organ recovery and improved survival

Histological analysis was used to evaluate the level of inflammation in tissues isolated from experiments conducted in **Figure 4**. H&E staining of five different organs including liver, kidney, spleen, lung, and heart are shown in **Figure 5**. LPS-treated liver showed severe immune cell infiltration, tubular necrosis was detected in the kidney, focal necrosis, and dysregulation in white pulp of spleen was observed, and extensive clots in the alveoli of lung appeared, but no significance of heart injury was measured. On the contrary, the IV and IP administered MDNP groups displayed notable improvements in histology scores when compared to the LPS group (**Figure 5B-F**). Based on the overall safety profile and reductions in pro-inflammatory cytokines, MDNP were used to further evaluate survival *in vivo*. Mice were challenged with lethal dose of 20 mg/kg LPS in sterile PBS intraperitoneally. Then, mice were monitored after two doses of 2 mg MDNP

Figure 5. Representative images of organ recovery and survival effect by treatment of MDNP during LPS-induced sepsis. (A) H&E staining of liver, kidney, spleen, lung, and heart. The LPS treated group images shows presence of neutrophil infiltration (yellow arrow) in liver, tubular necrosis (white arrow) in the tubules in kidney, focal necrosis and dysregulation in white pulp of spleen, appearance of extensive clots (red arrow) in the alveoli in lungs, while no noticeable damage in heart. On the contrary, mice organs administered with IP and IV displayed notably reduced level of organ damage of organ damage. (B-F) Quantification of H&E staining. Levels of injury scores were calculated. Data are expressed as mean ± SD (n=3). *P<0.05, **p<0.05 versus LPS control group. (G) A schematic of survival study of LPS-treated and LPS+MDNP-treated C57BL/6J mice (n=10/group). (H) The survival analysis graph displays a significant difference between LPS only and the MDNP treatment group. Kaplan-Meier curves and a log-rank test was performed to compare the survival rates. ***p<0.0005.

341 (**Figure 5G**). As a result, 60% of MDNP treated mice survived, whereas 0% of the LPS
342 treated (control) mice survived (**Figure 5H**).
343
Proteomic profiling of MDNP protein corona
The ability of MDNP to reduce systemic cytokines and promote LPS mouse
survival prompted an exploration of the composition and types of proteins sequestered
by the MDNP. Adsorbed proteins were characterized by forming a protein corona on
MDNP *ex vivo* by incubating with either healthy or LPS-treated mouse plasma (MDNP-H
or MDNP-LPS, respectively) (**Figure 6A**). Proteomics analysis of these coronas
identified 297 total proteins (**Supplementary Table S1**), with 270 common between the
two conditions, 1 distinctive for healthy, and 26 for LPS-treated (**Figure 6B**). **Table 1**
presents a list of uniquely adsorbed proteins from each group. Among these, CD14,
LCN2 (lipocalin-2), APCS (Serum Amyloid P component), FABP4 (Fatty Acid Binding
Protein 4), NGP (Neutrophilic Granule Protein), and HSP90AA (Heat Shock Protein 90
alpha) are known to play roles in inflammation and categorized as APP excluding NGP.
The heatmap in **Figure 6C** shows the differential expression of total 297 detected
proteins. Comparing MDNP-H to MDNP-LPS, highlighted significant changes in protein
abundance, with 49 proteins being significantly upregulated and 100 proteins
downregulated (**Figure 6D**). Haptoglobin (Hp), which is known for its pro-inflammatory
properties(Shen et al., 2011), was found to be the most differentially abundant protein in
the MDNP-LPS corona. The top 15 proteins (over 20-fold change) identified in the
MDNP-LPS group are shown in **Supplementary Figure S8**. Among these, Hp,
Seprina3n, and Saa1 are well-known for their pro-inflammatory roles in regulating
inflammatory process and promoting the release of inflammation cytokines. Their levels
can increase dramatically during inflammation(Massart et al., 2020). **Figure 6E** lists the
canonical pathways of MDNP-LPS coronas from ingenuity pathway analysis (IPA). APR
signaling was most highly associated with MDNP-LPS corona. The network diagram in
Figure 6F illustrates the complex but close interaction of APP and these upstream pro-
inflammatory regulators within the APR signaling pathway. The highlighted connections
between APR shows its direct interaction with the APP sequestered on MDNP-LPS
coronas (Hp, Saa1, SerpinA3N) and a subset of upstream regulators (IL-6, TNF, IL1B,
AGT, HNF1A). Importantly, IL-6, TNF and IL1B play pivotal roles in either indirectly or
directly activating these APP. **Figure 6G** illustrates the result of *ex vivo* cytokine
sequestration assays using MDNP. It demonstrates that MDNP retains the ability to bind
and remove pro-inflammatory cytokines, such as IL-6 and TNF-α, from the LPS plasma.
It is also important to point out that protein corona formed on the surface of MDNP did
not activate any pro-inflammatory responses when treated on BMDM, supporting that
the formation of the corona effectively neutralizes the pro-inflammatory responses of
these various mediators (**Supplementary Figure S9**). **Figure 6H** summarizes the
proposed mechanism of MDNP by forming a multimodal protein corona comprised of

Figure 6. Mapping and identification of protein corona adsorbed onto the surface of MDNP. (A) A schematic showing the process of evaluating the MDNP protein corona. (B) Venn diagram illustrating a total of 296 proteins detected including unique (1 in healthy and 26 in LPS) and 270 shared proteins in the coronas. (C) Heatmap shows abundance of 49 upregulated and 100 downregulated proteins in MDNP-LPS group compared to MDNP-H healthy group. (D) Volcano plot of significantly upregulated and downregulated corona proteins. Hp, Serpina3n and Saa1 (red dots) are categorized as pro-inflammatory and acute phase response signaling proteins. (E) Top 10 canonical pathways associated with the significantly upregulated protein corona were identified, with acute phase response signaling being the most prominent pathway. (F) A network representation of acute phase response signaling proteins and major upstream regulators. (G) *Ex vivo* sequestration of pro-inflammatory cytokines by incubating LPS plasma with MDNP. Both IL-6 and TNF-α was significantly reduced by 8 hr treatment. (H) A graphical summary illustrating how the cytokine and acute phase protein corona-coated MDNP sequesters and neutralizes inflammatory proteins, thereby reducing overall inflammation.

382 pro-inflammatory cytokines and APP and neutralization. This inhibits the key
383 inflammatory pathways, blocking the production of pro-inflammatory upstream
384 regulators, ultimately reducing pro-inflammatory cytokines, improving organ function,
385 and increasing survival.
386

MDNP treatment improved survival in polymicrobial sepsis model

The survival benefit of MDNP was also tested in a clinically relevant severe
polymicrobial cecal ligation and puncture (CLP) sepsis model (**Figure 7A**). Mice
underwent the CLP procedure and were treated with three doses of 2 mg MDNP,
followed by monitoring for 5 days. To confirm MDNP could sequester pro-inflammatory
cytokines in CLP plasma, we first incubated MDNP with plasma isolated at 3 hr post-
CLP and observed an almost complete reduction in IL-6 and TNF-α (**Figure 7B**). In the
survival study, MDNP treated mice showed significant improvement in survival (40%)
compared to control group (0%) (**Figure 7C**). Taken together, the ability of MDNP to
increase the survival of septic mice in the absence of antibiotics, highlights their
translational potential for treating inflammatory diseases, including sepsis.

Figure 7. Survival effects of MDNP treatment on severe CLP model. (A) Schematic timeline of survival study of cecal ligation and puncture (CLP) C57BL/6J murine model with MDNP treatment (n=5/group) based on the complement cascade pathway from proteomics. Mice underwent CLP surgery to induce severe polymicrobial inflammation and three separate doses of MDNP were administered intraperitoneally (0.5, 2, 24 hr timepoint, 2 mg/injection) to analyze survival. (B) *Ex vivo* sequestration test of CLP plasma and MDNP showed that both IL-6 and TNF-α was significantly reduced. Kaplan-Meier curves and a log-rank test was performed to compare the survival rates. (C) The survival test displayed a significant improvement (40%) by the treatment of MDNP **p< 0.005. Data are expressed as mean ± SD (n=5). ****p<0.0005 versus PBS control group.

Discussion

A tremendous amount of research has been implemented to enhance the clinical management of inflammatory diseases caused by excessive stress (Miller and Raison, 2016), infections(Ikoba et al., 2015), autoimmune disorders (Miller et al., 2007), or secondary diseases (Dufour et al., 2022). Despite many clinical trials aimed at blocking specific inflammatory mediators, most have been unsuccessful due to the complexity and multifactorial nature of inflammatory diseases. This highlights the need for multimodal approaches that target multiple inflammatory mediators simultaneously to improve treatment outcomes. The core pathophysiology of sepsis is known to involve the disruption of the finely tuned balance between inflammatory and anti-inflammatory immune responses (Jarczak et al., 2021). Recognition of pathogen- or damage-associated molecular patterns (PAMPs/DAMPs) by innate immune cells functions as the trigger to initiate inflammatory cell signaling and subsequent cytokine release, leading to multiorgan damage, and death. Notably, studies have shown that increased plasma IL-6 and TNF-α levels in sepsis patients are correlated with mortality (Mera et al., 2011; Unsinger et al., 2010; Wu et al., 2009). Our MDNP efficiently reduced systemic pro-inflammatory cytokines including IL-6, TNF-α, among others (**Figure 4E, 6G, 7B**), while simultaneously neutralizing additional pro-inflammatory mediators (APP) (**Figure 6D**). Strategies for reducing pro-inflammatory mediators include conventional adsorption resins like Cytosorb®, or more sophisticated membrane-coated nanoparticles (Thamphiwatana et al., 2017), telodendrimer nanotraps (Shi et al., 2020), or abiotic histone-capturing hydrogel nanoparticles (Koide et al., 2021b). Although these approaches have shown success in certain cases, the complex structural composition and manufacturing processes may hinder rapid development. In contrast, we developed a MDNP, isolated from maca root, as a novel approach to mitigate broad pro-inflammatory responses from abundant and readily available plant material, facilitating convenient access, and long-term storage potential (**Figure 1**). We further demonstrated their ability to reduce organ damage and improve survival as reflected in two representative and lethal sepsis models (**Figure 5A, 5H, 7C**) through the formation of a multimodal protein corona (**Figure 6**).

Sepsis endotypes include hyperinflammatory (early) and immunosuppressive (late) states (Hotchkiss et al., 2013). A clinical study indicated that early-identified sepsis is associated clinically with higher mortality than late-identified sepsis (Jee et al., 2020). Thus, achieving control over the cytokine storm holds great potential to improve patient survival. Modes of cytokine sequestration have generally relied on charge-based interactions, where most pro-inflammatory cytokines are negatively charged, and anti-inflammatory cytokines are positively charged at neutral pH (Messina et al., 2024). Interestingly, the zeta potential of MDNP was -9.6 mV (**Figure 1E**), yet cytokines like IL-6 (pI 6.96) and TNF-α (pI 5.01) were strongly sequestered but IL-1β (pI 4.96) and IFN-γ

(pI 8.25) were not (**Figure 2K, L, S6**). These differences suggest the possibility of a
combination of forces driving the binding of cytokines to the surface of MDNP. Our
lipidomics analysis of MDNP (**Figure 1H, I**) showed the presence of multiple charged or
ionizable lipid species like Cer (Phyto), HexCer (Phyto), FFA, LPC, and PC
(**Supplementary Figure S2**). Therefore, it is possible that a combination of local charge
and hydrophobic interactions are contributing to the apparently selective sequestering of
specific pro-inflammatory cytokines. Although we have not fully elaborated how MDNP
facilitate a selective binding interaction for certain cytokines, which is a limitation of our
study, our results provide the basis for future studies to investigate the role of specific
lipid components and their ability to alter binding interactions.
Upon exposure to biological fluids, nanoparticles dynamically adsorb
biomolecules onto their surface, forming a protein corona that dictates how
nanoparticles interact with host cells. These interactions can alter both drug delivery
efficiency (Corbo et al., 2016) and biological function (Cedervall et al., 2007) of the
particles. The analysis of MDNP protein coronas revealed significant differences
between MDNP-H and MDNP-LPS groups, highlighting that unique protein fingerprints
are associated with MDNP under inflammatory conditions. This is consistent with the
concept of the 'personalized protein corona', which demonstrates that unique protein
fingerprints are formed on the surface of nanoparticles as a function of disease state (Ju
et al., 2020; Shaw and Pearson, 2022). The ability of MDNP to bind cytokines in LPS- or
CLP-treated mouse plasma confirmed its ability to directly modulate inflammatory
responses in a disease setting (**Figure 6G, 7B**). An unexpected finding from our
proteomics analysis was the identification of multiple pro-inflammatory APP being
upregulated in the coronas of MDNP-LPS like Hp, SerpinA3N, and Saa1 (**Figure 6D**).
Hp activates immune cells, promotes secretion of pro-inflammatory cytokines, and
interacts with complement system, potentially intensifying inflammatory responses
(Torres et al., 2022). SerpinA3N is involved in regulating inflammatory responses by
controlling protease activity to prevent excessive tissue damage, but it's upregulated
under inflammatory conditions (Liu et al., 2023). Saa1 plays direct role in recruiting
immune cells and modulating pro-inflammatory cytokines (Ye and Sun, 2015).
Supporting that MDNP neutralized the pro-inflammatory functions of corona constituents,
MDNP-LPS was cultured with BMDM and no significant increase in IL-6 or TNF-α was
detected (**Supplementary Figure S9**). These results highlighted that MDNP formed
multifaceted protein corona under inflammatory conditions that could mitigate aberrant
immune activation caused by individual components, representing a novel approach to
the treatment of inflammatory diseases.
Here, we demonstrated that MDNP sequestered and neutralized multiple pro-
inflammatory cytokines and APP in its protein corona to reveal a highly effective anti-
inflammatory plant-derived therapeutic strategy for managing severe inflammatory
responses. Our data indicates that MDNP exhibit negligible toxicity, are efficiently

internalized by macrophages, display immunomodulatory activities, and demonstrate
significant therapeutic efficacy. Administration of MDNP *via* IP or IV routes effectively
reduced pro-inflammatory cytokines and promoted improved survival in two
representative sepsis mouse models, LPS-induced endotoxemia and CLP polymicrobial
sepsis. This research underscores the broad therapeutic potential of the MDNP, which
leverages a unique and never previously examined mechanism of action through the
formation of a multimodal protein corona that effectively binds and neutralizes pro-
inflammatory cytokines and APP from propagating systemic inflammation.
486
Materials and methods
Isolation, purification, and characterization of maca-derived lipid nanoparticles
(MDNP)
A fine ground organic powder of 100% *Lepidium meyenii* Walp (maca) (Family:
Brassicaceae, Order: Brassicas) was purchased from an organic producer, Happy
Andes (https://www.happyandes-usa.com/). Maca powder (30 g) was mixed in 400 mL
of autoclaved deionized water and stirred for 12 hrs at room temperature to make maca
juice. The obtained maca juice was transferred to 50 mL tubes and centrifuged at 3,000
x g for 20 min at 20°C then the supernatant was centrifuged again in new 50 mL tubes
at 10,000 x g for 2 hrs at 20°C to remove small and large fibers. Next, the clear light
brown colored supernatant was collected in 70 mL polycarbonate ultracentrifuge bottles
(Beckman Coulter, Brea, CA) and was ultracentrifuged with using a 45 Ti rotor
(Beckman Coulter, Brea, CA) at 150,000 x g for 2 hrs at 4°C. Then, the collected pellets
were suspended in 2 mL of phosphate-buffered saline (PBS) through dispersion with an
ultrasonic processor. PBS-suspended maca pellets were transferred to a sucrose
gradient (8%, 20%, 35% [g/v]) and ultracentrifuged at 150,000 x g for 2 hrs at 4°C, then
the cloudy band between 8% and 20% sucrose was collected as previously described
(Sung et al., 2019). The concentration of MDNP was measured using a Bio-Rad
quantification assay. The quantified MDNP were stored at 4°C. The size and zeta
potential of MDNP was measured by dynamic light scattering (DLS) using a Zetasizer
Nano ZSP (Malvern, UK). MDNP were also tested under different storage temperature
conditions (room temperature (RT), 4°C, -20°C, and -80°C) to determine their stability.
Transmission electron microscopy (TEM) images of MDNP were acquired using a
formvar-coated copper grid and staining using 1% uranyl acetate. Differential Scanning
Calorimeter (DSC) 2500 (TA Instruments, New Castle, DE) was used to analyze the
temperature and heat flow associated with thermal transition in the MDNP. 5 µL of
sample in liquid form was added into a hermetic aluminum pan and sealed with metal lid
(DSC Consumables, Austin, MN) with Tzero sample press kit (TA Instruments, New
Castle, DE). The following thermal procedure was used to scan: ramp 2°C/min from
25°C to 80°C. The heat flow transition was analyzed on a plot by TA analysis software
Trios (Version 5.4.0.300). For lyophilization, frozen MDNP in 1.5 mL tubes with holes on

top of screw caps were placed in a benchtop freeze-dryer (Labconco, Kansas City, MO)
overnight. Completely dried samples were collected and reconstituted in PBS for
stability testing.
521
Lipid extraction and composition analysis
A total lipid extract was prepared using a modified methyl-tert-butyl ether (MTBE)
lipid extraction protocol (Matyash et al., 2008). Briefly, 400 µL of cold methanol and 10
µL of internal standard mixture (EquiSPLASH lipidomix) were added to each sample
followed by incubation at 4°C, 650 RPM shaking for 15 min. Next, 500 µL of cold MTBE
was added followed by incubation at 4°C for 1 hr with 650 RPM shaking. 500 µL of cold
water was added slowly and resulting extract was maintained 4°C, 650 rpm shaking for
15 min. Phase separation was completed by centrifugation at 8,000 RPM for 8 min at
4°C. The upper, organic phase was removed and set aside on ice. The bottom,
aqueous phase was re-extracted with 200 µL of MTBE followed by 15 min of incubation
at 4°C with 650 RPM shaking. Phase separation was completed by centrifugation at
8,000 RPM for 8 min at 4°C. The organic extract was dried under a steady stream of
nitrogen at 30°C. The recovered lipids were reconstituted in 200 µL of
chloroform:methanol (1:1, v/v) containing 200 µM of butylated hydroxytoluene. Prior to
analysis, samples were further diluted with acetonitrile:isopropanol:water (1:2:1, v/v/v).
The total lipid extract was analyzed by liquid chromatography coupled to high resolution
tandem mass spectrometry (LC-MS/MS). The LC-MS/MS analyses were performed on
an Agilent 1290 Infinity LC coupled to an Agilent 6560 Quadrupole Time-of-Flight (Q-
TOF) mass spectrometer. The separation was achieved using a C18 CSH (1.7 µm; 2.1
x 100 mm) column (Waters, Milford, MA). Mobile phase A was 10 mM ammonium
formate with 0.1% formic acid in water/acetonitrile (40:60, v/v) and mobile phase B was
10 mM ammonium formate with 0.1% formic acid in acetonitrile/isopropanol (10:90, v/v).
The gradient was ramped from 40% to 43% B in 1 min, ramped to 50% in 0.1 min,
ramped to 54% B in 4.9 min, ramped to 70% in 0.1 min, and ramped to 99% B in 2.9
min. The gradient was returned to initial conditions in 0.5 min and held for 1.6 min for
column equilibration. The flow rate was 0.4 mL/min. The column was maintained at
55°C and the auto-sampler was kept at 5°C. A 2 µL injection was used for all samples.
LC-MS data from the iterative MS/MS workflow was analyzed for lipid identification via
Agilent's Lipid Annotator (version 1.0). Positive and negative ion mode adducts included
$[M+H]^+$, $[M+Na]^+$, $[M+NH_4]^+$, $[M-H]^-$, and $[M+CH_3CO_2]^-$, respectively. The LC-MS data
from the MS[1] workflow were processed using Agilent's MassHunter Profinder (version
10.0).
554
Characterization of MDNP protein composition and MDNP protein corona
fingerprints in healthy and LPS plasma

To form the MDNP protein corona, purified MDNP were incubated with plasma
isolated from healthy and LPS-treated C57BL/6J mice for 4 hrs at 37°C. MDNP bound
to proteins were washed and centrifuged 3 times at 13,000 xg, 4°C with 1x cold PBS.
Then it was evaluated using the below protocol. Prepared samples were lysed in a lysis
buffer containing 5% sodium dodecyl sulfate (Sigma, St. Louis, MO), 50 mM
triethylammonium bicarbonate (1M, pH 8.0) (Sigma, 7408). Proteins were extracted and
digested using S-trap micro columns (ProtiFi, Farmingdale, NY). The eluted peptides
from the S-trap column were dried, and peptide concentration was determined using a
BCA assay kit (Thermo Fisher Scientific, 23275), after reconstitution in 0.1% formic
acid. All tryptic peptides were separated on a nanoACQUITY Ultra-Performance Liquid
Chromatography analytical column (BEH130 C18, 1.7 µm, 75 µm x 200 mm; Waters
Corporation, Milford, MA, USA) over a 185-min linear acetonitrile gradient with 0.1%
formic acid on a nanoACQUITY Ultra-Performance Liquid Chromatography system
(Waters Corp, Milford, MA) and analyzed on a coupled Orbitrap Fusion Lumos Tribrid
mass spectrometer (Thermo Scientific, San Jose, CA). Full scans were acquired at
240,000 resolutions, and precursors were fragmented by high-energy collisional
dissociation at 35% for up to 3 sec. MS/MS raw files were processed using Thermo
Proteome Discoverer (PD, version 3.0.0.757) with the Sequest HT search engine
against the plant metagenome database and UniProt mouse reference proteome.
Trypsin was used with a maximum of two missed cleavages, and peptide lengths were
restricted to 6-144 amino acids. Label-free quantification was performed using the
Minora feature detector. Protein identification was filtered at a 1% false discovery rate
across PSM, peptide, and protein levels using the Percolator algorithm in PD. Exported
protein abundances were analyzed with Perseus software (version 1.6.14.0), with
further filtering to include only proteins identified without missing values across all
samples. The quantitative protein data were log_2 transformed and further normalized
using median centering. Two-tailed student's t-test (adjusted p-value < 0.05) was
applied to determine the differentially expressed proteins (DEPs). Plant data was
analyzed with library from Uniprot and the Plant Proteome Database (Lepidium meyenii:
taxonomy ID 153348, entry: 92 sequences; Plant metagenome: taxonomy ID 1297885,
entry: 27312 sequences; Zingiberales: taxonomy ID 4618, entry: 357900) Ingenuity
Pathway Analysis (Qiagen) was utilized to identify canonical pathways, biological
function, upstream regulators and disease association.
590
Cell culture
The generation of bone marrow-derived macrophages (BMDM) followed a
previously published method(Truong et al., 2024). As established, the tibia and femur
from a C57BL/6J mouse was isolated by removing bulk muscles and connective tissues.
RPMI media (supplemented with L-glutamine, penicillin (100 units/mL), streptomycin
(100 µg/mL), 10% heat-inactivated FBS, and 20% L929 cell-conditioned media) drawn

needles were inserted into bones to flush the marrow into a 10 cm petri dish. The
collected bone marrow was filtered to grow in uncoated 10 cm cell culture plates.
BMDM were cultured in RPMI media conditioned with L929 at 37°C, 5% CO_2. The media
was replaced every 3 days. On Day 8-10, BMDM were lifted using Versene (Gibco,
Grand Island, NY), to be used for subsequent experiments. Trypan blue solution was
used to determine the cell number and viability with an EVE^{TM} Automated Cell Counter
(NanoEntek, Waltham, MA). RAW-blue cells were also cultured to confluency in 75 cm^2
flasks in identical incubation condition with Dulbecco's Eagle Medium (DMEM)
supplemented with penicillin (100 U/mL), streptomycin (100 U/mL), and heat inactivated
fetal bovine serum (10%).
607
Mice
Male C57BL/6 (6 to 8-week-old) purchased from the Jackson Laboratory (Bar
Harbor, ME) were maintained in cages at ambient temperature, 55% relative humidity,
and under a 12 hr dark/light cycle. LPS-induced endotoxemia model: Mice were
challenged with 5 mg/kg LPS intraperitoneally (IP) prior to treatment with two doses of 2
mg MDNP at 30 min and 2 hr *via* both IP and intravenous (IV) injection. Cecal ligation
and punction (CLP) model: Mice were lightly anesthetized with a mixture of ketamine
(75 mg/kg) and xylazine (15 mg/kg) at 1:1 ratio. After abdominal fur was removed, a
small incision was made to expose the cecum. The cecum was then ligated with a silk
suture and perforated with 19-gauge needle. A small amount of feces was extruded by
gently squeezing the cecum, the cecum was replaced, and the abdomen was sutured
after the bowel was repositioned. Experimental groups were IP injected with three
doses of 2 mg MDNP at 30 min, 2 hr, and 24 hr timepoint. All experiments were
performed in compliance with the protocol by the University of Maryland, Baltimore
Institutional Animal Care and Use Committee (IACUC) as well as the ARRIVE
guidelines.
624
Cytotoxicity assay
To measure the cytotoxicity of MDNP, a PI/Annexin V-FITC apoptosis assay was
conducted using BMDM. The assay utilizes propidium iodide (PI) staining to distinguish
between early and late-stage apoptotic cells. FITC labeling allow for the visualization of
Annexin V-bound cells using flow cytometry. Cultured BMDM at a density of 1.0 x 10^5
per well were treated with MDNP at 1 mg/mL for 8 hrs, then the cells were harvested
with Versene after washing with PBS. Collected cells were centrifuged at 500 x g for 5
min and cell pellets were resuspended in 1x binding buffer. 100 µL of cell suspension in
flow cytometry tubes were added with Annexin V-FITC and PI to be analyzed by flow
cytometry. The data was analyzed by De Novo software FCS Express 7 (Dotmatics,
Boston, MA).
636

In vitro cellular internalization

637
BMDM were seeded overnight in Falcon Culture slides at a density of 0.5 x 10^5
per well. Cy5.5 was incubated with MDNP under shaking for 30 min in the dark to label
the MDNP and excessive Cy5.5 dye was removed by 20 min centrifugation at 13,000 x
g prior to use. Then, 50 µL of Cy5.5-labled MDNP were added to 0.5 mL of culture
medium and incubated with cells for 0, 4, 8 hrs. Negative control, untreated cells were
used to establish the 0 hr timepoint. At the determined timepoints, cells were washed
with PBS three times, fixed with 4% paraformaldehyde (PFA) for 10 min, and
dehydrated with acetone for 5 min at -20°C. After blocking the culture with 1% BSA in 1x
PBS 30 min, the cells were washed again with 1x PBS and treated with 100 µL of
fluorescein isothiocyanate (FITC)-labeled phalloidin (1:50 dilution in PBS) for 30 min to
stain F-actin. The cells were then washed two times with 1x PBS and dried in the dark
condition and glass cover slips were mounted with mounting medium containing 4',6-
diamidino-2-phenylindole (DAPI) after carefully removing the plastic chambers. The final
fluorescence images were captured using an Olympus fluorescence microscope (Tokyo,
Japan) equipped with Hamamatsu Digital Camera ORCA-03G and Nikon software was
used to analyze the image data.

Flow cytometric phenotyping of MDNP-treated macrophages

655
BMDM were seeded at a density of 1.0 x 10^5 per well were in a sterile 24 well
plate. Cells were treated with 100 ng/mL of LPS in complete media for 24 hrs to induce
inflammation. Then, 100 µg/mL and 200 µg/mL MDNP was treated for 8 hrs. Cells were
resuspended in MACS buffer (PBS pH 7.2 supplemented with 1% FBS and 0.4% 0.5 M
EDTA, Quality Biological, Gaithersburg, MD) and transferred to the flow cytometry tubes.
FcR blocking (CD16/32, Biolegend, San Diego, CA) was performed and cells were
stained with following surface marker antibodies: Live/Dead fixable green, F4/80,
CD11b, MHCII, CD206, CD80, and CD86 (Biolegend, San Diego, CA). Samples were
analyzed using Cytek Aurora 3 (Fremont, CA). FCS Expression 7 Flow Cytometry De
Novo Software was used for flow data processing. Mean fluorescence intensity (MFI)
was generated by GraphPad Prism Software 10.1.1.

In vitro pro-inflammatory cytokine sequestration and effects on NF-κB activity

668
Anti-inflammatory properties of MDNP were investigated in two ways;
prophylactic and therapeutic. For prophylactic treatment, BMDM at a density of 1.0 x
10^5 per well were treated with 100 µg/mL MDNP for 8 hrs and 100 ng/mL of LPS in
complete media in 24 well plates (Corning, Corning, NY) were incubated for 4 hrs. For
therapeutic study, BMDM in identical culture condition as above were treated with 100
ng/mL of LPS for 4 hrs and 100 µg/mL of MDNP was subsequently added and
incubated for 8 hrs. Supernatants were then collected to measure the reduction of pro-
inflammatory cytokines including IL-6, TNF-α, IL-1β, and IFN-γ by enzyme-linked

immunosorbent assay (ELISA) (BioLegend, San Diego, CA). We also assessed the
ability of MDNP to modulate cytokine levels in the absence of cells. The supernatants
from another set of LPS-treated BMDM were collected and incubated with 100 µg/mL of
MDNP in a time dependent manner (0, 0.5, 1, 4, 8, 12, and 24 hrs). Lastly, we confirmed
the direct ability of MDNP to sequester IL-6 by incubating 5 ng/mL of IL-6 recombinant
protein in DMEM with 100 µg/mL of MDNP for 8 hrs. Remaining IL-6 levels were
measured using ELISA.
An NF-κB reporter cell line, RAW-Blue, was used to determine the ability of
MDNP to alter NF-κB activity due to LPS stimulation. The cells were plated at a density
of 1.0 x 10^5 per well and were stimulated with LPS for 4 hrs and treated with 100 µg/mL
MDNP for 8 hrs. Then, QUANTI-Blue was added to the collected supernatant in flat
bottom 96 well plate. It was incubated at 37ºC for 1 hr, then secreted embryonic alkaline
phosphatase (SEAP) reporter was detected using a spectrophotometer at 620-655 nm.
Poly(lactic-co-glycolic acid) (PLGA) particles of similar size as MDNP were used as a
control and fabricated using a microfluidics device as we previously described(Truong et
al., 2021). The size and zeta potential of the PLGA was measured 163.8 nm and -39.3
mV, respectively.
694
***Ex vivo* cytokine sequestration assay**
To evaluate the cytokine sequestration ability of MDNP in mouse plasma *ex vivo*,
100 µg/mL of MDNP in PBS was prepared and added to 100 µL of diluted LPS-treated
or CLP mouse plasma to ultracentrifuge tubes. The mixture tubes were incubated at
37ºC for 8 hrs. After incubation, samples were centrifuged at 2,000 x g for 10 min to
pellet the cytokine bound MDNP. Then the supernatant was collected to measure IL-6
and TNF-α levels using ELISA (Biolegend, San Diego, CA).
702
***In vivo* biodistribution and anti-inflammatory effect following LPS challenge**
The *in vivo* biodistribution study was performed by intraperitoneal (IP) or
intravenous (IV) injection of MDNP that were labeled with the near-infrared fluorescent
Cy5.5 dye. Mice were administered 2 mg of Cy5.5-MDNP twice (0.5 and 2 hr time
point)-post LPS challenge (5 mg/kg) and euthanized after 4 hrs to collect blood by
cardiac puncture and organs including liver, lung, spleen, heart, and kidneys. The
fluorescence of the organs was imaged using an *in vivo* imaging system (IVIS) with
emission (720 nm) upon laser excitation (675 nm). Blood samples were centrifuged at
1,000 x g for 20 min at 4°C in microtainer capillary blood collection plasma tubes. The
plasma samples were used to measure the alteration of systematic cytokines levels
using a Luminex MAGPIX System (Luminex, Austin, TX). The 7-plex panel included
murine IL-6, TNF-α, MCP-1, IL-10, IL-1β, IFN-β, and IFN-γ. Blood biochemistry was also
measured for alanine aminotransferase (ATL/SGPT), aspartate aminotransferase
(AST/SGOT), creatine (CREA), blood urea nitrogen (BUN), total protein (TP), and

globulin, and albumin (VRL Animal Health Diagnostics, Gaithersburg, MD). The
collected organs were fixed in 4% formalin immediately after euthanasia and paraffin-
embedded for sectioning. Hematoxylin & eosin (H&E) staining was performed using
standard procedures by the Pathology Biorepository Shared Services Core at the
University of Maryland, Baltimore. The level of injury scores was calculated based on
the scoring criteria: score 0, no damage; score 1, >10% slight inflammation; score 2, 10-
25% mild; score 3, 26-50% moderate; score 4, 51-75% severe; score 5, >75% necrosis.
724

In vivo sepsis survival study

The *in vivo* survival study was carried out using two representative sepsis models:
a lethal LPS-induced endotoxemia and CLP polymicrobial mouse model. For
endotoxemia, C57BL/6J mice were challenged with 20 mg/kg LPS and two doses
MDNP (2 mg) administered *via* IP injection. For CLP model, the cecum of anesthetized
C57BL/6J mice was perforated to release fecal materials into peritoneal cavity to
generate an exacerbated polymicrobial infection. Three doses of MDNP (2 mg/injection)
were administered via IP route. Survival for each model was evaluated over the course
of 5 days. Body temperature was also recorded every day for CLP mice.
734

Statistical analysis

Data evaluation was performed using GraphPad Prism Software 10.1.1 (San
Diego, CA). One-way ANOVA and t-test for unpaired data were employed to evaluate
statistical significance along with Tukey's multiple comparison test. Kaplan-Meier
survival curve and statistical significance of mouse survival were determined with a
long-rank (Mantel-Cox) X^2 test. Significant differences were indicated as $p<0.05$,
$p<0.005$, $p<0.0005$ related to control unless otherwise stated.
742

743 ### Data availability statement

744   All lipidomics and proteomics raw data are included in the supporting information.
745 Additional datasets supporting the findings are available upon reasonable request from
746 the corresponding author.
747

Acknowledgements

This work was supported by Startup funds from the University of Maryland
School of Pharmacy, the National Institute of General Medical Sciences of the National
Institute of Health under Award number R35GM142752 awarded to R.M.P. A.L.C. was
supported by a pharmaceutical sciences (PSC) departmental fellowship. This
publication was supported by funds through the Maryland Department of Health's
Cigarette Restitution Fund Program and the National Cancer Institute – Cancer Center
Support Grant (CCSG) – P30CA134274. University of Maryland School of Medicine's &
Greenebaum Comprehensive Cancer Center's Flow Cytometry Core–Baltimore,
Maryland. University of Maryland School of Medicine's Center for Translational

Author contributions
Junsik J. Sung: Conceptualization, Formal analysis, Investigation, Methodology,
Visualization, Writing – original draft, Writing – review & editing. **Jacob R. Shaw:**
Methodology, Writing – review & editing. **Josie D. Rezende:** Investigation, Methodology.
Shruti Dharmaraj: Investigation, Methodology. **Andrea L. Cottingham:** Investigation,
Methodology. **Mehari M. Weldemariam:** Investigation, Methodology. **Jace W.
Jones:** Investigation, Methodology, Resources, Supervision. **Maureen A. Kane:**
Investigation, Methodology, Resources, Supervision. **Ryan M. Pearson:**
Conceptualization, Funding acquisition, Methodology, Resources, Supervision, Writing –
original draft, Writing – review & editing.
773
774 **Declaration of interests**
775   J.J.S. and R.M.P are inventors on a patent application that describes the
776 isolation and use of MDNP for the treatment of inflammatory diseases.
777
778
779

References

Ansar, W., Ghosh, S., 2016. Inflammation and Inflammatory Diseases, Markers, and
Mediators: Role of CRP in Some Inflammatory Diseases, in: Ansar, W., Ghosh, S.
(Eds.), Biology of C Reactive Protein in Health and Disease. Springer India, New
Delhi, pp. 67–107.

Cao, F., Zhang, H., Yan, Y., Chang, Y., Ma, J., 2023. Extraction of polysaccharides from
Maca enhances the treatment effect of 5-FU by regulating CD4+T cells. Heliyon 9,
e16495.

Castellheim, A., Brekke, O.-L., Espevik, T., Harboe, M., Mollnes, T.E., 2009. Innate
Immune Responses to Danger Signals in Systemic Inflammatory Response
Syndrome and Sepsis. Scand J Immunol 69, 479–491.

Cedervall, T., Lynch, I., Lindman, S., Berggård, T., Thulin, E., Nilsson, H., Dawson, K.A.,
Linse, S., 2007. Understanding the nanoparticle–protein corona using methods to
quantify exchange rates and affinities of proteins for nanoparticles. Proc Natl Acad
Sci 104, 2050–2055.

Chafe, S.C., 1969. The fine structure of the collenchyma cell wall. Planta 90, 12–21.

Chen, L., Deng, H., Cui, H., Fang, J., Zuo, Z., Deng, J., Li, Y., Wang, X., Zhao, L., 2018.
Inflammatory responses and inflammation-associated diseases in organs.
Oncotarget 9, 7204–7218.

Corbo, C., Molinaro, R., Parodi, A., Toledano Furman, N.E., Salvatore, F., Tasciotti, E.,
2016. The Impact of Nanoparticle Protein Corona on Cytotoxicity, Immunotoxicity
and Target Drug Delivery. Nanomedicine 11, 81–100.

Dufour, J.-F., Anstee, Q.M., Bugianesi, E., Harrison, S., Loomba, R., Paradis, V., Tilg, H.,

Wong, V.W.-S., Zelber-sagi, S., 2022. Current therapies and new developments in

NASH. Gut 71, 2123.

Ehlting, C., Wolf, S.D., Bode, J.G., 2021. Acute-phase protein synthesis: a key feature

of innate immune functions of the liver. Biol Chem 402, 1129–1145.

Fang, X., Feng, J., Zhu, X., Feng, D., Zheng, L., 2024. Plant-derived vesicle-like

nanoparticles: A new tool for inflammatory bowel disease and colitis-associated

cancer treatment. Mol Ther 32, 890–909.

Fei, W., Hou, Y., Yue, N., Zhou, X., Wang, Y., Wang, L., Li, A., Zhang, J., 2020. The

effects of aqueous extract of Maca on energy metabolism and immunoregulation.

Eur J Med Res 25, 24.

Fei, W., Zhang, J., Yu, S., Yue, N., Ye, D., Zhu, Y., Tao, R., Chen, Yan, Chen, Yawen, Li,

A., Wang, L., 2022. Antioxidative and Energy Metabolism-Improving Effects of

Maca Polysaccharide on Cyclophosphamide-Induced Hepatotoxicity Mice via

Metabolomic Analysis and Keap1-Nrf2 Pathway. Nutrients 14.

Hirano, T., 2021. IL-6 in inflammation, autoimmunity and cancer. Int Immunol 33, 127–

148.

Hotchkiss, R.S., Monneret, G., Payen, D., 2013. Sepsis-induced immunosuppression:

from cellular dysfunctions to immunotherapy. Nat Rev Immunol 13, 862–874.

Ikoba, U., Peng, H., Li, H., Miller, C., Yu, C., Wang, Q., 2015. Nanocarriers in therapy of

infectious and inflammatory diseases. Nanoscale 7, 4291–4305.

Jarczak, D., Kluge, S., Nierhaus, A., 2021. Sepsis—Pathophysiology and Therapeutic

Concepts. Front Med (Lausanne) 8.

Jee, W., Jo, S., Lee, J.B., Jin, Y., Jeong, T., Yoon, J.C., Park, B., 2020. Mortality

difference between early-identified sepsis and late-identified sepsis. Clin Exp

Emerg Med 7, 150–160.

Ju, Y., Kelly, H.G., Dagley, L.F., Reynaldi, A., Schlub, T.E., Spall, S.K., Bell, C.A., Cui, J.,

Mitchell, A.J., Lin, Z., Wheatley, A.K., Thurecht, K.J., Davenport, M.P., Webb, A.I.,

Caruso, F., Kent, S.J., 2020. Person-Specific Biomolecular Coronas Modulate

Nanoparticle Interactions with Immune Cells in Human Blood. ACS Nano 14,

15723–15737.

Khatoon, N., Zhang, Z., Zhou, C., Chu, M., 2022. Macrophage membrane coated

nanoparticles: a biomimetic approach for enhanced and targeted delivery. Biomater

Sci 10, 1193–1208.

Koide, H., Okishima, A., Hoshino, Y., Kamon, Y., Yoshimatsu, K., Saito, K., Yamauchi, I.,

Ariizumi, S., Zhou, Y., Xiao, T.-H., Goda, K., Oku, N., Asai, T., Shea, K.J., 2021a.

Synthetic hydrogel nanoparticles for sepsis therapy. Nat Commun 12, 5552.

Koide, H., Okishima, A., Hoshino, Y., Kamon, Y., Yoshimatsu, K., Saito, K., Yamauchi, I.,

Ariizumi, S., Zhou, Y., Xiao, T.-H., Goda, K., Oku, N., Asai, T., Shea, K.J., 2021b.

Synthetic hydrogel nanoparticles for sepsis therapy. Nat Commun 12, 5552.

Lasola, J.J.M., Kamdem, H., McDaniel, M.W., Pearson, R.M., 2020. Biomaterial-Driven

Immunomodulation: Cell Biology-Based Strategies to Mitigate Severe Inflammation

and Sepsis. Front Immunol 11.

Li, X., Bao, H., Wang, Z., Wang, M., Fan, B., Zhu, C., Chen, Z., 2018. Biogenesis and

function of multivesicular bodies in plant immunity. Front Plant Sci 9.

Lima, A.C., Cunha, C., Carvalho, A., Ferreira, H., Neves, N.M., 2018. Interleukin-6

Neutralization by Antibodies Immobilized at the Surface of Polymeric Nanoparticles

as a Therapeutic Strategy for Arthritic Diseases. ACS Appl Mater Interfaces 10,

13839–13850.

Liu, C., Zhao, X.-M., Wang, Q., Du, T.-T., Zhang, M.-X., Wang, H.-Z., Li, R.-P., Liang, K.,

Gao, Y., Zhou, S.-Y., Xue, T., Zhang, J.-G., Han, C.-L., Shi, L., Zhang, L.-W., Meng,

F.-G., 2023. Astrocyte-derived SerpinA3N promotes neuroinflammation and

epileptic seizures by activating the NF-κB signaling pathway in mice with temporal

lobe epilepsy. J Neuroinflammation 20, 161.

Liu, T., Zhang, L., Joo, D., Sun, S.-C., 2017. NF-κB signaling in inflammation. Signal

Transduct Target Ther 2, 17023.

Markanday, A., 2015a. Acute Phase Reactants in Infections: Evidence-Based Review

and a Guide for Clinicians. Open Forum Infect Dis 2, ofv098.

Markanday, A., 2015b. Acute Phase Reactants in Infections: Evidence-Based Review

and a Guide for Clinicians. Open Forum Infect Dis 2, ofv098.

Massart, I.S., Paulissen, G., Loumaye, A., Lause, P., Pötgens, S.A., Thibaut, M.M.,

Balan, E., Deldicque, L., Atfi, A., Louis, E., Gruson, D., Bindels, L.B., Meuwis, M.-A.,

Thissen, J.-P., 2020. Marked Increased Production of Acute Phase Reactants by

Skeletal Muscle during Cancer Cachexia. Cancers (Basel) 12.

Matyash, V., Liebisch, G., Kurzchalia, T. V, Shevchenko, A., Schwudke, D., 2008. Lipid

extraction by methyl-tert-butyl ether for high-throughput lipidomics. J Lipid Res 49,

1137–1146.

Mera, S., Tatulescu, D., Cismaru, C., Bondor, C., Slavcovici, A., Zanc, V., Carstina, D.,

Oltean, M., 2011. Multiplex cytokine profiling in patients with sepsis. APMIS 119,

155–163.

Messina, J.M., Luo, M., Hossan, M.S., Gadelrab, H.A., Yang, X., John, A., Wilmore, J.R.,

Luo, J., 2024. Unveiling cytokine charge disparity as a potential mechanism for

immune regulation. Cytokine Growth Factor Rev 77, 1–14.

Miller, A.H., Raison, C.L., 2016. The role of inflammation in depression: from

evolutionary imperative to modern treatment target. Nat Rev Immunol 16, 22–34.

Miller, S.D., Turley, D.M., Podojil, J.R., 2007. Antigen-specific tolerance strategies for

the prevention and treatment of autoimmune disease. Nat Rev Immunol 7, 665–677.

Nyati, K.K., Masuda, K., Zaman, M.M.-U., Dubey, P.K., Millrine, D., Chalise, J.P., Higa,

M., Li, S., Standley, D.M., Saito, K., Hanieh, H., Kishimoto, T., 2017. TLR4-induced

NF-κB and MAPK signaling regulate the IL-6 mRNA stabilizing protein Arid5a.

Nucleic Acids Res 45, 2687–2703.

Parisi, L., Gini, E., Baci, D., Tremolati, M., Fanuli, M., Bassani, B., Farronato, G., Bruno,

A., Mortara, L., 2018. Macrophage Polarization in Chronic Inflammatory Diseases:

Killers or Builders? J Immunol Res 2018, 8917804.

Pearson, R.M., Juettner, V. V, Hong, S., 2014. Biomolecular corona on nanoparticles: a

survey of recent literature and its implications in targeted drug delivery. Front Chem

2.

Perut, F., Roncuzzi, L., Avnet, S., Massa, A., Zini, N., Sabbadini, S., Giampieri, F.,

Mezzetti, B., Baldini, N., 2021. Strawberry-Derived Exosome-Like Nanoparticles

Prevent Oxidative Stress in Human Mesenchymal Stromal Cells. Biomolecules 11.

Popa, C., Netea, M.G., van Riel, P.L.C.M., van der Meer, J.W.M., Stalenhoef, A.F.H.,

2007. The role of TNF-α in chronic inflammatory conditions, intermediary

metabolism, and cardiovascular risk. J Lipid Res 48, 751–762.

Robards, a.w., kidwai, p., 1969. vesicular involvement in differentiating plant vascular

cells. New Phytologist 68, 343–349.

Shao, H., Im, H., Castro, C.M., Breakefield, X., Weissleder, R., Lee, H., 2018. New

Technologies for Analysis of Extracellular Vesicles. Chem Rev 118, 1917–1950.

Shaw, J., Pearson, R.M., 2022. Nanoparticle personalized biomolecular corona:

implications of pre-existing conditions for immunomodulation and cancer. Biomater

Sci 10, 2540–2549.

Shen, H., Colangelo, C., Wu, T., Maffei, M., Goldstein, D.R., 2011. 235 Haptoglobin: A

Novel Innate Ligand that Induces Inflammation after Tissue Necrosis. J Heart Lung

Transplant 30, S84.

Shen, S., Han, F., Yuan, A., Wu, L., Cao, J., Qian, J., Qi, X., Yan, Y., Ge, Y., 2019.

Engineered nanoparticles disguised as macrophages for trapping

lipopolysaccharide and preventing endotoxemia. Biomaterials 189, 60–68.

Shi, C., Wang, X., Wang, L., Meng, Q., Guo, D., Chen, L., Dai, M., Wang, G., Cooney,

R., Luo, J., 2020a. A nanotrap improves survival in severe sepsis by attenuating

hyperinflammation. Nat Commun 11, 3384.

Shi, C., Wang, X., Wang, L., Meng, Q., Guo, D., Chen, L., Dai, M., Wang, G., Cooney,

R., Luo, J., 2020b. A nanotrap improves survival in severe sepsis by attenuating

hyperinflammation. Nat Commun 11, 3384.

Sung, J., Yang, C., Viennois, E., Zhang, M., Merlin, D., 2019. Isolation, Purification, and

Characterization of Ginger-derived Nanoparticles (GDNPs) from Ginger, Rhizome

of Zingiber officinale. Bio Protoc 9, e3390.

Tanaka, T., Narazaki, M., Kishimoto, T., 2014. IL-6 in Inflammation, Immunity, and

Disease. Cold Spring Harb Perspect Biol 6, a016295–a016295.

Tenci, B., Di Cesare Mannelli, L., Maresca, M., Micheli, L., Pieraccini, G., Mulinacci, N.,

Ghelardini, C., 2017. Effects of a water extract of Lepidium meyenii root in different

models of persistent pain in rats. Zeitschrift für Naturforschung C 72, 449–457.

Thamphiwatana, S., Angsantikul, P., Escajadillo, T., Zhang, Q., Olson, J., Luk, B.T.,

Zhang, S., Fang, R.H., Gao, W., Nizet, V., Zhang, L., 2017a. Macrophage-like

nanoparticles concurrently absorbing endotoxins and proinflammatory cytokines for

sepsis management. Proc Natl Acad Sci 114, 11488–11493.

Torres, A., Vivanco, S., Lavín, F., Pereda, C., Chernobrovkin, A., Gleisner, A., Alcota, M.,

Larrondo, M., López, M.N., Salazar-Onfray, F., Zubarev, R.A., González, F.E., 2022.

Haptoglobin Induces a Specific Proteomic Profile and a Mature-Associated

Phenotype on Primary Human Monocyte-Derived Dendritic Cells. Int J Mol Sci 23.

Truong, N., Black, S.K., Shaw, J., Scotland, B.L., Pearson, R.M., 2021. Microfluidic-

Generated Immunomodulatory Nanoparticles and Formulation-Dependent Effects

on Lipopolysaccharide-Induced Macrophage Inflammation. AAPS J 24, 6.

Truong, N., Cottingham, A.L., Dharmaraj, S., Shaw, J.R., Lasola, J.J.M., Goodis, C.C.,

Fletcher, S., Pearson, R.M., 2024. Multimodal nanoparticle-containing modified

suberoylanilide hydroxamic acid polymer conjugates to mitigate immune

dysfunction in severe inflammation. Bioeng Transl Med 9, e10611.

Unsinger, J., McGlynn, M., Kasten, K.R., Hoekzema, A.S., Watanabe, E., Muenzer, J.T.,

McDonough, J.S., Tschoep, J., Ferguson, T.A., McDunn, J.E., Morre, M., Hildeman,

D.A., Caldwell, C.C., Hotchkiss, R.S., 2010. IL-7 Promotes T Cell Viability,

Trafficking, and Functionality and Improves Survival in Sepsis. The J Immunology

184, 3768–3779.

Wang, Q., Zhuang, X., Mu, J., Deng, Z.-B., Jiang, H., Zhang, L., Xiang, X., Wang, B.,

Yan, J., Miller, D., Zhang, H.-G., 2013. Delivery of therapeutic agents by

nanoparticles made of grapefruit-derived lipids. Nat Commun 4, 1867.

Wang, S., Zhu, F., 2019. Chemical composition and health effects of maca (Lepidium

meyenii). Food Chem 288, 422–443.

Wu, H.-P., Chen, C.-K., Chung, K., Tseng, J.-C., Hua, C.-C., Liu, Y.-C., Chuang, D.-Y.,

Yang, C.-H., 2009. Serial cytokine levels in patients with severe sepsis. Inflamm

Res 58, 385–393.

Yang, C., Long, D., Sung, J., Alghoul, Z., Merlin, D., 2021. Orally Administered Natural

Lipid Nanoparticle-Loaded 6-Shogaol Shapes the Anti-Inflammatory Microbiota and

Metabolome. Pharmaceutics 13.

Yang, J., Cho, H., Gil, M., Kim, K.E., 2023. Anti-Inflammation and Anti-Melanogenic

Effects of Maca Root Extracts Fermented Using Lactobacillus Strains. Antioxidants

12.

Ye, R.D., Sun, L., 2015. Emerging functions of serum amyloid A in inflammation. J

Leukoc Biol 98, 923–929.

Zhang, Q., Hu, Chengyi, Feng, J., Long, H., Wang, Y., Wang, P., Hu, Chenglu, Yue, Y.,

Zhang, C., Liu, Z., Zhou, X., 2024. Anti-inflammatory mechanisms of neutrophil

membrane-coated nanoparticles without drug loading. J Control Release 369, 12–

24.

Zhang, Y., Ning, B., 2021. Signaling pathways and intervention therapies in sepsis.

Signal Transduct Target Ther 6, 407.

Zhu, H., Wang, R., Hua, H., Qian, H., Du, P., 2022. Deciphering the potential role of

Maca compounds prescription influencing gut microbiota in the management of

exercise-induced fatigue by integrative genomic analysis. Front Nutr 9.

(CD206) markers (G) A timeline for prophylactic treatment. Prophylactic approach did

not make significant change in levels of (H) IL-6 and (I) TNF-▯. (J) A timeline for

therapeutic treatment. Therapeutic treatment displayed significant reduction of (K) IL-6

and (L) TNF-▯. Data are expressed as mean ± SD (n=5). Data are expressed as mean

± SD (n=3). *P<0.05, **p<0.05 versus LPS control group.

Figure 3. *In vitro* sequestration of pro-inflammatory cytokines from LPS-induced bone

marrow derived macrophages. (A) Time course reduction of LPS-stimulated pro-

inflammatory cytokine by treatment of MDNP. 8 hr time point exhibited the most

significant reduction for both (B) IL-6 and (C) TNF-α. (D) Recombinant IL-6 was

sequestered through incubation with MDNP ex vivo without cellular interaction. (E) 100

µg/mL MDNP was the most effective concentration to remove IL-6 protein. (F)

Expression of NF-κB in in RAW-Blue 264.7 cell is suppressed by the treatment of MDNP.

(G) NF-κB activity level is downregulated after treatment of MDNP. PLGA was used as a

control to show different effect on NF-κB. All data are expressed as means ± SD (n=5).

**p<0.05, ***p<0.005, and ****p<0.0005 versus LPS or NT control group.

1009

Figure 4. Route of administration biodistribution and anti-inflammatory effects of MDNP

(A) Timeline of MDNP treatments following LPS challenge. The schematic illustrates the

IP and IV injection performed on randomly selected C57BL/6J mice groups (LPS only,

LPS+IP injection, and LPS+IV injection, n=3 per group). (B) Biodistribution of

fluorescently labeled MDNP in mouse organs (liver, kidney, spleen, lung, and heart). (C)

Fluorescence intensity measurement of MDNP biodistribution. (D) Heatmap of pro-

inflammatory cytokine expression including: (E) IL-6, (F) TNF-α, (G) MCP-1, (H) IL-10, (I)

IL-1β, (J) IFN-β, (K) GROα from mice plasma samples. Data are expressed as mean ±

SD (n=3). *P<0.05, **p<0.05, ***p<0.005, and ****p<0.0005 versus PBS control group.

Figure 5. Representative images of organ recovery and survival effect by treatment of

MDNP during LPS-induced sepsis. (A) H&E staining of liver, kidney, spleen, lung, and

heart. The LPS treated group images shows presence of neutrophil infiltration (yellow

arrow) in liver, tubular necrosis (white arrow) in the tubules in kidney, focal necrosis and

dysregulation in white pulp of spleen, appearance of extensive clots (red arrow) in the

alveoli in lungs, while no noticeable damage in heart. On the contrary, mice organs

administered with IP and IV displayed notably reduced level of organ damage of organ

damage. (B-F) Quantification of H&E staining. Levels of injury scores were calculated.

Data are expressed as mean ± SD (n=3). *P<0.05, **p<0.05 versus LPS control group.

(G) A schematic of survival study of LPS-treated and LPS+MDNP-treated C57BL/6J

mice (n=10/group). (H) The survival analysis graph displays a significant difference

between LPS only and the MDNP treatment group. Kaplan-Meier curves and a log-rank

test was performed to compare the survival rates. ***p<0.0005.

Figure 6. Mapping and identification of protein corona adsorbed onto the surface of

MDNP. (A) A schematic showing the process of evaluating the MDNP protein corona. (B)

Venn diagram illustrating a total of 296 proteins detected including unique (1 in healthy

and 26 in LPS) and 270 shared proteins in the coronas. (C) Heatmap shows abundance

of 49 upregulated and 100 downregulated proteins in MDNP-LPS group compared to

MDNP-H healthy group. (D) Volcano plot of significantly upregulated and downregulated corona proteins. Hp, Serpina3n and Saa1 (red dots) are categorized as pro-inflammatory and acute phase response signaling proteins. (E) Top 10 canonical pathways associated with the significantly upregulated protein corona were identified, with acute phase response signaling being the most prominent pathway. (F) A network representation of acute phase response signaling proteins and major upstream regulators. (G) *Ex vivo* sequestration of pro-inflammatory cytokines by incubating LPS plasma with MDNP. Both IL-6 and TNF-α was significantly reduced by 8 hr treatment. (H) A graphical summary illustrating how the cytokine and acute phase protein corona-coated MDNP sequesters and neutralizes inflammatory proteins, thereby reducing overall inflammation.

Figure 7. Survival effects of MDNP treatment on severe CLP model. (A) Schematic timeline of survival study of cecal ligation and puncture (CLP) C57BL/6J murine model with MDNP treatment (n=5/group) based on the complement cascade pathway from proteomics. Mice underwent CLP surgery to induce severe polymicrobial inflammation and three separate doses of MDNP were administered intraperitoneally (0.5, 2, 24 hr timepoint, 2 mg/injection) to analyze survival. (B) *Ex vivo* sequestration test of CLP plasma and MDNP showed that both IL-6 and TNF-α was significantly reduced. Kaplan-Meier curves and a log-rank test was performed to compare the survival rates. (C) The survival test displayed a significant improvement (40%) by the treatment of MDNP **$p < 0.005$. Data are expressed as mean ± SD (n=5). ****$p<0.0005$ versus PBS control group.

1062
1063
1064
1065
1066   **Tables**
1067
1068   **Table 1.** A list of unique proteins that are part of the MDNP protein corona formed from LPS and healthy plasma.
1069   A total 26 proteins are derived from the LPS plasma protein corona, while only one unique protein is found in
1070   healthy plasma corona. Acute phase proteins are highlighted with circles to underscore the potential presence of
      pro-inflammatory proteins associated with MDNP.

Protein Corona	Accession	Description	Gene Symbol	Function	Acute Phase Protein
LPS Plasma	Q8BMT4	Transforming growth factor beta activator LRRC33	Nrros	Key regulator of TGFB1 required for microglia function in nervous system	
	Q9Z1T2	Thrombospondin-4	Thbs4	Adhesive glycoprotein that mediates cell-to-cell interaction	
	P39876	Metalloproteinase inhibitor 3	Timp3	Regulates hematopoietic stem cell proliferation, differentiation, and trafficking	
	P02089	Hemoglobin subunit beta-2	Hbb-b2	Oxygen, CO2 transport and pH regulation	
	P62806	Histone H4	H4	Central role in transciption regulation, DNA repair and replication	
	P10810	Monocyte differentiation antigen CD14	Cd14	Co-receptor for TLR4 and recognize LPS and PAMPs	O
	P11672	Neutrophil gelatinase-associated lipocalin	Lcn2	Regulates neutrophil recruitment and modulatess pro-inflammatory cytokines	O
	P14152	Malate dehydrogenase, cytoplasmic	Mdh1	Involved in Krebs cycle and plays critical role in cellular energy production	
	P12246	Serum amyloid P-component	Apcs	Acute-phase reactant that bind to complement protetins and damaged cell	O
	P04117	Fatty acid-binding protein, adipocyte	Fabp4	Regulates choesterol ester accumulation and inflammatory responses	O
	P32848	Parvalbumin alpha	Pvalb	Calcium-binding protein involved in regulation of Ca2+ levels in neurons to muscle	
	P70296	Phosphatidylethanolamine-binding protein 1	Pebp1	Regulate the trafficking and distribution of phosphotidylethanolamine within cells	
	Q99PT1	Rho GDP-dissociation inhibitor 1	Arhgdia	Inhibits GDP/GTP exchange activity of Rho GTPases	
	O08692	Neutrophilic granule protein	Ngp	Enzymatic activity that break down bacterial wall that damaged bacterial DNA	
	Q64339	Ubiquitin-like protein ISG15	Isg15	Antiviral defense by degradating viral proteins and modulate immune response	
	P09411	Phosphoglycerate kinase 1	Pgk1	Catalyze the reversible transfer of a phosphate group between 1,3-BPG and ADP	
	P48774	Glutathione S-transferase Mu 5	Gstm5	Catalyze the conjugation of GSH to electrophilic compound such as toxin and ROS	
	P70290	55 kDa erythrocyte membrane protein	Mpp1	Facilitate the exchange of Cl- and HCO2- across the erythrocyte membrane	
	P07901	Heat shock protein HSP 90-alpha	Hsp90aa1	Involved in protein folding, stability and trafficking	O
	P54116	Stomatin	Stom	Maintain mechanical stability of erythrocyte membrane	
	O89103	Complement component C1q receptor	Cd93	Initiate the classical pathway of complement activation	
	P28666	Murinoglobulin-2	Mug2	Involved in cell adhesion, cell signalig regulation and contribute to barrier function	
	P49222	Protein 4.2	Epb42	Maintain stability and integrity of erythrocyte membrane	
	Q8BFZ3	Beta-actin-like protein 2	Actbl2	Involved in assembly and regulation of actn cytoskeleton	
	Q8VDD5	Myosin-9	Myh9	Transport vesicles and organelles within the cell	
	P63242	Eukaryotic translation initiation factor 5A-1	Eif5a	Facilitate the intiation of protein translation on ribosomes	
Healthy Plasma	Q14B46	Rhotekin-2	Rtkn2	Key regulator of actin cytoskeleton, cell signaling and cell adhesion	