Identification of Ephrin type-B receptor 4 as a critical mediator of tissue fibrosis

Authors: Brian Wu[1,2,3], Starlee S. Lively[1,2], Noah Fine[1,2], Chiara Pastrello[1,2], Anca Maglaviceanu[1,2,3], Osvaldo Espin-Garcia[1,2,4,5,6], Evan Pollock-Tahiri[1,2], Sayaka Nakamura[1,2], Paramvir Kaur[1,2], Jason S. Rockel[1,2], Pratibha Potla[1,2], Himanshi Gupta[1,2], Poulami Datta[1,2], Laura Tang[1,2,3], Shabana Vohra[1], Akihiro Nakamura[1,2,7], Rajiv Gandhi[1,2,8], Jiangping Wu[9], Boris Hinz[10,11], Igor Jurisica[1,2,11,12,13] and Mohit Kapoor[1,2,3,8]*

Affiliations:
[1] Schroeder Arthritis Institute, University Health Network; Toronto, Ontario, Canada.
[2] Krembil Research Institute, University Health Network; Toronto, Ontario, Canada.
[3] Department of Laboratory Medicine and Pathobiology, University of Toronto; Toronto, Ontario, Canada.
[4] Department of Epidemiology and Biostatistics, Western University. London, Ontario, Canada
[5] Department of Biostatistics, University Health Network, Toronto, Ontario, Canada
[6] Dalla Lana School of Public Health and Department of Statistical Sciences, University of Toronto, Toronto, Ontario, Canada
[7] Department of Medicine, Division of Rheumatology, Queen's University, Kingston, ON, Canada.
[8] Department of Surgery, University of Toronto; Toronto, Ontario, Canada.
[9] Department of Medicine, University of Montreal Hospital Research Centre (CRCHUM); Montreal, Québec, Canada.
[10] Keenan Research Institute for Biomedical Science of the St. Michael's Hospital, Toronto, M5B 1T8, Ontario, Canada.
[11] Faculty of Dentistry, University of Toronto, Toronto, M5S 3E2 Ontario, Canada.
[12] Departments of Medical Biophysics and Computer Science, University of Toronto, Toronto, Ontario, Canada.
[13] Institute of Neuroimmunology, Slovak Academy of Sciences, Bratislava, Slovakia.

*Mohit.kapoor@uhnresearch.ca

Key Words: Pulmonary fibrosis, fibroblast, EphB4, idiopathic pulmonary fibrosis, cell biology

ABSTRACT

Pulmonary fibrosis (PF) is a pathology associated with interstitial lung diseases (ILDs), including idiopathic pulmonary fibrosis (IPF). Fibrosis promotes continual secretion of extracellular matrix (ECM), producing non-functional scar tissue and causing organ failure. This study investigated the tyrosine kinase receptor Ephrin type-B receptor 4 (EphB4) as a mediator of PF. To this end, we generated mice with conditional *Col1a2*-driven deletion of *Ephb4* and used a preclinical mouse model of PF, total and single nuclei RNA sequencing, NanoString, publicly available single cell data, computational analysis, and functional assays of mouse and human healthy control and IPF lung fibroblasts. *Col1a2*-Cre^{ERT}-driven *Ephb4* deletion, or EphB4 inhibition via NVP-BHG712, markedly protected against bleomycin-induced PF. Total RNA-sequencing of fibroblasts isolated from *Ephb4*-deficient fibrotic mouse lungs exhibited reduced expression of ECM, ER Cargo, and protein trafficking-related genes. NVP-BHG712 reduced expression of these identified genes in mouse lung fibroblasts under fibrotic conditions *in vitro*. Single nuclei RNA sequencing of mouse lungs treated with NVP-BHG712 identified transcriptomic changes of ECM genes in specific fibroblast subpopulations. Total RNA sequencing, computational, and functional assays using mouse and human IPF fibroblasts identified elastin as a key mediator involved in EphB4 signaling. Overall, we identified EphB4 as a crucial mediator of PF.

INTRODUCTION

Pathological overactivation of tissue repair facilitates fibrosis, causing irreversible tissue damage which can lead to organ failure and patient mortality[1]. Pulmonary fibrosis (PF) is characterized by excessive extracellular matrix (ECM) deposition that severely impairs oxygen exchange and deteriorates health. Idiopathic pulmonary fibrosis (IPF) and systemic sclerosis-related interstitial lung disease (SSc-ILD) are two diseases where PF is a primary pathology causing patient mortality[2]. Advancements in PF research have led to the approval of growth factor receptor inhibitors nintedanib and pirfenidone for the treatment of IPF, along with the subsequent repurposing of nintedanib for the treatment of progressing fibrotic ILDs[3-5]. Both nintedanib and pirfenidone inhibit multiple receptors in epidermal and connective tissue cells ('fibroblasts'), which has accelerated the investigation of epithelial and connective tissue in fibrogenesis [3-5]. Despite significant therapeutic advancements, current interventions neither halt nor reverse PF. Deciphering disease mechanisms underlying fibrosis will help uncover novel therapeutic avenues to improve patient outcomes.

Erythropoietin-producing hepatocellular receptor interacting protein (Ephrin) and Eph receptors are unique protein families that undergo bidirectional signaling, where ligand-receptor interactions initiate downstream signaling from both the receptor (forward signaling) and ligand (reverse signaling)[6]. The Ephrin-B2 ligand and EphB4 receptor signaling axis mediates angiogenesis and vasculogenesis in prenatal development by regulating cell migration via actin cytoskeletal remodeling pathways[6]. Further evidence suggests that Ephrin-B2 may play a key role in pulmonary, cardiac, skin, and hepatic fibrosis[7-9]. Release of the soluble Ephrin-B2 active ectodomain into lung tissue, modulated by a disintegrin and metalloproteinase domain-containing protein 10 (ADAM10), was shown to promote fibrosis[7]. Further, siRNA targeting of EphB4 abrogates soluble Ephrin-B2-mediated fibrotic responses in pulmonary fibroblasts *in vitro*[7]. While EphB4 was proposed as a target of Ephrin-B2-

mediated pro-fibrotic responses *in vitro*, to the best of our knowledge, the exact role of EphB4 signaling in PF is yet to be investigated.

To study the role of EphB4 receptor in PF, we used a preclinical PF mouse model, total and single nuclei RNA sequencing, NanoString, publicly available single cell data, integrative computational biology analysis, and functional assays using primary mouse and human lung fibroblasts (**Figure 1**). We generated mice with conditional deletion of *Ephb4* in *Col1a2*-expressing cells, including fibroblasts, to show that *Ephb4* deletion reduced the severity of bleomycin-induced PF. This protective antifibrotic phenotype was replicated using the pharmacological EphB4 inhibitor NVP-BHG712. High-throughput RNA sequencing of lung fibroblasts derived from bleomycin-treated mice showed that *Ephb4* deletion induced downregulation of key genes associated with ECM regulation, protein trafficking and endoplasmic reticulum (ER) cargo concentration. NanoString analysis on mouse lung fibroblasts treated with EphB4 inhibitor also indicated a reduction in the expression of key ECM, protein trafficking and ER cargo genes. Single nuclei RNA sequencing identified 4 fibroblast subclusters in the lungs of bleomycin-treated mice. Elastin (*ELN*) was identified as one of the top genes associated with fibroblast subcluster 1 and its expression was reduced with EphB4 inhibition in select fibroblast subsets under bleomycin condition. In alignment with our findings in an IPF mouse model, total RNA sequencing identified *ELN* as the most differentially expressed ECM gene in primary human lung fibroblasts isolated from IPF patients, which also exhibited significant downregulation in response to EphB4 inhibition. Computational analysis of integrated mouse and human fibroblast total RNA sequencing data identified *ELN* as a crucial downstream target of EphB4 signaling. TGF-β1 induced elastin expression was prominently reduced by inhibition or silencing of EphB4 in human IPF fibroblasts. Similarly, elastin was reduced in bleomycin-treated mouse lungs following

pharmacological inhibition or genetic *Ephb4* deletion. Overall, this study suggests a novel

mechanism by which EphB4 signaling contributes to PF.

RESULTS

Genetic deletion of *Ephb4* in *Col1a2*-expressing cells: generation of *Ephb4* conditional

knockout mice

As part of the pathology of PF, fibroblasts are responsible for the deposition of

excessive ECM components, like type I collagen. To examine whether *Ephb4* expression in

Col1a2-expressing cells (such as fibroblasts) contributes to PF development, floxed *Ephb4*

(*Ephb4*$^{tm1.1Jwu}$) mice were crossed with B6.Cg-Tg(*Col1a2*-Cre/ERT,-ALPP)7Cpd mice to

generate *Ephb4*$^{f/f}$ *Col1a2-Cre*^{ERT} mice (detailed pedigree for the generation of the mouse line

is included in **Supplementary Figure 1**). *Col1a2*-driven expression of Cre in mice has been

used in previous studies to ablate specific genes in fibroblasts[7,10,11]. PCR-genotyping was

performed to confirm the presence of *Col1a2-Cre*^{ERT} in mice exhibiting homozygosity of the

floxed *Ephb4* sites (500 bp) compared to the WT *Ephb4* gene (400 bp) (**Figure 2A**). At 4-5

weeks of age, *Ephb4*$^{f/f}$; *Col1a2-Cre*^{ERT} mice were treated with 4-hydroxytamoxifen (1 mg/day

for 5 days) to induce *Ephb4* gene deletion in *Col1a2*-expressing cells, including fibroblasts, or

with corn oil (vehicle). Sustained knockdown of *Ephb4* expression was confirmed at both

mRNA and protein levels in primary lung fibroblasts isolated 7 days after the final 4-

hydroxytamoxifen administration (**Figure 2B**). No significant reduction in the expression of

EphB4 was observed in other cell types examined, such as macrophages and endothelial cells

(**Supplementary Figure 2**). Thus, *Ephb4* conditional knockout mice permit 4-

hydroxytamoxifen-inducible deletion of *Ephb4* in *Col1a2*-expressing cells (herein referred to

as *Ephb4* CKO mice), including fibroblasts, compared to control mice treated with corn oil

(*Ephb4* control mice).

Ephb4 CKO mice are protected against the development of PF

Ephb4 CKO and _Ephb4_ control mice were subjected to intratracheal instillation of bleomycin to induce PF, with PBS used as a corresponding bleomycin control (**Figure 2C**). Mouse lungs were collected 14 days later to assess collagen content using Gomori's One Step Trichrome blue collagen stain. As shown in **Figure 2D**, collagen content was comparable in both _Ephb4_ CKO and _Ephb4_ control lungs treated with PBS. After bleomycin challenge, _Ephb4_-expressing control mice showed a striking increase in collagen staining, indicating significant development of PF. Interestingly, this effect was not readily observed in _Ephb4_ CKO mice treated with bleomycin. Instead, deletion of _Ephb4_ resulted in a marked decrease in PF severity and reduced the number of lobes exhibiting fibrosis, as determined by Ashcroft scoring and collagen staining quantification (**Figure 2D; Supplementary Table 1**)[12]. Next, immunohistochemical staining of alpha smooth muscle actin (α-SMA, gene name _Acta2_), a marker of activated fibroblasts associated with fibrosis[13], was examined. As expected, PBS control treatment yielded minimal α-SMA staining in mouse lungs of _Ephb4_ CKO and control mice. Bleomycin treatment induced strong α-SMA staining in _Ephb4_ control mice with substantially weaker staining in _Ephb4_ CKO mice (**Figure 2E**). Intriguingly, the antifibrotic phenomenon of _Ephb4_ deletion in lung fibroblasts after bleomycin challenge was maintained _in vitro_. Primary lung fibroblasts derived from bleomycin-treated _EphB4_ CKO mice also exhibited reduced α-SMA (_Acta2_) and type I collagen (_Col1a1_) gene expression compared to bleomycin-treated _EphB4_ control mice (**Figure 2F**). Taken together, it appears that _Ephb4_ in lung fibroblasts plays an important role in PF development.

We next wanted to determine if _Ephb4_ CKO had a similarly altered response of fibroblasts in another form of fibrosis. To this aim, we examined a skin fibrosis model. We first confirmed knockdown of _Ephb4_ expression at both mRNA and protein levels in skin fibroblasts from _Ephb4_ CKO and _Ephb4_ control mice (**Supplementary Figure 3A**). After

Ephb4 knockdown, mice received daily subcutaneous bleomycin injections (0.5 U/mL) for 21

days (**Supplementary Figure 3B**). Skin samples were then collected 22 days after *Ephb4*

knockdown and subjected to histological Gomori's trichrome staining. No differences were

observed in fibrosis severity and dermal thickening (**Supplementary Figure 3C**). Thus, the

antifibrotic effect of *Ephb4* CKO observed in PF was not observed in the bleomycin-induced

model of skin fibrosis, suggesting that the effects of *Ephb4* ablation in fibroblasts may depend

on the type of tissue/organ and the tissue microenvironment.

**Total RNA sequencing identifies downregulation of ECM, endoplasmic reticulum (ER)
cargo concentration and protein trafficking-related transcriptome in lung fibroblasts
from *Ephb4* CKO mice**

To elucidate the effects of *Ephb4* ablation on the lung fibroblast transcriptome in the

context of PF, total RNA sequencing was performed on lung fibroblasts isolated from *Ephb4*

CKO (or *Ephb4* control mice) 2 weeks after bleomycin (or PBS) challenge (**Figure 3A**). First,

the effects of bleomycin on lung fibroblast transcriptome were examined by comparing *Ephb4*

control mice (bleomycin vs. PBS) two weeks after bleomycin treatment. A total of 257

differentially expressed genes (DEGs; false discovery rate (FDR) q < 0.05) were identified

(155 upregulated and 102 downregulated) in bleomycin-treated *Ephb4* control mice compared

to PBS-treated *Ephb4* control mice. No DEGs were identified in lung fibroblasts from *Ephb4*

control and *Ephb4* CKO mice after PBS treatment. We next examined the effect of *Ephb4*

ablation on bleomycin-induced gene expression in lung fibroblasts. When Ephb4 CKO and

Ephb4 control mice challenged with bleomycin were compared, 317 DEGs were identified

(103 upregulated and 214 downregulated) in lung fibroblasts from bleomycin-challenged

Ephb4 CKO mice (**Figures 3B/C**), suggesting that the effects of *Ephb4* ablation in lung

fibroblasts are unique to fibrotic conditions. The full list of DEGs identified can be found in

Supplementary Table 2A-D and the full sequencing dataset can be accessed online (GSE205612).

Next, we used pathDIP[14] to identify unique and common pathways enriched in the up and downregulated DEG lists of *Ephb4* CKO lung fibroblasts relative to their *Ephb4* control counterparts under bleomycin challenge. A subset of 31 upregulated DEGs were uniquely enriched for 8 pathways, and a subset of 57 downregulated DEGs were uniquely enriched for 11 pathways (**Figure 3D**). Upregulated DEGs were uniquely linked to pathways associated with phospholipid and lipid metabolism, cytokine signaling and interleukin signaling. Among the downregulated genes, 11 were associated with ECM organization, 7 with protein trafficking, 4 with ER cargo concentration, and 12 with class I mediated antigen processing and presentation. The unique enriched gene-pathway associations were visualized in an interaction map using NAViGaTOR (**Figures 3E**)[15]. For the full list of enriched pathways see **Supplementary Table 3**.

The list of gene-pathway associations was used to identify cellular processes that *Ephb4* might regulate under fibrotic conditions. Unique DEGs downregulated in lung fibroblasts from *EphB4* CKO mice challenged with bleomycin were associated with ECM-relevant processes, including collagen formation and crosslinking, elastic fiber formation, fibronectin matrix formation or ECM organization as well as genes related to protein trafficking and ER cargo concentration related processes (**Figure 3F**). Interestingly, 11 downregulated DEGs associated with ECM organization identified, 9 have been documented as prominent profibrotic mediators, including: prolyl 4-hydroxylase beta polypeptide (*P4bh*), integrin b1 (*Itgb1*), integrin a5 (*Itga5*), peroxidasin (*Pxdn*), lysl oxidase like 2 (*Loxl2*), lysl oxidase (*Lox*), type XI collagen (*Col11a1*), tolloid like 1 (*Tll1*), and elastin (*Eln*)[16-21]. Among the ECM, ER cargo and protein trafficking genes, 20/22 exhibited a >1.5 fold change in gene expression in bleomycin-

challenged *Ephb4* CKO animals compared to bleomycin-challenged *Ephb4* control mice, with elastin (*Eln*) having the greatest fold change (**Figure 3G**).

Overall, total RNA sequencing showed that under fibrotic stimulation (bleomycin treatment), fibroblasts from *Ephb4* CKO mice compared to *Ephb4* control mice exhibit reduced expression of genes related to fibrosis, ECM organization, ECM remodeling, ER cargo concentration and protein trafficking.

Pharmacological EphB4 inhibition reduces mouse lung fibroblast expression of genes related to ECM organization, ER cargo concentration and protein trafficking

We next examined whether pharmacological inhibition of EphB4 elicited comparable changes in gene expression (as observed in total RNA sequencing) in primary mouse lung fibroblasts. WT primary lung fibroblasts were isolated from C57BL/6J mice and stimulated with TGF-β1 (or PBS as control) in combination with the EphB4 small molecule inhibitor NVP-BHG712 (or vehicle) (**Figure 4A)**. NVP-BHG712 has previously been shown to decrease EphB4 activity and phosphorylation *in vitro* and *in vivo,* including in mouse lung tissue[22]. We first confirmed that NVP-BHG712 effectively inhibits phosphorylation of EphB4 in human lung fibroblasts at 250 nM (**Supplementary Figure 4**) and used this concentration for our experiments. A NanoString CodeSet of 47 gene targets was designed using our previously identified downregulated DEGs involved in ECM organization, protein trafficking, ER cargo concentration, MHC Class I presentation and post-translational protein modification (remainders were excluded, including genes related to translation). TGF-β1 significantly increased the expression of 25 of the 47 selected genes, with EphB4 inhibition significantly reducing the expression of 19 out of 25 of these genes (see **Supplementary Table 4** for the full dataset). Notably, EphB4 inhibition downregulated many genes associated with cellular processes involved in ECM organization (9/11), ER cargo concentration (3/4), and protein trafficking (6/7) (**Figure 4B,** see **Supplementary Figure 5** for graphical representation of the

remaining dataset). Thus, pharmacological inhibition of EphB4 in mouse lung fibroblasts recapitulated reduced expression of most genes identified in our *in vivo Ephb4* CKO mouse lung fibrosis model.

NVP-BHG712 attenuates bleomycin-induced PF in mice

Since NVP-BHG712 attenuated the expression of key genes involved in ECM organization, ER cargo concentration and protein trafficking in cultured mouse lung fibroblasts, we next tested its anti-fibrotic potential *in vivo* using the bleomycin lung fibrosis mouse model. 7-week-old WT C57BL/6J mice were intratracheally challenged with bleomycin or PBS (control); and 5 days later, received NVP-BHG712 (10 µg/kg daily, *p.o.*) or vehicle (control; N-methyl-2-pyrrolidone plus polyethylene glycol 300 at ratio of 1:9) for 5 days. Lungs were harvested for Gomori's blue collagen stain histological assessment 14 days after the start of the bleomycin challenge (**Figure 5A**). As observed in *Ephb4* CKO mice, WT mice treated with NVP-BHG712 also exhibited reduced severity in the degree of bleomycin-induced lung fibrosis (**Figure 5B**). Ashcroft lung fibrosis score of affected areas showed reduced fibrosis severity in NVP-BHG712-treated mice, with a mean ± SD disease severity score of 2.48±0.55 compared to 4.05±0.61 in bleomycin challenged vehicle-treated lungs (**Figure 5C, left.** Histomorphometric analysis of lung section images indicated that lung collagen content in NVP-BHG712 treated lungs decreased from 47.2±10.8% to 37.8±8.2% per lobe with NVP-BHG712 treatment and 49.7±8.8% to 37.81±6.8% (mean ± SD) per NVP-BHG712-treated animal (**Figure 5C, middle/right**).). Notably, there was no significant change in Ashcroft scoring or lung collagen quantification in NVP-BHG712 treated mice compared to vehicle treated mice in the PBS-challenged groups. α-SMA staining was also markedly reduced in NVP-BHG712-treated lungs compared to vehicle-treated controls after bleomycin challenge, with no difference found between PBS-treated controls (**Figure 5D**). Collectively, these results

show that EphB4 pharmacological inhibition in mice confers protection in bleomycin-induced

lung fibrosis, similar to the genetic deletion in *Ephb4* CKO mice.

Transcriptomic differences in fibroblast subclusters using single nuclei RNA sequencing

of mouse lungs treated with NVP-BHG712 in the bleomycin model

Since NVP-BHG712 treatment resulted in reduced PF severity *in vivo* and reduced

expression of ECM genes in mouse lung fibroblasts *in vitro*, we next conducted single nuclei

RNA sequencing and bioinformatic analyses on bleomycin or PBS instilled mouse lungs

treated with NVP-BHG712 or vehicle to identify lung fibroblast subsets and interrogate the

differences in the ECM transcriptome of these fibroblast subsets in response to EphB4

inhibition. A total of 12 mouse lungs (n=3 PBS+vehicle, n=3 PBS+NVP-BHG712, n=3

bleomycin+vehicle and n=3 bleomycin+NVP-BHG712) were subjected to single nuclei RNA

sequencing. 34,816 nuclei were sequenced and, after filtering, 28,721 nuclei were analyzed

(**Figure 6A**). Multiple cell types, including fibroblasts, macrophages, endothelial cells, type I

alveolar cells, and type II alveolar cells, were identified based on canonical cell markers

(**Figure 6B&C**). We next subclustered fibroblasts across all 12 samples, identifying 4

fibroblast subclusters (**Figure 6D**). Top DEGs (p adj <0.05 and log2FC>0.5) associated with

each fibroblast subcluster are shown in **Supplementary Table 5**. Previous studies using single

cell RNA sequencing in the bleomycin lung fibrosis model have identified transcriptomic

profiles associated with activated fibroblasts, *Col13a1*-expressing fibroblasts, and *Col14a1*-

expressing fibroblasts[23,24]. Using the markers identified in these studies, we mapped the key

transcriptomic profiles to our four fibroblast subclusters. The majority of genes associated with

activated fibroblasts[23] mapped to fibroblast subcluster 1 (included *Fbn1*, *Eln* and *Tpm1*) or

subcluster 3 (included *Col1a1*, *Col1a2* and *Col3a1*) in our dataset (**Supplementary Figure 6**).

Key marker genes associated with *Col13a1* matrix fibroblasts[24] mapped primarily to subcluster

and 2, while the majority of the key marker genes associated with *Col14a1* expressing matrix fibroblasts mapped to fibroblast subcluster 3.

To identify differences in the transcriptome of fibroblast subclusters in response to EphB4 inhibition under bleomycin condition, we compared the fibroblast subclusters between NVP-BHG712 and vehicle control groups (**Figure 6E**). We identified the top expressing genes associated with each fibroblast sub-cluster under bleomycin condition (**Figure 6F**). Interestingly, *Eln* was identified as one of the top genes associated with fibroblast subcluster 1. Thus, we next investigated the expression of *Eln* in fibroblast sub-cluster 1, which showed downregulation in *Eln* expression in NVP-BHG712-treated bleomycin group compared vehicle-treated bleomycin group.In addition to fibroblast sub-cluster 1, *Eln* expression was also downregulated in sub-cluster 3 (**Supplementary Figure 7A**). We also investigated the expression of *Col1a1* in our fibroblast subclusters in response to NVP-BHG712 treatment under bleomycin condition (**Figure 6G**). Subcluster 3 fibroblasts exhibited highest expression of *Col1a1* and its expression was reduced with NVP-BHG712 treatment when compared to vehicle under fibrotic conditions. In addition to fibroblast sub-cluster 3, *Col1a1* expression was also downregulated in sub-cluster 1 (**Supplementary Figure 7A**). We also conducted gene ontology (GO) analysis on the DEGs that were upregulated or downregulated in NVP-BHG712-bleomycin group compared to vehicle-bleomycin group to identify the biological processes impacted by EphB4 inhibition under bleomycin conditions. Interestingly, biological processes such as elastic fiber formation, tissue remodeling and ECM assembly were identified as putative biological processes associated with downregulated genes in NVP-BHG712-bleomycin group compared to vehicle-bleomycin group (**Figure 6H**).

Further, Cell Chat was used to investigate the signaling interactions between each fibroblast subcluster in NVP-BHG712 compared to vehicle group under bleomycin condition. Interestingly, inhibition of EphB4 by NVP-BHG712 under bleomycin condition exhibited

reduced ligand-receptor signaling interactions in multiple fibroblast subclusters. Particularly, *Col1a1* associated ligand-receptor interactions (such as *Col1a1-Cd44* and *Col1a1*-integrins) were among the key downregulated signaling interactions across in fibroblast subclusters 1 and 3 in NVP-BHG712 versus vehicle under bleomycin condition (**Figure 6I** and **Supplementary Figure 7B/C**). Overall, our data have identified ECM transcriptomic changes associated with distinct fibroblast subsets in response to NVP-BHG712 treatment during fibrosis.

Transcriptomic differences in control donor vs IPF human lung fibroblasts treated with NVP-BHG712: identifying elastin as an EphB4 signaling partner

To investigate the relevance of EphB4 inhibition in human fibroblasts, we treated primary human lung fibroblasts from control donors (HLF) and IPF subjects (IPF) (**Supplementary Table 6)** with NVP-BHG712 or vehicle for 24 h and subjected each sample to total RNA sequencing (total 16 samples; n=4/group; **Figure 7A**). When IPF and HLF fibroblasts were compared under vehicle conditions, 3,108 DEGs (1,569 downregulated and 1,539 upregulated) were identified in the IPF group (**Figure 7B**). Notably, expression of *COL1A1* and some of the identified ECM genes were significantly increased, with *ELN* being determined as the most upregulated ECM gene. We also investigated another published IPF dataset[25] and found the expression of some of these genes to be elevated in IPF group (**Supplementary Figure 8)**.

Next, the transcriptomic differences elicited by EphB4 inhibition were determined by comparing IPF fibroblasts treated with NVP-BHG712 compared to corresponding vehicle controls. In total, 85 DEGs (24 upregulated and 61 downregulated) were identified (**Figure 7C**; the full list of DEGs is provided in **Supplementary Table 7**). Notably, expression of *ELN* (among other ECM genes) was significantly downregulated by NVP-BHG712 inhibition in IPF fibroblasts.

To identify common signaling partners involved in EphB4 signaling in mouse and human fibroblasts during lung fibrosis, the DEG list from human IPF RNA sequencing analysis was compared to the DEG list from *Ephb4* CKO mouse RNA sequencing. Four common gene targets were identified that included 2 upregulated genes [low density lipoprotein receptor (*LDLR*) and transmembrane protein 38b (*TMEM38B*)], and 2 downregulated genes [*ELN* and glycyl-TRNA synthetase 1 (*GARS1*)] **(Supplementary Table 8)**. In our pathDIP analysis, *LDLR* and *TMEM38B* did not associate with any pathways, *ELN* associated with ECM organization and elastin synthesis pathways, and *GARS1* associated with protein translation. Among these 4 genes, *ELN* stood out as a prime target for investigating EphB4 signaling since *ELN* was consistently downregulated with *Ephb4* CKO (total RNA sequencing) and pharmacological EphB4 targeting in both mouse (single nuclei RNA sequencing and NanoString) and human IPF fibroblasts (total RNA sequencing).

To further elucidate a potential relationship between EphB4 signaling and elastin, we queried physical protein-protein interactions (PPIs) between the two entities in the Integrated Interaction Database (**Figure 7D; Supplementary Table 9)**. We identified 4 proteins that can physically interact with EphB4 and elastin, including amyloid precursor protein (APP), fibronectin (FN1), perlecan (HSPG2), and TGF-β1. It is notable that a key pro-fibrotic cytokine TGF-β1 linked EphB4 to elastin[13]. Indeed, GO analysis also indicated that elastin, among 16 other DEGs from the human IPF total RNA sequencing data, was involved in TGF-β regulation of ECM (**Supplementary Figure 9** and **Supplementary Table 10)**. Overall, PPI analysis coupled with our sequencing data suggest that elastin may be a crucial signaling partner involved in EphB4 signaling during lung fibrosis.

Inhibition of EphB4 mitigates fibrotic upregulation of elastin in lung fibroblasts and fibrotic lung tissue

To further investigate the regulatory relationship between EphB4 and elastin, and their potential link to TGF-β1 signaling, lung fibroblasts from IPF patients were subjected to TGF-β1 stimulation followed by EphB4 inhibition using NVP-BHG712 or *EPHB4* siRNA treatment. As shown in **Figure 8A&B**, TGF-β1 stimulation markedly increased the expression of elastin in lung fibroblasts, while subsequent treatment with NVP-BHG712 or *EPHB4* siRNA moderately or significantly attenuated TGF-β1-induced elastin expression, respectively.

As a key regulatory cytokine for fibrogenesis, TGF-β1 typically relies on canonical signaling associated with phosphorylation of the transcription factors SMAD2/3. To assess whether SMAD2/3 activity impacts the regulatory effect of EphB4 on elastin expression, human lung fibroblasts were treated with TGF-β1 or control for 24 h, then *EPHB4* siRNA or control for an additional 24 h (**Figure 8C**). As predicted, TGF-β1 treatment increased phosphorylation of SMAD2/3 in human lung fibroblasts; however, silencing of *EPHB4* did not decrease phosphorylation levels of SMAD2/3, suggesting that EphB4 likely does not modify fibrotic pathways through canonical TGF-β1 signaling.

To assess the link between EphB4 and elastin *in vivo*, elastin staining, assessed using Verhoeff's Van Gieson elastin stain, was examined in the lungs of bleomycin-challenged *Ephb4* CKO and *Ephb4* control mice (**Figure 8D**) and NVP-BHG712-treated mice (**Figure 8E**). Both *Ephb4* CKO and NVP-BHG712 mouse lung tissue showed qualitative decreases in elastin protein after bleomycin challenge with no changes observed in PBS controls. Thus, our data suggest that EphB4 expression is crucial for elastin expression in the context of fibrosis in both mouse tissue and human lung fibroblasts. Overall, our data supports a mechanism involving EphB4-elastin signaling in lung fibrosis.

DISCUSSION

Ephrins and Ephs are key regulators of embryogenesis, prenatal development, and wound healing[26]. While Ephrin-B2 is known to participate in the dysregulation of tissue repair

during fibrosis[7,9,27], the role of EphB4 signaling in fibrosis remains unclear. To comprehensively explore the role of EphB4 receptor in PF development, we employed a multi-faceted approach that included the generation of *Ephb4* CKO mice in conjunction with a preclinical animal model of lung fibrosis, total and single nuclei RNA sequencing, NanoString, publicly available single cell data, advanced computational biology analysis, and functional assays using mouse and human (HLF and IPF) lung fibroblasts.

To our knowledge, we are the first to demonstrate that pharmacologic inhibition or genetic deletion of EphB4 reduces the expression of fibrotic ECM-related transcriptome and attenuates the development of PF. We showed that *Col1a2*-driven deletion of *Ephb4* CKO or EphB4 pharmacological inhibition using NVP-BHG712 attenuated bleomycin-induced PF *in vivo*. Our results indicated that fibrotic injury caused transcriptomic changes in lung fibroblasts, along with the transcriptomes of specific fibroblast subpopulations. Consistent with the reduced pathological histomorphometry, we also found that genetic or pharmacological targeting of EphB4 ameliorated fibrotic injury-induced changes through reduction in select ECM genes in specific fibroblast subpopulations. A subset of genes identified in our mouse studies were also elevated in human lung fibroblasts from IPF patients when compared to HLFs, with ELN identified as a prominent interaction partner of EphB4 signaling in mouse and human lung fibroblasts, and across genetic and pharmacological EphB4 targeting.

To date, the role of EphB4 in injury and fibrosis remain elusive, particularly in lung tissue. EphB4 mutations causing increased expression in lung tumors has been linked to critical cell processes including tumor cell survival and proliferation[28,29]. Interestingly, we found that reduction of EphB4 in lung tissue markedly protects against development of PF, and that this antifibrotic phenotype garners profound changes in the transcriptome (317 DEGs) of mouse pulmonary fibroblasts as determined though total RNA sequencing. A subset of the 214 downregulated DEGs were linked to ECM organization, protein transport, or ER cargo

regulation in fibroblasts. While the exact mechanisms of EphB4 signaling remains to be fully elucidated, downstream targets of EphB4 mediate cytoskeletal rearrangements via actin GTPase activity, which is typically associated with cell motility[27,30]. EphB4 deficiency has also been linked to accumulation of type IV collagen in endothelial cell ER through impaired protein trafficking, illuminating the potential of EphB4 as a regulator of protein transport and secretion[31]. However, our in vivo data also suggests that the role of EphB4 may be context specific, as we did not observe protection from bleomycin-induced skin fibrosis in *Ephb4* CKO mice. Overall, the cytoskeletal role of EphB4 signaling may intersect with ECM secretion and protein trafficking genes observed in our study, however further investigation is required to parse the biological and pathological signaling mechanisms for EphB4 in various tissues.

Recent single cell studies using a bleomycin-induced fibrosis model have identified activated fibroblast cell subsets and transcriptomic profiles that may play a key role in PF development[32-34]. Using single nuclei RNA sequencing, we identified four fibroblast subclusters within mouse lung tissues. Mapping of transcriptomic profiles associated with activated fibroblasts, *Col13a1*-expressing fibroblasts, and *Col14a1*-expressing fibroblasts, identified by previous studies, showed enrichment of these transcriptomic profiles in distinct fibroblast subsets[32-34]. Fibroblast subclusters 1 and 3 from our dataset were enriched with the majority of genes associated with the activated fibroblasts, while key transcriptomic profile associated with *Col14a1*-expressing fibroblasts were enriched in subcluster 3 of our dataset. Interestingly, *Col1a1,* considered a prominent ECM component of tissue fibrosis, mapped primarily to subcluster 3 fibroblasts and its expression was reduced with EphB4 inhibition. Furthermore, *Eln* was identified as one of top genes found to be associated with fibroblast subcluster 1 and its expression was reduced with EphB4 inhibition in this fibroblast subcluster. It's interesting to note the subclusters 1 and 3 which showed highest expression of *Eln* and *Col1a1*, respectively map with the transcriptome associated with previously activated

fibroblasts and *Col14a*, further indicating the importance of these subclusters in PF, which need to be further investigated. Furthermore, these ECM genes are also reduced with EphB4 inhibition in fibroblast subcluster 1 and 3, suggesting the effect of EphB4 inhibition on select ECM transcriptomes in distinct fibroblast populations.

Among the ECM genes identified in mouse lung tissue, a subset was also elevated in lung fibroblasts from IPF patients compared to donor controls; however, when IPF fibroblasts were treated with EphB4 inhibitor, *ELN* was the only ECM gene that was significantly downregulated. *ELN* also emerged as a gene that was consistently responsive to EphB4 deletion and inhibition across total RNA sequencing of IPF and control donor fibroblasts, total and single nuclei RNA sequencing of mouse lung fibroblasts, and NanoString in mouse lung fibroblasts. Our PPI results highlighted elastin as a key signaling interaction partner of EphB4 during PF. PF pathological impairment of tissue function is largely driven by excessive ECM accumulation, to which elastin reorganization and protein deposition are known indicators of disease in PF patients[19]. PPI and GO analyses suggested that EphB4 and elastin signaling both interact with TGF-β1. TGF-β1 is well established to increase elastin expression in lung fibroblasts[35]; however, we show that this induction is ameliorated by *EPHB4* silencing. Furthermore, both genetic and pharmacological blockade of EphB4 was sufficient to reduce elastin in mouse lung tissue following bleomycin-induced PF, further demonstrating a crucial link between ephB4 and elastin signaling during PF

In this study, we used the bleomycin-induced model of lung fibrosis. While the bleomycin preclinical model of lung fibrosis is important in understanding mechanisms of fibrosis, it does not fully capture all clinical manifestations of human IPF. For example, strong acute inflammatory phase observed in the bleomycin model is typically not present in human IPF patients[1,2]; thus, future studies should be directed towards understanding the role of Ephb4 signaling in other relevant animal models to further understand its role in PF.

Together, our *Ephb4* CKO, small molecule EphB4 inhibitor NVP-BHG712, and

EPHB4 siRNA data demonstrate an important role of EphB4 signaling in PF (**Figure 8F**).

Targeting EphB4 signaling requires extensive additional investigation due to the vital functions

of this molecule in developmental and physiological processes, including angiogenesis[36-38],

and blood pressure and heart rate regulation[39]. Additional studies are also warranted to

elucidate the fibrotic signaling pathways from EphB4 to the expression of its effector molecules

(e.g., elastin). Signaling molecules in these pathways may provide alternative therapeutic

targets for the treatment of PF and other pathological fibrotic conditions in which EphB4

signaling is implicated.

METHODS

Sex as a biological variable

For our animal studies, we used male and female animals for the conditional knockout mice

subjected to the bleomycin model of pulmonary fibrosis. For our human cell studies involving

RNA sequencing and functional assays, we obtained lung fibroblasts from both male and

female subjects.

Generation of *Ephb4* CKO mouse

Ephb4$^{tm1.1Jwu}$ mice, kindly donated by Dr. Jianping Wu (University of Montreal), were

crossed with B6.Cg-Tg(*Col1a2*-cre/ERT,-ALPP)7Cpd mice (The Jackson Laboratory) to

generate *Ephb4$^{loxP/loxP}$ Col1a2-Cre^{ERT}* mice on a C57B/6J background. Mouse lines were

backcrossed for 5 generations prior to the initiation of mouse experiments. To induce

conditional *Ephb4* knockdown, 4-hydroxytamoxifen was administered by intraperitoneal (IP)

injection (1 mg in 100 µL 20% ethanol/80% corn oil vehicle) for 5 consecutive days. Control

mice were administered vehicle comprising 20% ethanol and 80% corn oil. Lung fibroblasts

were isolated and cultured from *Ephb4* CKO mice and *Ephb4* control mice, and qRT-PCR and

Western blotting were performed to assess successful deletion of EphB4.

Mouse genotyping

Ear punches were collected, and alkaline lysis was performed at 90°C for 1 h to extract

DNA. DNA was subjected to PCR against the *Ephb4* and *Cre* genes (100 V for 35 min) and

gels were imaged using Sybr Safe in a BIO-RAD Chemidoc apparatus. Primers used can be

found in **(Supplementary Table 11)**.

Lung fibrosis model

To induce PF, bleomycin sulphate (1.2 U/kg in 50 µL PBS; Bioshop) was

intratracheally instilled in 6–7-week-old mice that were anesthetized with isoflurane (1.5%)

and xylazine/ketamine cocktail (10 mg/kg xylazine:100 mg/kg ketamine). Bleomycin control mice were administered PBS only. Instillation was initiated 7 min following xylazine/ketamine administration. Mice were sacrificed by anesthetization and CO_2 asphyxiation 14 days following bleomycin challenge. For experiments assessing the effects of NVP-BHG712 on the development of bleomycin-induced PF, mice were intratracheally administered bleomycin sulphate (1.2 U/kg) at 7 weeks of age. At 5 days post-bleomycin challenge, the treatment with NVP-BHG712 (Sigma Aldritch; cat # SML0333; 10 mg/kg in 150 µL N-methyl-2-pyrrolidone and polyethylene glycol 300 at ratio of 1:9) or vehicle control via oral gavage was initiated. Mice were given NVP-BHG712 once daily for a total of 5 days. Mice were sacrificed and lungs were collected 2 weeks after bleomycin challenge.

Skin Fibrosis Model

To induce skin fibrosis, mice were treated with bleomycin sulphate by subcutaneous injection. Briefly, bleomycin sulphate was reconstituted to 1 U/mL in PBS, aliquoted, and stored at -80°C. *Ephb4* CKO and *Ephb4* control mice at 8 weeks of age were anesthetized with xylazine and ketamine cocktail (10 mg/kg xylazine:100 mg/kg ketamine), and back hair was removed using Nair. Bleomycin stock solution was diluted to working concentration (0.1 U/mL in 100 µL volume) and mice were subcutaneously treated with bleomycin daily for 21 days. Mice were sacrificed 1 day following treatment end point and four skin sections were collected for each animal.

Lung histology

After mice were sacrificed, the heart was carefully removed and the lungs were inflated with 1 mL 10% formalin via the trachea, which was then tied off. The entire lung was excised, rinsed in PBS under gentle shaking for 1 min, and fixed in 10% formalin at room temperature with shaking for 48 h. After rinsing the lungs in PBS for 30 min, lobes were individually

excised, dehydrated in reagent alcohol for 1 min with vigorous agitation, and incubated for 24

h in a minimum of 50 mL JFC solution (Milestone) per mouse lung for completion of tissue

dehydration and tissue clearing. Processed lobes were paraffin-embedded with the ventral side

facing the surface of the paraffin block. For staining, 4 μm lung sections were collected after

removal of 80−150 μm of tissue. Gomori's One Step Trichrome (Thermo) was performed

according to manufacturer's instructions to assess lung collagen content. To assess the degree

of PF in affected lung tissue, Ashcroft scoring was performed across total lung sections with a

minimum of 50 non-overlapping fields of view[12]. For quantification of lung collagen staining

across total lung tissue, lung sections were imaged at 4X magnification and stitched together

using Autostitch[40] to form full lobe images. Lobes were colour corrected in GIMP image

editing software V.2.10.34. Quantification of lung collagen staining was performed in Fiji V.

2.9.0 [41]. Colour deconvolution was performed selecting the preset Mason's Trichrome filter to

separate blue, green and red colour channels[42]. The blue colour channel was selected and

thresholding was performed using the filter "mean". Normalization to total lung tissue was

performed by converting total lobe images to RGB stacks and quantifying the total pixel count

using the threshold filter "default". Lung collagen was expressed as percent lung collagen (%)

by normalizing the blue colour channel to the total lobe. Lung sections from C57BL/6J mice

treated with bleomycin, then NVP-BHG712 or vehicle were normalized to total tissue and

collagen content was expressed as a percentage. Verhoeff Van Gieson's stain (Abcam) was

performed according to manufacturer's instructions to assess elastin in lung tissue. Lung

sections were assigned elastin expression scores on a scale of 0–3 (absent, mild, moderate, or

severe staining) based on the level of black staining of elastin fibers against pink and yellow

counterstain.

Immunohistochemistry

Lung sections (4 μm) were deparaffinized in xylene followed by rehydration using a

graded series of alcohol washes. Antigen retrieval was performed using proteinase K (20 ug/mL) for 15 min, and endogenous peroxide was blocked for 30 min using 3% H_2O_2. Nonspecific IgG binding was blocked by incubating sections with 2% goat serum in PBS for 30 min. Sections were then incubated with primary antibodies for α-SMA (Sigma-Aldrich; catalog A2547) at 1:1000 overnight at 4°C. Sections were incubated with biotinylated secondary anti-mouse antibody (Vector Laboratories; catalog BA-9200) for 30 min, followed by incubation using the Vectastain Elite ABC kit (Vector Laboratories), as per manufacturer's direction, and counterstained with hematoxylin. Lung sections were assigned α-SMA positivity expression scores on a scale of 0–3 (absent, mild, moderate, or severe staining) based on the level of brown staining of α-SMA against the blue counterstain.

Skin histology

Mouse epidermis and dermis sections were collected using surgical scissors and forceps. Mouse skin was fixed in 10% formalin at room temperature for 48 h, then incubated for 24 h in a minimum of 15 mL JFC solution per 4 mouse skin sections. Skin sections were embedded in paraffin and 4 μm sections were collected. Mouse skin sections were stained with Gomori's Onse Step Trichrome. Skin fibrosis scoring and dermal thickness analyses were performed as previously described[7].

Lung fibroblast isolation

Mouse lungs were perfused with 10 mL PBS at 37°C through the right ventricle of the heart and out the abdominal aorta until completely blanched. Lungs were excised and rinsed in PBS. After blotting away excess PBS, lung tissue was minced by cutting with scissors 50 times. 1 mL collagenase (1 mg/mL in DMEM; Sigma) was then added, and tissue pieces were cut an additional 50 times until they were no larger than 1 mm in diameter. The lung tissue + collagenase slurry was collected and volume adjusted with collagenase to 5 mL. Tissue pieces

were digested by incubating at room temperature for 15 min. Tissue digest was deactivated with 1 mL FBS and the cells were immediately passed through a 70-µm cell filter and rinsed with 10 mL DMEM. Cells were centrifuged at 300 g for 8 min and resuspended in 1 mL complete DMEM (10% FBS, 1% penicillin/streptomycin). Cells were seeded in a 6 cm plate in 4 mL complete DMEM for 24 h, at which point 50% of the media was changed. Cells were lifted with trypsin (0.05%) and passaged when they reached 80% confluence (3−7 days). For the preparation of RNA for NanoString gene analysis, mouse lung fibroblasts from C57BL-6J mice were treated with TGF-β1 (20 ng/mL) or PBS (control), and NVP-BHG712 (250 nM) or DMSO (vehicle control), for 24 h.

Skin fibroblast isolation

Ephb4 control (n=6) and *Ephb4* CKO (n=6) mice were used to isolate dermal fibroblast cells. After sacrifice, mouse skin was sterilized with 70% ethanol and fur was removed with Nair. The skin was excised, minced, and digested in collagenase for 2 h at 37°C. Cell suspension was collected and centrifuged at 300 g for 8 min. Cell pellets were resuspended in DMEM (10% FBS, 1% penicillin/streptomycin) and cultured in 60 mm culture dishes for 24 h. Mouse fibroblasts were passaged to P1 and cell supernatants were collected for RNA and protein collection.

Primary human lung fibroblast culture and treatment

Primary human lung fibroblasts from either IPF patients (n=4) or donors with no history of lung fibrosis (n=5) were purchased from Lonza at P2 (**Supplementary Table 6**). Fibroblasts were seeded, expanded and passaged to P4. The cells were starved overnight and subsequently treated with TGF-β1 or PBS (control) for 48 h. After 24 h of TGF-β1 treatment, NVP-BHG712 (250 nM) or DMSO vehicle was spiked into the fibroblast cultures for an additional 24 h, and RNA was isolated for qRT-PCR. For RNA sequencing, P4 primary IPF lung fibroblasts were treated with NVP-BHG712 or DMSO vehicle for 24 h. IPF lung fibroblasts were alternatively

treated with silencer select siRNA (Thermo) targeted to EphB4 using RNAiMax lipofectamine according to manufacturer's instruction. siRNA treatment was performed for 24 h at the recommended concentration of 25 pmol per well in a 6-well plate.

Western blot

Protein lysates were generated from mouse lung fibroblasts using RIPA buffer with protease and phosphatase inhibitor tablets. Lysates were applied to SDS-acrylamide gels (7.5 or 10%) for electrophoresis. Electroblotting was performed on PVDF membranes and blocked with 1% TWEEN20 10 mM Tris-buffered saline (TTBS) containing 2.5% skim milk, and 2.5% bovine serum albumin (BSA). Probing was performed for EphB4 (R&D, catalog #AF446-SP) at 1:500, phospho-EphB4 (ThermoFisher, catalog #PA5-64792) at 1:500, elastin (Abcam, ab213720) at 1:300, phospho-SMAD 2/3 (Abcam, catalog #ab272332), SMAD 2/3 (Abcam, catalog #ab217553), and GAPDH (Abcam, catalog #ab9485) at 1:1000 for 2 h. Membranes were washed with TTBS 3 times for 5 min each and incubated in HRP-conjugated anti-rabbit (1:5,000; Sigma-Aldrich, catalog #SAB3700843) or anti-mouse (1:10,000; Sigma-Aldrich, catalog #A2179) secondary antibodies for 1 h at room temperature. Membranes were washed multiple times in TTBS before visualizing with chemiluminescence substrate (Clarify Western ECL Substrate, BIO-RAD, or SuperSignal West Pico, Thermo Science) and scanned using a BIO-RAD Chemidoc apparatus. Membrane signal intensity was quantified using ImageJ (National Institutes of Health, USA).

Flow cytometry

Lungs were recovered from *Ephb4* control and *Ephb4* CKO mice after vascular perfusion with PBS, then minced and digested for 30 min at 37°C in 0.5 mg/ml collagenase VIII (Sigma). Single cells were filtered through a 70 µm filter and 1×10^6 cells were labeled with a live/dead marker (eFlour780, Fisher) in PBS according to the manufacturer's instructions. Cells were

washed once, resuspended in 50 µl of FACS buffer and labeled with CD45-PE-Cy7 (Biolegend, 30F11), F4/80-APC (Biolegend, BM8), CD31-PE (Biolegend, 390) and goat anti-EphB4 (R&D, AF446, 12 µl), followed by donkey anti-goat-AF488 secondary (Invitrogen, A11055). At least 100,000 events were acquired using a BD FACS-Canto II flow cytometer. Single cells, live cells, and either macrophages (CD45$^+$/F4/80$^+$) or endothelial cells (CD45$^-$/CD31$^+$) were gated. Geometric mean fluorescence intensity (gMFI) of EphB4 was determined for each gated population using FlowJo_v10.8.4. Cell sorting of similarly labeled cell populations was performed on a BD FACS ARIA III cell sorter, and cells were lysed in RIPA buffer for Western blot analysis. Total EphB4 was detected by incubating the membrane with an anti-EphB4 antibody (R&D, AF446) at 1/250 dilution, followed by a donkey anti-goat HRP secondary (1:10,000).

RNA isolation and qRT-PCR

RNA isolation was performed using the TRIzol-chloroform method. Nanodrop was used to evaluate the quality of the RNA, and cDNA was generated using the QuantiTect™ Reverse Transcription kit (catalog 205311; Qiagen) according to manufacturer's instruction. QRT-PCR was performed using SYBR™ Green PCR Master Mix (catalog 4309155; ThermoFisher) according to the manufacturer's instructions. qRT-PCR primer sequences can be found in **Supplementary Table 12**.

NanoString

Mouse lung fibroblasts treated with TGF-β1 and NVP-BHG712, or human lung fibroblasts from IPF donors treated with NVP-BHG712 were lysed in Trizol reagent and frozen at -80°C. Total RNA was isolated using RNeasy MinElute columns (catalog 74204; Qiagen), according to standard protocols. RNA was quantified by Nanodrop and 100 ng of total RNA was hybridized overnight at 65°C on the nCounter instrument (NanoString Technologies,

Seattle, WA), using custom designed panels for mouse or human transcripts of interest. Primers used for the NanoString gene panels can be found in **Supplementary Table 13a/b**. Results were assessed using the nCounter analysis system on log transformed expression counts with normalization performed using the standard housekeeping controls *Acad9* and *Rplp0*. Paired t-test and generalized least squared regression were performed for unadjusted and adjusted (age and sex) analyses. The resulting p-values were further adjusted for multiple comparisons according to the Benjamini-Hochberg method with q<0.05 being determined as significant.

Total RNA sequencing and analysis

Quality of total RNA extracted from either cultured lung fibroblasts of *Ephb4* CKO mice treated with bleomycin (n=4) or PBS (n=5), *Ephb4* control mice treated with bleomycin (n=4) or PBS (n=4), or human IPF lung fibroblasts treated with NVP-BHG712 or vehicle (n=4 per treatment group) were assessed using an RNA Nano chip on an Agilent Bioanalyzer (Agilent, Santa Clara, CA, USA). RNA integrity numbers for all samples were >9.5. Samples were fluorometrically quantified using the Qubit RNA BR assay (ThermoFisher) on the Denovix DS-11 spectrophotometer (Denovix, Wilmington, DE, USA). For each sample, 200 ng RNA was used to prepare sequencing libraries using the TruSeq Stranded Total RNA with RiboZero following the low sample protocol as per manufacturer's recommendations (Illumina, San Diego, CA, USA), as previously described[43]. Library quality was assessed on a high sensitivity DNA Bioanalyzer chip (Agilent). In total, 17 mouse libraries or 8 human libraries were volumetrically pooled and sequenced on an Illumina NextSeq 550 sequencer for 75 paired end read-cycles at the Centre for Arthritis Diagnostic and Therapeutic Innovation (CADTI, Schroeder Arthritis Institute, Krembil, Toronto, ON, Canada). Sequencing data are deposited in the gene expression omnibus (GEO) data repository under the accession codes GSE205612 and GSE205613 for the mouse and human lung fibroblasts, respectively.

Demultiplexing of samples was performed using bcl2fastq conversion tool

(v2.19.1.403). Quality assessment of each sample revealed high quality of reads. To maintain

minimum read length of 25 bp post trimming of adapters, Cutadapt (v2.5) software was

executed (very few reads being contaminated ~3%) along with trimming of Ns. Splice-aware

alignment of reads using a Hierarchical Graph FM index (HGFM) method was performed using

HISAT2 software (v2.1.0 with parameters –rna-strandness RF –dta) against mouse reference

genome (vGRCm38) and human reference genome (vGRCh38) for mouse and human samples

respectively. To populate the abundance of transcripts based on the reference genome and

transcriptome, StringTie (v2.0.3 with parameters -e -B -G referencetranscripts.gtf) was run to

generate output as table format files. These files were processed using the script 'prepDE.py'

from StringTie software to collect the raw gene expression levels for downstream processing.

For mouse lung fibroblast RNA sequencing analyses, raw gene expression levels

comprised 55,421 genes measured across fibroblasts obtained from 17 mice. Lowly expressed

genes were filtered out based on genes with less than 2 samples with a count per million less

or equal to 10. This filter kept 10,481 genes for further analysis. For human IPF fibroblast RNA

sequencing analyses, 60,675 genes were measured across 8 samples in the original matrix of

counts. Lowly expressed genes were filtered out based on genes with less than 2 samples with

a count per million less or equal to 10. This filter kept 9,619 genes for further analysis.

Changes in gene expression were evaluated using the method proposed by Anders and

Huber[44], which assumes a negative binomial distribution. Function DESeq in R package

DESeq2 was used with default parameters to perform unadjusted and (age) adjusted differential

expression analyses[45]. Pairwise comparisons among groups of interest were calculated and

tested for statistical differences in expression levels. All analyses include batch number as a

categorical covariate in the model to control for potential batch effects. P-values were adjusted

for multiple comparisons via FDR correction proposed by Benjamini and Hochberg[46]. An FDR

of 5% (q<0.05) was statistically significant. All analyses were performed in R version 3.5.0[47].

Computational analysis

Identified DEGs were analyzed using pathDIP[14] (https://ophid.utoronto.ca/pathDIP) v.4 selecting mouse as organism, Reactome as a source and Ortholog Pathways members as annotation, retaining only pathways with q-value<0.01[46]. Up- and down-regulated genes were subjected to pathway analysis and unique pathways were further considered. For network annotation, each gene was annotated with the pathway with the lowest q-value it belonged to, and a network with the gene-pathway interactions was built using NAViGaTOR v.3.0.16[15]. After validation through qRT-PCR of genes involved in pathways linked to ECM, PPIs were retrieved using mouse Integrated Interaction Database (IID; https://ophid.utoronto.ca/iid) v.2021-05[47], and shortest paths between *Ephb4* and the confirmed genes were calculated using igraph_1.2.6 in R 4.0.3[48].

EPHB4 and *ELN* were used to query IID v.2021-05 to retrieve their direct, human, physical protein interactions[47]. We used NAViGaTOR v.3.0.16 to visualize the resulting network, and highlight connecting proteins and those confirmed in RNAseq data (red label)[15]. Node color depicts Gene Ontology Cellular Component. Network image was exported in SVG format and finalized in Adobe Illustrator version 26.3 to produce the final Figure 3A with legend. Pathway enrichment analysis was performed using Enrichr web portal (June 1, 2022) and BioPlanet 2019 pathway source[49].

Single Nuclei RNA Sequencing (snRNA-seq) and bioinformatics analysis

C57B/6J mice were subjected to intratracheal instillation of bleomycin or PBS control, then treated with NPV-BHG712 or vehicle control as described above. Mouse lungs were perfused with PBS, isolated, snap frozen, and stored at -80°C. Nuclei were isolated from frozen

mouse lungs (n=12; n=3/group) by physical dissociation using a glass dounce homogenizer

and were sorted using fluorescence-activated nuclei sorting (FANS) based on DAPI-positive

fluorescence to obtain a single nuclei population. Nuclei concentration was determined using

DAPI staining and hemocytometer, then adjusted to 1,000 nuclei/μL. Single nuclei RNA

sequencing libraries were generated using the 10X Genomics Chromium Next GEM Single

Cell 3' Reagent Kits (v3.1 Dual Index). Sequencing was performed on the Illumina NextSeq

550 system (2x150 bp High Output Kit) at the CADTI. For data analysis of the single nuclei

sequencing experiments, sequencing data were processed using Cell Ranger pipeline (v 7.2.0)

by 10x Genomics (https://www.10xgenomics.com). Reads were aligned to the mouse

transcriptome (mm10), followed by filtering and correction of cell barcodes and Unique

Molecular Identifiers (UMIs). Reads associated with retained barcodes were quantified and

used to build expression matrices. The standard procedures of filtering, normalization,

dimensionality reduction and clustering were performed using an R package Seurat (v5.0.3)[50].

Low quality cells expressing less than 200 genes and low expressed genes present in less than

3 cells were removed. Cells with high mitochondrial DNA, high number of unique feature

counts (potential doublets or multiplets) were excluded. We used DoubletFinder for the

detection of doublets [SV1][51]. Samples were merged for subsequent clustering and

visualization. The batch correction between samples was done with the Harmony R package

(v1.0)[52].

Data were log-normalized and highly variable genes were detected using variance-

stabilizing transformation (vst). The normalized data were centered and scaled. Dimensionality

reduction was performed using principal component analysis (PCA) and the most significant

PCs were chosen for subsequent clustering. Graph-based clustering was implemented by

calculating k-nearest neighbors, followed by modularity optimization to clusters cells. Non-

linear dimensionality reduction and visualization was performed using UMAP (Uniform

Manifold Approximation and Projection). Clusters were annotated on the basis of canonical markers and differential gene expression testing was used to determine a gene set signature for each cluster using the Wilcoxon Rank Sum test. GO enrichment analysis was performed on differentially expressed genes using enrichGO as implemented in cluster Profiler package (version 4.4.4)[53].

To explore potential cell-cell interactions across fibroblast subclusters in bleomycin samples, we employed CellChat v2.0 R package where each fibroblast subcluster was chosen as 'receiver' cell type and the other subclusters as 'sender' cell type[54]. The "netVisual_chord_gene" function in CellChat was used to generate a chord diagram to visualize the complex ligand-receptor interactions between source and target cell types. We identified the upregulated and down-regulated signaling ligand-receptor pairs based on the differential expression analysis.

Analysis of publicly available single-cell RNA sequencing data

We generated dotplots to look at the expression of gene signatures for activated fibroblasts from Peyser et al[23], and the expression of gene signatures for Col13a1 matrix and Col14a1 matrix fibroblasts from Xie et al[24].

The analysis of the published scRNA-seq dataset (GSE128033) was performed using R package Seurat (v4.2.1)[50]. The UMI matrix was filtered to remove low quality cells expressing very few genes and potential doublets or multiplets expressing an aberrantly high number of genes. Samples were merged for subsequent clustering and visualization. Batch correction between samples was done with Harmony R package (v1.0)[52].

Data were log-normalized and highly variable genes were selected using variance-stabilizing transformation (vst) method. Most significant principal components from dimensionality reduction using PCA were chosen for subsequent clustering. Graph-based clustering was implemented by calculating k-nearest neighbors, followed by modularity

optimization to clusters cells. Non-linear dimensionality reduction and visualization was performed using UMAP (Uniform Manifold Approximation and Projection). Clusters were annotated using canonical markers from the literature.

Statistical Analysis

qRT-PCR and Western blot data are presented as interleaved scatter plots with horizontal lines representing the mean. The upper and lower bounds in the scatter plots correspond to the standard deviation of the data set. Statistical analysis identifying the effects of the following utilized 2-Way Anova followed by Tukey's honest significant difference post-hoc test: determining the severity of fibrosis using either the Ashcroft scoring system or through ImageJ computational analysis, quantification of IHC scoring for α-SMA in lung tissue, evaluating the mRNA expression of *Col1a1* and *Acta2* after treatment with PBS vs bleomycin and *Ephb4* control vs *Ephb4* CKO in mouse lung fibroblasts via qRT-PCR, evaluating the protein expression of elastin via Western blot after PBS vs TGF-β1 and Vehicle vs NVP-BHG712/*EPHB4* siRNA, and quantification of elastin stain scoring for the Van Gieson stain in lung tissue. Statistical analysis identifying the effects of the following utilized student's T test: determining EphB4 expression or phosphorylation of EphB4 either through qRT-PCR or Western blot comparing *Ephb4* control to *Ephb4* CKO. In all comparison tests, a value of $p<0.05$ was considered statistically significant. Graphs and analysis were generated in GraphPad Prism version 10. Full access to all data points, means, p-values, and sample size numbers can be found in the **Supporting Data Values** file.

Study Approval

Animal experiments were approved by the Animal Resources Centre under the animal use protocol number #3730, University Health Network.

Data Availability

All datasets for RNA sequencing can be accessed online as part of the GEO super series GSE273468. Datasets for total RNA sequencing of mouse lung fibroblasts can be accessed online under accession number GSE205612. Datasets for single nuclei RNA sequencing of mouse lung fibroblasts can be accessed online under accession number GSE273467. Datasets for RNA sequencing of human lung fibroblasts can be accessed online under accession number GSE205613.

In this study, the following publicly available dataset was accessed: GSE128033.

Acknowledgments and Affiliations: This study is supported by grants to MK by National Science and Engineering Research Council (NSERC), Canada Research Chairs Program, Tony and Shari Fell Platinum Chair in Arthritis Research and University Health Network Foundation. BW is the recipient of the Ontario Graduate Scholarship, The Arthritis Society PhD Salary Award, and the CIHR Frederick Banting and Charles Best Canada Graduate Scholarship. IJ was supported in part by funding from Natural Sciences Research Council (NSERC RGPIN-2024-04314), Canada Foundation for Innovation (CFI #225404, #30865), and Ontario Research Fund (RDI #34876, RE010-020). The research of BH is supported by a foundation grant from the Canadian Institutes of Health Research (#375597) and support from the John Evans Leadership funds (#36050, #38861, and 38430) and innovation funds ('Fibrosis Network, #36349') from the Canada Foundation for Innovation (CFI) and the Ontario Research Fund (ORF). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The authors would like to kindly thank Dr. Evgeny Rossomacha for their hep with the histology protocols and KP and JG for assisting in animal care and lab management for the duration of this study.

Competing interests: Authors declare that they have no competing interests.

Figure 1. A methodological approach to assessing the role of EphB4 in pulmonary fibrosis

Figure 1. **Figure 1. Methodological approaches used to assess the role of EphB4 in pulmonary fibrosis.** A schematic representation of the methods and experimental approach used in this study.

Figure 2

Figure 2. *Ephb4* **conditional knockout mice exhibit a protective phenotype against the development of pulmonary fibrosis.** *Ephb4^{f/f}* mice were crossed with *Col1a2-Cre^{ERT}* mice to generate a line with *Col1a2*-driven deletion of *Ephb4* (*Ephb4* CKO). (A) Genotyping was performed to confirm the presence of homozygous floxed *Ephb4* alleles and *Cre*. (B) Mice treated with 4-hydroxytamoxifen to induce the CKO were sacrificed and primary lung fibroblasts were cultured to first passage. RNA was collected and *Ephb4* was quantified by qRT-PCR (n=6–7). Protein lysates were collected from cell cultures and EphB4 protein was quantified by Western blotting (n=5). (C) A mouse model for inducing fibrosis in *Ephb4* CKO mice was performed using the intratracheal bleomycin challenge or PBS (control) following 4-hydroxytamoxifen-induced deletion. (D) Mouse lungs were collected at indicated end points and stained using the Gomori's One Step collagen stain (blue). Representative histology images from two separate animals/group are shown. Fibrosis was quantified either by modified Ashcroft scoring or with ImageJ total lung collagen content quantification (n=8 mice/group). (E) α-SMA staining was performed in lung tissue sections to assess fibrotic activity in *Ephb4* CKO and control mice with and without bleomycin challenge (n=5). Semi-quantitative scoring (0-3) of the staining intensity was assigned to each mouse lung. (F) Primary lung fibroblasts were isolated from *Ephb4 Col1a2-Cre^{ERT}* mice treated with 4-hydroxytamoxifen or corn oil, then bleomycin. RNA was isolated from fibroblasts and qRT-PCR was performed against *Col1a1*, and *Acta2* (n=4). For qRT-PCR (B, F), ΔCT values were used to represent relative expression of the gene of interest normalized to *Gapdh*. For all graphs, data are expressed as mean ± SD where differences (*, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p<0.0001$) were

1   identified using either Student's t-test (comparison between 2 groups) or 2-way ANOVA with

2   Tukey's post-hoc tests (comparison of 2 variables).

Figure 3

Figure 3. *Ephb4* **CKO fibrotic mouse lung fibroblasts exhibit unique transcriptomic**
profiles. (A) *Ephb4^{f/f} Col1a2-Cre^{ERT}* mice were treated with 4-hydroxytamoxifen (1 mg/day
for 5 days) or corn oil control at 4 weeks of age, then intratracheally instilled with PBS or
bleomycin (1.2U/kg) at 6 weeks of age. Mice were sacrificed 14 days after bleomycin or PBS

challenge and lung fibroblasts were isolated and cultured (n=4 mice per group). Fibroblasts were enriched with a single passage and RNA was extracted for total RNA sequencing. (B) Pairwise comparisons were made between bleomycin and PBS treated *Ephb4* control mice, or between *Ephb4* CKO and control mice under bleomycin condition (n=4 mice/group). Differentially expressed genes (DEGs) were determined using an FDR cutoff of 5% (red). In the bleomycin vs PBS comparison 257 DEGs were identified in fibroblasts, including 155 upregulated and 102 downregulated DEGs. In the *Ephb4* CKO vs control comparison, 317 DEGs were identified in fibroblasts, including 103 upregulated and 214 downregulated DEGs. (C) DEGs from both comparisons were visualized in heat maps. (D) Gene-pathway associations included 8 unique pathways in the upregulated DEGs, and 11 unique pathways associated with downregulated DEGs. 57 downregulated genes were associated with 11 pathways (down arrows) compared to 31 upregulated genes, which were associated with 8 pathways (up arrows). Among the 57 downregulated genes, 11 were associated with ECM organization, 7 were associated with protein trafficking, 4 were associated with ER cargo concentration, and 12 were associated with class I mediated antigen processing and presentation. (E) Lists of both upregulated and downregulated DEGs in the RNA sequencing dataset of *Ephb4* CKO (n=4) vs *Ephb4* control mice (n=4) treated with bleomycin were separately analyzed using pathDIP to identify subsets of genes that enriched for unique pathways in the Reactome database. Gene-pathway associations were visualized in NAViGaTOR v.3.0.16. (F) Unique gene-pathway enrichment from the downregulated gene list visualized by dotplot with significance determined using the Benjamini and Hochberg method with an adjusted q-value cutoff of <0.01. (G) Expression of downregulated DEGs associated with ECM organization, protein transport and ER cargo concentration visualized by dotplot.

Figure 4
A

Figure 4. NVP-BHG712 downregulates TGF-β1-induced expression of genes related to ECM organization, protein trafficking, and ER cargo concentration regulation. (A) Lungs were isolated from WT mice and fibroblasts cultured. Fibroblasts were simultaneously treated with the pro-fibrotic cytokine TGF-β (20 ng/mL; or PBS as control) and the EphB4 inhibitor NVP-BHG712 (250 nM; or DMSO as inhibitor control) for 24 h (n=9). RNA was isolated and NanoString was performed on a 47 gene CodeSet to examine the expression on the downregulated DEGs identified by RNA-seq in bleomycin-challenged *Ephb4* CKO lung fibroblasts. (B) Transcript levels of a subset of the CodeSet of genes that were significantly upregulated with TGF-β1 treatment and downregulated with NVP-BHG712 treatment and were associated with ECM organization, protein trafficking, and ER cargo concentration regulation (remaining genes can be found in **Supplementary Figure 5**). Gene expression was normalized to housekeeping genes *Rplp0* and *Acad9* displayed as log2 transformed gene expression. Wilcoxon signed rank test and false discovery rate (FDR) with a 5% cut-off were performed to determine q-values to identify significant gene expression. Legend colours indicate the gene-associated pathDIP-identified pathways. *, FDR adjusted q-value < 0.05.

Figure 5

Figure 5. Pharmacological inhibition of EphB4 protects against development of PF in mice. (A) Bleomycin sulphate (1.2 U/kg) or PBS (control) was intratracheally administered to WT C57B/6 mice. 5 days later, mice were given either NVP-BHG712 (10 μg/kg; p.o.) or vehicle (control) daily for 5 days (n=6/group). Lungs were harvested 14 days post-bleomycin challenge and histology was performed. (B) To assess fibrosis, collagen fibers in mouse lungs were stained via Gomori's One Step Trichrome (blue). Representative histology images from 2 separate animals/group are shown. (C) Mouse lung section images (n=5 animals/group; 5 lobes/animal) were graded for severity of fibrosis using the Ashcroft scoring system (left graph). Images of lung sections underwent color deconvolution in Fiji V. 2.9.0 to isolate the blue channel as a representative of total lung collagen. Total lung collagen was normalized to total lung tissue and expressed as percent total lung collagen. Graphical representations indicate individual lobes (middle graph) or averages of 5 lobes per animal (right graph). (D) α-SMA

immunohistochemistry performed on mouse lung sections (n=5 animals/group). Semi-quantitative scores were assigned from 0–3 based on degree of staining. For all graphs, data expressed as mean ± SD and analyzed by 2-way ANOVA followed by Tukey's post-hoc tests for multiple comparisons, where *, $p < 0.05$; **, $p < 0.01$; ****, $p < 0.0001$.

Figure 6. SnRNA-seq profiling of mouse lung tissues. (A) Schematic detailing the single nuclei RNA sequencing of mouse lungs treated with bleomycin or PBS as control, and NVP-BHG712 or vehicle (n=12 lung tissues; n=3/group). (B) UMAP projection of 15 distinct cell populations identified in mouse lung tissue. (C) Dotplot showing expression of canonical markers used to annotate the clusters. (D) UMAP of fibroblast subclusters from 4 treatment groups (bleomycin/PBS and inhibitor/vehicle). (E) UMAP visualization of bleomycin-treated fibroblasts with inhibitor vehicle or NVP-BHG712. (F) Dotplot showing the expression of top genes associated with each fibroblast subcluster comparing bleomycin challenged NVP-BHG712 treatment group to vehicle treatment group. (G) Expression of elastin (*Eln*) in fibroblast subcluster 1 and *Col1a1* in fibroblast subcluster 3 in bleomycin challenged NVP-BHG712 treatment group compared to vehicle treatment group. (H) GO analysis showing biological processes associated with upregulated and downregulated DEGs between bleomycin vehicle and bleomycin NVP-BHG712 groups. (I) Cell-cell communication in NVP-BHG712 and vehicle bleomycin-treated groups showing ligand-receptor interactions between sender and receiver fibroblast subclusters (0–3) in collagen signaling pathway.

Figure 7

Figure 7. Elastin is a key signaling partner of EphB4 in human IPF. (A) Schematic
detailing total RNA sequencing pipeline of human lung fibroblasts (HLF) from control donors
or IPF patients subjected to NVP-BHG712 (250 nM) or vehicle (n=4/group). (B) Differentially
expressed genes (DEGs) comparing IPF to HLF fibroblasts using an FDR cutoff of 5% (q-
value<0.05) (red). Heat map depicting 3,108 DEGs (1,569 downregulated and 1,539
upregulated) identified in IPF vs HLF fibroblasts. Dotplot showing expression of top ECM
genes in IPF vs HLF fibroblasts. (C) DEGs comparing IPF fibroblasts treated with NVP-
BHG712 or vehicle (red). Heat map depicting 85 DEGs (61 downregulated and 24 upregulated)
identified in response to EphB4 inhibition of human IPF fibroblasts. Dotplot showing
expression of ELN comparing NVP-BHG712 vs vehicle treatment in IPF fibroblasts. (D)

EphB4 was queried with elastin, as identified across human and mouse RNA sequencing, using the Integrated Interaction Database to find direct human PPIs and visualized using NAViGaTOR.

Figure 8

2
Figure 8. EphB4 signaling in PF. (A/B) Human IPF lung fibroblasts were treated with TGF-
β1 or PBS for 24 h. Cells (total 16 samples; n=4 patients/group) were then treated with either
NVP-BHG712 (250 nM) (A) or *Ephb4* siRNA (B), or their respective controls, for an
additional 24 h. Protein lysates were collected and Western blotting was performed to detect
elastin protein levels. ImageJ densitometry was performed and normalized to GAPDH. *,
p<0.05; **, p<0.01, as determined by paired two-way ANOVA followed by Tukey's post-hoc
test adjusted for multiple comparisons. (C) Western blotting of siRNA treated IPF cells to
detect p-SMAD 2/3 and SMAD 2/3 (n=4/group). (D/E) Verhoeff Van Gieson staining was
performed on lung sections from (D) mice with *Ephb4* conditional knockout or respective
control mice to visualize elastin (n=5/group) following bleomycin or PBS instillation, or (E)
mice with bleomycin or PBS instillation treated with EphB4 inhibitor NVP-BHG712 (10
mg/kg) or vehicle (n=5/group). Elastin staining was semi-quantitatively scored (0-3). ****, p-
value<0.0001 as determined by 2-way ANOVA followed by Tukey's post-hoc tests adjusted
for multiple comparisons. (F) Schematic depicting the role of EphB4 signaling in PF.
17

REFERENCES

1.   Wu B, Tang L, Kapoor M. Fibroblasts and their responses to chronic injury in pulmonary fibrosis. *Semin Arthritis Rheum*. Feb 2021;51(1):310-317. doi:10.1016/j.semarthrit.2020.12.003

2.   Herzog EL, Mathur A, Tager AM, Feghali-Bostwick C, Schneider F, Varga J. Review: interstitial lung disease associated with systemic sclerosis and idiopathic pulmonary fibrosis: how similar and distinct? *Arthritis Rheumatol*. Aug 2014;66(8):1967-78. doi:10.1002/art.38702

3.   Flaherty KR, Wells AU, Cottin V, et al. Nintedanib in Progressive Fibrosing Interstitial Lung Diseases. *N Engl J Med*. Oct 31 2019;381(18):1718-1727. doi:10.1056/NEJMoa1908681

4.   Richeldi L, du Bois RM, Raghu G, et al. Efficacy and safety of nintedanib in idiopathic pulmonary fibrosis. *N Engl J Med*. May 29 2014;370(22):2071-82. doi:10.1056/NEJMoa1402584

5.   Distler O, Highland KB, Gahlemann M, et al. Nintedanib for Systemic Sclerosis-Associated Interstitial Lung Disease. *N Engl J Med*. Jun 27 2019;380(26):2518-2528. doi:10.1056/NEJMoa1903076

6.   Kullander K, Klein R. Mechanisms and functions of Eph and ephrin signalling. *Nat Rev Mol Cell Biol*. Jul 2002;3(7):475-86. doi:10.1038/nrm856

7.   Lagares D, Ghassemi-Kakroodi P, Tremblay C, et al. ADAM10-mediated ephrin-B2 shedding promotes myofibroblast activation and organ fibrosis. *Nat Med*. Dec 2017;23(12):1405-1415. doi:10.1038/nm.4419

8.   Mimche PN, Lee CM, Mimche SM, et al. EphB2 receptor tyrosine kinase promotes hepatic fibrogenesis in mice via activation of hepatic stellate cells. *Sci Rep*. Feb 7 2018;8(1):2532. doi:10.1038/s41598-018-20926-9

9.   Su SA, Yang D, Wu Y, et al. EphrinB2 Regulates Cardiac Fibrosis Through Modulating the Interaction of Stat3 and TGF-β/Smad3 Signaling. *Circ Res*. Sep 1 2017;121(6):617-627. doi:10.1161/circresaha.117.311045

10.   Jin W, Luo H, Wu J. Effect of reduced EPHB4 expression in thymic epithelial cells on thymocyte development and peripheral T cell function. *Mol Immunol*. Mar 2014;58(1):1-9. doi:10.1016/j.molimm.2013.10.008

11.   Zheng B, Zhang Z, Black CM, de Crombrugghe B, Denton CP. Ligand-dependent genetic recombination in fibroblasts : a potentially powerful technique for investigating gene function in fibrosis. *Am J Pathol*. May 2002;160(5):1609-17. doi:10.1016/s0002-9440(10)61108-x

12.   Hübner RH, Gitter W, El Mokhtari NE, et al. Standardized quantification of pulmonary fibrosis in histological samples. *Biotechniques*. Apr 2008;44(4):507-11, 514-7. doi:10.2144/000112729

13.   Meng XM, Nikolic-Paterson DJ, Lan HY. TGF-β: the master regulator of fibrosis. *Nat Rev Nephrol*. Jun 2016;12(6):325-38. doi:10.1038/nrneph.2016.48

14.   Rahmati S, Abovsky M, Pastrello C, et al. pathDIP 4: an extended pathway annotations and enrichment analysis resource for human, model organisms and domesticated species. *Nucleic Acids Res*. Jan 8 2020;48(D1):D479-d488. doi:10.1093/nar/gkz989

15.   Brown KR, Otasek D, Ali M, et al. NAViGaTOR: Network Analysis, Visualization and Graphing Toronto. *Bioinformatics*. Dec 15 2009;25(24):3327-9. doi:10.1093/bioinformatics/btp595

16.   Wan XL, Zhou ZL, Wang P, et al. Small molecule proteomics quantifies differences between normal and fibrotic pulmonary extracellular matrices. *Chin Med J (Engl)*. May 20 2020;133(10):1192-1202. doi:10.1097/cm9.0000000000000754

17.   Péterfi Z, Donkó A, Orient A, et al. Peroxidasin is secreted and incorporated into the extracellular matrix of myofibroblasts and fibrotic kidney. *Am J Pathol*. Aug 2009;175(2):725-35. doi:10.2353/ajpath.2009.080693

18.   Chen L, Li S, Li W. LOX/LOXL in pulmonary fibrosis: potential therapeutic targets. *J Drug Target*. Aug 2019;27(7):790-796. doi:10.1080/1061186x.2018.1550649

19.   Hoff CR, Perkins DR, Davidson JM. Elastin gene expression is upregulated during pulmonary fibrosis. *Connect Tissue Res*. 1999;40(2):145-53. doi:10.3109/03008209909029110

20.   Bayoumi A, Jalil I, Metwally M, et al. Genetic variation in the TLL1 gene is not associated with fibrosis in patients with metabolic associated fatty liver disease. *PLoS One*. 2020;15(12):e0243590. doi:10.1371/journal.pone.0243590

21.   Fiore VF, Wong SS, Tran C, et al. αvβ3 Integrin drives fibroblast contraction and strain stiffening of soft provisional matrix during progressive fibrosis. *JCI Insight*. Oct 18 2018;3(20)doi:10.1172/jci.insight.97597

22.   Martiny-Baron G, Holzer P, Billy E, et al. The small molecule specific EphB4 kinase inhibitor NVP-BHG712 inhibits VEGF driven angiogenesis. *Angiogenesis*. Sep 2010;13(3):259-67. doi:10.1007/s10456-010-9183-z

23.   Peyser R, MacDonnell S, Gao Y, et al. Defining the Activated Fibroblast Population in Lung Fibrosis Using Single-Cell Sequencing. *Am J Respir Cell Mol Biol*. Jul 2019;61(1):74-85. doi:10.1165/rcmb.2018-0313OC

24.   Xie T, Wang Y, Deng N, et al. Single-Cell Deconvolution of Fibroblast Heterogeneity in Mouse Pulmonary Fibrosis. *Cell Rep*. Mar 27 2018;22(13):3625-3640. doi:10.1016/j.celrep.2018.03.010

25.   Morse C, Tabib T, Sembrat J, et al. Proliferating SPP1/MERTK-expressing macrophages in idiopathic pulmonary fibrosis. *Eur Respir J*. Aug 2019;54(2)doi:10.1183/13993003.02441-2018

26.   Wu B, Rockel JS, Lagares D, Kapoor M. Ephrins and Eph Receptor Signaling in Tissue Repair and Fibrosis. *Curr Rheumatol Rep*. Apr 12 2019;21(6):23. doi:10.1007/s11926-019-0825-x

27.   Frye M, Stritt S, Ortsäter H, et al. EphrinB2-EphB4 signalling provides Rho-mediated homeostatic control of lymphatic endothelial cell junction integrity. *Elife*. Sep 8 2020;9doi:10.7554/eLife.57732

28.   Ferguson BD, Liu R, Rolle CE, et al. The EphB4 receptor tyrosine kinase promotes lung cancer growth: a potential novel therapeutic target. *PLoS One*. 2013;8(7):e67668. doi:10.1371/journal.pone.0067668

29.   Ferguson BD, Carol Tan YH, Kanteti RS, et al. Novel EPHB4 Receptor Tyrosine Kinase Mutations and Kinomic Pathway Analysis in Lung Cancer. *Sci Rep*. Jun 15 2015;5:10641. doi:10.1038/srep10641

30.   Yang NY, Pasquale EB, Owen LB, Ethell IM. The EphB4 receptor-tyrosine kinase promotes the migration of melanoma cells through Rho-mediated actin cytoskeleton reorganization. *J Biol Chem*. Oct 27 2006;281(43):32574-86. doi:10.1074/jbc.M604338200

31.   Xiao Z, Carrasco R, Kinneer K, et al. EphB4 promotes or suppresses Ras/MEK/ERK pathway in a context-dependent manner: Implications for EphB4 as a cancer target. *Cancer Biol Ther*. Jun 2012;13(8):630-7. doi:10.4161/cbt.20080

32.   Tsukui T, Sun KH, Wetter JB, et al. Collagen-producing lung cell atlas identifies multiple subsets with distinct localization and relevance to fibrosis. *Nat Commun*. Apr 21 2020;11(1):1920. doi:10.1038/s41467-020-15647-5

33.   Zepp JA, Zacharias WJ, Frank DB, et al. Distinct Mesenchymal Lineages and Niches Promote Epithelial Self-Renewal and Myofibrogenesis in the Lung. *Cell*. Sep 7 2017;170(6):1134-1148.e10. doi:10.1016/j.cell.2017.07.034

34.   Valenzi E, Bulik M, Tabib T, et al. Single-cell analysis reveals fibroblast heterogeneity and myofibroblasts in systemic sclerosis-associated interstitial lung disease. *Ann Rheum Dis*. Oct 2019;78(10):1379-1387. doi:10.1136/annrheumdis-2018-214865

35.   Kuang PP, Zhang XH, Rich CB, Foster JA, Subramanian M, Goldstein RH. Activation of elastin transcription by transforming growth factor-beta in human lung fibroblasts. *Am J Physiol Lung Cell Mol Physiol*. Apr 2007;292(4):L944-52. doi:10.1152/ajplung.00184.2006

36.   Chen D, Hughes ED, Saunders TL, et al. Angiogenesis depends upon EPHB4-mediated export of collagen IV from vascular endothelial cells. *JCI Insight*. Feb 22 2022;7(4)doi:10.1172/jci.insight.156928

37.   Wang HU, Chen ZF, Anderson DJ. Molecular distinction and angiogenic interaction between embryonic arteries and veins revealed by ephrin-B2 and its receptor Eph-B4. *Cell*. May 29 1998;93(5):741-53. doi:10.1016/s0092-8674(00)81436-1

38.   Wang Y, Nakayama M, Pitulescu ME, et al. Ephrin-B2 controls VEGF-induced angiogenesis and lymphangiogenesis. *Nature*. May 27 2010;465(7297):483-6. doi:10.1038/nature09002

39.   Wang Y, Thorin E, Luo H, et al. EPHB4 Protein Expression in Vascular Smooth Muscle Cells Regulates Their Contractility, and EPHB4 Deletion Leads to Hypotension in Mice. *J Biol Chem*. May 29 2015;290(22):14235-44. doi:10.1074/jbc.M114.621615

40.   Brown MA, Lowe DG. Recognising panoramas. *Proceedings Ninth IEEE International Conference on Computer Vision*. 2003:1218-1225 vol.2.

41.   Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for biological-image analysis. *Nature Methods*. 2012/07/01 2012;9(7):676-682. doi:10.1038/nmeth.2019

42.   Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for biological-image analysis. *Nat Methods*. Jun 28 2012;9(7):676-82. doi:10.1038/nmeth.2019

43.   Endisha H, Datta P, Sharma A, et al. MicroRNA-34a-5p Promotes Joint Destruction During Osteoarthritis. *Arthritis Rheumatol*. Mar 2021;73(3):426-439. doi:10.1002/art.41552

44.   Anders S, Huber W. Differential expression analysis for sequence count data. *Genome Biol*. 2010;11(10):R106. doi:10.1186/gb-2010-11-10-r106

45.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*. 2014;15(12):550. doi:10.1186/s13059-014-0550-8

46.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*. 2018;57(1):289-300. doi:10.1111/j.2517-6161.1995.tb02031.x

47.   *R: A Language and Environment for Statistical Computing*. 2021. https://www.R-project.org/

48.   Kotlyar M, Pastrello C, Ahmed Z, Chee J, Varyova Z, Jurisica I. IID 2021: towards context-specific protein interaction analyses by increased coverage, enhanced annotation and enrichment analysis. *Nucleic Acids Res*. Jan 7 2022;50(D1):D640-d647. doi:10.1093/nar/gkab1034

49.   Xie Z, Bailey A, Kuleshov MV, et al. Gene Set Knowledge Discovery with Enrichr. *Curr Protoc*. Mar 2021;1(3):e90. doi:10.1002/cpz1.90

50.   Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal single-cell data. *Cell*. Jun 24 2021;184(13):3573-3587.e29. doi:10.1016/j.cell.2021.04.048

51.   McGinnis CS, Murrow LM, Gartner ZJ. DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst*. Apr 24 2019;8(4):329-337.e4. doi:10.1016/j.cels.2019.03.003

52.   Korsunsky I, Millard N, Fan J, et al. Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat Methods*. Dec 2019;16(12):1289-1296. doi:10.1038/s41592-019-0619-0

53.   Wu T, Hu E, Xu S, et al. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innovation (Camb)*. Aug 28 2021;2(3):100141. doi:10.1016/j.xinn.2021.100141

54.   Jin S, Guerrero-Juarez CF, Zhang L, et al. Inference and analysis of cell-cell communication using CellChat. *Nat Commun*. Feb 17 2021;12(1):1088. doi:10.1038/s41467-021-21246-9