Early Life Stress induces brain-wide electrical network predisposition to migraine

Micah Johnson[1,2], Maureen Eberle[1,2], Ian Hultman[3], Xinyu Zhang[3], Yassine Filali[1,2,] Benjamin

Hing[1], Molly Matkovich[1], Alli Jimenez[1], Sara Mitchell [1,2], Anjayooluwa Adegboyo[5], Radha

Velamuri[1], Julia Miller[1], Kung-Sik Chan[3], Sanvesh Srivastava[3], Rainbo Hultman[1,4*]

1 Department of Molecular Physiology and Biophysics, University of Iowa, Iowa City, IA, USA

2 Interdisciplinary Graduate Program in Neuroscience, University of Iowa, Iowa City, IA, USA

3 Department of Statistics and Actuarial Science, University of Iowa, Iowa City, IA, USA

4 Department of Psychiatry, University of Iowa, Iowa City, IA, USA

5 University of Maryland Baltimore County, Baltimore, MD, USA

[*] Correspond with Dr Hultman, lead contact, (Dept. of Molecular Physiology & Biophysics,

Iowa Neuroscience Institute, 51 Newton Rd., 6-510 Bowen Science Building, Iowa City, IA,

52242, USA; 319-335-2855 (o), 919-451-6906 (c); rainbo-hultman@uiowa.edu

ABSTRACT

Background: Migraine is a disorder of severe, recurrent headaches and debilitating sensory, cognitive and affective symptoms, often triggered by stress. Early life stress in childhood has been shown to increase the likelihood of migraine in adulthood in humans. Calcitonin-gene relate peptide (CGRP) has been shown to reliably and acutely induce migraine or migraine-like behavior in both humans and rodent models. Here we investigate the impact of early life stress and CGRP on migraine-related neural circuitry, as well as the impact of early life stress on CGRP-mediated migraine-like activity in order to better understand the mechanisms by which early life stress predisposes neural circuitry to migraine brain activity.

Methods: We implemented an early life stress paradigm in the outbred strain of mice, CD1. We evaluated the impact of peripheral CGRP on migraine-like behavior and employed multi-site in vivo neurophysiology in freely behaving mice. A changepoint analysis was used to dissect differences in individual CGRP-induced responses.

Results: We found that early life stress exacerbated migraine-related behavioral and network physiology. CGRP alone caused disruptions in neural oscillatory activity across a network of brain regions including the anterior cingulate cortex (ACC), amygdala (AMY), thalamus (Po, VPM, and MDthal), and parabrachial nucleus (PBN). We found that power across the network was lowered within 10 minutes of peripheral CGRP exposure, which was sustained for ~40-50 min. Coherence was mostly disrupted in amygdalar brain region pairings, and took on a shorter timecourse, with partial rescue of these responses by migraine abortive, sumatriptan. We found that early life stress exacerbated most of these responses, especially AMY-thalamic coherence pairings, although early life stress in the absence of CGRP demonstrated no impact on the network overall. We further identified individual mice with brain-network activity hypersusceptible to migraine.

Conclusions: Our findings demonstrate that early life stress confers vulnerability to migraine, simultaneously impacting behavior and brain network activity responses to peripheral CGRP.

KEYWORDS

migraine, CGRP (calcitonin-gene related peptide), brain networks, early life stress, local field potentials (LFPs)

BACKGROUND

Migraine is an intricate and incapacitating neurological disorder and the second leading cause of
global disability [1]. Migraine is characterized by recurrent headache episodes spanning 4 to 72
hours with various sensory anomalies including hypersensitivities such as photophobia,
phonophobia, osmophobia, and allodynia [2]. It is important to recognize that migraine is far more
involved than just headache, and frequently occurs over characteristic stages including
prodrome, aura (impacts ~20% of individuals with migraine), headache, postdrome, and
interictal phases [3]. In addition to sensory hypersensitivities, many of these phases of migraine
can include changes in perception, cognition, autonomic nervous system regulation, and mood,
pointing toward multiple sites of regulation and impact, especially across the brain. While the
underlying origin of migraine has been historically debated in terms of central and peripheral
mechanisms, there is growing consensus of both peripheral and central contributors to migraine
[4]. Regardless of origin, it is clear that migraine engages multiple functional domains (cognition,
perception, emotion/affect etc.) that are known to have causal origins in the brain.

Beyond the complex symptomology of migraine, environmental factors, especially those that
have been experienced early in life, can shape susceptibility to migraine in adulthood. Early life
stress (ELS), which is defined by adverse events experienced during childhood, is associated
with increased risk for chronic pain conditions, including migraine [5]. Several epidemiological
studies have indicated that experiencing ELS leads to a greater risk of migraine and headache
severity during adulthood [6-8]. ELS is also one of many possible mechanisms linking the
comorbidity of migraine and mood disorders, such as major depressive disorder [9]. Finally, ELS
has been associated with altered connectivity in brain regions associated with emotional
regulation and stress, which are also implicated in migraine (especially amygdala)[10,11]. However,
the mechanisms by which ELS predisposes one to migraine remains unknown.

Clues to brain mechanisms engaged in migraine stem largely from a combination of
symptomology and imaging studies. Imaging studies have revealed altered functional
connectivity across brain networks in migraine patients as well as structural differences in pain-
related regions of migraine patients (Figure 8A). These changes have been observed not only
during migraine attacks but also in the interictal period [3,12-14]. A number of studies have indicated
that a large part of what makes migraine so severe and debilitating comes from the aversive
aspect of pain, which is greater for pain in the head compared with pain in the periphery [15-17].

Head pain is ranked as more emotionally salient and severe than other types of peripheral pains.
For example, a study in humans demonstrated that repeated application of noxious heat to the
face induces sensitization whereas the same application of noxious heat to the hand induces
habituation [15].

Thus, we focused our initial efforts toward understanding endogenous neural activity of migraine
across several brain regions with an emphasis on regions involved in the emotional salience of
head pain and mood aspect of migraine and involvement in ELS: anterior cingulate cortex
(ACC), amygdala (AMY), posterior thalamus (Po), ventroposteromedial thalamus (VPM),
medial dorsal thalamus (MDthal), and parabrachial nucleus (PBN), all of which have been
implicated in the pathophysiology of migraine [3,18]. The ACC is a key region in pain processing
and more specifically, emotional processing of pain [19]. Reduction in ACC grey matter is
correlated with frequency of migraine attacks in migraine patients [20]. AMY is implicated in
affective pain processing and the perception of such pain [21,22]. Resting functional connectivity
patterns in a network involving the amygdala have been shown to discern migraine and healthy
control patients [23]. Further, increased connectivity patterns between the nucleus accumbens and
AMY were found during the preictal phase compared to the ictal phase in specific phases of
migraine[24]. The thalamus is broadly implicated in the integration of sensory information and has
been associated with migraine-related allodynia, photophobia , and hypersensitivity to negative
emotional stimuli [25-28]. Additionally, altered thalamo-cortical functional connectivity patterns
have been reported in migraine [29]. Optogenetic stimulation of the posterior thalamus was found
to induce photophobia-like behavior in the absence of anxiety [30]. Finally, the brainstem as a
whole, and the PBN in particular, appear to play an important role in migraine[31,32]. PBN is a
critical sensory relay center and is important for homeostatic function, arousal, cravings, thirst,
nociception and regulation of affective responses [33,34]. Altered connectivity patterns across
subcortical-cortical networks that include the PBN have been found in migraine patients during
the preictal phase [12,18,35,36]. Importantly, the PBN contains CGRP projections to the central
amygdala, which has been demonstrated to play an important role in the emotional aspect of pain
[21,33,37] as well as projections to the thalamus, which convey nociceptive information to higher
order cortical regions [38] . Overall, human imaging studies point to brain-wide connectivity
changes in migraine, however, ultimately endogenous multi-region brain network studies in

disease models are required to understand the ways in which brain regions communicate on a
timescale of neuronal activity.

The ways in which endogenous brain-network activity is engaged and coordinated in mouse
models of migraine remains largely unknown. Preclinical models of migraine allow for probing
of neuronal circuitry across time in a way that can help better understand the actual mechanisms
underlying the interconnected circuitry that leads to migraine and to directly test the impact of
prior exposure of ELS on the development of migraine-like brain activity and behavior.
Developing an understanding of systems-level brain activity in preclinical models of migraine
holds great promise for a new era of migraine research and therapeutic development [39].
One of the best characterized and most robust animal models of migraine makes use of calcitonin
gene-related peptide (CGRP) to induce a migraine-like phenotype in mice. This model is
especially important given CGRP therapeutics have played such an important role in migraine
treatment and prevention, being the first pharmacologics designed specifically for migraine[40].
The CGRP model demonstrates validity to the human disorder in three important ways: face
validity (the model displays a similar disease phenotype as in humans) [41] construct validity (the
model shares the same underlying cause of the disease in humans) [42,43], and predictive validity
(treatments that are effective in humans are effective in the model) [42,44]. CGRP has emerged as a
neuropeptide of prominent importance in migraine pathophysiology. Key evidence for the
involvement of CGRP in migraine is that CGRP levels are elevated during migraine attacks [45-47].
Blocking CGRP signaling can also alleviate migraine symptoms in both humans and rodent
models [48]. Additionally, intravenous infusion of CGRP (either central or peripheral) leads to a
delayed migraine-like headache in humans [43] and preclinical migraine-like phenotypes (e.g., light
aversion, spontaneous pain, and allodynia) in rodents [41]. There has been mounting evidence that
CGRP acts on receptors peripherally, with indirect impacts that reach the brain [4,39]. CGRP has
also been shown in preclinical models to evoke a migraine-like response when injected directly
into the brain [30]. However, the nature of the impact of peripheral CGRP on the brain remains one
of the fundamental questions in solving the mystery of migraine.

In this manuscript, we aim to investigate the impact of ELS on CGRP-induced migraine-like
activity in mice. To better understand brain network dynamics underlying migraine, we employ
multi-site *in vivo* neurophysiological recordings in freely behaving animals to monitor changes in

brain activity and connectivity in response to peripheral CGRP administration over time,
providing both high spatial and temporal resolution. We then further probe brain network activity
with anti-migraine medication, sumatriptan, and ELS, in order to understand how ELS may
predispose brain networks to migraine. Finally, we applied a changepoint analysis in order to
identify network-wide changes associated with migraine-like phenotypes.

153
METHODS

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Animal Care and Use

All experiments were performed with approved protocols by the Institutional Animal Care and
Use Committee (IACUC) of the University of Iowa. CD1 mice purchased from Charles River
Laboratory were utilized for all experiments. For neurophysiological experiments (excluding
maternal separation stress), male and female mice aged 9 weeks of age were group housed 2-3
animals per cage for 1 week prior to surgery. Mice were maintained on a 12 h light/dark cycle
with ad libitum access to food and water. All testing was performed between 9:00 AM and 9:00
PM.

Maternal Separation Stress with Early Weaning and Limited Nesting

CD1 mice were obtained from Charles River Laboratory and bred in-house to generate breeding
pairs and their offspring. All experimental animals were a minimum of one generation removed
from shipping stress. Behavioral testing was conducted across several cohorts based on
parturition date. The maternal separation stress paradigm was adapted from previously described
methods [49,50]. All mice were housed on beta chip bedding with only 1/3 of the supplemental
nesting material provided to controls (Enviro-Dri nesting material). Litters were culled to 8 pups
on P1 with balanced sex distribution. Starting at P2, pups were separated from dams for 4 hours
daily until P5 then 8 hours daily until P16. Supplemental heat was provided during separation
periods. Pups were weaned early at P17 to eliminate the likelihood of overcompensation of

maternal care behavior upon dams reunion with pups [51]. Prior to experiments, animals were

weighed weekly and placed into balanced treatment groups based on weights and cohort.

Neurophysiological Experiments

Drug Administration

All injections were performed intraperitoneally (IP) with a 25G needle. We administered

calcitonin gene-related peptide (CGRP) (Sigma-Aldrich, Saint Louis, MO) at 0.1 mg/kg and

sumatriptan at 0.6 mg/kg (AuroMedics Pharma LLC, E. Windsor, NJ). We utilized phosphate-

buffered saline (PBS) as a diluent and vehicle for all experiments. We co-injected sumatriptan

and CGRP by doubling the total volumes of each and diluting them such that our co-injections

were the same solution volume as the single CGRP and PBS injections. All solutions were

prepared 1 hour prior to the start of experiments and placed on ice for the remainder of the

experiment.

Electrode Implantation

Animals aged 10-12 weeks were anesthetized with isoflurane, placed in a stereotaxic device, and

craniotomized for electrode implantation. Metal ground screws were secured above the olfactory

bulb and cerebellum. Tungsten wires (California Fine Wires) were implanted in the anterior

cingulate cortex (ACC), ventral posteromedial thalamus (VPM), medial dorsal thalamus

(MDthal), posterior thalamus (Po), central amygdala (AMY), and left and right parabrachial

nucleus (LPBN/RPBN). Coordinates were adapted for CD1 animals and verified histologically.

Male surgical coordinates for the center of bundles placed (not actual electrode tips) are as

follows: ACC, 0.8mm A/P, 0.25mm M/L, 1.8mm D/V; Amy (BLA/CeA) -1.58mm A/P, 2.75mm

M/L, 3.9mm D/V; thal (MDthal/Po/VPM) -1.82mm A/P, 1.5mm M/L, 3.1mm D/V; PBN -

5.5mm A/P, +/-1.5mm M/L, 2.3mm D/V. Female are: 0.78mm A/P, 0.24mm M/L, 1.8mm D/V;

Amy (BLA/CeA) -1.54mm A/P, 2.68mm M/L, 3.8mm D/V, thal (MDthal/Po/VPM) -1.77mm

A/P, 1.46mm M/L, 3.02mm D/V; PBN -5.36mm A/P, +/-1.46mm M/L, 2.24mm D/V.

Light-Aversion Test

To assess light-aversive behavior, we performed a light/dark assay. In this assay, animals were

placed in a transparent testing chamber (MedAssociates, East Fairfield, VT) consisting of a dark

or light zone divided by a dark insert. We utilized infrared beam tracking and Activity Monitor
software to analyze activity across a 30-minute time period under 20,000 lux light intensity.
While a range of time windows for the light/dark assay have been used across the literature, in
our hands, 15 minutes has always provided the best metric of CGRP response, even when
compared against a 30-minute test (Figure S2), which is consistent with other studies in the
literature [42,52]. Because some studies also report CGRP-induced behavioral changes across 30
min, we also collected additional data beyond the 15-minute assessment window in order to be
thorough. On the first day, animals underwent a baseline assessment in which they are assessed
for a 15-minute period. Mice were injected with vehicle the following day and assessed for
another 15-minute period. Lastly, animals were injected with CGRP and activity was assessed
for 15 minutes in the testing chamber. For ELS experiments, testing occurred at P81-85.

Neurophysiological recordings

Mice were connected to Mu headstages (Blackrock Microsystems, UT, USA) under isoflurane
anesthesia prior to recordings. During recordings, mice were placed in a homecage environment
consisting of standard cages with corn cob bedding and recorded for 30 minutes under baseline
conditions. Neurophysiological data was captured using the Cereplex Direct acquisition system
and sampled at 30kHz. Local field potentials (LFPs) were stored at 1000 Hz and a bandpass filter
was applied at 0.5-250 Hz.

Timeline of neurophysiological recordings during peripheral administration of CGRP

Two weeks following recovery from surgery, animals underwent three days of homecage
recordings. On Day 1, animals were recorded for 30 minutes then received an IP PBS vehicle
injection and recorded for an additional 2 hours post-injection. Day 2, we administered IP CGRP
injections. To ensure any observed changes in brain activity on Day 2 could be attributed to
changes induced by CGRP and not the injection itself, we repeated an IP PBS vehicle injection
recording on Day 3.

Histology/Electrode Placement

All animals were euthanized upon completion of experiments and perfused with 4%
paraformaldehyde (PFA). Brains were stored in 30% sucrose overnight and submerged in an
OCT compound then frozen on dry ice and stored at -80°C. Brains were sliced in 35µm sections

then washed in PBS prior to being stained with red Nissl (ThermoFisher, Waltham, MA). Slices
were imaged at 5x magnification using a Leica fluorescence microscope. Each individual slice
was aligned to the Paxinos and Franklin brain atlas for verification of electrode placement.
Overall, electrode placements had a ~97% success rate. Two animals were removed from
analysis due to failure to hit targets.

Early Life Stress Behavioral Experiments

Elevated Plus Maze

To assess anxiety-like behavior in mice, we utilized an elevated plus maze (EPM) apparatus
consisting of two opposing open and closed arms. A video camera was placed directly above the
center of the maze to capture activity during each trial. Each animal was placed in the center of
the EPM facing the same closed arm at the start of each trial and tested for a total of 5 minutes.
All mice were tested at P66-69.

Plantar von Frey Test

To assess mechanical allodynia in mice, we utilized the plantar von Frey test. Over a total of 5
days, mice were placed in acrylic chambers over a grid support (Bioseb, France). Mice
habituated to the chambers for a total of 2 hours the first two days. For each of the following
days, mice habituated to the chambers for 1 hour prior to the start of experiments. Mice were
considered well habituated when standing on all four paws, calm, and non-responsive to the
habituation filament . Starting on Day 3 (Baseline day), a von Frey filament was applied to the
plantar surface of the right hindpaw. We followed the "up-down" method to obtain a paw
withdrawal threshold for each animal [53,54]. On Day 4, all animals received an IP PBS (vehicle)
injection and were allowed to rest for 30 minutes prior to testing. On Day 5, animals received
either CGRP or vehicle injections with the administering experimenter blinded to conditions, and
were allowed to rest for 30 minutes prior to testing. Animals were tested at P72-79.

Open Field Test

To determine whether the light aversive behavior was independent of anxiety-like behavior in
mice, we performed an open field test. Animals were tested at the end of the experiments at P83-
86 the day after light aversion experiments. Mice were placed in a large open field box for 20

minutes at a time following administration of vehicle or CGRP, with the administering
experimenter blinded to conditions. We utilized ANY-maze software to monitor the total amount
of time animals spent in the center zone of the open field box.

Light aversion test for ELS was carried out as described above with the administering
experimenter blinded to conditions.

QUANTIFICATION AND STATISTICAL ANALYSIS

Implanted vs. Unimplanted Behavioral Statistics

We fit a three factor mixed effects model using the *lme4 R* library [55] with fixed effects terms for
the model intercept, CGRP, day, implant and the interaction between CGRP and implant. We
employed a random intercept for each mouse. We obtained the reported p-values for the fixed
effects terms using the *lmerTest R* library [56].

Directionality Analysis

LFP signal directionality between pairs of brain regions was estimated using a temporal lag
approach introducing phase offsets as previously described [57]. Butterworth bandpass filters were
applied to isolate LFP oscillations within 1Hz windows using a 1Hz step in the 1□50Hz range.
The Hilbert transform was used to determine the instantaneous phase of the filtered LFPs and the
instantaneous phase offset time series was calculated for each LFP brain region pair. The mean
resultant length (MRL) for the phase offset time series was calculated, yielding the phase
coherence between pairs of LFPs. Temporal offsets were introduced between each pair of LFPs
ranging from □250ms to 250ms in 2 ms steps and the phase coherence was recalculated between
each LFP pair.

Frequency Band Identification

In order to determine the specific frequency bands at which pairs of brain regions demonstrated
clear temporal offset (leading or lagging), we used a fused lasso approach within the $1Hz$ to
$50Hz$ range obtained from the phase offsets data for 34 mice and for each pair of brain regions
from the set: ACC, LPBN, RPBN, CeA, BLA, MD_thal, Po, VPM. Lasso (Least absolute
shrinkage and selection operator), is a model parameter selection and regularization method. The
lasso method minimizes the least squares loss over a set of parameters in a linear regression
problem by imposing the condition that the sum of the absolute values of the parameter estimates

(that is, the ℓ_1 norm of the parameters) is less than some value [58]. This method generally results in

some parameter estimates being zeroed out, in effect selecting the most important covariates that

are more predictive of the response than those with zeroed out regression coefficients. The fused

lasso accounts for temporal or spatial dependence amongst covariates by further imposing a

condition that the sum of the absolute values of the lag one differences between subsequent

parameter estimates be less than some value [59]. Modeling the response under fused lasso then

$$\underset{\underline{\beta}\in\mathbb{R}^p}{\text{argmin}}\|y - X\underline{\beta}\|_2^2 + \lambda\sum_{i=1}^{p}|\beta_i| + \gamma\sum_{j=2}^{p}|\beta_j - \beta_{j-1}|.$$

amounts to finding a set of parameter estimates $\hat{\underline{\beta}} \in \mathbb{R}^p$ as a solution of

The parameters $\underline{\lambda}$ and $\underline{\gamma}$ are positive tuning parameters. The solution of the optimization problem

in (1) depends on the chosen value of $\underline{\lambda}$ and $\underline{\gamma}$, that is $\hat{\underline{\beta}}_{\lambda,\gamma}$, which are chosen via cross validation

in such a way that the test mean squared error is minimized.

We hypothesized that a small number of frequencies are active in these measurements and an

active frequency implies that neighboring frequencies are also active. We applied the application

of fused lasso for signal approximation [60]. We first computed the mean over all observations for

each pair of brain regions and for each mouse. This resulted in 168 observations (8 mice times 21

brain region pairs) each with 50 elements corresponding to the mean phase offset at each

frequency from $1Hz$ to $50Hz$. This 168×50 matrix was then reshaped into a 400×21 response

matrix **Y** in which each column contained the means across frequencies for each mouse for a

particular pair of brain regions. The design matrix **X** was comprised of eight 50×50 identity

matrices stacked on top of each other (one for each mouse) resulting in a 400×50 matrix. The $\hat{\boldsymbol{\beta}}$

matrix of parameter estimates was a 50×21 matrix for which each column contained the model

estimates for the phase offsets over the $1Hz$ to $50Hz$ range for a given pair of brain regions. We

then used the genlasso library (*genlasso: Path Algorithm for Generalized Lasso Problems* ,R

package 1.5) in R to obtain estimates $\hat{\boldsymbol{\beta}}$ under the fused lasso model with the tuning parameters $\underline{\lambda}$

and $\underline{\gamma}$ set to 3 and 0.5 respectively. This resulted in 21 different vectors (the columns of $\hat{\underline{\beta}}$) with

bands of non-zero values corresponding to significant phase offset frequencies and zeros

indicating non-significant frequencies for each pair of brain regions.

Frequency bands of interest

The Po-CeA relationship was the only brain region pairing that included a different, 1-7 Hz

band. We accounted for 1-3 Hz and 1-7 Hz in our multiple comparisons corrections, however

we did not observe substantial evidence for the biological relevance of the 1-3 Hz band across all

brain regions we observed given how small the differences in this band were relative to the other

frequency bands and the fact that it was only significant in one region pairing and not all

(Supplemental Figure S3) so we focused on the 4-7 Hz and 8-13 Hz frequency bands.

Analysis of Power and Coherence over time

We applied a log transformation to the recordings to preprocess the power data. For every brain

region and every mouse on each day of the experiment, we averaged the log-transformed power

across all channels within each brain region and all frequencies within specific frequency bands.

This resulted in a single log-power time series for every frequency band and for every brain

region being associated with each mouse on each day of an experiment. We computed the mean

log-power over baseline recordings for each frequency band and brain region for each mouse on

each day of a given experiment. We normalized the post-injection log-power recordings relative

to the baseline by subtracting these means from the corresponding log-power post-injection

recordings.

We preprocessed the coherence data similarly. For each pair of brain regions and all frequencies

within specific frequency bands, the coherence recordings were averaged across all the channel

pairs, resulting in a single coherence time series for every frequency band across each mouse's 28

pairs of brain regions on each day. We then applied a log-odds transformation to the averaged

coherence data to normalize the post-injection log-odds coherence recordings using the same

approach as we did for the power data. Specifically, we calculated the mean of the baseline log-

odds coherence for each frequency band and brain region pair and subtracted these means from

the corresponding post-injection log-odds coherence values.

After preprocessing, we used these data to fit the FOS model using generalized additive model

(GAM) function in the mgcv library in R to plot 30 second rolling means in order to identify

underlying trends over time [61]. We first computed 30-second rolling means over each mouse's

log-power post-injection time series for each day of the experiment, for each frequency band and

each brain region, using the *tf_smooth* function in the *R* library *tf* [62]. Next, we averaged the

vehicle 30-second rolling mean time series across days. This resulted in each mouse having one log-power 30 second rolling mean post-injection time series for each treatment (vehicle and CGRP), frequency band, and brain region. We used these series for all the mice to compute the average and one standard deviation and obtained one mean CGRP log-power time series, one mean vehicle log-power time series and their respective one standard deviation time series for each frequency band for each brain region. For each frequency band in each brain region, we plotted these 30-second rolling mean time series together to facilitate comparison between the CGRP and vehicle conditions. We used the *R* packages *ggplot2*[63] and *tidyfun*[64] to plot these figures.

We applied the same procedure to the coherence data, resulting in one mean CGRP log-odds coherence 30-second rolling mean time series, one mean vehicle log-odds coherence 30-second rolling mean time series, and their respective one standard deviation time series for each frequency band across each pair of brain regions. We plotted these 30 second rolling mean time series together for comparison in the same manner as the log-power analyses.

For the power GAM analysis, we computed the mean of each mouse's log-power post-injection vehicle time series, resulting in one log-power post-injection time series for vehicle and one for CGRP for each frequency band in each brain region. We obtained one pairwise difference time series for each frequency band in each brain region for each mouse by subtracting the mouse's log-power vehicle time series from the corresponding CGRP time series. For each frequency band in each brain region, we used 90 minutes of the post-injection recordings for the power data analysis.

The coherence analysis is performed similarly, except we computed the mean log-odds coherence instead of the mean log-power. This resulted in each mouse having one log-odds coherence post-injection time series for the vehicle and one for CGRP for each frequency band across each pair of brain regions. We then subtracted each mouse's log-odds coherence vehicle time series from the corresponding CGRP time series to obtain one pairwise difference time series for each frequency band across each pair of brain regions for each mouse. For each frequency band in each brain region, we used 60 minutes of the post-injection recordings data for the coherence data analysis.

The coherence analysis is performed similarly, except we computed the mean log-odds

coherence instead of the mean log-power. This resulted in each mouse having one log-odds

coherence post-injection time series for the vehicle and one for CGRP for each frequency band

across each pair of brain regions. We then subtracted each mouse's log-odds coherence vehicle

time series from the corresponding CGRP time series to obtain one pairwise difference time

series for each frequency band across each pair of brain regions for each mouse. For each

frequency band in each brain region, we used 60 minutes of the post-injection recordings data for

the coherence data analysis.

For the power and coherence analyses, we used the *gam* function in the *R* package *mgcv* (Wood,

2011) for FOS regression. Each model had the following form:

$$y_{r,f}(t) = \beta_{r,f} + s_{r,f}(t) + \varepsilon_{r,f}(t), \qquad\qquad (1)$$

where $y_{r,f}(t)$ denotes the pairwise difference in log-power or log-odds coherence at time point t,

brain region (for power) or pair of regions (for coherence) r and frequency band f; $\beta_{r,f}$ denotes

the (r, f)-specific model intercept; $s_{r,f}(t)$ denotes the (r, f)-specific model smooth function-over-

time effect at time point t; and $\varepsilon_{r,f}(t)$ denotes (r, f)-specific independent centered Gaussian

noise at time point t. This FOS regression model assumes that the pairwise differences in log-

power or log-odds coherence are frequency band and region- or pair-specific functions of time.

We fit this model in *R* as follows. For the mouse i, $X_{i,r,f}$ denotes the *R* data.frame with columns

representing time and response, where t is the vector of evenly spaced time steps and y denotes

the observed pairwise difference in log-power or log-odds coherence for mouse i in brain region

or pair of brain regions r and frequency band f. Let *X_rf* denote the data.frame formed by

stacking the elements of $X_{i,r,f}$ on top of each other for $i = 1, ...,34$. We fit the FOS regression

model using the *mgcv* package in *R* as follows:

model_rf = gam(y ~ s(t), data=X_rf);

The fit model_rf contains the fitted values indicating the mean effect of CGRP over time along

with p-values for both the intercept and function-over-time components $\beta_{r,f}$ and $s_{r,f}(t)$

respectively. We plotted each model's fitted values along with a two-standard error region.

When these two-standard error confidence regions contain zero, we interpret this as indicating no

significant CGRP effect at that point in time.

Analysis of Sumatriptan over time

For 10 of the 34 mice from the previous experiment, we collected additional power and

coherence data over three more days. This n was determined for a repeated measures design in

keeping with similar studies[65,66]. Each mouse received a vehicle injection on the first day, a

Sumatriptan injection on the second day, and an injection of both CGRP and Sumatriptan on the

third day. This resulted in each of these 10 mice having a total of three vehicle injection days,

one CGRP-only injection day, one Sumatriptan-only injection day and one CGRP-with-

Sumatriptan injection day. The recorded power and coherence data were preprocessed as

described earlier, and each set of vehicle time series' were averaged into one vehicle time series

such that each mouse had a vehicle time series, a CGRP-only time series, a Sumatriptan-only

time series and a CGRP-with-Sumatriptan time series associated with each frequency band and

each brain region for power or pair of brain regions for coherence.

For both power and coherence, we again computed the 30-second rolling means for the post-

injection time series using the *tf_smooth* function from the R library *tf* (Scheipl & Goldsmith,

2024)[62] and plotted them in the same manner as in the previous experiment. Now, however, each

plot has four lines: one for vehicle, one for CGRP, one for Sumatriptan and one for CGRP-with-

Sumatriptan.

For the power and coherence FOS GAM analyses, we focused only on comparing CGRP to

CGRP-with-Sumatriptan. These analyses follow the previous section. For every mouse, we

obtained a pairwise power difference time series for each frequency band in each brain region by

subtracting each mouse's log-power CGRP-with-Sumatriptan time series from the corresponding

CGRP-only time series. Similarly, for the coherence analysis, we obtained the pairwise

difference time series for each frequency band across each pair of brain regions for each mouse

by subtracting each mouse's log-odds coherence CGRP-with-Sumatriptan time series from the

corresponding CGRP-only time series. We used 45 minutes of the post-injection recordings data

for our power analyses, and we used 20 minutes of the post-injection recordings data for our

coherence analyses.

For the power and coherence analyses, we used a function-on-scalar regression using the *gam*

function in the *R* package *mgcv* (Wood, 2011)[67]. Each model had the same form as equation (1)

above and was fit in the same manner as the previous experiment. We plotted the fitted values

along with two-standard error confidence bands and obtained each model's p-values

corresponding to the intercept and function-over-time components $\beta_{r,f}$ and $s_{r,f}(t)$ respectively.

Analysis of Early Life Stress Data over Time

We collected power and coherence data from 11 different mice over three days. Similar to the

experiment comparing CGRP to vehicle, each mouse received vehicle injections on the first and

third days and a CGRP injection on the second day. Of these 11 mice, six were ELS mice and

five were control mice. We preprocessed the power and coherence data as described above. This

experiment differed from the previous two experiments in that each treatment group consisted of

different mice.

We computed the 30 second rolling mean plots in a similar manner as the previous two

experiments using the *tf_smooth* function in the *R* library *tf* (Scheipl & Goldsmith, 2024)[62]. For

the power analysis we first computed 30 second rolling means over each mouse's log-power

post-injection time series. Next, we averaged each mouse's vehicle 30-second rolling mean time

series across days. This resulted in each mouse having one log-power 30 second rolling mean

post-injection time series for each treatment (vehicle and CGRP), frequency band, and brain

region. We grouped the observations by ELS vs. control and vehicle vs. CGRP and computed the

average and one standard deviation log-power 30 second rolling mean time series across the

observations within each group. This resulted in separate mean and one standard deviation log-
power time series for each group (ELS-with-CGRP, ELS-with-vehicle, control-with-CGRP and
control-with-vehicle) for each frequency band for each brain region. For each frequency band for
each brain region, we plotted these 30 second rolling mean time series together for comparison in
the same manner as the previous experiments.
466
467 We processed the coherence data in the same way so that we had separate mean log-odds
468 coherence 30 second rolling mean time series for ELS-with-CGRP, ELS-with-vehicle, control-
469 with-CGRP and control-with-vehicle along with their respective one standard deviation time
470 series for each frequency band for each pair of brain regions. We plotted these 30 second rolling
471 mean time series together for comparison in the same manner as the log-power analyses.
472
The different experimental design implied we use a different GAM model. We performed FOS
GAM analyses for each frequency band in each brain region for power or pair of brain regions
for coherence based on a two-factor model with interaction, where the first factor was ELS vs.
control and the second factor was CGRP vs. vehicle. Because each mouse had observations over
three different days, we used a mixed effects model with random intercept and smooth function-
over-time terms for each mouse. Each model used either log-power or log-odds coherence as the
response and had the following terms:
● Overall model intercept
● Mouse-specific random intercept
● Mouse-specific random smooth function-over-time
● Overall smooth function-over-time
● ELS effect intercept
● CGRP effect intercept
● ELS:CGRP interaction effect intercept
● ELS effect smooth function-over-time
● CGRP effect smooth function-over-time
● ELS:CGRP interaction effect smooth function-over-time

This FOS regression model assumes that the log-power or log-odds coherence are frequency

band and region- or pair-specific functions of time. For both the power and coherence analyses

we used 45 minutes of the post-injection recordings.

We fit this model in R as follows. For the mouse i, $X_{i,r,f}$ denotes the R data.frame with columns

representing time, ELS vs. control labels, CGRP vs. vehicle labels, the interaction between ELS

and CGRP labels, the mouse ID and response. Let t denote the vector of evenly spaced time steps

repeated for each day's recordings, els denote the vector of factor labels indicating whether an

observation is associated with an ELS or control mouse, $cgrp$ denote the vector of factor labels

indicating whether an observation is associated with CGRP or vehicle injection, els_w_cgrp

denote the ELS:CGRP interaction dummy variable vector indicating whether an observation is

both CGRP and ELS or not, $mouse$ denote the mouse ID associated with the observation and y

denote the observed log-power or log-odds coherence for mouse i in brain region or pair of brain

regions r and frequency band f. Let X_rf denote the data.frame formed by stacking the elements

of $X_{i,r,f}$ on top of each other for $i = 1, \ldots, 11$. We fit the FOS using the *mgcv* package in R as

follows:

model_rf = gam(y ~ s(mouse, bs='re') + s(mouse, t, bs='re') +

508   s(t) + els + cgrp + els_w_cgrp + s(t, by=els) +

509   s(t, by=cgrp) + s(t, by=els_w_cgrp), data=X_rf);

From the fit model_rf, we obtained the smooth function-over-time components of the fitted

values which provide the estimated ELS effects over time and the estimated ELS:CGRP

interaction effects over time. We plotted these estimated effects over time along with a two

standard error confidence region. We also obtained from the model the p-values associated with

the ELS effect intercept, the ELS effect smooth function-over-time, the ELS:CGRP interaction

effect intercept and the ELS:CGRP interaction effect smooth function-over-time.

We took all of the p-values obtained from each experiment's analyses and computed FDR

corrections on them in order to perform multiple comparisons hypothesis tests of whether or not

there was a significant difference in treatment effects over time for a given brain region or pair of

brain regions in a particular frequency band.

Changepoint analysis

Changepoint analysis was carried out as originally described in Zhang & Chan (2024) [68]. For this

analysis, we evaluated the data in the two frequency bands we found to be most impacted by

peripheral CGRP (4-7Hz and 8-13 Hz).

For conciseness, we shall simply describe the case of detecting changes in the 4-7 Hz component

of the LFP, as the approach is the same for both frequency bands. Specifically, for each mouse,

the 7-dimensional LFP times series are blocked into non-overlapping 10-second blocks, over

each of which (i) we computed its smooth spectral density matrix estimate (as a function of

frequency with 1 Hz steps) via a Bartlett window, (ii) normalized the spectral density matrix

estimates so that their diagonal elements are identically equal to 1 and the off-diagonal elements

measuring the LFP coherency between pairs of brain regions, and (iii) summarized the

normalized spectral density matrix function by averaging over the 4-7 Hz.

The matrices per the 10-second blocks are 7 by 7 complex valued matrices, and their real part

can be interpreted as proportional to the correlation matrix of the 7-dimensional 4-7 Hz

(periodic) components of the LFP time series, with the latter denoted as Ξ_t for the t-th block

(section 7.1 in Brillinger (2001)[69]. The change-point detection then proceeds to find the epochs

(i.e. the change points, if they exist) across which Ξ_t's undergo a change, as follows. Let ν

denote the projection vector, of unit length, pointing the principal direction of the change in

Ξ_t, i.e., the quadratic form $\nu^\top \Xi_t \nu$ changes the most across the change point, among any unit

directional vectors. The projection ν is then the principal direction along which the pre- and

post-change-point correlation matrices of the 4-7 Hz periodic components of LFP differ from

each other to the greatest extent.

The projection vectors reveal the structure of the change of the coherency matrix over the

detected change point. Larger values in magnitude indicates larger changes for the corresponding

brain regions. Opposite signs within the projection vector are indicative of two major types of

organizational matrix changes, which can be identified as blocks within the change in coherency

matrix across the changepoint. These blocks can segregate coherency into activity within a
subgroup of brain regions as two groups and activity between the subgroups (as in Fig. 7E). If
the projection comprises coefficients of opposite signs, the brain regions are split into two groups
according to the signs and may signify that across the change point, the inter-brain-region
correlations between the two groups change differently, i.e., a variably heightened or lessened
synchrony between the two groups, while there are no within-group changes. In the case that all
coefficients of the projection vector are of the same sign, the correlations between brain regions
of the 4-7 Hz periodic LFP components change in a more uniform way to an extent proportional
to the product of their coefficients in the projection vector, i.e., a variably heightened or lessened
synchrony across the 8 brain regions. The projection then comprises the direction of change in
each brain region's LFPs along which the coherency structure of the LFPs sustains a change after
induction by CGRP.

Changepoint Clustering

For each possible combination pair of mice, we calculated the cosine distance between their
projection directions. We then used this distance for k-medoids clustering. The number of
clusters is estimated by optimum average silhouette width. And the clustering procedure is
implemented on each of the three days for each frequency band.

RESULTS

ELS exacerbates migraine-related behavioral phenotypes in CD1 mice

Stress, especially stress experienced early in life, is a naturally-occurring event in the life of both
humans and rodents that has been demonstrated to play an important role in migraine and has
been implicated as altering plasticity in several affect-related brain regions including: the anterior
cingulate cortex (ACC), amygdala (AMY), posterior thalamus (Po), ventroposteromedial
thalamus (VPM), medial dorsal thalamus (MDthal), and parabrachial nucleus (PBN). [10,11,70,71].
Specifically, adverse childhood events, also referred to as early life stress (ELS), are associated
with migraine diagnosis in adulthood [6,7]. In preclinical models, previous studies have found that
ELS enhances migraine-like phenotypes in rats [72] and female C57Bl/6J mice [73], but importantly,
this relationship hasn't previously been identified in the context of CGRP.

We induced ELS in the outbred strain, CD1, mice by utilizing a maternal separation stress

paradigm that also included early weaning and limited nesting (Figure 1A). A total of 52 litters

were utilized for experiments, and animals were split into balanced groups according to weight

and cohort prior to experiments (Figure 1A). Animals were randomly divided in an age-balanced

way such that half of the litters underwent ELS and half remained group-housed (control). To

assess the effect of ELS on migraine-related phenotypes, we administered peripheral PBS or

CGRP to both groups (Control-Veh/Control-CGRP and ELS-Veh/ELS-CGRP). Following our

stress paradigm (see Methods), we waited until mice reached adulthood (P66-69) and performed

behavioral assays to assess anxiety-like and migraine-like behavior. We hypothesized that ELS

would exacerbate migraine-related phenotypes in mice treated with CGRP compared to control

mice.

First, we assessed anxiety-like behavior at P66-69 using the elevated plus maze and determined

that animals exhibited an anxiety-like phenotype after undergoing ELS. We found that mice that

underwent ELS spent less time in the open arms of the elevated plus maze (p = 0.04, t(91) =

2.11, unpaired t test) which indicates that the ELS stressed the animals sufficiently to

demonstrate an affect-related phenotype in adulthood (Figure 1B).

Cutaneous allodynia is present in 60% of patients with migraine [74]. We assessed cutaneous

allodynia in CD1 mice at P72-79 by measuring mechanical sensitivity on the plantar surface of

the hindpaw using the von Frey test. We log-transformed the data and utilized a mixed-effects

analysis and found the interaction between ELS and CGRP treatment was significant (p<0.001).

Using a Bonferroni's multiple comparisons post-hoc test, we found that both Control-CGRP (p

<0.001, t(94) = 5.92) and ELS-CGRP treated mice (p<0.001, t(94) = 8.18) exhibited increased

mechanical sensitivity (50% threshold) compared to baseline sensitivity thresholds (Figure 1C).

As expected, neither the Control-Veh (p =0.08, t(94) = 1.80) nor ELS-Veh (p = 0.16, t(94) =

1.41) groups exhibited significant differences in sensitivity thresholds.

At P81-85, we injected CGRP or PBS 30 minutes prior to assessing light-aversive behavior.

Light aversion, or sensitivity to light, is present in nearly all patients with migraine, and about

half of patients reported photophobia (light aversion) as the most bothersome symptom of

migraine [75]. The light/dark assay has been utilized to assess migraine-like behavior during the

CGRP mouse model of migraine [41]. Light aversion was measured by the average amount of time

spent in the light zone of the light/dark box during the first 15 minutes of the testing period
(Figure 1D). Using a Two-way ANOVA followed by Bonferroni's multiple comparisons post-
hoc test, we found that the interaction between ELS and CGRP treatment was significant
($p<0.0001$, $F(3, 93) = 7.90$). CGRP-treated control animals spent less time in the light following
CGRP compared to PBS ($p<0.001$, $t (93) = 4.55$). CGRP-treated control animals also spent less
time in light following CGRP compared to PBS ($p<0.001$, $t(93) = 5.18$). As expected, both
Control-Veh ($p = 0.74$, $t(93) = 0.337$ and ELS-Veh ($p = 0.78$, $t(93) = 0.283$) did not show
significant differences in time spent in light. Because the light/dark test can also be used to
assess anxiety-like behavior, the standard in the field is to run an open field test under the same
conditions as the light/dark test and verify that animals do not spend more time in center [76]. We
tested the animals at P83-86. We found that there were no significant differences in time spent in
the center during the open field test ($p= 0.76$, $F(3, 97) = 0.394$) (Figure 1E), thus leading us to
conclude that CGRP-treated animals did in fact demonstrate light aversion not attributable to
anxiety during the test. Overall, we found that ELS predisposes CD1 mice to enhanced
peripheral CGRP-induced migraine-related phenotypes.

Peripheral injection of CGRP induces migraine-like behavior in electrode-implanted animals

To gain insight into brain mechanisms of migraine, and the ways that ELS impacts such
mechanisms, we implanted electrodes and recorded neurophysiological activity from multi-site
electrode wires in the ACC, AMY, MDthal, Po, VPM, and LPBN/RPBN (Figure 2A,
Supplementary Figure S1). Such an approach allows us to quantify spatiotemporal dynamics
following peripheral CGRP administration based on local field potential (LFP) activity and
subsequently perform behavioral studies to phenotypically validate the effect of peripheral
CGRP (Figure 2B). We first determined that our implanted animals exhibit similar light aversive
behavior following CGRP administration as unimplanted animals (Supplemental Figure S2).
There was no significant interaction between CGRP treatment and the electrode-implanted
condition ($p = 0.79$). However, as expected, CGRP treatment did result in decreased time spent
in the light on the CGRP treatment day compared to the vehicle day ($p = 0.01$, Supplemental
Figure S2). We found no significant effect of repeated injection ($p = 0.67$, Supplemental Figure
S2), nor due to the day itself ($p = 0.77$). Overall, we demonstrated that our electrode-implanted
animals exhibit the same phenotype as unimplanted animals, consistent with phenotypes

previously published [42]. We also analyzed the total amount of time spent in light and resting time
in dark across all implanted animals utilized in subsequent experiments (Figure S3). Again,
implanted animals injected with CGRP exhibited decreased time spent in light during all 30
minutes (n = 31, p <0.001, Figure S2) and the first 15 minutes of the light/dark test (n = 31,
p<0.001, Figure S2).

LFP directionality points to Amygdala-centric subnetwork

To identify LFP oscillation frequencies of interest, we calculated directionality of information
flow from one brain region to another between brain region pairs based on a temporal lag
approach in which phase offsets are introduced. Multiple studies have demonstrated important
biology and behavioral predictions in frequencies where one region of a circuit clearly leads
another between coherent brain regions [57,65,77-81]. The temporal offset at maximal synchrony
indicates which region is leading and lagging at each frequency [57,79,82-84]. In order to specify
precise frequency bands in which information flow appears to be directional from one region to
another, we applied a fused lasso approach, enforcing a lasso penalty on absolute value
differences between contiguous frequencies in phase offset (see methods, Figure 2).

Given the roles of the amygdala and thalamic nuclei in affect and migraine, and that they appear
to play a hub role in migraine neural circuitry (Figure 8A), we focused on the directional
relationships between these regions. To determine frequency bands of interest, we utilized a
lasso regression on the greatest mean temporal phase offsets within these pairings (Figure 2). We
found that AMY leads all of the thalamic (MD/Po/VPM) and bilateral parabrachial
(LPBN/RPBN) nuclei in the 4-7 Hz frequency range (Figure 2). Continuing in this subnetwork,
we found that all other regions led the AMY in the 8-13 Hz range (Figure 2). As such, we
focused our migraine mechanistic studies on these frequency bands.

CGRP induces decreases in LFP power across migraine-related brain regions

Our analysis indicates that 4-7 Hz is a frequency band in which the AMY is a major leading
region. This frequency range has been identified in various brain regions to be involved in
functions like sleep, memory, and anxiety [83,85-90]. We hypothesized that we would observe
significant differences in LFP power across each of the brain regions we implanted in the 4-7 Hz
frequency. All data were normalized to the baseline period before plotting, and the time series

was evaluated starting from the 30-minute mark, corresponding to the injection time for each
animal. Because we're interested in both the magnitude of response as well as the time-
dependence, we employed a Function on Scalar General Additive Model analysis (FOS)
followed by an FDR correction, testing the hypothesis that there are differences in LFP power
across time in response to peripheral CGRP within each brain region. This analysis revealed that
LFP power in the 4-7 Hz frequency band was significantly different across all regions we
implanted over time when animals were treated with CGRP compared to when treated with PBS
(Figure 3). Across all seven brain regions (ACC, AMY, MDthal, Po,VPM, LPBN, and RPBN)
there was a decrease in LFP mean logpower following CGRP administration compared to
vehicle control initially following the injection point (Figure 3A) and gradually increasing until
reaching a similar mean logpower under the vehicle condition (Figure 3B-H). ACC (Figure 3B)
contained the greatest relative changes immediately post-injection. The peak changes in power
induced by CGRP occurred around 10-30 minutes post-injection, with the effects primarily
resolving by 45 minutes post-injection (Figure 3, Supplemental Figures S6-7). Some regions
leveled off in change by 60-min post injection or beyond but do not reach zero difference
between CGRP and vehicle within the timeframe observed (e.g. Fig 3B,F,H, Supplemental
Figures S6-7). The general timeframe of activity following peripheral CGRP injection from these
data are highly consistent with what has been reported in terms of behavioral response to
peripheral CGRP using facial grimace[66]. Notably, the VPM appears to be impacted first and
resolve first in both frequency ranges, with the PBNs peaking and resolving on the later side
(Supplemental Figures S6-7). ACC also was the last or among the last to resolve at both
frequencies (Supplemental Figures S6-7).

We also hypothesized that the 8-13 Hz frequency band was a relevant frequency band as all of
the other brain regions led the AMY in the 8-13 Hz frequency band (Figure 2). Similar to the 4-7
Hz frequency band, we observed changes in mean logpower in all regions of our network,
though minimal in the 1-3 Hz band (Supplemental Figures S4-5). All three frequency bands were
taken into account for our FDR corrections (see methods). In the 8-13 Hz band, the ACC
(Supplemental Figure S5) contained the greatest relative difference, followed by the AMY and
LPBN and overall power changes resolved by around 45-60 minutes post-injection
(Supplemental Figure S5). Overall, we observed changes in LFP power across all brain regions

of the network in two different frequency bands in response to peripheral CGRP, supporting

centrally-mediated activity during a peripherally-induced migraine brain state.

CGRP induces disruptions in LFP coherence across migraine-related brain regions

Next, we wanted to evaluate how functional connectivity across the network was altered, if at all,

during a CGRP- induced migraine brain state, especially between AMY-brain region pairings.

To answer the question of how brain regions communicate with each other across time during a

migraine-like brain state, we analyzed coherence among brain region pairings following CGRP

administration compared to vehicle conditions.

High mean coherence was observed across the major pairings of our network (e.g. pairings with

AMY and Po hubs) across all frequencies of interest (i.e. 1-13 Hz) +/- CGRP (Figure 4A). All

LFP data was normalized to baseline periods and plotted starting at the 30-minute point onward

providing a stabilizing transform while preserving direction, as log odds coherence. Similar to

CGRP-induced changes in power, the disruptions in coherence in response to CGRP were most

pronounced in the first 10 minutes post-injection, but unlike the power, they largely resolved

much faster, within ~20 minutes post-injection (Figure 4B). The FOS model demonstrated that

there are significant disruptions in coherence across network brain region pairings in response to

CGRP compared to vehicle (Figure 4C).

In both the 4-7 Hz band and 8-13 Hz frequency band, we observed significant coherence

disruption in all AMY brain region pairings, the most of any of the regions (Figure 4C). Brain

regions in which we observed the greatest magnitudes of CGRP-induced change in coherence

(i.e. difference in mean log-odds coherence) were: the AMY-LPBN (0.218), AMY-VPM

(0.213), and ACC-LPBN (0.223; all p= 3.74e-16, Figure 4D-G, Supplemental Figure S8-9).

Notably, AMY-VPM coherence exhibited the most temporally sustained disruptions across time

(~35 minutes post-injection, p= 3.74e-16, Figure 4E-F). These findings support the critical role

of electrical activity alterations in the amygdala during migraine attacks. Disruption of AMY-

related LFP patterns appear to underlie the endogenous migraine brain state, leading to the

painful and affective symptoms associated with migraine.

Sumatriptan partially reverses the effect of CGRP

To contextualize the clinical relevance of these CGRP-induce brain network dynamics, we
evaluated how CGRP-induced migraine-related brain network dynamics respond to a common
acute migraine treatment, sumatriptan[91]. Sumatriptan is a serotonin 1B/D agonist that was
originally thought to have peripheral mechanisms of action but recently has been found to act in
the brain as well[92,93]. Brain network dynamics measured by fMRI of thalamus and amygdala
have been demonstrated to be altered in response to sumatriptan [94]. In other rodent models
sumatriptan has had somewhat limited efficacy but does at least partially reduce CGRP-induced
migraine-like phenotypes [42,66]. We hypothesized that coadministration of sumatriptan and CGRP
would attenuate some or all of the changes in power and coherence induced by CGRP.

We compared responses between CGRP alone and sumatriptan+CGRP and included vehicle and
sumatriptan-alone conditions to assess the direction of the sumatriptan+CGRP relative to CGRP
and vehicle (Figure 5, Supplemental Figures S10-11). Interestingly, ACC, and AMY in the 8-13
Hz frequency band were the only power features in which co-administration of
sumatriptan+CGRP attenuated the CGRP response, with AMY demonstrating the most clear
response (Figure 5B-C, p=3.74e-16). With regard to coherence in the 20 minutes following
injection most disrupted by CGRP, we found only modest effects of sumatriptan+CGRP that
occurred late in the timeframe (Supplemental Figure S10). Interestingly, the brain region with
the greatest number of pairings reaching statistical significance and the greatest effect of
sumatriptan blunting CGRP response was ACC, including ACC-AMY 8-13 Hz (p= 1.62e-13,
Figure 5F), ACC-VPM 4-7Hz (p=7.5e-7, Figure 5H), 8-13Hz (p=3.74e-16, Figure 5G), ACC-Po
8-13Hz (p=4.68e-8, Figure S11), ACC-RPBN 4-7Hz (p=1e-6, Figure S10-11).

Sumatriptan also appeared to impact the network in additional ways beyond blunting the
response to CGRP. The coherence changes between the LPBN and Po in the 8-13 Hz band, while
significant, were not in the direction of CGRP reversal (p= 3.74e-16, Supplement Figure S10-
11).

Overall, coadministration of sumatriptan and CGRP blunted the impact of CGRP on LFP power
in the ACC, and AMY, and coherence across much of the network, especially in ACC-related
region pairings and at later time points where the impact of CGRP is beginning to wane. These
findings appear largely consistent with the timing and degree of sumatriptan blunting of CGRP
in behavioral studies in mice[42,66].

ELS induces changes in LFP power and coherence in response to CGRP

Having identified a peripheral CGRP-induced migraine brain network response, we next set out to determine whether such network components could explain the exacerbating impact of ELS on CGRP-induced migraine-like phenotypes. Given the role of the AMY in stress and affect and the changes in LFP power and coherence we observed in response to CGRP, we hypothesized that animals exposed to ELS would exhibit significant differences in LFP power in the AMY and AMY-related coherence in response to CGRP, compared to animals that remained group-housed during early life. ELS and group-housed control animals were treated with vehicle on Days 1 and 3, and CGRP on Day 2. Overall, the ELS-CGRP exhibited the strongest decrease from baseline of relative power and coherence across regions and pairings and the greatest amount of changes from baseline compared to control groups (ELS-vehicle, Control-vehicle, Control-CGRP, Figure 6). We examined whether there was a significant interaction between ELS and CGRP response across time in these groups, as well as network-wide impacts of ELS itself in the absence of CGRP. We found significant interaction between ELS and CGRP with regard to LFP power in either the 4-7 Hz or 8-13 Hz to varying degrees across all brain regions of the network (i.e., a significant interaction, Figure 6, Supplemental Figure S-9).

With regard to ELS x CGRP interaction in power, in the 8-13Hz band, ACC, AMY, Po and VPM were the most significant features (Figure 6), whereas in the 4-7Hz range, the thalamic regions were (Po, VPM, MDthal, Figure 6, S12-13), implying likely AMY x thalamus mechanisms. There were no significant effects of ELS alone in any brain region or frequency (Supplemental Figure S14).

Given the differences in power reactivities during the ELSxCGRP interaction, we examined whether coherence patterns in the network were also impacted (Figure 6G). Many significant differences in coherence were observed between control and ELS groups after CGRP administration, including AMY-related pairings, which were consistently altered. Specifically, coherence was most disrupted in the ELS group in response to CGRP for AMY-thalamus pairings; AMY-Po 8-13Hz (ELS x CGRP interaction p= 3.74e-16, ELS effect p= 0.79, Figure 6I) and AMY-MDthal 4-7Hz, 8-13Hz (ELS x CGRP interaction p= 3.74e-16, ELS effect p= 0.68 Figure 6H, K) pairings, and AMY-VPM 4-7 Hz (interaction p=3.57e-10, ELS effect p=0.86, Figure 6J). When considering coherence pairings that had been most-impacted by CGRP, beyond

AMY-VPM, AMY-LPBN showed no significant interaction of ELS and CGRP (4-7 Hz p=0.18;
8-13Hz p=0.22), ACC-LPBN demonstrated no interaction effect at 4-7Hz (p=0.66) but a strong
interaction at 8-13 Hz (3.74e-16), and LPBN-Po demonstrated a moderate interaction (ELS x
CGRP interaction p= 5.65e-5, ELS effect p= 0.67, Supplemental Figure S13).

Overall, these findings indicate that ELS induces changes across the network, and especially
AMY and thalamic-related brain circuits in the context of CGRP across critical neural features
that are responsive to CGRP. Given that there is no effect of ELS alone on any of these features
suggests that ELS sensitizes the brain's responsiveness to CGRP. Several coherence pairings that
are most responsive to CGRP (AMY-Po and AMY-VPM) are rendered even more responsive
with pretreatment with ELS. Our data thus demonstrate that ELS predisposes mice to a CGRP-
susceptible brain state, resulting in compromised brain circuitry within CGRP-related pain
pathways in the presence of CGRP. Such findings establish a novel model that predisposes
rodents to a migraine-susceptible state (behaviorally and neurophysiologically), enabling
rigorous investigation of dysfunctional migraine-related circuitry and providing a framework for
studying the CGRP-vulnerable brain state (Figure 8).

Changepoint Analysis identifies differing brain network responses to peripherally delivered
CGRP

Up to this point, we have provided a thorough demonstration of the average changes in brain
activity across a population of mice over time during peripheral induction of migraine with
CGRP and the ways in which ELS impacts these network components toward a vulnerability to
CGRP. In order to investigate the possibility of individual vulnerability to CGRP, we employed a
change-point analysis to examine the evolution of important network-wide events in each
individual in the CGRP-induction of a migraine-like state. This spectral-based method is used for
detecting changes in the correlation structure of specific frequency components in the LFP (time
series), over time, and the "principal component" of the LFPs, along which a change occurs. The
change-point detection identifies the epochs (i.e. the change points) across which a data-driven
linear combination of the periodic components of LFP in our frequency bands of interest undergo
a variance change. Thus, a changepoint reflects a point where coherence across multiple brain
regions have a coordinated and significant change. We reasoned that the first changepoint after
the injection is likely to be the most informative, as this likely reflects an animal's initial CNS

response to the peripheral CGRP. The first changepoint after the injection (at 30 min), occurred

within 40 minutes post-injection across all three days with no obvious relationship between light

aversion behaviors and the timing of the first changepoint (Figure 7A). It does seem, however,

that the time intervals between the change point and injection are smaller on day 2, the day

CGRP was injected, than the vehicle days, 1 and 3. In order to ensure robustness of these

findings, an additional robustness step was implemented, where we repeated the change point

detection algorithm 50 times with different initialization random seeds, and evaluated the output

change point locations by the mode and the projections by the median.

For a given changepoint, there exists a coefficient "projection" vector describing the brain region

relationships that characterize the changepoint, which could be due to an altered synchronization

strength among those brain regions in a given frequency band (see Fig. 7 and below). In order to

determine whether there are common features regarding connectivity changes at the changepoint

across animals in response to CGRP, we conducted a clustering analysis of the mice. We

clustered in an unsupervised way based on the LFP features involved in each animals'

changepoint, by using a cosine distance between the projection vectors of the mice. We reasoned

that different CD1 mice might respond differently at the level of the CNS in response to

peripheral CGRP (i.e. have different projections at the first change point) and we therefore used

a k-medoids clustering approach to identify groups of mice that respond with similar projections.

We used a silhouette coefficient to assess the strength of the clustering structure with average

silhouette width >0.8 for each frequency (4-7Hz and 8-13 Hz). We found four (4-7Hz) and three

(8-13Hz) clusters this way, suggesting different brain network activity responses to CGRP in the

CD1 mouse brain (Figure 7B, S15).

In order to understand the functional differences in the nature of changes in connectivity

associated with these four different types of changepoints, we plotted the projection vectors for

all of the animals grouped by cluster and ordered them from groups that are most to least

photophobic in response to CGRP (Figure 7 C,D). As shown in Figure 7C, different coherency

organizational structures, as indicated by the projection vectors (see Methods), define groups that

are more or less susceptible to the migraine-inducing effects of peripheral CGRP injection. In

other words, the network activity around the first changepoint explains the behavioral phenotype.

To quantify the extent to which network activity can be used to predict hypersusceptibility to

migraine in the mice, we calculated an AUC for the most highly susceptible group compared to
the rest of the animals for 4-7Hz and 8-13Hz (AUC=0.92, p=0.028 4-7Hz; AUC=0.61, p=0.48,
8-13 Hz). In order to assess whether this strong relationship at 4-7Hz between brain network
activity and behavior was specific for CGRP we did the same comparison for days 1 and 3 and
found much weaker associations with behavior (day 1 AUC=0.72, p=0.06, day 3
AUC=0.68,p=0.24). Altogether, these data demonstrate that there is not one global response in
connectivity of all animals to CGRP, that is to say that subgroups of different animals have
differing connectivity response to peripheral CGRP injection across the brain. These data also
demonstrate that coherence changes at the first timepoint following CGRP injection can be used
to predict animals with extreme migraine-like responses based on 4-7HZ LFP activity.

In order to provide a more direct biological interpretation of the network features defining this
network of migraine hypersusceptibility, we examined the coherency matrices immediately
before and immediately following the changepoint (Supplemental Fig. S16-S17). In Figure 7E,
we show the cluster mean of difference of coherency matrix on 4-7 Hz over the first change
point after injection by cluster. For the 4-7 Hz frequency band, clear block-wise structure is
observed for cluster 1 and 3. Interestingly, cluster 1, the cluster in which the animals have the
strongest vulnerability to CGRP as measured by photophobia, is notably the cluster in which
coherence is most increased, rather than decreased, across the network at the first changepoint
following CGRP injection. This is consistent with the idea of the migraine response being
concerted across multiple synchronized brain regions during specific events, while
desynchronizing networks overall.

DISCUSSION

868
Migraine is a complex and profound disorder, which includes experiences of pain and episodic
disabling brain function, impacting up to 20% of the population at some point (~1 billion people
worldwide). Stress is a common trigger for migraine; high levels of stress are reported in
migraine patients and, simultaneously, migraine is comorbid with numerous brain disorders
associated with stress [95,96]. In humans, early life adversity has a well-documented association
with migraine and headache in adulthood[97]. Studies using rodent models have begun to

investigate mechanisms by which this occurs [98]. In this study, we sought to contribute
substantially to our understanding of this mystery of how early life events can result in migraine
symptoms in adulthood and found that neural activity within the amygdala and its functional
connectivity network play important roles.

To understand the development of such a complex disorder as migraine, which induces systemic
and coordinated malfunction of brain systems spanning emotion, sensation, autonomic nervous
system regulation, and cognition, a systems neuroscience approach is needed [39]. The field's
understanding of migraine has evolved significantly, shifting from thinking of it as a primarily
neurovascular disorder to a complex neurological condition involving multiple brain regions,
with different regions engaged during distinct phases [99,100]. Several pieces of evidence point
toward both central and peripheral pathophysiology, including the complex symptomatology
experienced by patients with migraine, differences in functional connectivity between patients
and healthy controls, and varied pharmacological responses; how exactly brain networks
coordinate over time to produce a migraine brain state remain unknown. In this study, we have
implemented a systems-based approach combining multi-site in vivo neurophysiology with
peripheral CGRP injection, and migraine-relevant behavioral assays, in order to understand a
peripherally induced migraine brain state, and the ways in which ELS impacts such a brain state.

We used peripheral injection of CGRP to best parallel human studies and account for the most
universally understood mechanisms of how CGRP initiates a migraine episode. There are several
plausible explanations of how peripheral CGRP ultimately regulates brain function, beginning
with nociceptive primary afferents expressing and releasing CGRP providing input to second-
order neurons of the spinal dorsal horn and trigeminal nucleus caudalis [33,101]. This input is then
transmitted to the parabrachial (PBN) and thalamic nuclei. A monosynaptic connection between
trigeminal afferents and the PBN also exists, which can transduce craniofacial affective pain and
exist in the dura [37] [32]. CGRP is also expressed in projections from PBN neurons to the thalamus
and amygdala [102]. Additionally, CGRP can also directly or indirectly modulate neuronal activity
and synaptic plasticity in several brain regions implicated in migraine, including the cortex,
amygdala and thalamus [102-104]. These data all contribute to our hypothesis that peripheral CGRP
facilitates nociceptive transmission into the brain and contributes to the sensitization of both
primary afferent sensory and second-order pain neurons to contribute to a migraine brain state.

Importantly, by identifying frequency bands with directionality to the flow of information
(Figure 2), our results are interpretable in terms of AMY-leading (4-7Hz) and AMY-receiving
(8-13 Hz) oscillations. Quantifying spatiotemporal dynamics following peripheral CGRP
administration resulted in identification of coordinate changes in power and coherence, which
are particularly interesting when considering the context of migraine as a phasic disorder. In the
4-7 Hz range, we found that CGRP injection led to a decrease in power across all 8 regions
which peaked at 10-30 minutes post-injection, primarily resolved or leveled off by 45 minutes
post-injection (Figure 3). A similar time course of CGRP-induced behavioral sensitization was
found in a study where peripherally administered CGRP induced a significantly greater
spontaneous facial pain response than controls starting 10 min post-injection, peaking at 30
minutes, and lasting for about 75-90 minutes [66]. This is consistent with other preclinical mouse
models of migraine, and, to some extent, human studies of photophobia [42,105]. A recent fMRI
study of patients across the prodrome to pain state transition also reported a large drop in
widespread brain network connectivity from the transition between prodrome and the
pain/headache attack phase as well[106]. While a lot of heterogeneity exists with regard to human
fMRI studies broadly, globally, thalamic connectivity entering a headache attack is both
strengthened and weakened, our findings are largely consistent with human studies of the
migraine attack phase as a whole[100,107]. We observed that VPM was the first region to reach its
maximal activity and the first region for CGRP-induced changes in LFP power to resolve. This
suggests that VPM plays an important role in the initiation of the migraine response across the
brain, perhaps initiating through projections it receives from SpV [25,108]. The ACC was a region
that was last to resolve, extending beyond the timeframe of most observed migraine-like
behaviors[66]. Given the role of the ACC in processing the emotional aspects of pain, and that LTP
has been found to be involved in the ACC in pain and migraine, this finding may point to a
possible role for the ACC in post-drome or interictal migraine phases[109]. Imaging studies of the
postdrome phase have understandably been quite limited in numbers of patients and brain
regions examined[100,110]. Additionally, in light of the field's ongoing debate about peripheral vs.
central mechanisms of CGRP, and the fact that CGRP largely does not cross the blood-brain
barrier, it is especially important that we have begun to map out specific activity across the brain
in response to a peripheral inject of CGRP. Notably, the half-life of CGRP is ~6 minutes and yet

we see sustained activity across the brain, pointing to a peripherally triggered process that gets

sustained at much higher levels of processing.

Given our overall network connectivity hypotheses, we were interested in whether CGRP

disrupted coherence between AMY, ACC, and thalamic regions, and as these regions form key

circuits involved in chronic pain [111] and pain perception [112]. Coherence was disrupted across the

network shortly after injection with CGRP and across the board followed a shorter time course

than power disruptions. Significant disruptions in coherence among brain region pairings were

especially prevalent in the first 10 minutes immediately post–injection of CGRP, and resolved

around 20 minutes post-injection. This time course gives us insight into how information flow

between brain regions is disrupted by a migraine trigger to orchestrate the migraine brain state.

Because CGRP enhances sensitivity to sensory input in both the periphery and central nervous

system, it is possible the changes in coherence are a reflection of a transition into a

hypersensitized brain state [113].

As a well-established therapeutic for migraine, sumatriptan provided an important test of the

impact of a clinically relevant perturbant on the CGRP-responsive network. Mechanistically,

sumatriptan is part of a triptan class of $5\text{-HT}_{1B/1D}$ receptor agonists that can directly suppress

migraine-related increases in CGRP secretion from trigeminal neurons and normalize cranial

CGRP levels, however, the site of action is debated [92,94,114-116]. It can prevent the induction of

sensitization in central trigeminovascular neurons but also has recently been shown to reach

brainstem and hypothalamic structures quite rapidly[92,117,118]. Sumatriptan has partially reversed

behavioral phenotypes of peripheral CGRP injection previously[66]. Coadministration of CGRP

and sumatriptan reversed the changes in power in the ACC and AMY in the 8-13 Hz frequency

band especially at 25 minutes post-injection (Figure 5, Supplemental Figures S10-11).

Sumatriptan reversed coherence disruptions in brain region pairings involving the AMY and

ACC, especially within the first 20 minutes post-injection. An fMRI study with normal, healthy

volunteers demonstrate that peripheral CGRP-induced decreases in BOLD-signal in the caudate

nuclei, thalamus, and cingulate cortex were reversed by sumatriptan at 40 minutes post-CGRP

injection and 15 minutes post-sumatriptan injection. [119]. Our data seem to recapitulate at least the

cingulate aspect of this finding in our mouse model. Given that sumatriptan doesn't ablate all

CGRP-induced effects and even introduces some additional connectivity patterns, such brain

connectivity metrics may be useful to consider in therapeutic development moving forward with

regard to optimizing network responsivity. Future studies should also investigate the effect of

CGRP-targeting pharmacologics (e.g., CGRP monoclonal antibodies, CGRP antagonists) on

CGRP-induced brain states to assess whether network activity is more strongly attenuated.

While other studies in rodent models have investigated a role for ELS in migraine, to the best of

our knowledge, ELS had not yet been investigated in a CGRP mouse model of migraine, nor in

CD1 mice. In our study, ELS significantly enhanced the CGRP-induced photophobia and

allodynia (Figure 1). Further, we investigated the impact of ELS on brain network activity and

found that ELS significantly enhanced the CGRP-induced drop in power found across the

network, especially with regard to the amygdala. Coherence patterns demonstrated that most, but

not all, of the network connectivity pairings that increase most with CGRP are significantly

exacerbated by ELS (AMY-VPM, AMY-Po), providing a plausible mechanism by which ELS

sensitizes the brain to CGRP response. Importantly, AMY-MDthal connectivity also displayed

significantly increased CGRP response in ELS animals, which is especially interesting given the

strong relationship between both of these regions and stress response in the context of mood

disorders[120,121]. Globally, ELS appears to especially predispose the network so that CGRP drives

amygdalar-thalamic coherence even further. Given the role of the thalamus in sensory

gain/sensitivity and the role of the amygdala in the emotional aspect of pain, CGRP-dependent

dysregulation of the functional connectivity between these two regions in the context of ELS

appears to be an important aspect of the way that ELS impacts migraine in adulthood. Given that

ELS alone has no significant impact on the network activity, the interaction of ELS and CGRP

must derive from some network property that is not actively engaged until it gets unmasked upon

stimulation with CGRP. In short, ELS predisposes mice to a CGRP-vulnerability brain state.

Now that we have determined a way to induce and measure such a state, future studies can now

more deeply investigate the detailed cellular and molecular contributors of this CGRP-vulnerable

brain state (Figure 8).

These global findings provide an overall picture of brain network responses generalized across

the population that we measured and ways in which this network is altered by ELS, however, our

changepoint analysis best helps us understand what brain network activity might be most

contributing to individualized migraine-like responses. Given the heterogeneity of migraine

experience, we thought this was especially important to pursue. We identified changepoints,
which are time epochs where LFP signals within specific frequency bands undergo a substantial
and coordinated change in variance (Figure 7). Our change-point analysis clusters suggest
different brain network activity responses to CGRP in the CD1 mouse brain. These clusters,
when ordered from most to least photophobic in response to CGRP, revealed different coherency
organizational structures which defined groups that are more or less susceptible to peripheral
CGRP injection-induced migraine. Importantly, the network activity around the first changepoint
– 10 minutes post-injection – provides sufficient information to explain or even to some extent
predict, the behavioral phenotype. Thus, coherence changes at this first timepoint following
CGRP injection can be used to predict animals with extreme migraine-like response to light. The
cluster of animals with the most photophobic response to CGRP is the only cluster in which
coherence in the 4-7 Hz frequency range is increased at the first changepoint, suggesting an
important transient transition of brain network activity specifically in highly migraine susceptible
animals that differs from the overall population. Pinpointing the differences between migraine
susceptible and resilient brain network activity opens up exciting possibilities for studying
migraine resilience and the ways in which environmental contributors such as ELS make some
individual open to migraine vulnerability.

Our findings provide an important roadmap of endogenous brain network activity in response to
arguably the most clinically relevant mouse model of migraine, induction by peripheral CGRP.
In a context where central and peripheral mechanisms of CGRP's influence in migraine have
been debated for so long, this map with both global (multi-region connectivity) and neural
timescale information provides the needed framework for the next era of work to understanding
brain response to peripheral physiology. Moreover, we have demonstrated that such metrics can
provide important answers about clinically relevant problems such as the ways in which ELS
predisposes the brain to respond to CGRP. Our results provide unique insights into the neural
basis of migraine and provide a reasonable launching point for the development of novel
therapies targeting brain-wide networks and potential targets for neuromodulation.

1022

1023   **CONCLUSIONS**

We describe a brain-wide network neural activity response to peripheral induction of migraine-like activity by CGRP. This is an important step forward in the field because CGRP does not cross the blood brain barrier and the degree to which peripheral induction of CGRP impacts brain activity has been an ongoing debate in the field. Furthermore, we provide evidence for a mechanism by which ELS predisposes the brain to respond to this common migraine trigger of peripheral CGRP. We anticipate that these findings open doors for future brain-network based therapeutic development for migraine.

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Dr. Rainbo Hultman (rainbo-hultman@uiowa.edu)

Materials availability

There were no materials generated during this study.

Data and code availability

- All original code has been deposited at GitLab and will be publicly available as of the date of publication.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

FUNDING DECLARATIONS

We would like to thank our funding agencies who supported this work: NIH Director's New Innovator 1DP2MH126377-01 (RH), the McKnight Foundation McKnight Neurobiology of Brain Disorders Award (RH), the Roy J. Carver Charitable Trust (RH), NINDS T32NS007124 (MJ, SM, and YF), Ramon D. Buckley Graduate Student Award (MJ) and NSF Graduate Research Fellowship 1945994 (ME).

OTHER ACKNOWLEDGMENTS

We would also like to thank Andy Russo and Levi Sowers for helpful conversations and pharmacological expertise and Kiriana Williams for technical assistance. We would like to acknowledge Biorender, which was used to make some of the figures.

AUTHOR CONTRIBUTIONS

Conceptualization, R.H., S.S., and M.J. methodology, M.J., M.E., I.H., X.Z., ;Investigation, M.J. M.E.,Y.F., ; technical support, M.J., M.E.,Y.F., A.J., M.M., S.M., R.V., and J.M.; writing—original draft, R.H., M.J., A.A., and M.E.; writing—review & editing, M.J., M.E., I.H., and X.Z.,; funding acquisition, R.H.. and S.S..; resources, I.H., X.Z., S.S.. and K.S.C..; supervision, R.H., S.S., B.H., and K.S.C.

DECLARATION OF INTERESTS

The authors have no interests to declare.

DECLARATION OF GENERATIVE AI AND AI-ASSISTED TECHNOLOGIES

The authors have nothing to declare regarding AI.

SUPPLEMENTAL INFORMATION

Document S1.
Figures S1–S14

 FIGURE TITLES AND LEGENDS

 Figure 1. ELS exacerbates migraine-related behavior

A) Timeline of ELS experiment with behavior and recordings. (n= Control Veh:24, Control-
CGRP: 24, ELS-Veh: 28, ELS-CGRP: 27) B) Animals spend less time in the open arms of the
elevated plus maze following ELS C) CGRP-treated animals exhibit increased 50% threshold
compared to baseline. D) CGRP-treated animals spend less time in the light during the light/dark
assay. E) There were no significant differences in time spent in the center of the open field test
following the light/dark assay.

Figure 2. Multi-site in vivo Neurophysiological recordings identify AMY-centric network

A) Brain regions targeted in 32 CD1 mice (15 M, 17 F): ACC, AMY, VPM, MDthal, Po, and
bilateral PBN (see also Figure S1). B) Experimental timeline of recordings and behavior C-H)
LFP directional timing between brain region pairings across migraine-related subnetwork. The
mean temporal offset at which each pair of LFPs optimally phase synchronized is plotted for
each frequency 1-50 Hz. The solid blue lines indicated the mean temporal offset where lighter
shaded blue represents standard deviations from the mean (1-3 standard deviations). C) MD–
AMY D) Po–AMY E) VPM–AMY F) LPBN–AMY G) RPBN–AMY H) Summary: The
mean temporal offset across the MD, Po, VPM, AMY, and PBN pairings are the greatest in the
1-3 Hz (green), AMY leads Po; 4-7 Hz (orange), AMY leads all components of the network; and
8-13 Hz (purple), all other regions lead AMY; frequency bands are indicated by the dashed lines.
F) Summarized directionality relationships across all regions. Overall, LFP directionality in a
migraine-related subnetwork is the greatest at the 4-7 and 8-13 Hz frequency bands.

Figure 3. CGRP alters LFP power in the 4-7 Hz frequency band

LFP mean logpower plotted across time (n=32, see also Figure S3-4). A) Representative spectral
power plot of the ACC across 1-13 Hz. Power is represented in decibels (dB) (left). Heat map
showing all p-values across brain regions of interest (right). B-H) Plots showing the mean log-
power across time following CGRP treatment (pink) compared to vehicle PBS (blue) (left) for
each brain region. Function on scalar generalized additive model analysis (FOS) plots showing
the difference between the mean log-power for CGRP and vehicle conditions immediately post-
injection for each brain region (right). Data are represented as 30-second rolling means ± SEM.

Figure 4. CGRP induces disruptions in LFP coherence across migraine-related brain regions

A) High mean coherence across all animals was observed during the 1-13 Hz frequency range (top) AMY-Po (bottom) AMY LPBN. B) Representative coherence plots of the (top) AMY-Po and (bottom) AMY-LPBN across 1-13 Hz. Red arrow indicates injection point. C) Heat map of p-values for differences between CGRP and vehicle mean log-odds coherence. D-F) For each brain region pairing, LFP mean log-odds coherence plotted across time for vehicle and CGRP conditions (left), and difference between vehicle and CGRP conditions (right) (see also Figure S5-6). Data are represented as 30-second rolling means ± SEM.

Figure 5. Coadministration of CGRP and Sumatriptan partially reverses the effect of CGRP

A) Experimental timeline of sumatriptan and CGRP experiments. B) Heat map showing -log10 p-values summarizing the effect of sumatriptan+CGRP on LFP power. Red boxes indicate significant differences in which reversal of the CGRP effect occurred. C) The mean log-power is plotted across time for the CGRP, sumatriptan+CGRP, and vehicle conditions (left) and difference between sumatriptan+CGRP and CGRP conditions (right). (top) ACC (bottom) AMY Power data are represented as 30-second rolling.

Figure 6. ELS alters AMY and Po-related circuitry in the presence of CGRP

A) Heat map of -log10 p-values reflecting the interaction between ELS x CGRP in ELS on LFP power (see also Figure S10) (n=11), B-G) the mean logpower in the AMY and Po are plotted across time for CGRP and vehicle conditions in ELS and control mice. Power data are represented as 30-second rolling means mean ± SEM. I) Heat map of -log10 p-values reflecting the interaction between ELS x CGRP in ELS on LFP coherence J-L) Mean log-odds coherence is plotted across time for Po-related region pairings for CGRP and vehicle conditions in ELS and control mice (see also Figure S12-13). Coherence data are represented as 60-second rolling means ± SEM.

Figure 7. Changepoint Analysis identifies differing brain network responses to peripherally delivered CGRP

A) Time of 1st change point after injection (at 30min) is shown on the y-axis, with mice shown in decreasing order of their behavior from greatest to least photophobia following CGRP. B) 4-7Hz k-medoids clustering of initial changepoint projection features following injection with

CGRP, silhouette width on the x axis, and each mouse on the y axis, the counts and average

silhouette width in each cluster listed on the right, and the overall average silhouette width on the

bottom. C) 4-7Hz projection features for each mouse by cluster (colors to left). Brain regions are

represented across the bottom with each mouse representing a different row, ordered by the

cluster average of highest relative dark ratio (most photophobic) at the top. The legend shows the

color of values of each cluster in the heatmap. D) light-aversive behavior (y-axis) for 4-7 Hz

clusters demonstrate relationship between brain network feature clustering and behavior on the

light/dark test, represented by the relative dark ratio= Treatment time in dark/ Vehicle time in

dark. E) The cluster mean of difference of coherency matrix for 4-7 Hz over the first change

point (i.e. the change in coherency matrices from the segment preceding changepoint to the

segment after changepoint) after injection (in clusters with at least 2 mice and ordered by the

average relative dark ratio) (see also Figure S15-18).

Figure 8. Model of Migraine Network Activity

Top: Brain network components implicated in migraine either by human fMRI or by rodent pain

studies. *Abbreviations*: ACC- anterior cingulate cortex, Amy- amygdala, BNST- bed nucleus of

the stria terminalis, Hypothal- hypothalamus, M1- motor cortex, NTS- nucleus tractus solitarius,

PAG- periaqueductal gray, PFC- prefrontal cortex, PBN- parabrachial nucleus, SS-

somatosensory cortex, TNC- trigeminal nucleus caudalis, V1- visual cortex, VTA- ventral

tegmental area. Figure created by first organizing human migraine imaging literature to identify

key brain regions involved in migraine and then organizing network connectivity by reviewing

preclinical migraine and related circuits. Finally, circuit connections were filled in between the

major hubs and network components. [3,11,21,28,33,37-39,100,101,122-147]. Map is not meant to be

comprehensive, especially between cortical regions. Also not shown are spinal inputs, which can

directly impact thalamic and hypothalamic regions, among others [25,148]. Bottom: Model of brain

network activity in peripheral CGRP-induced migraine state.

DATA AVAILABILITY

The data used in this article will be shared on reasonable request to the corresponding author.
1160
REFERENCES

1.   Steiner TJ, Stovner LJ, Jensen R, Uluduz D, Katsarava Z, Lifting The Burden: the Global Campaign
against H. Migraine remains second among the world's causes of disability, and first among young
women: findings from GBD2019. *J Headache Pain*. Dec 2 2020;21(1):137. doi:10.1186/s10194-020-
01208-0
2.   Headache Classification Committee of the International Headache Society (IHS) The
International Classification of Headache Disorders, 3rd edition. *Cephalalgia*. Jan 2018;38(1):1-211.
doi:10.1177/0333102417738202
3.   Goadsby PJ, Holland PR, Martins-Oliveira M, Hoffmann J, Schankin C, Akerman S.
Pathophysiology of Migraine: A Disorder of Sensory Processing. *Physiol Rev*. Apr 2017;97(2):553-622.
doi:10.1152/physrev.00034.2015
4.   Russo AF, Hay DL. CGRP physiology, pharmacology, and therapeutic targets: migraine and
beyond. *Physiol Rev*. Apr 1 2023;103(2):1565-1644. doi:10.1152/physrev.00059.2021
5.   Groenewald CB, Murray CB, Palermo TM. Adverse childhood experiences and chronic pain
among children and adolescents in the United States. *Pain Rep*. Sep-Oct 2020;5(5):e839.
doi:10.1097/PR9.0000000000000839
6.   Fuller-Thomson E, Baker TM, Brennenstuhl S. Investigating the association between childhood
physical abuse and migraine. *Headache*. May 2010;50(5):749-60. doi:10.1111/j.1526-4610.2010.01626.x
7.   Anda R, Tietjen G, Schulman E, Felitti V, Croft J. Adverse childhood experiences and frequent
headaches in adults. *Headache*. Oct 2010;50(9):1473-81. doi:10.1111/j.1526-4610.2010.01756.x
8.   Sikorski C, Mavromanoli AC, Manji K, Behzad D, Kreatsoulas C. Adverse Childhood Experiences
and Primary Headache Disorders: A Systematic Review, Meta-analysis, and Application of a Biological
Theory. *Neurology*. Nov 21 2023;101(21):e2151-e2161. doi:10.1212/WNL.0000000000207910
9.   Johnson M, Filali Y, Adegboyo A, et al. Mechanistic intersections between migraine and major
depressive disorder. *J Headache Pain*. Jul 9 2025;26(1):157. doi:10.1186/s10194-025-02097-x
10.   Anda RF, Felitti VJ, Bremner JD, et al. The enduring effects of abuse and related adverse
experiences in childhood. A convergence of evidence from neurobiology and epidemiology. *Eur Arch
Psychiatry Clin Neurosci*. Apr 2006;256(3):174-86. doi:10.1007/s00406-005-0624-4
11.   Jaramillo AA, Brown JA, Winder DG. Danger and distress: Parabrachial-extended amygdala
circuits. *Neuropharmacology*. Oct 15 2021;198:108757. doi:10.1016/j.neuropharm.2021.108757
12.   Schwedt TJ, Chiang CC, Chong CD, Dodick DW. Functional MRI of migraine. *Lancet Neurol*. Jan
2015;14(1):81-91. doi:10.1016/S1474-4422(14)70193-0
13.   Schwedt TJ. Multisensory integration in migraine. *Curr Opin Neurol*. Jun 2013;26(3):248-53.
doi:10.1097/WCO.0b013e328360edb1
14.   Schwedt TJ, Schlaggar BL, Mar S, et al. Atypical resting-state functional connectivity of affective
pain regions in chronic migraine. *Headache*. May 2013;53(5):737-51. doi:10.1111/head.12081
15.   Schmidt K, Schunke O, Forkmann K, Bingel U. Enhanced short-term sensitization of facial
compared with limb heat pain. *J Pain*. Aug 2015;16(8):781-90. doi:10.1016/j.jpain.2015.05.003
16.   Smith JG, Elias LA, Yilmaz Z, et al. The psychosocial and affective burden of posttraumatic
neuropathy following injuries to the trigeminal nerve. *J Orofac Pain*. Fall 2013;27(4):293-303.
doi:10.11607/jop.1056

17.   Schmidt K, Forkmann K, Sinke C, Gratz M, Bitz A, Bingel U. The differential effect of trigeminal vs.
peripheral pain stimulation on visual processing and memory encoding is influenced by pain-related
fear. *Neuroimage*. Jul 1 2016;134:386-395. doi:10.1016/j.neuroimage.2016.03.026
18.   Charles A. The pathophysiology of migraine: implications for clinical management. *Lancet*
Neurol. Feb 2018;17(2):174-182. doi:10.1016/S1474-4422(17)30435-0
19.   Baliki MN, Chialvo DR, Geha PY, et al. Chronic pain and the emotional brain: specific brain
activity associated with spontaneous fluctuations of intensity of chronic back pain. *J Neurosci*. Nov 22
2006;26(47):12165-73. doi:10.1523/JNEUROSCI.3576-06.2006
20.   Valfre W, Rainero I, Bergui M, Pinessi L. Voxel-based morphometry reveals gray matter
abnormalities in migraine. *Headache*. Jan 2008;48(1):109-17. doi:10.1111/j.1526-4610.2007.00723.x
21.   Han S, Soleiman MT, Soden ME, Zweifel LS, Palmiter RD. Elucidating an Affective Pain Circuit that
Creates a Threat Memory. *Cell*. Jul 16 2015;162(2):363-374. doi:10.1016/j.cell.2015.05.057
22.   Corder G, Ahanonu B, Grewe BF, Wang D, Schnitzer MJ, Scherrer G. An amygdalar neural
ensemble that encodes the unpleasantness of pain. *Science*. Jan 18 2019;363(6424):276-281.
doi:10.1126/science.aap8586
23.   Chong CD, Gaw N, Fu Y, Li J, Wu T, Schwedt TJ. Migraine classification using magnetic resonance
imaging resting-state functional connectivity data. *Cephalalgia*. Aug 2017;37(9):828-844.
doi:10.1177/0333102416652091
24.   Schulte LH, Menz MM, Haaker J, May A. The migraineur's brain networks: Continuous resting
state fMRI over 30 days. *Cephalalgia*. Dec 2020;40(14):1614-1621. doi:10.1177/0333102420951465
25.   Noseda R, Jakubowski M, Kainz V, Borsook D, Burstein R. Cortical projections of functionally
identified thalamic trigeminovascular neurons: implications for migraine headache and its associated
symptoms. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. Oct 5
2011;31(40):14204-17. doi:10.1523/JNEUROSCI.3285-11.2011
26.   Noseda R, Kainz V, Jakubowski M, et al. A neural mechanism for exacerbation of headache by
light. *Nat Neurosci*. Feb 2010;13(2):239-45. doi:10.1038/nn.2475
27.   Wilcox SL, Veggeberg R, Lemme J, et al. Increased Functional Activation of Limbic Brain Regions
during Negative Emotional Processing in Migraine. *Frontiers in human neuroscience*. 2016;10:366.
doi:10.3389/fnhum.2016.00366
28.   Roy DS, Zhang Y, Halassa MM, Feng G. Thalamic subnetworks as units of function. *Nat Neurosci*.
Feb 2022;25(2):140-153. doi:10.1038/s41593-021-00996-1
29.   Chen Z, Xiao L, Liu H, et al. Altered thalamo-cortical functional connectivity in patients with
vestibular migraine: a resting-state fMRI study. *Neuroradiology*. Jan 2022;64(1):119-127.
doi:10.1007/s00234-021-02777-w
30.   Sowers LP, Wang M, Rea BJ, et al. Stimulation of Posterior Thalamic Nuclei Induces Photophobic
Behavior in Mice. *Headache*. Oct 2020;60(9):1961-1981. doi:10.1111/head.13917
31.   Holland PR, Saengjaroentham C, Vila-Pueyo M. The role of the brainstem in migraine: Potential
brainstem effects of CGRP and CGRP receptor activation in animal models. *Cephalalgia*. Mar
2019;39(3):390-402. doi:10.1177/0333102418756863
32.   Uddin O, Anderson M, Smith J, Masri R, Keller A. Parabrachial complex processes dura inputs
through a direct trigeminal ganglion-to-parabrachial connection. *Neurobiol Pain*. Jan-Jul 2021;9:100060.
doi:10.1016/j.ynpai.2021.100060
33.   Palmiter RD. The Parabrachial Nucleus: CGRP Neurons Function as a General Alarm. *Trends in*
neurosciences. May 2018;41(5):280-293. doi:10.1016/j.tins.2018.03.007
34.   Vila-Pueyo M, Hoffmann J, Romero-Reyes M, Akerman S. Brain structure and function related to
headache: Brainstem structure and function in headache. *Cephalalgia*. Nov 2019;39(13):1635-1660.
doi:10.1177/0333102418784698

35.	Karsan N, Bose PR, O'Daly O, Zelaya FO, Goadsby PJ. Alterations in Functional Connectivity
During Different Phases of the Triggered Migraine Attack. *Headache*. Jul 2020;60(7):1244-1258.
doi:10.1111/head.13865
36.	Jia Z, Chen X, Tang W, Zhao D, Yu S. Atypical functional connectivity between the anterior
cingulate cortex and other brain regions in a rat model of recurrent headache. *Mol Pain*. Jan-Dec
2019;15:1744806919842483. doi:10.1177/1744806919842483
37.	Rodriguez E, Sakurai K, Xu J, et al. A craniofacial-specific monosynaptic circuit enables
heightened affective pain. *Nat Neurosci*. Dec 2017;20(12):1734-1743. doi:10.1038/s41593-017-0012-1
38.	Krout KE, Loewy AD. Parabrachial nucleus projections to midline and intralaminar thalamic
nuclei of the rat. *J Comp Neurol*. Dec 18 2000;428(3):475-94. doi:10.1002/1096-
9861(20001218)428:3<475::aid-cne6>3.0.co;2-9
39.	Brennan KC, Pietrobon D. A Systems Neuroscience Approach to Migraine. *Neuron*. Mar 7
2018;97(5):1004-1021. doi:10.1016/j.neuron.2018.01.029
40.	Versijpt J, Paemeleire K, Reuter U, MaassenVanDenBrink A. Calcitonin gene-related peptide-
targeted therapy in migraine: current role and future perspectives. *Lancet*. Mar 22
2025;405(10483):1014-1026. doi:10.1016/S0140-6736(25)00109-6
41.	Wattiez AS, Wang M, Russo AF. CGRP in Animal Models of Migraine. *Handb Exp Pharmacol*.
2019;255:85-107. doi:10.1007/164_2018_187
42.	Mason BN, Kaiser EA, Kuburas A, et al. Induction of Migraine-Like Photophobic Behavior in Mice
by Both Peripheral and Central CGRP Mechanisms. *The Journal of neuroscience : the official journal of
the Society for Neuroscience*. Jan 04 2017;37(1):204-216. doi:10.1523/JNEUROSCI.2967-16.2016
43.	Hansen JM, Hauge AW, Olesen J, Ashina M. Calcitonin gene-related peptide triggers migraine-
like attacks in patients with migraine with aura. *Cephalalgia*. Oct 2010;30(10):1179-86.
doi:10.1177/0333102410368444
44.	Asghar MS, Hansen AE, Kapijimpanga T, et al. Dilation by CGRP of middle meningeal artery and
reversal by sumatriptan in normal volunteers. *Neurology*. Oct 26 2010;75(17):1520-6.
doi:10.1212/WNL.0b013e3181f9626a
45.	Ashina M, Bendtsen L, Jensen R, Schifter S, Olesen J. Evidence for increased plasma levels of
calcitonin gene-related peptide in migraine outside of attacks. *Pain*. May 2000;86(1-2):133-8.
doi:10.1016/s0304-3959(00)00232-3
46.	Cernuda-Morollon E, Larrosa D, Ramon C, Vega J, Martinez-Camblor P, Pascual J. Interictal
increase of CGRP levels in peripheral blood as a biomarker for chronic migraine. *Neurology*. Oct 1
2013;81(14):1191-6. doi:10.1212/WNL.0b013e3182a6cb72
47.	Goadsby PJ, Edvinsson L, Ekman R. Vasoactive peptide release in the extracerebral circulation of
humans during migraine headache. *Ann Neurol*. Aug 1990;28(2):183-7. doi:10.1002/ana.410280213
48.	Edvinsson L, Haanes KA, Warfvinge K, Krause DN. CGRP as the target of new migraine therapies -
successful translation from bench to clinic. *Nat Rev Neurol*. Jun 2018;14(6):338-350.
doi:10.1038/s41582-018-0003-1
49.	Gracia-Rubio I, Moscoso-Castro M, Pozo OJ, Marcos J, Nadal R, Valverde O. Maternal separation
induces neuroinflammation and long-lasting emotional alterations in mice. *Prog Neuropsychopharmacol
Biol Psychiatry*. Feb 4 2016;65:104-17. doi:10.1016/j.pnpbp.2015.09.003
50.	Pena CJ, Smith M, Ramakrishnan A, et al. Early life stress alters transcriptomic patterning across
reward circuitry in male and female mice. *Nature communications*. Nov 8 2019;10(1):5098.
doi:10.1038/s41467-019-13085-6
51.	Kikusui T, Isaka Y, Mori Y. Early weaning deprives mouse pups of maternal care and decreases
their maternal behavior in adulthood. *Behav Brain Res*. Jul 30 2005;162(2):200-6.
doi:10.1016/j.bbr.2005.03.013

52.   Kaiser EA, Kuburas A, Recober A, Russo AF. Modulation of CGRP-induced light aversion in wild-
type mice by a 5-HT(1B/D) agonist. *The Journal of neuroscience : the official journal of the Society for*
Neuroscience. Oct 31 2012;32(44):15439-49. doi:10.1523/JNEUROSCI.3265-12.2012

53.   Chaplan SR, Bach FW, Pogrel JW, Chung JM, Yaksh TL. Quantitative assessment of tactile
allodynia in the rat paw. *J Neurosci Methods*. Jul 1994;53(1):55-63. doi:10.1016/0165-0270(94)90144-9

54.   Dixon WJ. Efficient analysis of experimental observations. *Annu Rev Pharmacol Toxicol*.
1980;20:441-62. doi:10.1146/annurev.pa.20.040180.002301

55.   Bates D, Maechler, M., Bolker, B., Walker, S. Fitting Linear Mixed-Effects Models Using lme4.
Journal of Statistical Software. 2015;67((1)):1-48. doi:doi:10.18637/jss.v067.i01.

56.   Kuznetsova A, Brockhoff, P.B., Christensen, R.H.B. . lmerTest Package: Tests in Linear Mixed
Effects Models. *Journal of Statistical Software*. 2017;82(13):1-26. doi:doi: 10.18637/jss.v082.i13

57.   Kumar S, Hultman R, Hughes D, Michel N, Katz BM, Dzirasa K. Prefrontal cortex reactivity
underlies trait vulnerability to chronic social defeat stress. *Nature communications*. 2014;5:4537.
doi:10.1038/ncomms5537

58.   Tibshirani R. Regression Shrinkage and Selection via the Lasso. *J R Stat Soc Ser B*.
1996;58(1):267–288.

59.   Tibshirani R SM, Rosset S, Zhu J, Knight K. . Sparsity and smoothness via the fused lasso. *J R Stat*
Soc Ser B (Statistical Methodol). 2005;67(1):91–108. doi:doi:https://doi.org/10.1111/j.1467-
9868.2005.00490.

60.   Tibshirani RJ TJ. THE SOLUTION PATH OF THE GENERALIZED LASSO. *Ann Stat*. 2011;39(3):1335–
1371.

61.   Hastie T, & Tibshirani, R. . Generalized Additive Models. *Statistical Science*. 1986;1((3)):297–310.

62.   Scheipl FaJG, J. tf: S3 Classes and Methods for Tidy Functional Data. 2024;

63.   Wickham H. *ggplot2: Elegant Graphics for Data Analysis.* Springer-Verlag; 2016.

64.   Scheipl F, Goldsmith, J. and Wrobel, J. . tidyfun: Tools for Tidy Functional Data. 2024.
https://github.com/tidyfun/tidyfun

65.   Hultman R, Ulrich K, Sachs BD, et al. Brain-wide Electrical Spatiotemporal Dynamics Encode
Depression Vulnerability. *Cell*. Mar 22 2018;173(1):166-180 e14. doi:10.1016/j.cell.2018.02.012

66.   Rea BJ, Wattiez AS, Waite JS, et al. Peripherally Administered CGRP Induces Spontaneous Pain in
Mice: Implications for Migraine. *Pain*. Jul 9 2018;doi:10.1097/j.pain.0000000000001337

67.   Wood SN. Fast Stable Restricted Maximum Likelihood and Marginal Likelihood Estimation of
Semiparametric Generalized Linear Models. *Journal of the Royal Statistical Society Series B: Statistical*
Methodology. 2011;73(1):3-36. doi:10.1111/j.1467-9868.2010.00749.x

68.   Zhang X, Chan, K. Spectral Change Point Estimation for High Dimensional Time Series by Sparse
Tensor Decomposition. 2024;2305.10656v2Located at: arxiv.

69.   Brillinger DR. *Time Series: Data Analysis and Theory*. Society for Industrial and Applied
Mathematics; 2001.

70.   Zhai ZW, Yip SW, Lacadie CM, Sinha R, Mayes LC, Potenza MN. Childhood trauma moderates
inhibitory control and anterior cingulate cortex activation during stress. *Neuroimage*. Jan 15
2019;185:111-118. doi:10.1016/j.neuroimage.2018.10.049

71.   Holschneider DP, Guo Y, Mayer EA, Wang Z. Early life stress elicits visceral hyperalgesia and
functional reorganization of pain circuits in adult rats. *Neurobiol Stress*. Jun 1 2016;3:8-22.
doi:10.1016/j.ynstr.2015.12.003

72.   Raoof M, Amanpour S, Roghani A, et al. The effects of neonatal maternal deprivation and
chronic unpredictable stresses on migraine-like behaviors in adult rats. *Neurosci Lett*. Feb 16
2022;772:136444. doi:10.1016/j.neulet.2022.136444

73.   Eller OC, Yang X, Fuentes IM, et al. Voluntary Wheel Running Partially Attenuates Early Life
Stress-Induced Neuroimmune Measures in the Dura and Evoked Migraine-Like Behaviors in Female
Mice. *Front Physiol*. 2021;12:665732. doi:10.3389/fphys.2021.665732
74.   Minguez-Olaondo A, Quintas S, Morollon Sanchez-Mateos N, et al. Cutaneous Allodynia in
Migraine: A Narrative Review. *Front Neurol*. 2021;12:831035. doi:10.3389/fneur.2021.831035
75.   Munjal S, Singh P, Reed ML, et al. Most Bothersome Symptom in Persons With Migraine: Results
From the Migraine in America Symptoms and Treatment (MAST) Study. *Headache*. Feb 2020;60(2):416-
429. doi:10.1111/head.13708
76.   Wang M, Mason BN, Sowers LP, Kuburas A, Rea BJ, Russo AF. Investigating Migraine-Like
Behavior Using Light Aversion in Mice. *J Vis Exp*. Aug 11 2021;(174)doi:10.3791/62839
77.   Hultman R, Mague, S.D., Li, Q., Katz, B.M., Michel, N., Lin, L., Wang, J., David, L.K., Blount,
Chandy, R., Carlson, D., Ulrich, K., Carin, L., Dunson, Kumar, S., Deisseroth, K. Moore, S.D., Dzirasa, K.
Prefrontal Cortex-Mediated Slow-Evolving Limbic Dynamics Drives Stress-Induced Emotional Pathology.
Neuron. 2016;In Press
78.   Adhikari A, Topiwala MA, Gordon JA. Synchronized activity between the ventral hippocampus
and the medial prefrontal cortex during anxiety. *Neuron*. Jan 28 2010;65(2):257-69.
doi:10.1016/j.neuron.2009.12.002
79.   Hultman R, Mague SD, Li Q, et al. Dysregulation of Prefrontal Cortex-Mediated Slow-Evolving
Limbic Dynamics Drives Stress-Induced Emotional Pathology. *Neuron*. Jul 20 2016;91(2):439-52.
doi:10.1016/j.neuron.2016.05.038
80.   Perge JA, Zhang S, Malik WQ, et al. Reliability of directional information in unsorted spikes and
local field potentials recorded in human motor cortex. *J Neural Eng*. Aug 2014;11(4):046007.
doi:10.1088/1741-2560/11/4/046007
81.   Schaich Borg J, Srivastava S, Lin L, et al. Rat intersubjective decisions are encoded by frequency-
specific oscillatory contexts. *Brain Behav*. Jun 2017;7(6):e00710. doi:10.1002/brb3.710
82.   Nolte G, Ziehe A, Nikulin VV, et al. Robustly estimating the flow direction of information in
complex physical systems. *Phys Rev Lett*. Jun 13 2008;100(23):234101.
doi:10.1103/PhysRevLett.100.234101
83.   Dzirasa K, Fuentes R, Kumar S, Potes JM, Nicolelis MA. Chronic in vivo multi-circuit
neurophysiological recordings in mice. Research Support, N.I.H., Extramural

Research Support, Non-U.S. Gov't. *Journal of neuroscience methods*. Jan 30 2011;195(1):36-46.
doi:10.1016/j.jneumeth.2010.11.014
84.   Adhikari A, Sigurdsson T, Topiwala MA, Gordon JA. Cross-correlation of instantaneous
amplitudes of field potential oscillations: a straightforward method to estimate the directionality and lag
between brain areas. *J Neurosci Methods*. Aug 30 2010;191(2):191-200.
doi:10.1016/j.jneumeth.2010.06.019
85.   Buzsàki G, Eidelberg E. Phase relations of hippocampal projection cells and interneurons to theta
activity in the anesthetized rat. *Brain research*. May 1983;266(2):334-9. doi:10.1016/0006-
8993(83)90665-0
86.   GREEN JD, ARDUINI AA. Hippocampal electrical activity in arousal. *J Neurophysiol*. Nov
1954;17(6):533-57. doi:10.1152/jn.1954.17.6.533
87.   Gordon JA, Lacefield CO, Kentros CG, Hen R. State-dependent alterations in hippocampal
oscillations in serotonin 1A receptor-deficient mice. *The Journal of neuroscience : the official journal of
the Society for Neuroscience*. Jul 2005;25(28):6509-19. doi:10.1523/jneurosci.1211-05.2005
88.   Winson J, Abzug C. Neuronal transmission through hippocampal pathways dependent on
behavior. *J Neurophysiol*. May 1978;41(3):716-32. doi:10.1152/jn.1978.41.3.716

89.   Padilla-Coreano N, Canetta S, Mikofsky RM, et al. Hippocampal-Prefrontal Theta Transmission
Regulates Avoidance Behavior. *Neuron*. 11 2019;104(3):601-610.e4. doi:10.1016/j.neuron.2019.08.006
90.   Adhikari A, Topiwala MA, Gordon JA. Synchronized Activity between the Ventral Hippocampus
and the Medial Prefrontal Cortex during Anxiety. *Neuron*. 28 January 2010 2010;65(2):257-269.
91.   Cameron C, Kelly S, Hsieh SC, et al. Triptans in the Acute Treatment of Migraine: A Systematic
Review and Network Meta-Analysis. *Headache*. Jul-Aug 2015;55 Suppl 4:221-35.
doi:10.1111/head.12601
92.   Holland PR, Sureda-Gibert P, Vila-Pueyo M. Rapid uptake of sumatriptan into the brain: An
ongoing question of blood-brain barrier permeability. *Cephalalgia*. Apr 2020;40(4):327-329.
doi:10.1177/0333102420905131
93.   Tfelt-Hansen PC. Does sumatriptan cross the blood-brain barrier in animals and man? *J
Headache Pain*. Feb 2010;11(1):5-12. doi:10.1007/s10194-009-0170-y
94.   Becerra L, Bishop J, Barmettler G, et al. Triptans disrupt brain networks and promote stress-
induced CSD-like responses in cortical and subcortical areas. *J Neurophysiol*. Jan 1 2016;115(1):208-17.
doi:10.1152/jn.00632.2015
95.   Buse DC, Reed ML, Fanning KM, et al. Comorbid and co-occurring conditions in migraine and
associated risk of increasing headache pain intensity and headache frequency: results of the migraine in
America symptoms and treatment (MAST) study. *J Headache Pain*. Mar 2 2020;21(1):23.
doi:10.1186/s10194-020-1084-y
96.   Maleki N, Becerra L, Borsook D. Migraine: maladaptive brain responses to stress. *Headache*. Oct
2012;52 Suppl 2(Suppl 2):102-6. doi:10.1111/j.1526-4610.2012.02241.x
97.   Hammond NG, Colman I. The Role of Positive Health Behaviors in the Relationship Between
Early Life Stress and Migraine. *Headache*. Jun 2020;60(6):1111-1123. doi:10.1111/head.13808
98.   Avona A, Mason BN, Lackovic J, et al. Repetitive stress in mice causes migraine-like behaviors
and calcitonin gene-related peptide-dependent hyperalgesic priming to a migraine trigger. *Pain*. Nov
2020;161(11):2539-2550. doi:10.1097/j.pain.0000000000001953
99.   Marciszewski KK, Meylakh N, Di Pietro F, et al. Changes in Brainstem Pain Modulation Circuitry
Function over the Migraine Cycle. *The Journal of neuroscience : the official journal of the Society for
Neuroscience*. Dec 5 2018;38(49):10479-10488. doi:10.1523/JNEUROSCI.1088-18.2018
100.   Messina R, Gollion C, Christensen RH, Amin FM. Functional MRI in migraine. *Curr Opin Neurol*.
Jun 1 2022;35(3):328-335. doi:10.1097/WCO.0000000000001060
101.   Iyengar S, Ossipov MH, Johnson KW. The role of calcitonin gene-related peptide in peripheral
and central pain mechanisms including migraine. *Pain*. Apr 2017;158(4):543-559.
doi:10.1097/j.pain.0000000000000831
102.   Iyengar S, Johnson KW, Ossipov MH, Aurora SK. CGRP and the Trigeminal System in Migraine.
Headache. May 2019;59(5):659-681. doi:10.1111/head.13529
103.   Han JS, Li W, Neugebauer V. Critical role of calcitonin gene-related peptide 1 receptors in the
amygdala in synaptic plasticity and pain behavior. *The Journal of neuroscience : the official journal of the
Society for Neuroscience*. Nov 16 2005;25(46):10717-28. doi:10.1523/JNEUROSCI.4112-05.2005
104.   Eftekhari S, Salvatore CA, Calamari A, Kane SA, Tajti J, Edvinsson L. Differential distribution of
calcitonin gene-related peptide and its receptor components in the human trigeminal ganglion.
Neuroscience. Aug 25 2010;169(2):683-96. doi:10.1016/j.neuroscience.2010.05.016
105.   Guo S, Vollesen ALH, Olesen J, Ashina M. Premonitory and nonheadache symptoms induced by
CGRP and PACAP38 in patients with migraine. *Pain*. Dec 2016;157(12):2773-2781.
doi:10.1097/j.pain.0000000000000702
106.   Schulz E, Mayr A, Jahn P, Stankewitz A. Brain connectivity in individuals with migraine resets
during the headache phase: a whole-brain connectivity study. *Brain Commun*. 2025;7(1):fcaf045.
doi:10.1093/braincomms/fcaf045

107. Puledda F, Vigano A, Sebastianelli G, et al. Electrophysiological findings in migraine may reflect
abnormal synaptic plasticity mechanisms: A narrative review. *Cephalalgia*. Aug
2023;43(8):3331024231195780. doi:10.1177/03331024231195780
108. Oh SW, Harris JA, Ng L, et al. A mesoscale connectome of the mouse brain. *Nature*. Apr 10
2014;508(7495):207-14. doi:10.1038/nature13186
109. Liu RH, Zhang M, Xue M, et al. Inhibiting neuronal AC1 for treating anxiety and headache in the
animal model of migraine. *iScience*. Jun 16 2023;26(6):106790. doi:10.1016/j.isci.2023.106790
110. Schulte LH, May A. The migraine generator revisited: continuous scanning of the migraine cycle
over 30 days and three spontaneous attacks. *Brain*. Jul 2016;139(Pt 7):1987-93.
doi:10.1093/brain/aww097
111. Valentinova K, Acuna MA, Ntamati NR, Nevian NE, Nevian T. An amygdala-to-cingulate cortex
circuit for conflicting choices in chronic pain. *Cell Rep*. Oct 31 2023;42(10):113125.
doi:10.1016/j.celrep.2023.113125
112. Veinante P, Yalcin I, Barrot M. The amygdala between sensation and affect: a role in pain. *J Mol
Psychiatry*. 2013;1(1):9. doi:10.1186/2049-9256-1-9
113. Russo AF. Calcitonin gene-related peptide (CGRP): a new target for migraine. *Annu Rev
Pharmacol Toxicol*. 2015;55:533-52. doi:10.1146/annurev-pharmtox-010814-124701
114. Brar Y, Hosseini SA, Saadabadi A. Sumatriptan. *StatPearls*. 2024.
115. Durham PL, Russo AF. Regulation of calcitonin gene-related peptide secretion by a serotonergic
antimigraine drug. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. May
1 1999;19(9):3423-9. doi:10.1523/JNEUROSCI.19-09-03423.1999
116. Benemei S, Cortese F, Labastida-Ramirez A, et al. Triptans and CGRP blockade - impact on the
cranial vasculature. *J Headache Pain*. Oct 10 2017;18(1):103. doi:10.1186/s10194-017-0811-5
117. Levy D, Jakubowski M, Burstein R. Disruption of communication between peripheral and central
trigeminovascular neurons mediates the antimigraine action of 5HT 1B/1D receptor agonists.
Proceedings of the National Academy of Sciences of the United States of America. Mar 23
2004;101(12):4274-9. doi:10.1073/pnas.0306147101
118. Muzzi M, Zecchi R, Ranieri G, et al. Ultra-rapid brain uptake of subcutaneous sumatriptan in the
rat: Implication for cluster headache treatment. *Cephalalgia*. Apr 2020;40(4):330-336.
doi:10.1177/0333102419896370
119. Asghar MS, Becerra L, Larsson HB, Borsook D, Ashina M. Calcitonin Gene-Related Peptide
Modulates Heat Nociception in the Human Brain - An fMRI Study in Healthy Volunteers. *PloS one*.
2016;11(3):e0150334. doi:10.1371/journal.pone.0150334
120. Zhang X, Li X, Steffens DC, Guo H, Wang L. Dynamic changes in thalamic connectivity following
stress and its association with future depression severity. *Brain Behav*. Dec 2019;9(12):e01445.
doi:10.1002/brb3.1445
121. Carlson D, David LK, Gallagher NM, et al. Dynamically Timed Stimulation of Corticolimbic
Circuitry Activates a Stress-Compensatory Pathway. *Biological psychiatry*. Dec 15 2017;82(12):904-913.
doi:10.1016/j.biopsych.2017.06.008
122. Messina R, Rocca MA, Goadsby PJ, Filippi M. Insights into migraine attacks from neuroimaging.
Lancet Neurol. Sep 2023;22(9):834-846. doi:10.1016/S1474-4422(23)00152-7
123. Flook EA, Feola B, Avery SN, et al. BNST-insula structural connectivity in humans. *Neuroimage*.
Apr 15 2020;210:116555. doi:10.1016/j.neuroimage.2020.116555
124. Krout KE, Loewy AD. Periaqueductal gray matter projections to midline and intralaminar
thalamic nuclei of the rat. *J Comp Neurol*. Aug 14 2000;424(1):111-41. doi:10.1002/1096-
9861(20000814)424:1<111::aid-cne9>3.0.co;2-3
125. Torres-Rodriguez JM, Wilson TD, Singh S, et al. The parabrachial to central amygdala pathway is
critical to injury-induced pain sensitization in mice. *Neuropsychopharmacology : official publication of*

the American College of Neuropsychopharmacology. Feb 2024;49(3):508-520. doi:10.1038/s41386-023-
01673-6
126.   Pietrobon D, Moskowitz MA. Pathophysiology of migraine. *Annu Rev Physiol.* 2013;75:365-91.
doi:10.1146/annurev-physiol-030212-183717
127.   Wu Q, Clark MS, Palmiter RD. Deciphering a neuronal circuit that mediates appetite. *Nature.*
Mar 14 2012;483(7391):594-7. doi:10.1038/nature10899
128.   Cheng W, Gonzalez I, Pan W, et al. Calcitonin Receptor Neurons in the Mouse Nucleus Tractus
Solitarius Control Energy Balance via the Non-aversive Suppression of Feeding. *Cell Metab.* Feb 4
2020;31(2):301-312 e5. doi:10.1016/j.cmet.2019.12.012
129.   Xu WJ, Barisano G, Phung D, et al. Structural MRI in Migraine: A Review of Migraine Vascular and
Structural Changes in Brain Parenchyma. *J Cent Nerv Syst Dis.* 2023;15:11795735231167868.
doi:10.1177/11795735231167868
130.   Robert C, Bourgeais L, Arreto CD, et al. Paraventricular hypothalamic regulation of
trigeminovascular mechanisms involved in headaches. *The Journal of neuroscience : the official journal*
of the Society for Neuroscience. May 15 2013;33(20):8827-40. doi:10.1523/JNEUROSCI.0439-13.2013
131.   Yu W, Pati D, Pina MM, et al. Periaqueductal gray/dorsal raphe dopamine neurons contribute to
sex differences in pain-related behaviors. *Neuron.* Apr 21 2021;109(8):1365-1380 e5.
doi:10.1016/j.neuron.2021.03.001
132.   Rikhye RV, Gilra A, Halassa MM. Thalamic regulation of switching between cortical
representations enables cognitive flexibility. *Nat Neurosci.* 12 2018;21(12):1753-1763.
doi:10.1038/s41593-018-0269-z
133.   Wolff M, Halassa MM. The mediodorsal thalamus in executive control. *Neuron.* Mar 20
2024;112(6):893-908. doi:10.1016/j.neuron.2024.01.002
134.   Ong WY, Stohler CS, Herr DR. Role of the Prefrontal Cortex in Pain Processing. *Mol Neurobiol.*
Feb 2019;56(2):1137-1166. doi:10.1007/s12035-018-1130-9
135.   Wang M, Tutt JO, Dorricott NO, Parker KL, Russo AF, Sowers LP. Involvement of the cerebellum
in migraine. *Front Syst Neurosci.* 2022;16:984406. doi:10.3389/fnsys.2022.984406
136.   Waung MW, Margolis EB, Charbit AR, Fields HL. A Midbrain Circuit that Mediates Headache
Aversiveness in Rats. *Cell Rep.* Sep 10 2019;28(11):2739-2747 e4. doi:10.1016/j.celrep.2019.08.009
137.   Li JN, Sheets PL. The central amygdala to periaqueductal gray pathway comprises intrinsically
distinct neurons differentially affected in a model of inflammatory pain. *The Journal of physiology.* Dec
2018;596(24):6289-6305. doi:10.1113/JP276935
138.   Noseda R, Burstein R. Migraine pathophysiology: anatomy of the trigeminovascular pathway
and associated neurological symptoms, CSD, sensitization and modulation of pain. *Pain.* Dec 2013;154
Suppl 1doi:10.1016/j.pain.2013.07.021
139.   Todd AJ. Neuronal circuitry for pain processing in the dorsal horn. *Nature reviews Neuroscience.*
Dec 2010;11(12):823-36. doi:10.1038/nrn2947
140.   Gauriau C, Bernard JF. Pain pathways and parabrachial circuits in the rat. *Exp Physiol.* Mar
2002;87(2):251-8. doi:10.1113/eph8702357
141.   Boucher MN, Aktar M, Braas KM, May V, Hammack SE. Activation of Lateral Parabrachial
Nucleus (LPBn) PACAP-Expressing Projection Neurons to the Bed Nucleus of the Stria Terminalis (BNST)
Enhances Anxiety-like Behavior. *J Mol Neurosci.* Mar 2022;72(3):451-458. doi:10.1007/s12031-021-
01946-z
142.   Chiang MC, Nguyen EK, Canto-Bustos M, Papale AE, Oswald AM, Ross SE. Divergent Neural
Pathways Emanating from the Lateral Parabrachial Nucleus Mediate Distinct Components of the Pain
Response. *Neuron.* Jun 17 2020;106(6):927-939 e5. doi:10.1016/j.neuron.2020.03.014
143.   Wang F, Chen Y, Lin Y, et al. A parabrachial to hypothalamic pathway mediates defensive
behavior. *Elife.* Mar 17 2023;12doi:10.7554/eLife.85450

144.   Ruggiero DA, Anwar S, Kim J, Glickstein SB. Visceral afferent pathways to the thalamus and
olfactory tubercle: behavioral implications. *Brain research*. Jul 13 1998;799(1):159-71.
doi:10.1016/s0006-8993(98)00442-9
145.   Janak PH, Tye KM. From circuits to behaviour in the amygdala. *Nature*. Jan 15
2015;517(7534):284-92. doi:10.1038/nature14188
146.   Gehrlach DA, Weiand C, Gaitanos TN, et al. A whole-brain connectivity map of mouse insular
cortex. *Elife*. Sep 17 2020;9doi:10.7554/eLife.55585
147.   Guillamon-Vivancos T, Anibal-Martinez M, Puche-Aroca L, Martini FJ, Lopez-Bendito G. Sensory
modality-specific wiring of thalamocortical circuits. *Nature reviews Neuroscience*. Jul 30
2025;doi:10.1038/s41583-025-00945-y
148.   Kang SJ, Liu S, Kim JH, et al. Thalamic CGRP neurons define a spinothalamic pathway for affective
pain. *Proceedings of the National Academy of Sciences of the United States of America*. Jul 15
2025;122(28):e2505889122. doi:10.1073/pnas.2505889122

1545

Figure 2

A. Experimental Day Procedure

30 min Baseline → IP injection → 2 hours Post-injection

CGRP Experiments

Day 1 +PBS
Day 2 +sumatriptan
Day 3 +CGRP & sumatriptan

Day 1 +PBS
Day 2 +CGRP
Day 3 +PBS

Pharmacological Scheme

+ CGRP →
+ Sumatriptan →
+ CGRP & Sumatriptan →

B. Sumatriptan + CGRP vs CGRP only Power

4-7 Hz
8-13 Hz

Acc AMY LPBN MD_thal Po RPBN VPM

-log10(p)

C. AMY (8-13 Hz)

CGRP
CGRP + Sumatriptan
Sumatriptan
Vehicle

Mean log-power

time (min)

CGRP- (CGRP + Suma)

time (min)

A.

Brain systems/functionality

- ● Trigeminal (pain/nociception)
- ● Sensory gain/photophobia
- ● Structural alterations
- ● Cognition
- ● Affect
- ● Homeostasis
- ● Cortical Spreading Depression

PFC ACC SS V1 M1 Auditory Cortex Piriform Cortex

Insular cortex Thalamus TNC

Hippocampus Amygdala PBN

Hypothal

BNST PAG VTA

Cerebellum

Trigeminal Nerve

Retinal ganglion cells

B.

CGRP

Sumatriptan + CGRP

ELS + CGRP

control ELS

Power

Coherence

- PFC
- ACC
- SS
- V1
- Insular Cortex
- Amy
- BNST
- NTS
- PBN
- PAG
- Thalamus