UV-Induced DNA Repair Mechanisms and Their Effects on Mutagenesis and Culturability in *Escherichia coli*

Sreyashi Ghosh, and Mehmet A. Orman*

William A. Brookshire Department of Chemical and Biomolecular Engineering, University of Houston, Houston, TX, USA

*Correspondence to: S222 Engineering Bldg 1, 4226 Martin Luther King Boulevard, Houston, TX 77204-4004, Phone: 713-743-6785, Email: morman@central.uh.edu

Abstract

Mutagenic processes drive evolutionary progress, with ultraviolet (UV) radiation significantly affecting evolution. Despite extensive research on SOS response-mediated mutagenesis, UV-induced repair mechanisms remain complex, and their effects on cell survival and mutagenesis are not fully understood. We previously observed a near-perfect correlation between RecA-mediated SOS response and mutation levels in *Escherichia coli* following UV treatment. However, prolonged UV exposure caused transient non-culturability and impaired SOS-mediated mutagenesis. Using fluorescent reporters, flow cytometry, promoter-reporter assays, single-gene deletions, knockouts, and clonogenic assays, we found that excessive UV exposure disrupts cellular translation, reducing SOS gene expression, albeit with minimal impact on membrane permeability or reactive oxygen species levels. While our findings underline the abundance of repair mechanisms in *E. coli* cells, enabling them to compensate when specific genes are disrupted, they also highlighted the differential impacts of gene deletions on mutagenesis versus culturability, leading to three major outcomes: (i) Disruption of proteins involved in DNA polymerase for trans-lesion synthesis (UmuC and UmuD) or Holliday junction resolution (RuvC) results in significantly decreased mutagenesis levels while maintaining a transient non-culturability pattern after UV exposure. (ii) Disruption of proteins involved in homologous recombination (RecA and RecB) and nucleotide excision repair (UvrA) leads to both significantly reduced mutagenesis and a more severe transient non-culturability pattern after UV exposure, making these cells more sensitive to UV. (iii) Disruption of DNA Helicase II (UvrD), which functions in mismatch repair, does not affect mutagenesis levels from UV radiation but results in a very pronounced transient non-culturability pattern following UV exposure. Overall, our results further advance our understanding of bacterial adaptation mechanisms and the role of DNA repair pathways in shaping mutagenesis.

Author Summary

Ultraviolet (UV) radiation has been a significant force in driving genetic variation and adaptation throughout billions of years of evolution. By directly damaging DNA and triggering repair mechanisms, UV radiation is a powerful tool for studying mutagenesis. This study aims to provide new insights into the complex processes behind bacterial mutagenesis, a critical topic in microbiology and public health. Although the connection between the SOS response and DNA repair in UV-treated cells is well known, two key phenomena — UV-induced bacterial cell dormancy and mutagenesis — remain poorly understood. Our findings reveal a highly unusual,

42 SOS response-dependent transient non-culturability in *Escherichia coli* cells following prolonged
43 UV exposure. However, the downstream mechanisms of this phenomenon and its links to
44 mutagenesis remain unclear. This study seeks to thoroughly investigate these phenomena, offering
45 new insights into their underlying molecular processes.

Introduction

Mutagenic processes are key to evolutionary progress [1–3], as they generate genetic diversity. This
is essential for organisms to adapt to and endure in dynamic settings that are continuously
changing. Understanding the mechanisms underlying mutagenesis helps us better comprehend the
evolutionary forces shaping organisms [4,5]. This knowledge also aids in developing more potent and
effective medical treatments to combat many diseases, given that pathogenic microorganisms and
tumorigenic cells develop resistance through mutagenesis[6–9].

Ultraviolet (UV) radiation has continually shaped the trajectory of evolution over billions of years,
from contributing to the origins of life to driving genetic variation and adaptation[10,11]. This is
because UV radiation directly damages DNA and induces certain repair mechanisms that are
highly mutagenic[12–15]. The UV light spectrum, ranging from 100 to 400 nanometers (nm), can be
categorized into four groups: UV-A (315-400 nm), UV-B (280-315 nm), UV-C (100-280 nm), and
vacuum-UV (100-200 nm)[16]. UV-C is more readily absorbed by nucleic acids compared to UV-B
and UV-A[17]. When bacteria are exposed to UV radiation, [6,7] including UV-C, UV-B, and UV-A,
it triggers the formation of pyrimidine dimers and other DNA lesions that disrupt the normal
structure and function of DNA molecules[18,19]. These UV lesions are removed by various effective
bacterial systems, such as the photoreactivation repair system, (i.e. dimer monomerization
under visible   light by photolyase enzymes[20]),   base excision   repair   (BER)   (removal   of
cyclobutane-thymine dimers using DNA glycosylases and AP endonucleases)[21], and the primary
pathway for removal of bulky DNA lesions, i.e. the nucleotide excision repair (NER) (for rapid
repair of photo-lesions using the *uvr* genes, *polA* and ligase enzyme) [22]. However, an excessive
amount of dimers that cannot be removed by the repair systems can accumulate and interfere with
the cellular replication process, resulting in the formation of "secondary" lesions in the form of
ssDNA fragments[23]. Therefore, to deal with this severe DNA damage, a post-replication damage
repair mechanism known as the bacterial SOS response pathway is induced, preventing premature
cell division [12,24,25] and providing cells with sufficient time to repair the damaged DNA[14,26–28]. This
SOS response in bacteria is controlled by the multifunctional RecA protein, a key component that
cleaves the transcriptional repressor LexA[29–31]. This cleavage initiates the expression of more than
40 SOS genes. RecA, involved in the recombination repair of ssDNA gaps[32], helps provide an
error-free damage removal route. However, excessive DNA damage leads to the induction of the
mutagenic phase of the SOS response, mediated by bacterial DNA polymerase enzymes, including
Pol V [33,34], which is encoded by the *umuD* and *umuC* genes [35]. Pol V can bypass template lesions
during DNA replication through a process known as translesion DNA synthesis (TLS) [36]. Two
additional TLS DNA polymerases, Pol II (encoded by *polB*) and Pol IV (encoded by *dinB*),
facilitate replication past blocking lesions, albeit at the cost of introducing mutations. Another
DNA damage repair strategy is the mismatch repair (MMR) pathway [37], which recognizes base-
base mismatches and insertion/deletion loops, including those introduced by DNA polymerases
and as part of TLS during the SOS response. However, with increased mutagenic stress, the
efficiency of MMR decreases, as observed in the case of fluoroquinolone treatment [38].

UV radiation can serve as a valuable experimental tool for investigating mutagenesis because it
induces a wide range of mutations without an apparent sequence preference[39,40]. Additionally, UV

radiation is of great interest to scientists for its applications in biotechnology, particularly in
directed evolutionary strategies to engineer proteins or organisms[41,42] . Moreover, there is growing
interest in using UV radiation as a disinfectant, including for sterilizing air, equipment, and
surfaces; reducing the transmission of airborne diseases; and even treating wound infections[43,44].
Despite their increased applications in clinical settings, we believe that two phenomena— UV-
induced bacterial cell dormancy and mutagenesis— need to be fully understood. Bacterial cells
are highly heterogeneous, and they can enter a growth-arrested state through stochastic
mechanisms or environmental factors[45], including UV treatment, which can make these cells
highly tolerant. Also, the SOS-response-mediated cell dormancy not only promotes cell survival
but also sustains mutagenic processes, as dormant cells may still harbor DNA damage[46,47].
Additionally, UV radiation itself is highly mutagenic; it can accelerate the emergence of more
resilient mutants, which could pose a serious threat to public health.

UV-induced repair mechanisms are highly complex as UV can damage many cellular components,
including DNA, RNA, lipids, and proteins[48,49]. Although the role of the SOS response in
mutagenesis has been extensively studied (refer to the following review articles[12,26,28] for more
details), our understanding of the downstream repair mechanisms and their contributions to cell
survival, culturability, and mutagenesis remains limited. In our recent study[50], we observed a
significant correlation between the RecA-mediated SOS response and mutation levels.
Additionally, prolonged UV exposure was found to impair SOS-mediated mutagenesis and induce
a transient non-culturability state [50], which may represent a potential survival strategy for cells to
evade UV-induced mutagenesis. However, the underlying mechanisms remain unclear. Our
current study aims to investigate these complex mechanisms further, providing new insights into
the molecular processes underlying mutagenesis and dormancy.

Results and Discussion

The transient unculturability is largely driven by the global regulator RecA, crucial for the SOS response and recovery

To study UV-mediated mutagenesis and the SOS response, we used our previously established
methodology[50]. We cultured *E. coli* MG1655 wild-type cells in 2 mL Lysogeny Broth (LB)
medium in test tubes until they reached the mid-exponential phase with an optical density (OD_{600})
of ~0.5. Subsequently, the cells were transferred to petri dishes, forming a thin film of cultures that
increased the surface area for UV exposure. This film was then subjected to UV-B light (302 nm
thin-line transilluminator, UV intensity ~2000 $\mu W/cm^2$, UVP ChemStudio, Analytik Jena) for
various durations: 2, 4, 8, 16, 24, and 32 minutes (min). Following exposure, the cells were
promptly transferred back to test tubes and cultured in a shaker for 24-hour (h) recovery (refer to
the Materials and Methods section for details). Cultures that did not receive UV-B treatment served
as controls. During recovery, colony-forming units (CFU) were quantified at the indicated time
points for each condition, as shown in **Fig. 1a**. At the beginning of recovery (t~0), CFU levels
decreased in cultures exposed to longer UV-B treatments. A 4-8 min exposure caused a 10-fold
reduction, while 16 min led to a ~100-fold drop compared to the control. Exposure times of 24 and
32 min resulted in a ~10,000-fold reduction (**Fig. 1a**). After 15 minutes, CFU levels notably
increased in cultures with 24 and 32 min of UV-B exposure, likely due to mechanisms making
cells temporarily unculturable. This drastic increase in CFU levels is not due to cell division, given
that the *E. coli* doubling time is about 20-25 minutes, which is too short for significant growth
within that period. By the 24-hour mark, all conditions showed similar CFU levels (**Fig. 1a**).

To measure mutagenesis, we quantified rifampicin (RIF)-resistant colonies after 24 hours of
recovery. Cultures were plated on LB agar with 500 µg/ml RIF, and colony counts were reported
per 10^8 cells (**Fig. 1b**). Although this method does not directly measure mutation levels, it is widely
used [40,51–53] due to the strong correlation between mutagenesis and resistant colony formation [54,55].
We chose a 24-hour recovery period to ensure accurate colony counts, consistent with previous
studies[51,56,57], as mutation levels generally plateau by this time[50]. Our results showed that
mutagenesis increased with UV-B exposure, peaking at 16 minutes, but longer exposures (24-32
min) significantly reduced RIF-resistant colonies (**Fig. 1b**). This pattern, consistent with our
previous findings[50], suggests a link between UV-B exposure, mutagenesis, and temporary
unculturability.

In our previous study, we found that excessive UV-B treatment (24 and 32 minutes) impaired the
SOS response, including the expression level of the RecA protein[50]. Therefore, the observed
transient unculturability is likely more complex and may involve various mechanisms, given that
the SOS response is intricate, involving multiple proteins, with RecA serving as the global
regulator. We hypothesized that if the observed transient unculturability is linked to the impaired
SOS response, then genetic perturbation of this response through *recA* deletion would further
decrease cell culturability. Our results verified that the knockout strain of *recA* exhibited a drastic
reduction in culturability immediately after UV treatment across all conditions (**Fig. 1c**). Initially,
CFU levels in the Δ*recA* strain were below the limit of detection (1 CFU) for most UV treatment
conditions; however, this was transient, as we observed a rapid increase in CFU levels (from 1 to
10^7) within 15 minutes of recovery (**Fig. 1c**). This finding is highly interesting and, to our
knowledge, has not been previously reported. As we have already shown the deletion *recA*
eliminates the mutagenesis following UV treatment[50], the impairment of the RecA-mediated SOS
response during prolonged UV exposures[50] should significantly affect downstream processes.
However, it remains unclear how these downstream processes impact cell culturability and
mutagenesis; thus, we aim to investigate these aspects in the subsequent sections of our study.

Genes significantly expressed during the SOS response have a moderate or negligible effect on
cell culturability and mutagenesis

To identify the downstream mechanisms, we screened a subset of the *E. coli* promoter library
containing 1900 promoter reporters. In this library, the promoters of *E. coli* genes are fused to a
fast-folding green fluorescent protein (GFP) gene in low-copy-number plasmids, allowing
accurate and reproducible measurement of gene expression[58]. We focused on approximately 40
reporters containing promoters that regulate the key SOS response and associated genes. These
promoters are under the control of the RecA-LexA regulatory system, chosen based on a
comprehensive analysis of *E. coli* databases[59]. In our experiments, *E. coli* MG1655 cells containing
the selected promoters were cultured individually until the mid-exponential phase in 96-well
plates. Subsequently, they were exposed to UV-B radiation for 16 min. This duration was chosen
as it induces maximum SOS upregulation and mutagenesis without drastically impacting cell
culturability (**Fig. 1a,b**). Also, 16-min UV treatments were used as positive controls for the
subsequent sections. GFP levels were measured after 24 h of recovery, allowing adequate time for
UV-induced promoters to express GFP. High cell densities ensured reliable GFP measurements,
given that we used a plate reader in this screening assay.

Our specific focus was on genes showing significant upregulation following UV treatment (**Fig.
2a**), as they likely play pivotal roles in the downstream mechanisms of RecA-mediated SOS
response. Several promoters showed drastic upregulation, including P*recA* and P*lexA* (promoters of

genes encoding SOS response global regulators), P*recN* and P*rmuC* (promoters of genes involved in
DNA recombinational repair), P*sulA* (promoter of a gene encoding cell division inhibitor), P*polB* and
P*dinB* (promoters of genes encoding DNA polymerase enzymes), P*ftsK* (promoter of a gene encoding
essential cell division protein), P*sbmC* (promoters of a gene encoding inhibitor of DNA gyrase-
mediated DNA supercoiling), and P*ybfE* (promoter of a *lexA*-regulated gene whose function is not
well characterized)[60] (**Fig.2a**). To determine whether the genes associated with the identified
promoters are involved in SOS response-mediated culturability and mutagenesis, we generated
single deletions in *E. coli* MG1655 for these genes (except for *lexA* and *ftsK*, as they are essential
and could not be deleted[61–63]). The knockout strains were exposed to UV-B for 16 min or 32 min
or untreated, and their CFU profiles during recovery, as well as their RIF-resistant colony levels,
were similarly measured. For all knockout strains, the temporal CFU profiles remained the same
as that of the wild-type, showing a transient non-culturability during the first 15 min of recovery
after a high UV-B exposure duration of 32 min (**Fig. 2b**). As expected, the deletion of *recA*
completely eliminated mutagenesis (**Fig. 2c**). Although 32 min of UV-B exposure significantly
reduced mutagenesis in all strains, including wild type, the two DNA recombination gene
knockouts, Δ*recN* and Δ*rmuC*, showed a 3-fold and 1.5-fold decrease in RIF-resistant mutant
levels, respectively, compared to WT cells following 16 min of UV-B treatment (**Fig. 2c**).
Surprisingly, other single deletion strains, including those for the genes encoding DNA polymerase
II and IV, *dinB* and *polB*, did not show significant changes (**Fig. 2c**).

Overall, our results showed that single deletions of the downstream genes of RecA did not
drastically impact cellular culturability and mutagenesis, except for RecN and RmuC deletions,
which slightly reduced mutagenesis. RecN is involved in the recombinational repair of DNA
double-strand breaks, and its mutation makes cells sensitive to mitomycin C and ionizing
radiation[64–66]. Although RmuC is thought to be an inner membrane protein with a nuclease
domain[67,68], its function is not well understood, and our data indicate it only moderately impacts
UV-B-mediated mutagenesis. It is quite surprising that the deletion of DNA polymerase II and IV,
two key proteins involved in indirect mutagenesis[69], did not impact UV-B-induced mutagenesis.
As mutagenesis is a multifaceted process regulated by multiple mechanisms, testing multi-deletion
strains might be necessary to show the complex interplay between these mechanisms. Furthermore,
deleting a single polymerase enzyme may have minimal impact due to the abundance of other
polymerase enzymes. We first generated Δ*recN*Δ*rmuC* and showed that the double deletion
resulted in a cumulative decrease (~5 fold) in RIF-resistant colony levels compared to the wild
type following 16 min of UV-B treatment (**Fig. 2c**). To assess the impact of DNA polymerase II
and IV, we generated a Δ*polB*Δ*dinB* mutant strain, which did not affect RIF-resistant colony levels
(**Fig. 2c**). Additionally, we generated Δ*recN*Δ*polB* and Δ*recN*Δ*dinB* double knockout strains and
demonstrated that they exhibited approximately a 5-fold reduction in RIF-resistant colony levels
compared to wild-type cells (**Fig. 2c**). Deletion of DNA polymerase genes *polB* and *dinB* from
Δ*recN*Δ*rmuC* individually did not show significant changes compared to the Δ*recN*Δ*rmuC* strain
in RIF-resistant mutant levels (**Fig. 2c**), suggesting that the observed reduction in RIF-resistant
colonies in multi-deletion strains is primarily due to the effects of *recN* and/or *rmuC*.

Screening the knockout strains reveals the redundancy of repair mechanisms in UV-induced
mutagenesis and culturability

Although our initial screening yielded intriguing results, the critical downstream mechanisms
underlying our observed phenomenon still remain elusive. Thus, we conducted a second screening
using the *E. coli* BW25113 Keio knockout collection to assess the impact of UV on mutagenesis

and culturability of the knockout strains. This screening focused on all potential repair mechanisms
involved[12], particularly those (i) that are not regulated by RecA and (ii) whose expression was not
significantly upregulated in our initial screening (**Fig. 2a**). For this screening, the knockout strains
in 96-well plates were treated with UV-B for 16 min during mid-exponential-phase growth, and
RIF-resistant colonies were quantified after a 24-hour recovery period. We identified several genes
whose deletion resulted in a significant reduction in RIF-resistant colony formation (**Fig. 3a**),
including *recB*, a component of Exonuclease V or the RecBCD complex that promotes
homologous recombination in the repair of double-strand DNA breaks[70,71]; *umuC* and *umuD*,
encoding error-prone DNA polymerase V[72]; *ruvC*, encoding an endonuclease that binds to and
cleaves Holliday junctions[73]; and *katE*, encoding *E. coli* catalase enzyme[74]. Additionally, we
observed several genes that showed an increase in mutagenesis, such as *uvrD* and *uvrA*, which are
involved in the nucleotide excision repair pathway[75,76], and *mutY*, involved in the base excision
repair pathway[77] (**Fig. 3a**).

To better understand how these genes affect mutagenesis and culturability, and to see if similar
trends occur in other *E. coli* strains, we deleted these genes in *E. coli* MG1655 and analyzed the
knockout strains after 16 and 32 min of UV-B exposure. Deletion of *recB*, *ruvC*, *umuC*, and *umuD*
significantly reduced mutagenesis (**Fig. 3c**), consistent with our screening data. The absence of
UmuC and UmuD proteins, specialized for translesion synthesis, was shown to sensitize *E. coli*
cells to UV[78,79], aligning with our findings. While Δ*uvrA* in the *E. coli* BW25113 strain from
screening showed higher mutant levels compared to wild-type cells, its deletion in the MG1655
background resulted in a significant reduction in mutant formation (**Fig. 3c**), suggesting that the
role of certain genes in mutagenesis can vary depending on the genetic background of the *E. coli*
strain. Also, our results indicate that KatE, UvrD, and MutY may not be directly involved in the
mutagenic response to UV radiation under the conditions tested here, as their deletions did not
clearly impact UV-B-induced mutagenesis in *E. coli* MG1655 (**Fig. 3c**). Interestingly, Δ*uvrD* and
Δ*mutY* strains exhibited increased RIF-resistant colonies in cultures not treated with UV-B (**Fig.**
3c). UvrD is involved in various DNA repair pathways, including NER and MMR processes[37,80,81].
MutY is a glycosylase enzyme that corrects adenine mismatches resulting from DNA replication
errors, primarily in the BER pathway[77]. While the specific roles of UvrD and MutY in UV-B-
induced mutagenesis are not evident in this study, the increased RIF-resistant colonies in untreated
cultures might highlight their potential role in maintaining genomic stability under normal growth
conditions, possibly by mitigating spontaneous mutations.

The temporal CFU profiles revealed interesting trends as well. Δ*ruvC*, Δ*umuC*, and Δ*umuD* strains
showed no significant change in transient non-culturability immediately after UV-B treatment
compared to the wild type (**Fig. 3b**). However, Δ*recB* exhibited a notable decrease in CFU levels,
while Δ*uvrA* and Δ*uvrD* mutants showed even more drastic reductions following UV-B treatment
compared to the wild type (**Fig. 3c**). This reduction in CFU levels was transient, as their levels
increased approximately 10^7 -fold within 15 min of recovery. UvrA and UvrD together form key
components of the NER pathway[80]. UvrA forms a complex with UvrB to detect DNA lesions,
while UvrD functions as a helicase to unwind and remove damaged DNA segments[81,82]. As both
UvrA and UvrD are regulated by the RecA protein[28], it is not surprising that the deletion of *recA*
resulted in drastic reductions in CFU levels following UV treatment (**Fig. 1c**). Since RecA is
essential for inducing the SOS response, its absence means that many DNA repair genes, including
uvrA and *uvrD*, are not upregulated in response to UV-induced DNA damage.

Excessive UV exposure impairs cellular translation processes

Although our results highlight specific SOS proteins involved in mutagenesis and cell culturability, it remains unclear why excessive UV exposure impairs *recA* expression and, consequently, the SOS response. This impairment could be due to the activation of cellular mechanisms, such as growth arrest proteins, which may help limit excessive DNA damage and mutagenesis. Additionally, the production of toxic molecules, such as reactive oxygen species, or the inhibition of certain cellular functions, like translation, may also contribute to this effect. These mechanisms—namely the involvement of growth arrest proteins, reactive oxygen species, or the inhibition of translation—will be thoroughly explored in the subsequent section of our study.

Within the SOS response, two genes, *sulA* and *tisB*, play key roles in cell growth inhibition. The *sulA* gene in *E. coli* encodes a cell-division inhibitor that temporarily halts the cell cycle during the SOS response, preventing damaged DNA from prematurely segregating into daughter cells[83]. On the other hand, *tisB* encodes a toxin protein that reduces cellular ATP levels[46,84]. This protein may also induce reversible dormancy by suppressing cell metabolism. To investigate the individual and combined effects of these genes, we constructed knockout strains of *E. coli* MG1655 Δ*sulA*, Δ*tisB*, and Δ*sulA*Δ*tisB*. Similarly, these strains, at the mid-exponential phase, were exposed to varying durations of UV-B radiation followed by a 24-hour recovery period. *E. coli* MG1655 WT cells were used as the control group for comparison. Deleting *sulA* and *tisB* individually did not alter the trend of transient non-culturability observed in our experiments. Specifically, the CFU profiles initially showed a significant decrease after prolonged UV-B exposure of 24 and 32 min, followed by a sharp increase within the first 15 min of recovery, similar to the wild-type strain (**Fig. 1a vs Fig. 4a-c**). However, CFU levels of Δ*sulA*Δ*tisB* immediately after 24 and 32 min of UV-B exposure (corresponding to the t~0 point on the CFU plots) were nearly 10-fold higher compared to those of the wild-type or single knockout strains (**Fig. 1a vs Fig. 4a-c**). We observed approximately a 10-fold increase in CFU levels within the first 15 min of recovery in the double knockout strain, although this increase was not as pronounced as in the control or single knockout strains (**Fig. 4a-c**). While deleting both genes enhanced the culturability of cells following UV-B treatment, the phenomenon observed in the wild-type strain was not completely eliminated in the double knockout strain, highlighting the complexity of the underlying mechanisms. No distinct profiles of RIF-resistant mutants were observed after 24 hours of recovery in either single or double-knockout strains compared to the wild-type strain (**Fig. 1b vs Fig. 4d-f**), indicating that these genes are not directly involved in mutagenesis, although they may affect cellular culturability to some extent.

UV radiation has the potential to damage cellular membranes and compromise essential cellular functions[15]. This damage can lead to increased membrane permeability, leakage of cellular contents, and, ultimately, cell death[85,86]. Additionally, prolonged exposure to UV radiation can trigger excessive production of reactive oxygen species (ROS) within cells[87–89]. ROS are highly reactive molecules that can cause oxidative damage to cellular components such as DNA, proteins, and lipids, impairing cellular functions. It is possible that the 24-minute and 32-minute UV-B treatments studied here may not be severe enough to kill the bacterial cells completely, but they are likely sufficient to hinder cell growth and mutagenic processes by disrupting vital metabolic processes. To characterize cellular membrane permeability, cells treated with UV-B were immediately collected and incubated with propidium iodide (PI), a fluorescent dye that enters cells with compromised membranes. Flow cytometry data indicated no significant increase in permeabilization following prolonged UV-B treatment (32 min) up to 1 hour of recovery (**Fig. 5a**),

compared to both untreated cells and moderately UV-treated cells (16 min), which did not
markedly reduce CFU levels. Furthermore, when UV-treated samples were subjected to Amplex
Red reagent (see Materials and Methods for details) to detect hydrogen peroxide (H_2O_2), a primary
reactive oxygen species, no significant changes were observed in the 32-minute UV-treated cells
compared to untreated or 16-minute UV-treated cells (**Fig. 5b**), indicating H_2O_2 may not be a key
player. In our previous study, we found that ROS significantly affects cell culturability only when
their concentrations are much higher than physiological levels, and their impact on cell
culturability was observed to be non-transient [50].

UV exposure is known to interfere with ribosomal function by inducing crosslinks in ribosomal
RNAs[90], which can inhibit overall translation processes and impair the expression of SOS response
genes. To assess the effects of UV treatments on cellular translation, we analyzed GFP expression
under the control of the *recA* promoter (P_{recA}) at the single-cell level using *E. coli* MG1655 cells
carrying the pUA66 P_{recA}-*gfp* plasmids. The cells were subjected to UV-B treatments for 16 min
and 32 min, and *recA* expression was measured using flow cytometry during the early recovery
phase (**Fig. 5c**). UV-B treatment should activate promoters of SOS response genes and induce
recA expression. As expected, the 16-min UV-B treatment resulted in significantly higher *recA*
expression compared to untreated cells (**Fig. 5c, e**). However, the 32-min UV-B exposure resulted
in more heterogeneous and lower *recA* expression compared to the 16-min UV-B exposure (**Fig.
5c, e**), potentially implying an impairment in the translational process. This difference between the
16- and 32-min treatments cannot be solely attributed to increased cell death following 32 min of
UV-B exposure. Despite the observed transient culturability in the prolonged UV-exposure
culture, CFU levels in both conditions are nearly similar after the brief transient period, around 15
minutes of recovery culture (see the 0.25-hour time point in **Fig. 1a**). To substantiate the hypothesis
regarding the impairment of the translational process, we employed a GFP-expressing plasmid
where the GFP gene is tightly regulated by an isopropyl β-D-1-thiogalactopyranoside (IPTG)-
inducible T5 promoter and a strong LacI^q repressor[91]. Our rationale was that if excessive UV-B
treatment indeed impairs translational processes, the levels of GFP expression in cells treated with
32 min of UV-B should be lower compared to those in the no-UV control or 16-minute UV-treated
cells, despite the presence of the inducer. We supplemented cell cultures with 0.1 mM IPTG
immediately before UV-B treatments. As anticipated, the 32-min UV-B treatment resulted in a
lower level of GFP expression compared to the no-UV control or the 16-minute UV-B treatment
(**Fig. 5d, f**). However, the 16-min UV-B treatment exhibited the highest GFP expression levels
(**Fig. 5d, f**), suggesting that moderate UV treatment (16 min) enhances translational activity,
whereas excessive UV-B treatment (32 min) slightly inhibits it. This moderate inhibitory effect of
excessive UV-B treatment on cellular translation, including RecA expression, likely impacts
overall cellular processes and compromises the SOS response. This was further validated when we
measured the expression levels of *uvrD* and *uvrA*—two critical genes involved in mutagenesis and
culturability. Their expression levels were slightly reduced under 32 minutes of UV treatment
compared to 16 minutes (**Fig. 6a-d**).

Altogether, our findings suggest that while prolonged UV-B treatments may not directly impact
membrane integrity or ROS levels, the observed unculturability of the cells likely results from a
combined effect of multiple factors, including cell growth inhibitors, impaired translation
processes, and downstream repair pathways. The impairment of cellular processes such as
translation may hinder the SOS response, as evidenced by the decreased expression of RecA in
cells treated with 32 min of UV-B (**Fig. 5e**). This interpretation is further supported by our findings
with the Δ*recA* strain, which exhibited near-complete unculturability immediately after UV

treatments. The immediate reduction in CFU levels in the knockout strains ($\Delta recA$, $\Delta uvrA$ and
$\Delta uvrD$), as well as in the wild-type strain exposed to prolonged UV, highlights the importance of
the SOS response in the initial phases of DNA damage repair. The transient nature of CFU
reduction in all these mutants also suggests that *E. coli* possesses redundant or numerous repair
mechanisms[12] that can partially restore culturability even when certain DNA repair pathways are
disrupted.

Conclusions

In this study, we investigated the effects of UV-B treatments on *E. coli* mutagenesis and
culturability and provided important insights into bacterial survival mechanisms under stress.
Mutagenesis, measured by RIF-resistant colony formation, peaked at 16 min of UV-B exposure
but decreased with longer exposures (24 and 32 min). The longer UV-B exposures also led to a
significant decrease in CFU levels, but this phenomenon was observed to be a transient state, as
we saw a drastic jump in CFU levels after a short recovery period. We also explored the role of
key SOS response genes and demonstrated that knockout strains lacking *sulA* and *tisB* genes
exhibited similar patterns of transient non-culturability as wild-type cells, although double
knockouts exhibited higher initial CFU levels post-exposure, indicating a partial role in
culturability. Further investigation revealed that excessive UV exposure did not significantly affect
membrane permeability or reactive oxygen species (ROS) levels but moderately impaired
translation processes, as evidenced by reduced expression of SOS response genes like *recA* or
IPTG-inducible fluorescent proteins. This impaired translation likely contributed to the observed
reduction in mutagenesis and transient non-culturability following prolonged UV exposure, given
the significant impact of *recA* deletion on cell culturability and mutagenesis.

Although our screening of promoter reporters highlighted several genes significantly upregulated
following UV treatment, the knockout strains of most of these genes generally had minimal impact
on culturability or mutagenesis, except for $\Delta recN$ and $\Delta rmuC$, which moderately reduced
mutagenesis. In our second screening with the Keio knockout collection, we identified several
critical genes, such as *recB, umuC, umuD,* and *ruvC*, whose deletions led to decreased
mutagenesis, highlighting their crucial roles in facilitating error-prone repair mechanisms during
UV-B stress. Furthermore, we observed differential impacts of gene deletions on mutagenesis
versus culturability, providing insights into additional functions of these genes. Particularly in
strains lacking *uvrA*, or *uvrD* showed a drastic but transient reduction in their culturability
following UV treatment, highlighting their role in maintaining genomic stability. Our findings also
underline the abundance of repair mechanisms in *E. coli*, demonstrating that multiple independent
pathways can significantly impact cell mutagenesis and culturability. Overall, this study
contributes to understanding bacterial responses to environmental stressors and the complex
interactions of DNA repair pathways in maintaining genomic integrity, culturability, and survival.

Materials and Methods

Bacterial strains and plasmids

Escherichia coli K-12 MG1655 wild type (WT), and pUA66 plasmids with genes encoding a green
fluorescent protein (*gfp*) under the control of promoter, P_{recA} were obtained from Dr. Mark P.
Brynildsen at Princeton University. *E. coli* K-12 MG1655 $\Delta sulA$, $\Delta tisB$, $\Delta recA$ was constructed in
our previous studies [50,92,93]. The promoter strain collection of *E. coli* MG1655 in a 96-well plate
format used for the screening assay was obtained from Horizon Discovery, Lafayette, CO, USA.
The Keio knockout strain collection (a single-gene deletion library of E. coli K-12 BW25113) was

obtained from Dharmacon Keio Collection (Dharmacon, Cat# OEC4988). The strains from both the collections used in this study are listed in **Supplementary Table 1**. The knockout *E. coli* MG1655 strains generated for this study are detailed in **Supplementary Table 2**. The method of Datsenko and Wanner[94] was used to generate these strains and the oligonucleotides used to delete the genes are provided in **Supplementary Table 3.**

Chemicals, media, and culture conditions

Unless otherwise specified, all chemicals were procured from Fisher Scientific (Atlanta, GA), VWR International (Pittsburgh, PA), or Sigma Aldrich (St. Louis, MO). PI staining kit was purchased from Promega Corporation (Madison, WI). *E. coli* cells were cultured in liquid Lysogeny-Broth (LB) medium. LB agar medium was utilized for enumerating colony-forming units (CFU) of *E. coli*. The liquid LB medium was prepared by dissolving 5 g yeast extract, 10 g tryptone, and 10 g sodium chloride in 1 L of deionized (DI) water. LB Agar media were prepared by dissolving pre-mixed 40 g LB agar in 1 L of DI water. Both solid and liquid media were subjected to autoclaving for sterilization.

Kanamycin (50 μg/mL) was included in the liquid LB media for plasmid selection and retention. IPTG at 0.1 mM was used to induce *gfp* expression. For PI staining, sterile 0.85% sodium chloride solution was used. When necessary, cells were washed with phosphate-buffered saline (PBS, 1X). Stock solutions for rifampicin (RIF; 50 mg/mL) was prepared by dissolving in DI water using 0.01 N sodium hydroxide. The IPTG stock solution (1 mM) was dissolved in DI water. All chemical solutions were sterilized using 0.2 μm VWR syringe filters.

To prepare RIF-agar plates, the stock RIF solution was added to autoclaved LB agar, resulting in a final plate concentration of 500 μg/mL RIF. Unless specified otherwise, overnight pre-cultures were generated in 14-mL Falcon test tubes containing 2 mL of liquid media. These pre-cultures were inoculated from a 25% glycerol cell stocks stored at −80 °C and cultivated for 24 h at 37 °C with shaking at 250 revolutions per min (rpm). Experimental cell cultures were prepared by diluting the overnight pre-cultures (1:100) into 2 mL of fresh LB medium in 14-mL Falcon test tubes. Bacterial cells in this study reached the mid-exponential phase ($OD_{600} \sim 0.5$) after around 3 h, attaining an average cell density of 7×10^8 CFUs/ml. All treatments involving UV were administered at this stage.

UV treatment and cell recovery

Overnight pre-cultures of *E. coli* MG1655 cells were diluted 100-fold in 2 mL fresh LB media in test tubes and grown at 37°C with shaking (250 rpm). Cell growth was monitored by measuring optical density at 600 nm wavelength (OD_{600}) with a plate reader (Varioskan LUX Multimode Microplate Reader, Thermo Fisher, Waltham, MA, United States). When the cell density reached $OD_{600} \sim 0.5$, cultures from the test tubes were transferred to petri dishes (the diameter of petri dishes= 100 mm, catalog no. FB0875713, Fisher Scientific). This procedure created a thin film of culture in the dish with a height of about 0.25 mm. Then, these cultures were exposed to a light source emitting UV-B (302 nm) at approximately 2000μW (UVP ChemStudio, catalog no. 849-97-0928-02; Analytik Jena, Jena, Germany) for varying exposure times (0, 2, 4, 8, 16, 24 and 32 min). After UV exposure, cells were transferred back to test tubes and recovered for 24 h. During the recovery period, 10 μl samples were collected at specified time points from each test tube, serially diluted in PBS in round-bottom 96-well plates, and then, plated on LB agar media. The plates were incubated at 37 °C for 16 h to enumerate CFU. We note that new colonies were not formed when incubated beyond 16 h. A similar procedure was followed for the other strains.

Assessing mutagenesis

To assess the extent of mutant cell formation induced by UV exposure, cells were exposed to UV radiation for the specified durations, followed by a 24-h recovery period. After this recovery period, 500 μL of cells were collected and spread onto agar plates containing 500 μg/mL of RIF. These plates were then incubated at 37 °C for 16 h to enumerate RIF-resistant colonies. To determine clonogenic survival, CFU levels were determined, and mutant levels were normalized by dividing the number of RIF-resistant colonies by the total number of colonies in a 1 mL culture volume. Unless specified otherwise, mutant formation was reported as the count of RIF-resistant colonies per 10^8 cell population.

PI Staining

Mid-exponential *E. coli* cells were treated with UV for 16 min and 32 min duration following the protocol mentioned above. At the beginning of the recovery and 1h after the recovery (t=0 h and 1 h), UV-treated cells in LB media were diluted 20-fold in 1.0 ml 0.85% NaCl solution in flow cytometry tubes (5 ml round bottom Falcon tubes, size: 12×75 mm) to achieve the desired cell density ($\sim 10^6$-10^7 cells/ml) for flow cytometry analysis. The resulting cell suspensions were treated with 20 μM PI dye. PI produces red fluorescence upon binding DNA; however, it can only penetrate cells with damaged membranes. The samples were incubated in the dark at 37 °C for 15 min before analyzing them with a conventional bench-top flow cytometer (NovoCyte 3000RYB, ACEA Biosciences Inc., San Diego, CA, United States). For flow cytometry analysis, we chose a slow sample flow rate (14 μl/ min) to have a sample stream diameter (i.e., core diameter) of 7.7 μm. The instrument has a constant sheath flow rate of 6.5 ml/min. The flow cytometer utilizes low-power solid-state lasers. Cells were excited at a 561 nm wavelength and red fluorescence was detected with a 615/20-nm bandpass filter. At least 30,000 events were recorded for each sample. NovoExpress software was used to collect the data. PI-stained dead cells, obtained after ethanol (70% v/v) treatment, were used as a positive control. PI-stained live cells were used as a negative control. Forward and side scatter signals of untreated live cells were used to determine the cells on flow cytometry diagrams; the positive and negative controls were used to gate PI positive (+) and PI negative (-) cell populations **(Supplementary Fig. S1)**

H₂O₂ measurement

The Amplex™ Red Hydrogen Peroxide/Peroxidase Assay Kit (Invitrogen, Thermo fischer scientific, Catalog number: A22188) was used to assess the amount of H_2O_2 formed in the UV-treated cells. The Amplex™ Red Hydrogen Peroxide/Peroxidase Assay Kit contains a sensitive, one-step assay that uses the Amplex™ Red reagent (10-acetyl-3,7-dihydroxyphenoxazine) in combination with horseradish peroxidase (HRP) to detect hydrogen peroxide (H_2O_2). In the presence of peroxidase, the Amplex™ Red reagent reacts with H_2O_2 in a 1:1 stoichiometry to produce the red-fluorescent oxidation product, resorufin. Resorufin has excitation and emission maxima of approximately 571 nm and 585 nm. For preparing the stock solutions, 10 mM Amplex® Red reagent was prepared by dissolving the contents of the vial of Amplex® Red reagent in 60 μL of DMSO. Reaction buffer was diluted 5-fold to prepare 1X reaction buffer and 10 U/mL Horseradish Peroxidase (HRP) was prepared by dissolving one vial of HRP in 1.0 mL of 1X Reaction Buffer. The working solution of 100 μM Amplex® Red reagent and 0.2 U/mL HRP was prepared by adding 50 μL of 10 mM Amplex® Red reagent stock solution and 100 μL of 10 U/mL HRP stock solution in 4.85 mL of 1X Reaction Buffer. The assay volume for this experiment was

100 µL. Mid-exponential *E. coli* cells were treated with UV for 16 min and 32 min duration and
right after the treatment, 50 µL of sample for each condition was serially diluted in 1X Reaction
Buffer to determine the optimal amount of sample for the assay. 50 µL of the Amplex® Red
reagent/HRP working solution was added to each microplate well containing the samples. The
samples were incubated at room temperature for 30 min, and protected from light. At the end, the
fluorescence was measured using a microplate reader equipped for excitation in the range of 530–
560 nm and fluorescence emission detection at ~590 nm. For the standard curve, 20 mM Hydrogen
Peroxide (H_2O_2) working solution was prepared by dissolving 3.0% (0.88 M) H_2O_2 in 1X Reaction
Buffer and then further diluting 20 mM H_2O_2 working solution into 1X Reaction Buffer to produce
H_2O_2 concentrations of 0 to 10 µM, each in a volume of 50 µL. 50 µL of the Amplex® Red
reagent/HRP working solution was added to each microplate well containing the standards
followed by 30 min of incubation and then measuring the fluorescence following the same way
(Supplementary Fig. S2).

Fluorescent protein expression assay for reporter genes

Overnight pre-cultures of *E. coli* MG1655 cells with reporter genes fused to the SOS gene
promoters (P_{recA}, P_{uvrA}, and P_{uvrD}) were diluted 1:100 in 2 mL of LB media within test tubes and
incubated at 37°C with shaking (250 rpm). Upon reaching the mid-exponential phase, the cells
were subjected to UV radiation for specified durations (16 min and 32 min). An untreated culture
was used as a control. At the beginning of the recovery, 20 µM PI dye was added to the cultures
for continuous monitoring of the membrane permeability. At specified time points during the
recovery (t=0, 0.25, 0.5, 1, 2, 3 and 4 h), UV-treated cells were diluted 20-fold in 1.0 ml 0.85%
NaCl solution in flow cytometry tubes (5 ml round bottom Falcon tubes, size: 12×75 mm) to
achieve the desired cell density (~10^6-10^7 cells/ml). The samples were analyzed using the same
flow cytometry method described above (see the "PI staining" section); however, cells were
analyzed using two lasers. For measuring the red fluorescence from PI dye, cells were excited at a
561 nm wavelength, and red fluorescence was detected with a 615/20-nm bandpass filter. For
measuring the green fluorescence, cells were excited at a 488 nm wavelength, and the green
fluorescence was detected with a 530/30 bandpass filter.

Transcription/translation activities of pUA66-*gfp* plasmid

To assess the effect of the UV treatment on transcription/translation, the amount of GFP produced
by *E. coli* strains from the low-copy plasmid was measured. The plasmid, pUA66-*gfp*, has a *gfp*
gene under the control of a strong, IPTG-inducible T5 promoter and a strong LacI^q repressor.
Overnight pre-cultures of *E. coli* MG1655 cells carrying pUA66-*gfp* were diluted 100-fold in 2
mL fresh LB media in test tubes and grown at 37°C with shaking (250 rpm). At the mid-
exponential phase (OD_{600}~0.5), 0.1mM IPTG was added to the test tubes and then immediately
treated with UV for 16 min and 32 min duration. Untreated cultures having IPTG only (without
UV exposure) served as controls. At specified time points during the recovery (t=0, 0.25, 0.5, 1,
2, 3, and 4 h), UV-treated cells were diluted 20-fold in 1.0 ml PBS in flow cytometry tubes (5 ml
round bottom Falcon tubes, size: 12×75 mm) to achieve the desired cell density (~10^6-10^7
cells/ml). The samples were analyzed using the same flow cytometry method described above (see
the "PI staining" section); however, cells were analyzed with a laser emitting light at 488 nm and
the green fluorescence was detected with a 530/30 bandpass filter.

Promoter library screening

Overnight precultures were prepared by inoculating the strains from the promoter library into the wells of 96-well plates containing 200 µl LB medium and 50 µg/ml kanamycin (for plasmid retention). The plates were sealed with a sterile, oxygen-permeable membrane (Breathe-Easier, Cat# BERM-2000, VWR International) and cultured for 24 h at 37°C with shaking at 250 rpm. Overnight precultures were diluted 1:40 in LB medium with kanamycin in a new 96-well plate, sealed, and incubated at 37°C with shaking at 250 rpm. At the mid-exponential phase (OD_{600} = 0.5), cultures in 96 well plates were exposed to UV light (UVP ChemStudio, catalog no. 849-97-0928-02; Analytik Jena, Jena, Germany) for 16 min, and then, recovered for 24 h at 37°C with shaking at 250 rpm. GFP was measured with a Varioskan LUX Multimode Microplate Reader (Thermo Fisher, Waltham, MA, USA) at the indicated times with untreated cultures as a control. The excitation and emission wavelengths for GFP measurement were 485 nm and 511 nm, respectively. Fold changes for GFP (treated/untreated cultures after 24 h treatment) were used to report the promoter activity. The top 10 promoters showing the highest expression were selected for subsequent experiments.

Screening *E. coli* BW25113 Keio knockout collection

Overnight cultures of single mutants **(Supplementary Table 1)** carrying kanamycin resistance genes were diluted 40-fold into flat-bottom 96-well plates, with each well containing 200 µL of cell culture. Kanamycin (50 µg/ml) was added to overnight and treatment cultures to prevent contamination. The plates were securely sealed using a sterile, oxygen-permeable membrane (Breathe-Easier, Cat# BERM-2000, VWR International) and incubated at 37°C with agitation at 250 rpm. When the cultures reached the mid-exponential phase (OD_{600} ~ 0.5), the cells in the 96-well plates were exposed to UV light (UVP ChemStudio, catalog no. 849-97-0928-02; Analytik Jena, Jena, Germany) for 16 min. Subsequently, they were allowed to recover for 24 h at 37°C with shaking at 250 rpm. To quantify UV-induced mutant cells, 10 µL samples were collected from each well and spotted onto LB agar plates, containing 500 µg/mL RIF, at the end of the 24-h recovery period. The plates were then incubated at 37°C for 16 h to enumerate RIF-resistant colonies. The mutant formation was reported as the count of RIF-resistant colonies per 10 µL of cell culture.

Statistical analysis and reproducibility

For all pairwise comparisons, one-way ANOVA with Dunnett's post-test was utilized. A minimum of four independent biological replicates (unless otherwise specified) were conducted for experiments involving UV exposure. In all figures, data corresponding to each time point represents the mean value ± standard deviation. Regarding statistical significance analysis, the threshold values were set as follows: *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ****$P < 0.0001$. All figures were created using GraphPad Prism 10.0.2, and the statistical analyses were carried out using GraphPad Prism 10.0.2 statistical functions. FlowJo V 10.7.1 was used to analyze the data obtained from flow cytometry.

Acknowledgment

The authors would like to thank the members of Orman Lab for their help. This study was supported by NIH/NIAID R01-AI143643.

Contributions

S.G. and M.A.O. conceived and designed the study. S.G. performed the experiments. S.G. and M.A.O. analyzed the data and wrote the paper.

Corresponding author

Correspondence to Mehmet A. Orman (morman@central.uh.edu)

Declaration of interests

The authors declare no competing interests.

Data availability

All data in this manuscript are available in either the main text or supplementary files. All raw data have been deposited in the Figshare repository (doi.org/10.6084/m9.figshare.27701493.v1).

References

1. Hershberg, R. Mutation—The Engine of Evolution: Studying Mutation and Its Role in the Evolution of Bacteria. *Cold Spring Harb Perspect Biol* **7**, (2015).
2. MacLean, R. C., Torres-Barceló, C. & Moxon, R. Evaluating evolutionary models of stress-induced mutagenesis in bacteria. *Nature Reviews Genetics 2013 14:3* **14**, 221–227 (2013).
3. Zimmermann, A., Prieto-Vivas, J. E., Voordeckers, K., Bi, C. & Verstrepen, K. J. Mutagenesis techniques for evolutionary engineering of microbes – exploiting CRISPR-Cas, oligonucleotides, recombinases, and polymerases. *Trends Microbiol* (2024) doi:10.1016/J.TIM.2024.02.006.
4. Sung, W., Ackerman, M. S., Miller, S. F., Doak, T. G. & Lynch, M. Drift-barrier hypothesis and mutation-rate evolution. *Proc Natl Acad Sci U S A* **109**, 18488–18492 (2012).
5. Wielgoss, S. *et al.* Mutation rate dynamics in a bacterial population reflect tension between adaptation and genetic load. *Proc Natl Acad Sci U S A* **110**, 222–227 (2013).
6. Fitzgerald, D. M., Hastings, P. J. & Rosenberg, S. M. Stress-Induced Mutagenesis: Implications in Cancer and Drug Resistance. *Annu Rev Cancer Biol* **1**, 119 (2017).
7. Morreall, J. F., Petrova, L. & Doetsch, P. W. Transcriptional mutagenesis and its potential roles in the etiology of cancer and bacterial antibiotic resistance. *J Cell Physiol* **228**, 2257 (2013).
8. Russo, M., Sogari, A. & Bardelli, A. Adaptive evolution: How bacteria and cancer cells survive stressful conditions and drug treatment. *Cancer Discov* **11**, 1886–1895 (2021).
9. Lambert, G. *et al.* An analogy between the evolution of drug resistance in bacterial communities and malignant tissues. *Nat Rev Cancer* **11**, 375 (2011).
10. Rothschild, L. J. The Influence of UV Radiation on Protistan Evolution. *Journal of Eukaryotic Microbiology* **46**, 548–555 (1999).
11. Rozema, J. *et al.* The role of UV-B radiation in aquatic and terrestrial ecosystems—an experimental and functional analysis of the evolution of UV-absorbing compounds. *J Photochem Photobiol B* **66**, 2–12 (2002).

12. Rastogi, R. P., Richa, Kumar, A., Tyagi, M. B. & Sinha, R. P. Molecular Mechanisms of
Ultraviolet Radiation-Induced DNA Damage and Repair. *J Nucleic Acids* **2010**, 32 (2010).
13. Sinha, R. P. & Häder, D. P. UV-induced DNA damage and repair: a review.
Photochemical & Photobiological Sciences 2002 1:4 **1**, 225–236 (2002).
14. Krishna, S., Maslov, S. & Sneppen, K. UV-Induced Mutagenesis in Escherichia coli SOS
Response: A Quantitative Model. *PLoS Comput Biol* **3**, e41 (2007).
15. Kciuk, M., Marciniak, B., Mojzych, M. & Kontek, R. Focus on UV-Induced DNA
Damage and Repair—Disease Relevance and Protective Strategies. *Int J Mol Sci* **21**, 1–33
(2020).
16. Gray, N. F. Ultraviolet Disinfection. *Microbiology of Waterborne Diseases:*
Microbiological Aspects and Risks: Second Edition 617–630 (2014) doi:10.1016/B978-0-
12-415846-7.00034-2.
17. Hessling, M., Haag, R., Sieber, N. & Vatter, P. The impact of far-UVC radiation (200-230
nm) on pathogens, cells, skin, and eyes - a collection and analysis of a hundred years of
data. *GMS Hyg Infect Control* **16**, Doc07 (2021).
18. Black, H. S. *et al.* Photocarcinogenesis: an overview. *J Photochem Photobiol B* **40**, 29–47
(1997).
19. Douki, T., Court, M., Sauvaigo, S., Odin, F. & Cadet, J. Formation of the main UV-
induced thymine dimeric lesions within isolated and cellular DNA as measured by high
performance liquid chromatography-tandem mass spectrometry. *J Biol Chem* **275**, 11678–
11685 (2000).
20. Rupert, C. S. Enzymatic Photoreactivation: Overview. *Basic Life Sci* **5 A**, 73–87 (1975).
21. Seawell, P. C. *et al.* den V gene of bacteriophage T4 determines a DNA glycosylase
specific for pyrimidine dimers in DNA. *J Virol* **35**, 790–796 (1980).
22. Van Houten, B. Nucleotide excision repair in Escherichia coli. *Microbiol Rev* **54**, 18–51
(1990).
23. Rupp, W. D. & Howard-flanders, P. Discontinuities in the DNA synthesized in an
Excision-defective strain of Escherichia coli following ultraviolet irradiation. *J Mol Biol*
31, 291–304 (1968).
24. Wendel, B. M., Hollingsworth, S., Courcelle, C. T. & Courcelle, J. UV-induced DNA
damage disrupts the coordination between replication initiation, elongation and
completion. *Genes to Cells* **26**, 94–108 (2021).
25. Majchrzak, M., Bowater, R. P., Staczek, P. & Parniewski, P. SOS Repair and DNA
Supercoiling Influence the Genetic Stability of DNA Triplet Repeats in Escherichia coli. *J*
Mol Biol **364**, 612–624 (2006).
26. Simmons, L. A., Foti, J. J., Cohen, S. E. & Walker, G. C. The SOS Regulatory Network.
EcoSal Plus **3**, (2008).
27. Little, J. W. & Mount, D. W. The SOS regulatory system of Escherichia coli. *Cell* **29**, 11–
22 (1982).
28. Maslowska, K. H., Makiela-Dzbenska, K. & Fijalkowska, I. J. The SOS system: A
complex and tightly regulated response to DNA damage. *Environ Mol Mutagen* **60**, 368–
384 (2019).
29. Walker, G. C. Mutagenesis and inducible responses to deoxyribonucleic acid damage in
Escherichia coli. *Microbiol Rev* **48**, 60–93 (1984).
30. Montelone, B. A. DNA Repair and Mutagenesis. Second Edition. By Errol C Friedberg,
Graham C Walker, Wolfram Siede, Richard D Wood, Roger A Schultz, and, Tom

Ellenberger. Washington (DC): ASM Press. $179.95. xxix + 1118 p + 11 pl; ill.; index. ISBN: 1-55581-319-4. 2006. . *https://doi.org/10.1086/509407* **81**, 273–273 (2015).

31. Martins-Pinheiro, M., Marques, R. C. P. & Menck, C. F. M. Genome analysis of DNA repair genes in the alpha proteobacterium Caulobacter crescentus. *BMC Microbiol* **7**, 17 (2007).

32. Kuzminov, A. Recombinational Repair of DNA Damage in Escherichia coli and Bacteriophage λ . *Microbiology and Molecular Biology Reviews* **63**, 751–813 (1999).

33. Burckhardt, S. E., Woodgate, R., Scheuermann, R. H. & Echols, H. UmuD mutagenesis protein of Escherichia coli: overproduction, purification, and cleavage by RecA. *Proceedings of the National Academy of Sciences* **85**, 1811–1815 (1988).

34. Woodgate, R., Rajagopalan, M., Lu, C. & Echols, H. UmuC mutagenesis protein of Escherichia coli: purification and interaction with UmuD and UmuD'. *Proceedings of the National Academy of Sciences* **86**, 7301–7305 (1989).

35. Kato, T. & Shinoura, Y. Isolation and characterization of mutants of Escherichia coli deficient in induction of mutations by ultraviolet light. *MGG Molecular & General Genetics* **156**, 121–131 (1977).

36. Wang, Z. Translesion synthesis by the UmuC family of DNA polymerases. *Mutation Research/DNA Repair* **486**, 59–70 (2001).

37. Liu, J. *et al.* MutL sliding clamps coordinate exonuclease-independent Escherichia coli mismatch repair. *Nat Commun* **10**, (2019).

38. Lin, T. *et al.* Contribution of the SOS response and the DNA repair systems to norfloxacin induced mutations in E. coli. *Mar Life Sci Technol* **5**, 538–550 (2023).

39. Shibai, A. *et al.* Mutation accumulation under UV radiation in Escherichia coli. *Scientific Reports 2017 7:1* **7**, 1–12 (2017).

40. Badran, A. H. & Liu, D. R. Development of potent in vivo mutagenesis plasmids with broad mutational spectra. *Nature Communications 2015 6:1* **6**, 1–10 (2015).

41. Tillich, U. M., Lehmann, S., Schulze, K., Dühring, U. & Frohme, M. The Optimal Mutagen Dosage to Induce Point-Mutations in Synechocystis sp. PCC6803 and Its Application to Promote Temperature Tolerance. *PLoS One* **7**, e49467 (2012).

42. Hashimoto, S. *et al.* Isolation of auxotrophic mutants of diploid industrial yeast strains after UV mutagenesis. *Appl Environ Microbiol* **71**, 312–319 (2005).

43. Hijnen, W. A. M., Beerendonk, E. F. & Medema, G. J. Inactivation credit of UV radiation for viruses, bacteria and protozoan (oo)cysts in water: A review. *Water Res* **40**, 3–22 (2006).

44. Reed, N. G. The History of Ultraviolet Germicidal Irradiation for Air Disinfection. *Public Health Reports* **125**, 15 (2010).

45. Bergkessel, M., Basta, D. W. & Newman, D. K. The physiology of growth arrest: uniting molecular and environmental microbiology. *Nat Rev Microbiol* **14**, 549 (2016).

46. Dörr, T., Vulić, M. & Lewis, K. Ciprofloxacin causes persister formation by inducing the TisB toxin in Escherichia coli. *PLoS Biol* **8**, 29–35 (2010).

47. Völzing, K. G. & Brynildsen, M. P. Stationary-phase persisters to ofloxacin sustain DNA damage and require repair systems only during recovery. *mBio* **6**, (2015).

48. Santos, A. L. *et al.* Effects of UV radiation on the lipids and proteins of bacteria studied by mid-infrared spectroscopy. *Environ Sci Technol* **47**, 6306–6315 (2013).

49. Wurtmann, E. J. & Wolin, S. L. RNA under attack: Cellular handling of RNA damage. *Crit Rev Biochem Mol Biol* **44**, 34 (2009).

50. Ghosh, S. & Orman, M. A. Exploring the links between SOS response, mutagenesis, and
resistance during the recovery period. *Antimicrob Agents Chemother* **68**, (2024).

51. Barrett, T. C., Mok, W. W. K., Murawski, A. M. & Brynildsen, M. P. Enhanced antibiotic
resistance development from fluoroquinolone persisters after a single exposure to
antibiotic. *Nature Communications 2019 10:1* **10**, 1–11 (2019).

52. Krašovec, R. *et al.* Mutation rate plasticity in rifampicin resistance depends on
Escherichia coli cell-cell interactions. *Nat Commun* **5**, (2014).

53. Reynolds, M. G. Compensatory evolution in rifampin-resistant Escherichia coli. *Genetics*
156, 1471 (2000).

54. Pope, C. F., O'Sullivan, D. M., McHugh, T. D. & Gillespie, S. H. A Practical Guide to
Measuring Mutation Rates in Antibiotic Resistance. *Antimicrob Agents Chemother* **52**,
1209 (2008).

55. Rosche, W. A. & Foster, P. L. Determining Mutation Rates in Bacterial Populations.
Methods **20**, 4 (2000).

56. Lagage, V., Chen, V. & Uphoff, S. Adaptation delay causes a burst of mutations in
bacteria responding to oxidative stress. *EMBO Rep* **24**, (2023).

57. Rodríguez-Rojas, A. *et al.* Non-lethal exposure to H2O2 boosts bacterial survival and
evolvability against oxidative stress. *PLoS Genet* **16**, e1008649 (2020).

58. Zaslaver, A. *et al.* A comprehensive library of fluorescent transcriptional reporters for
Escherichia coli. *Nat Methods* **3**, 623–628 (2006).

59. Keseler, I. M. *et al.* The EcoCyc database: reflecting new knowledge about Escherichia
coli K-12. *Nucleic Acids Res* **45**, D543–D550 (2017).

60. Fernández De Henestrosa, A. R. *et al.* Identification of additional genes belonging to the
LexA regulon in Escherichia coli. *Mol Microbiol* **35**, 1560–1572 (2000).

61. Goodall, E. C. A. *et al.* The Essential Genome of Escherichia coli K-12. *mBio* **9**, (2018).

62. Yu, X.-C., Weihe, E. K. & Margolin, W. Role of the C Terminus of FtsK in Escherichia
coli Chromosome Segregation. *J Bacteriol* **180**, 6424 (1998).

63. Wang, L. & Lutkenhaus, J. FtsK is an essential cell division protein that is localized to the
septum and induced as part of the SOS response. *Mol Microbiol* **29**, 731–740 (1998).

64. Meddows, T. R., Savory, A. P., Grove, J. I., Moore, T. & Lloyd, R. G. RecN protein and
transcription factor DksA combine to promote faithful recombinational repair of DNA
double-strand breaks. *Mol Microbiol* **57**, 97–110 (2005).

65. Picksley, S. M., Attfield, P. V. & Lloyd, R. G. Repair of DNA double-strand breaks in
Escherichia coli K12 requires a functional recN product. *Mol Gen Genet* **195**, 267–274
(1984).

66. Wang, T. C. & Smith, K. C. Different effects of recJ and recN mutations on the
postreplication repair of UV-damaged DNA in Escherichia coli K-12. *J Bacteriol* **170**,
2555–2559 (1988).

67. Kosinski, J., Feder, M. & Bujnicki, J. M. The PD-(D/E)XK superfamily revisited:
identification of new members among proteins involved in DNA metabolism and
functional predictions for domains of (hitherto) unknown function. *BMC Bioinformatics* **6**,
(2005).

68. Lomize, A. L., Lomize, M. A., Krolicki, S. R. & Pogozheva, I. D. Membranome: a
database for proteome-wide analysis of single-pass membrane proteins. *Nucleic Acids Res*
45, D250–D255 (2017).

69.   Napolitano, R., Janel-Bintz, R., Wagner, J. & Fuchs, R. P. P. All three SOS-inducible
DNA polymerases (Pol II, Pol IV and Pol V) are involved in induced mutagenesis. *EMBO*
J **19**, 6259 (2000).

70.   Hickson, I. D., Robson, C. N., Atkinson, K. E., Hutton, L. & Emmerson, P. T.
Reconstitution of RecBC DNase activity from purified Escherichia coli RecB and RecC
proteins. *J Biol Chem* **260**, 1224–9 (1985).

71.   Payne-Dwyer, A. L., Syeda, A. H., Shepherd, J. W., Frame, L. & Leake, M. C. RecA and
RecB: probing complexes of DNA repair proteins with mitomycin C in live Escherichia
coli with single-molecule sensitivity. *J R Soc Interface* **19**, (2022).

72.   Reuven, N. B., Arad, G., Maor-Shoshani, A. & Livneh, Z. The mutagenesis protein UmuC
is a DNA polymerase activated by UmuD', RecA, and SSB and is specialized for
translesion replication. *J Biol Chem* **274**, 31763–31766 (1999).

73.   Takahagi, M., Iwasaki, H., Nakata, A. & Shinagawa, H. Molecular analysis of the
Escherichia coli ruvC gene, which encodes a Holliday junction-specific endonuclease. *J*
Bacteriol **173**, 5747–5753 (1991).

74.   Loewen, P. C. Isolation of catalase-deficient Escherichia coli mutants and genetic
mapping of katE, a locus that affects catalase activity. *J Bacteriol* **157**, 622–626 (1984).

75.   Epshtein, V. *et al.* UvrD facilitates DNA repair by pulling RNA polymerase backwards.
Nature **505**, 372–377 (2014).

76.   Orren, D. K. & Sancar, A. The (A)BC excinuclease of Escherichia coli has only the UvrB
and UvrC subunits in the incision complex. *Proc Natl Acad Sci U S A* **86**, 5237–5241
(1989).

77.   de Oliveira, A. H. S., da Silva, A. E., de Oliveira, I. M., Henriques, J. A. P. & Agnez-
Lima, L. F. MutY-glycosylase: An overview on mutagenesis and activities beyond the GO
system. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* **769**,
119–131 (2014).

78.   Shurtleff, B. W., Ollivierre, J. N., Tehrani, M., Walker, G. C. & Beuning, P. J. Steric Gate
Variants of UmuC Confer UV Hypersensitivity on Escherichia coli. *J Bacteriol* **191**, 4815
(2009).

79.   Wang, T. chien V. & Smith, K. C. Role of the umuC gene in postreplication repair in UV-
irradiated Escherichia coli K-12 uvrB. *Mutation Research/DNA Repair Reports* **145**, 107–
112 (1985).

80.   Moolenaar, G. F., Moorman, C. & Goosen, N. Role of the Escherichia coli Nucleotide
Excision Repair Proteins in DNA Replication. *J Bacteriol* **182**, 5706 (2000).

81.   Orren, D., Selby, C., Hearst, J. & Sancar, A. Post-incision steps of nucleotide excision
repair in Escherichia coli. Disassembly of the UvrBC-DNA complex by helicase II and
DNA polymerase I. *J Biol Chem* **267**, 780–788 (1992).

82.   Bertrand-Burggraf, E., Selby, C. P., Hearst, J. E. & Sancar, A. Identification of the
different intermediates in the interaction of (A)BC excinuclease with its substrates by
DNase I footprinting on two uniquely modified oligonucleotides. *J Mol Biol* **219**, 27–36
(1991).

83.   Higashitani, A., Ishii, Y., Kato, Y. & Horiuchi, K. Functional dissection of a cell-division
inhibitor, SulA, of Escherichia coli and its negative regulation by Lon. *Molecular and*
General Genetics **254**, 351–357 (1997).

84.   Unoson, C. & Wagner, E. G. H. A small SOS-induced toxin is targeted against the inner
membrane in Escherichia coli. *Mol Microbiol* **70**, 258–270 (2008).

85. McKenzie, K. *et al.* The effects of 405 nm light on bacterial membrane integrity determined by salt and bile tolerance assays, leakage of UV-absorbing material and SYTOX green labelling. *Microbiology (N Y)* **162**, 1680 (2016).

86. Yeo, S. K. & Liong, M. T. Effects and applications of sub-lethal ultrasound, electroporation and UV radiations in bioprocessing. *Ann Microbiol* **63**, 813–824 (2013).

87. Cadet, J., Douki, T. & Ravanat, J. L. Oxidatively generated base damage to cellular DNA. *Free Radic Biol Med* **49**, 9–21 (2010).

88. Winterbourn, C. C. Reconciling the chemistry and biology of reactive oxygen species. *Nature Chemical Biology 2008 4:5* **4**, 278–286 (2008).

89. Schieber, M. & Chandel, N. S. ROS Function in Redox Signaling and Oxidative Stress. *Curr Biol* **24**, R453 (2014).

90. Zwieb, C., Ross, A., Rinke, J., Meinke, M. & Brimacombe, R. Evidence for RNA-RNA cross-link formation in Escherichia coli ribosomes. *Nucleic Acids Res* **5**, 2705 (1978).

91. Orman, M. A. & Brynildsen, M. P. Inhibition of stationary phase respiration impairs persister formation in E. coli. *Nature Communications 2015 6:1* **6**, 1–13 (2015).

92. Mohiuddin, S. G., Nguyen, T. V. & Orman, M. A. Pleiotropic actions of phenothiazine drugs are detrimental to Gram-negative bacterial persister cells. *Communications Biology 2022 5:1* **5**, 1–15 (2022).

93. Mohiuddin, S. G., Massahi, A. & Orman, M. A. lon Deletion Impairs Persister Cell Resuscitation in Escherichia coli. *mBio* **13**, (2022).

94. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**, 6640 (2000).

Figures

**Fig. 1: Excessive UV exposure resulted in transient loss of culturability and decreased
mutagenesis.** (a) Exponential-phase *E. coli* MG1655 WT cells were exposed to UV-B light for 0,
2, 4, 8, 16, 24, and 32 minutes, followed by a 24-h recovery period. At specific time points during
recovery (t = 0 h, 0.25 h, 0.5 h, 1 h, 2 h, 4 h, 6 h, and 24 h), cells were collected and plated to
determine colony-forming units (CFU). (b) Levels of UV-induced rifampicin (RIF) resistance
mutations were measured by counting RIF-resistant colonies (per 10^8 cells) in the WT cultures
after recovery for the indicated UV exposure times. (c) The temporal CFU profiles of *E. coli*
MG1655 Δ*recA* cells were monitored during recovery following UV treatment. n=4. Statistical
analysis was performed using one-way ANOVA with Dunnett's post-test, where *$P < 0.05$,
**$P < 0.001$, ***$P < 0.01$, ****$P < 0.0001$. Data corresponding to each time point represent mean
value ± standard deviation.

Fig. 2: UV-induced upregulation of SOS genes and its impact on mutagenesis and cell
culturability. (a) A library of *E. coli* MG1655 strains with promoter reporters for SOS response
genes was UV-treated during the mid-exponential phase and allowed to recover for 24 h. GFP
levels were measured after recovery and normalized to untreated controls. WT cells without any
promoter reporters were used as background controls. (b) Genes upregulated following UV
exposure were individually or combinatorially deleted in *E. coli* MG1655, and a mutagenesis assay
was performed. Mid-exponential phase cells were exposed to UV for 0, 16, and 32 min and
recovered for 24 h. At specified time points during recovery ($t = 0$ h, 0.25 h, 0.5 h, 1 h, 2 h, 4 h, 6
h, and 24 h), cells were collected and plated to determine CFU. (c) RIF-resistant cells were
quantified by plating samples on RIF-agar plates after 24 h of recovery, with results reported as
RIF-resistant colony counts per 10^8 cells for different UV exposure durations. n=4. Statistical
analysis was performed using one-way ANOVA with Dunnett's post-test, where $*P < 0.05$,
$**P < 0.001$, $***P < 0.01$, $****P < 0.0001$. Data corresponding to each time point represents mean
value $\pm$ standard deviation.

897 **Fig. 3: Knockout library screening to elucidate their effects on mutagenesis and cell**
898 **culturability**. (a) The *E. coli* BW25113 Keio knockout collection was screened following 16 min
899 of UV treatment. RIF-resistant cells were quantified by spotting 10 μL samples from each well
900 onto RIF-agar plates after a 24-hour recovery period. (b) Selected genes were deleted in *E. coli*
901 MG1655 and the mutagenesis assay was performed. Mid-exponential phase cells were exposed to

UV for 0, 16 and 32 min and recovered for 24 h. At indicated time points (t = 0 h, 0.25 h, 0.5 h, 1 h, 2 h, 4 h, 6 h and 24 h) during recovery, cells were collected and plated to enumerate CFU. (c) RIF-resistant cells were quantified by plating the samples on RIF-agar plates after 24 h of recovery. The values were reported as RIF-resistant colony count per 10^8 cell population for different UV exposure durations. n=4. Statistical analysis was performed using one-way ANOVA with Dunnett's post-test, where $*P < 0.05$, $**P < 0.001$, $***P < 0.01$. Data corresponding to each time point represents mean value ± standard deviation.

Fig. 4: Perturbation of both SulA and TiB proteins moderately increased cell culturability during recovery. (a-c) Exponential-phase *E. coli* MG1655 Δ*sulA*, Δ*tisB*, and Δ*sulAΔtisB* cells were exposed to UV-B light for 0, 2, 4, 8, 16, 24, and 32 minutes, followed by a 24-h recovery period. At specific time points during recovery, cells were collected and plated to determine their CFU levels. (d-f) Levels of UV-induced RIF resistance mutations were measured by counting RIF-resistant colonies (per 10^8 cells) in the cultures of three knockout strains after recovery for the indicated UV exposure times. n=4. Statistical analysis was performed using one-way ANOVA with Dunnett's post-test, where *$P < 0.05$, **$P < 0.001$, ***$P < 0.01$, ****$P < 0.0001$. Data corresponding to each time point represent mean value ± standard deviation.

Fig. 5: Excessive UV exposure did not compromise cell membrane integrity or increase hydrogen peroxide levels, but it did impair cellular translation. (a) Membrane integrity of UV-treated cells (0, 16, and 32 min of UV exposure) was assessed via flow cytometry with PI staining (see **Supplementary Fig. S1** for controls). FSC-H: Forward light scatter. (b) H_2O_2 concentrations were measured using the Amplex Red Hydrogen Peroxide/Peroxidase Assay Kit. Reaction mixtures containing Amplex Red reagent, horseradish peroxidase, and UV-treated cells in sodium phosphate buffer (pH 7.4) were incubated for 30 minutes at room temperature, followed by fluorescence measurement with a microplate reader. H_2O_2 levels were calculated using a standard curve (**Supplementary Fig. S2**). (c) Expression of P$_{recA}$-*gfp* in UV-treated *E. coli* MG1655 cells was analyzed via flow cytometry for the indicated time points during recovery (P$_{recA}$: the *recA* promoter). (d) IPTG-inducible GFP expression in UV-treated *E. coli* MG1655 pUA66-*gfp* cells

was also assessed by flow cytometry for the indicated time points during recovery. (e) GFP expression profiles of UV-treated cells harboring the *recA* promoter and (f) IPTG-inducible GFP expression systems were evaluated using mean GFP values from flow cytometry data. Note that the vertical lines in the flow diagrams roughly represent the upper, lower, and peak fluorescence levels of the overall cell population exposed to the 16-min UV-B treatment (corresponding to the 4-h treatment time point). These lines were included to help readers more easily distinguish the outcomes of different experimental conditions. $n \geq 4$. Statistical analysis was performed using one-way ANOVA with Dunnett's post-test, where $*P < 0.05$, $**P < 0.001$, $***P < 0.01$, $****P < 0.0001$. Data corresponding to each time point represent mean value $\pm$ standard deviation.

Fig. 6: Flow cytometry analysis of nucleotide excision repair genes followed by UV-treatment
(a, b) The GFP profiles of UV-treated cells harboring P_{uvrA}-*gfp*, and (c, d) the GFP profiles of UV-
treated cells harboring P_{uvrD}-*gfp* were assessed using flow cytometry after UV exposure (for the
indicated time points during recovery). Note that the vertical lines in the flow diagrams roughly
represent the upper, lower, and peak fluorescence levels of the overall cell population exposed to
the 16-min UV-B treatment (corresponding to the 4-h treatment time point). These lines were
included to help readers more easily distinguish the outcomes of different experimental conditions.
n=4. Statistical analysis was performed using two-way ANOVA, where $*P < 0.05$, $**P < 0.001$,
$***P < 0.01$, $****P < 0.0001$. Data corresponding to each time point represent mean
value $\pm$ standard deviation.

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035
Supplementary Information
1037
UV-Induced DNA Repair Mechanisms and Their Effects on Mutagenesis and Culturability in *Escherichia coli*

1040 **Sreyashi Ghosh[1], and Mehmet A. Orman[1]***

1041 [1]William A. Brookshire Department of Chemical and Biomolecular Engineering, University of
1042 Houston, Houston, TX, USA

1043 *Correspondence to: S222 Engineering Bldg 1, 4226 Martin Luther King Boulevard, Houston,
1044 TX 77204-4004, Phone: 713-743-6785, Email: <u>morman@central.uh.edu</u>

1045

Supplementary figures
1047

1048
Fig. S1. PI staining of live and dead cells. Live cells (Red) and ethanol-treated dead cells (70% v/v, Grey) were stained with PI and analyzed by flow cytometry to assess live and dead cell populations. The figure shows a representative flow cytometry diagram. Similar results were observed across all independent biological replicates, n=4.

1053

1054
1055
Fig. S2. Standard curve for H_2O_2 measurement using the Amplex Red Hydrogen
Peroxide/Peroxidase Assay Kit. The standard curve was generated using a 20 mM Hydrogen
Peroxide (H_2O_2) working solution, prepared by dissolving 3.0% (0.88 M) H_2O_2 in 1X Reaction
Buffer. This solution was then diluted in 1X Reaction Buffer to achieve H_2O_2 concentrations
ranging from 0 to 10 μM, in 50 μL aliquots. To each microplate well containing the standards, 50
μL of Amplex® Red reagent/HRP working solution was added. After a 30-minute incubation,
fluorescence was measured using a microplate reader with excitation in the range of 530–560 nm
and emission detection at ~590 nm.
1064

Supplementary tables

Table S1: Promoter library and single-deletion knockout library used in this study.

Bacterial Strains	Source
E. coli K-12 MG1655 *pMSs201* P$_{uvrA}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{dps}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{soxS}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{recX}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{molR}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{polA}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{dinG}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{mutS}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{mutH}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{mutY}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{ybfE}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{sbmC}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{mutM}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{yjiW}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{mutT}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{dinB}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{ftsK}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{sodB}$-*gfp*	Promoter library from Horizon Discovery, Lafayette, CO, USA
E. coli K-12 MG1655 *pMSs201* P$_{recN}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{rmuC}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{dinJ}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{yebG}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{sulA}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{soxR}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{ahpC}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{katE}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{sodC}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{ahpF}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{yhiL}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{uvrC}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{oxyR}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{polB}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{uvrD}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{sodA}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{lexA}$-*gfp*	
E. coli K-12 MG1655 *pMSs201* P$_{recA}$-*gfp*	

E. coli K-12 MG1655 *pMSs201* P*lon*-*gfp*	
E. coli K-12 MG1655 *pMSs201* P*umuD*-*gfp*	
E. coli K-12 MG1655 *pMSs201* P*ssb*-*gfp*	
E. coli K-12 BW25113 Δ*fnr*	
E. coli K-12 BW25113 Δ*soxR*	
E. coli K-12 BW25113 Δ*soxS*	
E. coli K-12 BW25113 Δ*umuD*	
E. coli K-12 BW25113 Δ*recA*	
E. coli K-12 BW25113 Δ*ruvB*	
E. coli K-12 BW25113 Δ*uvrD*	
E. coli K-12 BW25113 Δ*sodB*	
E. coli K-12 BW25113 Δ*sodC*	
E. coli K-12 BW25113 Δ*ahpF*	
E. coli K-12 BW25113 Δ*katG*	
E. coli K-12 BW25113 Δ*sodA*	
E. coli K-12 BW25113 Δ*ygjF*	
E. coli K-12 BW25113 Δ*polA*	
E. coli K-12 BW25113 Δ*ruvA*	
E. coli K-12 BW25113 Δ*ruvC*	
E. coli K-12 BW25113 Δ*umuC*	
E. coli K-12 BW25113 Δ*uvrA*	Keio collection, Cat# OEC4987
E. coli K-12 BW25113 Δ*alkA*	
E. coli K-12 BW25113 Δ*ahpC*	
E. coli K-12 BW25113 Δ*sulA*	
E. coli K-12 BW25113 Δ*katE*	
E. coli K-12 BW25113 Δ*uvrB*	
E. coli K-12 BW25113 Δ*nei*	
E. coli K-12 BW25113 Δ*dinB*	
E. coli K-12 BW25113 Δ*recB*	
E. coli K-12 BW25113 Δ*nth*	
E. coli K-12 BW25113 Δ*uvrC*	
E. coli K-12 BW25113 Δ*fur*	
E. coli K-12 BW25113 Δ*ligB*	
E. coli K-12 BW25113 Δ*ung*	
E. coli K-12 BW25113 Δ*tag*	
E. coli K-12 BW25113 Δ*recN*	
E. coli K-12 BW25113 Δ*mutY*	
E. coli K-12 BW25113 Δ*ycaQ*	
E. coli K-12 BW25113 Δ*mutM*	

E. coli K-12 BW25113 Δ*recC*	
E. coli K-12 BW25113 Δ*polB*	
E. coli K-12 BW25113 Δ*recD*	
E. coli K-12 BW25113 Δ*oxyR*	
E. coli K-12 BW25113 Wild type	

1068

1069

1070 **Table S2: Bacterial plasmid and strains used in this study.**

Bacterial Strains	Source
E. coli K-12 MG1655 pUA66 P*recA-gfp*	Gift from Dr. Mark P. Brynildsen
E. coli K-12 MG1655 pUA66-*gfp*	Gift from Dr. Mark P. Brynildsen
E. coli K-12 MG1655 Wild type	Gift from Dr. Mark P. Brynildsen
E. coli K-12 MG1655 Δ*sulA*	Previous study
E. coli K-12 MG1655 Δ*tisB*	Previous study
E. coli K-12 MG1655 Δ*recA*	Previous study
E. coli K-12 MG1655 Δ*sulA*Δ*tisB*	This study
E. coli K-12 MG1655 Δ*recN*	This study
E. coli K-12 MG1655 Δ*rmuC*	This study
E. coli K-12 MG1655 Δ*polB*	This study
E. coli K-12 MG1655 Δ*dinB*	This study
E. coli K-12 MG1655 Δ*sbmC*	This study
E. coli K-12 MG1655 Δ*ybfE*	This study
E. coli K-12 MG1655 Δ*polB*Δ*dinB*	This study
E. coli K-12 MG1655 Δ*recN*Δ*rmuC*	This study
E. coli K-12 MG1655 Δ*recN*Δ*polB*	This study
E. coli K-12 MG1655 Δ*recN*Δ*dinB*	This study
E. coli K-12 MG1655 Δ*recN*Δ*rmuC*Δ*polB*	This study
E. coli K-12 MG1655 Δ*recN*Δ*rmuC*Δ*dinB*	This study
E. coli K-12 MG1655 Δ*recB*	This study
E. coli K-12 MG1655 Δ*ruvC*	This study
E. coli K-12 MG1655 Δ*umuC*	This study
E. coli K-12 MG1655 Δ*umuD*	This study
E. coli K-12 MG1655 Δ*katE*	This study
E. coli K-12 MG1655 Δ*uvrA*	This study
E. coli K-12 MG1655 Δ*uvrD*	Previous study
E. coli K-12 MG1655 Δ*mutY*	This study

1071

1072

 Table S3: Oligonucleotides used the generating single and multi-deletion mutants

	Oligonucleotides to generate gene deletions		
Mutation	**Forward Primer (5' to 3'):**	**Reverse Primer (5' to 3')**	**Source**
Δ*recN*::KAN^R	ACACAATAACAGTAATGGTTTTTCATACAGGAAAACGACTGTGTAGGCTGGAGCTGCTTC	CGTTTTGCTGTTTACTCTGACCGTGAAGCAGGAAAAAAGTTTAACGGCTGACATGGGAAT	Integrated DNA Technologies, Inc.
Δ*rmuC*::KAN^R	CAGGAAATGCCTTTCCAACTGGACGTTTGTACAGCACAATTCTATTTTGTGCGGGTAAGTTGTTGCGTCAGGAGGCGTTGTGTAGGCTGGAGCTGCTTC	TCAAAAAATTGTTCCAGAAGTGTAACAGATTGGGCGTCGATGCCCTAGATTTCTACCCGGCTTAACTACTCCCAATGGGTTAACGGCTGACATGGGAAT	Integrated DNA Technologies, Inc.
Δ*polB*::KAN^R	CAGGCTATACTCAAGCCTGGTTTTTTGATGGATTTTCAGCGTGTAGGCTGGAGCTGCTTC	TCACGCATCAAAATGGTATCTGGCGAACTCTTTTTTTTGCTTAACGGCTGACATGGGAAT	Integrated DNA Technologies, Inc.
Δ*dinB*::KAN^R	ACGCGTTAAATGCTGAATCTTTACGCATTTCTCAAACCCTGAAATCACTGTATACTTTACCAGTGTTGAGAGGTGAGCAGTGTAGGCTGGAGCTGCTTC	CCGATTTTTCAGCGAGAATTCGATGCATACAGTGATACCCTCATAATAATGCACACCAGAATATACATAATAGTATACATTAACGGCTGACATGGGAAT	Integrated DNA Technologies, Inc.
Δ*sbmC*::KAN^R	CAACTATACTGTATATAAAAACAGTATCAATGGAGGCGTCGTGTAGGCTGGAGCTGCTTC	AAAGAGTGGTCATCGCGTTAACACACCGCCCTGAGATGAATTAACGGCTGACATGGGAAT	Integrated DNA Technologies, Inc.
Δ*ybfE*::KAN^R	AGAGGCCTGGCTGATTGTTTCCCCCGAAGTCACCAAGATCGTGTAGGCTGGAGCTGCTTC	ACGAGCAGGACTGCACACTGTGCTACATGAAAGTGGAAATTTAACGGCTGACATGGGAAT	Integrated DNA Technologies, Inc.
Δ*recB*::KAN^R	AGCGCGTTGCAGCAAACAATGCCCCTGATGAGTGAAAAGAGTGTAGGCTGGAGCTGCTTC	GCGGGCGTAGCTGTTTGTGCTCCACAGCTTCCAGTAATTGCTTTTGCAATTTCATTAACGGCTGACATGGGAAT	Integrated DNA Technologies, Inc.
Δ*ruvC*::KAN^R	CTCTGATGAGGCCTGCTAAACAGCAAAACGGAGACGCGTGGTGTAGGCTGGAGCTGCTTC	GTGGAAACGCCTCAGCCGGAACTGACCGAGGCGGTATAACTTA	Integrated DNA Technologies, Inc.

		ACGGCTGACATGGGAAT	
Δ*umuC*::KAN^R	TCTTTGGTGTGGTGATCCACGTCGTTAAGGCGATGCGCTGGTGTAGGCTGGAGCTGCTTC	TCGGCGCTCCTGCGGGAGCGCTTTTTCCTGCCGCTATATTTAACGGCTGACATGGGAAT	Integrated DNA Technologies, Inc.
Δ*umuD*::KAN^R	GAACAGACTACTGTATATAAAAACAGTATAACTTCAGGCAGATTATTGTGTAGGCTGGAGCTGCTTC	CAGCTGGCATAAAACGCGTTACATCACAGAGGGCAAACATTTAACGGCTGACATGGGAAT	Integrated DNA Technologies, Inc.
Δ*katE*::KAN^R	ACAGCGGCCCTTTCAGTAATAAATTAAGGAGACGAGTTCAGTGTAGGCTGGAGCTGCTTC	ATGTAAATCATTTGAGGCGGCGCAATTGCGCCGCCTCCCATTAACGGCTGACATGGGAAT	Integrated DNA Technologies, Inc.
Δ*uvrA*::KAN^R	ATGCCACCGGGCAAAAAAGCGTTAATCCGGGAAAGGTGAGTGTAGGCTGGAGCTGCTTC	CTCTGAAAGGAAAAGGCCGCTCAGAAAGCGGCCTTAACGATTAACGGCTGACATGGGAAT	Integrated DNA Technologies, Inc.
Δ*mutY*::KAN^R	TGCTGCAATCTTGCCCCCAACAACAGTGAATTCGGTGACCGTGTAGGCTGGAGCTGCTTC	TCGTTCTGCTCATAAATCATCCTCTTTATCGACTCACGCGTTAACGCTGACATGGGAAT	Integrated DNA Technologies, Inc.

Oligonucleotides to verify gene deletions					
Mutation	**External Forward Primer (5' to 3')**	**External Reverse Primer (5' to 3')**	**Internal Forward Primer (5' to 3')**	**Internal Reverse Primer (5' to 3')**	**Source**
Δ*recN*::KAN^R	GATTCGTCGCTGTGATTACCATC	GCTCTTCGTCCAGATCATCCT	GCACAACTGACCATCAGCAA	CAGTTGCTGCTGTTCTTCCAGTAG	Integrated DNA Technologies, Inc.
Δ*rmuC*::KAN^R	AAAGCCATGCGGTGAAAATC	GCTCTTCGTCCAGATCATCCT	GTTATTGCGTTGGTGGGTGT	GGGTTCAACGGGAATAAACA	Integrated DNA Technologies, Inc.
Δ*polB*::KAN^R	GTCAGTTAGCGCCGCAGTTA	GCTCTTCGTCCAGATCATCCT	GCGCAGGCAGGTTTTATCTT	GGTTTATCTTCGGCGAAACG	Integrated DNA Technologies, Inc.
Δ*dinB*::KAN^R	GACCAAAAGTGCGTCCGATA	GCTCTTCGTCCAGATCATCCT	GTGGATATGGACTGCTTTTTCG	GGCGTTCATCCCAGGTTTTA	Integrated DNA Technologies, Inc.

Δ*sbmC*::KAN^R	GGTTACCC TGGAATCT GTCG	GCTCTTCG TCCAGATC ATCCT	GAAGAGA AACGTACC GTTGCAG	GCTGCAC CGCAACA TACATT	Integrated DNA Technologies, Inc.
Δ*ybfE*::KAN^R	TACGCGCT ATCCGTCG CTAT	GCTCTTCG TCCAGATC ATCCT	AACAAAC GGACCGT ACGACA	GCCAGTT GCTGCAT TAACATC T	Integrated DNA Technologies, Inc.
Δ*recB*::KAN^R	TTCTTCCA TCAGGCG GTGGT	GCTCTTCG TCCAGATC ATCCT	ATCCACTG TACGAAC GCCTG	AAATTCG GTACTGC TGGGGG	Integrated DNA Technologies, Inc.
Δ*ruvC*::KAN^R	GAAACCG CACCGAA ACTGAT	GCTCTTCG TCCAGATC ATCCT	GGCTATTA TTCTCGGC ATTGATCC	AACGTGG CAGTGGG TGATAG	Integrated DNA Technologies, Inc.
Δ*umuC*::KAN^R	CTGTTGAC GGCGAGT TTACG	GCTCTTCG TCCAGATC ATCCT	AGCTGTG AGACGGT GTTTCG	GATCGCG TAGCAGC GTTAAT	Integrated DNA Technologies, Inc.
Δ*umuD*::KAN^R	TTTTCATT TCTGCTGG ATGAGC	GCTCTTCG TCCAGATC ATCCT	TCTTGTTC AGTGTGG CTTTCC	TGGATCA CCACACC AAAGACA	Integrated DNA Technologies, Inc.
Δ*katE*::KAN^R	GTTTAGCC GATTTAGC CCCTG	GCTCTTCG TCCAGATC ATCCT	ATCAGCCG CTCACGGT TATT	TATGGTA AGGGCA GGTCGGA	Integrated DNA Technologies, Inc.
Δ*uvrA*::KAN^R	F_Ex_sodA: CACAACA CTCCGGGT AATGC	GCTCTTCG TCCAGATC ATCCT	TGAATCCC TTTCCGCC TACG	ACGGATG ACGACGA ATGGAG	Integrated DNA Technologies, Inc.
Δ*mutY*::KAN^R	CAAATTCC GGTGAAAT GACG	GCTCTTCG TCCAGATC ATCCT	GCCCAGG TTCTGGAC TGGTA	CGACACG GGAAGCC ACATAG	Integrated DNA Technologies, Inc.

1074
1075

1076

1077

1078

1079

1080