UV-B-Induced DNA Repair Mechanisms and Their Effects on Mutagenesis and Culturability in *Escherichia coli*

Sreyashi Ghosh, Jenet Narzary, and Mehmet A. Orman*

William A. Brookshire Department of Chemical and Biomolecular Engineering, University of Houston, Houston, TX, USA

*Correspondence to: S222 Engineering Bldg 1, 4226 Martin Luther King Boulevard, Houston, TX 77204-4004, Phone: 713-743-6785, Email: morman@central.uh.edu

Abstract

Ultraviolet-B (UV-B) radiation, intensified by ozone depletion, induces DNA damage and promotes mutagenesis, shaping evolution. While UV-induced SOS responses are well characterized in bacteria, the cellular consequences of prolonged UV-B exposure remain less clear. Prolonged UV-B exposure disrupts translation and RecA-mediated SOS induction without major changes in membrane permeability or reactive oxygen species. This impairs mutagenesis and induces a reversible loss of culturability. Genetic analysis reveals both redundant and differential roles for DNA repair pathways: homologous recombination (RecA, RecB), nucleotide excision repair (UvrA), and translesion synthesis (UmuC/D) are essential for maintaining mutagenesis and culturability, while others (RecN, RmuC) have limited impact. Notably, deletion of UvrD (a repair-associated helicase) intensifies transient non-culturability without affecting mutagenesis, underscoring complexity in repair networks. Overall, our findings reveal a dose-dependent trade-off: moderate UV-B promotes mutagenesis with minimal viability loss, whereas prolonged exposure suppresses mutagenesis via transient dormancy, reflecting an adaptive strategy with significant evolutionary implications.

Introduction

Mutagenic processes are key to evolutionary progress [1–3], as they generate genetic diversity. This is essential for organisms to adapt to and endure in dynamic settings that are continuously changing. Understanding the mechanisms underlying mutagenesis helps us better comprehend the evolutionary forces shaping organisms [4,5]. This knowledge also aids in developing more potent and effective medical treatments to combat many diseases, given that pathogenic microorganisms and tumorigenic cells develop resistance through mutagenesis [6–9].

Ultraviolet (UV) radiation, particularly UV-B, plays a dual role in Earth's ecosystems; it acts as both a developmental cue and an environmental stressor. While moderate UV-B levels may regulate processes like plant defense and microbial interactions, elevated UV-B due to ozone depletion has been linked to ecological disruptions and even mass extinction events, such as the end-Permian and Devonian–Carboniferous extinctions [10–13]. UV-B is of particular interest because it is thought to reach the Earth's surface in greater intensity during periods of severe ozone depletion caused by catastrophic events. On a microbial scale, UV radiation has continually shaped the trajectory of evolution over billions of years, from contributing to the origins of life to driving

genetic variation and adaptation[14,15]. This is because UV radiation directly damages DNA and
induces certain repair mechanisms that are highly mutagenic[16–19]. The UV light spectrum, ranging
from 100 to 400 nanometers (nm), can be categorized into four groups: UV-A (315-400 nm), UV-
B (280-315 nm), UV-C (100-280 nm), and vacuum-UV (100-200 nm)[20]. UV-C is more readily
absorbed by nucleic acids compared to UV-B and UV-A[21]. When bacteria are exposed to UV
radiation, [6,7] it triggers the formation of pyrimidine dimers and other DNA lesions that disrupt the
normal structure and function of DNA molecules[22,23]. These UV lesions are removed by various
effective bacterial systems, such as the photoreactivation repair system (i.e. dimer monomerization
under visible light by photolyase enzymes, such as Phr)[24], base excision repair (BER) (removal
of cyclobutane-thymine dimers using DNA glycosylases and AP endonucleases)[25], and the
primary pathway for removal of bulky DNA lesions, i.e. the nucleotide excision repair (NER) (for
rapid repair of photo-lesions using the *uvr* genes, *polA* and ligase enzyme) [26]. However, an
excessive amount of dimers that cannot be removed by the repair systems can accumulate and
interfere with the cellular replication process, resulting in the formation of "secondary" lesions in
the form of ssDNA fragments[27]. Therefore, to deal with this severe DNA damage, a post-
replication damage repair mechanism known as the bacterial SOS response pathway is induced,
preventing premature cell division [16,28,29] and providing cells with sufficient time to repair the
damaged DNA[18,30–32]. This SOS response in bacteria is controlled by the multifunctional RecA
protein, a key component that cleaves the transcriptional repressor LexA[33–35]. This cleavage
initiates the expression of more than 40 SOS genes. RecA, involved in the recombination repair of
ssDNA gaps[36], helps provide an error-free damage removal route. However, excessive DNA
damage leads to the induction of the mutagenic phase of the SOS response, mediated by bacterial
DNA polymerase enzymes, including Pol V [37,38], which is encoded by the *umuD* and *umuC* genes
[39]. Pol V can bypass template lesions during DNA replication through a process known as
translesion DNA synthesis (TLS) [40]. Two additional TLS DNA polymerases, Pol II (encoded by
polB) and Pol IV (encoded by *dinB*), facilitate replication past blocking lesions, albeit at the cost
of introducing mutations. Another DNA damage repair strategy is the mismatch repair (MMR)
pathway [41], which recognizes base-base mismatches and insertion/deletion loops, including those
introduced by DNA polymerases. However, with increased mutagenic stress, the efficiency of
MMR decreases, as observed in the case of fluoroquinolone treatment [42]. Such stress can
potentially activate the SOS response and be addressed by SOS-mediated repair mechanisms, as
the accumulation of DNA lesions and stalled replication forks are known to trigger this global
DNA damage response.

UV radiation can serve as a valuable experimental tool for investigating mutagenesis because it
induces a wide range of mutations without an apparent sequence preference[43,44]. Additionally, UV
radiation is of great interest to scientists for its applications in biotechnology, particularly in
directed evolutionary strategies to engineer proteins or organisms[45,46]. Moreover, there is growing
interest in using UV radiation as a disinfectant, including for sterilizing air, equipment, and
surfaces; reducing the transmission of airborne diseases; and even treating wound infections[47,48].
Despite their increased applications in clinical settings, we believe that two phenomena, UV-
induced bacterial cell dormancy and mutagenesis, need to be fully understood. Bacterial cells are
highly heterogeneous, and they can enter a growth-arrested state through stochastic mechanisms
or environmental factors[49], including UV treatment, which can make these cells highly tolerant.
Also, the SOS-response-mediated cell dormancy not only promotes cell survival but also sustains
mutagenic processes, as dormant cells may still harbor DNA damage[50,51]. Additionally, UV

radiation itself is highly mutagenic; it can accelerate the emergence of more resilient mutants,
which could pose a serious threat to public health.

UV-induced repair mechanisms are highly complex as UV can damage many cellular components,
including DNA, RNA, lipids, and proteins[52,53]. While the role of the SOS response in mutagenesis
has been extensively studied (refer to the following review articles[16,30,32] for more details), our
understanding of the downstream repair mechanisms and their contributions to cell survival,
culturability, and mutagenesis remains limited. Although our recent study showed an almost
perfect correlation between RecA levels and mutation frequency in *E. coli* following UV-B
treatment, prolonged UV-B exposure was found to impair SOS-mediated mutagenesis and induce
a transient non-culturability state[54]. While this may represent a potential survival strategy for cells
to evade UV-B-induced mutagenesis, the underlying mechanisms remain unclear. Our current
study aims to investigate these complex mechanisms further, providing new insights into the
molecular processes underlying mutagenesis and dormancy.

Results and Discussion

The transient unculturability is largely driven by the global regulator RecA, crucial for the SOS
response and recovery

To study UV-mediated mutagenesis and the SOS response, we used our previously established
methodology[54]. We cultured *E. coli* MG1655 wild-type cells in 2 mL Lysogeny Broth (LB)
medium in test tubes until they reached the mid-exponential phase with an optical density (OD_{600})
of ~0.5. Subsequently, the cells were transferred to petri dishes, forming a thin film of cultures that
increased the surface area for UV exposure. This film was then subjected to UV-B light (302 nm
thin-line transilluminator, UVP ChemStudio, Analytik Jena) for various durations: 2, 4, 8, 16, 24,
and 32 minutes (min), ensuring a wide range of UV intensity from 120 J/m^2 to as high as 1920
J/m^2 as shown in **Fig 1a** (see **Supplementary Fig. S1,** and **Materials and Methods**). We used a
broad UV-B dosage range to capture dose-dependent effects, as UV-B is less energetic than UV-
C and requires higher exposure to induce cellular stress. This approach aligns with prior studies
using UV-B, where treatment durations and doses vary widely depending on strain and
experimental conditions[55–59]. Following exposure, the cells in LB were promptly transferred back
to test tubes and cultured in a shaker for a 24-hour recovery (refer to the Materials and Methods
section for details). Cultures that did not receive UV-B treatment served as controls. During
recovery, colony-forming units (CFU) were quantified at the indicated time points for each
condition, as shown in **Fig. 1b**. At the beginning of recovery (t~0), CFU levels decreased in
cultures exposed to longer UV-B treatments. A 4-8 min exposure caused a 10-fold reduction, while
16 min led to a ~100-fold drop compared to the control. Exposure times of 24 and 32 min resulted
in a ~10,000-fold reduction (**Fig. 1b**). After 15 minutes, CFU levels notably increased in cultures
with 24 and 32 min of UV-B exposure, likely due to mechanisms making cells temporarily
unculturable. This drastic increase in CFU levels is not due to cell division, given that the *E. coli*
doubling time is about 20-25 minutes, which is too short for significant growth within that period.
By the 24-hour mark, all conditions showed similar CFU levels (**Fig. 1b**).

To measure mutagenesis, we quantified rifampicin (RIF)-resistant colonies after 24 hours of
recovery. Cultures were plated on LB agar with 500 µg/ml RIF, and colony counts were reported
per 10^8 cells (**Fig. 1c**). Although this method does not directly measure mutation levels, it is widely
used [44,60–62] due to the strong correlation between mutagenesis and resistant colony formation [63,64].

We chose a 24-hour recovery period to ensure accurate colony counts, consistent with previous
studies[60,65,66], as mutation levels generally plateau by this time[54]. Our results showed that
mutagenesis increased with UV-B exposure, peaking at 16 minutes, but longer exposures (24-32
min) significantly reduced RIF-resistant colonies (**Fig. 1c**). This pattern, consistent with our
previous findings[54], suggests a link between UV-B exposure, mutagenesis, and temporary
unculturability.

Our initial experiments were conducted without controlling for light exposure. Given the role of
photoreactivation in bacterial recovery, we investigated whether the observed transient loss and
restoration of culturability were driven by photolyase-mediated repair. The *phr* gene encodes DNA
photolyase, which reverses UV-induced cyclobutane pyrimidine dimers in the presence of blue
light[67,68]. As we were unable to obtain viable Δ*phr* knockouts in *E. coli* MG1655 using the λ-Red
recombinase technique (see Materials and Methods)[69], a result consistent with a previous report [70],
we evaluated the role of photoreactivation by repeating the UV-B exposure and recovery
experiments under strict dark conditions. Importantly, despite minor fluctuations in mutation
frequency, expected due to altered photolyase-mediated repair mechanisms, we observed the same
core features under dark conditions: a pronounced but transient decline in culturability, followed
by recovery, and a dose-dependent mutation frequency (**Fig. 1d, e**). To directly test whether *phr*
function is necessary for these outcomes, we conducted parallel experiments using a Δ*phr* strain
from the Keio collection (*E. coli* BW25113 Δ*phr*), as this strain can have a viable knockout *phr*
strain. Upon UV-B treatment and recovery, this strain exhibited a comparable transient
unculturability and dose-dependent mutation frequency (**Supplementary Fig. S2**), mirroring the
responses seen in MG1655. While some quantitative differences were observed in BW25113
(likely due to the distinct genetic background), these results indicate that the observed phenomena
were not solely contingent on light-driven DNA repair. This is plausible, as UV-induced mutations
likely arise from error-prone repair mechanisms rather than error-free photoreactivation by Phr.

In our previous study, we found that excessive UV-B treatment (24 and 32 minutes) impaired the
SOS response, including the expression level of the RecA protein[54]. Therefore, the observed
transient unculturability is likely more complex and may involve various mechanisms, given that
the SOS response is intricate, involving multiple proteins, with RecA serving as the global
regulator. We hypothesized that if this transient unculturability is linked to the impaired SOS
response, then genetic perturbation of this response through *recA* deletion would further decrease
cell culturability. Our results verified that the knockout strain of *recA* exhibited a drastic reduction
in culturability immediately after UV treatment across all conditions (**Fig. 1f**). Initially, CFU levels
in the Δ*recA* strain were below the limit of detection (1 CFU) for most UV treatment conditions;
however, this was transient, as we observed a rapid increase in CFU levels (from 1 to 10^7) within
15 minutes of recovery (**Fig. 1f**). This observation is highly interesting, and to our knowledge,
such a dramatic recovery pattern in a *recA*-deficient background has not been previously reported.
Although liquid-holding recovery has been well documented since the 1960s (primarily involving
excision repair and photoreactivation during non-nutritive incubation in wild-type strains)[71,72], our
findings are distinct, given that the marked, transient increase in culturability following UV-B
treatment occurs in a genetic background lacking a functional SOS response. While we previously
demonstrated that prolonged UV-B exposure impairs the RecA-mediated SOS response and that
recA deletion completely eliminates mutagenesis across all exposure durations (and thus did not
repeat those experiments here)[54], it remains unclear how downstream processes governed by RecA
influence cell culturability and mutagenesis. We therefore aim to investigate these aspects in the
following sections of this study.

Genes significantly expressed during the SOS response have a moderate or negligible effect on cell culturability and mutagenesis

To identify the downstream mechanisms, we screened a subset of the *E. coli* promoter library containing 1900 promoter reporters. In this library, the promoters of *E. coli* genes are fused to a fast-folding green fluorescent protein (GFP) gene in low-copy-number plasmids, allowing accurate and reproducible measurement of gene expression[73]. We focused on approximately 40 reporters containing promoters that regulate the key SOS response and associated genes. These promoters are under the control of the RecA-LexA regulatory system, chosen based on a comprehensive analysis of *E. coli* databases[74]. In our experiments, *E. coli* MG1655 cells containing the selected promoters were cultured individually until the mid-exponential phase in 96-well plates. Subsequently, they were exposed to UV-B radiation for 16 min. This duration was chosen as it induces maximum SOS upregulation and mutagenesis without drastically impacting cell culturability (**Fig. 1b,c**). Also, 16-min UV treatments were used as positive controls for the subsequent sections. GFP levels were measured after 24 h of recovery, allowing adequate time for UV-induced promoters to express GFP. High cell densities ensured reliable GFP measurements, given that we used a plate reader in this screening assay.

Our specific focus was on genes showing significant upregulation following UV treatment (**Fig. 2a**), as they likely play pivotal roles in the downstream mechanisms of RecA-mediated SOS response. Several promoters showed drastic upregulation, including P_{recA} and P_{lexA} (promoters of genes encoding SOS response global regulators, as expected), P_{recN} and P_{rmuC} (promoters of genes involved in DNA recombinational repair), P_{sulA} (promoter of a gene encoding cell division inhibitor), P_{polB} and P_{dinB} (promoters of genes encoding DNA polymerase enzymes), P_{ftsK} (promoter of a gene encoding essential cell division protein), P_{sbmC} (promoters of a gene encoding inhibitor of DNA gyrase-mediated DNA supercoiling), and P_{ybfE} (promoter of a *lexA*-regulated gene whose function is not well characterized)[75] (**Fig.2a**). To determine whether the genes associated with the identified promoters are involved in SOS response-mediated culturability and mutagenesis, we generated single deletions in *E. coli* MG1655 for these genes (except for *lexA* and *ftsK*, as they are essential and could not be deleted[76–78]). The knockout strains were exposed to UV-B for 16 min or 32 min or untreated, and their CFU profiles during recovery, as well as their RIF-resistant colony levels, were similarly measured. For all knockout strains, the temporal CFU profiles remained the same as that of the wild-type, showing a transient non-culturability during the first 15 min of recovery after a high UV-B exposure duration of 32 min (**Fig. 2b, Supplementary Fig. S3**). As expected, the deletion of *recA* completely eliminated mutagenesis (**Fig. 2c**). Although 32 min of UV-B exposure significantly reduced mutagenesis in all strains, including wild type, the two DNA recombination gene knockouts, Δ*recN* and Δ*rmuC*, showed a 3-fold and 1.5-fold decrease in RIF-resistant mutant levels, respectively, compared to WT cells following 16 min of UV-B treatment (**Fig. 2c**). Surprisingly, other single deletion strains, including those for the genes encoding DNA polymerase II and IV, *dinB* and *polB*, did not show significant changes (**Fig. 2c**).

Overall, our results showed that single deletions of the downstream genes of RecA did not drastically impact cellular culturability and mutagenesis, except for RecN and RmuC deletions, which slightly reduced mutagenesis. RecN is involved in the recombinational repair of DNA double-strand breaks, and its mutation makes cells sensitive to mitomycin C and ionizing radiation[79–81]. Although RmuC is thought to be an inner membrane protein with a nuclease domain[82,83], its function is not well understood, and our data indicate it only moderately impacts

UV-B-mediated mutagenesis. It is quite surprising that the deletion of DNA polymerase II and IV,
two key proteins involved in indirect mutagenesis[84], did not impact UV-B-induced mutagenesis.
As mutagenesis is a multifaceted process regulated by multiple mechanisms, testing multi-deletion
strains might be necessary to show the complex interplay between these mechanisms. Furthermore,
deleting a single polymerase enzyme may have minimal impact due to the abundance of the other
polymerase enzyme. Therefore, we first generated ΔrecNΔrmuC and showed that the double
deletion resulted in a cumulative decrease (~5 fold) in RIF-resistant colony levels compared to the
wild type following 16 min of UV-B treatment (**Fig. 2c**). To assess the impact of DNA polymerase
II and IV, we generated a ΔpolBΔdinB mutant strain, which did not affect RIF-resistant colony
levels (**Fig. 2c**). Additionally, we generated ΔrecNΔpolB and ΔrecNΔdinB double knockout strains
and demonstrated that they exhibited approximately a 5-fold reduction in RIF-resistant colony
levels compared to wild-type cells (**Fig. 2c**). Deletion of DNA polymerase genes *polB* and *dinB*
from ΔrecNΔrmuC individually did not show significant changes compared to the ΔrecNΔrmuC
strain in RIF-resistant mutant levels (**Fig. 2c**), suggesting that the observed reduction in RIF-
resistant colonies in multi-deletion strains is primarily due to the effects of *recN* and/or *rmuC*.

Screening the knockout strains reveals the redundancy of repair mechanisms in UV-induced
mutagenesis and culturability

Although our initial screening yielded intriguing results, the critical downstream mechanisms
underlying our observed phenomenon still remain elusive. Thus, we conducted a second screening
using the *E. coli* BW25113 Keio knockout collection to assess the impact of UV on mutagenesis
and culturability of the knockout strains. This screening focused on all potential repair mechanisms
involved[16], particularly (i) those that are not regulated by RecA (such as, MMR genes) and (ii)
those that are regulated by RecA but whose expression was not significantly upregulated in our
initial screening (**Fig. 2a**). For this second screening, the knockout strains in 96-well plates were
treated with UV-B for 16 min during mid-exponential-phase growth, and RIF-resistant colonies
were quantified after a 24-hour recovery period. We identified several genes whose deletion
resulted in a significant reduction in RIF-resistant colony formation (**Fig. 3a**), including *recB*, a
component of Exonuclease V or the RecBCD complex that promotes homologous recombination
in the repair of double-strand DNA breaks[85,86]; *umuC* and *umuD*, encoding error-prone DNA
polymerase V[87]; *ruvC*, encoding an endonuclease that binds to and cleaves Holliday junctions[88];
and *katE*, encoding *E. coli* catalase enzyme[89]. Additionally, we observed several genes that showed
an increase in mutagenesis, such as *uvrD* and *uvrA*, which are involved in the nucleotide excision
repair pathway[90,91], and *mutY*, involved in the base excision repair pathway[92] (**Fig. 3a**). Knockouts
of MMR genes *(mutS, mutL, mutH,* and *exoI)*, which function in replication error correction, were
included in our screen but did not show significant reductions in mutagenesis, indicating MMR
may play a minimal role under our UV-B exposure conditions (**Fig. 3a**).

To further investigate genes that showed the most pronounced impact on UV-induced mutagenesis
in our screening, and to see if similar trends occur in other *E. coli* strains, we deleted these genes
in *E. coli* MG1655 and analyzed the knockout strains after 16 and 32 min of UV-B exposure.
Deletion of *recB*, *ruvC*, *umuC*, and *umuD* significantly reduced mutagenesis (**Fig. 3c**), consistent
with our screening data. The absence of UmuC and UmuD proteins, specialized for translesion
synthesis, was shown to sensitize *E. coli* cells to UV[93,94], aligning with our findings. While ΔuvrA
in the *E. coli* BW25113 strain from screening showed higher mutant levels compared to wild-type
cells, its deletion in the MG1655 background resulted in a significant reduction in mutant
formation (**Fig. 3c**), suggesting that the role of certain genes in mutagenesis can vary depending

on the genetic background of the *E. coli* strain. Also, our results indicate that KatE, UvrD, and
MutY may not be directly involved in the mutagenic response to UV radiation under the conditions
tested here, as their deletions did not clearly impact UV-B-induced mutagenesis in *E. coli* MG1655
(**Fig. 3c**). Interestingly, Δ*uvrD* and Δ*mutY* strains exhibited increased RIF-resistant colonies in
cultures not treated with UV-B (**Fig. 3c**). UvrD is involved in various DNA repair pathways,
including NER and MMR processes[41,95,96]. MutY is a glycosylase enzyme that corrects adenine
mismatches resulting from DNA replication errors, primarily in the BER pathway[92]. While the
specific roles of UvrD and MutY in UV-B-induced mutagenesis are not evident in this study, the
increased RIF-resistant colonies in untreated cultures might highlight their potential role in
maintaining genomic stability under normal growth conditions, possibly by mitigating
spontaneous mutations.

The temporal CFU profiles revealed interesting trends as well. Δ*ruvC*, Δ*umuC*, and Δ*umuD* strains
showed no significant change in transient non-culturability immediately after UV-B treatment
compared to the wild type (**Fig. 3b, Supplementary Fig. S4**). However, Δ*recB* exhibited a notable
decrease in CFU levels, while Δ*uvrA* and Δ*uvrD* mutants showed even more drastic reductions
following UV-B treatment compared to the wild type (**Fig. 3c**). This reduction in CFU levels was
transient, as their levels increased approximately 10^7 -fold within 15 min of recovery. UvrA and
UvrD together form key components of the NER pathway[95]. UvrA forms a complex with UvrB to
detect DNA lesions, while UvrD functions as a helicase to unwind and remove damaged DNA
segments[96,97]. As both UvrA and UvrD are regulated by the RecA protein[32], it is not surprising that
the deletion of *recA* resulted in drastic reductions in CFU levels following UV treatment (**Fig. 1f**).
Since RecA is essential for inducing the SOS response, its absence means that many DNA repair
genes, including *uvrA* and *uvrD*, are not upregulated in response to UV-induced DNA damage.

Excessive UV exposure impairs cellular translation processes

Although our results highlight specific SOS proteins involved in mutagenesis and cell culturability,
it remains unclear why excessive UV exposure impairs *recA* expression and, consequently, the
SOS response. This impairment could be due to the activation of cellular mechanisms, such as
growth arrest proteins, which may help limit excessive DNA damage and mutagenesis.
Additionally, the production of toxic molecules, such as reactive oxygen species, or the inhibition
of certain cellular functions, like translation, may also contribute to this effect. These
mechanisms—namely the involvement of growth arrest proteins, reactive oxygen species, or the
inhibition of translation—will be thoroughly explored in the subsequent section of our study.

Within the SOS response, two genes, *sulA* and *tisB*, play key roles in cell growth inhibition. The
sulA gene in *E. coli* encodes a cell-division inhibitor that temporarily halts the cell cycle during
the SOS response, preventing damaged DNA from prematurely segregating into daughter cells[98].
On the other hand, *tisB* encodes a toxin protein that reduces cellular ATP levels[50,99]. This protein
may also induce reversible dormancy by suppressing cell metabolism. To investigate the individual
and combined effects of these genes, we constructed knockout strains of *E. coli* MG1655 Δ*sulA*,
Δ*tisB*, and Δ*sulA*Δ*tisB*. Similarly, these strains, at the mid-exponential phase, were exposed to
varying durations of UV-B radiation followed by a 24-hour recovery period. *E. coli* MG1655 WT
cells were used as the control group for comparison. Deleting *sulA* and *tisB* individually did not
alter the trend of transient non-culturability observed in our experiments. Specifically, the CFU
profiles initially showed a significant decrease after prolonged UV-B exposure of 24 and 32 min,
followed by a sharp increase within the first 15 min of recovery, similar to the wild-type strain
(**Fig. 1b vs Fig. 4a-c**). However, CFU levels of Δ*sulA*Δ*tisB* immediately after 24 and 32 min of

UV-B exposure (corresponding to the t~0 point on the CFU plots) were nearly 10-fold higher compared to those of the wild-type or single knockout strains (**Fig. 1b vs Fig. 4a-c**). We observed approximately a 10-fold increase in CFU levels within the first 15 min of recovery in the double knockout strain, although this increase was not as pronounced as in the control or single knockout strains (**Fig. 4a-c, Supplementary Fig. S5**). While deleting both genes enhanced the culturability of cells following UV-B treatment, the phenomenon observed in the wild-type strain was not completely eliminated in the double knockout strain, highlighting the complexity of the underlying mechanisms. No distinct profiles of RIF-resistant mutants were observed after 24 hours of recovery in either single or double-knockout strains compared to the wild-type strain (**Fig. 1c vs Fig. 4d-f**), indicating that these genes are not directly involved in mutagenesis, although they may affect cellular culturability to some extent.

UV radiation has the potential to damage cellular membranes and compromise essential cellular functions[19]. This damage can lead to increased membrane permeability, leakage of cellular contents, and, ultimately, cell death[100,101]. Additionally, prolonged exposure to UV radiation can trigger excessive production of reactive oxygen species (ROS) within cells[102–104]. ROS are highly reactive molecules that can cause oxidative damage to cellular components such as DNA, proteins, and lipids, impairing cellular functions. It is possible that the 24-minute and 32-minute UV-B treatments studied here may not be severe enough to kill the bacterial cells completely, but they are likely sufficient to hinder cell growth and mutagenic processes by disrupting vital metabolic processes. To characterize cellular membrane permeability, cells treated with UV-B were immediately collected and incubated with propidium iodide (PI), a fluorescent dye that enters cells with compromised membranes. Flow cytometry data indicated no significant increase in permeabilization following prolonged UV-B treatment (32 min) up to 1 hour of recovery (**Fig. 5a**), compared to both untreated cells and moderately UV-treated cells (16 min), which did not markedly reduce CFU levels. Furthermore, when UV-treated samples were subjected to Amplex Red reagent (see Materials and Methods for details) to detect hydrogen peroxide (H_2O_2), a primary reactive oxygen species, no significant changes were observed in the 32-minute UV-treated cells compared to untreated or 16-minute UV-treated cells (**Fig. 5b**), indicating H_2O_2 may not be a key player. In our previous study, we found that ROS significantly affects cell culturability only when their concentrations are much higher than physiological levels, and their impact on cell culturability was observed to be non-transient [54].

UV exposure is known to interfere with ribosomal function by inducing crosslinks in ribosomal RNAs[105], which can inhibit overall translation processes and impair the expression of SOS response genes. To assess the effects of UV treatments on cellular translation, we analyzed GFP expression under the control of the *recA* promoter (P_{recA}) at the single-cell level using *E. coli* MG1655 cells carrying the pUA66 P_{recA}-*gfp* plasmids. The cells were subjected to UV-B treatments for 16 min and 32 min, and *recA* expression was measured using flow cytometry during the early recovery phase (**Fig. 5c**). UV-B treatment should activate promoters of SOS response genes and induce recA expression. As expected, the 16-min UV-B treatment resulted in significantly higher *recA* expression compared to untreated cells (**Fig. 5c, e**). However, the 32-min UV-B exposure resulted in more heterogeneous and lower *recA* expression compared to the 16-min UV-B exposure (**Fig. 5c, e**), potentially implying an impairment in the translational process. This difference between the 16- and 32-min treatments cannot be solely attributed to increased cell death following 32 min of UV-B exposure. Despite the observed transient culturability in the prolonged UV-exposure culture, CFU levels in both conditions are nearly similar after the brief transient period, around 15 minutes of recovery culture (see the 0.25-hour time point in **Fig. 1b**).

To substantiate the hypothesis regarding the impairment of the translational process, we employed a GFP-expressing plasmid where the GFP gene is tightly regulated by an isopropyl β-D-1-thiogalactopyranoside (IPTG)-inducible T5 promoter and a strong LacI[q] repressor[106]. Our rationale was that if excessive UV-B treatment indeed impairs translational processes, the levels of GFP expression in cells treated with 32 min of UV-B should be lower compared to those in the no-UV control or 16-minute UV-treated cells, despite the presence of the inducer. We supplemented cell cultures with 0.1 mM IPTG immediately before UV-B treatments. As anticipated, the 32-min UV-B treatment resulted in a lower level of GFP expression compared to the no-UV control or the 16-minute UV-B treatment (**Fig. 5d, f**). However, the 16-min UV-B treatment exhibited the highest GFP expression levels (**Fig. 5d, f**), suggesting that moderate UV treatment (16 min) enhances translational activity, whereas excessive UV-B treatment (32 min) slightly inhibits it. This moderate inhibitory effect of excessive UV-B treatment on cellular translation, including RecA expression, likely impacts overall cellular processes and compromises the SOS response. This was further validated when we measured the expression levels of *uvrD* and *uvrA*—two critical genes involved in mutagenesis and culturability. Their expression levels were slightly reduced under 32 minutes of UV treatment compared to 16 minutes (**Supplementary Fig. S6a-d**).

To further validate the role of RecA-mediated SOS response in recovery and mutagenesis following UV-B exposure, particularly under conditions where transcription and translation are impaired, we constructed an IPTG-inducible *recA* expression system. This was achieved by cloning *recA* under the control of a T5 promoter (**Supplementary Fig. S7**) and introducing the plasmid into the *E. coli* MG1655 Δ*recA* background. In the absence of IPTG, the strain behaves as a *recA* knockout, showing impaired recovery and loss of UV-induced mutagenesis (**Fig. 5g**). To ensure that RecA protein was present at the time of UV treatment, IPTG was added approximately 30 minutes prior to exposure. This timing was chosen to avoid possible interference from UV-induced translation defects that could impair *recA* expression if induction occurred during or after treatment. Furthermore, we tested a range of IPTG concentrations to evaluate whether RecA-dependent outcomes were dose-responsive, as suggested by our data here and in our previous study [54]. As shown in **Fig. 5g**, increasing IPTG levels led to a corresponding restoration of both CFU levels and mutation frequency following 16- and 32-minute UV-B exposures, indicating a clear correlation between RecA expression and the recovery of culturability and mutagenesis. These findings suggest that translational capacity directly impacts SOS activation by limiting RecA availability.

Altogether, our findings suggest that while prolonged UV-B treatments may not directly impact membrane integrity or ROS levels, the observed unculturability of the cells likely results from a combined effect of multiple factors, including cell growth inhibitors, impaired translation processes, and downstream repair pathways. The impairment of cellular processes such as translation may hinder the SOS response, as evidenced by the decreased expression of RecA in cells treated with 32 min of UV-B (**Fig. 5e**). This interpretation is further supported by our findings with the Δ*recA* strain, which exhibited near-complete unculturability immediately after UV treatments. The immediate reduction in CFU levels in the knockout strains (Δ*recA*, Δ*uvrA* and Δ*uvrD*), as well as in the wild-type strain exposed to prolonged UV, highlights the importance of the SOS response in the initial phases of DNA damage repair. The transient nature of CFU reduction in all these mutants also suggests that *E. coli* possesses redundant or numerous repair mechanisms[16] that can partially restore culturability even when certain DNA repair pathways are disrupted.

Conclusions

In this study, we investigated the effects of UV-B treatments on *E. coli* mutagenesis and culturability and provided important insights into bacterial survival mechanisms under stress. Mutagenesis, measured by RIF-resistant colony formation, peaked at 16 min of UV-B exposure but decreased with longer exposures (24 and 32 min). The longer UV-B exposures also led to a significant decrease in CFU levels, but this phenomenon was observed to be a transient state, as we saw a drastic jump in CFU levels after a short recovery period. We also explored the role of key SOS response genes and demonstrated that knockout strains lacking *sulA* and *tisB* genes exhibited similar patterns of transient non-culturability as wild-type cells, although double knockouts exhibited higher initial CFU levels post-exposure, indicating a partial role in culturability. Further investigation revealed that excessive UV exposure did not significantly affect membrane permeability or reactive oxygen species (ROS) levels but moderately impaired translation processes, as evidenced by reduced expression of SOS response genes like *recA* or IPTG-inducible fluorescent proteins. This impaired translation likely contributed to the observed reduction in mutagenesis and transient non-culturability following prolonged UV exposure, given the significant impact of *recA* deletion on cell culturability and mutagenesis. Importantly, reintroducing *recA* expression in the knockout strain restored culturability and increased mutagenesis in proportion to *recA* expression levels, further supporting the link between translational capacity, *recA*-mediated DNA repair, and the observed phenotypes.

Although our screening of promoter reporters highlighted several genes significantly upregulated following UV treatment, the knockout strains of most of these genes generally had minimal impact on culturability or mutagenesis, except for Δ*recN* and Δ*rmuC*, which moderately reduced mutagenesis. In our second screening with the Keio knockout collection, we identified several critical genes, such as *recB, umuC, umuD*, and *ruvC*, whose deletions led to decreased mutagenesis, highlighting their crucial roles in facilitating error-prone repair mechanisms during UV-B stress. Furthermore, we observed differential impacts of gene deletions on mutagenesis versus culturability, providing insights into additional functions of these genes. Particularly in strains lacking *uvrA*, or *uvrD* showed a drastic but transient reduction in their culturability following UV treatment, highlighting their role in maintaining genomic stability. Our findings also underline the abundance of repair mechanisms in *E. coli*, demonstrating that multiple independent pathways can significantly impact cell mutagenesis and culturability.

Overall, this study enhances our understanding of bacterial responses to environmental stressors, particularly UV-B radiation, by uncovering the complex interplay between DNA repair pathways, mutagenesis, and transient non-culturability. Our findings reveal a dose-dependent balance between survival and genetic variability, suggesting that bacteria modulate their repair and stress responses to optimize adaptability under UV-B exposure. This regulatory balance has important implications for understanding how UV-B, especially during periods of ozone depletion, may act as a powerful evolutionary force.

Materials and Methods

Bacterial strains and plasmids

Escherichia coli K-12 MG1655 wild type (WT), the pUA66 plasmid containing an IPTG-inducible *gfp* (green fluorescent protein gene) expression system, and the pUA66 plasmid with *gfp* under the

control of the *recA* promoter, P*recA*, were obtained from Dr. Mark P. Brynildsen at Princeton University. *E. coli* K-12 MG1655 Δ*sulA, ΔtisB, ΔrecA* was constructed in our previous studies [54,107,108]. The promoter strain collection of *E. coli* MG1655 in a 96-well plate format used for the screening assay was obtained from Horizon Discovery, Lafayette, CO, USA. The Keio knockout strain collection (a single-gene deletion library of *E. coli* K-12 BW25113) was obtained from Dharmacon Keio Collection (Dharmacon, Cat# OEC4988). Since the knockout strains carry a kanamycin resistance gene, high-throughput screening was performed in the presence of kanamycin to prevent contamination; thus, Δ*lacI* was used as the reference control, given that the parental BW25113 strain lacks kanamycin resistance. We selected the Δ*lacI* mutant as it is not involved in DNA repair pathways and serves as a neutral background. Furthermore, its response to UV-B exposure closely mirrors that of the MG1655 wild-type strain (**Supplementary Fig. S2a**), supporting its suitability as a functional reference for our experimental comparisons. The strains from both the promoter and knockout collections used in this study are listed in **Supplementary Table 1**. The knockout *E. coli* MG1655 strains generated for this study are detailed in **Supplementary Table 2**. The method of Datsenko and Wanner[69] was used to generate these strains and the oligonucleotides used to delete the genes are provided in **Supplementary Table 3.** We also attempted to generate a Δ*phr* mutant in *E. coli* MG1655 using the same approach; however, no viable recombinants were obtained despite testing multiple primer sets with varying homology arm lengths (40–60 bp) (**Supplementary Table 3**). An IPTG-inducible *recA* overexpression plasmid was generated using a commercial cloning service (Synbio Technologies, USA). The *recA* coding sequence from *E. coli* MG1655 was cloned downstream of a T5 promoter into a low-copy plasmid backbone (pUA66) containing a strong mutated *lacI* repressor for tight regulation and a kanamycin resistance marker[108]. The resulting plasmid was verified by sequencing and subsequently transformed into MG1655 Δ*recA* cells for inducible expression studies. The plasmid map is shown in **Supplementary Fig. S7.**

Chemicals, media, and culture conditions

Unless otherwise specified, all chemicals were procured from Fisher Scientific (Atlanta, GA), VWR International (Pittsburgh, PA), or Sigma Aldrich (St. Louis, MO). PI staining kit was purchased from Promega Corporation (Madison, WI). *E. coli* cells were cultured in liquid Lysogeny-Broth (LB) medium. LB agar medium was utilized for enumerating colony-forming units (CFU) of *E. coli*. The liquid LB medium was prepared by dissolving 5 g yeast extract, 10 g tryptone, and 10 g sodium chloride in 1 L of deionized (DI) water. LB Agar media were prepared by dissolving pre-mixed 40 g LB agar in 1 L of DI water. Both solid and liquid media were subjected to autoclaving for sterilization.

Kanamycin (50 μg/mL) was included in the liquid LB media for plasmid selection and retention. IPTG at 0.1 mM was used to induce *gfp* expression. For PI staining, sterile 0.85% sodium chloride solution was used. When necessary, cells were washed with phosphate-buffered saline (PBS, 1X). Stock solutions for rifampicin (RIF; 50 mg/mL) was prepared by dissolving in DI water using 0.01 N sodium hydroxide. The IPTG stock solution (1 mM) was dissolved in DI water. All chemical solutions were sterilized using 0.2 μm VWR syringe filters.

To prepare RIF-agar plates, the stock RIF solution was added to autoclaved LB agar, resulting in a final plate concentration of 500 μg/mL RIF. Unless specified otherwise, overnight pre-cultures were generated in 14-mL Falcon test tubes containing 2 mL of liquid media. These pre-cultures were inoculated from a 25% glycerol cell stocks stored at −80 °C and cultivated for 24 h at 37 °C with shaking at 250 revolutions per min (rpm). Experimental cell cultures were prepared by

490 diluting the overnight pre-cultures (1:100) into 2 mL of fresh LB medium in 14-mL Falcon test
491 tubes. Bacterial cells in this study reached the mid-exponential phase ($OD_{600} \sim 0.5$) after around 3
492 h, attaining an average cell density of 7×10^8 CFUs/ml. All treatments involving UV were
493 administered at this stage.

UV treatment and cell recovery

Overnight pre-cultures of *E. coli* MG1655 cells were diluted 100-fold in 2 mL fresh LB media in
test tubes and grown at 37°C with shaking (250 rpm). Cell growth was monitored by measuring
optical density at 600 nm wavelength (OD_{600}) with a plate reader (Varioskan LUX Multimode
Microplate Reader, Thermo Fisher, Waltham, MA, United States). When the cell density reached
$OD_{600} \sim 0.5$, cultures from the test tubes were transferred to petri dishes (the diameter of petri
dishes= 100 mm, catalog no. FB0875713, Fisher Scientific). This procedure created a thin film of
culture in the dish with a height of about 0.25 mm. This configuration resulted in a uniformly
distributed liquid film without observable surface tension-driven droplet formation (i.e., beading).
The cultures were exposed to a light source emitting UV-B at 302 nm (UVP ChemStudio, catalog
no. 849-97-0928-02; Analytik Jena, Jena, Germany) for varying exposure times (0, 2, 4, 8, 16, 24
and 32 min). In this setup, UV-B light was emitted from below using a transilluminator and passed
through a UV-permeable plastic Petri dish placed directly on the transilluminator surface. This
configuration ensured uniform and efficient bottom-up exposure of bacterial cells within the thin
culture film. To verify the consistency and intensity of UV-B irradiation, direct irradiance
measurements were performed using a Vernier™ UV-B Sensor (Fisher Scientific, Cat. No.
S16273ND), calibrated for peak sensitivity at 315 nm (detection range: 290–320 nm), in
combination with a LabQuest™ 3 data acquisition platform. Measurements were taken under four
conditions to assess the impact of the experimental setup on UV transmission:(a) sensor placed
directly on the transilluminator surface without a Petri dish, (b) sensor on an empty Petri dish, (c)
sensor inside a Petri dish containing LB medium (not in contact with the liquid), and
(d) sensor in direct contact with the LB medium inside the dish. In all cases, UV-B irradiance was
consistently measured at approximately 1000 mW/m² (equivalent to 1 W/m²), indicating that
neither the Petri dish nor the medium significantly interfered with UV transmission (see
Supplementary Fig. S1). The energy dose (in J/m²) for each exposure condition was calculated
using the formula:

Energy dose (J/m²) = Irradiance (W/m²) × Exposure time (s)

After UV exposure for the indicated time points, cells with the LB media were transferred back to
test tubes and recovered for 24 h. During the recovery period, 10 μl samples were collected at
specified time points from each test tube, serially diluted in PBS in round-bottom 96-well plates,
and then, plated on LB agar media. The plates were incubated at 37 °C for 16 h to enumerate CFU.
We note that new colonies were not formed when incubated beyond 16 h. A similar procedure was
followed for the other strains. For dark recovery experiments, all procedures prior to UV treatment
were identical to those described above. Following UV exposure, samples were incubated in a
large, dark incubator-shaker, with the front window fully covered in aluminum foil to prevent light
exposure and photoreactivation. Additionally, the test tubes containing the cultured cells were
completely wrapped in aluminum foil to ensure total darkness during the recovery period.

Assessing mutagenesis

To assess the extent of mutant cell formation induced by UV exposure, cells were exposed to UV
radiation for the specified durations, followed by a 24-h recovery period. After this recovery

period, 500 μL of cells were collected and spread onto agar plates containing 500 μg/mL of RIF.
These plates were then incubated at 37 °C for 16 h to enumerate RIF-resistant colonies. To
determine clonogenic survival, CFU levels were determined, and mutant levels were normalized
by dividing the number of RIF-resistant colonies by the total number of colonies in a 1 mL culture
volume. Unless specified otherwise, mutant formation was reported as the count of RIF-resistant
colonies per 10^8 cell population.

PI Staining

Mid-exponential *E. coli* cells were treated with UV for 16 min and 32 min duration following the
protocol mentioned above. At the beginning of the recovery and 1h after the recovery (t=0 h and
1 h), UV-treated cells in LB media were diluted 20-fold in 1.0 ml 0.85% NaCl solution in flow
cytometry tubes (5 ml round bottom Falcon tubes, size: 12×75 mm) to achieve the desired cell
density ($\sim 10^6$-10^7 cells/ml) for flow cytometry analysis. The resulting cell suspensions were treated
with 20 μM PI dye. PI produces red fluorescence upon binding DNA; however, it can only
penetrate cells with damaged membranes. The samples were incubated in the dark at 37 °C for
15 min before analyzing them with a conventional bench-top flow cytometer (NovoCyte
3000RYB, ACEA Biosciences Inc., San Diego, CA, United States). For flow cytometry analysis,
we chose a slow sample flow rate (14 μl/ min) to have a sample stream diameter (i.e., core
diameter) of 7.7 μm. The instrument has a constant sheath flow rate of 6.5 ml/min. The flow
cytometer utilizes low-power solid-state lasers. Cells were excited at a 561 nm wavelength and red
fluorescence was detected with a 615/20-nm bandpass filter. At least 30,000 events were recorded
for each sample. NovoExpress software was used to collect the data. PI-stained dead cells, obtained
after ethanol (70% v/v) treatment, were used as a positive control. PI-stained live cells were used
as a negative control. Forward and side scatter signals of untreated live cells were used to
determine the cells on flow cytometry diagrams; the positive and negative controls were used to
gate PI positive (+) and PI negative (-) cell populations **(Supplementary Fig. S8).**

H₂O₂ measurement

The Amplex™ Red Hydrogen Peroxide/Peroxidase Assay Kit (Invitrogen, Thermo fischer
scientific, Catalog number: A22188) was used to assess the amount of H_2O_2 formed in the UV-
treated cells. The Amplex™ Red Hydrogen Peroxide/Peroxidase Assay Kit contains a sensitive,
one-step assay that uses the Amplex™ Red reagent (10-acetyl-3,7-dihydroxyphenoxazine) in
combination with horseradish peroxidase (HRP) to detect hydrogen peroxide (H_2O_2). In the
presence of peroxidase, the Amplex™ Red reagent reacts with H_2O_2 in a 1:1 stoichiometry to
produce the red-fluorescent oxidation product, resorufin. Resorufin has excitation and emission
maxima of approximately 571 nm and 585 nm. For preparing the stock solutions, 10 mM Amplex®
Red reagent was prepared by dissolving the contents of the vial of Amplex® Red reagent in 60 μL
of DMSO. Reaction buffer was diluted 5-fold to prepare 1X reaction buffer and 10 U/mL
Horseradish Peroxidase (HRP) was prepared by dissolving one vial of HRP in 1.0 mL of 1X
Reaction Buffer. The working solution of 100 μM Amplex® Red reagent and 0.2 U/mL HRP was
prepared by adding 50 μL of 10 mM Amplex® Red reagent stock solution and 100 μL of 10 U/mL
HRP stock solution in 4.85 mL of 1X Reaction Buffer. The assay volume for this experiment was
100 μL. Mid-exponential *E. coli* cells were treated with UV for 16 min and 32 min duration and
right after the treatment, 50 μL of sample for each condition was serially diluted in 1X Reaction
Buffer to determine the optimal amount of sample for the assay. 50 μL of the Amplex® Red
reagent/HRP working solution was added to each microplate well containing the samples. The

samples were incubated at room temperature for 30 min, and protected from light. At the end, the
fluorescence was measured using a microplate reader equipped for excitation in the range of 530–
560 nm and fluorescence emission detection at ~590 nm. For the standard curve, 20 mM Hydrogen
Peroxide (H_2O_2) working solution was prepared by dissolving 3.0% (0.88 M) H_2O_2 in 1X Reaction
Buffer and then further diluting 20 mM H_2O_2 working solution into 1X Reaction Buffer to produce
H_2O_2 concentrations of 0 to 10 µM, each in a volume of 50 µL. 50 µL of the Amplex® Red
reagent/HRP working solution was added to each microplate well containing the standards
followed by 30 min of incubation and then measuring the fluorescence following the same way
(Supplementary Fig. S9).

Fluorescent protein expression assay for reporter genes

Overnight pre-cultures of *E. coli* MG1655 cells with reporter genes fused to the SOS gene
promoters (P_{recA}, P_{uvrA}, and P_{uvrD}) were diluted 1:100 in 2 mL of LB media within test tubes and
incubated at 37°C with shaking (250 rpm). Upon reaching the mid-exponential phase, the cells
were subjected to UV radiation for specified durations (16 min and 32 min). An untreated culture
was used as a control. At the beginning of the recovery, 20 µM PI dye was added to the cultures
for continuous monitoring of the membrane permeability. At specified time points during the
recovery (t=0, 0.25, 0.5, 1, 2, 3 and 4 h), UV-treated cells were diluted 20-fold in 1.0 ml 0.85%
NaCl solution in flow cytometry tubes (5 ml round bottom Falcon tubes, size: 12×75 mm) to
achieve the desired cell density (~10^6-10^7 cells/ml). The samples were analyzed using the same
flow cytometry method described above (see the "PI staining" section); however, cells were
analyzed using two lasers. For measuring the red fluorescence from PI dye, cells were excited at a
561 nm wavelength, and red fluorescence was detected with a 615/20-nm bandpass filter. For
measuring the green fluorescence, cells were excited at a 488 nm wavelength, and the green
fluorescence was detected with a 530/30 bandpass filter.

Transcription/translation activities of pUA66-*gfp* plasmid

To assess the effect of the UV treatment on transcription/translation, the amount of GFP produced
by *E. coli* strains from the low-copy plasmid was measured. The plasmid, pUA66-*gfp*, has a *gfp*
gene under the control of a strong, IPTG-inducible T5 promoter and a strong LacI^q repressor.
Overnight pre-cultures of *E. coli* MG1655 cells carrying pUA66-*gfp* were diluted 100-fold in 2
mL fresh LB media in test tubes and grown at 37°C with shaking (250 rpm). At the mid-
exponential phase (OD_{600}~0.5), 0.1mM IPTG was added to the test tubes and then immediately
treated with UV for 16 min and 32 min duration. Untreated cultures having IPTG only (without
UV exposure) served as controls. At specified time points during the recovery (t=0, 0.25, 0.5, 1,
2, 3, and 4 h), UV-treated cells were diluted 20-fold in 1.0 ml PBS in flow cytometry tubes (5 ml
round bottom Falcon tubes, size: 12×75 mm) to achieve the desired cell density (~10^6-10^7
cells/ml). The samples were analyzed using the same flow cytometry method described above (see
the "PI staining" section); however, cells were analyzed with a laser emitting light at 488 nm and
the green fluorescence was detected with a 530/30 bandpass filter.

Promoter library screening

Overnight precultures were prepared by inoculating the strains from the promoter library into the
wells of 96-well plates containing 200 µl LB medium and 50 µg/ml kanamycin (for plasmid
retention). The plates were sealed with a sterile, oxygen-permeable membrane (Breathe-Easier,

Cat# BERM-2000, VWR International) and cultured for 24 h at 37°C with shaking at 250 rpm.
Overnight precultures were diluted 1:40 in LB medium with kanamycin in a new 96-well plate,
sealed, and incubated at 37°C with shaking at 250 rpm. At the mid-exponential phase (OD_{600} =
0.5), cultures in 96 well plates were exposed to UV light (UVP ChemStudio, catalog no. 849-97-
0928-02; Analytik Jena, Jena, Germany) for 16 min, and then, recovered for 24 h at 37°C with
shaking at 250 rpm. GFP was measured with a Varioskan LUX Multimode Microplate Reader
(Thermo Fisher, Waltham, MA, USA) at the indicated times with untreated cultures as a control.
The excitation and emission wavelengths for GFP measurement were 485 nm and 511 nm,
respectively. Fold changes for GFP (treated/untreated cultures after 24 h treatment) were used to
report the promoter activity. The top 10 promoters showing the highest expression were selected
for subsequent experiments.

Screening *E. coli* BW25113 Keio knockout collection

Overnight cultures of single mutants **(Supplementary Table 1)** carrying kanamycin resistance
genes were diluted 40-fold into flat-bottom 96-well plates, with each well containing 200 μL of
cell culture. Kanamycin (50 μg/ml) was added to overnight and treatment cultures to prevent
contamination. The plates were securely sealed using a sterile, oxygen-permeable membrane
(Breathe-Easier, Cat# BERM-2000, VWR International) and incubated at 37°C with agitation at
250 rpm. When the cultures reached the mid-exponential phase (OD_{600} ~ 0.5), the cells in the 96-
well plates were exposed to UV light (UVP ChemStudio, catalog no. 849-97-0928-02; Analytik
Jena, Jena, Germany) for 16 min. Subsequently, they were allowed to recover for 24 h at 37°C
with shaking at 250 rpm. To quantify UV-induced mutant cells, 10 μL samples were collected
from each well and spotted onto LB agar plates, containing 500 μg/mL RIF, at the end of the 24-
h recovery period. The plates were then incubated at 37°C for 16 h to enumerate RIF-resistant
colonies. The mutant formation was reported as the count of RIF-resistant colonies per 10 μL of
cell culture.

Overexpression of *recA* Using an IPTG-Inducible Plasmid

Overnight pre-cultures of *E. coli* MG1655 Δ*recA* cells harboring the pUA66-*recA* plasmid were
diluted 1:100 into 2 mL of fresh LB medium in test tubes and incubated at 37 °C with shaking at
250 rpm. After 2.5 hours of growth, IPTG was added to final concentrations of 0.01, 0.1, and 1
mM. One tube was maintained without IPTG as a non-induced control. Cultures were then
incubated for an additional 30 minutes to allow *recA* expression prior to UV exposure.
Subsequently, the cultures were exposed to UV for either 16 or 32 minutes. As a control, *E. coli*
MG1655 Δ*recA* cells carrying the pUA66 empty vector were included for both UV exposure
durations. During the recovery period, 10 μL samples were collected at defined time points from
each culture, serially diluted in PBS using round-bottom 96-well plates, and plated on LB agar to
determine CFU. Plates were incubated at 37 °C for 16 hours. RIF-resistant colonies were
enumerated following the same procedure.

Statistical analysis and reproducibility

For all pairwise comparisons, one-way ANOVA with Dunnett's post-test was utilized. A minimum
of four independent biological replicates (unless otherwise specified) were conducted for
experiments involving UV exposure. In all figures, data corresponding to each time point
represents the mean value ± standard deviation. Regarding statistical significance analysis, the
threshold values were set as follows: $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and $****P < 0.0001$. All

figures were created using GraphPad Prism 10.0.2, and the statistical analyses were carried out using GraphPad Prism 10.0.2 statistical functions. FlowJo V 10.7.1 was used to analyze the data obtained from flow cytometry.

Acknowledgment

The authors would like to thank the members of Orman Lab for their help. This study was supported by NIH/NIAID R01-AI143643.

Contributions

S.G. and M.A.O. conceived and designed the study. S.G., and J.N. performed the experiments. S.G. and M.A.O. analyzed the data and wrote the paper. All authors read and approved the paper.

Corresponding author

Correspondence to Mehmet A. Orman (morman@central.uh.edu)

Declaration of interests

The authors declare no competing interests.

Data availability

All data presented in this manuscript are available in the Main Text or the Supplementary File. All raw data have been published on Figshare:

https://doi.org/10.6084/m9.figshare.27701493.v2

References

1.   Hershberg, R. Mutation—The Engine of Evolution: Studying Mutation and Its Role in the Evolution of Bacteria. *Cold Spring Harb Perspect Biol* **7**, (2015).

2.   MacLean, R. C., Torres-Barceló, C. & Moxon, R. Evaluating evolutionary models of stress-induced mutagenesis in bacteria. *Nature Reviews Genetics 2013 14:3* **14**, 221–227 (2013).

3.   Zimmermann, A., Prieto-Vivas, J. E., Voordeckers, K., Bi, C. & Verstrepen, K. J. Mutagenesis techniques for evolutionary engineering of microbes – exploiting CRISPR-Cas, oligonucleotides, recombinases, and polymerases. *Trends Microbiol* (2024) doi:10.1016/J.TIM.2024.02.006.

4.   Sung, W., Ackerman, M. S., Miller, S. F., Doak, T. G. & Lynch, M. Drift-barrier hypothesis and mutation-rate evolution. *Proc Natl Acad Sci U S A* **109**, 18488–18492 (2012).

5.   Wielgoss, S. *et al.* Mutation rate dynamics in a bacterial population reflect tension between adaptation and genetic load. *Proc Natl Acad Sci U S A* **110**, 222–227 (2013).

6.   Fitzgerald, D. M., Hastings, P. J. & Rosenberg, S. M. Stress-Induced Mutagenesis: Implications in Cancer and Drug Resistance. *Annu Rev Cancer Biol* **1**, 119 (2017).

7.   Morreall, J. F., Petrova, L. & Doetsch, P. W. Transcriptional mutagenesis and its potential roles in the etiology of cancer and bacterial antibiotic resistance. *J Cell Physiol* **228**, 2257 (2013).

8.   Russo, M., Sogari, A. & Bardelli, A. Adaptive evolution: How bacteria and cancer cells survive stressful conditions and drug treatment. *Cancer Discov* **11**, 1886–1895 (2021).

9.   Lambert, G. *et al.* An analogy between the evolution of drug resistance in bacterial
communities and malignant tissues. *Nat Rev Cancer* **11**, 375 (2011).
10.   Marshall, J. E. A., Lakin, J., Troth, I. & Wallace-Johnson, S. M. UV-B radiation was the
Devonian-Carboniferous boundary terrestrial extinction kill mechanism. *Sci Adv* **6**,
eaba0768 (2020).
11.   Liu, F. *et al.* Dying in the Sun: Direct evidence for elevated UV-B radiation at the end-
Permian mass extinction. *Sci Adv* **9**, (2023).
12.   Benca, J. P., Duijnstee, I. A. P. & Looy, C. V. UV-B–induced forest sterility: Implications
of ozone shield failure in earth's largest extinction. *Sci Adv* **4**, (2018).
13.   Ries, G. *et al.* Elevated UV-B radiation reduces genome stability in plants. *Nature* **406**, 98–
101 (2000).
14.   Rothschild, L. J. The Influence of UV Radiation on Protistan Evolution. *Journal of
Eukaryotic Microbiology* **46**, 548–555 (1999).
15.   Rozema, J. *et al.* The role of UV-B radiation in aquatic and terrestrial ecosystems—an
experimental and functional analysis of the evolution of UV-absorbing compounds. *J
Photochem Photobiol B* **66**, 2–12 (2002).
16.   Rastogi, R. P., Richa, Kumar, A., Tyagi, M. B. & Sinha, R. P. Molecular Mechanisms of
Ultraviolet Radiation-Induced DNA Damage and Repair. *J Nucleic Acids* **2010**, 32 (2010).
17.   Sinha, R. P. & Häder, D. P. UV-induced DNA damage and repair: a review. *Photochemical
& Photobiological Sciences 2002 1:4* **1**, 225–236 (2002).
18.   Krishna, S., Maslov, S. & Sneppen, K. UV-Induced Mutagenesis in Escherichia coli SOS
Response: A Quantitative Model. *PLoS Comput Biol* **3**, e41 (2007).
19.   Kciuk, M., Marciniak, B., Mojzych, M. & Kontek, R. Focus on UV-Induced DNA Damage
and Repair—Disease Relevance and Protective Strategies. *Int J Mol Sci* **21**, 1–33 (2020).
20.   Gray, N. F. Ultraviolet Disinfection. *Microbiology of Waterborne Diseases:
Microbiological Aspects and Risks: Second Edition* 617–630 (2014) doi:10.1016/B978-0-
12-415846-7.00034-2.
21.   Hessling, M., Haag, R., Sieber, N. & Vatter, P. The impact of far-UVC radiation (200-230
nm) on pathogens, cells, skin, and eyes - a collection and analysis of a hundred years of
data. *GMS Hyg Infect Control* **16**, Doc07 (2021).
22.   Black, H. S. *et al.* Photocarcinogenesis: an overview. *J Photochem Photobiol B* **40**, 29–47
(1997).
23.   Douki, T., Court, M., Sauvaigo, S., Odin, F. & Cadet, J. Formation of the main UV-induced
thymine dimeric lesions within isolated and cellular DNA as measured by high performance
liquid chromatography-tandem mass spectrometry. *J Biol Chem* **275**, 11678–11685 (2000).
24.   Rupert, C. S. Enzymatic Photoreactivation: Overview. *Basic Life Sci* **5 A**, 73–87 (1975).
25.   Seawell, P. C. *et al.* den V gene of bacteriophage T4 determines a DNA glycosylase specific
for pyrimidine dimers in DNA. *J Virol* **35**, 790–796 (1980).
26.   Van Houten, B. Nucleotide excision repair in Escherichia coli. *Microbiol Rev* **54**, 18–51
(1990).
27.   Rupp, W. D. & Howard-flanders, P. Discontinuities in the DNA synthesized in an Excision-
defective strain of Escherichia coli following ultraviolet irradiation. *J Mol Biol* **31**, 291–304
(1968).
28.   Wendel, B. M., Hollingsworth, S., Courcelle, C. T. & Courcelle, J. UV-induced DNA
damage disrupts the coordination between replication initiation, elongation and completion.
Genes to Cells **26**, 94–108 (2021).

29. Majchrzak, M., Bowater, R. P., Staczek, P. & Parniewski, P. SOS Repair and DNA
Supercoiling Influence the Genetic Stability of DNA Triplet Repeats in Escherichia coli. *J*
Mol Biol **364**, 612–624 (2006).
30. Simmons, L. A., Foti, J. J., Cohen, S. E. & Walker, G. C. The SOS Regulatory Network.
EcoSal Plus **3**, (2008).
31. Little, J. W. & Mount, D. W. The SOS regulatory system of Escherichia coli. *Cell* **29**, 11–
22 (1982).
32. Maslowska, K. H., Makiela-Dzbenska, K. & Fijalkowska, I. J. The SOS system: A complex
and tightly regulated response to DNA damage. *Environ Mol Mutagen* **60**, 368–384 (2019).
33. Walker, G. C. Mutagenesis and inducible responses to deoxyribonucleic acid damage in
Escherichia coli. *Microbiol Rev* **48**, 60–93 (1984).
34. Montelone, B. A. DNA Repair and Mutagenesis. Second Edition. By Errol C Friedberg,
Graham C Walker, Wolfram Siede, Richard D Wood, Roger A Schultz, and, Tom
Ellenberger. Washington (DC): ASM Press. $179.95. xxix + 1118 p + 11 pl; ill.; index.
ISBN: 1-55581-319-4. 2006. . *https://doi.org/10.1086/509407* **81**, 273–273 (2015).
35. Martins-Pinheiro, M., Marques, R. C. P. & Menck, C. F. M. Genome analysis of DNA repair
genes in the alpha proteobacterium Caulobacter crescentus. *BMC Microbiol* **7**, 17 (2007).
36. Kuzminov, A. Recombinational Repair of DNA Damage in Escherichia coli and
Bacteriophage λ . *Microbiology and Molecular Biology Reviews* **63**, 751–813 (1999).
37. Burckhardt, S. E., Woodgate, R., Scheuermann, R. H. & Echols, H. UmuD mutagenesis
protein of Escherichia coli: overproduction, purification, and cleavage by RecA.
Proceedings of the National Academy of Sciences **85**, 1811–1815 (1988).
38. Woodgate, R., Rajagopalan, M., Lu, C. & Echols, H. UmuC mutagenesis protein of
Escherichia coli: purification and interaction with UmuD and UmuD'. *Proceedings of the*
National Academy of Sciences **86**, 7301–7305 (1989).
39. Kato, T. & Shinoura, Y. Isolation and characterization of mutants of Escherichia coli
deficient in induction of mutations by ultraviolet light. *MGG Molecular & General Genetics*
156, 121–131 (1977).
40. Wang, Z. Translesion synthesis by the UmuC family of DNA polymerases. *Mutation*
Research/DNA Repair **486**, 59–70 (2001).
41. Liu, J. *et al.* MutL sliding clamps coordinate exonuclease-independent Escherichia coli
mismatch repair. *Nat Commun* **10**, (2019).
42. Lin, T. *et al.* Contribution of the SOS response and the DNA repair systems to norfloxacin
induced mutations in E. coli. *Mar Life Sci Technol* **5**, 538–550 (2023).
43. Shibai, A. *et al.* Mutation accumulation under UV radiation in Escherichia coli. *Scientific*
Reports 2017 7:1 **7**, 1–12 (2017).
44. Badran, A. H. & Liu, D. R. Development of potent in vivo mutagenesis plasmids with broad
mutational spectra. *Nature Communications 2015 6:1* **6**, 1–10 (2015).
45. Tillich, U. M., Lehmann, S., Schulze, K., Dühring, U. & Frohme, M. The Optimal Mutagen
Dosage to Induce Point-Mutations in Synechocystis sp. PCC6803 and Its Application to
Promote Temperature Tolerance. *PLoS One* **7**, e49467 (2012).
46. Hashimoto, S. *et al.* Isolation of auxotrophic mutants of diploid industrial yeast strains after
UV mutagenesis. *Appl Environ Microbiol* **71**, 312–319 (2005).
47. Hijnen, W. A. M., Beerendonk, E. F. & Medema, G. J. Inactivation credit of UV radiation
for viruses, bacteria and protozoan (oo)cysts in water: A review. *Water Res* **40**, 3–22 (2006).

48.   Reed, N. G. The History of Ultraviolet Germicidal Irradiation for Air Disinfection. *Public*
Health Reports **125**, 15 (2010).

49.   Bergkessel, M., Basta, D. W. & Newman, D. K. The physiology of growth arrest: uniting
molecular and environmental microbiology. *Nat Rev Microbiol* **14**, 549 (2016).

50.   Dörr, T., Vulić, M. & Lewis, K. Ciprofloxacin causes persister formation by inducing the
TisB toxin in Escherichia coli. *PLoS Biol* **8**, 29–35 (2010).

51.   Völzing, K. G. & Brynildsen, M. P. Stationary-phase persisters to ofloxacin sustain DNA
damage and require repair systems only during recovery. *mBio* **6**, (2015).

52.   Santos, A. L. *et al.* Effects of UV radiation on the lipids and proteins of bacteria studied by
mid-infrared spectroscopy. *Environ Sci Technol* **47**, 6306–6315 (2013).

53.   Wurtmann, E. J. & Wolin, S. L. RNA under attack: Cellular handling of RNA damage. *Crit*
Rev Biochem Mol Biol **44**, 34 (2009).

54.   Ghosh, S. & Orman, M. A. Exploring the links between SOS response, mutagenesis, and
resistance during the recovery period. *Antimicrob Agents Chemother* **68**, (2024).

55.   Kumar, J., Babele, P. K., Singh, D. & Kumar, A. UV-B radiation stress causes alterations
in whole cell protein profile and expression of certain genes in the rice phyllospheric
bacterium Enterobacter cloacae. *Front Microbiol* **7**, 208904 (2016).

56.   Joux, F., Jeffrey, W. H., Lebaron, P. & Mitchell, D. L. Marine bacterial isolates display
diverse responses to UV-B radiation. *Appl Environ Microbiol* **65**, 3820–3827 (1999).

57.   Fernández Zenoff, V., Siñeriz, F. & Farías, M. E. Diverse responses to UV-B radiation and
repair mechanisms of bacteria isolated from high-altitude aquatic environments. *Appl*
Environ Microbiol **72**, 7857–7863 (2006).

58.   Xue, Y. & Nicholson, W. L. The two major spore DNA repair pathways, nucleotide excision
repair and spore photoproduct lyase, are sufficient for the resistance of Bacillus subtilis
spores to artificial UV-C and UV-B but not to solar radiation. *Appl Environ Microbiol* **62**,
2221–2227 (1996).

59.   Prada Medina, C. A., Aristizabal Tessmer, E. T., Quintero Ruiz, N., Serment-Guerrero, J.
& Fuentes, J. L. Survival and SOS response induction in ultraviolet B irradiated Escherichia
coli cells with defective repair mechanisms. *Int J Radiat Biol* **92**, 321–328 (2016).

60.   Barrett, T. C., Mok, W. W. K., Murawski, A. M. & Brynildsen, M. P. Enhanced antibiotic
resistance development from fluoroquinolone persisters after a single exposure to antibiotic.
Nature Communications 2019 10:1 **10**, 1–11 (2019).

61.   Krašovec, R. *et al.* Mutation rate plasticity in rifampicin resistance depends on Escherichia
coli cell-cell interactions. *Nat Commun* **5**, (2014).

62.   Reynolds, M. G. Compensatory evolution in rifampin-resistant Escherichia coli. *Genetics*
156, 1471 (2000).

63.   Pope, C. F., O'Sullivan, D. M., McHugh, T. D. & Gillespie, S. H. A Practical Guide to
Measuring Mutation Rates in Antibiotic Resistance. *Antimicrob Agents Chemother* **52**, 1209
(2008).

64.   Rosche, W. A. & Foster, P. L. Determining Mutation Rates in Bacterial Populations.
Methods **20**, 4 (2000).

65.   Lagage, V., Chen, V. & Uphoff, S. Adaptation delay causes a burst of mutations in bacteria
responding to oxidative stress. *EMBO Rep* **24**, (2023).

66.   Rodríguez-Rojas, A. *et al.* Non-lethal exposure to H2O2 boosts bacterial survival and
evolvability against oxidative stress. *PLoS Genet* **16**, e1008649 (2020).

67. Husain, I. & Sancar, A. Photoreactivation in phr mutants of Escherichia coli K-12. *J*
Bacteriol **169**, 2367 (1987).
68. Hays, J. B., Martin, S. J. & Bhatia, K. Repair of nonreplicating UV-irradiated DNA:
cooperative dark repair by Escherichia coli uvr and phr functions. *J Bacteriol* **161**, 602–608
(1985).
69. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**, 6640 (2000).
70. Gerdes, S. Y. *et al.* Experimental Determination and System Level Analysis of Essential
Genes in Escherichia coli MG1655. *J Bacteriol* **185**, 5673 (2003).
71. Witkin, E. M. Photoreversal and "dark repair" of mutations to prototrophy induced by
ultraviolet light in photoreactivable and non-photoreactivable strains of Escherichia coli.
Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis **1**, 22–36
(1964).
72. Castellani, A., Jagger, J. & Setlow, R. B. Overlap of Photoreactivation and Liquid Holding
Recovery in Escherichia coli B. *Science (1979)* **143**, 1170–1171 (1964).
73. Zaslaver, A. *et al.* A comprehensive library of fluorescent transcriptional reporters for
Escherichia coli. *Nat Methods* **3**, 623–628 (2006).
74. Keseler, I. M. *et al.* The EcoCyc database: reflecting new knowledge about Escherichia coli
K-12. *Nucleic Acids Res* **45**, D543–D550 (2017).
75. Fernández De Henestrosa, A. R. *et al.* Identification of additional genes belonging to the
LexA regulon in Escherichia coli. *Mol Microbiol* **35**, 1560–1572 (2000).
76. Goodall, E. C. A. *et al.* The Essential Genome of Escherichia coli K-12. *mBio* **9**, (2018).
77. Yu, X.-C., Weihe, E. K. & Margolin, W. Role of the C Terminus of FtsK in Escherichia
coli Chromosome Segregation. *J Bacteriol* **180**, 6424 (1998).
78. Wang, L. & Lutkenhaus, J. FtsK is an essential cell division protein that is localized to the
septum and induced as part of the SOS response. *Mol Microbiol* **29**, 731–740 (1998).
79. Meddows, T. R., Savory, A. P., Grove, J. I., Moore, T. & Lloyd, R. G. RecN protein and
transcription factor DksA combine to promote faithful recombinational repair of DNA
double-strand breaks. *Mol Microbiol* **57**, 97–110 (2005).
80. Picksley, S. M., Attfield, P. V. & Lloyd, R. G. Repair of DNA double-strand breaks in
Escherichia coli K12 requires a functional recN product. *Mol Gen Genet* **195**, 267–274
(1984).
81. Wang, T. C. & Smith, K. C. Different effects of recJ and recN mutations on the
postreplication repair of UV-damaged DNA in Escherichia coli K-12. *J Bacteriol* **170**,
2555–2559 (1988).
82. Kosinski, J., Feder, M. & Bujnicki, J. M. The PD-(D/E)XK superfamily revisited:
identification of new members among proteins involved in DNA metabolism and functional
predictions for domains of (hitherto) unknown function. *BMC Bioinformatics* **6**, (2005).
83. Lomize, A. L., Lomize, M. A., Krolicki, S. R. & Pogozheva, I. D. Membranome: a database
for proteome-wide analysis of single-pass membrane proteins. *Nucleic Acids Res* **45**, D250–
D255 (2017).
84. Napolitano, R., Janel-Bintz, R., Wagner, J. & Fuchs, R. P. P. All three SOS-inducible DNA
polymerases (Pol II, Pol IV and Pol V) are involved in induced mutagenesis. *EMBO J* **19**,
6259 (2000).

85. Hickson, I. D., Robson, C. N., Atkinson, K. E., Hutton, L. & Emmerson, P. T. Reconstitution of RecBC DNase activity from purified Escherichia coli RecB and RecC proteins. *J Biol Chem* **260**, 1224–9 (1985).

86. Payne-Dwyer, A. L., Syeda, A. H., Shepherd, J. W., Frame, L. & Leake, M. C. RecA and RecB: probing complexes of DNA repair proteins with mitomycin C in live Escherichia coli with single-molecule sensitivity. *J R Soc Interface* **19**, (2022).

87. Reuven, N. B., Arad, G., Maor-Shoshani, A. & Livneh, Z. The mutagenesis protein UmuC is a DNA polymerase activated by UmuD', RecA, and SSB and is specialized for translesion replication. *J Biol Chem* **274**, 31763–31766 (1999).

88. Takahagi, M., Iwasaki, H., Nakata, A. & Shinagawa, H. Molecular analysis of the Escherichia coli ruvC gene, which encodes a Holliday junction-specific endonuclease. *J Bacteriol* **173**, 5747–5753 (1991).

89. Loewen, P. C. Isolation of catalase-deficient Escherichia coli mutants and genetic mapping of katE, a locus that affects catalase activity. *J Bacteriol* **157**, 622–626 (1984).

90. Epshtein, V. *et al.* UvrD facilitates DNA repair by pulling RNA polymerase backwards. *Nature* **505**, 372–377 (2014).

91. Orren, D. K. & Sancar, A. The (A)BC excinuclease of Escherichia coli has only the UvrB and UvrC subunits in the incision complex. *Proc Natl Acad Sci U S A* **86**, 5237–5241 (1989).

92. de Oliveira, A. H. S., da Silva, A. E., de Oliveira, I. M., Henriques, J. A. P. & Agnez-Lima, L. F. MutY-glycosylase: An overview on mutagenesis and activities beyond the GO system. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* **769**, 119–131 (2014).

93. Shurtleff, B. W., Ollivierre, J. N., Tehrani, M., Walker, G. C. & Beuning, P. J. Steric Gate Variants of UmuC Confer UV Hypersensitivity on Escherichia coli. *J Bacteriol* **191**, 4815 (2009).

94. Wang, T. chien V. & Smith, K. C. Role of the umuC gene in postreplication repair in UV-irradiated Escherichia coli K-12 uvrB. *Mutation Research/DNA Repair Reports* **145**, 107–112 (1985).

95. Moolenaar, G. F., Moorman, C. & Goosen, N. Role of the Escherichia coli Nucleotide Excision Repair Proteins in DNA Replication. *J Bacteriol* **182**, 5706 (2000).

96. Orren, D., Selby, C., Hearst, J. & Sancar, A. Post-incision steps of nucleotide excision repair in Escherichia coli. Disassembly of the UvrBC-DNA complex by helicase II and DNA polymerase I. *J Biol Chem* **267**, 780–788 (1992).

97. Bertrand-Burggraf, E., Selby, C. P., Hearst, J. E. & Sancar, A. Identification of the different intermediates in the interaction of (A)BC excinuclease with its substrates by DNase I footprinting on two uniquely modified oligonucleotides. *J Mol Biol* **219**, 27–36 (1991).

98. Higashitani, A., Ishii, Y., Kato, Y. & Horiuchi, K. Functional dissection of a cell-division inhibitor, SulA, of Escherichia coli and its negative regulation by Lon. *Molecular and General Genetics* **254**, 351–357 (1997).

99. Unoson, C. & Wagner, E. G. H. A small SOS-induced toxin is targeted against the inner membrane in Escherichia coli. *Mol Microbiol* **70**, 258–270 (2008).

100. McKenzie, K. *et al.* The effects of 405 nm light on bacterial membrane integrity determined by salt and bile tolerance assays, leakage of UV-absorbing material and SYTOX green labelling. *Microbiology (N Y)* **162**, 1680 (2016).

101. Yeo, S. K. & Liong, M. T. Effects and applications of sub-lethal ultrasound, electroporation and UV radiations in bioprocessing. *Ann Microbiol* **63**, 813–824 (2013).

928 102. Cadet, J., Douki, T. & Ravanat, J. L. Oxidatively generated base damage to cellular DNA.
929    *Free Radic Biol Med* **49**, 9–21 (2010).
930 103. Winterbourn, C. C. Reconciling the chemistry and biology of reactive oxygen species.
931    *Nature Chemical Biology 2008 4:5* **4**, 278–286 (2008).
932 104. Schieber, M. & Chandel, N. S. ROS Function in Redox Signaling and Oxidative Stress.
933    *Curr Biol* **24**, R453 (2014).
934 105. Zwieb, C., Ross, A., Rinke, J., Meinke, M. & Brimacombe, R. Evidence for RNA-RNA
935    cross-link formation in Escherichia coli ribosomes. *Nucleic Acids Res* **5**, 2705 (1978).
936 106. Orman, M. A. & Brynildsen, M. P. Inhibition of stationary phase respiration impairs
937    persister formation in E. coli. *Nature Communications 2015 6:1* **6**, 1–13 (2015).
938 107. Mohiuddin, S. G., Nguyen, T. V. & Orman, M. A. Pleiotropic actions of phenothiazine
939    drugs are detrimental to Gram-negative bacterial persister cells. *Communications Biology*
940    *2022 5:1* **5**, 1–15 (2022).
941 108. Mohiuddin, S. G., Massahi, A. & Orman, M. A. lon Deletion Impairs Persister Cell
942    Resuscitation in Escherichia coli. *mBio* **13**, (2022).

Figures

Fig. 1: Excessive UV exposure resulted in transient loss of culturability and decreased mutagenesis. (a) The graph shows the relationship between UV-B exposure time (min) and calculated energy dosage (J/m²), based on measured irradiance (see Materials and Methods). (b) Exponential-phase *E. coli* MG1655 WT cells were exposed to UV-B light for 0, 2, 4, 8, 16, 24,

and 32 minutes, followed by a 24-h recovery period. At specific time points during recovery ($t = 0$
h, 0.25 h, 0.5 h, 1 h, 2 h, 4 h, 6 h, and 24 h), cells were collected and plated to determine colony-
forming units (CFU). (c) Levels of UV-induced rifampicin (RIF) resistance mutations were
measured by counting RIF-resistant colonies (per 10^8 cells) in the WT cultures after recovery for
the indicated UV exposure times. (d) CFU counts and (e) mutation frequency of *E. coli* MG1655
following UV-B exposure were determined under dark conditions. The same experimental setup
described above was used, but cultures were recovered in the absence of light to eliminate
photoreactivation. (f) The temporal CFU profiles of *E. coli* MG1655 Δ*recA* cells were monitored
during recovery following UV treatment. n=4. Statistical analysis was performed using one-way
ANOVA with Dunnett's post-test, where $*P < 0.05$, $**P < 0.001$, $***P < 0.01$, $****P < 0.0001$.
Data corresponding to each time point represent mean value ± standard deviation.

982
983

Fig. 2: UV-induced upregulation of SOS genes and its impact on mutagenesis and cell culturability. (a) A library of *E. coli* MG1655 strains with promoter reporters for SOS response genes was UV-treated during the mid-exponential phase and allowed to recover for 24 h. GFP levels were measured after recovery and normalized to untreated controls. WT cells without any promoter reporters were used as background controls. (b) Genes upregulated following UV exposure were individually or combinatorially deleted in *E. coli* MG1655, and a mutagenesis assay was performed. Mid-exponential phase cells were exposed to UV for 0, 16, and 32 min and

recovered for 24 h. At specified time points during recovery (t = 0 h, 0.25 h, 0.5 h, 1 h, 2 h, 4 h, 6
h, and 24 h), cells were collected and plated to determine CFU. Note that only the 0 and 0.25 h
time points are shown here; the full-time course is provided in **Supplementary Fig. S3**. (c) RIF-
resistant cells were quantified by plating samples on RIF-agar plates after 24 h of recovery, with
results reported as RIF-resistant colony counts per 10^8 cells for different UV exposure durations.
n=4. Statistical analysis was performed using one-way ANOVA with Dunnett's post-test, where
*$P < 0.05$, **$P < 0.001$, ***$P < 0.01$, ****$P < 0.0001$. Data corresponding to each time point
represents mean value ± standard deviation.
999

1000
1001 **Fig. 3: Knockout library screening to elucidate their effects on mutagenesis and cell**
1002 **culturability**. (a) The *E. coli* BW25113 Keio knockout collection was screened following 16 min

of UV treatment. RIF-resistant cells were quantified by spotting 10 μL samples from each well onto RIF-agar plates after a 24-hour recovery period. Δ*lacI* was used as the reference control (see Materials and Methods). (b) Selected genes were deleted in *E. coli* MG1655 and the mutagenesis assay was performed. Mid-exponential phase cells were exposed to UV for 0, 16 and 32 min and recovered for 24 h. At indicated time points (t = 0 h, 0.25 h, 0.5 h, 1 h, 2 h, 4 h, 6 h and 24 h) during recovery, cells were collected and plated to enumerate CFU. Note that only the 0 and 0.25 h time points are shown here; the full-time course is provided in **Supplementary Fig. S4.** (c) RIF-resistant cells were quantified by plating the samples on RIF-agar plates after 24 h of recovery. The values were reported as RIF-resistant colony count per 10^8 cell population for different UV exposure durations. n=4. Statistical analysis was performed using one-way ANOVA with Dunnett's post-test, where $*P < 0.05$, $**P < 0.001$, $***P < 0.01$. Data corresponding to each time point represents mean value ± standard deviation.

1022

Fig. 4: Perturbation of both SulA and TisB proteins moderately increased cell culturability
during recovery. (a-c) Exponential-phase *E. coli* MG1655 Δ*sulA*, Δ*tisB*, and Δ*sulA*Δ*tisB* cells
were exposed to UV-B light for 0, 2, 4, 8, 16, 24, and 32 minutes, followed by a 24-h recovery
period. At specific time points during recovery, cells were collected and plated to determine their
CFU levels. Note that only the 0 and 0.25 h time points are shown here; the full-time course is
provided in **Supplementary Fig. S5.** (d-f) Levels of UV-induced RIF resistance mutations were
measured by counting RIF-resistant colonies (per 10^8 cells) in the cultures of three knockout strains
after recovery for the indicated UV exposure times. n=4. Statistical analysis was performed using
one-way ANOVA with Dunnett's post-test, where $*P < 0.05$, $**P < 0.001$, $***P < 0.01$,
$****P < 0.0001$. Data corresponding to each time point represents mean value ± standard
deviation.

1034

1035

Fig. 5: Excessive UV exposure did not compromise cell membrane integrity or increase hydrogen peroxide levels, but it did impair cellular translation. (a) Membrane integrity of UV-treated cells (0, 16, and 32 min of UV exposure) was assessed via flow cytometry with PI staining (see **Supplementary Fig. S8** for controls). FSC-H: Forward light scatter. (b) H_2O_2 concentrations were measured using the Amplex Red Hydrogen Peroxide/Peroxidase Assay Kit. Reaction mixtures containing Amplex Red reagent, horseradish peroxidase, and UV-treated cells in sodium phosphate buffer (pH 7.4) were incubated for 30 minutes at room temperature, followed by fluorescence measurement with a microplate reader. H_2O_2 levels were calculated using a standard curve (**Supplementary Fig. S9**). (c) Expression of P_{recA}-*gfp* in UV-treated *E. coli* MG1655 cells was analyzed via flow cytometry for the indicated time points during recovery (P_{recA}: the *recA* promoter). (d) IPTG-inducible GFP expression in UV-treated *E. coli* MG1655 pUA66-*gfp* cells was also assessed by flow cytometry for the indicated time points during recovery. (e) GFP expression profiles of UV-treated cells harboring the *recA* promoter and (f) IPTG-inducible GFP expression systems were evaluated using mean GFP values from flow cytometry data. (g) Recovery (CFU counts) and mutation frequency of *E. coli* MG1655 Δ*recA* carrying an IPTG-inducible *recA* overexpression plasmid were assessed following 16- and 32-minute UV-B

1052　exposures, shown for the indicated IPTG concentrations. $n \geq 4$. Statistical analysis was performed
1053　using one-way ANOVA with Dunnett's post-test, where $*P < 0.05$, $**P < 0.001$, $***P < 0.01$,
1054　$****P < 0.0001$. Data corresponding to each time point represent mean value $\pm$ standard deviation.