Quantifying key parameters of environmental transmission and age-specific

susceptibility for *Mycobacterium avium* subspecies *paratuberculosis (MAP)*

Yuqi Gao[1,2], Piter Bijma[2], Nienke Hartemink[1], Mart C.M. de Jong[1]

[1] Infectious Disease Epidemiology Group, Department of Animal Sciences,

Wageningen University and Research, Wageningen, Netherlands

[2] Animal Breeding and Genomics Group, Department of Animal Sciences,

Wageningen University and Research, Wageningen, Netherlands

* Corresponding author

E-mail: yuqi.gao@wur.nl (YG)

These authors contributed equally to this work.

Abstract

Paratuberculosis is a chronic disease in cows and other ruminants, caused by

Mycobacterium avium subspecies paratuberculosis (*MAP*). We developed an age-

specific dose-response model and two environmental transmission models (Model A

and B) to estimate key parameters based on previously published experiments for

MAP in dairy cows. In the dose-response model, the age-specific susceptibility

decrease rate parameter was estimated at $0.0629\,\mathrm{wk}^{-1}$, suggesting that a

previously used parameter of $0.1\,\mathrm{wk}^{-1}$ may have underestimated the infection risks

with increasing age. For the transmission models, Model A represents infectivity

differences among transiently infectious (Itr), low shedding (Il), high shedding (Ih)

individuals by varying transmission rate parameters ($\widehat{\beta_{Itr}}$, $\widehat{\beta_{Il}}$, $\widehat{\beta_{Ih}}$ = 0.0299, 0.0150,

0.219 wk⁻¹) with a standardized constant shedding rate parameter of 2.10 wk⁻¹,

whereas Model B captures these differences by varying shedding rate parameters

($\widehat{sh_{Itr}}$, $\widehat{sh_{Il}}$, $\widehat{sh_{Ih}}$: 2.10, 1.05, 15.4 wk⁻¹) with one transmission rate parameter of

0.0299 wk⁻¹. Although both models have identical best estimates and AIC values,

Model B exhibited wider 95% confidence intervals (95% CI). In both models, the

MAP decay rate parameter was estimated at 0.150 wk⁻¹, corresponding to a half-life

of MAP of approximately 4.61 weeks, which aligns well with previously published

values. To better interpret these parameters and understand how different biological

assumptions about infectivity influence predicted exposure, we performed scenario

analyses examining environmental contamination over time, along with infection

probabilities over exposure timing, exposure duration, and the recipient's age at

exposure. All 95% CI are provided in the main text.

Introduction

Paratuberculosis, caused by *Mycobacterium avium subspecies paratuberculosis*

(*MAP*), is a chronic disease affecting cows and other ruminants (1). The disease is

known for its long incubation period, imperfect diagnostic tests (especially for early

stages), limited vaccination effect, and long survival in the environment, all of which

alone or in combination could be the reason why *MAP* is difficult to control and

eradicate in cattle(2-7). *MAP* can be transmitted horizontally via contaminated milk or

faeces and vertically from dam to offspring (8, 9). A major obstacle in eradicating

MAP is its environmental transmission, where contaminated environments can cause

infections even in the absence of infectious individuals at the location and time of the

recipients' presence.

Understanding environmental transmission is crucial for *MAP* dynamics; therefore,

we applied an advanced mathematical model developed by Chang and De Jong (10)

to study it. This model, which has proven effective for various diseases (10-13), was

not yet used for *MAP*. The model incorporates three key parameters: for, shedding

rate, decay rate, and transmission rate. Additionally, if spatial spread is considered, a

fourth parameter, for the diffusion rate, is necessary (12). For *MAP* on dairy farms,

spatial transmission was simplified by considering three different environments on a

dairy farm: the calving house, calf collective pen, and lactating adult pen. This

research focuses on estimating the shedding, decay and transmission rate

parameters, differentiated for individuals at various infectious states, following the

approach of Chang and De Jong (10). The model assumes that infectious animals

shed *MAP* into the environment at their shedding rate, and once present, *MAP*

contamination level in the environment decays exponentially according to *MAP*'s

natural decay rate. The cumulative exposure is determined by integrating

environmental contamination level. This approach allows us to capture transmission

both in the presence and absence of actively shedding individuals. By estimating all

necessary transmission parameters to support the application of this model to *MAP*,

our goal is to improve our understanding of *MAP* transmission dynamics and to

guide interventions aimed at effectively controlling and potentially eradicating *MAP*.

Accurate assessment of infection risk in *MAP* requires a thorough understanding of

age-specific susceptibility. Newborn calves, probably because of their heightened

intestinal permeability, are particularly vulnerable to *MAP* during the first 24 hours of

life (14, 15). Susceptibility decreases with age, with cows generally considered

susceptible only during their first year of life (16). However, infections in adult cows

have been observed, but only when exposed to very high levels of *MAP*, such as

through oral dosing of 180 mg of cultured MAP or housing with 15 high shedders for

4 years (14, 17). The commonly used exponential model for age-specific

susceptibility assumes a decrease rate parameter of 0.1 wk^{-1} (16, 18-20). However,

this parameter estimate is a broad estimate and could possibly be improved. Our

analysis addressed this gap by developing an age-specific dose-response model

and estimating the susceptibility decrease rate parameter using data from fourteen

inoculation experiments that include age at exposure. The study aimed to more

accurately capture how susceptibility to *MAP* varies with age and dose, and to

enhance the precision of infection risk calculation.

This research developed both an age-specific dose-response model and two

stochastic environmental transmission models (Model A and B) to estimate

parameters and their 95% confidence interval (95%CI) based on fourteen dose-

response experiments and two transmission experiments. The parameters include

the parameters for susceptibility decrease rate (g), median infective dose (ID_{50}),

transmission rate (β or β_{state}), shedding rate (sh_{state} or sh), *MAP* decay rate in the

environment (de), where $state$ is either transiently infectious (Itr), low shedding (Il)

or high shedding (Ih). This classification of shedding patterns is consistent with the

observed shedding patterns from transmission experiments. With the modes and the

estimated parameters, we then explored four scenarios: changes in contamination

levels over time, timing and duration of exposure, and the recipient's age at exposure.

Materials and Methods

Age-specific susceptibility decrease rate

Literature search was conducted focusing on *MAP* inoculation experiments

performed on calves with different age groups. Data were collected from

experiments that provided detailed information, including inoculation dose, bacterial

strain, number of experimental animals, age at inoculation, inoculation methods,

diagnostic methods, and binary results (infected/not infected) (2, 3, 21-35). The initial

susceptibility of new-borns is standardized to be 1 and it decreases exponentially

with age at a rate of g. ID_{50} is defined as the dose at which a newborn calf, when

exposed, would have a 50% probability of becoming infected. To estimate g and

ID_{50}, we constructed an age-specific dose-response model to calculate relative

susceptibility ($rS(age)$), based on exponential function (Equation 1). The maximum

likelihood method is applied on a stochastic model with the number of new infections

following a binomial distribution, with S trials and a dose/age dependent infection

probability $p(dose, age)$ (Equation 2). In the model, g is regarded as an adjustment

factor influencing ID_{50}, which implies that older individuals require a higher dose of

MAP to attain a 50% chance of infection compared to younger individuals.

$$rS(age) = e^{-g \times age} \#(1)$$

$$p(dose, age) = 1 - e^{-\left(rS(age) \times \frac{\ln(2)}{ID_{50}}\right) \times dose} \#(2)$$

117　Besides, to calculate the infection probability over an exposure period, where the

118　age increases simultaneously over time, the average relative susceptibility can be

119　calculated as $\overline{rS}(age_i, age_i + \tau) = \frac{1}{\tau} \int_{age_i}^{age_i+\tau} rS(\alpha)d\alpha$. This calculation provides the

120   mean susceptibility over the time interval $(i, i + \tau)$, accounting for how susceptibility

121   changes as the individual ages from age_i to $age_i + \tau$.

Transmission rate parameters

MAP experimental transmission data were obtained from experiments conducted by

van Roermund et al. (36), which involved 12 cows and 20 calves, and aimed to

replicate real-world cow-to-calf and calf-to-calf transmission scenarios. In the initial

phase of the experiment, six inoculated adult cows were placed in a row, and five 1-

week-old naive calves(S) were placed in between two successive cows. The

positions of calves were altered randomly every two weeks over the period of the

three-month experiment, ensuring each calf was equally exposed to all pairs of cows

in the row. Sampling of all individuals occurred once every two weeks, using three

diagnostic methods, i.e., serum ELISA, bacterial culture, and interferon-gamma tests.

Cows with at least one positive result (based on faecal culture) of more than 100

colonies of *MAP* per tube (0.16 g faeces) out of the seven tests were categorized as

high shedders (Ih), while those with at least one positive result or no positive result

were categorized as low shedders (Il) or non-infectious, respectively. This

classification was also applied throughout our modelling.

Subsequently, in the second phase of the experiment, the previously exposed five

calves (including susceptible, latently infected (L), and transiently infectious calves

(Itr)) were brought into contact with five new 1-week-old naive calves for a period of

three months. Within the calf pen, all 10 calves had unrestricted contact with each

other. Like before, the three diagnostic methods were conducted once every two

weeks on each calf. The whole experiment was repeated once, with again 6 cows, 5

naïve calves, and these were again brought in contact with 5 additional calves, to

verify consistency and accuracy of the findings. Regular pen cleaning was performed

weekly, without additional disinfection. All calves were fed artificial milk replacer,

eliminating the possibility of transmission through contaminated milk. A summary of

the original data from published experiments is provided in S1 File.

Based on data from these published experiments, we constructed two stochastic

environmental transmission models (A and B) to estimate three key parameters in

MAP transmission dynamics: i.e. those for shedding rate, transmission rate, and

MAP decay rate. According to the model assumptions, the environmental

contamination level ($E_{state}(t)$) is determined by the presence and shedding rate of

infectious individuals at different states ($state = Itr$, Il or Ih), as well as the

pathogen's decay rate in the environment (Equation 3). Specifically, Model A

captures infectivity differences through varying transmission rate parameters, while

Model B captures them through varying shedding rate parameters. The detailed

schemes for both models are provided in S2 File.

$$\frac{dE_{state}(t)}{dt} = sh_{state}\, I_{state,t} - de\, E_{state}(t) \#(3)$$

161   with $E_{state}(0) = 0$

163   By solving Equation 3, $E_{state}(t+\tau)$ at the end of the interval $(t, t+\tau)$ can be

164   expressed explicitly, when $E_{state}(t)$ and $I_{state,t}$ are known :

$$E_{state}(t+\tau) = \frac{(1 - e^{-de\,\tau})}{de} sh_{state}\, I_{state,t} + e^{-de\,\tau} E_{state}(t) \#(4)$$

In the environment, the total contamination is contributed by all infectious individuals

in the environment. As we assume individuals' infection states remain constant over

a time interval $(i, i + \tau)$, the total force of infection $\lambda tot_{i,i+\tau}$ is calculated by summing

contributions from each individual m in the infectious compartment I_i, where

$m \in \{1,2 \dots I_i\}$. Specifically, an individual m contributes $\int_i^{i+\tau} E_{state_m}(t) \, dt$ to the

environment, which reflects the amount of MAP shed during the interval $(i, i + \tau)$,

when the instantaneous contamination level $E_{state_m}(t)$ is given by Equation 4. For

the force of infection rate calculation, in the transmission-rate-based Model A, this

quantity (of accumulated contamination) is multiplied by the corresponding

transmission rate parameter β_{state_m}, representing the infectivity level of that shedding

state. Alternatively, in the shedding-rate-based Model B, shedding parameters vary

by $state_m$ and a constant transmission rate applies to all states.

$$\lambda tot_{i,i+\tau} = \sum_{m=1}^{I_i} \left(\beta_{state_m} \times \int_i^{i+\tau} E_{state_m}(t) \, dt \right) \#(5)$$

179   Then, the infection probability $(p_{i,i+\tau})$ for a recipient at age $(age_i, age_i + \tau)$, during

180   time interval $(i, i + \tau)$ equals:

$$p_{i,i+\tau} = 1 - e^{-\frac{\lambda tot_{i,i+\tau} \times \overline{rs}(age_i, age_i + \tau)}{N_i}} \#(6)$$

Here, N_i denotes the size of environment (37).

Although, each individual was tested every two weeks, we produced weekly model

outputs ($\tau = 1$ wk) to capture new infections while the underlying model runs in

continuous time. We defined the time of infection as the midpoint between the last

negative test and the first positive test. However, in experiments B1 and A2, six

calves remained negative throughout the contact period but tested positive only

during the subsequent individual housing period. In these cases, we assumed the

calves bypassed the transient shedding stage and entered a latent phase directly.

With no further data available, we assumed the midpoint of the experiment as the

most likely time of infection for these animals. The maximum likelihood method is

used to fit the model to observed infection events, with the estimates determined at

the minimal Akaike Information Criterion (AIC) value. The 95%CI are determined by

adding 2 to the minimum AIC value(38). Equations for maximum likelihood estimate

($\widehat{\mathcal{L}(\theta)}$) and AIC are listed below:

$$\widehat{\mathcal{L}(\theta)} = Max_\theta \left[\prod_i \binom{S_i}{case_i} \left(1 - p_{i,i+1}(\theta)\right)^{case_i} p_{i,i+1}(\theta)^{(S_i - case_i)} \right] \#(7)$$

$$AIC = 2k - 2\log\left(\widehat{\mathcal{L}(\theta)}\right) \#(8)$$

with $\theta = \{de, \beta_{Itr}, \beta_{Il}, \beta_{Ih}\}$ in model A, the shedding rate sh is a function of de as

follows: $sh(de) = \frac{de^2}{-1 + e^{-de\,\Delta t} + de\,\Delta t}$, with Δ$t$=1 wk (10). Or, in model B, with $\theta =$

$\{de, \beta, sh_{Il}, sh_{Ih}\}$ and $sh_{Itr}(de) = \frac{de^2}{-1 + e^{-de\,\Delta t} + de\,\Delta t}$, with Δ$t$=1 wk. The number of

unknow parameters k, equals 4.

As shown above, β_{Itr} in model A equals to β in model B, and sh in model A equals to

sh_{Itr} in model B. This is because Itr is the reference infectious state in both models,

which can be explained by the unit definition of the model. The unit of $E_{state}(t)$ differs

in scale from the dose unit (e.g. CFU for *MAP*). Instead, it is standardized such that,

in a clean environment, the $Exposure$ from one infectious individual at state Itr over

a week is set to one unit ($\int_0^1 E_{Itr}(t)\,dt$=1). This standardization ensures a consistent

interpretation of the transmission rate parameter and shedding rate parameter

across this research and other studies with different transmission mechanism

assumptions(10).

Model fit evaluation

Model fit is assessed by comparing the predicted infection probabilities with the

actual observations from the experiments. The experiment begins at time 0, and at

any subsequent time point t, $p_{0,t}$ can be calculated using the Equation 6. The total

infection probability over the whole experiment duration and the expected number of

new infections can be calculated ($E(New\ case) = S_0 \times p_{0,end\ of\ experiment}$). This

evaluation helps to determine how accurately the model reflects the true dynamics of

infection transmission and provide insights into how the underlying infection process

may look like in the real-world experiments.

Scenario study

In this research, a scenario study was included to show the interpretation of model A

and B, and the estimated parameters under more real-world conditions. The study

focuses on four key drivers of transmission: (1) Duration of contamination: examining

the impact of retaining an infectious individual (at state $Itr,\ Il$ or Ih) in a clean

environment, simulating changes in the contamination level over time ($E_{state}(t)$), the

force of infection rate over time($\lambda_{state}(t)$) and calculating time (t_{eq}) required to reach

equilibrium phase. (2) Timing of exposure: calculating infection probabilities

associated with short-term exposures before equilibrium (when $E_{state}(t)$ is still

increasing) and after equilibrium (when $E_{state}(t)$ has stabilized). To ensure

meaningful comparisons, the exposure interval is set to 1 week, with the susceptible

individual's age at 0 weeks at the start of exposure. (3) Duration of exposure:

Investigating how varying exposure durations influence infection probability over time.

Similarly to Scenario 2, the assumptions are that $E_{state}(t)$ is stabilized and that the

recipient is fully susceptible at the start. (4) Age of the recipient: Predicting how

changes in the recipient's age, along with corresponding variations in susceptibility,

affect infection probability, assuming $E_{state}(t)$ is stabilized and the exposure duration

is fixed at 1 week. For both Scenario 3 & 4, the total simulated duration was based

on the maximum susceptible age of 52 weeks. Table 1 provides a summary of the

scenario setups, variables, variation ranges, and corresponding equations.

Table 1. Overview of Scenario Study Parameters and Objectives.

Scenario	Infectious state	Exposure starts time (Week)	Length of exposure interval (Week)	Recipient's age at start of exposure (Week)	Output
1	$Itr,\ Il$ or Ih	$t,$ $t \in [0,52]$	-	-	$E_{state}(t)$ and $\lambda_{state}(t)$
2	$Itr,\ Il$ or Ih	$x,$ $x \in [0,52]$	1	0	$p_{x,x+1} = 1 - e^{-\beta_{state} \times \int_x^{x+1} E_{state}(t)dt \times \overline{rs}(0,1)}$
3	$Itr,\ Il$ or Ih	eq^*	$y,$ $y \in [0,52]$	0	$p_{eq,eq+y} = 1 - e^{-\beta_{state} \times \int_{eq}^{eq+y} E_{state}(t)dt \times \overline{rs}(0,y)}$
4	$Itr,\ Il$ or Ih	eq^*	1	$z, z \in [0,52]$	$p_{eq,eq+1} = 1 - e^{-\beta_{state} \times \int_{eq}^{eq+1} E_{state}(t)dt \times \overline{rs}(z,z+1)}$

243   * eq can be set to any time point after $E_{state}(t)$ reaches its equilibrium (t_{eq}), which will
244   be informed by the outcome of scenario 1.
245

246   All calculations and simulations were performed using Mathematica 14.0, with

247   machine precision applied throughout the computations to ensure numerical

248   accuracy. All data and code used for parameter estimation, simulation, model fitness

249   analysis and scenario analysis are available in a Git repository at

250   https://git.wur.nl/yuqi.gao/map_para_esti.

Results

Age-specific susceptibility decrease rate

The dose-response model was used to estimate infectious Dose 50 (ID_{50}) and age-specific decrease rate (g) parameters from *MAP* inoculation experiments in calves of different ages. The fit of the non-linear model, based on an exponential function, results in an ID_{50} estimate and its 95%CI of 2.75×10^6 (1.42×10^6, 4.08×10^6) CFU, and a g estimate and its 95%CI of 0.0629 (0.0446, 0.0811) wk^{-1}. We plotted the predicted age-specific dose-response plot, overlaid with the original observations (dot sizes proportional to sample sizes) (Fig. 1).

Fig 1. Age-Specific Dose-Response Model Prediction. This figure illustrates the predicted age-specific dose-response relationship (blue surface), overlaid with observed data points scaled by sample size (larger dots = larger recipients' group).

For model fit evaluation, the 95% prediction interval (PI)—accounting for both model uncertainty and the binary nature of infection—encompassed all observed data. Additionally, we achieved the coefficient of determination (R^2) of 0.797 and adjusted R^2 of 0.796, which means our model explains nearly 80% of the variance. The minimal gap between R^2 and adjusted R^2 further suggests minimal overfitting.

Relatively susceptibility ($rS(age)$) decreases exponentially with age at a rate of

0.0629 wk^{-1} and the average susceptibility

$(\overline{rS}(age_i, age_i + \tau) = \frac{1}{\tau} \int_{age_i}^{age_i+\tau} e^{-0.0629 \times \alpha} d\alpha)$ within age range $(age_i, age_i + \tau)$ is

quantified by integrating the continuous $rS(age)$ function and diving it by the interval

length.

As shown in Fig. 2, our model estimates a slower decrease rate in susceptibility with

age compared to the previously published rate of 0.1 wk^{-1}. This suggests that

susceptibility decrease rate with age has been overestimated, while the infection risk

has been underestimated.

Fig. 2. Comparison of Susceptibility Decrease with Age. The solid line and

shaded region represent the relative susceptibility decrease ($rS(age)$) with age

based on our model-estimated rate (g = 0.0629 wk^{-1}) and its 95% confidence

interval. The dashed line shows the relative susceptibility decrease using the

previously published rate of 0.1 wk^{-1}.

Transmission parameters estimation

The environmental transmission model parameters were estimated from

experimental cow-to-calf and calf-to-calf transmission data using maximum likelihood

estimation. These included the decay rate parameter estimate for *MAP* in the

environment ($\widehat{de}$); transmission rate parameter estimates, which were state-specific

($\widehat{\beta_{Itr}}$, $\widehat{\beta_{Il}}$, $\widehat{\beta_{Ih}}$) in Model A, and constant ($\hat{\beta}$) in Model B; and shedding rate estimates,

which were constant ($\widehat{sh}$) in Model A, and state-specific ($\widehat{sh_{Itr}}$, $\widehat{sh_{Il}}$, $\widehat{sh_{Ih}}$) in Model B.

In both models, the reference infectious state is Itr, which means that $\widehat{\beta_{Itr}}$ model A is

equivalent to $\hat{\beta}$ in model B, and $\widehat{sh}$ in model A is equivalent to $\widehat{sh_{Itr}}$ in model B.

Profile likelihood visualizations for each parameter are shown in Fig 3, with detailed

results in Table 2.

Fig 3. Profile-Likelihood Plots for Shedding, Transmission, and Decay Rate

Parameters. The upper panel represents results from Model A, where the lower

panel represent Model B. The best estimate for each parameter corresponds to the

lowest AIC value, marked by a red star. The black dots represent the critical values

for the 95% confidence interval, which is defined by minimum AIC+2 (the dashed

line).

Table 2. Profile Likelihood Estimates and 95% Confidence Intervals for

Transmission, Shedding, and Decay Rate Parameters in Model A and B

	Transmission Rate ($\hat{\beta}$) (Unit: wk^{-1})			Shedding Rate ($\widehat{sh}$) (Unit: wk^{-1})			Decay Rate ($\widehat{de}$) (Unit: wk^{-1})	AIC
	$\widehat{\beta_{Itr}}$	$\widehat{\beta_{Il}}$	$\widehat{\beta_{Ih}}$	$\widehat{sh_{Itr}}$	$\widehat{sh_{Il}}$	$\widehat{sh_{Ih}}$		
Model A	0.0299 (0.0141, 0.111)	0.0150 (0,0.134)	0.219 (0.178, 0.536)	2.10 (2.00, 2.29)			0.150 (0,0.416)	72.618
Model B	0.0299 (0.0189, 0.0445)			2.10 (2.01, 2.26)	1.05 (0, 5.92)	15.4 (8.50, 25.0)	0.150 (0.00795,0.376)	72.618

Model fit evaluation results

Model fit is assessed by comparing predicted infection probabilities with actual experimental observations. During Experiment A1, 3 high shedders and 3 low shedders were observed; during Experiment A2, 4 transient shedders were present. During Experiment B1, 1 high shedder, 3 low shedders, and 2 non-infected cows were observed, while during Experiment B2, no infectious individuals were present. As detailed in the Materials & Methods section, in Experiments A1 and B1, 6 cows were housed in individual pens, with 5 recipient calves placed between each pair of cows, and calves' positions were randomly switched every two weeks. For simplicity, this setup was treated as a "pairwise" experiment, where one cow and one calf are paired. As there was no specific information regarding the location of cows and calves in the experiment A1 or B1, the infectivity of the shedder in each pair was considered as the average across all 6 infected cows. In experiments A2 and B2, five shedders and five recipient calves were placed in the same pen, allowing for unrestricted contact between all animals. Based on the number and infectious state of individuals in each experiment, the environmental contamination levels ($E_{state}(t)$),

population force of infection ($\lambda tot_{0,t}$), and infection probabilities ($p_{0,t}$) were calculated

according to Equations 3-6.

As shown in Fig 4, the curve and shaded area (left axis) represent the predicted

infection probabilities over time (0, t), while the bar plot (right axis) displays the

observed data from the experiments. The visualization indicates a good fit of the

model, as the predictions aligns with the observed data, demonstrating that the

model effectively captures the dynamics of infection transmission. Besides, the total

infection probabilities over the whole experiment duration and the predicted number

of infections were calculated. In both model A and model B, to three decimal places,

the predicted overall infection probabilities for experiment A1, A2, B1, B2 are 97.8%,

45.2%, 76.3%, 0.00%. As there are five recipients in each experiment, the expected

number of new infections is calculated to be 4.89, 3.81, 2.26, 0, respectively, while 5,

4, 2, 0 infections were observed.

Fig 4. Comparison of Predicted Accumulative Infection Probabilities with

Experimental Observations in Model A and B. The upper panel represents results

from Model A, where the lower panel represent Model B. The four plots in each panel

show parallel cow-to-calf experiments (A1 & B1) and parallel calf-to-calf experiments

(A2 & B2). The solid curve and shaded area represent the predicted accumulative

infection probabilities and their 95% confidence interval from the start of

experiments(t=0) to time t. The blue bars indicate the number of newly infected

individuals observed in the experiments. The x-axis denotes time in weeks, with the

left y-axis showing infection probability and the right y-axis showing the number of

susceptible and infected individuals.

Scenario study results

To better interpret these parameters and explore how different biological

assumptions about infectivity influence predicted exposure, we conducted a scenario

study. Specifically, we simulated changes in environmental contamination level and

force of infection rate over time and evaluated the effects of exposure timing,

duration, and the recipient's age on infection probability (Table 2.). However, the

decay rate exhibited an extremely lower bound (0 for Model A and 0.00795 for Model

B), making it difficult to discern the impact of the other parameters. To focus on the

comparison of Model A and B, we therefore fixed the decay rate at its best-estimate

value. Consequently, the 95%CI present later for scenario analysis results reflect

only the uncertainty from the shedding, transmission, age-specific susceptibility

decrease rate parameters. Figures 5 and 6 present the results of this scenario study.

In Scenario 1, $E_{state}(t)$, $\lambda_{state}(t)$, and t_{eq} were calculated for environments with the

presence of an infectious individual in states $Itr, \ Il$ or Ih. The 95%CI were derived

from uncertainties in transmission and shedding rate parameters. As shown in Fig 5,

$E_{state}(t)$ rapidly increase and then stabilize at equilibrium, requiring 30.6 weeks to

reach 99% of the environmental contamination equilibrium. Following the methods in

Materials and Methods, we set teq =60 wk for subsequent analyses.

Fig 5. Result of Scenario 1: Changes in $E_{state}(t)$ and $\lambda_{state}(t)$ over time while

there is one Infectious individual in states Itr, Il or Ih: In Model A (left column),

infectious individuals in different states have the same shedding rate, while in Model

B (right), $E_{state}(t)$ increases at different rates due to varying shedding rates among

infectious states. The natural decay rate of MAP is constant in both models.

Scenario 2 assessed the infection probability for exposing a fully susceptible

individual to the previously defined three types of environments for 1 week, with the

variable being the timing of exposure x. As shown in Fig 6-upper row, the infection

probabilities firstly increased and then stabilized at 33.2% for Itr, 18.3% for Il, and

94.9% for Ih. In Scenario 3, extending the exposure time y results in a rapid increase

in infection probabilities. For Ih, the infection probability approaches 100% after

approximately 1.83 weeks. However, for Itr and Il, the probabilities do not reach 100%

due to the decreasing susceptibility over age/time but still reach 99.8% and 95.9%

respectively by the end of 52 weeks. In Scenario 4, infection probabilities decrease

with age

. By the end of calfhood (52 weeks old), after one week of exposure to a

stabilized contaminated environment, the predicted infection probabilities are: 1.53%

for , 0.767% for , 10.7% for . Therefore, we suggest heifer/cow above one

year should also be considered susceptible. In all four scenario analysis, although

model A and B produce the same mean prediction values, Model B has wider 95%CI

for low and high shedders but a smaller 95%CI for transient shedder.

Fig 6. Results of Scenarios 2,3,4: Upper: Predicted infection probabilities for a fully

susceptible individual exposed to the contaminated environment at time (for

[0,52]) after the start of contamination (=0), with each exposure lasting for a 1-

week interval; Middle: Predicted infection probabilities for a fully susceptible

individual exposed to the contaminated environment at equilibrium, across varying

exposure durations y (for $y \in$[0,52]); Lower: Predicted infection probabilities for

susceptible individuals of age z (for $z \in$[0,52]) exposed to the contaminated

environment at equilibrium, over a 1-week exposure interval. Solid, dashed, and

dotted lines represent the infectious individual at states Itr, Il, and Ih, respectively.

The shaded areas represent the corresponding 95% confidence intervals, calculated

based on the 95% confidence intervals of transmission rate parameters, shedding

rate parameter and the age-specific susceptibility decrease rate.

This research aims to improve understanding of *MAP* transmission dynamics

through a novel mathematical model, which has not been previously applied to *MAP*.

While comparisons with earlier studies are insightful, differences in transmission

mechanism assumptions do complicate direct comparisons. However, the force of

infection rates and infection probabilities can still be helpful for evaluating our

findings alongside previous research. Therefore, we used a different and commonly

used dose-response transmission model (19) and the parameters relevant to that

model to replicate the model fit evaluation and scenario analyse. Both models fit the

cow-to-calf experimental observations reasonably well; but a significant difference is

noticed in calf-to-calf transmission. The predicted number of new cases over the

entire period of experiments A1, A2, B1, and B2, based on dose-response model

were 5.00, 0.00, 3.98, and 0.00, respectively. These predictions differ from the

observed values (5, 2, 4, and 0, respectively), suggesting that the dose-response

transmission model may not adequately capture calf-to-calf transmission. Detailed

model descriptions, results, and discussions are provided in S4 File.

Discussion

The aims of this research were: firstly, to develop an age-specific dose-response

model to study changes in recipient susceptibility over age; secondly, to construct an

environmental transmission model to quantify the parameters for the transmission

dynamics of *MAP*, i.e. estimate the shedding rate parameter, transmission rate

parameter and *MAP* decay rate parameter. A scenario study followed, focusing on

four objectives: changes in contamination levels over time, the timing of exposure,

the duration of exposure, and the recipient's age. Maximum likelihood estimation was

applied to achieve the best estimates, and model fit was assessed. Additional details,

including a summary of original exposure data from published experiments, culture-

based data on *MAP* viability in cattle excretions, and comparisons with transmission

parameters from previous research, are provided in supplementary materials.

Research has suggested that calf-to-calf transmission may be underestimated (39-

42). In this study, we identified that this underestimation could be partly due to the

use of a rounded value of 0.1 for the age-specific susceptibility decrease rate. Based

on the exposure data of fourteen experiments, we estimated a decrease rate

parameter of 0.0629 wk^{-1} (95% CI: 0.0446, 0.0811 wk^{-1}). As shown in Fig. 2, starting

from a standardized average susceptibility of 1 for newborns, this value declines to

0.5 after 11 weeks and further drops to 0.0380 by 52 weeks of age. In contrast, the

commonly used rate of 0.1 wk^{-1} results in susceptibility dropping to 0.5 at 6.93

weeks and to 0.00552 by 52 weeks. This suggests that if a 52-week-old calf is

exposed to a stabilized contaminated environment from Ih for one week, the

expected infection probability would be 0.927%, which is significantly lower than our

estimate of 10.7%. Our prediction aligns more closely with observations of new

infections in heifers and older calves (2, 22), and could therefore improve the

evaluation of calf-to-calf transmission and inform more effective interventions.

Additionally, to verify the estimated $\widehat{de}$, we compare it with culture-based data on

MAP viability in cattle excretions. For instance, in faecal pellet samples collected

from boxes in the 100% shade treatment at 10-22℃, the decay rate was estimated to

be 0.189 wk^{-1}. Similarly, in cattle feces at site Armidale, Bathurst, Condo Bolin, under

temperatures of 3.6–19.7℃, 6.8–19.8℃, 10.2–24.4℃, the decay rates were 0.118

wk^{-1}, 0.242 wk^{-1}, and 0.242 wk^{-1}, respectively. These values provide a range for

comparison with the model's predictions(7, 14, 43-49). Further details can be found

in S3 File.

Transmission parameters were estimated using data from cow-to-calf and calf-to-calf

transmission experiments. The study applied two models—Model A and B—to

capture variations across different infectious states (Itr, Il, Ih) in *MAP* transmission.

Although Models A and B yield identical point estimates and AIC values, Model B

consistently shows wider 95% confidence intervals. One likely explanation is the

difference in parameter scaling. In Model A, infectivity is captured through

transmission rates with a standardized shedding rate, whereas in Model B, the

shedding rates vary while transmission remains constant. Because shedding rates in

Model B can span a broader numerical range, small relative changes in shedding

can produce large changes in the predicted outcomes, leading to wider confidence

intervals.

To interpret the estimates, a scenario study was conducted. The results indicate that

MAP accumulates quickly in the environment and takes approximately 30.6 weeks to

reach 99% of its equilibrium value. At equilibrium, the highest $E_{state}(t)$ is associated

with Ih, followed by Itr and Il, though the difference between the latter two is much

smaller. However, Il exhibits a larger 95% CI, and more importantly, Il could pose a

greater overall hazard due to its longer shedding period (50). Besides, the scenario 3

shows that the exposure duration greatly affects infection probability, assuming

exposing a fully susceptible recipient in a stabilized contaminated environment, the

infection probabilities reach 100% for Ih after 3.31 weeks, 95.9% for Il after 52

weeks and 99.8% for Itr after 52 weeks. The probabilities for Itr and Il didn't reach

1 because increasing exposure duration also means the recipient's age increases,

leading to a decrease in relative susceptibility.

In our study, the transmission rate in model A (or shedding rate in model B) for

transient shedders is roughly twice that of low shedders; however, experimental data

suggest only about half as many infected animals enter the transient shedding phase

compared to the low-shedding phase. For example, in Experiments A1 and B1,

nine calves were infected, but only four began shedding as Itr in Experiment A2.

Among all 11 infected calves (nine from A1 & B1 and two from A2 & B2), nine

eventually began shedding during the three-year individual housing period after

transmission experiments. These observations align with the transient transmission

period commonly assumed in modelling studies, where animals briefly shed before

entering latency and later resuming low or high shedding. However, when infection

histories are reconstructed solely from diagnostic results, distinguishing transient

shedders from low shedders—especially in animals infected later in life—can be

challenging. We therefore recommend that future backward inference models

carefully account for this concern.

A limitation of the current study is the absence of a dedicated environmental

transmission experiment—specifically, one that introduces susceptible individuals

into a contaminated environment without the presence of infectious individuals.

Conducting such an experiment in the future could improve the accuracy of

parameter estimation. Besides, the application of the estimated parameters in real

farm models is more complex than in controlled lab experiments. On farms, a larger

number of infected and susceptible animals at varying states and ages are involved,

alongside continuous contamination processes, limited diagnostic capabilities, and

more challenging hygiene management due to larger spaces. Next, multiple

interventions are widely applied, further complicating quantifying the transmission

dynamics. Future research should develop a continuous-time, age-structured

compartmental stochastic model that simulates *MAP* transmission at the farm level.

Although several simulation models already exist, the refined transmission

parameter estimates from this research have the potential to enhance simulation

accuracy and improve overall reliability. Such a model would incorporate factors like

age cohorts, animal grouping and movement, farm management, environmental

contamination, and real-world interventions.

Reference

1.   Johne H, Frothingham L. Ein eigenthuemlicher fall von tuberculose beim rind. Dtsch Z
Tiermed Pathol. 1895;21:438-54.

2.   Mortier RAR, Barkema HW, Bystrom JM, Illanes O, Orsel K, Wolf R, et al. Evaluation of
age-dependent susceptibility in calves infected with two doses of Mycobacterium avium
subspecies paratuberculosis using pathology and tissue culture. Veterinary Research.
2013;44(1):94.
3.   Mortier RAR, Barkema HW, De Buck J. Susceptibility to and diagnosis of
Mycobacterium avium subspecies paratuberculosis infection in dairy calves: A review.
Preventive Veterinary Medicine. 2015;121(3):189-98.
4.   Council NR. Diagnosis and Control of Johne's Disease. Washington, DC: The National
Academies Press; 2003. p. 243.
5.   Nielsen S, Ersbøll A. Age at occurrence of Mycobacterium avium subspecies
paratuberculosis in naturally infected dairy cows. Journal of dairy science. 2006;89(12):4557-
66.
6.   Nielsen SS. Transitions in diagnostic tests used for detection of Mycobacterium avium
subsp. paratuberculosis infections in cattle. Veterinary Microbiology. 2008;132(3):274-82.
7.   Eppleston J, Begg DJ, Dhand NK, Watt B, Whittington RJ. Environmental Survival of
Mycobacterium avium subsp. paratuberculosis in Different Climatic Zones of Eastern
Australia. Applied and Environmental Microbiology. 2014;80(8):2337-42.
8.   Aly SS, Thurmond MC. Evaluation of Mycobacterium avium subsp paratuberculosis
infection of dairy cows attributable to infection status of the dam. Journal of the American
Veterinary Medical Association. 2005;227(3):450-4.
9.   Whittington RJ, Windsor PA. In utero infection of cattle with Mycobacterium avium
subsp. paratuberculosis: A critical review and meta-analysis. The Veterinary Journal.
2009;179(1):60-9.
10.   Chang Y, de Jong MCM. A novel method to jointly estimate transmission rate and
decay rate parameters in environmental transmission models. Epidemics. 2023;42:100672.
11.   Chang Y, Hartemink N, Byrne AW, Gormley E, McGrath G, Tratalos JA, et al. Inferring
bovine tuberculosis transmission between cattle and badgers via the environment and risk
mapping. Front Vet Sci. 2023;10:1233173.
12.   Gamża AM, Hagenaars TJ, Koene MGJ, de Jong MCM. Combining a parsimonious
mathematical model with infection data from tailor-made experiments to understand
environmental transmission. Scientific Reports. 2023;13(1):12986.
13.   Gao Y, Boklund AE, Nielsen LH, Alban L, de Jong MCM. Estimating the impact of low
temperature on African swine fever virus transmission through contaminated environments.
Preventive Veterinary Medicine. 2023;219:105991.
14.   Larsen AB, Merkal RS, Cutlip RC. Age of cattle as related to resistance to infection
with Mycobacterium paratuberculosis. Am J Vet Res. 1975;36(3):255-7.
15.   Hagan W. Age as a Factor in Susceptibility to Johne's Disease. 34–40 p.
16.   Windsor PA, Whittington RJ. Evidence for age susceptibility of cattle to Johne's
disease. The Veterinary Journal. 2010;184(1):37-44.
17.   Rankin JD. The experimental infection of cattle with Mycobacterium johnei. IV. Adult
cattle maintained in an infectious environment. J Comp Pathol. 1962;72:113-7.
18.   Doyle T. Susceptibility to Johne's disease in relation to age. Vet Rec. 1953;65:363-5.
19.   Marcé C, Ezanno P, Seegers H, Pfeiffer DU, Fourichon C. Predicting fadeout versus
persistence of paratuberculosis in a dairy cattle herd for management and control purposes:
a modelling study. Veterinary Research. 2011;42(1):36.

20.   Biemans F, Tratalos J, Arnoux S, Ramsbottom G, More SJ, Ezanno P. Modelling
transmission of Mycobacterium avium subspecies paratuberculosis between Irish dairy
cattle herds. Veterinary Research. 2022;53(1):45.
21.   Sweeney RW, Whitlock RH, Hamir AN, Rosenberger AE, Herr SA. Isolation of
Mycobacterium paratuberculosis after oral inoculation in uninfected cattle. Am J Vet Res.
1992;53(8):1312-4.
22.   Payne JM, Rankin JD. A Comparison of the Pathogenesis of Experimental Johne's
Disease in Calves and Cows. Research in Veterinary Science. 1961;2(2):175-9.
23.   Payne JM, Rankin JD. The Pathogenesis of Experimental Johne's Disease in Calves.
Research in Veterinary Science. 1961;2(2):167-76.
24.   Rankin JD. The Experimental Infection of Cattle with Mycobacterium Johnei: I. Calves
Inoculated Intravenously. Journal of Comparative Pathology and Therapeutics. 1958;68:331-
IN24.
25.   Rankin JD. The Experimental Infection of Cattle With Mycobacterium Johnei: II. Adult
Cattle Inoculated Intravenously. Journal of Comparative Pathology and Therapeutics.
1961;71:6-IN1.
26.   Stuart P. Vaccination Against Johne's Disease in Cattle Exposed to Experimental
Infection. British Veterinary Journal. 1965;121(7):289-318.
27.   Gilmour NJL, Nisbet DI, Brotherston JG. Experimental oral infection of calves with
Mycobacterium johnei. Journal of Comparative Pathology. 1965;75(3):281-6.
28.   Gilmour NJL, Angus KW. The specificity and sensitivity of the fluorescent antibody
test in cattle experimentally infected with Mycobacterium avium and Mycobacterium johnei.
Research in Veterinary Science. 1976;20(1):6-9.
29.   Johnson D, Muscoplat CC, Larsen A, Thoen C. Skin testing, fecal culture, and
lymphocyte immunostimulation in cattle inoculated with Mycobacterium paratuberculosis.
Am J Vet Res. 1977;38(12):2023-5.
30.   Lepper A, Wilks C, Kotiw M, Whitehead J, Swart K. Sequential bacteriological
observations in relation to cell‐mediated and humoral antibody responses of cattle
infected with Mycobacterium paratuberculosis and maintained on normal or high iron intake.
Australian veterinary journal. 1989;66(2):50-5.
31.   Uzonna JE, Chilton P, Whitlock RH, Habecker PL, Scott P, Sweeney RW. Efficacy of
commercial and field-strain Mycobacterium paratuberculosis vaccinations with recombinant
IL-12 in a bovine experimental infection model. Vaccine. 2003;21(23):3101-9.
32.   Waters WR, Miller JM, Palmer MV, Stabel JR, Jones DE, Koistinen KA, et al. Early
Induction of Humoral and Cellular Immune Responses during Experimental *Mycobacterium*
avium subsp. *paratuberculosis* Infection of Calves. Infection and Immunity. 2003;71(9):5130-
8.
33.   Stabel J, Palmer M, Whitlock R. Immune responses after oral inoculation of weanling
bison or beef calves with a bison or cattle isolate of Mycobacterium avium subsp.
paratuberculosis. Journal of wildlife diseases. 2003;39(3):545-55.
34.   Sweeney RW, Whitlock RH, McAdams S, Fyock T. Longitudinal Study of ELISA
Seroreactivity to Mycobacterium Avium subsp. Paratuberculosis in Infected Cattle and
Culture-Negative Herd Mates. Journal of Veterinary Diagnostic Investigation. 2006;18(1):2-6.
35.   Stewart DJ, Vaughan JA, Stiles PL, Noske PJ, Tizard MLV, Prowse SJ, et al. A long-term
bacteriological and immunological study in Holstein-Friesian cattle experimentally infected
with Mycobacterium avium subsp. paratuberculosis and necropsy culture results for

Holstein-Friesian cattle, Merino sheep and Angora goats. Veterinary Microbiology.
2007;122(1):83-96.
36.   van Roermund HJW, Bakker D, Willemsen PTJ, de Jong MCM. Horizontal transmission
of Mycobacterium avium subsp. paratuberculosis in cattle in an experimental setting: Calves
can transmit the infection to other calves. Veterinary Microbiology. 2007;122(3):270-9.
37.   Jong MCMd, Diekmann O, Heesterbeek JAP. How does transmission of infection
depend on population size? : Cambridge University Press http://www.cambridge.org/uk;
1995.
38.   Burnham K, Anderson. Information and Likelihood Theory: A Basis for Model
Selection and Inference. In: Burnham KP, Anderson DR, editors. Model Selection and
Multimodel Inference: A Practical Information-Theoretic Approach. New York, NY: Springer
New York; 2002. p. 49-97.
39.   Corbett CS, de Jong MCM, Orsel K, De Buck J, Barkema HW. Quantifying transmission
of Mycobacterium avium subsp. paratuberculosis among group-housed dairy calves.
Veterinary Research. 2019;50(1):60.
40.   Nigsch A, Robbe-Austerman S, Stuber TP, Paulina DPB, Gröhn YT, Schukken YH. Who
infects whom?—Reconstructing infection chains of Mycobacterium avium ssp.
paratuberculosis in an endemically infected dairy herd by use of genomic data. PLoS One.
2021;16(5).
41.   Vass-Bognár B, Bakony M, Baumgartner W, Khol JL, Jurkovich V. Association between
calf rearing technology and farm-level paratuberculosis infection in Hungarian dairy farms.
Preventive Veterinary Medicine. 2022;207.
42.   Weber MF, Kelton D, Eisenberg SWF, Donat K. Progress in Paratuberculosis Control
Programmes for Dairy Herds. Animals. 2024;14(7):1127.
43.   Olsen JE, Jørgensen JB, Nansen P. On the reduction of Mycobacterium
paratuberculosis in bovine slurry subjected to batch mesophilic or thermophilic anaerobic
digestion. Agricultural Wastes. 1985;13(4):273-80.
44.   EAMENS G, SPENCE S, TURNER M. Survival of Mycobacterium avium subsp
paratuberculosis in amitraz cattle dip fluid. Australian Veterinary Journal. 2001;79(10):703-6.
45.   Collins CH, Grange JM, Yates MD. Mycobacteria in water. Journal of Applied
Bacteriology. 1984;57(2):193-211.
46.   Hanifian S. Behavior of Mycobacterium avium paratuberculosis in Lighvan cheese
tracked by propidium monoazide qPCR and culture. LWT. 2020;133:109886.
47.   Klanicova B, Slana I, Roubal P, Pavlik I, Kralik P. Mycobacterium avium subsp.
paratuberculosis survival during fermentation of soured milk products detected by culture
and quantitative real time PCR methods. International Journal of Food Microbiology.
2012;157(2):150-5.
48.   Whittington RJ, Marshall DJ, Nicholls PJ, Marsh IB, Reddacliff LA. Survival and
dormancy of Mycobacterium avium subsp. paratuberculosis in the environment. Appl
Environ Microbiol. 2004;70(5):2989-3004.
49.   Jorgensen JB. Survival of Mycobacterium paratuberculosis in slurry. Nord Vet Med.
1977;29(6):267-70.
50.   Mitchell RM, Whitlock RH, Gröhn YT, Schukken YH. Back to the real world:
Connecting models with data. Preventive Veterinary Medicine. 2015;118(2):215-25.
664

Supporting information

S1 File. Collected experimental data from previous research.

S2 File. Detailed Schemes of Transmission Rates Based Model A and

Shedding Rates Based Model B.

S3 File. Validation of estimated decay rate parameter using culture-based

experimental data on *MAP* viability in cattle excretions.

S4 File. A Comparative Analysis with A Previously Published Transmission

Model