1 Title

2 Single-cell transcriptomic analysis of macaque LGN neurons reveals novel subpopulations

4 Authors

5 Shi Hai Sun[1,2*], Kai Renshaw[1,3, *], John S. Pezaris[1,2]

6 * Equal Contribution

7 Correspondence: John S. Pezaris; Pezaris.John@mgh.harvard.edu

9 Affiliations

10 [1]Department of Neurosurgery, Massachusetts General Hospital, Boston, Mass., USA

11 [2]Department of Neurosurgery, Harvard Medical School, Boston, Mass., USA

12 [3]Tufts University School of Medicine, Boston, Mass., USA

14 Conflicts of Interest

15 The authors declare no competing interests.

16 Funding

17 Supported by Peter and Yayi Pezaris, Don Good, Michael Gersh, NIMA Foundation, NIH

18 EY027888, and the William M. Wood Foundation, Bank of America, Trustee.

19 **Abstract**

Neurons in the lateral geniculate nucleus (LGN) provide a pivotal role in the visual system by

modulating and relaying signals from the retina to the visual cortex. Although the primate LGN,

with its distinct divisions (magnocellular, M; parvocellular, P; and koniocellular, K), has been

extensively characterized, the intrinsic heterogeneity of LGN neurons has remained unexplained.

With the development of high-throughput single-cell transcriptomics, researchers can rapidly

isolate and profile large sets of neuronal nuclei, revealing a surprising diversity of genetic

expression within the nervous systems, such as two types of K neurons (Bakken *et al.*, 2021).

Here, we analyzed the transcriptomes of individual cells belonging to macaque LGN using raw

data from a public database to explore the heterogeneity of LGN neurons. Using statistical

analyses, we found additional subpopulations within the LGN transcriptomic population, whose

gene expressions imply functional differences. Our results suggest the existence of a more

nuanced complexity in LGN processing beyond the classic view of the three cell types and

highlight a need to combine transcriptomic and functional assessments. A complete account of

the cell type diversity of the primate LGN is critical to understanding how vision works.

Introduction

The lateral geniculate nucleus (LGN) of the thalamus serves as the main relay through which visual information travels from the retina to the cortex (Sherman & Guillery, 2006; Solomon & Lennie, 2007). Within LGN, visual input is segregated both spatially – distinct areas of the visual field correspond to distinct areas of LGN (Chen et al., 1999; K. A. Schneider et al., 2004) – and functionally (Denison et al., 2014; K. A. Schneider et al., 2004). A bulk of research has focused on determining the precise spatial, functional, and morphological organization of LGN, as a better understanding of LGN's organization can provide insight into visual processing and has important clinical relevance.

In primates, LGN is composed of distinct layers of functionally similar cell types: magnocellular (M), parvocellular (P), and koniocellular (K). Electrophysiological and morphological studies in primates have determined the specific characteristics of these layers: M neurons, which are large in size and receive rod photoreceptor-derived information, are well-suited for rapid temporal dynamics and detecting motion; P neurons, receiving cone photoreceptor-derived information, are attuned to chromatic changes, making them optimal for processing color and form; and less well-defined K neurons receive short-wavelength ("blue") photoreceptor-derived information (Hubel & Livingstone, 1990; Maunsell et al., 1999; Reid & Shapley, 2002; Tailby et al., 2008; Wiesel & Hubel, 1966).

As methods for electrophysiological investigation improve, evidence has been accumulating for functional heterogeneity within the three primary LGN cell types. For instance, *on*- and *off*-specific responses from the retina are preserved within distinct populations of M, P, and K cells (Ichinose & Habib, 2022; Rousso et al., 2016). K cells have also been shown to

express receptive fields with non-standard features, including orientation sensitivity, contrast

suppression, and many others (Cheong et al., 2013; Eiber et al., 2018; Solomon et al., 2010).

Lastly, a recent electrophysiological analysis identified visually responsive LGN cells that did

not exhibit any M, P, or K functional responses, suggesting a functional complexity beyond the

M, P, and K classification (Sun et al., 2024). All this evidence points toward the existence of

multiple subpopulations of LGN neurons within M, P, and K cell types.

Most recently, single-cell RNA-sequencing technologies have been developed for

transcriptomic profiling, enabling the identification of cellular subclasses of cells by comparing

the gene expression patterns across large numbers of individual cells within a tissue sample (Lein

et al., 2017; Lowe et al., 2017; Ståhl et al., 2016). While past attempts to understand the

organization of LGN have relied primarily on morphology and electrophysiology, we

hypothesize transcriptomic analysis of LGN neurons will reveal more details about their

organization and functional roles in visual processing. For example, single-cell RNA-seq studies

in macaque have identified 13 transcriptomically separable excitatory cell classes in V1 (Wei et

al., 2022) and more than 60 distinct cell types in the retina (Peng et al., 2019), while only four

excitatory cell types were reported in the LGN (Bakken et al., 2021). In the present study, we

explore the transcriptomic diversity of macaque LGN neurons using raw data obtained from a

published database (Bakken et al., 2021). Statistical analyses of the LGN transcriptomes revealed

seven subpopulations of excitatory LGN neurons: two magnocellular populations, two

parvocellular populations, and three koniocellular populations. We show that these

subpopulations differentially express gene profiles, which imply specific functional differences

between these putative subclasses. Our results suggest that there is further nuance in the

functional organization of LGN that may be reflected in its role in visual processing.

Methods

RNA data

The RNA-seq data consisting of 2,157 macaque nuclei and accompanying metadata were

sourced from the Allen Brain Map (Allen Institute for Brain Science, 2022), available at

https://portal.brain-map.org/atlases-and-data/rnaseq/comparative-lgn, and originally used in

(Bakken *et al.*, 2021). For single-cell/-nucleus processing for RNA sequencing and processing,

please see Tasic *et al.* (2018); Bakken *et al.* (2018, 2021); Hodge *et al.* (2019) for the complete

details. In summary, nuclei were isolated by Fluorescence-Activated Cell Sorting, enriched for

neurons with neuronal markers, processed with SMART-seq v4 (Clontech) and Nextera XT

(Illumina), and sequenced on HiSeq 2500 (Illumina). RNA-seq reads were aligned to

corresponding genomes using the STAR aligner (Dobin et al., 2013), and samples had a median

depth of 1.3 million reads/nucleus.

Preprocessing

Data preprocessing and analyses were performed in R using Seurat v5 (Butler et al., 2018; Hao et

al., 2024; Stuart et al., 2019) as Seurat consistently outperforms other modern algorithms (Fu et

al., 2025). We used a customized version of the standardized single-cell transcriptomics analysis

workflow (Clarke et al., 2021), as described below.

Using the supplied metadata from the (Allen Institute for Brain Science (2022) and

Bakken *et al.* (2021), low-quality cells (cells that did not pass quality control criteria: <100,000

total reads, <1,000 detected genes, < 75% of reads aligned to genome, or CG dinucleotide odds

102 ratio > 0.5; Tasic *et al.*, 2018) and genes (e.g., uncharacterized or non-informative genes) were

103 removed from all further analyses. All cells were prelabeled as M, P, K (Kap and Kp), GABA

104 (GABA 1-4), or pulvinar, as defined in the metadata. Since this study is a comparison of LGN

105 M, P, and K excitatory cells, GABA and pulvinar cells were removed from further analyses

106 (61% remaining, *n* = 1,309 / 2,157).

Clustering

M, P, and K cell sets were processed separately using cluster analysis to explore the

heterogeneity of LGN neurons. Before clustering, the data was preprocessed and normalized

using the Seurat function *SCTransform*, which uses a regularized negative binomial regression

model for normalization and analytic Pearson residuals to find top variable genes for

downstream analysis (Choudhary & Satija, 2022; Lause et al., 2021). For batch correction, we

used *IntegrateLayers* with *HarmonyIntegration*, which integrates cells from multiple datasets

(Korsunsky *et al.*, 2019). The processed data was then dimensionally reduced using principal

component analysis (PCA, via *RunPCA*). Clustering was performed using *FindNeighbors,* K-

nearest neighbor (KNN), and *FindClusters*, which uses the Louvain algorithm for cluster

identification. The clustering results were visualized using uniform manifold approximation and

projection (UMAP, via *RunUMAP*; McInnes *et al.*, 2020), and quantitatively evaluated using

silhouette scores to assess cluster separation.

To find the optimum parameters to use for the analysis, a global parameter optimization

strategy using silhouette scores as the objective metric was used. Specifically, for each of the M,

P, and K cell subsets, the optimum number of principal components (PCs), number of features

(highly variable genes in *SCTransform*), and clustering resolution (*FindClusters*) were

determined.

First, to obtain the optimal number of PCs, the standard deviation of each PC was plotted

across a range of feature counts (1,000 to 4,000 in increments of 500) using Seurat's *ElbowPlot*

function. The point where the curve plateaued was selected qualitatively as the optimal number

of PCs for each of the M, P, and K cell types. Because the location of the elbow is subjective, we

followed the recommendation of the Seurat developers to favor the higher side when choosing

this parameter.

Next, a global parameter search was conducted to identify the combination of feature

number and clustering resolution that maximized the average silhouette width. The number of

features ranged from 1,000 to 4,000 in steps of 100 and the clustering resolution values spanned

from 0.1 to 1.0 in increments of 0.02. For each combination, clustering was performed following

dimensionality reduction by PCA, using the optimum number of PCs as defined above, and

silhouette scores were computed. A Generalized Additive Model (GAM) was then fitted to

smooth the silhouette score surface and remove outliers. The parameters corresponding to the

peak predicted silhouette score were selected for each cell type group. The final clustering

parameters for M, P, and K cell types are specified in the Results section. See Figure 1B for a

visualized map of the described clustering workflow.

Clustering validation testing

To assess robustness, a sensitivity analysis was performed. This included random downsampling

of each dataset to 70-90% of the original cells, as well as ±20% variation in the number of highly

variable genes and ±1 variation in the number of dimensions used for clustering. This process

was bootstrapped and repeated 1,000 times, and the findings are referenced in the Results and

Discussion sections.

To assess the clustering stability we added Gaussian noise directly to the Harmony

embeddings, scaled to 20% of the standard deviation (SD) of each embedding dimension. For

each cluster group, we performed 1000 iterations of added noise while preserving the original

cluster labels, and computed the silhouette score for each perturbed embedding. An empirical *p*-

value was calculated as the proportion of noise-affected silhouette scores that were equal to or

greater than the silhouette score from the original data. Cluster degradation was defined as the

percentage decrease in silhouette relative to the original value.

As an external validation of the clustering approach, an in-silico benchmark was

conducted using a publicly available, cell-sorted peripheral blood mononuclear cell (PBMC)

single-cell RNA-seq dataset with established ground-truth labels from Wang *et al.* (2019),

downsampled from Zheng *et al.* (2017). The dataset was first clustered using the standard Seurat

workflow, consistent with methods used in Bakken *et al.* (2021), to broadly identify the immune

cell populations. Subsequently, the heterogeneous cluster from the initial analysis was re-

clustered using the optimized silhouette-based global parameter search pipeline, as described

above. To evaluate stability, the analysis was also repeated under downsampling conditions and

across multiple bootstrap iterations. The details and results of this validation experiment are

presented in Supplementary Figure 1 and referenced in the Results.

Differentially expressed genes

Each cluster was profiled through differential gene expression for comparative analysis.

Differentially expressed genes/markers (we will use *genes* and *markers* interchangeably) were

determined with the Seurat function *FindAllMarkers*, defined with an adjusted *p*-value < 0.05

(Wilcoxon Rank Sum test). We also defined these genes as biologically significant if the $\log_2$

fold change is greater than 1 (i.e., log2FC > 1 and adjusted *p*-value < 0.05). Gene expression

profiles were visualized using heat maps (*DoHeatMap*) and dot plots (*DotPlot*). For the gene

ontology analysis, ToppGene (https://toppgene.cchmc.org/enrichment.jsp) and the National

Center for Biotechnology Information (NCBI) Gene database

(https://www.ncbi.nlm.nih.gov/gene/) were used. As the interpretation of gene ontology results is

necessarily speculative, most of this analysis is presented in the Discussion section.

Results

We analyzed the transcriptomes of 1,309 nuclei from microdissected, anatomically defined

regions of the macaque lateral geniculate nucleus (LGN) using RNA-seq data sources from the

Allen Brain Map (Allen Institute for Brain Science, 2022; Bakken et al., 2021). The RNA-seq

data was preprocessed by removing low-quality cells and unnecessary genes (see Methods), as

per the metadata provided by the Allen Institute (Allen Institute for Brain Science, 2022). Like in

Bakken *et al.* (2021), the gene expression was quantified as the sum of intron and exon reads,

normalized as counts per million (CPM), and $\log_2$-transformed. Each cell was categorized into

one of the predefined clusters (M, P, K) based on the metadata associated with the published data

set (Bakken et al., 2021). These cells in their predefined labels are shown in Figure 1A,

illustrating the cell-to-cell genetic diversity in a dimensionally reduced UMAP (Uniform

Manifold Approximation and Projection) plot.

Figure 1. Analysis pipeline of LGN scRNA-seq data, sourced from the Allen Brain Map. **(A)**

Scatter plot visualization for each cell (n = 1309) in UMAP space, where the horizontal and

vertical axes represent the dimensionality-reduction UMAP coefficients. The magnocellular (M),

parvocellular (P), and koniocellular (K) cells are illustrated as red, blue, and green, respectively.

(B) Flow chart of the clustering and gene analysis. As labeled in the supplied metadata, sets of

the M, P, and K cells were individually preprocessed. Each group of cells went through

parameter optimization and then clustered using Seurat v5. Differential genes were collected and

annotated with each cell population.

Given the wide diversity within primate retinal ganglion cells (Dacey, 2004; Peng et al.,

2019), our increasing understanding of the complexity of the LGN (Sun et al., 2024) and the

visual cortex (Vries et al., 2020; Wei et al., 2022), we believe cellular subtypes within M, P, and

K are to be discovered and expect they can be distinguished using transcriptomics. Therefore,

each of the M ($n = 350$), P ($n = 625$), and K cell ($n = 334$) groups from the publicly sourced

dataset were reanalyzed using an adjusted Seurat analysis pipeline (Figure 1B). Notably, because

there are no universally accepted standards for the function parameter selection in the Seurat

analysis pipeline (Arbatsky et al., 2025) and the default parameters are inherently arbitrary (I.

Schneider et al., 2021), we employed a global optimization approach to optimize for cluster

separation quality (Silhouette score) to determine the optimum number of principal components

(PCs), number of features, and clustering resolution for each cell type (see Methods).

To confirm the robustness of the clustering pipeline, we first validated the workflow

using a ground-truth-labeled PBMC single-cell RNA-seq dataset. The clustering achieved a

90.7% accuracy match to the known labels, with a moderate silhouette score of 0.276

(Supplementary Figure 1). Stability testing through 100 iterations of downsampling and varying

the number of features and dimensions yielded a mean cluster identity match of 89.63% (SD =

1.23%), confirming that the pipeline reliably recovers biologically meaningful subpopulations

Our M, P, and K clustering results using the adjusted Seurat analysis pipeline will be

described below in threefold: (1) how many within-cluster subpopulations (i.e., clusters within

M, P, or K sets) are evident in the data along with the optimized parameters used; (2) what are

the significant genetic differences that define these subpopulations; and (3) what functions and

interpretations can we infer using genetic annotation from public resources. Much of the

interpretation from (3) will be in the Discussion section. Throughout the Results and Discussion,

we define significant genes as those with an adjusted *p*-value < 0.05, and biologically significant

genes as those with both an adjusted *p*-value < 0.05 and a fold-change > 2 (or log2FC > 1).

However, because the definition of biological significance is somewhat arbitrary (Dalman et al.,

2012), and informative genes can exhibit moderate fold changes of less than 2 (St. Laurent et al.,

2013), we will consider both significant and biologically significant genes, while explicitly

noting fold-change values throughout.

M cell clustering

Two subpopulations emerged from the set of M cells: M_1 ($n = 226$) and M_2 ($n = 124$).

The optimized parameters used were 4 PCs, 2800 top features, and a cluster resolution of 0.2

(Figure 2A). After GAM fitting (for smoothing and outlier removal), these parameters produced

a silhouette score of 0.231, indicating a low to moderate within-cluster cohesion and inter-cluster

separation between M_1 and M_2 cells. The two clusters using the optimized parameters can be

visualized in the UMAP plot of Figure 2B. Likewise, when only considering the significant

genes from these populations (n genes $= 185$), the UMAP plot in Figure 2C illustrates an ideal

separation between the two subpopulations with a silhouette score of 0.36, though we

acknowledge this is somewhat circular since gene selection was based on differential expression.

We also performed stability and sensitivity analyses to assess the robustness of the

clustering. Downsampling the number of cells (70-90%), varying the number of highly variable

genes (80-120%) and number of PCs (±1), consistently reproduced two M clusters in 71.5% of

bootstrap iterations ($n = 1000$), with an average cluster identity match of $86.0 \pm 6.0\%$. This

indicates that the optimized parameters are reasonably robust within typical parameter variations.

We also tested the clustering parameters under random noise conditions by introducing noise.

When adding Gaussian noise to the data (see Methods), no noise-iterated dataset ($n = 1,000$)

produced a silhouette score higher than the observed silhouette from the original clustering,

resulting in an empirical p-value of 0.008. This indicates that the cluster structure reflects

meaningful biological separation rather than random fluctuations. Notably, the silhouette score

exhibited only 6% degradation under this noise level, indicating cluster stability.

A total of 185 genes significantly differentiated between M_1 and M_2 cells (obtained from

Wilcoxon Rank Sum test; see Methods). The genes that contributed the most to the genetic

variance between M_1 and M_2 cells are illustrated in the heatmap of Figure 2D, showing the

expression levels of the most differential genes (top 10 genes from each population ranked by

ascending adjusted *p*-value) for each cell. Genes of particular interest from our ontology analysis

are shown in the dot and volcano plots (Figure 2E and 2F, respectively).

Generally, from our ontology, M_1 cells show upregulation of genes associated with action

potentials and signal transmission whereas M_2 cells express cytoskeletal/intracellular transport

proteins and genes involved in transcription and post-transcription regulation. Notably enriched

genes in M_1 cells include GRIK2 (ionotropic kainate-type glutamate receptor subunit) and

GRM1 (metabotropic glutamate receptor), alongside a diverse array of voltage-gated potassium

channel subunits (KCNC2, KCND2, KCNJ3, KCNH5) and the voltage-gated calcium channel

auxiliary subunit CACNA2D1. Conversely, M_2 cells express microtubule-associated proteins

such as MAP1A and MAPT, which stabilize dendritic and axonal microtubules respectively,

alongside cytoskeletal components SPTAN1, SPTBN2, and TTN. In addition, M_2 cells express

transcriptional regulators including KMT2B, a histone methyltransferase that modulates

chromatin architecture and gene expression; SRRT, an RNA-binding protein involved in miRNA

processing; and SNAPC4, a component of the snRNA-activating complex essential for snRNA

transcription and RNA processing. Finally, M_2 cells exhibit high levels of genes involved in

intracellular transport and cytoskeletal signaling. For example, KIF1A and KIFC2 encode

kinesin motor proteins that mediate anterograde and retrograde vesicular transport along

microtubules, while MAST1 encodes a microtubule-associated serine/threonine kinase

implicated in regulating synapse-associated cytoskeletal dynamics (Tripathy et al., 2018).

273   Overall, the mild degree of genetic differentiation is consistent with the modest silhouette

scores and the lack of sharply distinct clustering in UMAP space. This subtle separation could

suggest that M_1 and M_2 represent a gradient or continuum of transcriptional subtypes rather than

discrete populations. It is also noteworthy that the original work of Bakken et al. (2021) did not

identify subtypes within magnocellular neurons, suggesting that the M1/M2 split observed in the

current study likely arises from the increased sensitivity of our clustering optimization pipeline

(more in the Discussion), although only detecting modest transcriptomic divergence. In

summary, while transcriptomic evidence supports the existence of two subpopulations within M

cells, the limited biological separation indicates that these likely reflect subtle molecular

differences rather than robustly distinct cell types.

Figure 2. M cell clustering produced two clusters: M_1 (n = 226, red) and M_2 (n = 124, light-red).

(**A**) Top, elbow plot to qualitatively determine the number of PCs at the elbow (red dashed line).

Bottom, heat map of GAM-fitted silhouette scores from varying combinations of number of high

variable features and clustering resolution (bottom). The red circle in the heat map indicates the

maximum predicted silhouette score. The optimum number of PCs, high variable features, and

clustering resolution were used for downstream analysis. (**B**) Scatter plot visualization of M_1 and

M_2 clusters in UMAP space from the parameters determined in A. **(C)** The same UMAP plot but

only using the differentially expressed genes between M_1 and M_2 clusters (adjusted *p*-value <

0.05, *n* = 185). (**D**) Heatmap of RNA-seq expression for the top 10 differentially expressed genes

(ordered by increasing adjusted *p*-value; positive fold change only) for each cluster. The

horizontal and vertical axes represent individual cells and differentially expressed genes,

respectively. The colored bars on the top and left of the plot represent the cells and the top 10

differentially expressed for each cluster, respectively. The legend on the right describes the

expression levels for each gene and cell. (**E**) Gene expression dot plot showing the expression of

notable M_1 and M_2 significant markers (adjusted *p*-value < 0.05). Dot diameter indicates the

proportion of cells expressing the gene, and color intensity indicates average expression levels.

(F) Volcano plot of differential gene expression between M_1 and M_2 clusters. The x-axis

represents the fold change in expression, and the y-axis represents the –log10 adjusted p-value.

Genes passing significance threshold (adjusted p-value < 0.05) are represented as red data points.

The genes featured in E are labelled for reference.

P cell clustering

Similarly, two subpopulations were identified within P cells: P_1 (n = 334) and P_2 (n =

291). The optimized parameters were 4 PCs, 3200 top features, and a clustering resolution of

0.26 (Figure 3A) with a 0.225 silhouette score for this clustering, indicating a low to moderate

separation as visualized in the UMAP plot (Figure 3B). Restricting to the set of significant genes

(n = 692) improves the clustering to a moderate separation with a silhouette score of 0.329

(Figure 3C). Cluster sensitivity testing yielded a moderate cluster reproduction with 55% of

iterations matching two P clusters with an average cluster identity match of $82.3 \pm 9.4\%$.

Clustering stability testing, adding gaussian noise to the data, resulted in a silhouette score

degradation by only 5.88% with no noisy data iteration with a higher silhouette score than the

original (empirical *p*-value of 0.000), indicating that the clustering reflects meaningful biological

signal rather than noise.

318   A total of 692 genes were significantly differentially expressed between P_1 and P_2

319   (adjusted $p < 0.05$), with 625 genes enriched in P_1 and 67 in P_2. The most significant genetic

320   differences contributing to the P_1 and P_2 separation are illustrated in the heatmap of Figure 3D,

321   and genes of particular interest in our ontology are shown in the plots of Figure 3E and 3F.

322   From our ontology, P_1 cells express multiple components of the mitochondrial electron

transport chain complex I, including NDUFA11, NDUFV1, and NDUFS8, as well as glycolytic

enzymes such as TPI1 (triosephosphate isomerase) and GAPDH (glyceraldehyde-3-phosphate

dehydrogenase). The mitochondrial GTP-binding protein GTPBP6, implicated in proper

mitochondrial translation and oxidative phosphorylation (Lavdovskaia et al., 2020) and STK11, a

master regulator of cellular metabolism and energy homeostasis (Huang & Li, 2022), are also

highly expressed. In parallel, P_1 cells express a number of genes that afford protection against

oxidative and metabolic stress. These include HSPB1, HSP90AB1, and HSPA12A – heat shock

proteins known to stabilize cytoskeletal and synaptic proteins under stress (Dávila et al., 2014;

Toth et al., 2010; Wang et al., 2023); NUDT1, which prevents mutagenesis by hydrolyzing

oxidized nucleotides; BOK, a BCL-2 family protein involved in mitochondrial dynamics and

endoplasmic reticulum stress responses independent of apoptosis (D'Orsi et al., 2016); and

MAP4K2, an upstream activator of the JNK pathway, which is a 'cell death' pathway involved

in cellular stress signaling (Weston & Davis, 2007). Finally, P_1 cells express a number of ion

channels, including SCN2B (voltage-gated sodium channel subunit), KCNT1, KCNAB2,

KCNC1, KCNJ12, KCNK3, KCNQ1, KCNQ2 (potassium ion channel subunits), CACNA1G,

CACNA1A, CNCBA1B, CACNA1C (voltage-gated calcium channel subunits), and GRIN1

(glutamate NMDA receptor subunit).

By contrast, P_2 cells express genes such as NRXN1 (neurexin 1) and NLGN1

(neuroligin 1) which, as a pair, function as cell adhesion molecules which bind with each other to

establish excitatory/inhibitory synapse specificity (Arias-Aragón et al., 2025; Gomez et al.,

2021). NLGN1 specifically promotes excitatory glutamatergic synapse formation (Szíber et al.,

2024; Zeidan & Ziv, 2012). NRG3 (neuroregulin) is likewise involved in the formation of

excitatory synapses (Müller et al., 2018). CDH2, PCDH7, PCDH9 are all members of the

cadherin superfamily, playing roles in axon pathfinding, synaptic targeting, and circuit

refinement.

Overall, P_1 cells show high expression of genes associated with cellular metabolism and

energy production, especially aerobic metabolism, as well as a number of genes responsible for

protecting the cell from oxidative and metabolic stressors. P_2 cells, by contrast, express genes

important for synaptic regulation and specificity, cell adhesion, and calcium signaling –

suggesting an emphasis on connectivity and plasticity.

Figure 3. P cell clustering produced two clusters: P_1 (n = 334; dark blue) and P_2 (n = 291; light

blue). **(A)** Top, elbow plot to qualitatively determine the number of PCs at the elbow (red dashed

line). Bottom, heat map of GAM-fitted silhouette scores from varying combinations of number

of highly variable features and clustering resolution. The redcircle in the heat map marks the

optimal parameters used in downstream analysis. **(B)** Scatter plot visualization of P_1 and P_2

clusters in UMAP space using the parameters from A. **(C)** The same UMAP plot restricted to

differentially expressed genes between P_1 and P_2 (adjusted *p*-value < 0.05, *n* = 692). **(D)**

Heatmap of RNA-seq expression for the top 10 differentially expressed genes (ordered by

increasing adjusted *p*-value; positive fold change only) for each cluster. Horizontal and vertical

axes represent individual cells and genes, respectively. Colored bars denote cluster identity. **(E)**

Gene expression dot plot showing expression of notable significant markers in P_1 and P_2. Dot

diameter represents the proportion of cells expressing the gene; color intensity reflects average

expression. **(F)** Volcano plot showing fold change versus –log10 adjusted *p*-value for each gene.

Genes meeting both statistical and biological significance are highlighted in red, with markers

from **(E)** labeled.

K cell clustering

The K cell population exhibited a stronger degree of transcriptional heterogeneity in contrast to P

and M cells. Three subpopulations were identified: K_1 (n = 151), K_2 (n = 120), and K_3 (n = 63).

The optimized parameters we obtained for K clusters were 4 PCs, 1100 top features, and a

clustering resolution of 0.12 (Figure 4A), achieving a stronger silhouette score of 0.401. Using

only the significant genes (n = 642) yielded only a slightly higher silhouette score of 0.42 (Figure

4C). The K clustering also produced high stability with an average cluster reproduction match of

$92.1 \pm 4.5\%$ (*n* = 1000; 14.2% skipped due to cluster mismatch). Noise testing produced an

empirical *p*-value of 0.001, with a silhouette degradation of 6.96%, further confirming the

robustness of the clustering.

A total of 642 genes were significantly differentially expressed across the K clusters

(adjusted p < 0.05), with 66 enriched in K_1, 186 in K2, and 390 in K3. Of these, 361 genes met

biological significance criteria, including 16 in K_1, 76 in K2, and 269 in K3 (Figures 4D). This

distribution highlights the substantial heterogeneity within the K population, particularly the

strong separation of K_3. Genes of particular interest in our ontology are shown in the plots of

Figure 4E and 4F, and described below.

All three K cell subpopulations (K_1, K_2, K_3) express ion channels and glutamate receptors

needed for signal transmission and action potential generation, though with subtle functional

differences. In addition, coexpression of other genes (such as Ca^{2+}-ATPases in K_1, cytoskeletal

binding elements in K_2, and non-glutamate receptors in K_3) highlights distinct roles for each

subpopulation.

Key genes expressed by K_1 cells include ion channel subunits such as KCNC2, KCNN3,

KCNB2, and CACNA1I, as well as ATP1B2, a subunit of the Na^+/K^+-ATPase critical for

maintaining resting membrane potential. Notably, K_1 cells selectively express metabotropic

glutamate receptor subunits GRM5 and GRM7, G protein-coupled receptors that mediate slower,

modulatory synaptic responses via second messenger cascades. K_1 cells also express ATP2A3

and ATP2B1, which encode plasma membrane and endoplasmic reticulum Ca^{2+}-ATPases

involved in calcium extrusion and intracellular calcium homeostasis.

K_2 cells also express a number of proteins important in signal transmission,

including KCNH1, KCNJ3, KCNAB3, SCN1B, SCN9A (sodium and potassium ion channel

subunits) and GRIK4, GRIA4 (ionotropic kainate and AMPA glutamate receptors). However, in

addition to those proteins important for signal transmission are also a substantial number of

proteins involved in cytoskeletal binding, including TTN, TNS1, FMN1, MTUS1, SPTB,

MYLK, MYO5B, MYO16, MYO19, MYO14.

The K_3 population of cells expresses a large number of monoatomic ion channels,

channels important for electrical signaling in neurons. These include KCNIP1, KCNA3,

KCNQ1, KCNQ5, KCNC4, KCNJ6, KCNK2, KCNK3, KCNH3, KCNT2, KCNK10, KCNMA1

(subunits of different potassium channels, including voltage-gated, sodium-activated, two-pore

domain, and inwardly rectifying types), CACNA1E, CACNA1B, and CACNG4 (voltage-gated

calcium channel subunits). In addition to these ion channels, this cell population expresses a

substantial amount of neurotransmitter receptors for both glutamate – the primary excitatory

neurotransmitter in LGN – as well as other modulatory neurotransmitters. These include GRID1,

GRID2, GRIN3A, GRM2, GRIK5 (glutamate receptors; NMDA, delta, and kainate type), HRH1

(histamine H1 receptor), OPRK1 (opioid receptor kappa), MGLL (endocannabinoid related

lipase), TRPM3 (neurosteroid sensitive transient receptor potential channel; Webster et al.,

2020), CHRNA7 (nicotinic acetylcholine receptor subunit), and P2RX4 (purinergic receptor).

In the current analysis, the Kp and Kap cells from Bakken et al. (2021) were analyzed

and clustered together due to the relatively small number of Kap cells. Despite this, the

clustering from the current analysis consistently grouped Kap cells into K_3, while subdividing the

remaining Kp cells into K_1 and K_2. A supplementary figure (Supplementary Figure 2) confirms

that K_3 aligns with the Kap identity (90%, n = 66/73), indicating that the Kap population is

preserved and identifiable within the revised clustering framework.

Overall, the strong degree of genetic differentiation within the K population is reflected

in the ontology, higher silhouette scores, and clear separation in UMAP space. The dominant

transcriptomic signature of K_3, along with the high number of biologically significant genes,

suggests that K_1, K_2, and K_3 represent robust and distinct molecular subtypes. This result refines

the earlier findings of Bakken et al. (2021), which identified Kp and Kap subtypes, by further

subdividing Kp into K_1 and K_2 while preserving Kap as K_3. The transcriptomic evidence

supports the existence of three distinct subpopulations within K cells, indicating a greater degree

of heterogeneity in the koniocellular pathway compared to the magnocellular and parvocellular

pathways.

Figure 4. K cell clustering produced four clusters: K_1 ($n = 151$; dark green), K_2 ($n = 120$; light green), and K_3 ($n = 63$; white). **(A)** Top, elbow plot used to determine the number of PCs at the elbow (red dashed line). Bottom, heat map of GAM-fitted silhouette scores across combinations of feature number and clustering resolution. The red circle indicates the optimum parameter set.

(B) UMAP visualization of K_1, K_2, and K_3 clusters using the parameters in A. **(C)** UMAP using

only differentially expressed genes between clusters (adjusted *p*-value < 0.05, *n* = 642). **(D)**

Heatmap of RNA-seq expression for the top 10 differentially expressed genes (ordered by

increasing adjusted *p*-value; positive fold change only) per cluster. Axes indicate individual cells

and gene markers; color bars denote cluster identity. **(E)** Gene expression dot plot showing

notable markers in each K cluster. Dot size corresponds to expression frequency; color intensity

to average expression level. **(F)** Volcano plot of differential gene expression across clusters. X-

axis denotes fold change; Y-axis, –log10 adjusted *p*-value. Genes from **(E)** are labeled.

Interspecies and donor variability

To assess whether the observed subpopulations were confounded by interspecies differences or

individual subject variability, we performed a qualitative evaluation of UMAP plots for the M, P,

and K populations colored by individual animal (*n* = 3) and species (*M. fascicularis* and *M.*

nemestrina) (Supplementary Figure 3). As shown in these plots, the major subpopulations are

preserved across both subject and species. To quantify this observation, we performed Fisher's

exact tests to evaluate species and subject enrichment within each cluster. No species enrichment

was observed in any cluster after Bonferroni correction (*p* > 0.05), indicating that interspecies

differences did not drive the clustering. Similarly, no donor enrichment was detected in any of

the clusters, except for one koniocellular cluster, K3. K3 exhibited significant donor enrichment

(*p* = 0.00023; *p_adj* = 0.0033), likely reflecting the relatively small size of the K3 population.

Taken together, these results suggest that the M, P, and K subpopulations are not driven by

species- or subject-related differences, and reflect underlying biological distinctions.

Subtype	Key Gene Expression	Functional Implications
M_1	- Ionotropic (GRIK2) & metabotropic (GRM1) glutamate receptors - Voltage-gated potassium channels (KCNC2, KCND2, KCNJ3, KCNH5) - Voltage-gated calcium channel subunit (CACNA2D1)	- Optimized for rapid, high-fidelity signal transmission - Fast, phasic relay neurons - Fast excitability and spike timing regulation
M_2	- Cytoskeletal proteins (MAP1A, MAPT, SPTAN1, SPTBN2, TTN) - Transcriptional regulators (KMT2B, SRRT, SNAPC4) - Kinesin motor proteins (KIF1A, KIFC2) - Microtubule-associated kinase (MAST1)	- Structurally robust, adapted for synaptic plasticity - Modulatory role integrating feedforward and feedback inputs - Supports synaptic remodeling and network activity

P₁	- Mitochondrial ETC components (NDUFA11, NDUFV1, NDUFS8) - Glycolytic enzymes (TPI1, GAPDH) - Metabolic regulators (GTPBP6, STK11) - Heat shock proteins (HSPB1, HSP90AB1, HSPA12A) - Ion channels (SCN2B, KCNT1, KCNAB2, etc.) - NMDA receptor subunit (GRIN1)	- High metabolic activity and oxidative stress protection - Tonic firing, high-fidelity relay neurons - Sustained firing patterns suitable for detailed vision
P₂	- Synaptic adhesion molecules (NRXN1, NLGN1) - Cadherins (CDH2, PCDH7, PCDH9) - Calcium signaling genes (NRG3)	- Emphasis on synaptic specificity, plasticity, and connectivity - Modulatory role in circuit refinement and activity-dependent changes
K₁	- Ion channels (KCNC2, KCNN3, KCNB2) - Na$^+$/K$^+$-ATPase subunit - Metabotropic glutamate receptors (GRM5, GRM7) - Calcium ATPases (ATP2A3, ATP2B1)	- Slower synaptic response kinetics via metabotropic receptors - High-throughput synaptic activity with calcium regulation

K$_2$	- Signal transmission proteins (KCNH1, KCNJ3, SCN1B, GRIK4, GRIA4) - Cytoskeletal proteins (TTN, TNS1, FMN1, MTUS1, SPTB, MYLK, MYO family)	- Structurally robust with fast conduction properties - Possible projections to extrastriate areas for reflexive/feedforward processing
K$_3$	- Numerous potassium channel subunits (KCNIP1, KCNA3, etc.) - Glutamate receptors (GRID1, GRID2, GRIN3A, GRM2, GRIK5) - Non-glutamate receptors (HRH1, OPRK1, MGLL, TRPM3, CHRNA7, P2RX4) - PENK (proenkephalin)	- Highly electrically active, involved in visual modulation by multiple neurotransmitter systems - Respond to arousal, attention, pain, metabolic, and hormonal states

460   **Table 1.** Summary of ontology analysis.

Discussion

Our transcriptomic analysis revealed seven subpopulations of LGN neurons: two populations of magnocellular neurons (M_1 and M_2), two populations of parvocellular neurons (P_1 and P_2), and three populations of koniocellular neurons (K_1, K_2, and K_3). Although the same dataset was used as Bakken et al. (2021), our study employed several different analytical methods. Notably, we utilized analytic Pearson residuals for normalization (*SCTransform*, Lause *et al.*, 2021), Harmony integration for batch correction (*HarmonyIntegration*; Korsunsky *et al.*, 2019), and a silhouette-based global parameter optimization pipeline, all of which likely enabled the identification of the nuanced subpopulations in the LGN macaque data.

Examining the gene expression of M, P, and K subtypes elucidates some interesting themes (see Table 1 for summary). As for M and P cells, there appears to be two distinct populations of neurons: first, a population of neurons with genes optimizing for high-throughput signal transmission. In the high-throughput populations of M and P cells, distinct gene expression patterns suggest differences in signaling kinetics, matching the electrophysiologic response profiles observed for each population (Kaplan & Shapley, 1982; Lee & Sun, 2009; Maunsell et al., 1999). The second population of M and P cells shows a striking absence of proteins essential for signal transmission, including ion channels, glutamate receptors, and metabolic genes. Instead, they seem to serve a more modulatory role, expressing genes involved in synaptic plasticity, connectivity, and cytoskeletal structure.

K cells do not exhibit the clear high-throughput vs. modulatory division seen in M and P cells. This aligns with numerous studies describing K cells as a more heterogeneous, biochemically unique, and functionally distinct part of the visual system (Hendry & Reid, 2000;

Xu et al., 2001). All K neuron populations express proteins important for signal transmission,

however the specific signal transmission proteins upregulated in each population gives insight

into the response kinetics of each population and highlights the electrophysiologically observed

heterogeneity in response kinetics of K cells (Eiber et al., 2018; White et al., 2001; Xu et al.,

2001).

Of note, these functional differences of M, P, and K cell populations discussed below

come from gene ontology analyses based on a selected subset of expressed genes. While these

inferences provide valuable insights, the absence of electrophysiological, morphological, and

spatial data limits the ability to comprehensively characterize each cell type.

Magnocellular

The gene expression profile of M_1 cells suggests that they are specialized for rapid, high-fidelity

signal transmission, consistent with a role as fast, excitable relay neurons adapted to process

high-throughput glutamatergic input. The co-expression of both ionotropic (GRIK2) and

metabotropic (GRM1) glutamate receptors indicates that these cells are highly responsive to

glutamatergic input, likely from retinothalamic projections, through both fast inotropic and

slower metabotropic modulatory signaling mechanisms. Kainate receptors such as those encoded

by GRIK2 activate within <1 ms at high glutamate concentrations, generating rapid excitatory

currents, and rapidly deactivate or desensitize as glutamate levels decline or remain elevated,

enabling precise temporal transmission of retinal signals (Lerma et al., 1997). In contrast,

metabotropic glutamate receptors encoded by GRM1 are G protein-coupled receptors that

modulate downstream signaling pathways, potentially allowing for delayed dynamic tuning of

neuronal responsiveness over time (Sugiyama et al., 1987).

High expression of voltage-gated potassium and calcium channel subunits further

supports a role in feedforward, high-throughput synaptic transmission. The expression of

multiple potassium channel subtypes suggests finely tuned regulation of excitability and spike

timing (reviewed in Jan & Jan, 2012). For instance, KCNC2 encodes a high-threshold, fast-

activating potassium channel that enables brief, high-frequency action potentials (Yan et al.,

2005), while KCND2 contributes to A-type currents that regulate subthreshold excitability,

repetitive firing frequency, and back-propagation of action potentials (Connor & Stevens, 1971;

Shibata et al., 2000; Zhang & McBain, 1995). CACNA2D1, an auxiliary subunit of voltage-

gated calcium channels, enhances calcium channel trafficking and function, supporting efficient

neurotransmitter release and dendritic integration. Together, these expression patterns reinforce

the hypothesis that M_1 cells function as fast-acting, phasic relay neurons within the LGN,

aligning with the classical view of magnocellular pathways as mediators of high-speed visual

processing (Kaplan & Shapley, 1982; Maunsell et al., 1999).

M_2 cells, in contrast to M_1 cells, appear structurally robust and preferentially adapted for

synaptic plasticity rather than high-throughput signal relay. Their gene expression profile

suggests a potential role as modulatory hubs with context-dependent outputs, potentially

mediated through dynamic regulation of receptor composition, axonal targeting, or synaptic

scaffolding. Notably, M_2 cells show enriched expression of genes involved in cytoskeletal

architecture, intracellular transport, and transcriptional/post-transcriptional regulation. These

expression patterns suggest ongoing transcriptional remodeling, possibly in response to activity-

dependent plasticity or extra-thalamic modulation.

Principally, these findings support that M_2 cells are structurally robust. The concerted

expression of genes related to cytoskeletal organization, vesicular transport, and transcriptional

control – particularly in the relative absence of classical excitability markers such as ionotropic

receptors and voltage-gated ion channels – points to two possible, non-mutually exclusive

hypotheses regarding M_2 functionality: first, it's possible that the gene expression profile of M_2

cells indicates different action potential kinetics from M_1 cells. While M_1 cells are highly

excitable, phasic, and fast, M_2 cells may be more structurally robust, but exhibit slower, more

sustained kinetics. One source of evidence supporting this hypothesis comes from many

laboratories who have demonstrated that, while P cells demonstrate homogeneously sustained

kinetics, M cells contains both transient and sustained neurons (Blakemore & Vital-Durand,

1986; Kaplan & Shapley, 1982; Levitt et al., 2001; Xu et al., 2001). As a second hypothesis, it's

possible that these cells have a specialized role in coordinating synaptic plasticity, integrating

feedback and feedforward input, and regulating broader network activity across thalamocortical

circuits. M_2 cells may not be built structurally strong because of size, but instead due to a need to

coordinate traffic, form many synaptic connections, and modulate activity broadly. Given the co-

expression of many transcriptional regulatory proteins, these cells may serve a more integrative

role within the LGN, optimized for modulatory functions rather than rapid signal relay.

Overall, M cells appear to have two functionally distinct populations. M_1 cells are the

prototypically-understood definition of magnocellular neurons. They appear to be the work-

horses of electrical transmission, but simple relay neurons, transmitting information from retina

to cortex with minimal modulation or refinement. Their presumed response kinetics are the

classically understood fast response kinetics and high temporal frequencies associated with M

cells. M_2 cells seem to be more modulatory players. With an absence of proteins important for

excitability, action potentials, or synaptic transmission, these are structurally robust neurons,

possibly with extensive networks of dendrites or axons. Their high expression of proteins

involved in transcriptional and post-transcriptional suggests that they have a role in activity- or

state-dependent modulation of neural activity. As opposed to simple relay neurons, M_2 neurons

may be responsible for visual processing of retinofugal input to LGN.

Parvocellular

P_1 cells exhibit a gene expression profile marked by high metabolic activity and enhanced

resilience to oxidative and mitochondrial stress, suggesting specialization for sustained, energy-

demanding function. Among the most biologically significant markers are numerous enzymes

involved in core metabolic pathways and cellular stress regulation. Notably, P_1 cells express

multiple components of the mitochondrial electron transport chain complex I, as well as

glycolytic enzymes. Mitochondrial GTP-binding protein GTPBP6, implicated in proper

mitochondrial translation and oxidative phosphorylation (Lavdovskaia et al., 2020) and STK11, a

master regulator of cellular metabolism and energy homeostasis (Huang & Li, 2022) support the

interpretation that P_1 neurons have high, sustained energy demands, relying heavily on aerobic

respiration to maintain function under prolonged activity. Finally, P_1 cells express a number of

genes, such as heat shock proteins, NUDT1, BOK, and MAP4K2, that confer protection against

oxidative and metabolic stress.

To further support the hypothesis that P_1 cells function as a high-throughput population of relay

neurons, these cells exhibit robust expression of genes involved in action potential generation

and synaptic transmission, paralleling the molecular profile of M_1 cells. These include a voltage-

gated sodium channel subunit, potassium ion channel subunits, voltage-gated calcium channel

subunits, and glutamate NMDA receptor subunit.

This gene expression profile supports the hypothesis that P_1 cells are metabolically robust relay

neurons. Like M_1 neurons, P_1 neurons may serve high-throughput relay functions, however the

distinctive upregulation of genes associated with oxidative metabolism, mitochondrial integrity,

and stress resilience in P_1 cells suggests a tonic firing profile rather than the fast, phasic signaling

characteristic of M_1 cells. This aligns with classical physiological descriptions of parvocellular

neurons as exhibiting sustained firing patterns in support of detailed, high-fidelity, continuous

visual processing (Lee & Sun, 2009; Maunsell et al., 1999).

While P_1 cells express proteins suggestive of high-throughput signal transmission, P_2 cells

demonstrate gene expression important for synapse formation and specificity, cell adhesion, and

calcium signaling, with a notable absence of proteins related to signal transmission – such as ion

channels or neurotransmitter receptors. These features suggest a highly plastic and well-

connected population of neurons which, like M_2 neurons, may function as a connectional hub or

integrator, rather than a high-throughput information relay. Overall, genes such as NRXN1 and

NLGN1, as well as multiple members of the cadherin superfamily, suggest that P_2 cells play a

role in precision wiring and synaptic specificity, forming highly targeted connections. The

expression of activity-regulated structural proteins (e.g., neurexins and neuroligins) may indicate

a role in activity-dependent circuit adaptability (Jüngling et al., 2006).

Like M cells, P cells also have two functionally distinct populations. P_1 cells (similar to M_1 cells)

are the high-throughput signal transmitters with gene expression suggestive of substantial

metabolic needs, as well as protection from metabolic stress. These features suggest cells which

fire tonically, specific for high-fidelity, high-detail vision. P_2 cells, like M_2 cells, appear to play a

more modulatory, information-integration role. They seem to be well connected, with proteins

specific for synaptic specificity and involved in activity-dependent structural modulation and

regulation.

Koniocellular

Similar to M_1 and P_1 cell populations, K_1 cells exhibit enriched expression of genes

associated with action potential generation and synaptic transmission. However, the specific

repertoire of signal transmission proteins in K_1 cells offers insight into their distinct response

kinetics and functional specialization relative to M and P neurons. The absence of ionotropic

glutamate receptors and the exclusive expression of metabotropic receptors suggest that K_1 cells

may exhibit slower synaptic response kinetics, characteristic of the longer response latencies

associated with K cells (Pietersen et al., 2014; Xu et al., 2001). These studies also noted highly

heterogeneous response kinetics associated with K cells, which is demonstrated by the diversity

of different potassium, calcium, sodium, and glutamate receptor types expressed in K_1, K_2, and

K_3 cells. The expression of ATP2A3 and ATP2B1, calcium clearance pumps, supports the idea

that K_1 cells are equipped to handle high-throughput synaptic activity while minimizing calcium-

induced cytotoxicity.

In K_2 cells, the presence of cytoskeletal binding proteins suggests that these neurons, like

M_2 neurons, are structurally robust. With the coexpression of fast sodium and potassium channel

subunits and ionotropic glutamate receptors, one hypothesis is that the upregulation of structural

proteins reflects thicker axons with faster conduction velocities, perhaps with projections to

extrastriate areas. This implies that K_2 cells could have a role in reflexive behaviors or

feedforward signalling, projecting to areas requiring fast, high-bandwidth processing, and could

possibly be the basis for observed projections from K layers to MT (Casagrande, 1994; Sincich

et al., 2004).

The K_3 population of cells expresses many proteins important for electrical signaling in

neurons as well as a diverse population of glutamate and non-glutamate neurotransmitter

receptors. The increased expression of these genes suggests that this population of cells is highly

electrically active and may be the site of visual modulation by non-glutamate neurotransmitter

systems within LGN.

Like K_1 and K_2 cells, the expression of ion channels and glutamate receptors indicate an

active role in signal transmission. The diversity of glutamate receptor types (NMDA, kainate,

and delta type ionotropic, as well as metabotropic) and ion channels expressed in K_3 continue to

support the electrophysiological findings that K cells overall demonstrate highly heterogeneous

response characteristics. The increased expression of potassium ion channels relative to other K

subtypes implies that K_3 cells may demonstrate high temporal frequencies similar to M cells – a

point to which we will return. However, what is unique about K_3 cells is the vast expression of

non-glutamate neurotransmitter receptors. These include receptors specific for histamine,

opioids, acetylcholine, extracellular ATP, and lipids/steroids. The expression of these receptors

suggest that K_3 cells are under tight top-down and subcortical control; they respond to arousal,

attention, alertness, pain, and inflammation, dynamically shifting their output depending on

internal brain states (Javadi et al., 2015; Jin et al., 2002; Noseda et al., 2017; Yang et al., 2015).

In addition, these neurons may adapt their function in response to steroid hormones, metabolic

shifts, or stress, potentially linking vision or sensory processing with hormonal states.

The final remark regarding K_3 cells is the expression of PENK (proenkephalin). Bakken

et al. (2021) previously performed single-cell RNA sequencing on macaque LGN cells and

showed two populations of K cells, Kp and Kap. Kp cells selectively expressed PENK and were

enriched in ventral K layers K1 and K2, adjacent to M cells. This association suggests that the K_3

cells we identified were the same Kp cells identified by Bakken et al., or a subpopulation thereof,

and may help to explain the commonality in response characteristics between M and K_3 cells. In

fact, research in the K1 and K2 layers (where these Kp/K_3 cells are enriched) has shown that

these cells demonstrate large receptive fields with sensitivity to contrast and high temporal

frequencies – very similar to M cell receptive fields (Solomon et al., 2002; White et al., 2001; Xu

et al., 2001).

Conclusion

Overall, the transcriptomic evidence supports a refined view of LGN organization, extending

beyond the classical M, P, and K classification. The presence of modest subtypes in M and P

pathways and strong molecular heterogeneity in K cells suggests that LGN neurons may play a

more diverse and functionally specialized role in visual processing than traditionally held. This

heterogeneity could reflect differential involvement in non-visual modulation, temporal

dynamics, or attentional processes.

Future studies that combine transcriptomics with electrophysiological or anatomical data,

such as spatial transcriptomics or patch-seq, will be necessary to validate these subpopulations

and clarify their roles. Furthermore, matched functional-transcriptomic datasets would be

essential to determine whether any of the transcriptomic clusters observed here correspond to the

functionally distinct cells described in recent electrophysiology experiments, to draw further

conclusions about the physical organization of the LGN and its rich variety of cell types.

Supplementary Materials

Supplementary Figure 1. Testing the clustering analysis pipeline on ground truth PBMC data.

(A) UMAPs of clustering using the default Seurat parameters. This produced 5 clusters,

including one heterogeneous cluster (bottom-left), when compared to the true labels (bottom

UMAP). **(B)** The heterogeneous cluster in A was reclustered using the same clustering pipeline

as described in this study (top UMAP). Bottom, the same UMAP plot but labeled using the

growth truth labels. **(C)** Confusion matrix showing match (%) between clusters (y-axis) and true

label (x-axis). Overall, the clustering pipeline produced a 90.73% match with the ground truth

data.

Supplementary Figure 2. K clustering from the current study **(A)** and from the Bakken et al.

(2021) study **(B)**. K3 in the current study is analogous to Kp in the Bakken study with an 86.3%

match ($n = 63/73$).

Supplementary Figure 3. The UMAPs of M (left), P (middle), and K clusters (right) with the

data labelled by animal id (**A**, n = 3) and species (**B**, n = 2).

Data and Code Availability

The R scripts used for data preprocessing, clustering, and analysis in this study are available

online at https://github.com/shihaisun-scott/sun_LGN_macaque_transcriptomics.

References

Allen Institute for Brain Science. (2022). *Single-cell RNAseq of LGN of mouse, human, and nonhuman primate [dataset]* [Dataset]. knowledge.brain-map.org/data/MR9M3L8BU9PCOH3BHCV/summary

Arbatsky, M., Vasilyeva, E., Sysoeva, V., Semina, E., Saveliev, V., & Rubina, K. (2025). Seurat function argument values in scRNA-seq data analysis: Potential pitfalls and refinements for biological interpretation. *Frontiers in Bioinformatics*, *5*, 1519468. https://doi.org/10.3389/fbinf.2025.1519468

Arias-Aragón, F., Robles-Lanuza, E., Sánchez-Gómez, Á., Martinez-Mir, A., & Scholl, F. G. (2025). Analysis of neurexin-neuroligin complexes supports an isoform-specific role for beta-neurexin-1 dysfunction in a mouse model of autism. *Molecular Brain*, *18*(1), 20. https://doi.org/10.1186/s13041-025-01183-0

Bakken, T. E., Hodge, R. D., Miller, J. A., Yao, Z., Nguyen, T. N., Aevermann, B., Barkan, E., Bertagnolli, D., Casper, T., Dee, N., Garren, E., Goldy, J., Graybuck, L. T., Kroll, M., Lasken, R. S., Lathia, K., Parry, S., Rimorin, C., Scheuermann, R. H., … Tasic, B. (2018). Single-nucleus and single-cell transcriptomes compared in matched cortical cell types. *PLOS ONE*, *13*(12), e0209648. https://doi.org/10.1371/journal.pone.0209648

Bakken, T. E., van Velthoven, C. T., Menon, V., Hodge, R. D., Yao, Z., Nguyen, T. N., Graybuck, L. T., Horwitz, G. D., Bertagnolli, D., Goldy, J., Yanny, A. M., Garren, E., Parry, S., Casper, T., Shehata, S. I., Barkan, E. R., Szafer, A., Levi, B. P., Dee, N., … Tasic, B. (2021). Single-cell and single-nucleus RNA-seq uncovers shared and distinct

axes of variation in dorsal LGN neurons in mice, non-human primates, and humans.

eLife, *10*, e64875. https://doi.org/10.7554/eLife.64875

Blakemore, C., & Vital-Durand, F. (1986). Organization and post-natal development of the

monkey's lateral geniculate nucleus. *The Journal of Physiology*, *380*, 453–491.

https://doi.org/10.1113/jphysiol.1986.sp016297

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., & Satija, R. (2018). Integrating single-cell

transcriptomic data across different conditions, technologies, and species. *Nature*

Biotechnology, *36*(5), 411–420. https://doi.org/10.1038/nbt.4096

Casagrande, V. A. (1994). A third parallel visual pathway to primate area V1. *Trends in*

Neurosciences, *17*(7), 305–310. https://doi.org/10.1016/0166-2236(94)90065-5

Chen, W., Zhu, X. H., Thulborn, K. R., & Ugurbil, K. (1999). Retinotopic mapping of lateral

geniculate nucleus in humans using functional magnetic resonance imaging. *Proceedings*

of the National Academy of Sciences of the United States of America, *96*(5), 2430–2434.

https://doi.org/10.1073/pnas.96.5.2430

Cheong, S. K., Tailby, C., Solomon, S. G., & Martin, P. R. (2013). Cortical-Like Receptive

Fields in the Lateral Geniculate Nucleus of Marmoset Monkeys. *Journal of*

Neuroscience, *33*(16), 6864–6876. https://doi.org/10.1523/JNEUROSCI.5208-12.2013

Choudhary, S., & Satija, R. (2022). Comparison and evaluation of statistical error models for

scRNA-seq. *Genome Biology*, *23*(1), 27. https://doi.org/10.1186/s13059-021-02584-9

Clarke, Z. A., Andrews, T. S., Atif, J., Pouyabahar, D., Innes, B. T., MacParland, S. A., & Bader,

G. D. (2021). Tutorial: Guidelines for annotating single-cell transcriptomic maps using

automated and manual methods. *Nature Protocols*, *16*(6), 2749–2764.

https://doi.org/10.1038/s41596-021-00534-0

Connor, J. A., & Stevens, C. F. (1971). Voltage clamp studies of a transient outward membrane

current in gastropod neural somata. *The Journal of Physiology*, *213*(1), 21–30.

https://doi.org/10.1113/jphysiol.1971.sp009365

Dacey, D. (2004). Origins of perception: Retinal Ganglion Cell Diversity and the Creation of

Parallel Visual Pathways. In *Sensory Systems* (pp. 281–301).

Dávila, D., Jiménez-Mateos, E. M., Mooney, C. M., Velasco, G., Henshall, D. C., & Prehn, J. H.

M. (2014). Hsp27 binding to the 3'UTR of bim mRNA prevents neuronal death during

oxidative stress-induced injury: A novel cytoprotective mechanism. *Molecular Biology of*

the Cell, *25*(21), 3413–3423. https://doi.org/10.1091/mbc.E13-08-0495

Denison, R. N., Vu, A. T., Yacoub, E., Feinberg, D. A., & Silver, M. A. (2014). Functional

mapping of the magnocellular and parvocellular subdivisions of human LGN.

NeuroImage, *102*, 358–369. https://doi.org/10.1016/j.neuroimage.2014.07.019

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,

M., & Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner.

Bioinformatics, *29*(1), 15–21. https://doi.org/10.1093/bioinformatics/bts635

D'Orsi, B., Engel, T., Pfeiffer, S., Nandi, S., Kaufmann, T., Henshall, D. C., & Prehn, J. H. M.

(2016). Bok Is Not Pro-Apoptotic But Suppresses Poly ADP-Ribose Polymerase-

Dependent Cell Death Pathways and Protects against Excitotoxic and Seizure-Induced

Neuronal Injury. *The Journal of Neuroscience: The Official Journal of the Society for*

Neuroscience, *36*(16), 4564–4578. https://doi.org/10.1523/JNEUROSCI.3780-15.2016

Eiber, C. D., Rahman, A. S., Pietersen, A. N. J., Zeater, N., Dreher, B., Solomon, S. G., &

Martin, P. R. (2018). Receptive Field Properties of Koniocellular On/Off Neurons in the

Lateral Geniculate Nucleus of Marmoset Monkeys. *Journal of Neuroscience*, *38*(48),

10384–10398. https://doi.org/10.1523/JNEUROSCI.1679-18.2018

Fu, S., Wang, S., Si, D., Li, G., Gao, Y., & Liu, Q. (2025). Benchmarking single-cell multi-

modal data integrations. *Nature Methods*, 1–12. https://doi.org/10.1038/s41592-025-

02737-9

Gomez, A. M., Traunmüller, L., & Scheiffele, P. (2021). Neurexins: Molecular Codes for

Shaping Neuronal Synapses. *Nature Reviews. Neuroscience*, *22*(3), 137–151.

https://doi.org/10.1038/s41583-020-00415-7

Hao, Y., Stuart, T., Kowalski, M. H., Choudhary, S., Hoffman, P., Hartman, A., Srivastava, A.,

Molla, G., Madad, S., Fernandez-Granda, C., & Satija, R. (2024). Dictionary learning for

integrative, multimodal and scalable single-cell analysis. *Nature Biotechnology*, *42*(2),

293–304. https://doi.org/10.1038/s41587-023-01767-y

Hendry, S. H. C., & Reid, R. C. (2000). The Koniocellular Pathway in Primate Vision. *Annual*

Review of Neuroscience, *23*(Volume 23, 2000), 127–153.

https://doi.org/10.1146/annurev.neuro.23.1.127

Hodge, R. D., Bakken, T. E., Miller, J. A., Smith, K. A., Barkan, E. R., Graybuck, L. T., Close,

J. L., Long, B., Johansen, N., Penn, O., Yao, Z., Eggermont, J., Höllt, T., Levi, B. P.,

Shehata, S. I., Aevermann, B., Beller, A., Bertagnolli, D., Brouner, K., … Lein, E. S.

(2019). Conserved cell types with divergent features in human versus mouse cortex.

Nature, *573*(7772), Article 7772. https://doi.org/10.1038/s41586-019-1506-7

Huang, E., & Li, S. (2022). Liver Kinase B1 Functions as a Regulator for Neural Development

and a Therapeutic Target for Neural Repair. *Cells*, *11*(18), 2861.

https://doi.org/10.3390/cells11182861

Hubel, D. H., & Livingstone, M. S. (1990). Color and contrast sensitivity in the lateral geniculate

body and primary visual cortex of the macaque monkey. *The Journal of Neuroscience:*

The Official Journal of the Society for Neuroscience, *10*(7), 2223–2237.

https://doi.org/10.1523/JNEUROSCI.10-07-02223.1990

Ichinose, T., & Habib, S. (2022). On and off signaling pathways in the retina and the visual

system. *Frontiers in Ophthalmology*, *2*, 989002.

https://doi.org/10.3389/fopht.2022.989002

Jan, L. Y., & Jan, Y. N. (2012). Voltage-gated potassium channels and the diversity of electrical

signalling. *The Journal of Physiology*, *590*(11), 2591–2599.

https://doi.org/10.1113/jphysiol.2011.224212

Javadi, P., Bouskila, J., Bouchard, J.-F., & Ptito, M. (2015). The endocannabinoid system within

the dorsal lateral geniculate nucleus of the vervet monkey. *Neuroscience*, *288*, 135–144.

https://doi.org/10.1016/j.neuroscience.2014.12.029

Jin, C. Y., Kalimo, H., & Panula, P. (2002). The histaminergic system in human thalamus:

Correlation of innervation to receptor expression. *European Journal of Neuroscience*,

15(7), 1125–1138. https://doi.org/10.1046/j.1460-9568.2002.01951.x

Jüngling, K., Eulenburg, V., Moore, R., Kemler, R., Lessmann, V., & Gottmann, K. (2006). N-

Cadherin Transsynaptically Regulates Short-Term Plasticity at Glutamatergic Synapses

in Embryonic Stem Cell-Derived Neurons. *The Journal of Neuroscience*, *26*(26), 6968–

6978. https://doi.org/10.1523/JNEUROSCI.1013-06.2006

Kaplan, E., & Shapley, R. M. (1982). X and Y cells in the lateral geniculate nucleus of macaque

monkeys. *The Journal of Physiology*, *330*(1), 125–143.

https://doi.org/10.1113/jphysiol.1982.sp014333

Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y., Brenner,

M., Loh, P., & Raychaudhuri, S. (2019). Fast, sensitive and accurate integration of single-

cell data with Harmony. *Nature Methods*, *16*(12), 1289–1296.

https://doi.org/10.1038/s41592-019-0619-0

Lause, J., Berens, P., & Kobak, D. (2021). Analytic Pearson residuals for normalization of

single-cell RNA-seq UMI data. *Genome Biology*, *22*(1), 258.

https://doi.org/10.1186/s13059-021-02451-7

Lavdovskaia, E., Denks, K., Nadler, F., Steube, E., Linden, A., Urlaub, H., Rodnina, M. V., &

Richter-Dennerlein, R. (2020). Dual function of GTPBP6 in biogenesis and recycling of

human mitochondrial ribosomes. *Nucleic Acids Research*, *48*(22), 12929–12942.

https://doi.org/10.1093/nar/gkaa1132

Lee, B. B., & Sun, H. (2009). The chromatic input to cells of the magnocellular pathway of

primates. *Journal of Vision*, *9*(2), 15. https://doi.org/10.1167/9.2.15

Lein, E., Borm, L. E., & Linnarsson, S. (2017). The promise of spatial transcriptomics for

neuroscience in the era of molecular cell typing. *Science*, *358*(6359), 64–69.

https://doi.org/10.1126/science.aan6827

Lerma, J., Morales, M., Vicente, M. A., & Herreras, O. (1997). Glutamate receptors of the

kainate type and synaptic transmission. *Trends in Neurosciences*, *20*(1), 9–12.

https://doi.org/10.1016/S0166-2236(96)20055-4

Levitt, J. B., Schumer, R. A., Sherman, S. M., Spear, P. D., & Movshon, J. A. (2001). Visual

response properties of neurons in the LGN of normally reared and visually deprived

macaque monkeys. *Journal of Neurophysiology*, *85*(5), 2111–2129.

https://doi.org/10.1152/jn.2001.85.5.2111

Lowe, R., Shirley, N., Bleackley, M., Dolan, S., & Shafee, T. (2017). Transcriptomics

technologies. *PLOS Computational Biology*, *13*(5), e1005457.

https://doi.org/10.1371/journal.pcbi.1005457

Maunsell, J. H. R., Ghose, G. M., Assad, J. A., McADAMS, C. J., Boudreau, C. E., & Noerager,

B. D. (1999). Visual response latencies of magnocellular and parvocellular LGN neurons

in macaque monkeys. *Visual Neuroscience*, *16*(1), 1–14.

https://doi.org/10.1017/S0952523899156177

McInnes, L., Healy, J., & Melville, J. (2020). UMAP: Uniform Manifold Approximation and

Projection for Dimension Reduction. *arXiv:1802.03426 [Cs, Stat]*.

http://arxiv.org/abs/1802.03426

Müller, T., Braud, S., Jüttner, R., Voigt, B. C., Paulick, K., Sheean, M. E., Klisch, C.,

Gueneykaya, D., Rathjen, F. G., Geiger, J. R., Poulet, J. F., & Birchmeier, C. (2018).

Neuregulin 3 promotes excitatory synapse formation on hippocampal interneurons. *The*

EMBO Journal, *37*(17), e98858. https://doi.org/10.15252/embj.201798858

Noseda, R., Borsook, D., & Burstein, R. (2017). Neuropeptides and Neurotransmitters That

Modulate Thalamo-Cortical Pathways Relevant to Migraine Headache. *Headache*, *57*

Suppl 2(Suppl 2), 97–111. https://doi.org/10.1111/head.13083

Peng, Y.-R., Shekhar, K., Yan, W., Herrmann, D., Sappington, A., Bryman, G. S., Zyl, T. van,

Do, M. T. H., Regev, A., & Sanes, J. R. (2019). Molecular Classification and

Comparative Taxonomics of Foveal and Peripheral Cells in Primate Retina. *Cell*, *176*(5),

1222-1237.e22. https://doi.org/10.1016/j.cell.2019.01.004

Pietersen, A. N. J., Cheong, S. K., Solomon, S. G., Tailby, C., & Martin, P. R. (2014). Temporal

response properties of koniocellular (blue-on and blue-off) cells in marmoset lateral

geniculate nucleus. *Journal of Neurophysiology*, *112*(6), 1421–1438.

https://doi.org/10.1152/jn.00077.2014

Reid, R. C., & Shapley, R. M. (2002). Space and Time Maps of Cone Photoreceptor Signals in

Macaque Lateral Geniculate Nucleus. *Journal of Neuroscience*, *22*(14), 6158–6175.

https://doi.org/10.1523/JNEUROSCI.22-14-06158.2002

Rousso, D. L., Qiao, M., Kagan, R. D., Yamagata, M., Palmiter, R. D., & Sanes, J. R. (2016).

Two Pairs of ON and OFF Retinal Ganglion Cells Are Defined by Intersectional Patterns

of Transcription Factor Expression. *Cell Reports*, *15*(9), 1930–1944.

https://doi.org/10.1016/j.celrep.2016.04.069

Schneider, I., Cepela, J., Shetty, M., Wang, J., Nelson, A. C., Winterhoff, B., & Starr, T. K.

(2021). Use of "default" parameter settings when analyzing single cell RNA sequencing

data using Seurat: A biologist's perspective. *Journal of Translational Genetics and*

Genomics, *5*(1), 37–49. https://doi.org/10.20517/jtgg.2020.48

Schneider, K. A., Richter, M. C., & Kastner, S. (2004). Retinotopic Organization and Functional

Subdivisions of the Human Lateral Geniculate Nucleus: A High-Resolution Functional

Magnetic Resonance Imaging Study. *Journal of Neuroscience*, *24*(41), 8975–8985.

https://doi.org/10.1523/JNEUROSCI.2413-04.2004

Sherman, S. M., & Guillery, R. W. (2006). *Exploring the thalamus and its role in cortical*

function, 2nd ed (pp. xxi, 484). MIT Press.

Shibata, R., Nakahira, K., Shibasaki, K., Wakazono, Y., Imoto, K., & Ikenaka, K. (2000). A-

Type K+ Current Mediated by the Kv4 Channel Regulates the Generation of Action

Potential in Developing Cerebellar Granule Cells. *Journal of Neuroscience*, *20*(11),

4145–4155. https://doi.org/10.1523/JNEUROSCI.20-11-04145.2000

Sincich, L. C., Park, K. F., Wohlgemuth, M. J., & Horton, J. C. (2004). Bypassing V1: A direct

geniculate input to area MT. *Nature Neuroscience*, *7*(10), 1123–1128.

https://doi.org/10.1038/nn1318

Solomon, S. G., & Lennie, P. (2007). The machinery of colour vision. *Nature Reviews.*

Neuroscience, *8*(4), 276–286. https://doi.org/10.1038/nrn2094

Solomon, S. G., Tailby, C., Cheong, S. K., & Camp, A. J. (2010). Linear and Nonlinear

Contributions to the Visual Sensitivity of Neurons in Primate Lateral Geniculate Nucleus.

Journal of Neurophysiology. https://doi.org/10.1152/jn.01118.2009

Solomon, S. G., White, A. J. R., & Martin, P. R. (2002). Extraclassical Receptive Field

Properties of Parvocellular, Magnocellular, and Koniocellular Cells in the Primate Lateral

Geniculate Nucleus. *Journal of Neuroscience*, *22*(1), 338–349.

https://doi.org/10.1523/JNEUROSCI.22-01-00338.2002

St. Laurent, G., Shtokalo, D., Tackett, M. R., Yang, Z., Vyatkin, Y., Milos, P. M., Seilheimer, B.,

McCaffrey, T. A., & Kapranov, P. (2013). On the importance of small changes in RNA

expression. *Methods*, *63*(1), 18–24. https://doi.org/10.1016/j.ymeth.2013.03.027

Ståhl, P. L., Salmén, F., Vickovic, S., Lundmark, A., Navarro, J. F., Magnusson, J., Giacomello,

S., Asp, M., Westholm, J. O., Huss, M., Mollbrink, A., Linnarsson, S., Codeluppi, S.,

Borg, Å., Pontén, F., Costea, P. I., Sahlén, P., Mulder, J., Bergmann, O., … Frisén, J.

(2016). Visualization and analysis of gene expression in tissue sections by spatial

transcriptomics. *Science*, *353*(6294), 78–82. https://doi.org/10.1126/science.aaf2403

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., Hao, Y.,

Stoeckius, M., Smibert, P., & Satija, R. (2019). Comprehensive integration of single-cell

data. *Cell*, *177*(7), 1888-1902.e21. https://doi.org/10.1016/j.cell.2019.05.031

Sugiyama, H., Ito, I., & Hirono, C. (1987). A new type of glutamate receptor linked to inositol

phospholipid metabolism. *Nature*, *325*(6104), 531–533. https://doi.org/10.1038/325531a0

Sun, S. H., Killian, N. J., & Pezaris, J. S. (2024). *More than expected: Extracellular waveforms*

and functional responses in monkey LGN (p. 2023.11.22.568065). bioRxiv.

https://doi.org/10.1101/2023.11.22.568065

Szíber, Z., Drouet, A., Mondin, M., Levet, F., & Thoumine, O. (2024). Neuroligin-1 dependent

phosphotyrosine signaling in excitatory synapse differentiation. *Frontiers in Molecular*

Neuroscience, *17*, 1359067. https://doi.org/10.3389/fnmol.2024.1359067

Tailby, C., Solomon, S. G., & Lennie, P. (2008). Functional Asymmetries in Visual Pathways

Carrying S-Cone Signals in Macaque. *Journal of Neuroscience*, *28*(15), 4078–4087.

https://doi.org/10.1523/JNEUROSCI.5338-07.2008

Tasic, B., Yao, Z., Graybuck, L. T., Smith, K. A., Nguyen, T. N., Bertagnolli, D., Goldy, J.,

Garren, E., Economo, M. N., Viswanathan, S., Penn, O., Bakken, T., Menon, V., Miller,

J., Fong, O., Hirokawa, K. E., Lathia, K., Rimorin, C., Tieu, M., … Zeng, H. (2018).

Shared and distinct transcriptomic cell types across neocortical areas. *Nature*, *563*(7729),

Article 7729. https://doi.org/10.1038/s41586-018-0654-5

Toth, M. E., Gonda, S., Vigh, L., & Santha, M. (2010). Neuroprotective effect of small heat

shock protein, Hsp27, after acute and chronic alcohol administration. *Cell Stress &*

Chaperones, *15*(6), 807–817. https://doi.org/10.1007/s12192-010-0188-8

Tripathy, R., Leca, I., Dijk, T. van, Weiss, J., Bon, B. W. van, Sergaki, M. C., Gstrein, T.,

Breuss, M., Tian, G., Bahi-Buisson, N., Paciorkowski, A. R., Pagnamenta, A. T.,

Wenninger-Weinzierl, A., Martinez-Reza, M. F., Landler, L., Lise, S., Taylor, J. C.,

Terrone, G., Vitiello, G., … Keays, D. A. (2018). Mutations in MAST1 Cause Mega-

Corpus-Callosum Syndrome with Cerebellar Hypoplasia and Cortical Malformations.

Neuron, *100*(6), 1354-1368.e5. https://doi.org/10.1016/j.neuron.2018.10.044

Vries, S. E. J. de, Lecoq, J. A., Buice, M. A., Groblewski, P. A., Ocker, G. K., Oliver, M., Feng,

D., Cain, N., Ledochowitsch, P., Millman, D., Roll, K., Garrett, M., Keenan, T., Kuan,

L., Mihalas, S., Olsen, S., Thompson, C., Wakeman, W., Waters, J., … Koch, C. (2020).

A large-scale standardized physiological survey reveals functional organization of the

mouse visual cortex. *Nature Neuroscience*, *23*(1), 138–151.

https://doi.org/10.1038/s41593-019-0550-9

Wang, J., Agarwal, D., Huang, M., Hu, G., Zhou, Z., Ye, C., & Zhang, N. R. (2019). Data

denoising with transfer learning in single-cell transcriptomics. *Nature Methods*, *16*(9),

875–878. https://doi.org/10.1038/s41592-019-0537-1

Wang, J., Lu, T., Gui, Y., Zhang, X., Cao, X., Li, Y., Li, C., Liu, L., & Ding, Z. (2023).

HSPA12A controls cerebral lactate homeostasis to maintain hippocampal neurogenesis

and mood stabilization. *Translational Psychiatry*, *13*(1), 280.

https://doi.org/10.1038/s41398-023-02573-5

Webster, C. M., Tworig, J., Caval-Holme, F., Morgans, C. W., & Feller, M. B. (2020). The

Impact of Steroid Activation of TRPM3 on Spontaneous Activity in the Developing

Retina. *eNeuro*, *7*(2). https://doi.org/10.1523/ENEURO.0175-19.2020

Wei, J.-R., Hao, Z.-Z., Xu, C., Huang, M., Tang, L., Xu, N., Liu, R., Shen, Y., Teichmann, S. A.,

Miao, Z., & Liu, S. (2022). Identification of visual cortex cell types and species

differences using single-cell RNA sequencing. *Nature Communications*, *13*(1), 6902.

https://doi.org/10.1038/s41467-022-34590-1

Weston, C. R., & Davis, R. J. (2007). The JNK signal transduction pathway. *Current Opinion in*

Cell Biology, *19*(2), 142–149. https://doi.org/10.1016/j.ceb.2007.02.001

White, A. J. R., Solomon, S. G., & Martin, P. R. (2001). Spatial properties of koniocellular cells

in the lateral geniculate nucleus of the marmoset Callithrix jacchus. *The Journal of*

Physiology, *533*(2), 519–535. https://doi.org/10.1111/j.1469-7793.2001.0519a.x

Wiesel, T. N., & Hubel, D. H. (1966). Spatial and chromatic interactions in the lateral geniculate

body of the rhesus monkey. *Journal of Neurophysiology*, *29*(6), 1115–1156.

https://doi.org/10.1152/jn.1966.29.6.1115

Xu, X., Ichida, J. M., Allison, J. D., Boyd, J. D., Bonds, A. B., & Casagrande, V. A. (2001). A

comparison of koniocellular, magnocellular and parvocellular receptive field properties in

the lateral geniculate nucleus of the owl monkey (Aotus trivirgatus). *The Journal of*

Physiology, *531*(Pt 1), 203–218. https://doi.org/10.1111/j.1469-7793.2001.0203j.x

Yan, L., Herrington, J., Goldberg, E., Dulski, P. M., Bugianesi, R. M., Slaughter, R. S., Banerjee,

P., Brochu, R. M., Priest, B. T., Kaczorowski, G. J., Rudy, B., & Garcia, M. L. (2005).

Stichodactyla helianthus peptide, a pharmacological tool for studying Kv3.2 channels.

Molecular Pharmacology, *67*(5), 1513–1521. https://doi.org/10.1124/mol.105.011064

Yang, Y.-C., Hu, C.-C., & Lai, Y.-C. (2015). Non-additive modulation of synaptic transmission

by serotonin, adenosine, and cholinergic modulators in the sensory thalamus. *Frontiers in*

Cellular Neuroscience, *9*. https://doi.org/10.3389/fncel.2015.00060

Zeidan, A., & Ziv, N. E. (2012). Neuroligin-1 Loss Is Associated with Reduced Tenacity of

Excitatory Synapses. *PLOS ONE*, *7*(7), e42314.

https://doi.org/10.1371/journal.pone.0042314

Zhang, L., & McBain, C. J. (1995). Potassium conductances underlying repolarization and after-

hyperpolarization in rat CA1 hippocampal interneurones. *The Journal of Physiology*,

488(3), 661–672. https://doi.org/10.1113/jphysiol.1995.sp020998

Zheng, G. X. Y., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., Ziraldo, S. B.,

Wheeler, T. D., McDermott, G. P., Zhu, J., Gregory, M. T., Shuga, J., Montesclaros, L.,

Underwood, J. G., Masquelier, D. A., Nishimura, S. Y., Schnall-Levin, M., Wyatt, P. W.,

Hindson, C. M., … Bielas, J. H. (2017). Massively parallel digital transcriptional

profiling of single cells. *Nature Communications*, *8*(1), 14049.

https://doi.org/10.1038/ncomms14049

961