Descending locus coeruleus noradrenergic signaling to spinal astrocyte

subset is required for stress-induced mechanical pain hypersensitivity

Riku Kawanabe-Kobayashi[1,9], Sawako Uchiyama[1,9], Kohei Yoshihara[1,9], Keisuke Koga[2], Daiki

Kojima[1], Thomas McHugh[3], Izuho Hatada[4,5], Ko Matsui[6], Kenji F. Tanaka[7], Makoto Tsuda[1,8, *]

Affiliations

[1]Department of Molecular and System Pharmacology, Graduate School of Pharmaceutical

Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan

[2]Department of Neurophysiology, Hyogo Medical University, 1-1 Mukogawa, Nishinomiya,

Hyogo, Japan

[3]Laboratory for Circuit and Behavioral Physiology, RIKEN Center for Brain Science, 2-1

Hirosawa, Wako, Saitama 351-0198, Japan

[4]Laboratory of Genome Science, Biosignal Genome Resource Center, Institute for Molecular and

Cellular Regulation, Gunma University, 3-39-15 Showa-machi, Maebashi, Gunma 371-8512,

Japan

[5]Viral Vector Core, Gunma University Initiative for Advanced Research (GIAR), 3-39-22,

Showa-machi, Maebashi, Gunma, 371-8511, Japan

[6]Super-network Brain Physiology, Graduate School of Life Sciences, Tohoku University, 2-1-1

Katahira, Aoba-ku, Sendai, Miyagi 980-8577, Japan

[7]Division of Brain Sciences, Institute for Advanced Medical Research, Keio University School of

Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan

[8]Kyushu University Institute for Advanced Study, 744 Motooka, Nishi-ku, Fukuoka 819-0395,

Japan

[9]These authors contributed equally.

*Correspondence: Makoto Tsuda, Ph.D.;

TEL: 81-92-642-6628, **FAX:** 81-92-642-6566, **Email:** tsuda@phar.kyushu-u.ac.jp

Abstract

It is known that stress powerfully alters pain, but its underlying mechanisms remain elusive.

Here, we identified a circuit, locus coeruleus descending noradrenergic neurons projecting to the

spinal dorsal horn ($LC^{\rightarrow SDH}$-NA neurons), that is activated by acute exposure to restraint stress

and is required for stress-induced mechanical pain hypersensitivity in mice. Interestingly, the

primary target of spinal NA released from descending $LC^{\rightarrow SDH}$-NAergic terminals causing the

stress-induced pain hypersensitivity was α_{1A}-adrenaline receptors (α_{1A}Rs) in Hes5-positive

(Hes5$^+$) astrocytes located in the SDH, an astrocyte subset that has an ability to induce pain

sensitization. Furthermore, activation of Hes5$^+$ astrocytes reduced activity of SDH-inhibitory

neurons (SDH-INs) that have an inhibitory role in pain processing. This astrocytic reduction of

IN activity was canceled by an A_1-adenosine receptor (A_1R)-knockdown in SDH-INs, and the

A_1R-knockdown suppressed pain hypersensitivity caused by acute restraint stress. Therefore, our

findings suggest that $LC^{\rightarrow SDH}$-NA neuronal signaling to Hes5$^+$ SDH astrocytes and subsequent

astrocytic reduction of SDH-IN activity are essential for mechanical pain facilitation caused by

stress.

Introduction

Somatosensory information generated at the periphery, such as skin, is conveyed to the dorsal horn of the spinal cord (SDH) via primary afferent sensory neurons and further to the brain[1,2]. The SDH is the first region for processing the peripherally derived sensory information and thus is critical for its proper transmission to and perception in the brain[1-6]. The processing of somatosensory information is not only tightly regulated by complex neuronal circuits within the SDH but also powerfully modulated by descending neurons from the brain[7,8]. The locus coeruleus (LC), a small nucleus located in the brainstem, modulates pain information processing in the SDH via descending noradrenaline (NA) neurons[8-11]. Activation of SDH-projecting LC-NA (LC$^{\rightarrow SDH}$-NA) neurons leads to a release of NA in the SDH[12]. Spinal NA acts on α_{1A}-adrenaline receptors (α_{1A}Rs) expressed on inhibitory neurons (INs)[13,14] and facilitates γ-aminobutyric acid (GABA)-mediated inhibitory synaptic transmission[15,16]. This NA signal has been implicated in the inhibitory control of nociceptive information processing[9-11]. Besides neurons, glial cells in the SDH, such as microglia and astrocytes, also have a remarkable ability to modulate the processing and transmission of somatosensory information[17-19]. Glial modulation has been extensively studied in pathological settings[17-19], but studies using *in vivo* imaging have shown that SDH astrocytes respond to noxious stimuli applied to the periphery under normal conditions[20-23]. Furthermore, we have recently identified a subset of spinal astrocytes defined by expression of hairy and enhancer of split 5 (Hes5) as a new target of NA[22]. Astrocytes express various types of adrenaline receptors[24]; activation of LC$^{\rightarrow SDH}$-NA neurons induces an increase in intracellular Ca^{2+} ([Ca^{2+}]$_i$), an indicator of astrocyte activity[25], via α_{1A}Rs[22], which are the most abundant adrenaline receptors in astrocytes[26,27]. Hes5$^+$ astrocytes are selectively distributed in the superficial laminae of SDH where it receives primary afferent sensory fibers. Activation of α_{1A}Rs of Hes5$^+$ SDH astrocytes by intrathecal administration of NA induces pain sensitization to light mechanical stimuli applied to the skin[22]. Given that most of the varicosities of NAergic axons/terminals are located closer to astrocytic processes than to neuronal synapses[28], Hes5$^+$ astrocytes could be a critical NA target for modulating pain in the SDH.

LC-NA neurons are known to respond to several external and internal stressors[29]. In rodent models of stress, nociceptive stimuli (e.g., electric shock, animal bite, etc.) are frequently employed as stressors. LC-NA neurons respond not only to stimuli that directly excite nociceptors[30,31], but also to stressors that do not involve their direct stimulation[30,32]. Several studies using rodent stress models have shown that exposure to stressors with or without

nociceptor activation induces a remarkable change in pain-associated behavior[33,34]. Despite the

impact of stress on the pain system, the mechanism underlying a link between stress and pain,

especially pain exacerbation, at the neural circuit and cellular levels remains poorly understood,

although it has been proposed that activation of enkephalinergic and GABAergic INs in the SDH

is involved in stress-induced pain inhibition[35]. Since stressful events are known to have a

significant impact on chronic pain[36,37], providing new insights into the mechanisms by which

stress modulates pain is also of clinical importance.

In this study, we investigated how stress modulates pain with a focus on $LC^{\rightarrow SDH}$-NA neurons and

their target Hes5$^+$ astrocytes in the SDH, using multiple approaches. Our experiments using *in*

vivo Ca^{2+} imaging, chemogenetics and optogenetics, cell ablation, electrophysiology, biochemical

imaging for NA release and astrocytic Ca^{2+} responses, conditional gene knockout, and behavioral

analyses demonstrate for the first time that $LC^{\rightarrow SDH}$-NA neuronal signaling to Hes5$^+$ SDH

astrocytes and subsequent astrocytic reduction of SDH-IN activity are essential for stress-induced

mechanical pain hypersensitivity.

Results

$LC^{\rightarrow SDH}$-NA neurons mediate mechanical hypersensitivity after acute restraint stress.

To investigate the mechanism underlying stress-induced pain modulation, we subjected wild-type

(WT) mice to acute restraint stress exposure (Figure 1A), a well-known and frequently used stress

model[29]. Consistent with a previous study[38], restraint stress for 30 min and 1 hour induced

hypersensitivity to light mechanical stimuli (by applying von Frey filaments) to the plantar skin

of the hindpaw (Figure 1B). However, with longer exposure (2 hours) to restraint stress,

mechanical hypersensitivity was attenuated (Figure S1). In addition, mice subjected to 2-hour

restraint stress displayed antinociceptive responses to noxious heat stimuli (the hot-plate test)

(Figure S2), which is consistent with our previous finding[16] as well as numerous other reports

(Table S1). In contrast to restraint stress, forced swim stress, which has been reported to produce

antinociceptive effects on noxious heat[39], did not change mechanical pain sensitivity (Figure S2).

Thus, our data suggest that under our experimental conditions, single exposure of 1-hour restraint

stress causes behavioral pain hypersensitivity to mechanical stimuli, although stress-induced

modulation of pain is a complex phenomenon influenced by multiple factors, including the stress

model, intensity, and duration, as well as the sensory modality used for behavioral testing.

Since 1-hour exposure to restraint stress has also been reported to increase plasma corticosterone

levels[40], in subsequent experiments, we chose 1 hour as the exposure time to restraint stress. To

analyze LC-NA neuronal activity during restraint stress, we performed fiber photometry using

NET-Cre mice (Cre is expressed in NAergic neurons under the control of the promoter of NA

transporter [NET])[41] with microinjection into the LC of an adeno-associated viral (AAV) vector

designed to express GCaMP6s (genetic encoded fluorescent Ca^{2+} indicator) in a Cre-dependent

manner (AAV-FLEx[GCaMP6s]). Three weeks after the injection, GCaMP6s was expressed

selectively in tyrosine hydroxylase$^+$ (TH^+) LC-NA neurons (Figure 1C)[42]. We found that restraint

stress exposure increased the number of Ca^{2+} events in these neurons (Figure 1, D and E; Figure

S3; Movie). The number of Ca^{2+} events was highest during the first 20 min of the restraint stress

exposure and gradually decreased later.

To investigate the role of LC-NA neurons in mechanical hypersensitivity, we utilized a cell

ablation approach[43]; diphtheria toxin (DTX) receptors (DTR) were specifically expressed in LC-

NA neurons by injection of AAV-FLEx[DTR-EGFP] into the LC of *NET-Cre* mice (Figure 1F).

DTR^+ neurons (detected by GFP) mainly co-expressed TH, and these neurons were ablated by

DTX administration (10 µg/kg, two injections 24-hour apart), which resulted in a loss of TH

signal in the LC and reduction of NET signal in the SDH (Figure 1G). We performed behavioral

tests in these mice and observed that ablation of LC-NA neurons abolished mechanical

hypersensitivity after restraint stress exposure (Figure 1H). To specifically ablate $LC^{\rightarrow SDH}$-NA

neurons, a retrograde AAV vector expressing Cre (AAVretro-Cre) was injected into the L4 SDH,

followed by AAV-FLEx[DTR-EGFP] injection into the LC (Figure 1I). In these mice, TH^+

$LC^{\rightarrow SDH}$-NA neurons expressing DTR (detected by EGFP) were ablated after DTX administration

(Figure 1J; Figure S4, A and B) (4.4% of total TH^+ neurons in the LC (189 DTR^+ cells per 4398

TH^+ cells, n = 3 mice)). DTR^+ cells were not observed in A5 or A7 (Figure S4C), confirming the

specificity of $LC^{\rightarrow SDH}$-NAergic pathway targeting in our study. These mice also failed to induce

mechanical hypersensitivity after the restraint stress (Figure 1K). These results suggest that

descending $LC^{\rightarrow SDH}$-NA neurons are necessary for mechanical pain hypersensitivity after acute

restraint stress exposure. In contrast, motor behavior in the rotarod test and behavioral sensitivity

to heat stimuli in the paw-flick test were not affected by ablation of $LC^{\rightarrow SDH}$-NA neurons (Figure

S5), indicating that ablation of these neurons does not produce non-specific behavioral deficits in

motor function or other sensory modalities. Furthermore, we tested whether optogenetic

activation of $LC^{\rightarrow SDH}$-NA neurons could induce mechanical hypersensitivity, using ChrimsonR, a

well-established red-shifted channelrhodopsin[44]. *NET-Cre* mice were microinjected with AAV-

FLEx[ChrimsonR-tdTomato] into the LC (Figure 1L). Three weeks later, we confirmed

expression of ChrimsonR (detected by tdTomato) in TH^+ neurons in the LC and NET^+

axons/terminals in the SDH (Figure 1M; Figure S6). Optogenetic stimulation of $LC^{\rightarrow SDH}$-NA

neuronal axons/terminals in the SDH evoked mechanical hypersensitivity in these mice (Figure

1N). These results together demonstrate that $LC^{\rightarrow SDH}$-NA neurons are activated by acute stress

exposure and are not only necessary for stress-induced mechanical pain hypersensitivity but also

sufficient to phenocopy the behavioral response.

$LC^{\rightarrow SDH}$-NAergic signaling to $Hes5^+$ SDH astrocytes is required for stress-induced

mechanical hypersensitivity.

To investigate the mechanism by which activation of $LC^{\rightarrow SDH}$-NA neurons induces mechanical

hypersensitivity, we first examined if NA is released in the SDH when these neurons are

stimulated, using the genetically encoded fluorescent NA sensor $GRAB_{NE1m}$[45]. In spinal cord

slices from *NET-Cre* mice that had been injected AAV-FLEx[ChrimsonR-tdTomato] into the LC

and $AAV-gfaABC_1D-GRAB_{NE1m}$ (expressing $GRAB_{NE1m}$ under the control of the astrocyte-

selective promoter $gfaABC_1D$[46] into the SDH) (Figure 2A), optogenetic stimulation of $LC^{\rightarrow SDH}$-

NA axons/terminals increased $GRAB_{NE1m}$ fluorescence intensity (Figure 2B), confirming NA

release from $LC^{\rightarrow SDH}$-NA neuron terminals in our slice conditions. The fluorescence intensity

increased in a stimulation period-dependent manner (Figure 2C). Next, to determine the target

cells and receptors of $LC^{\rightarrow SDH}$-NA neurons, we focused on $\alpha_{1A}Rs$ in astrocytes because our

previous study showed that NA increases $[Ca^{2+}]_i$ in astrocytes via $\alpha_{1A}Rs$[22]. Using spinal cord

slices from *NET-Cre* mice that had been injected with AAV-FLEx[ChrimsonR-tdTomato] into

the LC and $AAV-gfaABC_1D-GCaMP6m$ into the SDH (Figure 2A), we found that optogenetic

stimulation of $LC^{\rightarrow SDH}$-NA axons/terminals evoked a rise in $[Ca^{2+}]_i$ in astrocytes (Figure 2, D and

E). Pretreatment with the $\alpha_{1A}R$-specific antagonist silodosin (40 nM) suppressed the Ca^{2+}

responses (Figure 2F), suggesting that NA released from $LC^{\rightarrow SDH}$-NA neurons induces Ca^{2+}

responses in SDH astrocytes via $\alpha_{1A}Rs$. Behaviorally, intrathecal injection of silodosin in *NET-*

Cre;AAV-ChrimsonR mice attenuated mechanical hypersensitivity by optogenetic stimulation of

the $LC^{\rightarrow SDH}$-NA neurons (Figure 2G). Consistent with our previous findings[22], intrathecal

administration of NA (0.1 nmol) evoked transient mechanical hypersensitivity, which was

completely abolished in *Hes5-CreERT2;Adra1a*flox/flox mice treated with tamoxifen ($Hes5^+$

astrocyte-selective conditional knockout of $\alpha_{1A}Rs$[22]: $Hes5^+$ astrocyte–$\alpha_{1A}R$ cKO) (Figure 2H).

Moreover, mechanical hypersensitivity after acute restraint stress was abolished in $Hes5^+$

astrocyte–α_{1A}R cKO mice (Figure 2I). In addition, intrathecal administration of 5,7-

dichlorokynurenic acid (DCK; 10 nmol), an antagonist of D-serine signaling on N-methyl-D-

aspartate (NMDA) receptors, also suppressed mechanical hypersensitivity (Figure 2J). Supporting

this finding, our previous report identified D-serine as a hypersensitivity-inducing factor released

from Hes5$^+$ astrocytes[22]. These results indicate that spinally-released NA from LC$^{\rightarrow SDH}$-NA

neurons after acute restraint stress activates α_{1A}Rs on Hes5$^+$ astrocytes, which causes mechanical

pain hypersensitivity.

Hes5$^+$ astrocytes mask IN function via adenosine A$_1$ receptors.

Besides astrocytes, *Adra1a* mRNA is also expressed in INs in the SDH[13,14]. However, *Vgat-*

Cre;*Adra1a*$^{flox/flox}$ mice (Vgat$^+$ IN-selective conditional knockout of α_{1A}Rs: Vgat$^+$ IN–α_{1A}R cKO

mice)[14,16] had no effect on the NA (0.1 nmol)-induced mechanical hypersensitivity (Figure 3A). A

similar behavioral phenotype was also observed after intrathecal administration of phenylephrine

(0.05 nmol), a selective agonist for α_1Rs (Figure 3B). Furthermore, stress-induced mechanical

hypersensitivity was not affected in Vgat$^+$ IN–α_{1A}R cKO mice (Figure 3C). Considering previous

data that SDH-INs (including Vgat$^+$) respond to NA via α_{1A}Rs at NA concentrations similar to

those that elicit Hes5$^+$ astrocyte response[16,22], it remains unclear why intrathecally administered

NA and spinally released NA from LC$^{\rightarrow SDH}$-NA neuron terminals (after acute restraint stress)

have no suppressive effect via α_{1A}R-mediated Vgat$^+$ IN activation on mechanical hypersensitivity.

We then hypothesized an interaction between Hes5$^+$ astrocytes and Vgat$^+$ INs as per a previous

report that optogenetic activation of rat SDH astrocytes leads to ATP release; ATP is converted to

adenosine extracellularly which subsequently suppresses the activity of SDH-INs via adenosine

A$_1$ receptors (A$_1$Rs)[47]. By performing whole-cell current-clamp recordings of Vgat$^+$ INs in spinal

cord slices from *Vgat-Cre*;*Rosa-tdTomato* mice, we found that bath application of NA (20 µM)

evoked depolarization in 22.2% (10/45 cells) of tdTomato$^+$ cells (Vgat$^+$ INs), with 30% of these

showing action potentials (3/10 cells) (Figure 3D). In our experiments, four antagonists for

AMPA and NMDA receptors and GABA$_A$ and glycine receptors were added to the aCSF to block

synaptic inputs from other neurons to the recorded Vgat$^+$ neurons[48,49], suggesting that NA directly

acts on the recorded SDH Vgat$^+$ neurons to produce excitation. The proportion of Vgat$^+$ INs

depolarized by NA is similar to that of *Adra1a* mRNA-expressing INs[14]. In 50% (5/10 cells) of

depolarized Vgat$^+$ INs, additional application of the A$_1$R-selective agonist N6-

cyclopentyladenosine (CPA; 1 µM) suppressed NA-induced depolarization/action potentials

(Figure 3, D and E). Supporting this finding, RNAscope *in situ* hybridization detected *Adora1*

mRNA fluorescence in 63.5 ± 6.9% of *Slc32a1* (*Vgat*)$^+$ INs (*n* = 497, from three mice) (Figure

3F). Because A_1Rs are also expressed in the dorsal root ganglion neurons[50], excitatory neurons

and microglia[51] in the SDH, we employed the CRISPR–Cas9 system using AAV vectors to

specifically knockdown A_1Rs in SDH-INs. AAV vectors designed to express the *Staphylococcus*

aureus Cas9 (SaCas9)[52] in a Cre-dependent manner (AAV-FLEx[SaCas9]) and single guide

RNA-expression vectors (AAV-FLEx[mCherry]-U6-sgAdora1 or -FLEx[mCherry]-U6-sgYFP

[as a control]) were co-microinjected into the SDH of *Vgat-Cre* mice (Figure 3G). SaCas9

(detected by HA-tag) and mCherry-labeled cells in the SDH were immunolabeled with paired box

2 (PAX2) (Figure 3G), a marker of INs[53]. In these SDH-Vgat$^+$ IN-selective A_1R conditional

knockdown mice (SDH-Vgat$^+$ IN–A_1R cKD mice), the inhibitory effect of CPA on NA-induced

depolarization/action potentials was completely abolished in mCherry$^+$ neurons (Figure 3, H and

I). Using a chemogenetic approach with modified human muscarinic G_q-protein-coupled

receptors (hM3Dq)[54], we further examined the effect of G_q-stimulated Hes5$^+$ astrocytes on

inhibitory synaptic transmission. *Hes5-CreERT2* mice were microinjected with AAV-

FLEx[hM3Dq-HA] into the SDH (*Hes5-CreERT2*;AAV-hM3Dq mice). hM3Dq (detected by

HA-tag) was expressed in the SDH, and these cells were double-labeled with glial fibrillary

acidic protein (GFAP) and SRY-related high-mobility group box 9 (SOX9), markers of astrocytes

(Figure 3J). Our previous studies using *Hes5-CreERT2* mice have confirmed that hM3Dq is not

expressed in other cell types (neurons, oligodendrocytes, or microglia)[22,55]. Using spinal cord

slices from these mice, we performed whole-cell voltage-clamp recordings of substantia

gelatinosa (SG) neurons. Bath application of hM3Dq agonist clozapine N-oxide (CNO; 100 µM)

reduced the frequency of spontaneous inhibitory postsynaptic currents (sIPSCs) in SG neurons

(Figure 3, K and L). The concentration of CNO has been used in our previous study[22] but is

relatively high; however, 100 µM CNO most effectively replicated the NA-induced Ca^{2+} signals

in astrocytes (Figure S7, A and B). In addition, spinal cord slices from mice that did not express

hM3Dq, 100 µM CNO had no effect on Ca^{2+} responses in SDH astrocytes (Figure S7A) and

sIPSCs in SG neurons (Figure S7C), confirming that the observed effect of CNO is not non-

specific. The reduction of sIPSCs by astrocytic hM3Dq activation was prevented by pretreatment

with the A_1R-specific antagonist, 8-cyclopentyl-1,3-dimethylxanthine (CPT; 1 µM) (Figure 3, M

and N). These results suggest that a signaling pathway from Hes5$^+$ astrocytes suppresses the

activity of Vgat$^+$ INs via A_1Rs in the SDH.

Hes5$^+$ astrocyte-mediated inhibitory signals to INs affect mechanical hypersensitivity.

We investigated whether A_1Rs in SDH-Vgat$^+$ INs contributed to the mechanical hypersensitivity

elicited by intrathecal NA (0.1 nmol). Acute inhibition of spinal A_1Rs by intrathecal

administration of CPT (3 nmol) suppressed NA-induced mechanical hypersensitivity (Figure 4A).

Similarly, SDH-Vgat$^+$ IN–A_1R cKD mice showed a significant reduction in the behavioral

hypersensitivity by NA (Figure 4B), indicating that A_1Rs in SDH-Vgat$^+$ INs contribute to

mechanical hypersensitivity by spinal NA. Furthermore, SDH-Vgat$^+$ IN–A_1R cKD mice and

intrathecal-CPT-pretreated mice exhibited a significant attenuation of the stress-induced

mechanical hypersensitivity (Figure 4, C and D). These results suggest that Hes5$^+$ astrocyte-

mediated negative control for SDH-Vgat$^+$ INs via A_1R-mediated signals is crucial for stress-

induced mechanical hypersensitivity. Moreover, we investigated the role of this interaction in

stress-induced alteration of somatosensory information processing in the SDH using c-FOS, a

neuronal activity marker, and phosphorylated extracellular signal-regulated kinase (pERK), a

neuronal nociceptive input marker in superficial SDH neurons[56]. Consistent with the results of a

previous report[57], Aβ fiber stimulation of the hindpaw alone did not increase the number of c-

FOS$^+$ (Figure 4, E and F) and pERK$^+$ neurons (Figure 4, G and H) in the superficial SDH

(laminae I–IIi using isolectin B4 [IB4], a marker of lamina IIi). Similarly, acute restraint stress

alone did not change the number of c-FOS$^+$ and pERK$^+$ neurons (Figure 4, E–H). Interestingly,

Aβ fiber stimulation following acute restraint stress exposure significantly increased the number

of c-FOS$^+$ and pERK$^+$ neurons in the superficial SDH (Figure 4, E–H). Intrathecal injection of

CPT prevented the stress-induced increase in c-FOS$^+$ and pERK$^+$ neurons (Figure 4, E–H). These

results suggest that the abnormal responsiveness of superficial SDH neurons to signals from Aβ

fibers in mice with acute restraint stress, involves adenosine signals that are important for

negatively controlling Vgat$^+$ INs by NA-stimulated Hes5$^+$ astrocytes and for mechanical

hypersensitivity.

Discussion

External and internal stressors elicit a variety of biological responses including pain modulation.

Acute exposure to a stressor inhibits or facilitates behavioral responses to thermal and mechanical

stimuli applied to the skin[33,38,58]. While the mechanism underlying pain inhibition has been

studied[33], the neural basis of pain facilitation by stress remains poorly understood. In this study,

we identified a circuit, the descending LC$^{→SDH}$-NA neurons, that is activated by acute restraint

stress exposure and demonstrated that the circuit is necessary for stress-induced mechanical pain

hypersensitivity. A notable finding of this study was that Hes5$^+$ astrocytes, a subpopulation of

spinal cord astrocytes[22], are the primary target of NA released from descending $LC^{\rightarrow SDH}$-NA

neurons for mechanical pain hypersensitivity caused by stress. Thus, this top-down signaling

pathway from $LC^{\rightarrow SDH}$-NA neurons to Hes5$^+$ SDH astrocytes is an essential link between stress

and mechanical pain sensitization.

Spinal NA derived from LC-NA neurons has been classically implicated in pain inhibitory

control[9-11]. The suggested mechanism to this effect is its action on SDH-INs via α_{1A}Rs[9,16]. In

contrast, recent our study has unveiled another facet of spinal NA signaling, which is

pronociceptive, by identifying Hes5$^+$ astrocytes in the SDH[22]. Mechanistically, spinal NA acts on

α_{1A}Rs expressed on Hes5$^+$ astrocytes, increases intracellular Ca^{2+} levels via a G_q/PLC/IP$_3$

pathway, and causes mechanical hypersensitivity via release of D-serine[22], an activator of NMDA

receptors. This study showed that the optogenetic activation of $LC^{\rightarrow SDH}$-NA neurons produced

mechanical hypersensitivity via α_{1A}Rs in SDH-Hes5$^+$ astrocytes, indicating the pronociceptive

effect of $LC^{\rightarrow SDH}$-NAergic signaling. In contrast, it has also been reported that a similar opto- or

chemogenetic activation produces an antinociceptive effect on thermal stimuli[59,60]. The reason for

these differing behavioral outcomes remains unclear; however, the stimulus modality used for the

behavioral assay (mechanical vs. thermal) may be responsible, since the SDH circuits for

mechanical and thermal information processing have been shown to be distinct[3,61-64].

Alternatively, the degree of $LC^{\rightarrow SDH}$-NA neuronal activity may also be involved. Although direct

comparisons between studies reporting pro- and anti-nociceptive effects are difficult, our previous

studies demonstrated that intrathecal administration of high doses of NA in naïve mice does not

induce mechanical pain hypersensitivity and produces an antinociceptive effect on noxious

heat[22,16]. Indeed, a continuous stimulation of descending LC-NA neurons over several minutes

has been reported to produce antinociceptive effect[65]. Collectively, these findings indicate that the

role of $LC^{\rightarrow SDH}$-NA neurons in modulating nociceptive signaling in the SDH is more complex

than previously appreciated, and these differences may be related to $LC^{\rightarrow SDH}$-NA neuron activity,

NA levels in the SDH, and the differential responses of SDH neurons in the superficial versus

deeper laminae.

Given that SDH-INs also express α_{1A}Rs[13,14], it is unclear why pain hypersensitivity is the

preferentially manifested behavioral phenotype after activation of $LC^{\rightarrow SDH}$-NAergic signal. This

study showed that NA-stimulated Hes5$^+$ astrocytes negatively regulate the activity of SDH-Vgat$^+$

INs. This astrocytic attenuation of SDH-Vgat$^+$ IN activity may involve adenosine signaling

because A$_1$R cKD in SDH-Vgat$^+$ INs canceled the suppressive effect of A$_1$R agonist on NA-

induced enhancement of Vgat$^+$-IN activity and NA-induced mechanical hypersensitivity. The

adenosine-mediated inhibition of SDH-INs is consistent with the data from a previous study[47]. In

addition, we also found that knockdown of A$_1$Rs in Vgat$^+$ neurons tended to increase the

proportion of Vgat$^+$ neurons with NA-induced depolarizing responses (Figure S8). Thus, these

results suggest that spinal NA acts on α_{1A}Rs expressed by both Hes5$^+$ astrocytes and Vgat$^+$ INs;

however, NA-stimulated Hes5$^+$ astrocytes suppress Vgat$^+$ IN activity by adenosine signaling via

A$_1$Rs; this induces a state where the Hes5$^+$ astrocyte-derived pain-facilitating effect predominates,

causing mechanical pain hypersensitivity (Figure 4I).

While a neuronal model for gating peripherally derived sensory signals in the SDH has been

previously established by identifying several subpopulations of INs[1-5], our study proposes a new

regulatory mechanism in this model in which Hes5$^+$ astrocytes act as non-neuronal gating cells in

the SDH for brain-derived NAergic signaling. This astrocytic gating control of INs may affect

processing and transmission of somatosensory information in the SDH. Indeed, a

pharmacological blockade of the adenosine signal to SDH-Vgat$^+$ INs suppressed the Aβ fiber-

mediated c-FOS and pERK expression in the superficial SDH of stressed mice.

Preproenkephalin$^+$ IN population may be among the SDH-Vgat$^+$ INs involved in stress-induced

pain modulation[35]. Interfering with GABA/glycine signaling in the SDH results in abnormal

synaptic inputs from primary afferent Aβ fibers onto superficial SDH neurons[66]. Our view is also

supported by previous findings from *in vivo* patch-clamp recordings, which show that, in about a

half of SG neurons tested, NA enhances excitatory synaptic responses evoked by tactile stimuli

(air puff to the skin)[67]. α_{1A}R expression has been reported to be detected in DRG neurons, but its

level seems to be much lower than in the spinal cord[68], and primary afferent glutamatergic

synaptic transmission has been shown to be unaffected by α_{1A}R agonists[69,70]. Additionally, our

hypothesis that Hes5$^+$ astrocytes co-release D-serine and ATP/adenosine in response to α_{1A}R

activation by NA is supported by previous findings in the neocortex[71]. Aβ fiber inputs into

neurokinin 1 receptor-positive projection neurons in SDH lamina I following inhibition of

GABA/glycine signaling is canceled by a pharmacological blockade of NMDA receptors[66],

supporting our data that the stress-induced mechanical hypersensitivity was suppressed by the

antagonist of D-serine site of NMDA receptors DCK. However, we cannot exclude the possibility

that D-serine also acts on NMDA receptors expressed by Vgat$^+$ INs. Nevertheless, given that

intrathecal injection of D-serine in naïve mice induces mechanical pain hypersensitivity[22], it

appears that the pronociceptive effect of D-serine in the SDH is primarily associated with

enhanced pain processing and transmission, presumably via NMDA receptors on excitatory

neurons. In addition, the effect of astrocytic adenosine on spinal pain processing has been

controversial: both, pronociceptive (consistent with our findings)[47] and antinociceptive[72]. Such

different effects might be related in part to the regionally restricted astrocytic subpopulations and

factors for their activation. Pro- and anti-nociceptive effects have been reported to be mediated by

astrocytes located in superficial and deeper laminae and stimulated by NA (via G protein-coupled

$\alpha_{1A}Rs$)[22] and ATP (via ionotropic P2X7 receptors)[72], respectively. In particular, the regional

differences in the role of astrocytes has been demonstrated by our previous study in which a

selective chemogenetic stimulation of astrocytes in deeper SDH did not cause mechanical

hypersensitivity[22]. Thus, it is conceivable that locally released adenosine from NA-stimulated

$Hes5^+$ astrocytes, following acute restraint stress, may suppress SDH-$Vgat^+$ IN function, which

results in mechanical pain hypersensitivity. A highly localized adenosine release from $Hes5^+$

astrocytes and the spatially restricted neuron–astrocyte communication via adenosine signaling

need to be investigated in future studies.

In this study, we showed mechanical hypersensitivity by acute exposure to restraint stress and

investigated its mechanisms; however, there are also many studies reporting acute stress-induced

antinociception (Table S1). A precise explanation for the bidirectional outcomes remains elusive.

However, it is widely recognized that the extent and direction of pain modulation may be due to

several factors, including the nature, duration, and intensity of the stressor, as well as the sensory

modality assessed in behavioral tests[33,34]. In fact, most studies reporting antinociceptive effects of

acute restraint stress assessed behavioral responses to heat stimuli (Table S1). This observation is

consistent with the findings in our previous study[16] and the present study (Figure S2). In contrast

to the robust and consistent antinociceptive effects observed with thermal stimuli, some studies

evaluating behavioral responses to mechanical stimuli have reported stress-induced

hypersensitivity[38] (Table S1), which aligns with our current findings. Nevertheless, acute and

short-term exposure ($\leq$1 hour) to restraint stress induces mechanical pain hypersensitivity, which

was not observed with longer exposure (2 hours) (Figure S1) (although a 2-hour restraint stress

exposure suppresses behavioral response to nociceptive thermal stimuli[16]). Thus, descending

$LC^{\rightarrow SDH}$-NA neurons, $Hes5^+$ SDH astrocytes, or their output signals involved in modulating

mechanical pain behavior may be attenuated by unidentified inhibitory signals evoked by longer

exposure to restraint stress. A similar bidirectional behavioral outcome on mechanical pain has

also been observed with intrathecal NA; low doses of NA induce mechanical pain

hypersensitivity while higher doses do not[22]. Thus, the bidirectional change may depend on the

level of NA neuronal activity and NA content in the SDH, which is supported by a previous
report[73].

Clinically, stressor-evoked psychological and physical responses are considered critical for the
development of chronic pain[36], especially nociplastic pain, a condition that arises from altered
nociceptive signaling without actual or threatened tissue damage, or disease or lesion of the
somatosensory system[37]. In fact, patients with chronic pain have a high incidence of stress-related
psychiatric disorders, such as anxiety, fear, and depression[11,74]; stressful life events are known to
be a key factor in establishing fibromyalgia and enhance pain in patients with irritable bowel
syndrome and headaches[37]. Thus, our findings provide a mechanistic insight into the link between
stress and pain modulation and may help in understanding and developing a new strategy for
chronic pain, including nociplastic pain.

There are several limitations to this study. First, under our experimental conditions, where AAV
vectors were injected into both the LC and SDH (L4) to express genes in $LC^{\rightarrow SDH}$ neurons, only a
small proportion (4.4%) of all LC-NA neurons could be manipulated (Figure 1, I–K; Figure S4, B
and C), although we confirmed specific expression of DTRs in NA neurons within the LC, but
not in A5 or A7 (Figure S4A). This low proportion is likely due to the selective targeting of LC-
NA neurons that project to the L4 segment of the SDH. The total number of SDH-projecting LC-
NA neurons across all spinal segments may be much higher. Given that primary afferent sensory
fibers from the plantar skin of the hindpaw predominantly terminate in the L4 segment of the
SDH, it is plausible that ablation of even a relatively small number of NA neurons in the LC can
have a significant impact on behavior. Second, in our model (Figure 4I), we proposed that NA
released by $LC^{\rightarrow SDH}$ NA neurons inhibits SDH-INs. Although we demonstrated that optogenetic
stimulation of $LC^{\rightarrow SDH}$ NA terminals in the SDH induces endogenous NA release, we were unable
to determine whether the released NA modulates the function of Hes5$^+$ astrocytes and Vgat$^+$ INs,
or their interactions. Addressing this question would require the use of *Vgat-Cre* and/or *Hes5-*
CreERT2 mice; however, employing these lines precludes the use of *NET-Cre* mice, which are
essential for specific and efficient expression of ChrimsonR in $LC^{\rightarrow SDH}$ NA neurons. Third, in our
in vivo optogenetic experiments showing that stimulation of $LC^{\rightarrow SDH}$-NAergic terminals in the
SDH causes mechanical pain hypersensitivity (Figure 1N), light could reach only the superficial
laminae, even though ChrimsonR expression in $LC^{\rightarrow SDH}$-NAergic terminals was also observed in
both superficial and deeper laminae of the SDH (Figure S6). Kucharczyk et al. have reported that
NA released from descending pathways reduces activity of wide-dynamic-range neurons in

deeper laminae[65]. Therefore, the roles of NA signaling in superficial versus deeper laminae

should also be further investigated.

An important future subject is to clarify the discrepancy for the pro- and antinociceptive role of

$LC^{\rightarrow SDH}$-NAergic signaling in the SDH. Since one possible candidate could be related to the

sensory modality used for behavioral testing, other assays (e.g., Randall-Selitto test, CatWalk gait

analysis, and weight-bearing test) should be tested. Furthermore, a difference between the

duration of behavioral responses and Ca^{2+} events after acute restraint stress should also be

addressed in further investigation. The behavioral effect of stress on PWT (Figure 1E) persisted

for 120 min after stress exposure, whereas Ca^{2+} events changes were only observed during stress

exposure (Figure 1B). A similar temporal difference is also observed following intraplantar

injection of capsaicin[22]; while $LC^{\rightarrow SDH}$-NA neuron-mediated astrocytic Ca^{2+} responses in SDH

astrocytes last for 5–10 min after injection, behavioral hypersensitivity peaks around 60 min post-

injection and gradually returns to baseline over the subsequent 60–120 min. These findings raise

the possibility that astrocyte-mediated mechanical pain hypersensitivity in the SDH may involve

a sustained alteration in spinal neural function, such as central sensitization.

Methods

Animals: Male C57BL/6J mice (CLEA Japan), male and female *NET-Cre* mice [Tg(*Slc6a2-cre*)#Stl] (kindly provided by Prof. Thomas McHugh, RIKEN Center for Brain Science)[41], *Hes5-CreERT2* mice [Tg(*Hes5-cre/ERT2*)2Vtlr] (kindly provided by Prof. Verdon Taylor)[75], *Adra1a*flox/flox mice[22], *Vgat-Ires-Cre* mice [*Slc32a1*tm2(cre)Lowl/J] (Stock No: 016962, The Jackson Laboratory)[76], and *Rosa26-tdTomato* mice [B6.Cg-*Gt(ROSA)26Sor*tm14(CAG-tdTomato)Hze/J] (Stock No: 007914, The Jackson Laboratory)[77] were used. To generate *Hes5-CreERT2;Adra1a*flox/flox mice[22] or *Vgat-Cre;Adra1a*flox/flox mice[16], *Adra1a*flox/flox mice were crossed with *Hes5-CreERT2* mice or *Vgat-Ires-Cre* (*Vgat-Cre*) mice, and the obtained *Hes5-CreERT2;Adra1a*flox/+ mice or *Vgat-Cre;Adra1a*flox/+ mice were further crossed with *Adra1a*flox/flox mice. Cre-negative *Adra1a*flox/flox littermate mice were used as controls. For induction of CreERT2 recombinase activity, *Hes5-CreERT2* mice were given an intraperitoneal (i.p.) injection of tamoxifen (#T5648, Sigma-Aldrich; 2 mg dissolved in 100 μl corn oil [#032-17016; Wako, Saitama, Japan]) once a day for 5–10 successive days. We used tamoxifen-injected mice for further analyses 7 days or more after the last injection. All mice used were 8–12 weeks of age at the start of each experiment and were housed at temperature and humidity ranges of 21–23°C and 40–60%, respectively, with a 12-hour light-dark cycle. All animals were fed food and water ad libitum. All animals were housed in standard polycarbonate cages in groups of same-sex littermates. All animal experiments were conducted according to relevant national and international guidelines contained in the 'Act on Welfare and Management of Animals' (Ministry of Environment of Japan) and 'Regulation of Laboratory Animals' (Kyushu University) and under the protocols approved by the Institutional Animal Care and Use committee review panels at Kyushu University.

Restraint stress model: According to the methods described in our previous study[16], mice were restrained by placing them in a Falcon® 50-ml conical tube (#352070; Corning) with a hole in the tip. The space remaining behind the mouse was filled with a paper towel. Mice were allowed to breathe but not turn around. Control mice were handled briefly and placed in home cages without water or food for 1 hour.

Measurement of behavioral response to mechanical stimuli using von Frey filaments: Mice were placed individually in an opaque plastic cylinder, which was placed on a wire mesh, and habituated for 0.5–1 hour to allow acclimatization to the new environment. After that, calibrated von Frey filaments (0.02–2.0 g; #NC12775; North Coast Medical) were applied to the plantar surface of the hindpaw of mice from below the mesh floor. 50% paw withdrawal threshold was

calculated using the up-down method[78]. The basal mechanical threshold was determined by

performing von Frey test before restraint stress exposure, AAV microinjection, DTX

administration, optogenetic stimulation, intrathecal injection of several drugs and tamoxifen

administration. Mechanical sensitivity was evaluated at 30, 60, 90, 120 and 180 min after restraint

stress or optogenetic stimulation, or at 30 min after intrathecal injection of NA or Phe.

Measurement of behavioral response to thermal stimuli in hot-plate and paw-flick tests: For

the hot-plate test (HOT/COLD PLATE ANALGESIA METER, MK-350HC, Muromachi Kikai,

Tokyo, Japan), the latency to exhibit nociceptive behaviors (licking or jumping) in response to

thermal stimulation (50°C)[16,79] was recorded. To prevent tissue damage, a cut-off time was set at

60 sec. For the paw-flick test (Tail Flick Unit, 7360, Ugo Basile, Gemonio, Italy), the latency to

elicit nociceptive responses (paw withdrawal) against heat stimulation at 40 V[22] was recorded,

with a cut-off time of 10 sec to avoid tissue damage.

Measurement of behavioral response to motor function by Rotarod test: Before AAV

injection, each mouse was trained on a rotarod (3 cm diameter, 8 r.p.m.; Rotarod, KN-75,

Natsume Seisakusho, Tokyo, Japan) until it could remain on the apparatus for 60 sec without

falling[80]. The mice then underwent a 60-s rotarod test both before and after DTX administration.

Recombinant AAV (rAAV) vector production: pAAV-CAG-FLEx[GCaMP6s]-WPRE

(#100842) and pAAV-synapsin (Syn)-FLEx[ChrimsonR-tdTomato]-WPRE (#62723) were

purchased from Addgene. The genes encoding GCaMP6m (#40754; Addgene), Cre, GRAB$_{NE1m}$

(#123308; Addgene) and mCherry were subcloned into the pENTR plasmid (Thermo Fisher

Scientific). To produce AAV vectors, we inserted GCaMP6m, GRAB$_{NE1m}$ and mCherry into

pZac2.1-gfaABC$_1$D-WPRE plasmid and Cre into pZac2.1-enhanced synapsin (ESYN)-WPRE

plasmid. To produce the AAV vector for the Cre-switch system, vectors containing the promoters

encoding EF1α were generated from pAAV-CA-FLEx (#38042; Addgene) by substituting the

promoter. We then inserted AcGFP, DTR-EGFP (kindly provided by Prof. Kenji Kohno, Nara

Institute of Science and Technology), mCherry and hM3Dq (#45547; Addgene) into pAAV-

EF1α-FLEx . The genes encoding U6-sgRNA (#61593; Addgene) and SaCas9 (#78601;

Addgene) were subcloned into the pENTR plasmid. Synthetic oligonucleotides including the

targeting sequence for *Adora1* (5'-AAGTTCCGGGTCACCTTTCTG-3') and non-targeting

sequence (5'-CCATGTGATCGCGCTTCTCGT-3') were replaced with the targeting site in the

original pENTR-U6-sgBsa1 plasmid. The resulting U6-sgRNA cassette was transferred into

pAAV-CMV-FLEx[mCherry]-WPRE plasmid to generate pAAV-CMV-FLEx[mCherry]-WPRE-

U6-sgAdora1/sgYFP (sgControl). The SaCas9 cassette was transferred into pAAV-CMV-FLEx-

WPRE plasmid to generate pAAV-CMV-FLEx[SaCas9]-WPRE. rAAV vectors were produced

from human embryonic kidney 293T (HEK293T) cells with triple transfection [pZac or pAAV,

cis plasmid; pAAV2/5 (University of Pennsylvania Gene Therapy Program Vector Core),

pAAV2/9 (University of Pennsylvania Gene Therapy Program Vector Core) or pAAV2/retro

(#81070; Addgene), trans plasmid; pAd DeltaF6, adenoviral helper plasmid (University of

Pennsylvania Gene Therapy Program Vector Core)] and purified by two cesium chloride density

gradient purification steps. The vector was dialyzed against phosphate-buffered saline (PBS)

containing 0.001% (v/v) Pluronic-F68 (#24040032; Thermo Fisher Scientific) using Vivaspin

Turbo 15 100,000 MWCO (#VS15T41; Sartorius). The genome titer of rAAV was determined by

Pico Green fluorometric reagent (#P7589; Thermo Fisher Scientific) following denaturation of

the AAV particles. Vector were stored at −80°C until use.

Intra-SDH and intra-LC injection of rAAV vectors: Viral vector injection was performed in

accordance with our previously described method[22,81]. Mice were deeply anesthetized by

subcutaneous injection of ketamine (100 mg/kg) and xylazine (10 mg/kg). For intra-SDH

injection, the skin was incised at Th11–L4 vertebrae, and custom-made clamps were attached to

the caudal sites of the vertebral column. Paraspinal muscles around the left side of the interspace

between Th13 and L1 vertebrae were removed, and the dura mater and arachnoid membrane were

carefully incised using the tip of a 30-G needle to make a small window allowing a glass

microcapillary to insert directly into the SDH. The microcapillary was inserted into the unilateral

SDH (around 120–150 μm in depth from the surface of the dorsal root entry zone). rAAV

solutions (approximately 500 nl) were injected using a Micro4 Micro Syringe Pump Controller

(World Precision Instrument). After microinjection, the glass microcapillary was removed, the

skin was sutured with 5-0 silk, and the mice were kept on a heating pad until recovery. For intra-

LC injection, rAAV solutions were injected (approximately 300 nl in one site) adjacent to the LC

(anteroposterior (AP): −5.4 mm from the bregma; mediolateral (ML): ±0.9 mm; dorsoventral

(DV): −3.2 mm from the dura). Only AAV2/9-CAG-FLEx[GCaMP6s]-WPRE was injected

unilaterally, and the other AAV vectors were injected bilaterally. To keep the bregma and lambda

in the same horizontal plane, tolerance was maintained at <50 μm in the dorsoventral axis

between the bregma/lambda. The microcapillary was withdrawn 3 min after ending the injection.

The skin was then sutured with 5-0 silk, and the mice were kept on a heating pad until recovery.

We used virus-injected mice for further analyses 3 weeks or more after the last injection. The

following viral titers were used: AAV2/9-CAG-FLEx[GCaMP6s]-WPRE, AAV2/9-gfaABC₁D-

GCaMP6m-WPRE, AAV2/9-EF1α-FLEx[AcGFP]-WPRE (for *NET-Cre* mice), AAV2/9-EF1α-

FLEx[DTR-EGFP]-WPRE (for *NET-Cre* mice), AAV2/9-EF1α-FLEx[mCherry], AAV2/9-hSyn-

FLEx[ChrimsonR-tdTomato], AAV2/9-gfaABC$_1$D-GRAB$_{NE1m}$-WPRE, AAV2/5-EF1α-

FLEx[hM3Dq], AAV2/9-CMV-FLEx[SaCas9]-WPRE and AAV2/5-gfaABC$_1$D-mCherry: $1.0 \times$

10^{12} genome copies (GC)/ml; AAV2/retro-ESYN-Cre-WPRE: 3.0×10^{12} GC/ml; AAV2/9-EF1α-

FLEx[AcGFP]-WPRE (for WT mice) and AAV2/9-EF1α-FLEx[DTR-EGFP]-WPRE (for WT

mice): 2.0×10^{12} GC/ml; AAV2/9-CMV-FLEx[mCherry]-WPRE-U6-sgYFP and AAV2/9-CMV-

FLEx[mCherry]-WPRE-U6-sgAdora1: 0.5×10^{12} GC/ml.

Immunohistochemistry: Mice were deeply anesthetized with an i.p. injection of pentobarbital

and transcardially perfused with PBS (#041-20211; Wako, Saitama, Japan) followed by ice-cold

4% paraformaldehyde (PFA; #162-16065; Wako, Saitama, Japan)/PBS. The transverse L4

segments of the spinal cord and brain were removed, postfixed in the same fixative for 3 hours

(spinal cord) or overnight (brain) at 4°C, and placed in 30% sucrose solution for 24–48 hours at

4°C. After incubation, the tissues were embedded in OCT compound (#4583; Sakura Finetek

Japan, Osaka, Japan) and stored at –25°C before use. Transverse spinal cord and brain sections

(30 µm) were incubated in blocking solution (3% normal goat serum [#S-1000; Vector

Laboratories] or normal donkey serum [#017-000-121; Jackson ImmunoResearch]) for 2 hours at

room temperature and then incubated for 48 hours at 4°C with primary antibodies: polyclonal

rabbit anti-tyrosine hydroxylase (TH; 1:1000; #AB152; Millipore); polyclonal sheep anti-TH

(1:1000; #AB1542; Millipore); monoclonal mouse anti-noradrenaline transporter (NET; 1:2000;

#NET05-2; Mab Technologies); polyclonal rabbit anti-green fluorescent protein (GFP; 1:1000;

#598; MBL International); monoclonal rabbit anti-hemagglutinin (HA)-tag (1:1000; #3724; Cell

Signaling); monoclonal rat anti-glial fibrillary acidic protein (GFAP; 1:2000; #13-0300;

Invitrogen); polyclonal goat anti-SRY-related high-mobility group box 9 (SOX9; 1:1000; #

AF3075; R&D Systems); polyclonal goat anti-paired box 2 (PAX2; 1:500; # AF3364; R&D

Systems); monoclonal rat anti-mCherry (1:2000; #M11217; Thermo Fisher Scientific); polyclonal

rabbit anti-phospho-p44/42 MAPK (ERK1/2) (Thr202/Tyr204) (pERK; 1:500; #9101; Cell

Signaling); anti-isolectin B4 (IB4)-biotin conjugate (1:1000; # I21414; Thermo Fisher Scientific);

monoclonal rabbit anti-c-FOS (1:5000; #226 008; Synaptic systems). After incubation, tissue

sections were washed and incubated for 3 hours at room temperature with secondary antibodies

(Alexa Fluor™ 488, 546 and/or 555; #A11001, A11008, A11035, A11056, A11081, S32351;

Thermo Fisher Scientific: #ab150178; Abcam: DyLight™ 405; #705-475-147; Jackson

ImmunoResearch). Then, tissue sections were washed, slide mounted, and subsequently placed

under coverslips with VECTASHIELD Hardmount (Vector Laboratories). Immunofluorescence

images were obtained with confocal laser microscopy (#LSM700 or LSM900; Carl Zeiss).

Fiber photometry: Immediately after microinjection of AAV vectors encoding GCaMPs, the

mice were implanted with an optic cannula. Cannulae were purchased from RWD (#R-FOC-

BL400C-50NA, ø1.25 mm ferrule, ø400 μm optic fiber, 0.50 NA). The length of the optic fiber

protruding from the tip of the ferrule was 4.0 mm. Optic cannula was implanted in the unilateral

LC (AP: −5.4 mm from the bregma; ML: 0.9 mm; DV: −3.0 mm from the dura). Next, the

cannula was secured using two types of dental glues (Super-bond; Sun Medical, Shiga, Japan:

UNIFAST III; GC, Tokyo, Japan). After implantation, the skin was sutured with 5-0 silk, and the

mice were kept on a heating pad until recovery. We used cannula-implanted mice for fiber

photometry 3 weeks or more after the surgery, and monitored Ca^{2+} events in LC-NA neurons

during restraint stress using the conical tube with a narrow slit (Movie). GCaMP fluorescent

signals were obtained with the fiber photometry apparatus from Doric Lenses, which includes a

fiber photometry console (FPC_V6), a 415-nm LED illumination (CLED_415), a 465-nm LED

illumination (CLED_465), an LED driver (LEDD_2), and a fluorescence mini cube with built-in

fluorescence photodetector amplifier (iFMC4-G2_IE(410-420)_E(460–490)_F(500–550)_S). A

415-nm light was modulated at 208 Hz, while a 465-nm light was modulated at 572 Hz. The

optic cannula implanted in mice and fluorescence mini cube were connected by a low-

autofluorescence patch cable (#MAF1L1; Thorlabs). The power output at the fiber tip was 30–

40 μW. After connecting to the patch cable, the mice were allowed to move freely in the home

cage for at least 10 min for habituation. After habituation, fluorescence recording was started.

Emitted signals from the tissue were collected through the same fiber and sampled at 12 kHz.

Processing and analysis of fiber photometry data: To avoid the effects of photobleaching, data

from the first 5 min of recording were discarded. The raw data were then decimated to 30 Hz

(down-sampled by 400) and low-pass Butterworth-filtered at 2 Hz. The 415-nm excitation

channel served as an isosbestic, Ca^{2+}-independent control wavelength for GCaMP, allowing for

bleaching and movement artifact corrections when directly fitted to the Ca^{2+}-dependent 465-nm

channel. The Correction Baseline function from Doric Neuroscience Studio (Doric Lenses) was

used to fit the 415-nm signal to the 465-nm signal using an adaptive iterative re-weighted

Penalized Least Squares algorithm[82]. The fluorescence change was expressed as a relative

change, $\Delta F/F = (F_{465}-F_{415})/F_{415}$, where F_{465} is the 465 nm-induced Ca^{2+}-dependent signal, and

F_{415} is the 415 nm-induced Ca^{2+}-independent signal. During restraint stress, the Ca^{2+} transients

and escape behaviors (struggling) occurred almost simultaneously (Movie), but the 415 nm-

induced Ca^{2+}-independent signals was unchanged. Robust Z-scores were then calculated as Z =

$(\Delta F/F - \text{Median } \Delta F/F)/\text{NIQR}$, where Median $\Delta F/F$ is the median of $\Delta F/F$ during the entire

measurement period, and NIQR is the normalized interquartile range ($0.7413 \times$ interquartile

range). Ca^{2+} events were detected with a threshold of Z-score >3. The frequency of Ca^{2+} events

was calculated as the number of Ca^{2+} events per minute in the 10 min before stress, 60 min during

stress (10 min $\times$ 6 times), and 10 min after stress.

Drug administration: To ablate DTR$^+$ neurons, DTX (10 µg/kg in PBS; #048-34371; Wako,

Saitama, Japan) was administered i.p. for two successive days 3 weeks after intra-LC injection of

AAV2/9-EF1α-FLEx[DTR-EGFP] or [AcGFP]-WPRE. We used DTX-injected mice for further

analyses 2 weeks or more after the last injection. For intrathecal injection, a 30-G needle attached

to a Hamilton microsyringe was inserted between the L5/L6 vertebrae and then punctured

through the dura. The following drugs and doses were used for intrathecal injection experiments:

silodosin (3 nmol in 5 µl PBS; #191-17591; Wako, Saitama, Japan; injection was given 30 min

before optogenetic stimulation); L-norepinephrine hydrochloride (0.1 nmol in 5 µl saline;

#74480; Sigma-Aldrich); 5,7-dichlorokynurenic acid (DCK; 10 nmol in 5 µl saline of 2%

dimethylsulfoxide (DMSO); #0286; Tocris; injection was given immediately before the start of

exposure to restraint stress); (R)-(-)-phenylephrine hydrochloride (Phe; 0.05 nmol in 5 µl saline;

#163-11791; Wako, Saitama, Japan); 8-cyclopentyl-1,3-dimethylxanthine (CPT; 3 nmol in 5 µl

saline of 2% DMSO; #C102; Sigma-Aldrich; injection was given immediately before the start of

exposure to restraint stress or 30 min before intrathecal injection of NA).

In vivo **optogenetic stimulation:** Immediately after microinjection of AAV vectors encoding

ChrimsonR, mice were implanted with an optic cannula. A cannula comprised a ceramic ferrule

(#CFLC230-10; Thorlabs; ø1.25 mm, 6.4 mm length) and an optic fiber (#FT200UMT; Thorlabs;

ø200 µm, 0.39 NA). The length of the optic fiber protruding from the tip of the ferrule was less

than 0.3 mm. The vertebrae were immobilized in place using a metal bar along the Th12–L1. A

small laminectomy was performed through the Th13 bone using a dental drill. The cannula was

secured above the dura mater using two types of dental glues (Super-bond; Sun Medical:

UNIFAST III; GC). In this procedure, the dura mater and spinal cord parenchyma were left intact

and unscathed. After implantation, the skin was sutured with 5-0 silk, and the mice were kept on a

heating pad until recovery. We used cannula-implanted mice for optogenetic stimulation 3 weeks

or more after the surgery. To activate ChrimsonR in LC$^{\rightarrow SDH}$-NA axons/terminals, 625-nm LED

light (#M625F2; Thorlabs) was delivered through a ferrule patch cable (#M83L01; Thorlabs).

Light stimulations (2 mW, 10 Hz, 5-ms pulse duration, 5-s light on, 15-s light off, 10 cycles) were

generated by an LED driver (#LEDD_2; Doric Lenses).

Fluorescent Ca^{2+} and NA imaging in spinal cord slices: Fluorescent imaging was performed in

accordance with our previously described method[22]. GCaMP6m- and GRAB$_{NE1m}$-expressing mice

were anesthetized with an i.p. injection of urethane (1.2–1.5 mg/kg), and lumbosacral

laminectomy was performed. The spinal cord (L1–S2) was removed and placed in a cold, high-

sucrose, artificial cerebrospinal (aCSF) fluid (27 mM NaHCO$_3$, 1.4 mM NaH$_2$PO$_4$, 2.5 mM KCl,

7.0 mM MgSO$_4$, 1.0 mM CaCl$_2$, 222 mM sucrose, and 0.5 mM ascorbic acid), which was bubbled

with 95% O$_2$ and 5% CO$_2$. After cutting all ventral and dorsal roots, transverse L4 spinal cord

slices (350 µm thick) were made using a vibrating microtome (#VT1200; Leica); the slices were

kept in oxygenated aCSF solution (125 mM NaCl, 1.25 mM NaH$_2$PO$_4$, 2.5 mM KCl, 1.0 mM

MgCl$_2$, 2.0 mM CaCl$_2$, 26 mM NaHCO$_3$, and 20 mM glucose) at room temperature (22–25°C) for

at least 30 min before use. The slices were placed in a recording chamber, and fluorescent signals

were measured using a confocal laser scanning microscope (#FV3000; Olympus, Tokyo, Japan).

The chamber was perfused with aCSF saturated with 95% O$_2$ and 5% CO$_2$ at 31–32°C and at 2–3

ml/min using peristaltic pumps. A 488-nm diode laser was used for the excitation of GCaMP6m

and GRAB$_{NE1m}$. The size of the image was 509×509 µm^2 (512×512 pixels). The recordings

were acquired at a rate of 1.083 s per image. Light stimulations (1 mW, 10 Hz, 5-ms pulse

duration, 1–20 s) were generated by an LED driver (#LEDD_2; Doric Lenses) and delivered

through an optic fiber (Thorlabs, FT200UMT, ø200 µm, 0.39 NA). For an experiment on α$_{1A}$R

inhibition, optogenetic stimulations (10 s) were performed twice (at least 15-min apart) in the

same slice. Silodosin (40 nM in aCSF) was continuously applied by bath application starting 5

min before the onset of 2nd optogenetic stimulation. After optogenetic stimulation, NA (1 or 10

µM in aCSF) was applied by bath application as the positive control.

Processing and analysis of Ca^{2+} and NA imaging data: Videos were imported into ImageJ Fiji

(https://imagej.net/software/fiji/), and motion artifacts were corrected using TurboReg[83]. The

fluorescence change was expressed as a relative percentage change, $\Delta F/F = 100 \times (F_t - F_0)/F_0$,

where F_t is the fluorescence at time t, and F_0 is the baseline average for 30 frames before the start

of stimulation. For Ca^{2+} imaging using GCaMP, the size of each region of interest (ROI) was set

to a circle with a diameter of 10 µm[22]. ROIs were manually identified based on the following

criteria: located within 200 μm from the surface of gray matter and ≥30% $\Delta F/F$ fluorescence

change after optogenetic stimulation or NA application. For the experiment on $\alpha_{1A}R$ inhibition,

normalized $\Delta F/F$ was calculated as ($\Delta F/F$ in 2nd stimulation)/($\Delta F/F$ in 1st stimulation). For NA

imaging using $GRAB_{NE1m}$, the raw image was binarized, and a single ROI was detected from such

a binarized image with the following parameters: contiguous pixel number >100 and

0□<□circularity□<□1[73]. Calculation of $\Delta F/F$ in ROIs was performed using the Time Series

Analyzer V3 plugin in ImageJ.

Whole-cell patch-clamp recordings using spinal cord slices: According to our previously

described method[16], mice were deeply anesthetized with urethane (1.2–1.5 mg/kg, i.p.), and the

lumbar spinal cord was removed and placed in a cold high-sucrose aCSF (250 mM sucrose, 2.5

mM KCl, 2 mM $CaCl_2$, 2 mM $MgCl_2$, 1.2 mM NaH_2PO_4, 25 mM $NaHCO_3$, and 11 mM glucose).

Parasagittal and transverse spinal cord slices (250–300 μm thick) were made with a vibrating

microtome (VT1200, Leica) and then the slices kept in oxygenated aCSF solution (125 mM

NaCl, 2.5 mM KCl, 2 mM $CaCl_2$, 1 mM $MgCl_2$, 1.25 mM NaH_2PO_4, 26 mM $NaHCO_3$, and 20

mM glucose) at room temperature (22–25°C) for at least 30 min. The spinal cord slice was then

put into a recording chamber, where it was continuously superfused with aCSF solution at 25–

28°C at a flow rate of 4–6 ml/min. Recordings were made with the Axopatch 700B amplifier and

pCLAMP 10.4 acquisition software (Molecular Devices). Data were digitized with an analog-to-

digital converter (Digidata 1550; Molecular Devices), stored on a personal computer with a data

acquisition program (ClampeX version 10.4; Molecular Devices), and analyzed with a software

package (Clampfit version 10.7; Molecular Devices). Spontaneous inhibitory postsynaptic

currents (sIPSCs) were recorded in the voltage-clamp mode at a holding potential of 0 mV. Patch

pipettes were filled with an internal solution (120 mM $CsMeSO_4$, 15 mM CsCl, 10 mM HEPES,

5 mM QX-314, 4 mM MgATP, 0.3 mM Na_2GTP, 0.2 mM EGTA, 10 mM TEA-Cl and 8 mM

NaCl [pH 7.28] adjusted with CsOH), and whole-cell patch-clamp recordings were made from

SG neurons. Resting membrane potentials (RMPs) were recorded in current-clamp mode. Patch

pipettes were filled with an internal solution (125 mM K-gluconate, 10 mM KCl, 0.5 mM EGTA,

10 mM HEPES, 4 mM ATP-Mg, 0.3 mM NaGTP, 10 mM phosphocreatine, pH 7.28 adjusted

with KOH), and whole-cell patch-clamp recordings were made from tdTomato[+] or mCherry[+]

neurons. The following drugs were used: NA (20 μM; #74480; Sigma-Aldrich), clozapine-N-

oxide (CNO; 100 μM; #BML-NS105; Enzo Life Sciences), CPT (1 μM), N6-

cyclopentyladenosine (CPA; 1 μM; #119135; Sigma-Aldrich), CNQX disodium salt hydrate

(CNQX; 10 μM; #C239; Sigma-Aldrich), (+)-MK-801 hydrogen maleate (MK-801; 20 μM;

#M107; Sigma-Aldrich), 1(S), 9(R)-(-)-bicuculline methiodide (10 µM; #14340; Sigma-Aldrich)

and strychnine (1 µM; #S0532; Sigma-Aldrich). All drugs were dissolved in aCSF solution. CNO

was superfused for 3 min. Bath application of CPT was continuously superfused from 4 min

before CNO application. NA was superfused with an antagonist cocktail (CNQX, MK-801,

bicuculline and strychnine) for 3–5 min to block synaptic inputs from other neurons to the

recorded Vgat$^+$ neurons[48,49]. If the recording neuron was depolarized by NA perfusion, then CPA

was co-perfused with NA and an antagonist cocktail for 3 min. The frequency of sIPSCs for 1

min of pre- and post-NA application were quantified using Clampfit version 10.7 (Molecular

Devices). We quantified averaged RMP for 1 min of pre- and post-drug application, and a change

in RMP (ΔRMP) of 5 mV or more was judged to be depolarization or hyperpolarization.

***In situ* hybridization:** Mice were deeply anesthetized with an i.p. injection of pentobarbital and

transcardially perfused with PBS followed by ice-cold 4% PFA/PBS. The L4 spinal cord was

removed and postfixed in the same fixative 24 hours at 4°C. After that, tissues were incubated

with 10%, 20% and 30% sucrose solutions at 4°C, embedded in OCT compound, and stored at

–25°C before use. The L4 segments were sectioned at a thickness of 14 µm. *In situ* hybridization

was performed using RNAscope® Multiplex Fluorescent Reagent Kit v2 (#323100; ACDbio),

according to the manufacture's protocol for fixed frozen tissue. The following probes were used:

Mm-Adora1 (#402261; ACDbio); Mm-Slc32a1-C3 (#319191-C3; ACDbio). Tissue sections were

analyzed using an LSM700 Imaging System. Cells were considered positive if three or more

punctate dots were present in the nucleus and/or cytoplasm[14].

Electrical stimulation of Aβ fibers and counting of c-FOS$^+$ and pERK$^+$ neurons in the SDH:

For pERK immunofluorescence, but not for c-FOS, after exposure to 1-hour restraint stress, mice

were deeply anesthetized with an i.p. injection of urethane (1.2–1.5 mg/kg). Thirty min after

stress, Aβ fiber stimulation was performed. The electrodes were attached to the plantar and dorsal

surfaces of the left hindpaw. Transcutaneous nerve stimuli were applied using a stimulus

generator (#STG4002-16□mA; Multi Channel Systems). The current intensity of the 2,000-Hz

stimuli was 1,000□µA[57]. Mice were fixed at 2 (for pERK) or 90□min (for c-FOS) after electrical

stimulation. For c-FOS immunostaining, mice were deeply anesthetized with an i.p. injection of

pentobarbital prior to fixation. For quantification of c-FOS$^+$ and pERK$^+$ cells, three sections from

the L4 spinal cord segments were randomly selected from each mouse, and the number of c-FOS$^+$

and pERK$^+$ neurons in the superficial laminae I–IIi (defined by staining of IB4) was counted.

Consistent with a previous study[57], we confirmed that electrical stimulation of Aδ and C fibers,

but not Aβ fibers, induced ERK phosphorylation in SDH neurons (data not shown).

Statistical analysis: Statistical analyses were performed using Prism (GraphPad). Quantitative

data were expressed as the mean ± SEM. Statistical significance of differences was determined by

using two-tailed paired *t*-test (Figure S1, S2E and S5), Mann-Whitney test (Figure 2F and S4B),

Wilcoxon signed-rank test (Figure 3L, 3N and S7C), Fisher's exact test (Figure S8), Friedman

test with Dunn's multiple comparisons test (Figure 1E and 2E), one-way ANOVA with Dunnett's

multiple comparisons test (Figure 2C), one-way ANOVA with Tukey's multiple comparisons test

(Figure 4F and 4H) and repeated measures two-way ANOVA with Bonferroni's multiple

comparisons test (Figure 1B, 1H, 1K, 1N, 2G, 2H, 2I, 2J, 3A, 3B, 3C, 4A, 4B, 4C, 4D, S2E and

S5), as appropriate, after determining the normality (Kolmogorov–Smirnov test or Shapiro–Wilk

test). P values are indicated as $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$.

Acknowledgments: We would like to thank Editage for editing a draft of this manuscript, and the
University of Pennsylvania vector core for providing pZac2.1, pAAV2/5, pAAV2/9 and pAd
DeltaF6 plasmid. Some figure elements were created with BioRender.com. This work was
supported by Japan Society for the Promotion of Science (JSPS) KAKENHI Grants JP19H05658
(M.T.), JP20H05900 (M.T.) and JP24H00067 (M.T.), by the Core Research for Evolutional
Science and Technology (CREST) program from AMED under Grant Number 25gm1510013h
(M.T.), and by Research Support Project for Life Science and Drug Discovery (Basis for
Supporting Innovative Drug Discovery and Life Science Research (BINDS)) from AMED under
Grant Number JP25ama121031 (M.T.). R.K-K., S.U., and K.Y. were JSPS research fellows
(JP22KJ2471, JP23KJ1729, and JP19J21063, respectively).

Author contributions:
Conceptualization: MT
Formal analysis: RKK, SU, KY
Funding acquisition: RKK, SU, KY, MT
Investigation: RKK, SU, KY, KK, DK
Methodology: RKK, SU, KY, KK, KM, KFT
Project administration: MT
Resources: TM, IH
Supervision: MT
Visualization: RKK, SU, MT
Writing–original draft: RKK, SU, KY, MT
Writing–review and editing: RKK, SU, MT

Declaration of interests: The authors declare no competing interests.

Data and materials availability: All data associated with this study are present in the paper or
the Supplemental Information.

Figures and figure legends

Figure 1. LC$^{\rightarrow SDH}$-NA neurons mediate stress-induced mechanical hypersensitivity.

(A) Schematic illustration of an experiment to investigate the effects of acute exposure to restraint stress (1

hour) on mechanosensory behavior in mice, using von Frey (vF) filaments. (B) Change in paw withdrawal

threshold (PWT) measured by vF filaments in wild-type mice after restraint stress ($n = 6$ mice per group;

two-way ANOVA with post hoc Bonferroni's multiple comparisons test; *$P < 0.05$, **$P < 0.01$, ***$P <$

0.001, ****P <0.0001 vs. no-restraint stress group). (C) Expression of GCaMP6s (GC; green) in the LC at

3 weeks after intra-LC injection of AAV-FLEx[GCaMP6s] in *NET-Cre* mice. TH immunofluorescence is

shown in magenta. Dashed line indicates the location of the implanted optic fiber. Scale bar, 100 μm. (D

and E) Representative traces and change in the frequency of GCaMP6s signals in LC-NA neurons ($n = 6$

mice; Friedman test with post hoc Dunn's multiple comparisons test; **$P < 0.01$ vs. the data of 'Before').

Traces shown at the top, middle, and bottom (D) indicate Ca^{2+} signals before, during, and after restraint

stress, respectively. (F) Schematic illustration of the strategy of ablating LC-NA neurons using AAV

vectors incorporating DTR (fused with EGFP) injected into the LC in *NET-Cre* mice. (G) TH

immunofluorescence (magenta) and GFP (green) in the LC (left) and NET immunofluorescence (magenta)

in the SDH (right) after administration of DTX (10 μg/kg, i.p., two injections 24 h-apart) in control mice

(top) and DTR-expressing mice (bottom). Scale bars, 100 μm. (H) Effect of ablation of LC-NA neurons on

PWT changes after acute restraint stress ($n = 7$ mice per group; two-way ANOVA with post hoc

Bonferroni's multiple comparisons test; **$P < 0.01$ vs. control group). (I) Schematic illustration of the

strategy of ablating LC$^{\rightarrow SDH}$-NA neurons using a retrograde AAV vector incorporating Cre injected into the

SDH and an AAV vector incorporating DTR (fused with EGFP) injected into the LC in wild-type mice. (J)

Representative images of LC$^{\rightarrow SDH}$-NA neurons in control or DTR-expressing mice treated with vehicle or

DTX administration, respectively. GFP (green) and TH (magenta). Scale bar, 100 μm. (K) Effect of

ablation of LC$^{\rightarrow SDH}$-NA neurons on PWT changes after restraint stress ($n = 11$ mice per group; two-way

ANOVA with post hoc Bonferroni's multiple comparisons test; **$P < 0.01$ vs. control group). (L)

Schematic illustration of the strategy for activating LC$^{\rightarrow SDH}$-NA neuronal axons/terminals using an AAV

vector incorporating ChrimsonR (fused with tdTomato) injected into the LC in *NET-Cre* mice and of an

optic cannula implanted in the SDH. (M) Representative images of TH (green) and tdTomato (magenta)

expression in the LC (top) and NET (green) and tdTomato (magenta) expression in the SDH (bottom) at 3

weeks after intra-LC injection of AAV-FLEx[ChrimsonR-tdTomato] in *NET-Cre* mice. Scale bars, 100 μm

(top) and 50 μm (bottom). (N) PWT before and after optogenetic stimulation (opto-stim.) in LC$^{\rightarrow SDH}$-NA

axons/terminals (625 nm, 2 mW, 10 Hz, 5-ms pulse duration, 5-s light on, 15-s light off, 10 cycles)

(Control, $n = 4$ mice; ChrimsonR, $n = 5$ mice; two-way ANOVA with post hoc Bonferroni's multiple

comparisons test; ****P <0.0001 vs. control group). Data represent mean ± SEM. See also Figures S1–S6.

Figure 2. α_{1A}Rs in Hes5$^+$ SDH astrocytes are required for stress-induced mechanical hypersensitivity.
(A) Schematic illustration of intra-SDH microinjection of AAV-gfaABC$_1$D-GRAB$_{NE1m}$ or -GCaMP6m and
intra-LC microinjection of AAV-FLEx[ChrimsonR-tdTomato] in *NET-Cre* mice. (B) Representative traces
of GRAB$_{NE1m}$ signals by fluorescence imaging using spinal cord slices. Each trace represents the
GRAB$_{NE1m}$ signal before and after optogenetic stimulation (625 nm, 1 mW, 10 Hz, 5 ms pulse duration, 1–
20 s). (C) Quantitative analysis of the peak amplitude of GRAB$_{NE1m}$ $\Delta F/F$ after optogenetic stimulation in
LC$^{\rightarrow SDH}$-NA axons/terminals (n = 4 slices; one-way ANOVA with post hoc Dunnett's multiple
comparisons test; **P < 0.01, ****P <0.0001). (D) Representative traces of astrocytic GCaMP6m signals
by fluorescence imaging using spinal cord slices. Each trace represents the GCaMP6m signal before and
after optogenetic stimulation (as described in B). (E) Quantitative analysis of the peak amplitude of
GCaMP6m $\Delta F/F$ after optogenetic stimulation in LC$^{\rightarrow SDH}$-NA axons/terminals (n = 133 cells, 4 slices, 4
mice; Friedman test with post hoc Dunn's multiple comparisons test; ****P <0.0001). (F) Effect of
silodosin (40 nM) on astrocytic Ca^{2+} responses in the SDH after optogenetic stimulation (10 s) in LC$^{\rightarrow SDH}$-
NA axons/terminals (Control, n = 83 cells, 4 slices, 4 mice; Silodosin, n = 53 cells, 4 slices, 4 mice; Mann-
Whitney test; ****P <0.0001). (G) Effect of intrathecal silodosin (3 nmol) on mechanical hypersensitivity
induced by optogenetic stimulation in LC$^{\rightarrow SDH}$-NA axons/terminals (Vehicle, n = 5 mice; Silodosin, n = 6
mice; two-way ANOVA with post hoc Bonferroni's multiple comparisons test; *P < 0.05, ***P < 0.001 vs.
vehicle group). (H) Change in PWT at 30 min after intrathecal injection of NA (0.1 nmol) in control
(*Adra1a*$^{flox/flox}$) and Hes5$^+$ astrocyte-selective α_{1A}R conditional knockout mice [*Hes5-
CreERT2;Adra1a*$^{flox/flox}$ mice treated with tamoxifen (TAM) (Hes5$^+$ astrocyte–α_{1A}R cKO mice) (n = 6 mice
per group; two-way ANOVA with post hoc Bonferroni's multiple comparisons test; ****P <0.0001). (I and
J) Stress-induced mechanical hypersensitivity in Hes5$^+$ astrocyte–α_{1A}R cKO mice [I: Control

801   ($Adra1a^{flox/flox}$), $n = 7$ mice; Hes5$^+$ astrocyte–α_{1A}R cKO, $n = 8$ mice] or wild-type mice with intrathecal

802   DCK (10 nmol) (J: $n = 5$ mice per group) (two-way ANOVA with post hoc Bonferroni's multiple

803   comparisons test; **$P < 0.01$, ***$P < 0.001$ vs. control or vehicle group). Data represent mean $\pm$ SEM.

Figure 3. Activation of Hes5$^+$ astrocytes reduces activity in Vgat$^+$ INs in the SDH.

(A and B) Intrathecal NA (A, 0.1 nmol) or Phe (B, 0.05 nmol)-induced mechanical hypersensitivity in

control (*Adra1a*$^{\text{flox/flox}}$) and Vgat$^+$ IN-selective α_{1A}R conditional knockout mice [*Vgat-Cre*;*Adra1a*$^{\text{flox/flox}}$

mice (Vgat$^+$ INs–α_{1A}R cKO mice)] (n = 6 mice per group; two-way ANOVA with post hoc Bonferroni's

multiple comparisons test; n.s., not significant). (C) Changes in PWT after acute exposure to restraint stress

in control (*Adra1a*$^{\text{flox/flox}}$; *n* = 6 mice) and Vgat$^+$ INs–α_{1A}R cKO mice (*n* = 9 mice) (two-way ANOVA with

post hoc Bonferroni's multiple comparisons test). (D) Representative trace of membrane potentials in

tdTomato$^+$ (Vgat$^+$) SDH neurons after application of NA (20 μM) to spinal cord slices from *Vgat-*

Cre;*Rosa-tdTomato* mice. A$_1$R agonist CPA (1 μM) was co-applied with NA. (E) Percentage of Vgat$^+$ SDH

neurons whose NA-evoked response was inhibited by CPA (*n* = 10 cells from 7 mice). (F) Representative

images of *Adora1* (A$_1$R) mRNA expression (green) in *Slc32a1* (Vgat)$^+$ INs (magenta). DAPI staining is

shown in gray. Arrowheads indicate A$_1$R-expressing Vgat$^+$ cells. Scale bar, 25 μm. (G) SaCas9 (yellow,

detected by HA-tag) and mCherry (magenta) expression in the PAX2$^+$ INs (cyan) at 3 weeks after intra-

SDH injection of AAV-FLEx[SaCas9] and AAV-FLEx[mCherry]-U6-sgAdora1 in *Vgat-Cre* mice.

Arrowheads indicate genome-editing Vgat$^+$ cells. Scale bar, 25 μm. (H) Representative traces of membrane

potentials in Vgat$^+$ INs after application of NA and CPA to spinal cord slices from *Vgat-Cre* mice with

conditional knockdown of A$_1$Rs in *Vgat$^+$* INs (SDH-Vgat$^+$ IN–A$_1$R cKD) and their controls (Control; *Vgat-*

Cre mice with intra-SDH injection of AAV-FLEx[mCherry]). (I) Percentage of mCherry$^+$ (Vgat$^+$) SDH

neurons whose NA-evoked response was inhibited by CPA (Control, $n = 10$ cells from 8 mice; SDH-Vgat$^+$

IN–A$_1$R cKD, $n = 8$ cells from 5 mice). (J) Expression of hM3Dq (green, detected by HA-tag) in the SDH

at 3 weeks after intra-SDH injection of AAV-FLEx[hM3Dq] in *Hes5-CreERT2* mice treated with TAM.

Dashed line indicates an outline of the gray matter of SDH. GFAP (magenta, bottom left), SOX9 (magenta,

bottom right). Arrowheads indicate HA-tag$^+$ astrocytes. Scale bars, 200 μm (top) and 25 μm (bottom). (K

and L) Representative traces of sIPSCs (K) and quantitative analysis of their frequency (L) in SG neurons

in spinal cord slices from *Hes5-CreERT2*;AAV-hM3Dq mice treated with TAM [Pre and CNO: before and

after bath application of CNO (100 μM), respectively] ($n = 7$ cells from 7 mice; Wilcoxon signed-rank test;

*$P < 0.05$). (M and N) Representative traces of sIPSCs (M) and quantitative analysis of their frequency (N)

in SG neurons in spinal cord slices from *Hes5-CreERT2*;AAV-FLEx[hM3Dq] mice treated with TAM [Pre

and CNO: before and after bath application of CNO with CPT (1 μM), respectively] ($n = 13$ cells from 13

mice; Wilcoxon signed-rank test; n.s., not significant). Data represent mean ± SEM. See also Figure S7 and

S8.

Figure 4. Hes5[+] astrocyte-mediated inhibitory signals to SDH-Vgat[+] INs contribute to stress-induced

mechanical hypersensitivity.

(A and B) PWT before and 30 min after intrathecal administration of NA (0.1 nmol) in wild-type mice

pretreated intrathecally with vehicle or CPT (3 nmol) (A: $n = 5$ mice per group) or in control (*Vgat-Cre*

mice with intra-SDH of AAV-FLEx[mCherry]) and SDH-Vgat[+] IN–A_1R cKD mice (B: $n = 9$ mice per

group) (two-way ANOVA with post hoc Bonferroni's multiple comparisons test; ***$P < 0.001$). (C and D)

PWT before and after acute restraint stress in wild-type mice pretreated intrathecally with vehicle or CPT

(C: Vehicle, $n = 6$ mice; CPT, $n = 5$ mice) or in control (*Vgat-Cre* mice with intra-SDH of AAV-

FLEx[mCherry]) and SDH-Vgat$^+$ IN–A$_1$R cKD mice (D: Control, $n = 6$ mice; SDH-Vgat$^+$ IN–A$_1$R cKD, n

= 7 mice) (two-way ANOVA with post hoc Bonferroni's multiple comparisons test; *$P < 0.05$, **$P < 0.01$,

***$P < 0.001$, ****$P < 0.0001$ vs. vehicle or control group). (E–H) Representative images of c-FOS (green,

E), pERK (green, G) and IB4 (magenta, E and G) immunofluorescence in the SDH with or without Aβ

fiber stimulation and/or restraint stress. CPT was intrathecally administered 30 min before stress exposure.

Quantitative analysis of the number of c-FOS$^+$ (F) and pERK$^+$ (H) cells in superficial laminae of the SDH

in each group ($n = 4$–5 mice per group; one-way ANOVA with post hoc Tukey's multiple comparisons test;

*$P < 0.05$, ***$P < 0.001$, ****$P < 0.0001$). (I) Schematic illustration of the mechanisms of stress-induced

mechanical pain facilitation highlighting NA signals from LC$^{\rightarrow SDH}$-NAergic terminals to Hes5$^+$ astrocytes

and Vgat$^+$ INs. Data represent mean ± SEM.

References

1   Todd, A. J. Neuronal circuitry for pain processing in the dorsal horn. *Nat Rev Neurosci*
11, 823-836, doi:10.1038/nrn2947 (2010).
2   Peirs, C. & Seal, R. P. Neural circuits for pain: Recent advances and current views.
Science **354**, 578-584, doi:10.1126/science.aaf8933 (2016).
3   Koch, S. C., Acton, D. & Goulding, M. Spinal Circuits for Touch, Pain, and Itch. *Annu*
Rev Physiol **80**, 189-217, doi:10.1146/annurev-physiol-022516-034303 (2018).
4   Moehring, F., Halder, P., Seal, R. P. & Stucky, C. L. Uncovering the Cells and Circuits of
Touch in Normal and Pathological Settings. *Neuron* **100**, 349-360,
doi:10.1016/j.neuron.2018.10.019 (2018).
5   Peirs, C., Dallel, R. & Todd, A. J. Recent advances in our understanding of the
organization of dorsal horn neuron populations and their contribution to cutaneous
mechanical allodynia. *J Neural Transm (Vienna)* **127**, 505-525, doi:10.1007/s00702-020-
02159-1 (2020).
6   Wang, L. H., Ding, W. Q. & Sun, Y. G. Spinal ascending pathways for somatosensory
information processing. *Trends Neurosci* **45**, 594-607, doi:10.1016/j.tins.2022.05.005
(2022).
7   Mercer Lindsay, N., Chen, C., Gilam, G., Mackey, S. & Scherrer, G. Brain circuits for
pain and its treatment. *Sci Transl Med* **13**, eabj7360, doi:10.1126/scitranslmed.abj7360
(2021).
8   Nguyen, E., Grajales-Reyes, J. G., Gereau, R. W. t. & Ross, S. E. Cell type-specific
dissection of sensory pathways involved in descending modulation. *Trends Neurosci* **46**,
539-550, doi:10.1016/j.tins.2023.04.002 (2023).
9   Yoshimura, M. & Furue, H. Mechanisms for the anti-nociceptive actions of the
descending noradrenergic and serotonergic systems in the spinal cord. *J Pharmacol Sci*
101, 107-117, doi:10.1254/jphs.crj06008x (2006).

10 Llorca-Torralba, M., Borges, G., Neto, F., Mico, J. A. & Berrocoso, E. Noradrenergic
Locus Coeruleus pathways in pain modulation. *Neuroscience* **338**, 93-113,
doi:10.1016/j.neuroscience.2016.05.057 (2016).
11 Suárez-Pereira, I. *et al.* The Role of the Locus Coeruleus in Pain and Associated Stress-
Related Disorders. *Biol Psychiatry* **91**, 786-797, doi:10.1016/j.biopsych.2021.11.023
(2022).
12 Li, J. *et al.* Activation of locus coeruleus-spinal cord noradrenergic neurons alleviates
neuropathic pain in mice via reducing neuroinflammation from astrocytes and microglia
in spinal dorsal horn. *J Neuroinflammation* **19**, 123, doi:10.1186/s12974-022-02489-9
(2022).
13 Häring, M. *et al.* Neuronal atlas of the dorsal horn defines its architecture and links
sensory input to transcriptional cell types. *Nat Neurosci* **21**, 869-880,
doi:10.1038/s41593-018-0141-1 (2018).
14 Shiraishi, Y. *et al.* α1A-adrenaline receptors in dorsal horn inhibitory neurons have an
inhibitory role in the regulation of chloroquine-induced itch in mice. *Mol Brain* **14**, 55,
doi:10.1186/s13041-021-00768-9 (2021).
15 Baba, H. *et al.* Norepinephrine facilitates inhibitory transmission in substantia gelatinosa
of adult rat spinal cord (part 2): effects on somatodendritic sites of GABAergic neurons.
Anesthesiology **92**, 485-492, doi:10.1097/00000542-200002000-00031 (2000).
16 Uchiyama, S. *et al.* Stress-induced antinociception to noxious heat requires α1A-
adrenaline receptors of spinal inhibitory neurons in mice. *Mol Brain* **15**, 6,
doi:10.1186/s13041-021-00895-3 (2022).
17 Inoue, K. & Tsuda, M. Microglia in neuropathic pain: cellular and molecular mechanisms
and therapeutic potential. *Nat Rev Neurosci* **19**, 138-152, doi:10.1038/nrn.2018.2 (2018).
18 Ji, R. R., Donnelly, C. R. & Nedergaard, M. Astrocytes in chronic pain and itch. *Nat Rev
Neurosci* **20**, 667-685, doi:10.1038/s41583-019-0218-1 (2019).
19 Tsuda, M., Masuda, T. & Kohno, K. Microglial diversity in neuropathic pain. *Trends
Neurosci* **46**, 597-610, doi:10.1016/j.tins.2023.05.001 (2023).
20 Sekiguchi, K. J. *et al.* Imaging large-scale cellular activity in spinal cord of freely
behaving mice. *Nat Commun* **7**, 11450, doi:10.1038/ncomms11450 (2016).
21 Yoshihara, K. *et al.* Astrocytic Ca(2+) responses in the spinal dorsal horn by noxious
stimuli to the skin. *J Pharmacol Sci* **137**, 101-104, doi:10.1016/j.jphs.2018.04.007
(2018).
22 Kohro, Y. *et al.* Spinal astrocytes in superficial laminae gate brainstem descending
control of mechanosensory hypersensitivity. *Nat Neurosci* **23**, 1376-1387,
doi:10.1038/s41593-020-00713-4 (2020).
23 Kawanabe, R., Yoshihara, K., Hatada, I. & Tsuda, M. Activation of spinal dorsal horn
astrocytes by noxious stimuli involves descending noradrenergic signaling. *Mol Brain* **14**,
79, doi:10.1186/s13041-021-00788-5 (2021).
24 Hertz, L., Lovatt, D., Goldman, S. A. & Nedergaard, M. Adrenoceptors in brain: cellular
gene expression and effects on astrocytic metabolism and [Ca(2+)]i. *Neurochem Int* **57**,
411-420, doi:10.1016/j.neuint.2010.03.019 (2010).
25 Bazargani, N. & Attwell, D. Astrocyte calcium signaling: the third wave. *Nat Neurosci*
19, 182-189, doi:10.1038/nn.4201 (2016).
26 Batiuk, M. Y. *et al.* Identification of region-specific astrocyte subtypes at single cell
resolution. *Nat Commun* **11**, 1220, doi:10.1038/s41467-019-14198-8 (2020).
27 Wahis, J. *et al.* The astrocyte α1A-adrenoreceptor is a key component of the
neuromodulatory system in mouse visual cortex. *Glia*, doi:10.1002/glia.24591 (2024).

28 Cohen, Z., Molinatti, G. & Hamel, E. Astroglial and vascular interactions of
noradrenaline terminals in the rat cerebral cortex. *J Cereb Blood Flow Metab* **17**, 894-
904, doi:10.1097/00004647-199708000-00008 (1997).
29 Valentino, R. J. & Van Bockstaele, E. Convergent regulation of locus coeruleus activity
as an adaptive response to stress. *Eur J Pharmacol* **583**, 194-203,
doi:10.1016/j.ejphar.2007.11.062 (2008).
30 Imbe, H. *et al.* Activation of ERK in the locus coeruleus following acute noxious
stimulation. *Brain Res* **1263**, 50-57, doi:10.1016/j.brainres.2009.01.052 (2009).
31 Chen, F. J. & Sara, S. J. Locus coeruleus activation by foot shock or electrical stimulation
inhibits amygdala neurons. *Neuroscience* **144**, 472-481,
doi:10.1016/j.neuroscience.2006.09.037 (2007).
32 McCall, J. G. *et al.* CRH Engagement of the Locus Coeruleus Noradrenergic System
Mediates Stress-Induced Anxiety. *Neuron* **87**, 605-620,
doi:10.1016/j.neuron.2015.07.002 (2015).
33 Butler, R. K. & Finn, D. P. Stress-induced analgesia. *Prog Neurobiol* **88**, 184-202,
doi:10.1016/j.pneurobio.2009.04.003 (2009).
34 Jennings, E. M., Okine, B. N., Roche, M. & Finn, D. P. Stress-induced hyperalgesia.
Prog Neurobiol **121**, 1-18, doi:10.1016/j.pneurobio.2014.06.003 (2014).
35 François, A. *et al.* A Brainstem-Spinal Cord Inhibitory Circuit for Mechanical Pain
Modulation by GABA and Enkephalins. *Neuron* **93**, 822-839.e826,
doi:10.1016/j.neuron.2017.01.008 (2017).
36 Abdallah, C. G. & Geha, P. Chronic Pain and Chronic Stress: Two Sides of the Same
Coin? *Chronic Stress (Thousand Oaks)* **1**, 2470547017704763,
doi:10.1177/2470547017704763 (2017).
37 Fitzcharles, M. A. *et al.* Nociplastic pain: towards an understanding of prevalent pain
conditions. *Lancet* **397**, 2098-2110, doi:10.1016/s0140-6736(21)00392-5 (2021).
38 Sousa, F. S. S. *et al.* α-(phenylselanyl) acetophenone abolishes acute restraint stress
induced-comorbid pain, depression and anxiety-related behaviors in mice. *Neurochem Int*
120, 112-120, doi:10.1016/j.neuint.2018.08.006 (2018).
39 Contet, C. *et al.* Dissociation of analgesic and hormonal responses to forced swim stress
using opioid receptor knockout mice. *Neuropsychopharmacology* **31**, 1733-1744,
doi:10.1038/sj.npp.1300934 (2006).
40 Kim, J. W. *et al.* Social support rescues acute stress-induced cognitive impairments by
modulating ERK1/2 phosphorylation in adolescent mice. *Sci Rep* **8**, 12003,
doi:10.1038/s41598-018-30524-4 (2018).
41 Wagatsuma, A. *et al.* Locus coeruleus input to hippocampal CA3 drives single-trial
learning of a novel context. *Proc Natl Acad Sci U S A* **115**, E310-E316,
doi:10.1073/pnas.1714082115 (2018).
42 Mulvey, B. *et al.* Molecular and Functional Sex Differences of Noradrenergic Neurons in
the Mouse Locus Coeruleus. *Cell Rep* **23**, 2225-2235, doi:10.1016/j.celrep.2018.04.054
(2018).
43 Saito, M. *et al.* Diphtheria toxin receptor-mediated conditional and targeted cell ablation
in transgenic mice. *Nat Biotechnol* **19**, 746-750, doi:10.1038/90795 (2001).
44 Klapoetke, N. C. *et al.* Independent optical excitation of distinct neural populations. *Nat*
Methods **11**, 338-346, doi:10.1038/nmeth.2836 (2014).
45 Feng, J. *et al.* A Genetically Encoded Fluorescent Sensor for Rapid and Specific In Vivo
Detection of Norepinephrine. *Neuron* **102**, 745-761.e748,
doi:10.1016/j.neuron.2019.02.037 (2019).

46   Lee, Y., Messing, A., Su, M. & Brenner, M. GFAP promoter elements required for
region-specific and astrocyte-specific expression. *Glia* **56**, 481-493,
doi:10.1002/glia.20622 (2008).
47   Nam, Y. *et al.* Reversible Induction of Pain Hypersensitivity following Optogenetic
Stimulation of Spinal Astrocytes. *Cell Rep* **17**, 3049-3061,
doi:10.1016/j.celrep.2016.11.043 (2016).
48   Wu, S. H., Ma, C. L. & Kelly, J. B. Contribution of AMPA, NMDA, and GABA(A)
receptors to temporal pattern of postsynaptic responses in the inferior colliculus of the rat.
J Neurosci **24**, 4625-4634, doi:10.1523/JNEUROSCI.0318-04.2004 (2004).
49   Liu, J., Wu, D. C. & Wang, Y. T. Allosteric potentiation of glycine receptor chloride
currents by glutamate. *Nat Neurosci* **13**, 1225-1232, doi:10.1038/nn.2633 (2010).
50   Metzner, K. *et al.* Lack of efficacy of a partial adenosine A1 receptor agonist in
neuropathic pain models in mice. *Purinergic Signal* **17**, 503-514, doi:10.1007/s11302-
021-09806-6 (2021).
51   Cui, X. *et al.* Enhancing spinal cord stimulation-induced pain inhibition by augmenting
endogenous adenosine signalling after nerve injury in rats. *Br J Anaesth* **132**, 746-757,
doi:10.1016/j.bja.2024.01.005 (2024).
52   Ran, F. A. *et al.* In vivo genome editing using Staphylococcus aureus Cas9. *Nature* **520**,
186-191, doi:10.1038/nature14299 (2015).
53   Cheng, L. *et al.* Tlx3 and Tlx1 are post-mitotic selector genes determining glutamatergic
over GABAergic cell fates. *Nat Neurosci* **7**, 510-517, doi:10.1038/nn1221 (2004).
54   Roth, B. L. DREADDs for Neuroscientists. *Neuron* **89**, 683-694,
doi:10.1016/j.neuron.2016.01.040 (2016).
55   Kagiyama, I., Uchiyama, S. & Tsuda, M. Hes5(+) astrocytes potentiate primary afferent
Adelta and C fiber-mediated excitatory synaptic transmission to spinal lamina I neurons.
Mol Brain **18**, 39, doi:10.1186/s13041-025-01212-y (2025).
56   Ji, R. R., Baba, H., Brenner, G. J. & Woolf, C. J. Nociceptive-specific activation of ERK
in spinal neurons contributes to pain hypersensitivity. *Nat Neurosci* **2**, 1114-1119,
doi:10.1038/16040 (1999).
57   Matsumoto, M., Xie, W., Ma, L. & Ueda, H. Pharmacological switch in Abeta-fiber
stimulation-induced spinal transmission in mice with partial sciatic nerve injury. *Mol*
Pain **4**, 25, doi:10.1186/1744-8069-4-25 (2008).
58   Kawabata, R. *et al.* The Anterior Cingulate Cortex is Critical for Acute Stress-induced
Hypersensitivity in Mice. *Neuroscience* **523**, 47-60,
doi:10.1016/j.neuroscience.2023.05.010 (2023).
59   Hickey, L. *et al.* Optoactivation of locus ceruleus neurons evokes bidirectional changes in
thermal nociception in rats. *J Neurosci* **34**, 4148-4160, doi:10.1523/jneurosci.4835-
13.2014 (2014).
60   Hirschberg, S., Li, Y., Randall, A., Kremer, E. J. & Pickering, A. E. Functional
dichotomy in spinal- vs prefrontal-projecting locus coeruleus modules splits descending
noradrenergic analgesia from ascending aversion and anxiety in rats. *Elife* **6**, e29808,
doi:10.7554/eLife.29808 (2017).
61   Duan, B. *et al.* Identification of spinal circuits transmitting and gating mechanical pain.
Cell **159**, 1417-1432, doi:10.1016/j.cell.2014.11.003 (2014).
62   Petitjean, H. *et al.* Dorsal Horn Parvalbumin Neurons Are Gate-Keepers of Touch-
Evoked Pain after Nerve Injury. *Cell Rep* **13**, 1246-1257,
doi:10.1016/j.celrep.2015.09.080 (2015).
63   Wang, H. *et al.* A novel spinal neuron connection for heat sensation. *Neuron* **110**, 2315-
2333.e2316, doi:10.1016/j.neuron.2022.04.021 (2022).

1031	64	Warwick, C. *et al.* Cell type-specific calcium imaging of central sensitization in mouse dorsal horn. *Nat Commun* **13**, 5199, doi:10.1038/s41467-022-32608-2 (2022).
1033	65	Kucharczyk, M. W., Di Domenico, F. & Bannister, K. Distinct brainstem to spinal cord noradrenergic pathways inversely regulate spinal neuronal activity. *Brain* **145**, 2293-2300, doi:10.1093/brain/awac085 (2022).
1036	66	Torsney, C. & MacDermott, A. B. Disinhibition opens the gate to pathological pain signaling in superficial neurokinin 1 receptor-expressing neurons in rat spinal cord. *J Neurosci* **26**, 1833-1843, doi:10.1523/jneurosci.4584-05.2006 (2006).
1039	67	Sonohata, M. *et al.* Noradrenaline modulates mechanically evoked responses in the rat spinal dorsal horn: an in vivo patch-clamp study. *J Pain Res* **12**, 1269-1278, doi:10.2147/jpr.s181210 (2019).
1042	68	Usoskin, D. *et al.* Unbiased classification of sensory neuron types by large-scale single-cell RNA sequencing. *Nat Neurosci* **18**, 145-153, doi:10.1038/nn.3881 (2015).
1044	69	Kawasaki, Y., Kumamoto, E., Furue, H. & Yoshimura, M. Alpha 2 adrenoceptor-mediated presynaptic inhibition of primary afferent glutamatergic transmission in rat substantia gelatinosa neurons. *Anesthesiology* **98**, 682-689, doi:10.1097/00000542-200303000-00016 (2003).
1048	70	Li, X. & Eisenach, J. C. alpha2A-adrenoceptor stimulation reduces capsaicin-induced glutamate release from spinal cord synaptosomes. *J Pharmacol Exp Ther* **299**, 939-944 (2001).
1051	71	Pankratov, Y. & Lalo, U. Role for astroglial α1-adrenoreceptors in gliotransmission and control of synaptic plasticity in the neocortex. *Front Cell Neurosci* **9**, 230, doi:10.3389/fncel.2015.00230 (2015).
1054	72	Xu, Q. *et al.* Astrocytes contribute to pain gating in the spinal cord. *Sci Adv* **7**, eabi6287, doi:10.1126/sciadv.abi6287 (2021).
1056	73	Oe, Y. *et al.* Distinct temporal integration of noradrenaline signaling by astrocytic second messengers during vigilance. *Nat Commun* **11**, 471, doi:10.1038/s41467-020-14378-x (2020).
1059	74	Lerman, S. F., Rudich, Z., Brill, S., Shalev, H. & Shahar, G. Longitudinal associations between depression, anxiety, pain, and pain-related disability in chronic pain patients. *Psychosom Med* **77**, 333-341, doi:10.1097/psy.0000000000000158 (2015).
1062	75	Lugert, S. *et al.* Homeostatic neurogenesis in the adult hippocampus does not involve amplification of Ascl1(high) intermediate progenitors. *Nat Commun* **3**, 670, doi:10.1038/ncomms1670 (2012).
1065	76	Vong, L. *et al.* Leptin action on GABAergic neurons prevents obesity and reduces inhibitory tone to POMC neurons. *Neuron* **71**, 142-154, doi:10.1016/j.neuron.2011.05.028 (2011).
1068	77	Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nat Neurosci* **13**, 133-140, doi:10.1038/nn.2467 (2010).
1071	78	Chaplan, S. R., Bach, F. W., Pogrel, J. W., Chung, J. M. & Yaksh, T. L. Quantitative assessment of tactile allodynia in the rat paw. *J Neurosci Methods* **53**, 55-63, doi:10.1016/0165-0270(94)90144-9 (1994).
1074	79	Atwal, N., Winters, B. L. & Vaughan, C. W. Endogenous cannabinoid modulation of restraint stress-induced analgesia in thermal nociception. *J Neurochem* **152**, 92-102, doi:10.1111/jnc.14884 (2020).
1077	80	Tsuda, M. *et al.* Behavioral phenotypes of mice lacking purinergic P2X4 receptors in acute and chronic pain assays. *Mol Pain* **5**, 28, doi:10.1186/1744-8069-5-28 (2009).

81   Kohro, Y. *et al.* A new minimally-invasive method for microinjection into the mouse
spinal dorsal horn. *Sci Rep* **5**, 14306, doi:10.1038/srep14306 (2015).
82   Zhang, Z. M., Chen, S. & Liang, Y. Z. Baseline correction using adaptive iteratively
reweighted penalized least squares. *Analyst* **135**, 1138-1146, doi:10.1039/b922045c
(2010).
83   Thévenaz, P., Ruttimann, U. E. & Unser, M. A pyramid approach to subpixel registration
based on intensity. *IEEE Trans Image Process* **7**, 27-41, doi:10.1109/83.650848 (1998).

1086