Assessing the conservation and targets of putative sRNAs in *Streptococcus pneumoniae*

Matthew C. Eichelman,[a] Michelle M. Meyer[a]#

[a]Department of Biology, Boston College, Chestnut Hill, Massachusetts, USA

Running Head: Conservation and targets of *S. pneumoniae* sRNAs

#Address correspondence to Michelle M. Meyer, m.meyer@bc.edu.

Abstract word count: 250

Abstract

RNA regulators are often found in regulatory networks and mediate growth and virulence in bacteria. Small RNAs (sRNAs) are non-coding RNAs that modulate translation initiation and mRNA degradation by base-pairing. To better understand the role of sRNAs in pathogenicity several studies identified sRNAs in *Streptococcus pneumoniae*; however, little functional characterization has followed. This study's goals are: 1) survey putative sRNAs in *S. pneumoniae*; 2) assess the conservation of these sRNAs; and 3) examine their predicted targets. Three previous studies in *S. pneumoniae* identified 287 putative sRNAs by high-throughput sequencing. This study narrows the candidates to a list of 58 putative sRNAs. BLAST analysis indicates that the 58 sequences are highly conserved across the *S. pneumoniae* pangenome, and 25 of them are identified sporadically in other Streptococcus species. However, only 2 have corresponding sequences identified across several Streptococcus species. We used four RNA-target prediction programs to predict targets for each of the 58 putative sRNAs. Across all probable predictions, six sRNAs have overlapping targets predicted by multiple programs, four of which target numerous transposase encoding transcripts. sRNAs targeting transposase genes display nearly identical and perfect base-pairing. One sRNA, M63 (Spd_sr37), has several probable targets in the CcpA regulon, a network responsible for global catabolite repression, suggesting a possible biological function in carbon metabolism control. Each M63-target interaction exhibits unique base-pairing increasing confidence in the biological relevance of the result. This study produces a list of S. *pneumoniae* putative sRNAs whose predicted targets suggest functional significance in transposon and carbon metabolism regulation.

Importance

Previous studies identified many small RNA candidates in *Streptococcus pneumoniae*, several of which were hypothesized to play a role in *S. pneumoniae* virulence. Due to the differing sequencing methods, diverse inclusion criteria, *S. pneumoniae* strain differences, as well as limited follow-up since, it is unclear to what extent candidates identified in different studies have overlapping sequences and functions, and their biological relevance remains ambiguous. This research aims to consolidate the candidate sRNAs across these studies and focuses attention on those that are likely to be regulatory and associated with virulence. This study's findings enhance our knowledge of the conservation of small regulatory RNAs across the many *Streptococcus pneumoniae* strains and highlights a handful that appear likely to have a role in growth or virulence.

Introduction

Streptococcus pneumoniae is a Gram-positive bacterium that causes various diseases including pneumonia, meningitis, bacteremia, otitis media and sinusitis. Invasive pneumococcal disease is particularly dangerous in children and the elderly (CDC et al., 2013), and in 2004 was responsible for approximately 4 million illness episodes, 445,000 hospitalizations, and 22,000 deaths in the United States (Huang et al., 2011). In 2016, *S. pneumoniae* was the leading cause of lower respiratory infection morbidity and mortality globally causing over a million deaths (GBD 2016 Lower Respiratory Infections Collaborators, 2016). Despite the threat *S. pneumoniae* poses, important components of regulation relating to metabolism and virulence remain less well characterized. Small regulatory RNAs (sRNA) are sequences of 40-500 nucleotides (nt) in length (Li et al., 2012) that can be transcribed by 5'-UTRs, 3'-UTRs, coding, and non-coding sequences

(Felden and Augagneur, 2021). However, studies seeking to identify sRNAs tend to focus on intergenic regions because RNAs transcribed in regions lacking an ORF are assumed to be more likely to be functional regulators, whereas those identified within ORFs may be an intermediate RNA decay product from a protein-encoding transcript. Among the different types of sRNAs are *trans*-encoded and *cis*-encoded RNAs. *Trans*-encoded sRNAs regulate genes from distant regions often with imperfect complementarity allowing them to interact with more than one target (Jabbour and Lartigue, 2021). *Cis*-encoded sRNAs act on the mRNA transcript encoded by the opposite DNA strand leading to perfect complementarity (Zorgani et al., 2016) (Fig. 1A). sRNAs modulate the expression of target mRNAs by base pairing to sequester a ribosome binding site or accelerate decay (Papenfort and Vanderpool, 2015). Some sRNAs are dependent on a chaperone like the Hfq or FinO family proteins that have a well-characterized role in aiding the formation of duplexes between sRNAs and their mRNA targets in Enterobacteriaceae. However, the role of RNA chaperones in Gram-positive bacteria is substantially less clear. Such proteins are frequently not present, and when they are present their functionality is often substantially different from that observed in gram-negative organisms (Christopoulou and Granneman, 2022). In *S. pneumoniae*, recent work shows KH domain proteins, like KhpA and KhpB, are associated with sRNAs (Olejiiczak et al., 2022), and sRNAs have been associated with the exonuclease Cbf1 (Hor et al. 2020). However, there are no confirmed sRNA chaperone homologs in the *S. pneumoniae* genome (Zhang et al., 2003).

Identification of putative sRNAs in *S. pneumoniae* has been performed several times, but follow-up characterization has been limited. The exception is a group of sRNAs called cia-dependent sRNAs (csRNAs) that have been studied across Streptococcus with considerable work done in *S. pneumoniae*. The csRNAs are controlled by the CiaRH two-component system (TCS) that is involved in natural competence and general virulence (Halfmann et al., 2007, Marx et al., 2010, Tsui et al., 2010). In *S. pneumoniae* the CiaRH TCS expresses five sRNAs, with experimentally verified targets, that prevent autolysis triggered by various conditions, like the presence of deoxycholate, to allow the maintenance of stationary growth phase (Mascher et al., 2006). The csRNAs have also recently been implicated in promoting Zn homeostasis (De Lay et al., 2024). Three previous studies identified hundreds of additional putative sRNAs using diverse inclusion criteria. Some of these sequences may be annotated as homologs of RNA families such as Pyr elements (RF00515) or TPP riboswitches (RF00059), BOX elements (AT-rich repeats that are highly transcribed), or ribosomal protein leaders (sequences in the 5'-UTR of ribosomal protein transcripts that control the concentration of the ribosomal protein) (Mann et al., 2012, Fu et al., 2013, Babina et al., 2015). However, it remains unclear which of the remaining sequences are regulatory. Here we find that 58 of the putative sRNAs are highly conserved in the *S. pneumoniae* pangenome and we predict the mRNA targets of these sRNAs. The predictions suggest 7 putative sRNAs, which have highly probable targets, interact with mRNAs coding for transposases and genes involved in carbon metabolism regulation.

Results and Discussion
S. pneumoniae genome contains +70 putative sRNAs
In assessing which previously identified sRNA candidates are likely to have a biological function and prioritize candidates for further investigation we examined a pool of 287 putative sRNAs originating from three studies (Acebo et al., 2012, Mann et al., 2012, and Sinha et al., 2019, Additional Datafile 1). We note 65% are functionally uncharacterized whereas the other 35%

may be annotated as homologs of *cis*-regulators, BOX elements, csRNAs, or overlap with small
open reading frames (sORF) (Fig. 1B). We further narrowed this pool to a list where each sRNA
has at least 1 of 3 attributes: 1) sequence identified in multiple studies, 2) expression confirmed
by Northern blot or RT-qPCR, 3) sequence characterized as a csRNA or overlapping with a
sORF (Laczkovich et al., 2022). *Cis*-regulators and BOX elements were also excluded from our
list. Thus, the new list contains 81 putative sRNAs that were assigned names "M1" through
"M81". However, 5 sequences are <50 nt and were subsequently removed from the list leaving a
final total of 76 candidate sRNAs (Fig. 1C). We observe that many of these sRNAs were found
in a single study emphasizing the different sequencing strategies and inclusion criteria of the
previous studies (Fig. 1D). After compiling the final list for further analysis, we conclude there
are over 70 putative sRNAs in the *S. pneumoniae* genome (Additional Datafile 2).
115

116
117 **Figure 1**: **A**) The *cis*-encoded sRNA, found on the strand opposite to the mRNA coding strand, binds the target
118 mRNA with perfect complementarity. The *trans*-encoded sRNA, expressed from a region distant to the target, binds
119 with imperfect complementarity. **B)** The classifications of the 287 putative sRNAs. **C)** Process of narrowing the 287
120 putative sRNAs to a list of 76 candidates for further analysis. **D)** Among the candidates, 7 were identified by all
121 three studies and 15 by two of the three studies.
122

Majority of sRNA candidates are conserved across the *S. pneumoniae* pangenome
To increase our confidence in the biological relevance of the putative sRNAs and prioritize them
for further investigation, we assessed the conservation of the candidate sRNAs across the
genomes of 385 *S. pneumoniae* strains. BLAST (Altschul et al., 1990) analysis indicated 70/76
candidates are present in the genomes for a majority of 385 *S. pneumoniae* strains (Fig. 2A)

(Cremers et al. 2015, Rosconi et al., 2022). Among these 70, only 60 candidates appear to be
non-repetitive sequences, all of which display average sequence identity >97% to the best hit in
each genome, indicating the sequences are highly conserved across the *S. pneumoniae*
pangenome (Table 1). Interestingly, all of the 6 sRNAs found in <12 strains were identified in the
S. pneumoniae strain TIGR4 (Fig. 2), highlighting the possibility for strain specific sRNAs in *S.*
pneumoniae. Two of the final candidates, M8 and M12, were identified as *cis*-regulatory
elements later during our analysis and removed, leaving a total of 58 sRNAs for further analysis.
From this final group of 58 candidates, we observed that synteny is preserved (see Methods)
across the *S. pneumoniae* strains in 43 of the 58 final candidates. To determine if any of the
sRNAs are conserved in species other than *S. pneumoniae* we also analyzed a range of other
species in the Streptococcus genus. Only 2 of the candidate sRNAs (M77 and M81) align with
sequence identities >65% to each of *S. pyogenes*, *S. mutans*, *S. suis*, *S. mitis*, *S. oralis*, and *S.*
gordonii, and 25 additional sRNA candidates can be identified in a subset of these organisms
(Supplemental Table S1). Thus, it appears over half of the sRNAs are unique to *S. pneumoniae*,
consistent with the narrow distribution of many sRNAs across other bacterial species (Peer and
Margalit, 2014).
144
Several sRNA candidates overlap with reported sORFs
Of the 58 final sRNAs candidates, 6 of them overlap with previously identified sORFs (Table 1)
(Laczkovich et al., 2022), and one (M64) is complementary to an sORF. Moreover, we see that
these 5 of the 7 sORFs are found in nearly all 385 strains with 1 alignment per genome and a
high degree of conservation with nucleotide and peptide sequence identities >98%. The first
exception, rio56, is a 6 amino acid peptide sequence and is too short for BLAST to produce
alignments with an e-value < 20, and the second, rio86, has an average peptide sequence identity
of 93.6%. Of particular interest is M61 (Spd_sr5/srf-02), which overlaps with rio3, whose
expression was shown to promote *in vivo* fitness (Laczkovich et al., 2022). However, dual-
function RNAs, sequences with coding and non-coding functions, are known to exist in other
bacteria like SgrT/SgrS and AzuC/AzuR in *Escherichia coli* (Aoyama and Storz, 2023) and Pel
RNA in *Streptococcus pyogenes* (Raina et al., 2018). Thus, we decided to retain sORF
overlapping sRNAs for target analysis.
158

A

76 sRNA candidates

1. <u>Conserved across pangenome</u>:
   Found in >338 strains?

2. <u>Non-repetitive sequence</u>:
   ≤15 alignments per genome?

3. <u>Synteny preserved</u>:
   Are the sequence identities of
   1000 nt up and downstream of
   the sRNA ≥75%

B

Conservation of sRNA candidates

Figure 2: **A)** The conservation of sRNA candidates across the *S. pneumoniae* pangenome. The * next to "60" highlights that 2 sRNAs were later removed to form the final list of 58 sRNAs (see Table 1 legend). **B)** The four degrees of conservation of the sRNA candidates. "Conserved", "non-repetitive sequence", and "synteny preserved" refer to the criteria for each level of the tree diagram in part A.

In-house ID	Other IDs	TIGR4 coordinates	D39 coordinates	Sequence identity	Genomes with match	Matches per genome	Overlap with sORF
M1	srn061/F7/CcnE	209768-209916	212278-212426	99.99%	385[1]	1	
M2	srn135	438151-438275		100.00%	8	1	
M3	srn151	501260-501363		96.32%	385	19	
M4	F15	501732-501843		99.04%	385[1]	1	
M5	srn206	781187-781304		99.77%	384[1]	1	
M6	srn231	853483-853605		97.97%	348[1]	1	
M7	srn235/F59/Spd_sr49	869478-869804	825803-826129	99.39%	385[1]	1	
M8**	srn241/F35	909029-909179		98.93%	385[1]	1	
M9	srn254/F38/Spd_sr17	956782-956927	912572-912717	97.90%	385[1]	1	rio56
M10	srn266/Spd_sr55	1035442-1035581	1697549-1697677	97.33%	385	15	
M11	srn277/F41	1071102-1071213		91.09%	4	1	

The 76 in-house sRNAs and their conservation

M12**	srn351/R21/Spd_sr78	1461066-1461188	1404039-1404161	99.93%	385[1]	1	
M13	srn477/Spd_sr105	1984829-1984968	1873277-1873416	99.86%	385[1]	1	
M14	trn0978	1986229-1986310		99.70%	385[1]	1	
M16	srn491/Spd_sr106	2005533-2005682	1892401-1892550	99.86%	385[1]	1	
M17	srn502/F66/Spd_sr109	2086085-2086325	1972860-1973100	99.55%	385	2	
M18	srn503/F67/Spd_sr111	2086380-2086628	1973155-1973403	98.41%	385	2	
M19	trn0012/CcnC	24164-24262	23967-24065	99.62%	385	2	
M20	trn0057	124989-125050		97.19%	385	1	
M21	Spd_sr18		134413-134576	96.75%	385	92	
M22	trn0157/F10	284239-284316		99.67%	374[1]	1	
M23	trn0332/F25/Spd_sr42	623244-623344	587440-587542	95.89%	378[1]	1	
M24	trn0485/F60	950115-950195		99.99%	385[1]	1	
M25	trn0696	1423633-1423713		93.05%	364	1	
M27	trn0830/R12	1731041-1731439		94.32%	385	209	
M28	trn0935/R9	1903548-1903637		99.82%	385	19	
M29	trn1025	2048577-2048643		99.99%	366[1]	1	
M30	R7	1791010-1791079		98.58%	385	6	
M31	R8	1892645-1892716		99.54%	9	1	
M32	R14	1034021-1034099		98.69%	385[1]	1	
M33	R17/Spd_sr69	1277241-1277387	1217391-1217537	97.32%	385	79	
M34	R18/Spd_sr72	1364575-1364763	1299979-1300167	98.30%	379[1]	1	
M36	F1	91593-91663		99.09%	361	1	
M37	F3	117143-117247		98.43%	385	1	
M39	F6	130439-130494		98.30%	383[1]	1	
M40	F8/CcnA	228626-228807	231143-231324	97.12%	385[1]	2	
M41	F11	286614-286707		99.13%	384[1]	1	
M42	F14	499570-499688		99.77%	339[1]	1	
M43	F18	538437-538491		97.84%	385	7	
M44	F19/Spd_sr33	543000-543149	508238-508388	99.11%	385[1]	1	rio39
M45	F22	592573-592711		99.44%	385[1]	1	
M46	F24	610528-610659		98.04%	385	12	
M47	F31	810811-810861		98.29%	385	12	
M48	F33	863736-863817		98.28%	385[1]	1	
M49	F36	941435-941486		99.66%	385[1]	1	
M50	F39	972498-972606		98.15%	385[1]	1	
M51	F40	1063101-1063150		100.00%	4	1	
M52	F43/Spd_sr63	1216148-1216245	1170289-1170386	99.87%	385[1]	1	
M53	F45	1408204-1408274		100.00%	384[1]	1	
M54	F48	1778293-1778426		99.27%	385	36	
M56	F53	588512-588589		99.92%	385	26	
M57	F55	1696066-1696161		97.13%	385	48	
M58	F56	158993-159089		98.09%	11	1	
M59	F61	972327-972383		98.82%	385[1]	1	
M60	F62	995726-995786		100.00%	9	1	
M61	Spd_sr5		39980-40081	97.40%	358[1]	1	rio3
M62	Spd_sr6		41494-41558	97.76%	384	7	
M63	Spd_sr37		131773-131841	99.83%	385[1]	1	rio16
M64	Spd_sr14		149223-149340	99.59%	385[1]	1	rio17

M65	CcnB		231331-231426	99.71%	385	3	
M66	Spd_sr24		231853-232034	99.34%	371	1	
M67	CcnD		233715-233808	99.89%	385[1]	1	
M68	Spd_sr31		476085-476234	98.76%	385[1]	1	
M69	Spd_sr47		825484-825544	99.38%	385[1]	2	
M70	Spd_sr60		1079136-1079199	99.99%	383[1]	1	
M71	Spd_sr67		1212230-1212526	98.77%	365[1]	2	
M72	Spd_sr71		1264469-1264569	99.06%	385	37	
M73	Spd_sr81		1464371-1464684	99.17%	385[1]	1	
M74	Spd_sr83		1528062-1528186	99.64%	385[1]	1	rio82
M75	Spd_sr84		1595446-1595563	98.99%	385[1]	1	
M76	Spd_sr89		1673201-1673322	99.92%	385[1]	1	rio86
M77	Spd_sr96		1759320-1759411	100.00%	385[1]	1	
M78	Spd_sr108		1913212-1913442	97.60%	385	52	
M79	Spd_sr110		1973001-1973113	99.16%	385	2	
M80	Spd_sr112		1973343-1973456	98.78%	385	2	
M81	Spd_sr116		2020113-2020228	99.98%	385[1]	1	

Table 1: The 76 candidate sRNAs and their conservation. In the "Other IDs" column, sRNAs with the prefix "srn", "trn", "R", and "F" were identified in TIGR4 (Acebo et al., 2012, Mann et al., 2012) and sRNAs with the prefix "Spd_sr" and "Ccn" were identified in D39W (Sinha et al., 2019). The TIGR4 coordinates indicate the sequence location in the NC_003028.3 genome. Likewise, D39 coordinates corresponds to the NC_008533.2 genome. The "Sequence identity" and "Matches per genome" are averages across the 385 strains. Preservation of synteny indicated by [1] in the "Genomes with match" column. **M8 later identified as a *cis*-regulator (pyrR element: RFAM:RF00515), and M12 identified as a Mn^{2+} responsive riboswitch (Martin et al. 2019, RFAM:RF000080). The "Overlap with sORF" column indicates the sRNA overlaps with a previously identified sORF (Laczkovich et al., 2022). White rows correspond to the 58 final sRNAs that are conserved in many *S. pneumoniae strains*, but have fewer than 15 instances per genome.

RNA target prediction programs struggle to correctly predict validated targets

Several RNA-RNA interaction prediction (RIP) programs have been developed to predict mRNA:sRNA interactions with newer models displaying the highest accuracies. These include IntaRNA (Busch et al., 2008, Wright et al., 2014, Mann et al., 2017, Raden et al., 2018), CopraRNA (Wright et al., 2013, Wright et al., 2014, Raden et al., 2018), sRNARFTarget (Naskulwar and Peña-Castillo, 2022), and TargetRNA3 (Tjaden, 2023). The programs take various approaches with newer RIP programs implementing machine learning algorithms. Despite the improvement over time, all the tools have a high false positive rate (Tjaden, 2023). Moreover, most of the data on which the models are validated and trained is from Hfq-dependent sRNA networks in *E. coli* that may not be reflective of sRNA-target interactions in organisms without Hfq like *S. pneumoniae*. This poses a challenge to determining the validity of a predicted target through computational methods alone. By examining the targets of multiple programs with different approaches we hope to increase the confidence in the validity of predicted sRNA-target pairs.

As a baseline evaluation of the RIP programs, we compared the predicted and known targets of the csRNAs using IntaRNA, sRNARFTarget, and TargetRNA3 (Schnorpfeil et al., 2013). None of the programs correctly predicted any of the known csRNA targets (SP_2237/SP_RS11435, SP_0090/ SP_RS00460, SP_0161/ SP_RS00830, SP_0626/SP_RS03070, and SP_1215/SP_RS05965) as the most likely target. If we include the top five most likely targets, then IntaRNA correctly predicts that csRNAs 2 and 3 target SP_RS00460 and both IntaRNA and

198 TargetRNA3 correctly predicted that csRNA4 targets SP_RS00460. We also confirmed the
199 sequences that we examined are consistent with reported 5' and 3' RNA-seq data in TIGR4
200 (Warrier et al., 2018, Furumo and Meyer, 2024) to ensure our inputs were not causing the low
201 accuracy. These results support the existing notion that even the best RIP programs suffer from a
202 high false positive rate, but do provide informative results.
203
RIP programs predict thousands of sRNA-target pairs
We used multiple programs to make target predictions for the candidates. For all 58 sRNAs we
used IntaRNA, sRNARFTarget, and TargetRNA3. Targets were predicted in six different *S.*
pneumoniae strains: TIGR4, D39, and four arbitrarily selected strains from PRJNA514780
(Rosconi et al., 2022) (see Methods). We also used CopraRNA, but only for the 3 sRNAs with
sequence identities >65% in at least 4 related Streptococcus species due to the algorithm's
comparative approach to target identification. CopraRNA was used to make predictions for *S.*
pneumoniae, *S. pyogenes*, *S. mutans*, *S. suis*, *S. mitis*, *S. oralis*, and *S. gordonii*. Each program
produces a variable number of outputs per sRNA per strain/species. IntaRNA and CopraRNA
made five predictions (a customizable parameter), sRNARFTarget predicted a probability for
every gene in the *S. pneumoniae* transcriptome (>2000 genes), and TargetRNA3 reported a
variable number of targets with a probability and p-value above a customizable threshold.
216
In total, we obtained thousands of predictions, the majority of which have low probabilities
($\leq$0.5) (See Additional Datafiles 3-7). To focus our attention on likely sRNA-target pairs without
excluding too many predictions we settled on targets with a predicted probability $\geq$0.7, referred
to as probable going forward. We also define the term MPT, most probable target, as the
prediction given the highest probability across all predictions for a given sRNA. Lastly, we
define a consensus target as a gene that was predicted to be the MPT for an sRNA in at least four
of the six *S. pneumoniae* strains. This term only pertains to the predictions made by IntaRNA,
sRNARFTarget, and TargetRNA3. We observed none of the sRNARFTarget predictions are
probable. This in combination with our baseline evaluation of the csRNAs led us to focus on the
predictions made by IntaRNA, TargetRNA3, and CopraRNA when applicable.
227
In many cases sRNA pairing can impact the internal base pairing of the mRNA to enable gene
expression changes (Papenfort and Storz 2024). Thus, we assessed whether the putative mRNA
target regions are likely to have internal base pairing that may be impacted sRNA interaction. We
used RNAfold (Lorenz et al., 2011) to predict the structure of mRNA target sequences including
25 nucleotides up/downstream of the binding region in the absence of any sRNA partner. We
consider a structured region to be a segment of the mRNA sequence displaying internal base
pairing (>2 consecutive bases). Across the MPTs predicted by IntaRNA and TargetRNA3, 51 out
of 58 sRNAs base pair with a region considered structured suggesting that the putative sRNA-
mRNA interaction may induce structural changes in the secondary or tertiary structures to enable
regulation (Tieng et al., 2023).
238
sRNAs may play a role in pathogenesis via their targets
Previous studies used transposon insertion mutants to conclude specific sRNAs may support
virulence in TIGR4 (Mann et al., 2012). We compared our predicted targets for these sRNAs
to evaluate these hypotheses. Previous work suggested 8 putative sRNAs (Table 2) play a
definitive role in pathogenesis, and some individual target loci were identified by microarray
analysis of attenuated sRNA mutants (Mann et al., 2012). Only 3 of these sRNAs met our criteria

for further investigation (Fig. 2A, Table 2). The others overlap with known *cis*-regulatory
elements (F20 and F44), transfer-messenger RNA (F32), or were removed following
conservation analysis (F41, F48). F41 is one of the sRNAs found in <12 strains, and F48 was
deemed a repetitive sequence (average of 36 copies per genome). Among the three remaining
candidates, one (M1/F7) is a csRNA with an established role in pathogenesis (DeLay 2024,
Patenge et al., 2012, Schnorpfeil et al., 2013). The other two, M45 and M23, are not
characterized. M45 is predicted to target type IV teichoic acid flippase TacF that is responsible
for transporting choline across the cytoplasmic membrane, a nutritional requirement of *S.*
pneumoniae (Damjanovic et al., 2007). M23 was originally reported to target SP_RS08340-50, a
putative carbohydrate transporter, based on microarray analysis, but we predict that it targets a
transposase encoding transcript (SP_RS13320) (Table 2). Without validating the targets, the role
of M45 and M23 in pathogenesis is unclear, however the predicted target (*tacF*) of M45 is
suggestive of such a role.
258

Mann study ID	In-house ID	Mann Putative Target	In-house Consensus Target
F7	M1, csRNA5		SP_RS06250
F20	T-box leader		
F22	M45		SP_RS06235
F25	M23	SP_RS08340-50	SP_RS13320
F32	tmRNA		
F41		SP_RS08340-50	
F44	PyrR binding site	SP_RS08340-50	
F48			

259
260   **Table 2**: A comparison of the in-house target predictions and the putative targets identified by microarray analysis.
261   The previously identified putative targets SP_RS08340-50 are three neighboring loci involved in carbohydrate
262   transport (carbohydrate ABC transporter permease) and proposed to be collectively regulated by three of the sRNAs
263   (Mann et al., 2012).
264

To further assess whether specific sRNAs are potentially regulating multiple targets in a
previously recognized regulatory response (Table 3), we investigated whether the predicted
targets belong to established operons or regulons in TIGR4. The regulons that appeared the most
often are PyrR, CodY, and CcpA and we noticed the targets belonging to established operons are
always the first or last gene in the operon with the first gene being more common. This suggests
that the sRNAs may be inhibiting translation, typically blocking the ribosome binding site (e.g.
start of an operon), or stabilizing the transcript by binding to the 3' end of the mRNA, depending
on the relative location of interaction (Papenfort and Vanderpool, 2015). Most notably one
sRNA, M63 (Fig. 3A), is predicted to target four different genes in the CcpA regulon (Table 3),
genes regulated by the catabolite control protein A (CcpA), an essential transcription factor in
Gram-positive bacteria that is responsible for mediating carbon catabolite repression and
activation. In *S. pneumoniae*, mutations in CcpA reduce virulence in mouse models
(Giammarinaro & Paton, 2002; Iyer et al., 2005). We also see that M63 interacts with the
different targets in various regions of the sRNA with unique base-pairing (Fig. 3B). Lastly, M63
is unique in that it is the only sRNA candidate where every reported target has a probability
≥0.7. Furthermore, this is true for all 6 *S. pneumoniae* strains assessed. M63 also overlaps with
the rio16 sORF, indicating that part of this sRNA is translated. The sORF has a conserved
sequence with average sequence identity of 99.7% across the pangenome. However, it remains
possible that M63 is a dual-function RNA both encoding a small protein and regulating the

members of the CcpA regulon. The observation that M63 has multiple probable targets acting on
a regulon associated with virulence (Iyer et al. 2005) makes this sRNA a high priority candidate
for further investigation.
287

sRNA	Target gene	Regulon	Position in operon	Target annotation
M63	*adhE*/SP_2026/SP_RS10245	CcpA	1 of 1	bifunctional acetaldehyde-CoA/alcohol dehydrogenase
M63	*adhE*/SP_2026/SP_RS10245	Rex	1 of 1	bifunctional acetaldehyde-CoA/alcohol dehydrogenase
M63	*hexA*/SP_0498/SP_RS02450	CcpA	1 of 1	bacterial Ig-like domain-containing protein
M63	*hexB*/SP_0057/SP_RS00325	CcpA	1 of 1	LPXTG-anchored beta-N-acetylhexosaminidase StrH
M63	*oppA2*/SP_1891/SP_RS09395	CodY	1 of 5	peptide ABC transporter substrate-binding protein
M63	*glgB*/SP_1121/SP_RS05550	CcpA	1 of 4	1,4-alpha-gltcan branching protein GlgB
M63	*ileS*/SP_1658/SP_RS08185	T-box(Ile)	1 of 1	isoleucine--tRNA ligase
M63	*carB*/SP_1275/SP_RS06250	PyrR	1 of 1	carbamoyl-phosphate synthase large subtnit

288
289 **Table 3**: The regulons, according to the RegPrecise database, in which all targets of M63 are involved. Predictions
290 made by TargetRNA3 in *S. pneumoniae* strain TIGR4.
291

292
293 **Figure 3**: **A)** The M63 sequence with underlined subsequences indicating the regions interacting in part B. Blue
294 corresponds to the *adhE* interaction, orange to *hexA*, and green to *glgB*. **B)** Three of the M63 interactions predicted
295 by TargetRNA3 in TIGR4. The gene names are predicted targets of M63 in the CcpA regulon. The numbers on
296 either end are relative positions of the interacting sequences within the full RNA sequence for the sRNA. For the
297 mRNAs, numbering is relative to the mRNA start codon. A ":" indicates a G-U pair.

298
Transposase associated sRNAs are frequent
Among the TargetRNA3 predictions, we noticed a large number of transposase associated
targets. Candidates targeting transposases encoding transcripts include M10, M47, M62, and
M69. These sRNAs are all encoded antisense to an annotated transposase (IL3 or IL30 family),
overlapping with the 5'-UTR or first few amino acids of the gene. A subset of these, M10, M47
and M62, show substantial sequence identity to each other, with M10 having a 3'-extention
compared to M47 and M62 (Fig. 4A). Notably, M10, M47, and M62 all have a large number of
BLAST hits in the genome, but these sequences did not exceed our threshold of >15 hits in the
genome to be considered repetitive sequences. M47 and M62 have consensus targets with both
IntaRNA and TargetRNA3 predicting transposase encoding targets. The collected targets for this
set of sRNAs (M10, M47, and M62) includes over 18 different transposase genes, with all but
two interactions showing high probability ≥0.92 (Supplemental Table S2). This large number of
targets results from the duplicated nature of the sequence immediately surrounding the

transposase gene (Fig. 4B). However, we note that all but one of the targets of M62 have a
predicted probability below our threshold of 0.7, potentially the result of the differences in length
between the M62 sequence and the M10 and M47 sequences (region accessibility due to RNA
folding is a factor in target prediction and may be influenced by the extra flanking sequence).
316
There are several well-characterized examples of transposon antisense encoded RNAs in bacteria
including RNA-OUT, inhibiting IS10 (Simons and Kleckner, 1983), art200, inhibiting IS200
(Ellis et al., 2015), and RNA-C inhibiting IS30 (Arini et al., 1997). Transposon associated
antisense RNAs that overlap the transposon coding sequence proximal to the start codon
typically act in trans, blocking the translation of the transposase (Simons and Kleckner 1983,
Ellis, 2015), but RNA-C, which is antisense to the transposase gene but not directly at the start
codon, only appears to act in cis (Arini, 1997). Thus, based on the position of these sRNAs
proximal and overlapping the ribosome-binding site, it is likely that they are *trans*-acting across
the many transposon copies present in the genome. The sequence of M69 is distinct from that of
M10, M47, and M62; however, its placement upstream and antisense to an annotated transposase
gene suggests a similar functionality.
328

A

```
M62    GTTCCATATGAATCTTTCTAATGAGTTGTTTGATCGCTTTTCATTATAGATCTTATGGGA    60
M47    --------------TTTCTAATGAGTTGTTTGGTCGCTTTTCATTATAGGTCATATGGGA    46
M10    GTTCCATATGAATCTTTCTAATGAGTTGTTTTGTCGCTTTTCATTATAGGTCATATGGGA    60
                     ***************** ***************** ** *******

M62    CTTTT-------------------------------------------------------    65
M47    CTTTT-------------------------------------------------------    51
M10    CTTTTTTCTACACAAAAATAGGCTCCATAATATCCATAGGGGATTTACCCACTACAAATA    120
       *****

M62    -------------------    65
M47    -------------------    51
M10    TTATAGAGTCATTTTTGCAC    140
```

B

```
                                                  A
M10         39    UUCGCUGUUUUGUUGAGUAAUCUUUCU AGUAUACCUUG    1
                  |||||||||||||||||||||||||||| |||||||||||
SP_RS06190  -32   AAGCGACAAAACAACUCAUUAGAAAGA UCAUAUGGAAC    7
                                                  A

M10         39    UUCGCUGUUUUGUUGAGUAAUCUUUCUAAGUAUACCUUG    1
                  |||||||||||||||||||||||||||||||||||||||
SP_RS12420  -32   AAGCGACAAAACAACUCAUUAGAAAGAUUCAUAUGGAAC    7

                       U U                        UA   U
M10         39    UUCGC G UUUGUUGAGUAAUCUUUC  AG AUACCUUG    1
                  |||||  |  ||||||||||||||||  ||  |||||||||
SP_RS02270  -32   AAGCG C AAACAACUCAUUAGAAAG   UC UAUGGAAC    7
                       C U                        UA   U
```

Figure 4: **A)** Clustal omega alignments of M10, M47, and M62 which are predicted to target numerous transposase encoding genes. '*' indicates a position in which all three sequences overlap. **B)** Three of the M10 interactions predicted by TargetRNA3 in TIGR4. The numbers on either end are relative positions of the interacting sequences within the full RNA sequence. A negative number indicates the sequence is upstream of the mRNA start codon. SP_RS### is the TIGR4 target locus. The underlined "AUG"s are the transposase gene start codons.

Cis-encoded sRNA candidates are less common

To identify other *cis*-encoded sRNA candidates we compared the genomic coordinates of the candidates and MPTs. Candidates suspected to be *cis*-encoded must have genomic coordinates overlapping with the target coordinates. IntaRNA shows 19 candidates may be *cis*-encoded, three of which exhibit probable interaction (Fig. 5A). In contrast, TargetRNA3 suggests only a single candidate to be *cis*-encoded, and it is probable and in common with IntaRNA's results (Fig. 5B). The possible *cis*-encoded sRNA predicted by TargetRNA3 is M66 (Fig. 5C). The M66 sequence appears antisense to the target with perfect binding across 40 nucleotides. M66 targets *ruvB* that codes for the Holliday junction branch migration DNA helicase RuvB, a subunit in the RuvABC complex. The complex processes Holliday junctions, nucleic acid structures that contain four joined double-stranded arms, during genetic recombination and DNA repair. The individual RuvB subunit is a hexameric ring helicase that acts like a motor to draw the DNA through the complex (Sharples et al., 1999). We believe that the overlap in *cis*-encoded predictions made by TargetRNA3 and IntaRNA suggest that this target is one of interest.

Figure 5: **A)** The distribution of sRNA locations relative to its targets predicted by IntaRNA for 58 candidate sRNAs **B)** The distribution of sRNA locations relative to its targets predicted by TargetRNA3. **C)** The M66 cis-interaction is predicted by TargetRNA3 and the sRNA is found on the complement strand and the target on the top strand. The numbers on either end are exact positions of the interacting sequences in the D39 genome (NC_008533.2).

Seven notable sRNAs for future experimental validation

Across the 58 sRNAs, 7 stood out for reasons that we believe warrant future work to experimentally validate this study's results. Each sRNA is highly conserved and targets a gene

with high probability. 4 of the notable sRNA candidates share a consensus target between at least two RIP programs (M18, M47, M62, M66) (Table 4). We believe a consensus target, a gene predicted to be the MPT for an sRNA in at least four of the six *S. pneumoniae* strains, is indicative of a highly likely true sRNA-target pair. Four of the notable candidates target multiple transposases (M10, M47, M62, M69). In addition to transposon associated sRNAs, there are also sRNA candidates with potential metabolic targets such as M63, which has 13 probable targets four of which are in the CcpA regulon. We also checked PneumoBrowse2 (Janssen et al., 2024), an interactive online platform with detailed annotations of *S. pneumoniae* genomes like D39V, D39W and TIGR4, for additional annotations on our 7 notable sRNAs. M18 is annotated as part of an anti-toxin/toxin system (Type I addiction module toxin, Fst family) and our predicted target is the ClC family H(+)/Cl(-) exchange transporter. From our analysis, we speculate that these 7 sRNAs are the most likely to lead to future validation of true sRNA-target pairs that may inform us about the role of sRNAs in *S. pneumoniae* metabolism and virulence.

In-house sRNA ID	Other IDs	Consensus	Predicted target(s)	Mechanism	Additional Comments
M63	Spd_sr37/srf-04		CcpA regulon	trans	Possibly influences virulence via CcpA regulon
M10	srn266/Spd_sr55		transposases	trans	Similar targets to M47, many transposon related
M47	F31	✓	transposases	trans	Similar targets to M10, many transposon related
M62	Spd_sr6/srf-03	✓	transposases	trans	RNA sequence related to M10 and M47
M69	Spd_sr47		transposases	trans	Targets 2 transposases with high probability
M66	Spd_sr24	✓	*ruvB*	cis	*ruvB* involved in recombination and DNA repair
M18	srn503/F67/Spd_sr111	✓	SP_RS05710	trans	Targets an H(+)/Cl(-) exchange transporter

Table 4: Notable sRNAs with a high degree of conservation and interesting predicted targets. Other IDs are the labels assigned in the data source studies. The "Consensus" column refers to whether a consensus target was found across multiple RIP programs for the sRNA. The "Mechanism" column refers to whether the sRNA is suspected to be *cis*-acting or *trans*-acting.

Conclusion

This study compiled a list of sRNAs in *S. pneumoniae*, then analyzed their conservation across the Streptococcus genus and within the *S. pneumoniae* pangenome and predicted mRNA targets of conserved sRNAs. BLAST indicates 58 sRNAs exhibit strong conservation across 385 strains. Four RNA-RNA interaction prediction programs made thousands of predictions for the 58 sRNAs. Ultimately, only the probable targets predicted by IntaRNA and TargetRNA3 were the focus of this study's target examination. From our analysis we propose that there are a handful of transposon associated sRNAs that target transposases encoding transcripts, likely acting in trans based on the position of the base-pairing. However, we also identified high probability targets for other sRNA candidates. For example, M63 (Spd_sr37/srf-04), has several predicted targets across the CcpA regulon implying a potential role in carbon metabolism. Through this work, we have identified a list of 7 sRNAs for which biological function can be hypothesized. Future work will strive to experimentally validate these hypotheses to reveal more regarding the nature of these sRNAs, their targets in *S. pneumoniae*, and roles in growth and virulence.

Methods

Compiling sRNA data sources

Previous studies identified putative sRNAs by high-throughput sequencing in *S. pneumoniae* strains TIGR4 (NC_003028.3) and D39W (CP000410.1). The putative sRNAs in TIGR4 (Acebo

399 et al., 2012, Mann et al., 2012) and D39W (Sinha et al., 2019) were narrowed down to a new list
400 of candidates for further analysis of conservation and target prediction (see Results and
401 Discussion).
402
403 **Creating the in-house list of putative sRNAs**
404 A list of in-house putative sRNAs was created from the candidates. Previous TIGR4 studies
405 identified different coordinates for transcription initiation and termination sites, so new in-house
406 coordinates were created by combining the smallest initiation and largest termination site
407 coordinates (Supplemental Fig. S1A). In-house Python scripts retrieved the sRNA sequences
408 from the TIGR4 and D39W genomes using the new coordinates. sRNAs identified in multiple
409 studies under different names were assigned an in-house ID and sequence using the new
410 coordinates (e.g. csRNA5/SN35/srn061/F7/CcnE becomes M1). Differences between the TIGR4
411 and D39W genomes forced the need to compare sequences rather than coordinates.
412 VectorBuilder (https://en.vectorbuilder.com/tool/sequence-alignment.html) aligned sRNAs to
413 confirm the sequences overlap. The largest possible sequence became the new in-house sRNA by
414 joining the overlapping subsequence and trailing sequences on either end (Supplemental Fig.
415 S1B). After forming the list of in-house sRNAs, sequences with length <50 nucleotides were
416 removed.
417
418 **Assessing conservation of sRNAs**
419 The in-house sRNAs were aligned to the genomes of 385 *S. pneumoniae* strains (Cremers et al.,
420 2015, Rosconi et al., 2022), *S. pyogenes* (NZ_LS483338.1), *S. mutans* (NZ_CP044221.1), *S. suis*
421 (NC_012926.1), *S. mitis* (NZ_GL397179.1), *S. oralis* (NZ_LR134336.1), and *S. gordonii*
422 (NZ_CP077224.1). Raw reads and the mapping results for 350 *S. pneumoniae* strains (Cremers
423 et al., 2015) available in BAM format were converted to consensus files in FASTA format with
424 the samtools consensus mode (Danecek et al., 2021). Then, a database, created with
425 makeblastdb, containing the sRNA sequences was aligned to each genome using blastn with the
426 task parameter set to megablast. Similarly, a database created with makeblastdb containing the
427 385 *S. pneumoniae* strains was aligned to each of the sORF nucleotide and peptide sequences
428 using blastn and tblastn. The e-value maximum threshold was raised from a default of 10 to 20
429 for the tblastn alignments. An in-house Python script retrieved the average number of
430 alignments, best sequence identity, and number of genomes with an alignment for each sRNA.
431 In-house sRNAs with an average number of alignments per strain >15 were classified as
432 potentially highly repetitive sequences and removed from the in-house list. sRNAs not appearing
433 in the majority of the 385 strains were also removed. A database containing the *S. pyogenes*, *S.*
434 *mutans*, *S. suis*, *S. mitis*, *S. oralis*, and *S. gordonii* genomes, created with makeblastdb, was used
435 to search for sRNA sequence alignments using blastn with the task parameter set to blastn.
436 Synteny of the sRNAs was evaluated by comparing the 1000 nt upstream and downstream of the
437 sequences to those in TIGR4. The 1000 nt on either end of the sRNA sequences for the other 384
438 strains were compared base-wise to obtain sequence identities. The average sequence identities,
439 both up and downstream, were averaged across all 384 strains, and if either average sequence
440 identity, up or downstream, is ≥75% then it was concluded that the synteny of the sRNA is
441 preserved.
442
443 **sRNA target prediction**

IntaRNA version 3.3.2, sRNARFTarget, and TargetRNA3 were used to predict targets for the 58
sRNA candidates. CopraRNA version 2.1.4 was only used to predict targets for 3 sRNAs with
significant alignments in *S. suis*, *S. pyogenes*, *S. mutans, S. mitis, S. oralis,* and *S. gordonii*.
Predictions for each sRNA were made in 6 different *S. pneumoniae* strains consisting of TIGR4
(NC_003028.3), D39 (NC_008533.2), TVO_Taiwan19F-14, TVO_1901920, TVO_1901934,
TVO_1902277 (Rosconi et al., 2022). IntaRNA was run with the IntaRNAsTar personality, the
number of predictions set to 5, and otherwise default parameters. sRNARFTarget was run with
the provided Docker container. sRNARFTarget requires a user input transcriptome, so an in-
house Python script created two transcriptome files for each of the six *S. pneumoniae* strains
using the coordinates from the respective GenBank files and retrieving the sequences from the
genome. The two transcriptomes are defined by the exact gene coordinates and coordinates
adjusted to include 100 nucleotides upstream of the start codon and 300 nucleotides downstream
of the stop codon or until the next gene's start codon, whichever comes first. TargetRNA3 was
run with the probability threshold lowered to 0.25 and otherwise default parameters. Note, the
six genomes were first added to the local user database by providing the accession identifier.
CopraRNA was run with default parameters in *S. pneumoniae* (NC_003028), *S. suis*
(NC_012926), *S. pyogenes* (NZ_LS483338), *S. mutans* (NZ_CP044221), *S. mitis*
(NZ_CP012646), *S. oralis* (NZ_LR134336), and *S. gordonii* (NZ_CP077224).
462
sRNA target analysis
An in-house Python script retrieved the sequences including 25 nucleotides upstream and
downstream of the mRNA interacting sequence. The structures of these sequences were predicted
using RNAfold version 2.6.4 with default parameters. If the sRNA interacting sequence overlaps
with a structured region of the mRNA, then the interaction was labeled as interacting with a
structured region. Only the MPTs of each sRNA in the six strains predicted by IntaRNA were
analyzed. To determine if the sRNAs are acting in regulons we searched the target loci against
the RegPrecise database (https://regprecise.lbl.gov/index.jsp). Only the MPTs and probable
targets predicted by TargetRNA3 in TIGR4 were searched seeing as TIGR4 is the only *S.*
pneumoniae strain for which the database contains information. The expression of sRNA
sequences was confirmed by checking if the transcription initiation and termination sites were
present in 5' and 3' RNA-end sequencing data (Warrier et al., 2018).
475

Additional Data Files

Additional_Data_1_source-sRNAs.xlsx – initial list of putative sRNAs from three studies
Additional_Data_2_In-house-sRNAs.xlsx – final in-house list after deduplication (Fig. 1C)
Additional_Data_3_CopraRNA.xlsx – CopraRNA target predictions
Additional_Data_4_IntaRNA.xlsx – IntaRNA target predictions
Additional_Data_5_sRNARFTarget-Transcriptome-Exact.xlsx – sRNARFT target predictions
(coding sequences only)
Additional_Data_6_sRNARFTarget-Transcriptome-100-300.xlsx – sRNARFT target predictions
(coding sequences and flanking regions)
Additional_Data_7_TargetRNA3.xlsx – TargetRNA3 target predictions
486

Supplemental Files

488 Eichelman_SupplementalFile1.pdf – Supplemental Figures and Tables

489   Eichelman_SupplementalFile2_python_scripts.zip – In-house python scripts
490

Acknowledgments

492   MCE was partially funded by a Boston College Undergraduate Research Fellowship and by
493   grants R21AI181123 and R01GM134259 from the US National Institutes of Health. We would
494   like to thank Quinlan Furumo for careful copyediting of the manuscript.
495

496   MCE roles: data curation, investigation, visualization, writing – original draft
497   MMM roles: conceptualization, data curation, funding acquisition, supervision, writing – review
498   and editing
499

References

Acebo, Paloma, Antonio J. Martin-Galiano, Sara Navarro, Ángel Zaballos, and Mónica Amblar.
2012. "Identification of 88 Regulatory Small RNAs in the TIGR4 Strain of the Human
Pathogen *Streptococcus Pneumoniae*." *RNA* 18 (3): 530–46.
https://doi.org/10.1261/rna.027359.111.
Altschul, Stephen F., Warren Gish, Webb Miller, Eugene W. Myers, and David J. Lipman. 1990.
"Basic Local Alignment Search Tool." *Journal of Molecular Biology* 215 (3): 403–10.
https://doi.org/10.1016/S0022-2836(05)80360-2.
Aoyama, Jordan J., and Gisela Storz. 2023. "Two for One: Regulatory RNAs That Encode Small
Proteins." *Trends in Biochemical Sciences* 48 (12): 1035–43.
https://doi.org/10.1016/j.tibs.2023.09.002.
Arini, Achille, Marcel P. Keller, and Werner Arber. 1997. "An Antisense RNA in IS30 Regulates
the Translational Expression of the Transposase." *Biological Chemistry* 378 (12).
https://doi.org/10.1515/bchm.1997.378.12.1421.
Babina, Arianne M., Mark W. Soo, Yang Fu, and Michelle M. Meyer. 2015. "An S6:S18
Complex Inhibits Translation of *E. Coli rpsF*." *RNA* 21 (12): 2039–46.
https://doi.org/10.1261/rna.049544.115.
Busch, Anke, Andreas S. Richter, and Rolf Backofen. 2008. "IntaRNA: Efficient Prediction of
Bacterial sRNA Targets Incorporating Target Site Accessibility and Seed Regions."
Bioinformatics 24 (24): 2849–56. https://doi.org/10.1093/bioinformatics/btn544.
CDC, National Center for Emerging Zoonotic and Infectious Diseases (U.S.), National Center
for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention (U.S.), and National Center for
Immunization and Respiratory Diseases (U.S.). 2013. *Antibiotic Resistance Threats in the
United States, 2013*. https://stacks.cdc.gov/view/cdc/20705/cdc_20705_DS1.pdf.
Christopoulou, Niki, and Sander Granneman. 2022. "The Role of RNA-Binding Proteins in
Mediating Adaptive Responses in Gram-Positive Bacteria." *The FEBS Journal* 289 (7):
1746–64. https://doi.org/10.1111/febs.15810.
Cremers, Amelieke J. H., Fredrick M. Mobegi, Marien I. De Jonge, Sacha A. F. T. Van Hijum,
Jacques F. Meis, Peter W. M. Hermans, Gerben Ferwerda, Stephen D. Bentley, and Aldert L.
Zomer. 2015. "The Post-Vaccine Microevolution of Invasive Streptococcus Pneumoniae."
Scientific Reports 5 (1): 14952. https://doi.org/10.1038/srep14952.
Damjanovic, Marlen, Arun S. Kharat, Alice Eberhardt, Alexander Tomasz, and Waldemar
Vollmer. 2007. "The Essential *tacF* Gene Is Responsible for the Choline-Dependent Growth

Phenotype of *Streptococcus Pneumoniae*." *Journal of Bacteriology* 189 (19): 7105–11. https://doi.org/10.1128/JB.00681-07.

Danecek, Petr, James K Bonfield, Jennifer Liddle, John Marshall, Valeriu Ohan, Martin O Pollard, Andrew Whitwham, et al. 2021. "Twelve Years of SAMtools and BCFtools." *GigaScience* 10 (2): giab008. https://doi.org/10.1093/gigascience/giab008.

De Lay, Nicholas R., Nidhi Verma, Dhriti Sinha, Abigail Garrett, Maximillian K. Osterberg, Daisy Porter, Spencer Reiling, David P. Giedroc, and Malcolm E. Winkler. 2024. "The Five Homologous CiaR-Controlled Ccn sRNAs of Streptococcus Pneumoniae Modulate Zn-Resistance." Edited by Rachel M. McLoughlin. *PLOS Pathogens* 20 (10): e1012165. https://doi.org/10.1371/journal.ppat.1012165.

Ellis, Michael J., Ryan S. Trussler, and David B. Haniford. 2015. "A *Cis* -Encoded sRNA, Hfq and mRNA Secondary Structure Act Independently to Suppress IS *200* Transposition." *Nucleic Acids Research* 43 (13): 6511–27. https://doi.org/10.1093/nar/gkv584.

Felden, Brice, and Yoann Augagneur. 2021. "Diversity and Versatility in Small RNA-Mediated Regulation in Bacterial Pathogens." *Frontiers in Microbiology* 12 (August):719977. https://doi.org/10.3389/fmicb.2021.719977.

Fu, Yang, Kaila Deiorio-Haggar, Jon Anthony, and Michelle M. Meyer. 2013. "Most RNAs Regulating Ribosomal Protein Biosynthesis in Escherichia Coli Are Narrowly Distributed to Gammaproteobacteria." *Nucleic Acids Research* 41 (6): 3491–3503. https://doi.org/10.1093/nar/gkt055.

Furumo, Quinlan, and Michelle M. Meyer. 2024. "PIPETS: A Statistically Informed, Gene-Annotation Agnostic Analysis Method to Study Bacterial Termination Using 3′-End Sequencing." *BMC Bioinformatics* 25 (1): 363. https://doi.org/10.1186/s12859-024-05982-5.

GBD 2016 Lower Respiratory Infections Collaborators. 2018. "Estimates of the Global, Regional, and National Morbidity, Mortality, and Aetiologies of Lower Respiratory Infections in 195 Countries, 1990-2016: A Systematic Analysis for the Global Burden of Disease Study 2016." *The Lancet. Infectious Diseases* 18 (11): 1191–1210. https://doi.org/10.1016/S1473-3099(18)30310-4.

Giammarinaro, Philippe, and James C. Paton. 2002. "Role of RegM, a Homologue of the Catabolite Repressor Protein CcpA, in the Virulence of *Streptococcus Pneumoniae*." *Infection and Immunity* 70 (10): 5454–61. https://doi.org/10.1128/IAI.70.10.5454-5461.2002.

Halfmann, Alexander, Márta Kovács, Regine Hakenbeck, and Reinhold Brückner. 2007. "Identification of the Genes Directly Controlled by the Response Regulator CiaR in *Streptococcus Pneumoniae* : Five out of 15 Promoters Drive Expression of Small Non-coding RNAs." *Molecular Microbiology* 66 (1): 110–26. https://doi.org/10.1111/j.1365-2958.2007.05900.x.

Hör, Jens, Geneviève Garriss, Silvia Di Giorgio, Lisa-Marie Hack, Jens T. Vanselow, Konrad U. Förstner, Andreas Schlosser, Birgitta Henriques-Normark, and Jörg Vogel. 2020. "Grad-Seq in a Gram-Positive Bacterium Reveals Exonucleolytic sRNA Activation in Competence Control." *The EMBO Journal* 39 (9): e103852. https://doi.org/10.15252/embj.2019103852.

Huang, Susan S., Kristen M. Johnson, G. Thomas Ray, Peter Wroe, Tracy A. Lieu, Matthew R. Moore, Elizabeth R. Zell, et al. 2011. "Healthcare Utilization and Cost of Pneumococcal Disease in the United States." *Vaccine* 29 (18): 3398–3412. https://doi.org/10.1016/j.vaccine.2011.02.088.

Iyer, Ramkumar, Nitin S. Baliga, and Andrew Camilli. 2005. "Catabolite Control Protein A (CcpA) Contributes to Virulence and Regulation of Sugar Metabolism in *Streptococcus Pneumoniae*." *Journal of Bacteriology* 187 (24): 8340–49. https://doi.org/10.1128/JB.187.24.8340-8349.2005.

Jabbour, Nancy, and Marie-Frédérique Lartigue. 2021. "An Inventory of CiaR-Dependent Small Regulatory RNAs in Streptococci." *Frontiers in Microbiology* 12 (May):669396. https://doi.org/10.3389/fmicb.2021.669396.

Janssen, Axel B, Paddy S Gibson, Afonso M Bravo, Vincent de Bakker, Jelle Slager, and Jan-Willem Veening. 2024. "PneumoBrowse 2: An Integrated Visual Platform for Curated Genome Annotation and Multiomics Data Analysis of *Streptococcus Pneumoniae*." *Nucleic Acids Research*, October, gkae923. https://doi.org/10.1093/nar/gkae923.

Laczkovich, Irina, Kyle Mangano, Xinhao Shao, Adam J. Hockenberry, Yu Gao, Alexander Mankin, Nora Vázquez-Laslop, and Michael J. Federle. 2022. "Discovery of Unannotated Small Open Reading Frames in Streptococcus Pneumoniae D39 Involved in Quorum Sensing and Virulence Using Ribosome Profiling." Edited by N. Luisa Hiller. *mBio* 13 (4): e01247-22. https://doi.org/10.1128/mbio.01247-22.

Li, Wuju, Xiaomin Ying, Qixuan Lu, and Linxi Chen. 2012. "Predicting sRNAs and Their Targets in Bacteria." *Genomics, Proteomics & Bioinformatics* 10 (5): 276–84. https://doi.org/10.1016/j.gpb.2012.09.004.

Lorenz, Ronny, Stephan H Bernhart, Christian Höner Zu Siederdissen, Hakim Tafer, Christoph Flamm, Peter F Stadler, and Ivo L Hofacker. 2011. "ViennaRNA Package 2.0." *Algorithms for Molecular Biology* 6 (1): 26. https://doi.org/10.1186/1748-7188-6-26.

Mann, Beth, Tim Van Opijnen, Jianmin Wang, Caroline Obert, Yong-Dong Wang, Robert Carter, Daniel J. McGoldrick, et al. 2012. "Control of Virulence by Small RNAs in Streptococcus Pneumoniae." Edited by Pascale Cossart. *PLoS Pathogens* 8 (7): e1002788. https://doi.org/10.1371/journal.ppat.1002788.

Mann, Martin, Patrick R. Wright, and Rolf Backofen. 2017. "IntaRNA 2.0: Enhanced and Customizable Prediction of RNA–RNA Interactions." *Nucleic Acids Research* 45 (W1): W435–39. https://doi.org/10.1093/nar/gkx279.

Martin, Julia E., My T. Le, Nabin Bhattarai, Daiana A. Capdevila, Jiangchuan Shen, Malcolm E. Winkler, and David P. Giedroc. 2019. "A Mn-Sensing Riboswitch Activates Expression of a Mn2+/Ca2+ ATPase Transporter in Streptococcus." *Nucleic Acids Research* 47 (13): 6885–99. https://doi.org/10.1093/nar/gkz494.

Marx, Patrick, Michael Nuhn, Martá Kovács, Regine Hakenbeck, and Reinhold Brückner. 2010. "Identification of Genes for Small Non-Coding RNAs That Belong to the Regulon of the Two-Component Regulatory System CiaRH in Streptococcus." *BMC Genomics* 11 (1): 661. https://doi.org/10.1186/1471-2164-11-661.

Mascher, Thorsten, Manuel Heintz, Dorothea Zähner, Michelle Merai, and Regine Hakenbeck. 2006. "The CiaRH System of *Streptococcus Pneumoniae* Prevents Lysis during Stress Induced by Treatment with Cell Wall Inhibitors and by Mutations in *Pbp2x* Involved in β-Lactam Resistance." *Journal of Bacteriology* 188 (5): 1959–68. https://doi.org/10.1128/JB.188.5.1959-1968.2006.

Naskulwar, Kratika, and Lourdes Peña-Castillo. 2022. "sRNARFTarget: A Fast Machine-Learning-Based Approach for Transcriptome-Wide sRNA Target Prediction." *RNA Biology* 19 (1): 44–54. https://doi.org/10.1080/15476286.2021.2012058.

Olejniczak, Mikolaj, Xiaofang Jiang, Maciej M. Basczok, and Gisela Storz. 2022. "KH Domain Proteins: Another Family of Bacterial RNA Matchmakers?" *Molecular Microbiology* 117 (1): 10–19. https://doi.org/10.1111/mmi.14842.

Papenfort, Kai, and Gisela Storz. 2024. "Insights into Bacterial Metabolism from Small RNAs." *Cell Chemical Biology* 31 (9): 1571–77. https://doi.org/10.1016/j.chembiol.2024.07.002.

Papenfort, Kai, and Carin K. Vanderpool. 2015. "Target Activation by Regulatory RNAs in Bacteria." *FEMS Microbiology Reviews* 39 (3): 362–78. https://doi.org/10.1093/femsre/fuv016.

Patenge, Nadja, Tomas Fiedler, and Bernd Kreikemeyer. 2012. "Common Regulators of Virulence in Streptococci." In *Host-Pathogen Interactions in Streptococcal Diseases*, edited by G. Singh Chhatwal, 368:111–53. Current Topics in Microbiology and Immunology. Berlin, Heidelberg: Springer Berlin Heidelberg. https://doi.org/10.1007/82_2012_295.

Peer, Asaf, and Hanah Margalit. 2014. "Evolutionary Patterns of *Escherichia Coli* Small RNAs and Their Regulatory Interactions." *RNA* 20 (7): 994–1003. https://doi.org/10.1261/rna.043133.113.

Raden, Martin, Syed M Ali, Omer S Alkhnbashi, Anke Busch, Fabrizio Costa, Jason A Davis, Florian Eggenhofer, et al. 2018. "Freiburg RNA Tools: A Central Online Resource for RNA-Focused Research and Teaching." *Nucleic Acids Research* 46 (W1): W25–29. https://doi.org/10.1093/nar/gky329.

Raina, Medha, Alisa King, Colleen Bianco, and Carin K. Vanderpool. 2018. "Dual-Function RNAs." *Microbiology Spectrum* 6 (5). https://doi.org/10.1128/microbiolspec.RWR-0032-2018.

Rosconi, Federico, Emily Rudmann, Jien Li, Defne Surujon, Jon Anthony, Matthew Frank, Dakota S. Jones, et al. 2022. "A Bacterial Pan-Genome Makes Gene Essentiality Strain-Dependent and Evolvable." *Nature Microbiology* 7 (10): 1580–92. https://doi.org/10.1038/s41564-022-01208-7.

Schnorpfeil, Anke, Miriam Kranz, Martá Kovács, Christian Kirsch, Julia Gartmann, Ines Brunner, Sabrina Bittmann, and Reinhold Brückner. 2013. "Target Evaluation of the Non-coding CSRNAs Reveals a Link of the Two-component Regulatory System CIARH to Competence Control in *S Treptococcus Pneumoniae* R 6." *Molecular Microbiology* 89 (2): 334–49. https://doi.org/10.1111/mmi.12277.

Sharples, Gary J., Stuart M. Ingleston, and Robert G. Lloyd. 1999. "Holliday Junction Processing in Bacteria: Insights from the Evolutionary Conservation of RuvABC, RecG, and RusA." *Journal of Bacteriology* 181 (18): 5543–50. https://doi.org/10.1128/JB.181.18.5543-5550.1999.

Simons, Robert W., and Nancy Kleckner. 1983. "Translational Control of IS10 Transposition." *Cell* 34 (2): 683–91. https://doi.org/10.1016/0092-8674(83)90401-4.

Sinha, Dhriti, Kurt Zimmer, Todd A. Cameron, Douglas B. Rusch, Malcolm E. Winkler, and Nicholas R. De Lay. 2019. "Redefining the Small Regulatory RNA Transcriptome in Streptococcus Pneumoniae Serotype 2 Strain D39." Edited by Victor J. DiRita. *Journal of Bacteriology* 201 (14). https://doi.org/10.1128/JB.00764-18.

Tieng, Francis Yew Fu, Muhammad-Redha Abdullah-Zawawi, Nur Alyaa Afifah Md Shahri, Zeti-Azura Mohamed-Hussein, Learn-Han Lee, and Nurul-Syakima Ab Mutalib. 2023. "A Hitchhiker's Guide to RNA–RNA Structure and Interaction Prediction Tools." *Briefings in Bioinformatics* 25 (1): bbad421. https://doi.org/10.1093/bib/bbad421.

Tjaden, Brian. 2023. "TargetRNA3: Predicting Prokaryotic RNA Regulatory Targets with Machine Learning." *Genome Biology* 24 (1): 276. https://doi.org/10.1186/s13059-023-03117-2.

Tsui, Ho-Ching Tiffany, Dhriti Mukherjee, Valerie A. Ray, Lok-To Sham, Andrew L. Feig, and Malcolm E. Winkler. 2010. "Identification and Characterization of Noncoding Small RNAs in *Streptococcus Pneumoniae* Serotype 2 Strain D39." *Journal of Bacteriology* 192 (1): 264–79. https://doi.org/10.1128/JB.01204-09.

Warrier, Indu, Nikhil Ram-Mohan, Zeyu Zhu, Ariana Hazery, Haley Echlin, Jason Rosch, Michelle M. Meyer, and Tim Van Opijnen. 2018. "The Transcriptional Landscape of Streptococcus Pneumoniae TIGR4 Reveals a Complex Operon Architecture and Abundant Riboregulation Critical for Growth and Virulence." Edited by Carlos Javier Orihuela. *PLOS Pathogens* 14 (12): e1007461. https://doi.org/10.1371/journal.ppat.1007461.

Wright, Patrick R., Jens Georg, Martin Mann, Dragos A. Sorescu, Andreas S. Richter, Steffen Lott, Robert Kleinkauf, Wolfgang R. Hess, and Rolf Backofen. 2014. "CopraRNA and IntaRNA: Predicting Small RNA Targets, Networks and Interaction Domains." *Nucleic Acids Research* 42 (W1): W119–23. https://doi.org/10.1093/nar/gku359.

Wright, Patrick R., Andreas S. Richter, Kai Papenfort, Martin Mann, Jörg Vogel, Wolfgang R. Hess, Rolf Backofen, and Jens Georg. 2013. "Comparative Genomics Boosts Target Prediction for Bacterial Small RNAs." *Proceedings of the National Academy of Sciences* 110 (37). https://doi.org/10.1073/pnas.1303248110.

Zhang, Aixia, Karen M. Wassarman, Carsten Rosenow, Brian C. Tjaden, Gisela Storz, and Susan Gottesman. 2003. "Global Analysis of Small RNA and mRNA Targets of Hfq." *Molecular Microbiology* 50 (4): 1111–24. https://doi.org/10.1046/j.1365-2958.2003.03734.x.

Zorgani, Mohamed A., Roland Quentin, and Marie-Frédérique Lartigue. 2016. "Regulatory RNAs in the Less Studied Streptococcal Species: From Nomenclature to Identification." *Frontiers in Microbiology* 7 (July). https://doi.org/10.3389/fmicb.2016.01161.