# Unbiased profiling of multipotency landscapes reveals spatial modulators of clonal fate biases

Alek G. Erickson[1]†, Sergey Isaev[2]†, Artem Artemov[2], Jingyan He[3], Bettina Semsch[3], Aliia Murtazina[1], Jia Sun[4], Katrin Mangold[3], Anthi Chalou[1], Jonas Frisen[3], Michael Ratz[3], Emma Andersson[3], Peter V. Kharchenko[5,6] * and Igor Adameyko[1,2] *

† - These authors contributed equally to this work
* - Corresponding authors. Email: igor.adameyko@ki.se and peter.kharchenko@post.harvard.edu

**Affiliations:**
[1] Department of Physiology and Pharmacology, Karolinska Institutet; 17177 Stockholm, Sweden.
[2] Department of Neuroimmunology, Center for Brain Research, Medical University of Vienna; 1090 Vienna, Austria.
[3] Department of Cell and Molecular Biology, Karolinska Institutet; 17177 Stockholm, Sweden.
[4] Department of Comparative Medicine, Karolinska Institutet; 17177 Stockholm, Sweden.
[5] San Diego Institute of Science, Altos Labs; San Diego, CA, USA.
[6] Department of Biomedical Informatics, Harvard Medical School; Boston, MA 02115, USA.

**Abstract:** Embryogenesis is commonly viewed through a tree model of cell differentiation, which fails to capture the spatiotemporal modulation of cell multipotency underlying morphogenesis. In this study we profile the multipotency landscape of the embryo, using *in vivo* single-cell clonal lineage tracing of mouse embryos traced from neurulation until mid-gestation, combined with a machine learning tool that categorizes individual clones into lineages based on shared transcriptional context. This revealed a previously unrecognized continuous, embryo-wide gradient of clonal fate biases, in which anatomical position and clonal composition are mutually predictive. Comparing clonal lineages revealed gene regulatory networks underlying the dynamic biasing of cells towards specific fates by spatial transcription factor programs. However, mosaic combinatorial perturbations targeting the Hedgehog pathway generated clones in which positional identity was mismatched with clonal composition, demonstrating that extrinsic signals can override the axial patterning system underlying clonal fate biases. Altogether, our work demonstrates an effective practical approach for dissecting mechanisms of lineage specification and has implications for stem cell engineering.

**Main Text:**

*Introduction*

Stochasticity during vertebrate embryonic development poses a challenge for understanding how the organism robustly emerges from a single cell, through an orderly sequence of cell fate decisions and mitotic divisions (Fig. 1a) [1]. Each tissue, organ, and body part, is formed from a defined set of progenitor cells that generate progressively elaborate clonal structures, varying in composition according to their unique lineage histories, each contributing just a part of the overall repertoire of the body's cell types (Fig. 1b). Consequently, the classic model of cell differentiation as a "tree-of-fates" represents a kind of aggregation of many individual clones at a given location. However, cell fate decisions and their underlying tree structure are thought to be highly influenced by anatomical position within the developing organism (Fig. 1c). Thus, a deeper understanding of organismal development requires massively parallel analysis of individual clonal fates in relation to instructive signals, timing, and locations.

Lineage tracing techniques have long been at the core of developmental biology, providing insights for identifying progenitor populations and their differentiation potentials [2]. Earlier clonal tracing approaches, such as those that rely on dye injections or multiplexed imaging of fluorescent proteins, have generally been limited in scale and throughput. Recent advances in multiplexed genetic tracing methods, such as Brainbow [3], scGESTALT [4], PolyLoxExpress [5], CARLIN/DARLIN [6, 7], and LINNAEUS [8], enabled construction of high-resolution lineage trees. These technologies have refined our understanding of cell differentiation pathways by revealing how cell progenitor fates vary systematically across biological contexts. Variability in cell fate can result from both intrinsic cellular biases and external environmental influences, though the molecular logic driving these variations remain incompletely understood, especially when it comes to understanding how the proportions of cell types are established [9, 10]. Quantifying the spatiotemporal variation of clonal fate biases, and identifying potential biasing factors, are important steps towards elucidating the genetic and epigenetic mechanisms that coordinate cell destinies in vivo.

Despite recent advancements in combining highly multiplexed tracing with analysis of cellular state, a few practical challenges remain. One such difficulty is sparse coverage, where only a small fraction of cells within a tissue or organism are effectively labeled and recovered, making it difficult to reconstruct comprehensive lineage trees. More fundamentally, the measurements of the cellular state that accompany tracing information are performed at the final timepoint. Consequently, molecular differences associated with fate decisions cannot be observed directly and must be inferred under additional assumptions, such as ergodicity of the differentiation process or persistence of past progenitor signatures at the final time point. The asynchronous nature of embryonic development, for instance in the case of the neural crest [11, 12] and neuro-mesodermal progenitor cells [13], represents a partial solution to these challenges by enabling faithful differentiation trajectories to be constructed from lineage-aware gene expression measurements.

While traditionally, lineage tracing has been used to establishing the range of cell types that can arise from progenitor populations (i.e. potency), here we focus on quantitatively mapping the relative frequencies of transcriptional states occurring among the progeny of a cell (clonal fate

biases). Even for progenitor cells with identical lineage potency, such frequencies will vary due to
the influence of external signals and stochastic differences in the progenitor state. Our aim is to
characterize such fate variation and identify factors biasing fate realization in embryonic
development. We revisit the framework of clonal analysis using a cellular barcoding approach,
combining highly multiplexed sparse tracing with unsupervised learning techniques and analysis
of expression states, to study the diversity of fate biases occurring in ectodermal and mesodermal
clones during mammalian embryonic development. This approach captured gradients of
continuous clonal fate variation along anatomical axes as well as sequential fate restrictions over
time, enabling computational inference of the regulatory gene programs likely responsible for
modulation of fates in neural and mesodermal progenitor cells. This system of axial clonal
patterning was also susceptible to disruption by multiplexed *in vivo* signaling receptor
perturbations, suggesting that positional regulatory factors cooperate with extrinsic signaling
during to ensure the correct clonal proportionality of cell types during organogenesis.
106
107
108   *Profiling of clonal behavior in mouse embryos*
109
To reconstruct joint patterns of transcriptional states and cell lineages in post-neurulation murine
development, we performed ultrasound-guided nano-injections [14] into amniotic fluid to
transduce mesoderm and ectoderm with heritable, transcribed genetic TREX barcodes [15, 16]
(Fig. 1d). The injections were carried out once or twice serially between days E7.5 and E8.5 (Fig.
1e-f, Extended Data Fig. 1) to capture rapid developmental shifts during embryonic development.
Trunk and cranial segments were harvested separately at E13.5 when most definitive cell types
manifest, and labeled cells were analyzed by scRNA-seq for linking clonal information to each
transcriptome (Fig. 1g-h, Extended Data Fig. 2). We optimized the study design to maximize the
fraction of cells captured from each individual clone. To accomplish this, we injected embryos
with low amounts of virus (in almost all cases 46-206 nL) such that only 10-40k cells would be
labeled in each harvested body segment – an amount that could be taken through downstream
processing steps without significant additional losses.
122
Our microinjection approach predominantly targets ectoderm [14-16], as well as some mesoderm.
This is explained by the fact that the ultrasound guided injections shower the exterior of the early
neurula with barcoded lentiviral particles, and the exposed cells residing in outermost positions
have the highest chance to become labeled [14]. As expected, in our experiments, we did not
recover traced endodermal derivatives. Following transduction of the embryos, retroviral
integration in dividing cells takes approximately eight hours [17]. Two days after the injection we
saw the formation of large clones by whole mount analysis of embryos injected with low volumes
of viral particles (Extended Data Fig. 3), and broad labeling thereafter (Extended Data Fig. 4),
indicating that the tracer system activates swiftly. Thus, this method facilitates rapid, accurate, and
broad lineage tracing of early ectodermal cells, as well as some mesodermal cells and axis-
extending neuromesodermal progenitors. Single-cell RNA-seq analysis of eight control embryos
(4 cranial, 7 trunk sections), recovered a total of 159,744 clonally barcoded cells. Specifically, we
recovered 4411 E7.5-traced clones and 6661 E8.5-traced clones. Of those, 3442 E7.5-traced clones
and 4722 E8.5-traced clones contained at least two cells. These informative clones encompassed
49% of all sequenced barcoded cells.
138

As expected, multicellular clones were distributed across the major embryonic ectodermal and
mesodermal derivatives, including keratinocytes of the skin, various mesenchymal cells, and
neuroglial cells of the central and peripheral nervous system (Fig. 1i-j, Extended Data Fig. 5 and
Extended Data Fig. 6). Most clones contained similar cells belonging to only one germ layer
derivative (Fig. 1k-l; for cranial, see Extended Data Fig. 7). Multipotent clones spanning more
than one major lineage were more abundant in the earlier developmental timepoint (1.7% in E8.5
vs. 5.2% in E7.5) indicating that early fate restrictions are already imposed in ectodermal and
mesodermal cell lineages prior to neurulation (Fig. 1l). The patterns of clonal co-occurrence
grouped similar cell types, recovering the expected consensus cell type lineage phylogeny (Fig.
1m and Extended Data Fig. 8). Reconstruction of such lineage structure from fixed barcode data,
however, is known to be challenging [18]. The resulting phylogeny would represent a population
average of cell fate decision structure, whereas the clones contributing to any given cell type can
vary greatly in their overall lineage distribution (Fig. 1b, k). We therefore reasoned that a new
computational approach is needed to analyze the processes underlying the substantial diversity in
clonal fate distributions.
154
155   *Clone2vec recapitulates spatiotemporal patterns of multipotency in the embryo*
156
To investigate variability among numerous sparsely sampled clonal populations, we developed a
computational method, called clone2vec, which adapts word2vec [19] word embeddings to lineage
tracing information (see Materials and Supplementary Note: Formal Description of clone2vec).
Clone2vec represents individual clones in a low-dimensional space that captures the variation of
clonal fates (Fig. 1n-o). In such an embedding, neighboring clones exhibit similar cell fate
distributions across the continuous expression space (Fig. 1p, Extended Data Fig. 9). By
establishing this metric space, clone2vec enables exploration of recurrent clonal patterns and
subsequent analysis of features associated with them. We characterized the performance of the
clone2vec method using simulated data (see Supplementary Note: Simulations), as well as
published datasets. We found that clone2vec embedding effectively reconstructs both discrete
clusters of clones, as well as patterns of continuous clonal variation within clusters. Importantly,
clone2vec performance remained robust under increasingly sparse sampling of cells
(Supplementary Note: Simulations). We compared clone2vec analysis with the more traditional
approach of representing clones in the space of cell types, where each clone is encoded as a vector
of cell type abundances, followed by PCA dimensionality reduction. This embedding results in
relatively poor reconstruction of the original clusters, with the advantages of clone2vec becoming
more evident under sparser clonal coverage (see Supplementary Note: Simulations). Also, much
like word embeddings, clonal embeddings are amenable to vector math operations (see
Supplementary Note: Simulations) – for instance, multipotent neuromesodermal progenitor
(NMP) clones are the vector sum of neuronal clones and mesodermal clones. Clone2vec re-
analysis of published hematopoietic lineage tracing data [20] was able to recover previously
described fate patterns with their associated genes, additionally uncovering more subtle continuous
axes of variation (see Supplementary Note: Real Data). These features make clone2vec a useful
tool for learning patterns of clonal behavior in the embryo.
181
Clone2vec analysis of our dataset revealed three major discrete compartments corresponding to
derivatives of the neural ectoderm, non-neural ectoderm, and mesoderm (see Data Availability for
an interactive visualization) and facilitated the assignment of clones into clonal clusters within

each compartment (Fig. 2a). The general structure of the clonal embedding reflected the observed
restricted nature of mesoderm, non-neural ectoderm, and neural ectoderm clones. However, two
clonal clusters exhibited neural-mesenchymal multipotency (Fig 2b, dotted box). Surveying the
specific combinations of epidermal, neuronal, and/or mesenchymal cell subtypes found in each
clonal cluster, revealed that nearby clusters often share many cell types, but differ chiefly in their
proportionality in which they are found, especially in mesenchymal clones (Fig. 2b). Reasoning
that adjacent clonal clusters represent points along a continuum of clonal behavior, we performed
principal components analysis (PCA) from discrete versus contiguous clonal clusters (Extended
Data Fig. 10a-b) and examined gene expression and cell fate distributions across a constructed
trajectory of clones following the top principal components (Extended Data Fig. 10c-d). The clones
in contiguous clusters exhibited a continuous gradient of gene expression (Extended Data Fig. 10e)
reflecting gradual transitions across mesenchymal cell types (Extended Data Fig. 10f-j).
197
To study the molecular drivers underlying continuous variation in clonal composition, we devised
a machine learning approach to identify genes whose expression, either in a specific cell type or
across all cell types, was strongly associated with clonal fate distributions. We first built a gradient
boosting model to predict the position of clones in the embedding based on cell type-stratified gene
expression in cells belonging to each clone (see Methods). We then scored gene importance by
quantifying each gene's predictive power using Shapley's values (measure of feature importance,
see Methods). Next, to exclude the most obvious predictors (cell type markers) we added an
additional categorical feature (cell type identity) to account for cell type-dependent effects.
Applying this approach to the trunk mesodermal clones revealed that many Hox transcription
factors were among the most important genes to predict clonal composition, suggesting that the
observed fate bias gradient was driven by clones arising from somites at different positions along
the body axis (Fig. 2c). We constructed a 'Hox score' to locate clones by aggregating their
expression levels of Hox genes (Fig. 2d, Extended Data Fig. 11a-d). Clone position along the
observed clone2vec fate gradient, and clone Hox score, were linearly related (Extended Data Fig.
11c). To validate that the clonal embedding captured the anterior-posterior axis of the embryo, we
separately dissected lower and middle trunk prior to sorting, confirming a striking difference in
their representation along the mesenchymal domain embedding (Fig 2d). We exploited this metric
to uncover several genes (beyond the Hox code) that are highly correlated with the body axis
(Extended Data Fig. 11e-h). This linear relationship was driven primarily by somite derived clones,
not other mesenchymal cells (Extended Data Fig. 11i-l) nor the cell cycle (Extended Data Fig.
11m). This analysis suggests that somite clonal composition and body position are mutually
predictive, and that knowledge of a few genes can reconstruct both, even though the underlying
cell types occur along the entire body axis. Additionally, the data suggests that application of these
models can help reveal new positional information genes.
222
To identify early biases during progenitor cell differentiation, we examined the predictors of
clone2vec embedding position of clones based exclusively on the expression profile of progenitor
cells (Extended Data Fig. 12a-b, e, please see embeddings of both neuronal and mesodermal
lineages). To annotate the effect of gene expression on fate distribution, we built an additional
model that predicts cell type composition of the clone based on the expression in progenitor cells
(see Methods and Extended Data Fig. 12c-d, f-g) and outlined programs biasing towards one or
another fate. Applying this approach revealed, for example, that during mesenchymal progenitor
cell differentiation, the most powerful predictor appeared to be the *Tbx3* gene rather than one of

the Hox genes (Extended Data Fig. 12d), showing that gene expression correlates of clonal
behavior are distinct when only considering progenitor cells, versus all cells (including
differentiated progeny). The same strategy was used on a publicly available dataset of *in vitro*
hematopoiesis (see Supplementary Note: Real Data) illustrating that this approach allows for
systematic analysis of gene expression associations with clonal behaviour.
236
Because the embryo develops from head to tail, caudal cells should exhibit greater developmental
potential than their anterior counterparts at a given moment. Neuromesodermal progenitors
(NMPs, clonal cluster 15 and 16), which are a cell population that contributes to secondary
neurulation and somitogenesis during axial extension, were found near the tail [13, 21]. We noted
that, while the posterior-most clones corresponding to NMPs exhibited the greatest multipotency
(Fig. 2e), levels of clone transcriptional diversity (measured via Shannon entropy) did not always
correspond with body position (compare feature plots of mesenchymal clonal embeddings in Fig.
2d versus 2e). For instance, tail clusters 15, 16, and 7 had rather similar Hox scores (bracket, Fig.
2d), but the most multipotent NMP cluster 16 was enriched with E7.5-labeled clones (Fig. 2f),
while other tail clusters (15, 7) contained a mix of E7.5 and E8.5 clones, suggesting a temporal,
rather than spatial, explanation for their divergent profiles.
248
To visualize the gradual posteriorizing of NMP-competent clones over time, we serially sectioned
littermate embryos double-injected with EGFP (at E7.5) and tdTomato (at E8.5) barcoded
lentivirus, analyzing the transduced somite derivatives such as vertebrae and other cells types in
every section along the anterior-posterior axis (Fig. 2g-h). Vertebrae were unlabeled in the mid-
trunk but labeled in the posterior trunk, while cells in neural structures such as the neural tube and
dorsal root ganglia showed both labels throughout the entire body. Labeling of somitic derivatives
spread longer along the body axis in E7.5 injected embryos as compared to E8.5 injected samples
(Fig. 2h), consistent with an expected coupling between the progressive spatial and temporal fate
restrictions in the NMP lineage.
258
We were intrigued that clonal cluster 15 behaved somewhat like NMPs by generating neural crest
cells and somitic derivatives, but not CNS cells (Fig. 2b, f). Consistent with this, we observed
some sections with labeling of both crest-derived neurons and vertebrae but few CNS neurons
(Fig. 2h). Together, this evidence suggests some NMPs can directly generate neural crest during
secondary neurulation without contributing to neural tube epithelialization, demonstrating
existence of a neural crest - mesodermal bipotent progenitor (NCC-MP, Fig. 2i). To demonstrate
the existence of this population, we analyzed the tail mesenchyme of four additional embryo
samples (Extended Data Fig. 13a-f). Indeed, we again recovered a clonal cluster corresponding to
NCC-MPs (Extended Data Fig. 13g-j). Compositional analysis of tail clones revealed that NCC-
MPs differ significantly from canonical NMP clones in the number of CNS cells, but both produce
similar amounts of neural crest cells; as a result, these clonal clusters are marked by a clear bias
towards either CNS or neural crest cells (Extended Data Fig. 13k-m). The NCC-MP population
has been predicted by earlier work [22, 23] but not yet reported. Further studies of NCC-MP
biology might prove relevant to the etiology of some *spina bifida* subtypes.
273
274
*Position-specific auxiliary transcriptional programs are superimposed on common developmental*
*paths of cell lineages to regulate clonal composition*

Cell fate determination involves coordinated induction of transcriptional programs and, often,
suppression of alternate fates [11]. By delineating sets of clones that demonstrate distinct fate
outcomes, we studied how context-dependent transcriptional programs subtly bias multipotency
among otherwise similar progenitor cell populations.

To illustrate how clonal embedding enables fine analysis of how positional factors influence cell
differentiation, we examined a group of trunk mesodermal lineages that build the axial skeleton
with the help of Hox-dependent patterning [24-26]. In the transcriptional embedding, trunk
mesenchymal populations form a large diffuse cluster (Fig. 1j), containing progenitor as well as
early differentiated populations (Fig. 3a, top left). While standard trajectory inference analyses on
such a diverse population would fail due to uncertainty, clone2vec neatly separated the
mesenchymal cells into a set of partially overlapping lineages (Fig. 3a). Ordering these lineages
according to their Hox score revealed their graded contribution to skeletal cells, dermis, meningeal
fibroblasts, and perivascular cells as a function of body position (Fig. 3b and boxed insets in Fig.
3a, Extended Data Fig. 10-11). We noted that cartilage cells were present in many of these lineages
and therefore may be a good model to study position-specific modulation of regulatory programs
in the body. We refer to such spatial regulatory factors that introduce variation to otherwise similar
cell differentiation events as "auxiliary programs".

To further understand the factors driving variation in multipotency profiles, compared two
chondrogenic lineages – Cluster 8, containing cartilage/meninge clones (that did not engage in
osteogenic differentiation, here referred to as 'chondrogenic'), and Cluster 2, containing
'osteochondral' cartilage/bone clones (Fig. 3a). To perform the analysis we extracted bone,
cartilage and progenitor cells from these lineages, filtered out meningeal cells, and re-embedded
them to compare their chondrogenic trajectories (Fig. 3c). Mapping lineage-specific cartilage gene
expression signatures to publicly available Stereo-seq data suggested that chondrogenic lineage 8
represent the forming vertebrae in the lower trunk (Fig 3d, Extended Data Fig. 14), while the
osteochondral lineage 2 is likely from anterior structures including early bone collar of ribs, which
starts forming from perichondrial cells at E13.5 (Fig. 3e) [27, 28]. We observed a full spectrum of
intermediate cell states at E13.5 due to continuous differentiation from mesenchymal progenitors
to mature chondrocytes from both clonal clusters (Fig. 3f-g), suggesting a *bona fide* developmental
trajectory in both cases, although osteochondral clones appear to maintain expression of progenitor
markers such as *Meox1* for longer (Fig. 3g). Additionally, the presence of meningeal cells in
Cluster 8 indicates that it is not simply a less-mature version of Cluster 2 (as might be predicted
based on the asynchrony of somite development).

Focusing on chondrogenic differentiation in both lineages, we found that both executed a
conserved transcriptional program, upregulating common chondrocyte gene markers (Fig. 3h and
Extended Data Fig. 15a-d) including *Col11a1*, *Sox6*, *Col9a3*, *Acan*, *Comp*, *Fgfr3*, *Col11a2*, and
*Matn3*. Expression of bone-related genes (Fig. 3i and Extended Data Fig. 15e-f) increased at the
beginning of differentiation in both trajectories, even in the clonal cluster without bone, suggesting
existence of competitive programs underlying fate selection even when one of the fates is absent.
Analysis of the bone-related genes with clear early activation in the osteochondral lineage revealed
gene expression of the well-known TF *Runx2* in mesenchymal progenitors as the strongest
indicator of overall clonal osteogenic potential (Fig. 3j), aligning with existing models [29].

Despite being downregulated in the chondrogenic-only lineage (Extended Data Fig. 15 g-k), *Runx2*
expression significantly correlated with the bone-related signature even there (Fig. 3k), suggesting
that molecular regulators of mesenchymal *Runx2* expression drive a pro-chondrogenic progenitor
cell's osteogenic versus fibroblast potential in vivo.
327
Differential expression analysis of the clonal lineages with distinct *Runx2* activation implicated
TFs related to the Hox code, mesenchymal fate commitment, and skeletal development (Extended
Data Fig. 16a-h), leading us to question the hierarchy of observed TFs. Using SCENIC [30], we
inferred transcriptional factor activity and likely gene regulatory networks acting during cartilage
development in both clonal clusters (Fig. 3l-m and Extended Data Fig. 16i-j). The analysis
suggested an upstream regulatory role of Hox TFs, with *Hoxa10/Hoxd10* (lumbosacral)
coordinating the regulatory network in chondrogenic clones, and *Hoxa7/Hoxc8* (thoracic) in
osteochondral clones. *Hoxa7* was positioned directly upstream of *Runx2* expression in the
reconstructed gene regulatory network (Fig. 3m), which could explain the positional differences
in clone skeletogenic potential. Additionally, *Hoxc8* (Fig. 3m) was situated upstream of
mesenchymal TF *Prrx1* (via intermediates *Meis1* and *Pbx3*), which like *Meox1* exhibited sustained
expression specifically in the bone-forming lineage 2, suggesting that positional factors might
regulate the tempo of early chondrogenesis by maintaining a mesenchymal progenitor status
(*Meox1/Prrx1* expression). These roles are consistent with a direct regulatory function of Hox
genes in mesenchymal progenitor fate decisions, in agreement with observations at later
developmental stages [31].
344
Overall, combining high-throughput lineage tracing with trajectory inference on cell manifolds
[10] uncovered mechanisms of clonal fate biasing in the embryo, by decoupling cell type-specific
gene modules from spatial gene modules (Fig. 3n). Comparing lumbar and thoracic mesoderm
revealed 1) the core transcriptional program of axial pro-chondrogenic differentiation, 2) early
molecular indicators of osteochondrogenic biopotency, and 3) evidence that spatial programs drive
bone collar formation in thoracic cells, explaining why ribs mineralize faster than the posterior
vertebral cartilages. Untangling how spatial regulatory mechanisms are influenced by
heterochrony, and how this directs specific clonal dynamics such as differentiation tempo and
proliferative potential, remains a future challenge.
354
*Positional codes in facial mesenchyme revealed through clonal analysis.*

In contrast to the trunk, the anterior cranial region is largely devoid of Hox expression. While
previous studies revealed many TFs that are critical for establishing the facial prominences [32,
33], how spatial programs influence multipotency of facial cells at a molecular level remains
insufficiently understood. We applied our clonal variation approach to this problem, again
focusing on chondrocytes, reasoning that transcriptional variation within a single cell type would
accentuate position-specific programs by minimizing cell-type-related artefacts. The cranial
dataset captured a rich spectrum of clonal fate biases (Fig. 4a-b), covering populations contributing
to central and peripheral nervous system, dermis, meninges, and mesenchymal derivatives of the
musculoskeletal system which was segregated into the major lineages corresponding the major
facial prominences (Fig. 4b-d). Examining the spatial distributions of lineage-specific genes in
Stereo-seq data, we found that they consistently mapped to different facial subregions (Fig. 4c).
Differential expression analysis of specific chondrogenic signatures within all three major lineages
revealed key transcription factors, which were sufficient to reconstitute the expected positional

identity (Fig. 4e). We then confirmed the cartilage positional signatures (including previously
reported markers *Alx1/Asb4/Hoxa2* [34, 35]) were also observed in progenitor cells originating
from the same clones (Fig. 4e). The true anatomical position of these signatures was validated by
re-analysis of Stereo-seq data (Fig. 4f). This suggests that these signatures likely represent
position-specific programs modulating progenitor cell fate choices and plasticity. Indeed, fate
profiles of cranial clonal clusters revealed that Hox-negative BA1 (branchial arch 1) progenitor-
derived clones were highly multi-fated, containing fates from nearly all mesenchymal clusters,
whereas Hoxa2$^+$ clones from BA2 (branchial arch 2) lacked several mesenchymal cell types (Fig.
4d). CatBoost regression-based numeric estimation of gene importance for predicting clone
composition and location resulted a list of top factors including known spatial regulatory factors
(Fig. 4g). Overall, these results demonstrate how analysis of continuous clonal variation can
suggest candidates for positional programs within the developing embryo.
381
382    *Cranial neural crest generates neuronal and mesenchymal fates from partly overlapping but*
383    *mostly distinct progenitor pools.*
384
Beyond position-specific effects, continuous clonal variation can also capture temporal
dependencies in multipotency regulation. Cranial neural crest cells (CNCCs) emigrate in waves
from the dorsal neural fold but face different microenvironmental contexts depending on their
exact timing of delamination, which makes for a good system to study temporal modulation of cell
fate choices in vivo [36, 37]. As a population, the cranial neural crest is highly multipotent, giving
rise to facial skeletal and other mesenchymal structures as well as to the peripheral nervous system
(trigeminal and sympathetic ganglia, including the glial cells), but whether such broad
multipotency can be attributed to individual CNCCs remains unclear [37-40]. For instance, each
wave of CNCCs might carry specific fate biases; alternatively, multipotent CNCCs would be
biased by the post-migratory environment.
395
Examining clonal variation of CNCC lineages, we found that facial mesenchymal derivatives and
trigeminal neuroglial cell types were overwhelmingly separated into distinct clonal subpopulations
(clusters 6+19 versus clusters 0-1,3-5,7-10,14-16, Fig. 4d) and very few clones shared trigeminal
neuronal and mesenchymal fates. Furthermore, these two clonal subpopulations were
distinguished by sequential E7.5 + E8.5 double injection, with neuroglial cell types labeled
predominantly at later E8.5 injections (Fig. 4h). This result suggests that early and late-migrating
CNCC waves become differentially biased.
403
To test the temporal separation between mesenchymal and neuroglial trigeminal cell types, we
performed lineage tracing experiments in *Sox10-CreERT2/R26Confetti* embryos, which enable
clonal tracing based on multicolor labelling of stochastically recombined cell populations (Fig 4i).
Recombination was triggered by tamoxifen injection at slightly different developmental times
(~12h differences at E8.5 stage). The resulting clonal tracing showed that, although early labeled
clones contribute broadly to all facial tissues, later-labeled clones were confined to trigeminal
ganglion and outgoing nerves, revealing a systematic difference in the cell type composition
depending on the timing of recombination (Fig 4j). This result is consistent with a temporal model
in which different migratory waves of CNCC undergo differential fate biasing (Fig 4k). This
finding demonstrates how clonal variation can help to resolve subtle temporal effects on lineage
separation and fate biasing among nearly uniform progenitor cell populations.

415
416     *Perturb-TREX: Joint clonal tracing with multiplexed signaling receptor perturbations.*

417
Hypotheses about programs modulating progenitor fate must be eventually tested by experimental
perturbations. In this regard, the highly orchestrated developmental process poses a challenge, as
disruption of pathways critical for cell fate determination, even with conditional genetic
perturbations can result in major distortions of the tissue and lead to widespread indirect effects.
Mosaic perturbations of individual clones, however, are unlikely to cause such tissue distortions,
presenting a valuable opportunity to study the role of genes with strong phenotypes *in vivo*.

424
To examine the role of extrinsic signals in early trunk development, we combined TREX tracing
with multiplexed CRISPR/Cas9 gene editing of individual clones (Fig. 5 and Extended Data Fig.
17) [41]. Taking advantage of detailed scRNA-seq data on differentiating neural crest and other
cell lineages, we extended the barcoded TREX vectors with a library of guide RNAs targeting
receptors of signals associated with early cell fate bifurcation steps (Fig. 5a) [11]. These
multiplexed lentiviral libraries were injected into murine embryos at E7.5 or E8.5 and examined
with scRNA-seq and clone2vec at E13.5 (Fig. 5b). In part because of the known relationship
between skeletogenesis and Hedgehog signaling, we included two gRNAs targeting the Hedgehog
pathway opportunity to study the role of extrinsic signals during axial patterning of skeletal cell
subtypes. While, as expected, none of the embryos showed major developmental abnormalities,
significant fate deviations were observed for mosaic perturbations at the clonal level, resulting in
ectopic clonal clusters (Fig. 5c). These clusters were enriched for *Ptch1* crispant clones - a gene
encoding sonic hedgehog (SHH) receptor [42], and likewise, *Ptch1* gRNA[+] clones were
significantly enriched within the ectopic clusters (Fig. 5d), which represented some variation on
neural tube clones (clonal cluster/lineage 12), ventral mesenchyme (lineage 9), and somitic clones
(lineage 11).

441
Hierarchical clustering to determine lineage similarity was used to find points of comparison for
the ectopic clusters (Fig. 5e). For example, in the developing CNS, *Ptch1*-crispant clones were
enriched in progenitor populations (Fig. 5f-g) compared to similar CNS lineages. *Ptch1* crispant
clones also upregulated a unique gene expression signature (i.e. *Hhip, Fgf15, Ptch2, Plagl1)*
related to cell cycle and Hedgehog signaling (Fig. 5h), and significantly increased cell counts per
clone (Fig. 5i). Thus, *Ptch1* editing in neuroectodermal clones results in an accumulation of stalled
intermediate progenitors prior to their cell cycle exit and neuronal differentiation, and Ptch1-
mediated inhibition of Hedgehog signaling is needed for CNS neural cells to mature beyond a
transit-amplifying progenitor state.

451
The ectopic clone type resulting from elimination of *Ptch1* from mesoderm-derived mesenchyme
was not found in any other analyzed conditions (Fig. 5j-k). This distinct clone type was enriched
with vertebral chondrocytes and meningeal cells (much like the lumbar path 8 in Fig. 3) but was
specifically depleted of undifferentiated mesenchyme and fibroblasts (Fig 5j, compare to control
clusters in Fig 5k). This suggests that in contrast to the CNS, the gain-of-function of Hedgehog
signaling in mesoderm speeds up the tempo of differentiation, depleting the mesenchymal
progenitor pool.

Given that somites occur along the body axis, we were surprised that *Ptch1* editing resulted in a
single perturbed somite lineage and questioned whether this was driven by mis-patterning or
differential local sensitivity. We compared the Hox score of mesenchymal-biased lineages
including *Ptch1* gRNA-enriched clusters 9 and 11 (Fig. 5l), finding that they drastically differed
in positional identity, suggesting an effect of *Ptch1* editing on mesodermal axial patterning.
Indeed, analyzing Hox score per gRNA condition revealed a bimodal distribution of Ptch1 clones
but no other condition including EGFP controls. Cluster 9, with anterior identity (and found
primarily in middle trunk samples), was primarily biased towards duct mesenchymal cell types
forming from the ventral part of the somite (Extended Data Fig. 17e, see location of gene
expression score in Extended Data Fig. 14h) suggesting that *Ptch1* perturbation caused
ventralization of anterior but not posterior mesoderm. On the other hand, the somite Cluster 11
(posterior identity, found primarily in lower trunk sample) contained clones from a wide range of
Hox scores, some significantly anterior compared to nearest neighbor lineages from lumbar and
tail somites (Fig. 5l), suggesting that the axial pattern of clonal fate biases was disrupted.
Accordingly, Hox scores of *Ptch1* gRNA[+] clones were significantly less predictable than clones
bearing other gRNAs, when using any nearest neighbor clone as reference (Fig. 5n), and even
when using the same gRNA as reference (Fig. 5o).
477
478    Taken together, while a highly predictable embryo-wide gradient of clonal fate biases is observed
479    under natural conditions, this can be overridden by the activation of cell signaling pathways. Future
480    studies focusing on the lineage-specific interactions of *Ptch1* with cell differentiation programs
481    should be performed using detailed comparison of the regulatory programs acting differentially
482    across lineages (as in Fig. 3) coupled with combinatorial perturbations along a time series.
483
484    *Biasing of cell fate decisions in neural crest revealed via clonal analysis.*
485
We next focused on the trunk neural crest, which has been hotly debated as being either a mixed
population of restricted progenitors, or a population of highly multipotent stem cells [12, 36, 43-
45]. Clonal variation analysis of wildtype embryos revealed that all detected neural crest sub-
lineages were highly fate biased. This was observed for sympathetic, melanocyte, somatosensory,
enteric, and glial progenies (Fig. 6a-c). Importantly, none of the neural crest lineages were clonally
linked with mesenchyme (Fig. 6c), and sparingly few crest-mesenchymal bipotent clones were
observed. Thus, although vagal neural crest gives rise to both cardiac mesenchyme and enteric
neurons, this multipotency is more accurately attributed to the collective population, and not to
individual neural crest cells. Instead, our results suggest that prior to clonal expansion, labeled
neural crest progenitors acquire strong fate biases, corresponding to the biases described for
different migratory waves of neural crest [36, 46]. Such progenitor differences may be too subtle
to be observed from transcriptional analysis alone [47]. For instance, we find that the progenitor
state of the clones destined to give rise to the *Phox2b*[+] autonomic nervous system and enteric
nervous system within the developing gut, both map to the same overall transcriptional state (Fig.
6d-e). The knowledge of clonal identity enables a more powerful way of identifying biasing
programs by directly contrasting the progenitors with distinct fates. In this way we find that these
early progenitors show robust upregulation of a set of biasing genes, including *Hand2* - a
transcription factor known to be involved in neural crest enteric differentiation [48]. Since such
genes reflect early progenitor biasing mechanisms that appear prior to overt differentiation, their

modulation may be helpful for cell type generation protocols and applications intended to cure
autonomic and enteric dysfunctions, including Hirschsprung's disease [49].
507
We also examined positional distribution of the neural crest multipotency along the anterior-
posterior and dorsoventral axes. Based on the *Hox* gene expression we find that along anterior-
posterior axis an expected strong anterior bias of the enteric-biased clones [50], as well as paucity
of sympathetic clones the posterior sacral and post-sacral part of the body (Fig. 6f). The remaining
clonal clusters showed broad distributions. To infer the distributions along the dorsoventral
position, we examined the co-occurrence of neural crest and neuronal CNS fates within the same
clones. Since neural crest is formed in the most dorsal part of neural tube, increasing contribution
of clones to CNS neurons would suggest a more ventral position - closer to the border between the
prospective crest domain and the future roof plate of the neural tube [48]. We find that the clones
biased towards somatosensory neurons and melanocytes shared a much higher proportion of
progeny with CNS neurons, suggesting that the originally transduced progenitor cells were
positioned more ventrally (Fig. 6g). These results are consistent with previous spatiotemporal fate
mapping of the dorsal neural tube in chick embryos [36].
521
Finally, to better understand the signals biasing neural crest clones, we examined the compositional
changes associated with CRISPR/Cas9 perturbations of different receptor genes. Our previous
work [11] suggested that neural crest cells undergo transcriptional bifurcation that determines
autonomic or sensory fates. We hypothesized this bifurcation point would be sensitive to the
signaling environment, leading to reciprocal effects on autonomic versus sensory clonal biases.
Receptor perturbation caused significant and opposing shifts in clonal composition of autonomic
vs somatosensory progenies in the neural crest lineage, suggesting that the corresponding receptors
and signaling pathways are indeed involved into defining proportions of fate outcomes (Figure 6h-
i). For instance, *Fzd2, Lrp2, Lgr4*-crispant clones had reductions in autonomic and enteric biases,
accompanied by increases in sensory and glial fates. The perturbation of *Fgfr1* showed exactly the
opposite effects on clonal biases and composition. These results expand our understanding of the
role of signaling receptors in neural crest specification by demonstrating the existence of a sensory-
autonomic clonal lineage bifurcation [11, 51] and its modulation by putative FGF and WNT
signaling molecules.
536
537  **Discussion**
538
Context-dependent regulation of cell fates is a fundamental aspect of embryogenesis, essential for
the proper formation of tissues. Tracing of clonal structures can reveal the spectrum of such fate
modulation and provide important clues about the mechanisms involved. Embryogenesis,
however, presents a particular challenge for modern lineage tracing approaches, since the large
number of generated cell types and the diverse position-specific nature of modulation demands
both increased resolution and scale of tracing. A resolution that would enable biological
discoveries has been difficult to achieve on an embryo scale, leaving high-throughput tracing
studies to focus on specific cell lineages such as hematopoiesis or forebrain development [15, 20].
547
Here, we presented an effective practical approach for studying mechanisms underlying fate
modulation in an embryo, by integrating multiplexed *in vivo* tracing, computational analysis of
transcriptional programs associated with clonal fate biases, and mosaic medium-throughput

multiplexed perturbations. The scale of the study, in this case profiling thousands of individually
barcoded ectodermal and mesodermal clones across developmental timepoints, is one enabling
aspect that allows study of relative levels of continuous versus discrete clonal variation across cell
lineages and spatial domains.
555
In the past, low-throughput clonal lineage tracing analysis has been used extensively to
successfully address hypotheses about cell multipotency and differentiation [44]. Although these
studies have established the classic paradigms about lineage splits in embryogenesis, they were
not sufficiently powered to investigate the continual variation of clonal fates. Even though
different developing structures are often made from the same cell types [52, 53], these structures
acquire distinct shapes and functions due to the fine regulation of cell fate proportions and spatial
positioning of similar cell progenitor populations [54, 55], developing within the same cell lineages
[56]. Insights into the continual variation of clonal structures help to explain how these
morphogenesis-related differences gradually arise in a coordinated and context-aware manner.
565
Even state-of-the-art high-throughput lineage tracing methods still sample only a small fraction of
cells in the embryo. The sparseness of the resulting clonal data represents a major challenge for
subsequent analysis. This has prompted us to develop a machine learning approach – clone2vec –
which represents the second enabling aspect of our study. Starting with a continuous representation
of expression state, clone2vec embeds sparse clonal data into a low dimensional space where
clones with similar or compatible fate profiles are grouped together. In contrast to the methods
aimed at analysis of factors associated with specific cell fate choices (CoSpar [57] or CellRank
[58]), clone2vec aims to provide an effective overview of clonal variation, enabling exploration of
recurrent patterns and subsequent analysis of associated molecular features. The continuous
representation of the expression space enables a shift away from a cell type-centric perspective of
clones' classification, allowing for a nuanced description of state patterns in clone distribution.
This approach, as we show, can recover complex clonal fate patterns in a manner highly robust to
dropouts (Supplementary Note Simulations), and permits generation of new hypotheses. In
addition to separating distinct progenitor lineages (*e.g.* mesenchymal and neural), clone2vec
embeddings capture continuous clonal variation within these lineages, which is much less
understood and is a central point of our study.
582
In contrast to tree representations [5-7, 18], clonal embeddings enable analysis of gradual variation
of clonal fates and inference of regulatory processes modulating clonal fates and proportions of
cell types across spatial locations. Clones within different spatiotemporal contexts display distinct
but overlapping regulatory and transcriptional signatures [59]. A good example is the population
of mesenchymal mesodermal derivatives which forms a large unstructured cluster of cells in the
transcriptional embedding, posing a challenge for the traditional methods of trajectory
reconstruction and other analysis. Lineage tracing and clone2vec analysis, however, effectively
partitions this population into a series of clonally connected sub-clusters, representing variation of
the fundamental differentiation process at different positional or temporal contexts (Fig. 3a). Such
nuanced clonal patterns cannot be easily discerned using traditional approaches such as
hierarchical clustering (Extended Data Figure 11n). Furthermore, the analysis of transcriptional
variation between clonal clusters can reveal more subtle "auxiliary programs" that regulate clonal
structures. For instance, we observed position-dependent variation in skeletogenic mesenchyme,
where additional programs do not appear to be essential for chondrogenic differentiation, but rather

modulate secondary features such as a probability of transiting into osteogenic fate or maturation
dynamics of progenitor cells. Such an approach should be useful for analyzing position-dependent
modulation of progenitor potency in other contexts such as during tissue regeneration [60]. Further,
distilling core versus auxiliary differentiation programs for more cell types might aid optimization
of efficient and parsimonious *ex vivo* differentiation methods needed for regenerative medicine.
602
In addition to resolving spatial modulation of clonal development, we also observed heterochrony
of clonal fate biases by introducing static barcodes at precise developmental stages. This can be
contrasted with other methods employing dynamic barcodes [7, 8], in which the timing of lineage
bifurcations can be difficult to pinpoint, complicating efforts to link lineage decisions to the
corresponding signals and developmental stages. Our comparison of cranial neural crest cells
traced from two successive time points showed temporal segregation of facial skeletogenic
mesenchyme and trigeminal ganglion from the cranial neural crest, which was noted in chick [40]
but not yet in mammals, to our knowledge. This finding suggests that temporal windows shaping
cell potency are essential for building an integrated head, as neural crest gives rise to structures as
distinct as peripheral nerves and facial bones [61]. The diversity and range of facial shapes
produced during vertebrate craniofacial morphogenesis likely benefits from an added layer of
dynamic temporal control that complements positional cues [62, 63].
615
The coupling of clonal lineage tracing with mosaic perturbations provides powerful means of
validating mechanistic and causative relationships that underlie cell fate regulation in various
spatiotemporal niches. Our focus in this study was on perturbations of extrinsic signaling axes, as
prior studies and our own correlative analysis of natural clonal patterns indicated that they play a
major role in modulating auxiliary programs and guiding progenitor fates. Indeed, we demonstrate
that perturbations of specific extrinsic signals alter positional biases in the neural crest lineage.
Furthermore, disrupting certain signaling axes, such as Hedgehog (via *Ptch1*), results in formation
of ectopic clonal clusters in which cell differentiation appears accelerated or retarded, suggesting
a new opportunity to study plasticity and robustness of development. Since mosaic perturbations
impact only a very small fraction of cells in the tissue, this approach can be used to evaluate the
impact of signaling axes that would otherwise severely distort the tissue or lead to unviable states.
In this way, mosaic multiplexed perturbations offer an effective strategy to study cell fate
regulation in native tissue environments under both normal and diseased conditions.
629
In addition to insight into signaling landscapes and positional cues, high-throughput analysis of
clonal biasing in the mammalian embryo uncovered previously unrecognized lineage relationships
or fate restrictions. For instance, that NMPs – multipotent mesoderm-neuro competent cells [13,
21] - coexist in the tail region with mesoderm-restricted progenitors. Our results yielded
transcriptional signatures that separate the E13.5 derivatives of NMPs from that of their fate-
restricted neighbors (*i.e. Ran, Hoxb13*) that could be helpful in determining whether a cellular
memory of NMP multipotency exists. This study also identified a population of murine NMPs that
contribute neural crest cells but not spinal cord neurons (so-called NCC-MPs) [23]. While classical
models assume that neural crest arises from the neuroepithelium, our findings suggest that a subset
of NMPs gives rise to neural crest independently of the CNS, which might represent a novel
mechanism of neural crest induction. This implies that axial progenitor pools are more diversified
than previously thought, and their regulation will influence the balance between CNS, neural crest,
and mesodermal cells during secondary neurulation and body extension. While our results provide

a gene signature to distinguish them (*i.e. Tspan18, Pde8a*), further studies will be needed to determine if neural crest cells derived from NCC-MPs bear functional differences from other sacral crest cells, and how differences in NCC-MP behavior affect spinal cord development and function.

On a larger developmental scale, our analysis reveals a picture of global fate restrictions at the stages of late gastrula and early neurula, which altogether opposes the idea of early plasticity of different germ layers as was suggested from classic experiments in vitro [64]. According to our data, most clones do not span the competence of more than a single germ layer, except for the NMPs and a handful of exceptional clones from the cranial neural crest. This does not rule out that cells from these stages might exhibit different multipotency profiles when subjected to different environments, as in the case of transplantation experiments that can reveal greater cell plasticity [1, 40].

Overall, we hope the presented approach will be helpful for further studies aimed at systematically dissecting the spatial and temporal correlates of clonal variation in different cell systems. The current clonal atlas, including its interactive interface (browsable online at https://clones2cells.streamlit.app) should provide a useful resource for a wide range of questions related to the formation of boundaries between developing embryonic structures [65]. Similarly, clone2vec analysis should aid in the identification of fine-grained positional and temporal codes. While the Hox code and other crude positional systems provide a general framework for the development of body axes and programming the large body domains [25], the fine-grained signals that guide specific cell behaviors within very local tissues are much harder to study due to the complexity and redundancy of embryonic signaling systems [66]. Clone2vec may assist investigations of how fine-grained positional information informs cell diversity within individual organs like the heart or lungs. This might shed light on other unresolved questions, for instance, the enigma of the anterior-posterior allocation of brain nuclei [67], or finer patterning of mesodermal populations building individual muscles [68]. Further, joining clone2vec clonal analysis with mosaic perturbations enables multiplexing simulated conditions that mimic human genetic abnormalities, including neurodevelopmental disorders and pediatric cancers (i.e. neuroblastoma). This might help reveal clonal compensatory behavior in modelled congenital diseases, as well as in regeneration, potentially revealing clonal mechanisms of embryonic resilience [69].

Despite the potential, we realize the limitations of the study. The presented multiplexed tracing and perturbation approach can be widely applied for focused analysis of specific tissues and compartments, however, the potential to refine these techniques is substantial. Current limitations include the constrained developmental time frames and specific cell types that can be effectively targeted using existing barcoding techniques. Inference of differentiation paths or the state of progenitor cells based on one measured endpoint poses well-known fundamental challenges, particularly when the underlying process is asynchronous and one cannot observe, for example, both progenitor and differentiated cells from the same clone in the snapshot provided by scRNA-seq data. While we show that successive barcoding rounds can help, incorporating dynamic barcoding strategies such as CRISPR scarring [5-8, 70] is likely to improve path reconstruction, and systematic variation of experimental endpoints might assist progenitor state inference.

689
690 **Acknowledgements:**
691
*General acknowledgements*
We thank Olga Kharchenko for the lovely illustrations. We thank Ruslan Soldatov for the
insightful discussions. We thank Daniel Wies, Adrian Martinez Martin, and Polina Kameneva for
their technical assistance. We acknowledge the National Genomics Infrastructure, Biomedicum
Flow Core, Biomedicum Imaging Core, Advanced Light Microscopy Core at SciLifeLab, and
Biomedicum Comparative Medicine for access to their facilities and services during the project.
698
*Funding*
AGE was supported by the National Institute of Dental and Craniofacial Research via the Ruth L.
Kirschstein National Research Service Award (F32DE029662). IA and SE were supported by the
ERC Synergy grant (KILL-OR-DIFFERENTIATE), Swedish Research Council, Paradifference
Foundation, Bertil Hallsten Research Foundation, Cancer Foundation in Sweden, Knut and Alice
Wallenberg Foundation, Austrian Science Fund. P.V.K. was founded by the NSF-14-532-
CAREER grant.
706
707
*Author contributions*
Conceptualization: IA, PVK, AGE, SI
Investigation: IA, PVK, AGE, SI, MR, JF, EA, BS, JH, JF
Validation: AGE, SI, AM, BS, EA, JH
Computational investigation and analysis: SI, AGE, MR, PVK
Writing – original draft: IA, PVK, AGE, SI, EA, AC, BS, JF, MR, AM
Funding and resources acquisition: IA, EA, AGE, JF, PVK
Supervision and project administration: IA, EA, AGE, JF, PVK
716
717 *Competing Interests*
718 The authors declare that they have no competing interests.

719
720
721 **Methods**
722
*CloneID and gRNA-CloneID library construction*
Barcoded lentiviral libraries were generated based on Ratz et al [15]. Briefly, a lentiviral backbone
was constructed containing human elongation factor 1 alpha (EF1a) promoter driving expression
of either tandem dimer tomato (tdTomato) or green fluorescent protein fused to histone 2b for
nuclear localization (H2B-EGFP). For perturbation libraries, a gRNA scaffold using capture
sequence "cs1" within the hairpin loop was inserted to LV-U6-tdTomato-NLS constructs using
Gibson assembly. Then, the resulting vector was digested using BsmBI and separately annealed
gRNA oligonucleotide sequences (which were chosen from the validated sequences in the Broad
Institute's Brie CRISPR knockout library) were cloned into the gRNA scaffold by ligation. The
resulting vectors were amplified by transformation by heat shock into ONE-SHOT STBL3
competent cells, isolated using a miniprep, and sequence-verified by Sanger sequencing. Next,
control backbones (or equimolar mixtures of gRNA-containing backbones) were used for insertion

of an amplified random 30N oligonucleotide library via Gibson assembly. The library was further
amplified via electroporation of Endura electrocompetent cells. The barcoded plasmids were then
subjected to third-generation lentiviral packaging with a VSV-G envelope and concentrated by
ultracentrifugation by KI Virus Tech facility or by GEGTech (Paris, France). The resulting ultra-
concentrated viruses had high functional titers greater than $10^9$ TU/ml.

*Animal experimentation*
All experiments were approved by the Swedish Animal Welfare Body Jordbruksverket De
regionala Djurförsöksetiska nämnderna and Stockholms Djurförsöksetiska nämnd, permit number
N15907-2019/18314-2021. Animals from non-genetically modified mouse lines CD-1 and
C57BL/6 were purchased from Charles River Laboratories and Janvier. Animals were kept in an
SPF animal facility with standardized conditions (24 °C, 12:12-h light–dark cycle, normal
humidity, food and water ad libitum). CD1 or Cas9-GFP mice were placed together into time-
controlled matings, and the next day the female was plug checked – if found positive, this was
counted as embryonic day (E) 0.5.

*Ultrasound guided injections (single injections)*
Injections for lentiviral gene delivery were performed according to previously published protocols
[14]. A glass capillary was pulled, and the tip was grinded to create an ultra-sharp needle. At
embryonic day E7.5-E8.5, the females were anesthetized with isoflurane (4% induction,
maintained at 1.5-2.5%) and placed on a heated plate for surgery. Depth of anesthesia was tested
through pinching the paw. Eye gel was applied to prevent dehydration. Pain relief e.g.
Buprenorphine 0.05-0.1 mg/kg was administered via subcutaneous injection shortly before the first
incision. All instruments were autoclaved and disinfected with chlorhexidine alcohol before
surgery. The skin was disinfected with chlorhexidine alcohol, and the hair on the abdomen was
removed with a shaver or hair removal agent such as Veet. A small incision was placed in the
abdomen and the uterus was exposed and pulled through an elastic bottom of a modified petri dish
with PBS. The uterus was kept moist with warm NaCl solution (37C). Embryos were visualized
using a Vevo2100 ultrasound imaging system. Using the glass capillary, lentivirus (from 23-206
nL) was injected through the uterine wall into the amniotic cavity. After the embryos were injected,
the uterus was brought back into the abdomen and the incisions in the muscle wall were sewn
together with absorbent suture thread, and the skin was closed with suture thread or staples. The
mice were taken off isoflurane and allowed to wake up in their own cage on a heating pad under
supervision. Postoperative pain relief was given in the form of a subcutaneous injection of
Buprenorphine (0.05-0.1 mg/kg). For embryo retrieval, the mother was euthanized via CO2
overdose and cervical dislocation, and the embryos were collected at E13.5 and euthanized via
decapitation.

*Ultrasound-guided double injections*
To determine temporal aspects of general fate restrictions in vivo, two sequential uterine injections
in pregnant CD1 mice were performed. A first injection of 206 nL (corresponding to ~313000
predicted unique CloneIDs) pMR532-LV-EGFP-bc lentivirus was done on mouse embryos from
embryonic day 7 (TS11). A second injection of 69nL (corresponding to ~72,000 predicted unique
CloneIDs) of pMR671-LV-TOM-bc was performed at an interval of 12-24 hours after the first.

There was no observable scar tissue formation on the day of the second injection. The ultrasound-
guided injection itself took place as described in the above section. The embryos survived without
visible developmental delays after the single or double injections. Embryos were euthanized by
decapitation at E13.5, and cells from the most highly labeled embryos were harvested for
downstream analysis as described above, while the remaining littermates were fixed overnight in
paraformaldehyde at 4C for histological analysis of cell labeling distribution. During and after the
second injection, the mice were given the following pain relief protocol: 5 mg/kg of the long-
acting anesthetic Meloxicam, 2 mg/kg of the local anesthetic Bupivacaine, and 0.1 mg
Buprenorphine. We tested this protocol for pain relief after double injections 24 hours apart.
Together with the veterinarian, we used 3 females to evaluate the effectiveness and impact of the
protocol. We monitored the females for pain and recovery from the procedure, assessed the healing
of the surgical wound, and checked the status of the fetuses toward the end of the procedure. All
females recovered quickly after the first and second injection (within 10 min of placing the cage
on a warm pad, the females were walking around the cage and eating normally). The animals
showed minimal signs of discomfort or pain after the procedure, on the injection days as well as
the day after.
797
*Single cell dissociation*
The embryos were removed from the uterus, sacrificed, and placed into a petri dish containing
Hank's Balanced Salt Solution (HBSS) on ice. Embryos were screened for expression of tdTomato
or EGFP using a benchtop fluorescence stereoscope. Limbs, liver, and the ventral skin were
removed. Body regions, primarily lower trunk, middle trunk, and cranial regions, were dissected
separately and placed into a small Petri dish containing 2.5 mL of a cell dissociation buffer
consisting of 2mg/mL Collagenase P (Roche, Sigma-Aldrich), 1X TrypLE, and 1uL/mL DNASe
I, fully dissolved in HBSS (without phenol red, including $Ca^{2+}$ and $Mg^{2+}$). Tissue was chopped
finely with scalpel and forceps under control of stereomicroscope and transferred to 15mL falcon
tubes. The tissue pieces were triturated 10X using a transfer pipette, and then 10X using a 1mL
pipette. Samples were then incubated for 20 minutes at 37 C in a preheated shaking incubator set
at 150 RPM, with the tubes tilted to ~60 degrees. During this time every 6-8 minutes, the samples
were vortexed and triturated using a 1mL pipette. After the tissue clumps disappeared, 12.5 mL of
2% Fetal Bovine Serum (FBS) diluted in ice-cold HBSS (without phenol red, without $Ca^{2+}$ and
$Mg^{2+}$, H6648 SIGMA) was added. The samples were centrifuged in a cold centrifuge (4C) for 15
minutes at 500 g. The supernatant was carefully removed and the pellet was resuspended in 1mL
HBSS + 2% FBS and kept on ice thereafter. After centrifugation and resuspension, the cells were
filtered into round-bottom polypropylene test tubes (Falcon) using a cell strainer with a 40
micrometer woven polyethylene terephthalate mesh (pluriSelect pluriStrainer SKU 43-10040-40)
for downstream cell sorting.
818
*FACS (Fluorescence activated cell sorting)*
FACSDiva software (BD), connected to FACSAria Fusion or FACSAria III flow cytometers (BD)
equipped with 100 micrometer nozzle and 1.5 neutral density filter, were used to isolate the
fluorescently labeled cells. During sorting the samples were refrigerated to 4 C and regularly
agitated by spinning. The event rate was kept below 4000 events/second. Debris, dead cells and
doublets were excluded by gating forward-scatter and side-scatter area versus width. Log EGFP
and tdTomato fluorescence was detected by excitation with 488 nm and 561 nm lasers, with
emissions filtered by 530/30 nm and 610/20 nm bandpass filters, respectively. FP-negative cells

served as an internal negative control for FACS sorting. Sorting precision mode was set to "four-
way purity". FP-positive cells were sorted directly into Low-binding tubes containing 50uL of
HBSS + 2% FBS. For each sample, all the sorted FP$^+$ cells were centrifuged in a cooled centrifuge
at 500g for 5 min and resuspended for cell counting with a hemocytometer. From FACS the
proportion of labeled cells was ~5% on average. Because tracing every cell in the embryo across
organogenesis would not be feasible (because of the massive number of cells in a single E13.5
embryo) we prioritized measuring fewer but more "complete" clones. We observe consistent cell
types and clone types across embryos, indicating that a broad range of accessible progenitor types
was sampled in all experiments.
836
*Single cell transcriptomics*
Cells were loaded into the Chromium machine (aiming for ~10,000 cells per reaction) using the
V3.1 single-cell reagent kit (10X Genomics). Following GEM creation and cell lysis, cDNA was
synthesized by reverse transcription, then amplified by PCR for 12 cycles as per the manufacturer′s
protocol (10X Genomics). A portion of the amplified cDNA was used to generate Gene expression
libraries and then indexed using the Single Index Plate T Set A. CloneID enrichment from cDNA
libraries was additionally performed based on [15] to increase the retrieval of the barcodes by ~5-
10%. Briefly, two nested PCR reactions were performed using Q5 error-correcting DNA
polymerase to amplify the CloneID-containing region, and a third reaction was used for sample
indexing using the Single Index Plate T Set A. In between each reaction, amplicons were purified
by 0.8X AMPure beads using a DynaMag Magnetic separator. Indexed GE and CloneID-
enrichment libraries were quantified using BioAnalyzer and/or Qubit, and then pooled prior to
Illumina sequencing on a NovaSeq 6000 S2-200 v1.5 flow cell with read setup 28-8-0-166.
850
*Confetti lineage tracing experiment*
Transgenic mouse line R26R-Confetti [37, 54], homozygous for the confetti allele, was crossed
with Sox10-CreERT2 or PLP-CreERT2 mice. At embryonic day E9 they were injected with 0.02
mg/g tamoxifen (TM) intraperitoneally. Approximately 42 h post TM injection, the pregnant mice
were sacrificed and embryos still in the uterus were collected in PBS on ice. Then, under the stereo
microscope (ZEISS Discovery V8) and in PBS, the muscle layers of the uterus were ripped
carefully with micro-dissecting forceps for the embryos to be extracted. Cre-positive R26R-
Confetti embryos were identified on the basis of multicolor fluorescence using a stereo microscope
fluorescence adapter. E10.5 R26R-Confetti Cre$^+$ mouse embryos were processed to be analyzed
histologically or in whole mount. Embryos were fixed in 4% paraformaldehyde (PFA) solution in
phosphate buffer saline (PBS) overnight at 4C on rotation. Afterwards, fixed samples headed for
histological analysis were washed three times in 1x PBS for 5 min each and were immersed in
30% sucrose in PBS overnight at 4oC on rotation. The next day, they were embedded in Optimal
Cutting Temperature (OCT)-compound on dry ice and stored at -20C until cryosectioning. An
NX70-Epredia cryostat was used to cut sections with a 14μm thickness which were serially
collected onto Superfrost® Plus slides (Thermo Scientific, J1800AMNZ/ground 90°). Prior to
storing at -20C, they were left to dry on the bench for 2 hours. Fixed embryos headed for whole
mount analysis were washed in PBS before immersed in CUBIC-Reagent I clearing solution for 1
day at 37 C, gently shaking. Then, they were carefully washed with PBS at room temperature (RT)
prior to being immersed in CUBIC Reagent II clearing solution at RT for approximately 6 h. The
embryos should be totally transparent in order to be placed onto glass bottom dishes (MatTek
Corporation) for imaging. The imaging media is CUBIC-Reagent II. This procedure was also used

for the whole mount imaging short-term E8.5 lentiviral transductions in E10.5 embryos, except
the samples were imaged using a light sheet microscope (Zeiss Z.1) equipped with a 5X objective.
Zen Black software was used during acquisition, and manual segmentation in Bitplane Imaris
software (v9) was used to reveal the clonal labeling patterns beneath the skin.
877
*Immunohistochemistry*
For histological analysis, tissue sections on slides were submerged in Target Retrieval Solution
(DAKO, S1699) for antigen retrieval and brought to boiling point using a microwave oven. The
solution was then left to cool down for 40-60 min. Sections were washed in PBS containing 0.1%
Tween-20 (PBST). A hydrophobic barrier was created with Super PAP Pen (Invitrogen, 008899)
surrounding the tissue. The sections were incubated in a humidified chamber overnight at 4 C with
primary antibodies diluted in PBST. The next day, sections were washed in PBST and incubated
with secondary antibodies diluted in PBST for 45-90 min. Afterwards, slides were washed with
PBST, excessive liquid was removed and the slides were mounted using Mowiol (Merck, 81381)
as mounting medium. Primary antibodies that were used against mouse at the indicated dilutions:
goat anti-Sox10 (1:800, R&D systems, AF2864), mouse anti-neurofilament medium chain (2H3)
(1:100, DSHB, AB_2314897), goat anti-CD31 (1:400, R&D systems, AF3628), mouse anti-
Myosin Heavy Chain (MyHC) (1:50, R&D systems, MAB4470), mouse anti-Tyrosine
Hydroxylase (TH) (1:1000, Sigma-Aldrich, T2928), chicken anti-GFP (1:500, Aves Labs, GFP-
1020), rabbit anti-tRFP (1:500, Invitrogen, PA5-79658), rabbit anti-Sox9 (1:500, Millipore).
Primary antibodies were detected with Alexa Fluor 405-, 488-, 555- and 647-conjugated secondary
antibodies produced in donkey against mouse, rabbit, goat or chicken antibodies (1:1,000,
Molecular Probes, Thermo Fisher Scientific).
896
*Microscopy and image analysis*
Confocal microscopes were used for the imaging of either tissue sections or whole mount samples.
Whole mount imaging of cleared E13.5 transduced embryos was performed using light sheet
microscopy on a Zeiss Z.1 microscope using a 5X objective. For the Sox10-CREERT2 Confetti
labeling, the ZEISS LSM 980 with Airyscan 2 confocal was used due to its capacity to effectively
distinguish the overlapping emissions between GFP and YFP. The images were acquired at a 20x
objective and 2056x2056 resolution with the use of a z-stack. The imaging of the sections from
lentiviral-injected samples was conducted at ZEISS LSM 800 and 800 Airy with a 20x objective
and 1024x1024 resolution. ImageJ 1.53t software was used for the analysis of tissue sections. The
different color combinations in Sox10-CREERT2 Confetti sections were counted manually with
the "Cell Counter" plugin. The determination of DRG units was based on the number of cells in
each DRG per embryo. The lentiviral-injected labeled cells in tissue sections were counted with
the Cell Counter plugin in compartments of interest like DRG and NT. Bitplane Imaris version
10.0 software was used for the analysis of whole Sox10-CREERT2 Confetti embryos. The four
channels were first masked based on the "Surface" to remove background, and a second surface
mask was used to distinguish double positive cells.
913
*scRNA-seq Data Preprocessing and Clone Identification*
In the course of the experiments, one to three libraries were generated for each sample: (1) 10x v3
Gene Expression, and (2-3) viral barcode-containing cDNA libraries (two in the case of double
injection). Detailed descriptions of each sample, corresponding libraries, and FP sequences are
provided in the Github repository of this project (please see Code Availability Statements). Reads

from experiments (1) and (2) were aligned to a chimeric mm39 and FP genome reference using
Cell Ranger 7.0.1. Subsequently, Gene Expression matrices underwent additional filtering based
on sequencing depth, UMI distribution per cell, genes per cell, and the percentage of mitochondrial
expression per cell. Specific thresholds are detailed in the Github repository (Code Availability
Statement). Clonal barcodes were identified using an algorithm suggested in [13] with
modifications available at https://github.com/serjisa/TREX.modified. Firstly, barcode sequence
corrections were performed individually for each cell, enhancing robustness against artificial
merging but slightly increasing clonal dropout rates. Secondly, cells were grouped into clones only
if they shared identical sets of viral barcodes, precluding the merging of cells with different
barcodes. Clonal identification was performed simultaneously on all samples corresponding to the
same mouse. Doublet filtering was conducted based on gene expression and clonal information
before grouping cells into clones. Scrublet v.0.2.3 [71] with manual thresholds was employed for
manual doublet filtering. Additionally, cells with two or more viral barcodes, where such
combinations existed in only one cell, were considered doublets. Despite these measures, a few
clusters likely representing doublets based on marker expression were observed in the final
embedding; these were retained in the analysis and labeled as "Doublets." Reference gene
expression embeddings were constructed using scanpy v.1.9.3 [72] in several steps. Gene
expression embeddings from all control samples (both head and trunk) were created to identify
major cell type domains (Mesenchyme, Neurons, and Other cells). Genes present in fewer than
five cells were filtered out. Subsequently, counts per cell were linearly transformed to sum to
10,000 counts per cell, followed by log-transformation of (counts + 1) [73]. The top 3,000 highly
variable genes were selected, and PCA was performed on gene-wise scaled expressions. Using the
top 30 PCs, dataset horizontal integration was performed with the harmonypy v.0.0.5 package
[74]. These corrected PCs were used to construct a kNN graph based on Euclidean distances,
followed by community detection with the Leiden algorithm [75] (leidenalg v.0.9.1) and
visualization with UMAP [76] (umap-learn v.0.5.3). These processing steps were repeated for each
of the larger cell subclusters in the head and trunk individually, with an additional step to mitigate
the effects of the cell cycle on clustering results. Cell cycle phases were assigned to each cell as
described in [77], and a "reference" was built with cells in interphase. This reference was used for
label and coordinate transfer using Symphony [78] (symphonypy v.0.2). The joint latent
representation post-assignment was used for further analysis.
950
*Lineage Relationships and clone2vec Algorithm*
For plotting correlational relationships between cell types, Pearson correlation-based hierarchical
clustering was employed. To create clonal latent embeddings, only clones with at least 10 cells
were selected. Cells from these clones, termed "whitelisted cells," were used to find 15 nearest
neighbors among the whitelisted cells based on batch-corrected Harmony coordinates using the
pynndescent v.0.5.7 package. Each whitelisted cell and its 15 nearest neighbors formed a "bag of
clones," consisting of 15 pairs of context for the target clone. Using a Skip-Gram architecture [19],
a neural network was trained to predict clonal labels of context cells based on the clonal label of
the target cell. Clonal labels were one-hot-encoded, and a fully connected neural network with one
hidden layer (size = 10) was trained with a negative log-likelihood loss. The matrix describing the
linear transformation between the input and hidden layers was used as the resulting embedding,
capturing similarities between clones. This embedding enabled the construction of a kNN graph
of clones, community detection, and UMAP visualization. Clonal embeddings were constructed
for both E7.5 and E8.5 clones simultaneously. Cells with multiple clonal labels from E7.5 and

E8.5 were duplicated prior to "bag of clones" construction to count them twice with different clonal
labels. For per-clone gene expression analysis, the expression of each cell within a clone was
averaged to obtain the final per-clone expression. For embedding exploratory purposes, PCA was
performed on the clone2vec embedding space, and gene expression was used to identify
associations with clone2vec PCs. Clonal cluster names were manually assigned by inspection of
the cell states and cell types that were present in them, and literature searches for embryonic
progenitor cell types that would give rise to each of those cell type ensembles. This information
was used in combination with analysis of the gene expression patterns indicating anatomical
position, and spatial validation in mouse embryos by mapping clone type-specific gene signatures
onto publicly available Stereo-seq datasets.
975
*Gene Importance Analysis*
To identify genes associated with clonal composition, we built models to predict either clone2vec
latent embedding coordinates or cell type proportions within each clone. For this purpose, we used
a gradient boosting model, as its applicability for gene regulatory network (GRN) inference has
been demonstrated [79]. Specifically, we employed the relatively overfitting-robust CatBoost
algorithm [80], following the strategy described below. First, we created pseudo-bulk profiles for
each clone within each cell type to mitigate learning of cell type–specific gene expression. To
further reduce this effect, we included cell type as a categorical predictor alongside gene
expression. Clones were then split into two subsets: a training set (80% of clones) and a validation
set (20%) to monitor overfitting during training. The validation set was selected using a geometric
sketching approach [81], choosing clones that best represent the clone2vec space. CatBoost
regression (for latent coordinate prediction) or classification (for clone composition prediction)
was applied using MultiRMSE or MultiCrossEntropy loss functions on both the training and
validation sets. Training was stopped if the validation loss did not improve for more than 100
iterations, and the best model was selected for further analysis. The final model was used to
estimate feature importance for each prediction across the dataset using Shapley values [82, 83].
To quantify the overall importance of each gene in clonal behavior, we averaged the L2-norm of
the SHAP vectors from the clone2vec model across all generated pseudo-bulks. To investigate
gene associations with cell types, we extracted SHAP value modules from the cell type model and
assigned them the sign of their correlation with gene expression. This sign correction allowed us
to distinguish between positive and negative markers of cell type proportions. Thus, the clone2vec
model captures the overall variance in clonal behavior, while the cell type model provides a
mechanistic interpretation of gene effects at the cell type level. Both models were trained using
the same train/validation split for consistency. For visualization, sign-corrected SHAP magnitudes
from each cell type can be min-max normalized with respect to their sign.
1001
1002
*Cartilage Trajectory Analysis*
To reconstruct differentiation trajectories within individual clonal clusters, cells were re-
embedded, and clusters linked to the cartilage/bone trajectory were selected from clonal clusters 2
and 8 (clustering resolution = 1, see Extended Data Fig. 18 for parameter sweeps of clustering
resolution). Velocity analysis with UniTVelo (v.0.2.5.2) confirmed the differentiation
directionality within the selected cells. Bifurcation points were identified using the scFates
package (v.1.0.7) [84]. Pseudotime was inferred using the diffusion pseudotime approach, and
cells with pseudotime value less than 0.3 were considered as progenitors [85]. Alignment between

two differentiation trajectories was performed with the DTW Python package (v.1.3.0) [86, 87]. The significance of the association between gene expression and pseudotime was determined using GAM fitting and LRT, comparing models with and without pseudotime-associated splines. Differences in associations with pseudotime across trajectories were identified by comparing GAMs with and without clonal cluster assignment as a covariate. Genes significantly associated with pseudotime ($p < 10^{-25}$) and not differing between clonal clusters (FDR > 0.05) were considered "common", at the same time genes with significant lineage association (FDR < $10^{-5}$) and fold-change > 1 between lineages were considered as "unique". Bone-related signature was gotten by differential expression analysis between cartilage and bone in clonal cluster 2 (top-20 highly expressed in cartilage trajectory genes with FDR < $10^{-20}$ and logFC > 4) using Student's t-test. Analysis of "early" genes was performed with standard scFates pipeline [84]. Transcription factor (TF) activity inference was conducted using pyscenic (v.0.12.1) [30] with arboreto (v.0.1.6). Differential activity analysis between clonal clusters followed the same procedure. Graph of the potential regulation was gotten from cisTarget step of pyscenic protocol. In all cases signature scores were calculated with sc.tl.score_genes scanpy function.

*Spatial Validation of Gene Expression Signatures*
Stereo-seq data from [88] was used to identify the spatial localization of gene expression signatures. Gene signatures were identified using Student's t-test between groups of cells (e.g., between different clonal clusters or within the same cell type across different clonal clusters) with a minimum log2FC of 2. Expression of gene signatures within one cell type was visualized as the product of gene and cell type signatures. Hox-score was calculated as a min-max normalized weighted sum of expressions of the following Hox-genes:

$Hox\ score_{raw} = (1 \times \text{expr}(Hoxb4) + 2 \times \text{expr}(Hoxb5) + 3 \times \text{expr}(Hoxc6) + 4 \times \text{expr}(Hoxa7) + 5 \times \text{expr}(Hoxc8) + 6 \times \text{expr}(Hoxd9) + 7 \times \text{expr}(Hoxc10) + 8 \times \text{expr}(Hoxd11) + 9 \times \text{expr}(Hoxc11) + 10 \times \text{expr}(Hoxc12) + 11 \times \text{expr}(Hoxc13)) / 66;$

$Hox\ score = (Hox\ score_{raw} - \min(Hox\ score_{raw})) / (\max(Hox\ score_{raw}) - \min(Hox\ score_{raw})).$

*Perturbation Analysis*
Perturbational screening experiments were mapped onto the control reference via Symphony, as previously described (separately for cycling and non-cycling cells). Each clone from the perturbational experiment was assigned to a gRNA based on the most abundant gRNA combination within the clone. Clones with multiple gRNAs were excluded from further analysis. New clonal embeddings were constructed as described in the Lineage Relationships and Clone2vec Algorithm section. Differential compositional testing between clones in different clonal clusters was conducted using scCODA [89] to analyze clone composition in the entire clonal embedding rather than individual cells within different cell types. An independent embedding focusing on NC-related clones (with at least three cells) was created using both control and perturbed cells. Compositional testing was performed for perturbations with multiple replicates using beta regression and LRT with FDR multiple testing correction.

**Code Availability**

All code that was used in the current study is readily available on Github: (https://github.com/kharchenkolab/clonal-atlas-paper/). The clone2vec Python package is also available on Github: (https://github.com/kharchenkolab/scLiTr). Additional documentation for the

scLiTr package can be found here: (https://sclitr.readthedocs.io/). The modified TREX pipeline is available here: (https://github.com/serjisa/TREX.modified). Also, the code for the clones2cells app is available via this link: (https://github.com/serjisa/clones2cells_app)

**Data Availability**

Raw data has been submitted to the NCBI Gene Expression Omnibus database (GEO) and is accessible via the following link: (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE269395). Data has also been uploaded to Zenodo: (https://zenodo.org/records/15397247). The data can be browsed via the clones2cells app via this link: (https://clones2cells.streamlit.app). Alternatively, browse the datasets on our CELLXGENE single-cell dataset repository: (https://adameykolab.hifo.meduniwien.ac.at/cellxgene_public/filecrawl/2024_Erickson_Isaev)

**References**

1.  Zechner, C., E. Nerli, and C. Norden, *Stochasticity and determinism in cell fate decisions.* Development, 2020. **147**(14).
2.  Kretzschmar, K. and F.M. Watt, *Lineage tracing.* Cell, 2012. **148**(1-2): p. 33-45.
3.  Livet, J., et al., *Transgenic strategies for combinatorial expression of fluorescent proteins in the nervous system.* Nature, 2007. **450**(7166): p. 56-62.
4.  McKenna, A., et al., *Whole-organism lineage tracing by combinatorial and cumulative genome editing.* Science (New York, N.Y.), 2016. **353**(6298): p. aaf7907-aaf7907.
5.  Pei, W., et al., *Resolving Fates and Single-Cell Transcriptomes of Hematopoietic Stem Cell Clones by PolyloxExpress Barcoding.* Cell Stem Cell, 2020. **27**(3): p. 383-395 e8.
6.  Bowling, S., et al., *An Engineered CRISPR-Cas9 Mouse Line for Simultaneous Readout of Lineage Histories and Gene Expression Profiles in Single Cells.* Cell, 2020. **181**(6): p. 1410-1422 e27.
7.  Li, L., et al., *A mouse model with high clonal barcode diversity for joint lineage, transcriptomic, and epigenomic profiling in single cells.* Cell, 2023. **186**(23): p. 5183-5199 e22.
8.  Spanjaard, B., et al., *Simultaneous lineage tracing and cell-type identification using CRISPR–Cas9-induced genetic scars.* Nature Biotechnology, 2018. **36**(5): p. 469-473.
9.  Raj, B., et al., *Simultaneous single-cell profiling of lineages and cell types in the vertebrate brain.* Nature Biotechnology, 2018. **36**(5): p. 442-450.
10. Wagner, D.E. and A.M. Klein, *Lineage tracing meets single-cell omics: opportunities and challenges.* Nature Reviews Genetics, 2020. **21**(7): p. 410-427.
11. Soldatov, R., et al., *Spatiotemporal structure of cell fate decisions in murine neural crest.* Science, 2019. **364**(6444).
12. Erickson, A.G., P. Kameneva, and I. Adameyko, *The transcriptional portraits of the neural crest at the individual cell level.* Semin Cell Dev Biol, 2023. **138**: p. 68-80.
13. Tzouanacou, E., et al., *Redefining the progression of lineage segregations during mammalian embryogenesis by clonal analysis.* Dev Cell, 2009. **17**(3): p. 365-76.

14. Mangold, K., et al., *Highly efficient manipulation of nervous system gene expression with*
*NEPTUNE.* Cell Rep Methods, 2021. **1**(4).
15. Ratz, M., et al., *Clonal relations in the mouse brain revealed by single-cell and spatial*
*transcriptomics.* Nature Neuroscience, 2022. **25**(3): p. 285-294.
16. Kameneva, P., et al., *Single-cell transcriptomics of human embryos identifies multiple*
*sympathoblast lineages with potential implications for neuroblastoma origin.* Nat Genet,
2021.
17. Roe, T., et al., *Integration of murine leukemia virus DNA depends on mitosis.* EMBO J,
1993. **12**(5): p. 2099-108.
18. Weinreb, C. and A.M. Klein, *Lineage reconstruction from clonal correlations.* Proc Natl
Acad Sci U S A, 2020. **117**(29): p. 17041-17048.
19. Mikolov, T., et al., *Efficient estimation of word representations in vector space.* arXiv
preprint arXiv:1301.3781, 2013.
20. Weinreb, C., et al., *Lineage tracing on transcriptional landscapes links state to fate*
*during differentiation.* Science, 2020. **367**(6479).
21. Henrique, D., et al., *Neuromesodermal progenitors and the making of the spinal cord.*
Development, 2015. **142**(17): p. 2864-2875.
22. Frith, T.J., et al., *Human axial progenitors generate trunk neural crest cells in vitro.*
Elife, 2018. **7**.
23. Lukoseviciute, M., S. Mayes, and T. Sauka-Spengler, *Neuromesodermal progenitor*
*origin of trunk neural crest <em>in vivo</em>.* bioRxiv, 2021: p. 2021.02.10.430513.
24. Hall, B.K. and T. Miyake, *All for one and one for all: condensations and the initiation of*
*skeletal development.* Bioessays, 2000. **22**(2): p. 138-47.
25. Carrasco, A.E. and S.L. López, *The puzzle of Hox genes.* Int J Dev Biol, 1994. **38**(4): p.
558-64.
26. McIntyre, D.C., et al., *Hox patterning of the vertebrate rib cage.* Development, 2007.
**134**(16): p. 2981-2989.
27. Flaherty, K. and J.T. Richtsmeier, *It's about Time: Ossification Center Formation in*
*C57BL/6 Mice from E12-E16.* J Dev Biol, 2018. **6**(4).
28. Maes, C., et al., *Osteoblast precursors, but not mature osteoblasts, move into developing*
*and fractured bones along with invading blood vessels.* Dev Cell, 2010. **19**(2): p. 329-44.
29. Loebel, C., et al., *In vitro osteogenic potential of human mesenchymal stem cells is*
*predicted by Runx2/Sox9 ratio.* Tissue Eng Part A, 2015. **21**(1-2): p. 115-23.
30. Aibar, S., et al., *SCENIC: single-cell regulatory network inference and clustering.* Nature
Methods, 2017. **14**(11): p. 1083-1086.
31. Bradaschia-Correa, V., et al., *Hox gene expression determines cell fate of adult periosteal*
*stem/progenitor cells.* Sci Rep, 2019. **9**(1): p. 5043.
32. Depew, M.J., et al., *Reassessing the Dlx code: the genetic regulation of branchial arch*
*skeletal pattern and development.* J Anat, 2005. **207**(5): p. 501-61.
33. Xu, P., et al., *Fox proteins are modular competency factors for facial cartilage and tooth*
*specification.* Development, 2018. **145**(12).
34. Gu, R., et al., *Single-cell transcriptomic signatures and gene regulatory networks*
*modulated by Wls in mammalian midline facial formation and clefts.* Development, 2022.
**149**(14).
35. Kessler, S., et al., *A multiple super-enhancer region establishes inter-TAD interactions*
*and controls Hoxa function in cranial neural crest.* Nat Commun, 2023. **14**(1): p. 3242.

1148 36. Krispin, S., et al., *Evidence for a dynamic spatiotemporal fate map and early fate*
1149   *restrictions of premigratory avian neural crest.* Development, 2010. **137**(4): p. 585-95.
1150 37. Kaucka, M., et al., *Analysis of neural crest-derived clones reveals novel aspects of facial*
1151   *development.* Sci Adv, 2016. **2**(8): p. e1600060.
1152 38. Dupin, E., et al., *The issue of the multipotency of the neural crest cells.* Dev Biol, 2018.
1153   **444 Suppl 1**: p. S47-S59.
1154 39. Weston, J.A. and J.P. Thiery, *Pentimento: Neural Crest and the origin of mesectoderm.*
1155   Developmental Biology, 2015. **401**(1): p. 37-61.
1156 40. Baker, C.V., et al., *Early- and late-migrating cranial neural crest cell populations have*
1157   *equivalent developmental potential in vivo.* Development, 1997. **124**(16): p. 3077-3087.
1158 41. Platt, Randall J., et al., *CRISPR-Cas9 Knockin Mice for Genome Editing and Cancer*
1159   *Modeling.* Cell, 2014. **159**(2): p. 440-455.
1160 42. Cohen, M., et al., *Ptch1 and Gli regulate Shh signalling dynamics via multiple*
1161   *mechanisms.* Nature Communications, 2015. **6**(1): p. 6709.
1162 43. Bronner-Fraser, M. and S.E. Fraser, *Cell lineage analysis reveals multipotency of some*
1163   *avian neural crest cells.* Nature, 1988. **335**(6186): p. 161-4.
1164 44. Baggiolini, A., et al., *Premigratory and migratory neural crest cells are multipotent in*
1165   *vivo.* Cell Stem Cell, 2015. **16**(3): p. 314-22.
1166 45. Krispin, S., E. Nitzan, and C. Kalcheim, *The dorsal neural tube: a dynamic setting for*
1167   *cell fate decisions.* Dev Neurobiol, 2010. **70**(12): p. 796-812.
1168 46. Tang, W., et al., *Clonal analysis and dynamic imaging identify multipotency of individual*
1169   *Gallus gallus caudal hindbrain neural crest cells toward cardiac and enteric fates.* Nat
1170   Commun, 2021. **12**(1): p. 1894.
1171 47. Yu, V.W.C., et al., *Epigenetic Memory Underlies Cell-Autonomous Heterogeneous*
1172   *Behavior of Hematopoietic Stem Cells.* Cell, 2017. **168**(5): p. 944-945.
1173 48. D'Autréaux, F., et al., *Hand2 is necessary for terminal differentiation of enteric neurons*
1174   *from crest-derived precursors but not for their migration into the gut or for formation of*
1175   *glia.* Development, 2007. **134**(12): p. 2237-2249.
1176 49. Montalva, L., et al., *Hirschsprung disease.* Nature Reviews Disease Primers, 2023. **9**(1):
1177   p. 54.
1178 50. Kam, M.K.M. and V.C.H. Lui, *Roles of Hoxb5 in the development of vagal and trunk*
1179   *neural crest cells.* Development, Growth & Differentiation, 2015. **57**(2): p. 158-168.
1180 51. Sutton, G., R.N. Kelsh, and S. Scholpp, *Review: The Role of Wnt/β-Catenin Signalling in*
1181   *Neural Crest Development in Zebrafish.* Frontiers in Cell and Developmental Biology,
1182   2021. **9**.
1183 52. Noden, D.M. and P.A. Trainor, *Relations and interactions between cranial mesoderm*
1184   *and neural crest populations.* Journal of Anatomy, 2005. **207**(5): p. 575-601.
1185 53. Wang, M., et al., *Distinct Gene Regulatory Dynamics Drive Skeletogenic Cell Fate*
1186   *Convergence During Vertebrate Embryogenesis.* bioRxiv, 2024: p. 2024.03.26.586769.
1187 54. Kaucka, M., et al., *Oriented clonal cell dynamics enables accurate growth and shaping of*
1188   *vertebrate cartilage.* eLife, 2017. **6**: p. e25902.
1189 55. Kaucka, M., et al., *Signals from the brain and olfactory epithelium control shaping of the*
1190   *mammalian nasal capsule cartilage.* Elife, 2018. **7**.
1191 56. Hirasawa, T. and S. Kuratani, *Evolution of the vertebrate skeleton: morphology,*
1192   *embryology, and development.* Zoological Lett, 2015. **1**: p. 2.

57. Wang, S.W., et al., *CoSpar identifies early cell fate biases from single-cell transcriptomic*
*and lineage information.* Nat Biotechnol, 2022. **40**(7): p. 1066-1074.
58. Weiler, P., et al., *CellRank 2: unified fate mapping in multiview single-cell data.* Nat
Methods, 2024. **21**(7): p. 1196-1205.
59. Bastide, S., et al., *TATTOO-seq delineates spatial and cell type–specific regulatory*
*programs in the developing limb.* Science Advances. **8**(50): p. eadd0695.
60. Otsuki, L. and E.M. Tanaka, *Positional Memory in Vertebrate Regeneration: A Century's*
*Insights from the Salamander Limb.* Cold Spring Harb Perspect Biol, 2022. **14**(6).
61. Adameyko, I. and K. Fried, *The Nervous System Orchestrates and Integrates*
*Craniofacial Development: A Review.* Frontiers in Physiology, 2016. **7**(49).
62. Usui, K. and M. Tokita, *Creating diversity in mammalian facial morphology: a review of*
*potential developmental mechanisms.* EvoDevo, 2018. **9**(1): p. 15.
63. Chen, Y.C. and N. Konstantinides, *Integration of Spatial and Temporal Patterning in the*
*Invertebrate and Vertebrate Nervous System.* Front Neurosci, 2022. **16**: p. 854422.
64. Waddington, C.H. and J. Taylor, *Conversion of Presumptive Ectoderm to Mesoderm in*
*the Chick.* Journal of Experimental Biology, 1937. **14**(3): p. 335-339.
65. Dahmann, C., A.C. Oates, and M. Brand, *Boundary formation and maintenance in tissue*
*development.* Nature Reviews Genetics, 2011. **12**(1): p. 43-55.
66. Tkačik, G. and T. Gregor, *The many bits of positional information.* Development, 2021.
**148**(2): p. dev176065.
67. Lipovsek, M. and R.J. Wingate, *Conserved and divergent development of brainstem*
*vestibular and auditory nuclei.* Elife, 2018. **7**.
68. Grimaldi, A. and S. Tajbakhsh, *Diversity in cranial muscles: Origins and developmental*
*programs.* Curr Opin Cell Biol, 2021. **73**: p. 110-116.
69. Solini, G.E., et al., *Xenopus embryos show a compensatory response following*
*perturbation of the Notch signaling pathway.* Developmental Biology, 2020. **460**(2): p.
99-107.
70. Alemany, A., et al., *Whole-organism clone tracing using single-cell sequencing.* Nature,
2018. **556**(7699): p. 108-112.
71. Wolock, S.L., R. Lopez, and A.M. Klein, *Scrublet: Computational Identification of Cell*
*Doublets in Single-Cell Transcriptomic Data.* Cell Systems, 2019. **8**(4): p. 281-291.e9.
72. Wolf, F.A., P. Angerer, and F.J. Theis, *SCANPY: large-scale single-cell gene expression*
*data analysis.* Genome Biology, 2018. **19**(1): p. 15.
73. Ahlmann-Eltze, C. and W. Huber, *Comparison of transformations for single-cell RNA-*
*seq data.* Nat Methods, 2023. **20**(5): p. 665-672.
74. Korsunsky, I., et al., *Fast, sensitive and accurate integration of single-cell data with*
*Harmony.* Nature Methods, 2019. **16**(12): p. 1289-1296.
75. Traag, V.A., L. Waltman, and N.J. van Eck, *From Louvain to Leiden: guaranteeing well-*
*connected communities.* Scientific Reports, 2019. **9**(1): p. 5233.
76. McInnes, L., J. Healy, and J. Melville, *Umap: Uniform manifold approximation and*
*projection for dimension reduction.* arXiv preprint arXiv:1802.03426, 2018.
77. Satija, R., et al., *Spatial reconstruction of single-cell gene expression data.* Nat
Biotechnol, 2015. **33**(5): p. 495-502.
78. Kang, J.B., et al., *Efficient and precise single-cell reference atlas mapping with*
*Symphony.* Nat Commun, 2021. **12**(1): p. 5890.

79.    Moerman, T., et al., *GRNBoost2 and Arboreto: efficient and scalable inference of gene*
1239           *regulatory networks.* Bioinformatics, 2019. **35**(12): p. 2159-2161.
1240    80.    Dorogush, A., *CatBoost: gradient boosting with categorical features support*, V. Ershov,
1241           Editor. 2018: Arxiv. p. arXiv:1810.11363.
1242    81.    Hie, B., et al., *Geometric Sketching Compactly Summarizes the Single-Cell*
1243           *Transcriptomic Landscape.* Cell Syst, 2019. **8**(6): p. 483-493 e7.
1244    82.    Lundberg, S., *Consistent Individualized Feature Attribution for Tree Ensembles*, G.
1245           Erion, Editor. 2019. p. arXiv:1802.03888.
1246    83.    Lundberg, S., *A unified approach to interpreting model predictions*, S.-I. Lee, Editor.
1247           2017: arXiv. p. arXiv:1705.07874.
1248    84.    Faure, L., et al., *scFates: a scalable python package for advanced pseudotime and*
1249           *bifurcation analysis from single-cell data.* Bioinformatics, 2023. **39**(1).
1250    85.    Haghverdi, L., et al., *Diffusion pseudotime robustly reconstructs lineage branching.* Nat
1251           Methods, 2016. **13**(10): p. 845-8.
1252    86.    Giorgino, T., *Computing and Visualizing Dynamic Time Warping Alignments in R: The*
1253           *dtw Package.* Journal of Statistical Software, 2009. **31**(7): p. 1 - 24.
1254    87.    Alpert, A., et al., *Alignment of single-cell trajectories to compare cellular expression*
1255           *dynamics.* Nat Methods, 2018. **15**(4): p. 267-270.
1256    88.    Chen, A., et al., *Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA*
1257           *nanoball-patterned arrays.* Cell, 2022. **185**(10): p. 1777-1792.e21.
1258    89.    Büttner, M., et al., *scCODA is a Bayesian model for compositional single-cell data*
1259           *analysis.* Nat Commun, 2021. **12**(1): p. 6876.
1260

Figure 1

**Clonal composition is driven by proliferation and differentiation**

**a**

Clonal proportions
30% 20% 50%

Cell geneaology

Transcriptional diversity

**Lineage trees are superpositions of clonal fate biases**

**b**

A B C D

100
50%
0

Clonal fraction per cell type

**How does anatomical location shape clonal fate biases?**

**c**

A B
A C D
C D
B

**d** Lentiviral library: 30 bp random sequence

EF1a H2B-eGFP 30N PA

CloneID_1  CloneID_2  ...  CloneID_n

**e** Ultrasound-guided injection

capillary
E8.0

Ultrasound system

**f** Genetic barcoding of exposed cells

Ectoderm  N ect

E7.5  E8.0  E8.5  Experiment timeline  E13.5

E8.0-8.25 single injection experiment

E7.5 + E8.5 double injection

harvest (trunk, head)

**g** Cell diversification over time

**h** FACS sorting + scRNAseq analysis

GEX UMAP2
GEX UMAP1

**Single-cell transcriptomics analysis of barcoded body regions**

**i** Head (50 568 cells)

Mesenchyme
Neurons and glia
Other

GEX UMAP2
GEX UMAP1

**j** Trunk (109 176 cells)

Mesenchyme
Glia
Neurons
Other

GEX UMAP2
GEX UMAP1

**Frequency of clonal behaviors observed in the dataset**

**k** E7.5 injection 5 (clone 209)

GEX UMAP2
GEX UMAP1

E7.5 injection 5 (clone 118)

GEX UMAP2
GEX UMAP1

**l**

Number of clones

1173
398 359 199 176 88 30 18 18 12 11 5 3 2

Trunk
E7.5 injection

Glia
Neurons
Other (skin, muscle, etc.)
Mesenchyme

Number of clones

1262  422  255  257  154  6  7  0  5  7  11  6  6
2095

Trunk
E8.5 injection

**m** Cell type lineage correlation

NT NCC MES NNE

NT
NCC
MES
NNE

**Clone2vec learns clonal relationships from transcriptional context**

**n**

Input: scRNAseq dataset

Clonal neighbourhood

Cell  Clonal context

GEX UMAP2
GEX UMAP1

**o** Skip-gram word2vec (clone2vec)

Target clone  Context clones

**Clonal embedding:**
• captures similarities between clones
• dropout-robust
• cluster-free

**p** Output: Clonal UMAP and clusters (each dot is a clone)

C2V UMAP2
C2V UMAP1

**Clones from one clonal cluster occupy similar transcriptional niches**

**Figure 1. Machine learning-based profiling of clonal behavior in mouse embryos.**

**(a)** Scheme of a single clone developing over time via mitotic divisions and cell maturation, determining the final clonal composition.
**(b)** Left, collective multipotency model, where partially overlapping fate sub-spectrums, from a population of three clones, collectively generate a fate spectrum (full repertoire of cell types).
**(c)** Anatomical position of the theoretical clones shown in (b).
**(d)** Clonal barcoding system.
**(e)** Experimental setup. Ultrasound is used to visualize the nanoinjector and embryo while injecting lentivirus into the amniotic cavity, once or twice, between the stages of E7.5-E8.5.
**(f)** Infection of the exposed virus-accessible cells in neural and non-neural ectoderm (uninfected cells beige, infected cells colorful) as well as some mesoderm.
**(g)** Clones are allowed to develop until E13.5 when tissue is harvested from head and trunk body regions (each clone color-coded; each cell type shaped differently).
**(h)** Barcoded cells are analyzed by single cell transcriptomics for clone reconstruction.
**(i-j)** UMAP embedding of the barcoded mouse embryonic E13.5 cells from head (i) and trunk (j), color coded by the major cell types retrieved.
**(k)** Examples of individual clones with partially overlapping compositions.
**(l)** Upset plot of major types of trunk clones resulting from injection at E7.5 (top) versus E8.5 (bottom).
**(m)** Pearson's correlation matrix based on clone sharing pairwise across all retrieved cell types.
**(n)** Schematic of clonal context neighborhood.
**(o)** Skip-gram production from clonal neighborhoods using word2vec architecture.
**(p)** Example of a clonal cluster within the resulting clone2vec UMAP embedding. (right) Example of similar clones constituting a clonal cluster.

**Figure 2**

**a** Clonal embedding of trunk

**b** Cell type distribution in clonal clusters

**c** Top predictors of clonal composition

**d** Spatial location of the clones

**e** Clone transcriptional diversity

**f** Timing of injection of the clones

**g** Labeled body regions

**h** Spatial distribution of labeled cells

**i** Updated model of axial progenitors

**Figure 2. Clone2vec captures spatiotemporal patterns of multipotency in the embryo.**

**(a-g)** Analysis of the Clone2vec UMAP embedding of E13.5 barcoded trunk cells.
**(a)** Embedding color-coded by clonal clusters (clustering resolution 2), labeled according to the fate biases predominantly observed in each of the three main super-clusters (corresponding to neural ectoderm, non-neural ectoderm, and mesodermal derivatives).
**(b)** Dot plot showing cell type composition among clonal clusters. Color represents average cells within each fate, per clone, within that cluster; dot size represents the fraction of the clones in the cluster containing that fate.
**(c)** Elbow plot showing the top twenty most important genes for predicting cell type composition of clones using the CatBoost model, color coded by the Shapley value (feature importance) for each gene for that model.
**(d)** Left, distribution of each clonal cluster Hox gene expression profile, plotted per clone against the known anteroposterior position of embryonic Hox domains (x-axis) and compared to ground-truth manually dissected body regions (horizontal bar plot). Right, feature plot of each mesenchymal clone's Hox score.
**(e)** Top, feature plot showing the number of fates per clone. Bottom, violin plot showing the distribution of Shannon entropy (here used as a metric for transcriptional diversity) per clone within each clonal cluster.
**(f)** Top left, mesenchyme embedding color-coded by injection time. Insets highlight the multipotent neuronal-mesenchymal clones in tail mesenchyme (see Extended Data Figures for validation of NCC-MP clones). Bottom: the composition of each clonal cluster by each injection time.
**(g)** Confocal micrographs of a dual injected (GFP, at E7.5; Tomato, at E8.5) E13.5 embryo at sacral (left) and thoracic (right) axial levels, stained for Sox10 (blue), GFP (green) and Tomato (red). Scale bar = 100 μm. Vb; vertebrae. DRG; dorsal root ganglion. NT; neural tube.
**(h)** Heat map displaying the labeled GFP$^+$ and Tomato$^+$ cells within the vertebra, neural tube, and sensory ganglia per section, ordered by axial position. Dotted box highlights the absence of labeled vertebral chondrocytes from anterior sections.
**(i)** Schematic of axial extension model featuring canonical as well as NCC-biased NMP clones (termed as NCC-MP, or "neural crest cell and mesodermal progenitor") in the caudal lateral epiblast. NMP; neuromesodermal progenitor. NCC; neural crest cell. UMAP; uniform manifold approximation projection.

**Figure 3**

**a** Mesenchymal cell types

Osteoblasts · Chondroblasts · Intervertebral disk (IVD) · Meningeal fibroblasts · Dermal mesenchyme

Fate biases of trunk mesenchyme clonal clusters (spatially-distinct somite-derived lineages)

Lineage 1 (tail) · Lineage 3 (sacral) · Lineage 4 (lumbar) · Lineage 8 (lumbar) · Lineage 2 (thoracic)

KDE density: High / Low

GEX UMAP2 / GEX UMAP1

**b** Lineage (posterior to anterior): 1 3 4 8 2

Dermal mesenchyme · Intervertebral disk (IVD) · Chondroblasts · Meningeal fibroblasts · Osteoblasts

Fraction of clones with fate (%): 20 40 60 80 100

Mean fraction of fate within clone (%): 0.05 0.10 0.15

**c** Chondrogenic clonal clusters with different behavior

Lumbar (lineage 8): meninge-forming clones · Thoracic (lineage 2): bone-forming clones

Chondroblasts · Chondroblasts

subselect + re-embed

Align trajectories (dynamic time warp)

Progenitor · Progenitor · Osteoblasts

Start / End

**d** Position of clonal clusters

Cartilage-specific DEGs

Lineage 8 · Lineage 2

High / Low

**e** Trunk bone deposition

Alizarin red stain

Thoracic ribs · Lumbar vertebrae

**f** Sox9 — chondrogenesis

Lineage 8 · Lineage 2

High / Low

**g** Meox1 — progenitors

Lineage 8 · Lineage 2

High / Low

**h** Common cartilage trajectory

Lineage 8 (meninge-forming) · Lineage 2 (bone-forming)

Gene expression

Abca9, Iqgap2, Col6a3, Col6a2, Col6a1, Meox1, Hic1, Lhfpl2, Sdk1, Tenm4, Egfr, Rbms3, Col5a2, Col3a1, Nfib, Nfia, Col1a1, Sox6, Col9a3, Sdc4, Col9a2, Wwp2, Sorbs2, Pde4dip, Scin, Acan, Comp, Fgfr3, Col1a2, Matn3, Fry, Lgals3

High / Low

pseudotime

**i** Bone signature in progenitors

Lineage 8 · Lineage 2

GEX UMAP2 / GEX UMAP1

Bone score: High / Low

**j** Bone signature

Lineage 2 · Lineage 8

Bone score

Progenitors

Pseudotime

**k** Correlation in progenitor cells

Bone signature score (without Runx2)

2 (r = 0.44, p = 10⁻¹⁶)
8 (r = 0.23, p = 0.006)

Runx2 expression

**l** Regulatory activity specific to Lineage 2

Lineage 8 (meninge-forming) · Lineage 2 (bone-forming)

TF expression: Hoxc8, Ets1, Nfatc4, Meis1, Hoxa7, Pbx1, Prrx1, Runx2, Pbx3, Tbx3, Etv5, Alx1, Runx1

Regulons activity: Hoxc8(+), Ets1(+), Nfatc4(+), Meis1(+), Hoxa7(+), Pbx1(+), Prrx1(+), Runx2(+), Pbx3(+), Tbx3(+), Etv5(+), Alx1(+), Runx1(+)

High / Low

**m** Reconstructed GRN specific to Lineage 2

Ets1, Alx1, Meis1, Pbx1, Hoxa7, Pbx3, Hoxc8, Prrx1, Etv5, Tbx3, Runx1, Runx2, Nfatc4

**n** Hox regulatory programs explain position-driven clonal fate biases

Cell fate restriction · Modulation of proportions

Auxiliary positional program · Clone-specific maturation path · Basic cell type differentiation program

**Figure 3. Position-specific programs are superimposed on common paths of cell maturation.**

**(a)** Kernel density estimate (KDE) plots of cells from each clonal cluster (clustering resolution 1) showing the proportions dedicated to each selected mesenchymal cell type (color-coded in the top left cell embedding). Insets mark the position of the clonal cluster on the mesenchymal portion of the clone2vec embedding (Figure 2a).

**(b)** A dot plot summarizing the fate distribution of each lineage shown in (a), ordered by their position along the body axis.

**(c)** Trajectory inference and alignment of the chondrocyte-meningeal clone type (lineage 8) and the osteochondral clone type (lineage 2) cells by dynamic time warp (DTW). Dark blue represents the origin of each trajectory (mesenchymal progenitors). Dotted insets show RNA velocity arrows denoting the directionality of cell maturation.

**(d)** E13.5 mouse Stereoseq feature plot of gene scores derived from transcriptional profiles of chondrocytes within fibro-chondral (left) versus osteochondral (right) clone types.

**(e)** Confocal fluorescence micrograph of a skeletal preparation of the trunk of an E14.5 embryo. Blue indicates cartilage; red indicates mineralized bone tissue.

**(f-g)** Feature plots displaying *Sox9* (f) and Meox1 (g) expression per cell within the selected chondrocyte-restricted (left) and bipotent osteochondral (right) clone types.

**(h)** Heatmap displaying common gene expression pattern along the chondrogenic trajectories of the restricted (left) and bipotent (right) clone types.

**(i)** Expression of bone-related genes in progenitors (pseudotime < 0.3) cells.

**(j)** Scatterplot of gene expression of the bone-related gene signature, as a function of pseudotime, in cells belonging to the chondral versus osteochondral clone types.

**(k)** Correlation between *Runx2* and bone-related genes' expression in progenitor cells from these two clone types.

**(l)** Heatmap displaying patterns of regulon activity (top) and transcription factor expression (bottom) unique to osteochondrogenic clones.

**(m)** Gene regulatory networks predicted within this lineage.

**(n)** Model of positional regulation of clone development, in which cell differentiation dynamics (driven by a core set of cell-type-specific genes) is under the influence of position-specific auxiliary programs (such as Hox genes).

Figure 4

Mapping of craniofacial clonal cluster signatures to transcriptional and anatomical space

**a** Clonal embedding of cranial region distinguishes facial prominences

**b** Craniofacial lineages mapped to mesenchymal cell UMAP

**c** Lineage density | Lineage marker enrichment
Cell embedding | Clonal embedding | Spatial transcriptomics

**d** Clonal cluster cell type composition

**e** Position-specific genes mark clonal clusters
Regions in cartilage | Cartilage regional markers

**g** Top predictors of clonal composition

**f** Spatial distribution of regional genes
Within cartilage | All cell types

**h** Injection time

**i** Temporal shift in cranial neural crest fate biases
Sox10 CreERT2; R26 Confetti: CFP YFP GFP RFP
0.5mg tamoxifen E8.5, harvest E13.5 | 0.5mg tamoxifen E8.5, harvest E10.5
Restricted clones (n=4 embryos) | Pattern 1 (n=4 embryos) | Pattern 2 (n=4 embryos)

**j** p = 0.00350

**k** Early - mesenchymal bias | Late - neuronal bias

**Figure 4. Collective multipotency of the cranial neural crest cell population.**

**(a)** Clone2vec UMAP embedding of E13.5 barcoded craniofacial cells, color-coded by clonal clusters (clustering resolution 2). The dotted lines indicate the major discovered mesenchymal lineages.

**(b)** Cell UMAP embedding of E13.5 barcoded craniofacial mesenchymal cells, color coded by the major lineages. Dotted box indicates the position of cartilage cells (later analyzed in (e).

**(c)** Mapping of nose (top), jaws (middle), and neck (bottom)-related clonal lineages by density plots for each lineage (left) and feature plots of aggregated lineage-specific gene expression scores on cell embeddings (second row), clonal embeddings (third row) and E13.5 Stereoseq mouse data (right).

**(d)** Dot plot summarizing the fate distribution of each clone type in (a). Dotted line highlights the chondrocyte cluster.

**(e)** Top left; isolated chondrocyte cell type colored by lineage. Right; Differential expression analysis of chondrocytes revealed lineage-enriched genes within mesenchymal cells (top dot plot) and chondrocytes (lower dot plot).

**(f)** Expression of Alx1, Asb4, Hoxa2 in cartilage (left) and in general (right).

**(g)** Elbow plot showing the Shapley value for the top 20 most important features (overall cell type ranked seventh) in a CatBoost model trained to predict cell type composition and position of facial clones.

**(h)** Embedding in (a) color-coded by timing of injection.

**(i-j)** Confocal micrographs of Sox10-CreERT2; R26R-Confetti embryos. Left; Sagittal section through the head of the embryo traced from E8.5 to E13.5. ERBB3 is labeled in magenta and 2H3 is labeled in dark blue. Cyan, yellow, green, and red colors derive from the confetti fluorescent protein labeling. Bottom left; an inset of trigeminal ganglia (outlined with the dotted line) showing rare double-positive GFP/RFP cells. Right panels; maximum intensity projection of whole mounts of Confetti-traced E10.5-E11.0 embryos traced from E8.5-E9. The proportion of observed tracing in the jaw versus trigeminal ganglia is quantified in (j).

**(k)** Model of evolving fate biases in neural crest based on the timing of emigration from the neural tube.

**Figure 5**

**Perturb-TREX experimental design**

**a** Experimental receptor perturbation

Morphogen signaling disrupted

Receptor expression

Fate re-biasing

Criteria for target gene selection

Neural tube

Neural crest

Major signaling pathway receptors
Expressed by progenitor cells
Required for normal development

Inject multiplexed barcoded lentiviral perturbation library

Cas9-eGFP embryos

Experiment 1  Experiment 2  Harvest embryos:
E7.5         E8.5          E13.5

Experiment 1: *Acvr2a, Fgfr1, Fzd2, Lrp2, Lgr4*

Experiment 2: *Acvrl1, Tgfbr3, Bmpr2, Fzd4, Hhip, Ptch1, Lrp4, Bmpr1a*

Trunk (expt. 1 & 2)
Face (expt. 1)

**Analysis of the perturbed clonal landscapes in trunk**

**b** Clonal clusters (lineages)

Skin 15

CNS neurons

NCC derivatives

Mesenchymal derivatives

**c** Perturbations

Novel clusters

Acvr2a, Acvrl1, Bmpr1a, Bmpr2, Fgfr1, Fzd2, Fzd4, Hhip, Lgr4, Lrp2, Lrp4, Tgfbr3

**d** Distribution of crispants among lineages

Proportion of clones in cluster

**e** Similarity of clonal clusters (via fate biases)

Somites

CNS

**Ptch1-gRNA arrests CNS cell maturation, resulting in larger clones**

**f** Trunk CNS neurons embedding

Lineage 12 (Ptch1-gRNA)

KDE density Low High

Dorsal inhibitory

Oligo.

CNS progenitors

**g** Lineage 1 Dorsal neurons

Dorsal excitatory neurons

Dorsal inhibitory lateral

Lineage 7 Ventral neurons

Ventral neurons

**h** Top-10 DEGs in CNS progenitors

Hhip, Igfbpl1, Fgf15, Tec, Ptch2, Ppp2r2b, Slitrk2, Glra1, Gm12239, Fndc5

Fraction of cells in group (%): 10 30 50 70 90

Mean Z-score: -0.7 0.0 0.7

**i** Effect of gRNA on clone size

CNS Progenitors

CNS Neurons

Mesenchyme

Cells per clone

**Ptch1-gRNA disrupts the somitic fate gradient**

**j** Trunk mesenchyme embedding

Lineage 11 (Ptch1-gRNA)

KDE density Low High

IVD, Cartilage, Bone, Meninges, Mesenchymal progenitor, Pericyte, Dermal, Visceral

**k** Lineage 2 tail somite

Lineage 0 lumbar somite

**l** Lineage positional identity

(tail) ... (head)

Hox Score

9 5 3 4 8 11 0 6 2

**m** gRNA effect on mesenchyme patterning

Kernel density estimation

Hox Score

(head) (tail)

Acvrl1, Bmpr1a, Bmpr2, EGFP-1, Fzd4, Hhip, Lrp4, Ptch1, Tgfbr3

**n** Hox 'mismatch score'

Prediction error

Tgfbr3, Bmpr2, Lrp4, Fzd4, Acvrl1, Bmpr1a, Hhip, **Ptch1**

**o** Prediction of Hox score from clone2vec (lower trunk)

R2-score (same gRNA is reference)

**Ptch1**, Acvrl1, Bmpr1a, EGFP, Lrp4, Fzd4

**Figure 5. Novel clonal behaviors caused by genetic perturbations.**

**(a)** Experimental design.
**(b)** Clone2vec UMAP embedding of E13.5 control and perturbed barcoded trunk cells, labeled and color-coded by clusters (clustering resolution 1).
**(c)** Same embedding as in (b), instead color-coded by perturbation. Dotted lines indicate clonal clusters highly enriched with Ptch1 gRNA+ clones. Note cluster 9 is also highly enriched with Ptch1 gRNA+ clones.
**(d)** Bar chart showing the proportion of clone types constituting each gRNA+ population, statistical significance was calculated with scCODA.

**(e)** Hierarchical clustering of clone types based on Euclidean distance between centroids in clone2vec space. Dotted boxes indicate the novel clone types.

**(f-g)** Density plot of the Ptch1-enriched neuronal clone type (f), and the two clone types most closely related to it (g), based on the clustering in (e).
**(h)** Dot plot displaying the top differentially expressed genes by the Ptch1 gRNA+ CNS neural progenitors.
**(i)** Box and whisker plot quantifying the distribution of the number of cells per clone within each gRNA+ population (line = median; box = IQR; whiskers = [Q1 - 1.5IQR; Q3 + 1.5IQR]).
**(j-k)** Density plot of the novel Ptch1-enriched mesenchymal clone type (j) and of the two clone types most closely related to it (k), based on the clustering in (e).

**(l)** Bar and whisker plot showing positional identity of each clone type, quantified by the Hox genes present in each clone. Higher score indicates a position closer to the tail. The label for each clone type is color coded according to the color scheme in (b) and (e). Blue and green bars represent Ptch1-gRNA-enriched mesenchymal clone types (9 and 11, respectively).

**(m)** Kernel density estimation plot showing the distribution of Hox scores per clone across different perturbation conditions. Note the bi-modal distribution in Ptch1 crispant clones.
**(n)** Error rate in predicting Hox score from nearest-neighbor clones (from clone2vec space). Low values indicate that the clones of a given condition consistently have a similar expression profile for Hox genes relative to neighboring clones, whereas high values indicate clones having different positional identities compared to cells with similar fate biases.

**(o)** R2-score (proportion of variance) for each gRNA, reflecting the Hox score predictability of gRNA+ clones, based on nearest neighbor clones (from clone2vec space) bearing the same gRNA. High score indicates high predictability (low variance) in positional identity within the condition, for clones with similar clonal fate biases. Note the low score (high variance) within Ptch1 clones.

Figure 6

**a** Clone type (control + perturbations)

C2V UMAP2 / C2V UMAP1

- Enteric bias
- Glial bias
- Melanocytes & sensory
- Sensory bias
- Sympathetic bias

**b** Neural crest cell types (control + perturbations)

GEX UMAP2 / GEX UMAP1

- Autonomic glia
- BCC
- Chromaffin cells
- Enteric neurons
- MNJ glia
- Melanocytes
- Myelinating glia
- Nonmyelinating glia
- Parasympathetic neurons
- SCP
- Sensory glia
- Sensory neurons
- Sympathetic neurons

**c** Clonal clusters' composition (only control)

Enteric bias
Melanocytes & sensory
Sensory bias
Glial bias
Sympathetic bias

Sensory neurons, Sensory glia, BCC, Parasympathetic neurons, SCP, Chromaffin cells, Myelinating glia, MNJ glia, Nonmyelinating glia, Sympathetic neurons, Melanocytes, Enteric neurons, Autonomic glia, Mesenchyme

Fraction of clones with fate: 20 40 60 80
log (# cells): 0 1 2

**Markers of clonal fate biases among autonomic progenitors subtypes**

**d**

GEX UMAP2 / GEX UMAP1

Enteric neurons
Chromaffin cells
Convergent Phox2b+ state

**e** Clone type fate bias:

Enteric
Glial
Sympathetic

Hand2os1, Hdnr, Plcb1, Hand2, Bnc2, Tlx2, Hoxb9, Robo2, Gse1, 2610035D17Rik

Fraction of cells in group (%): 20 40 60 80 100
Mean Z-score: -0.7 0.0 0.7

**Anteroposterior and dorsoventral characterization of clonal biases**

**f** Clone type fate bias:

Enteric
Sympathetic
Glial
Sensory
Melanocytes & sensory

(head) Hox score (tail)

Fraction of CNS-shared clones: 0.0 0.2 0.4

**g**

chance of becoming NCCs or CNS

undecided roof plate / NCC border at E7-8
cells here will result in CNS / NCC shared clones

**h** Each clone's type of perturbation

C2V UMAP2 / C2V UMAP1

- Acvr2a
- Acvrl1
- Bmpr1a
- Bmpr2
- EGFP-1
- EtOH
- Fgfr1
- Fzd2
- Fzd4
- Hhip
- Lgr4
- Lrp2
- Lrp4
- Ptch1
- Tgfbr3

**i** Effects of cell-autonomous signaling receptor perturbation on clonal biases

E7.5 perturbations (multiple replicates)

Glial
Sensory
Enteric
Sympathetic

Acvr2a, Fgfr1, Fzd2, Lgr4, Lrp2

log2FC between WT and % of clones within cluster

Control KDE

GEX UMAP2 / GEX UMAP1

Fzd2 KDE

**Figure 6. Multipotency landscape of trunk neural crest in normal and perturbed conditions.**

**(a)** Clone2vec UMAP embedding of E13.5 control and perturbed barcoded trunk neural crest-related cells (with more than 2 cells), color-coded by clone types.

**(b)** UMAP cell embedding of the same group of cells shown in (a), color coded by the cell type. Dotted line shows enteric and chromaffin trajectories.

**(c)** Dot plot showing cell type distribution of each clone type. Dot size corresponds to the proportion of clones including a given fate; Dot color corresponds to the average number of cells with that fate per clone.

**(d)** Cell UMAP of the diverging enteric and chromaffin trajectories. Highlighted region indicates the common Phox2b+ glial progenitor state.

**(e)** Dot plot showing gene expression profiles specific to enteric versus glial and sympathetic lineages within that shared state.

**(f)** Predicted anteroposterior positions based on Hox gene expression across all clones within each clone type (left) compared to the proportion of clones shared with CNS cell types shown as a vertical heatmap (right).

**(g)** Model of neural tube development in which dorsal NCC progenitors emigrate prior to ventral cells. This differential timing spent in the tube could affect the number of cell divisions, resulting in different proportions of CNS cells in the clone). This model results from the plotted data in panel f showing the fraction of neural crest clones shared with CNS.

**(h)** Embedding as in (a) color-coded by perturbation.

**(i)** Left; heat map showing fate distributions shifted by each perturbation. (Beta-regression LRT-based p-values with FDR multiple testing correction was used for significance estimation). Right; Density plots of control (left) and gRNA+ (right) clones on the NCC cell UMAP embedding. Arrows indicate the specific loss of enteric fates.