Negative regulation of miRNAs sorting in EVs: the RNA-binding protein PCBP2 impairs SYNCRIP-mediated miRNAs EVs loading.

Francesco Marocco[1,^], Sabrina Garbo[1,^], Claudia Montaldo[2], Alessio Colantoni[3,4], Luca Quattrocchi[1], Gioele Gaboardi[1], Carla Cicchini[1], Gian Gaetano Tartaglia[5], Cecilia Battistelli[1,^,,#] and Marco Tripodi[1,2,^,*]*

[1] Istituto Pasteur Italia-Fondazione Cenci Bolognetti, Department of Molecular Medicine, Department of Excellence 2023-2027, Sapienza University of Rome, Viale Regina Elena 324, 00161, Rome, Italy;

[2] National Institute for Infectious Diseases L. Spallanzani, IRCCS, Rome, Italy;

[3] Biology and Biotechnology Department C. Darwin, Sapienza University of Rome, P.le Aldo Moro 5, 00185, Rome, Italy;

[4] Center for Life Nano- and Neuro-Science, RNA Systems Biology Lab, Fondazione Istituto Italiano di Tecnologia (IIT), 00161 Rome, Italy;

[5] Center for Human Technologies, Istituto Italiano di Tecnologia, Via Enrico Melen, 83, 16152 Genova, Italy;

[^]These authors contributed equally.

∗Corresponding authors:

Francesco Marocco, francesco.marocco@uniroma1.it

Sabrina Garbo, sabrina.garbo@uniroma1.it

Claudia Montaldo, claudia.montaldo@inmi.it

Alessio Colantoni, alessio.colantoni@uniroma1.it

Luca Quattrocchi, luca.quattrocchi@uniroma1.it

Gioele Gaboardi, g.gaboardi@hotmail.it

Carla Cicchini, carla.cicchini@uniroma1.it

Gian Gaetano Tartaglia, gian.tartaglia@iit.it

Cecilia Battistelli, cecilia.battistelli@uniroma1.it

Marco Tripodi, marco.tripodi@uniroma1.it

ABSTRACT

While it is accepted that Extracellular Vesicles (EVs)-mediated transfer of microRNAs contributes to intercellular communication, the knowledge about molecular mechanisms controlling the selective and dynamic miRNA-loading in EVs is still limited to few specific RNA-binding proteins interacting with sequence determinants. Moreover, although mutagenesis analysis demonstrated the presence/function of specific intracellular retention motifs, the interacting protein/s remained unknown.

Here, PCBP2 was identified as a direct interactor of an intracellular retention motif: RIP coupled to RNA pull down and proteomic analysis demonstrated that it binds to miRNAs embedding this motif and mutagenesis proved the binding specificity. Notably, PCBP2 binding requires SYNCRIP, a previously characterized miRNA EV-loader as indicated by SYNCRIP knock-down.

SYNCRIP and PCBP2 may contemporarily bind to miRNAs as demonstrated by EMSA assays and PCBP2 knock-down causes EV-loading of intracellular microRNAs.

This evidence highlights that multiple proteins/miRNA interactions govern miRNA compartmentalization and identifies PCBP2 as a dominant inhibitor of SYNCRIP function.

Keywords: EVs, miRNAs, Poly-C Binding Protein 2, SYNCRIP, miR-122b-3p, miR-155-3p, miR-26b-3p, miR-23a-5p.

INTRODUCTION

Extracellular vesicles (EVs) budding from cellular plasma membrane and differing for size and origin (e.g. exosomes, of 30-150 nm diameter and microvesicles, which diameter is ≥ 50 nm) (*1*) are uptaken by neighboring as well as by distant recipient cells thus transferring their specific informational cargo of molecules. Indeed, EVs are now considered a well assessed route for donor cells to promote changes in gene expression and cell behavior of recipient cells in several pathophysiological processes (*2, 3*). Notably, concerning miRNAs, the EV-cargo does not simply reflect the cell of origin content, but rather is defined by dynamic and selective cell-specific loading mechanisms (*4-8*). The existence of multiple players controlling miRNA compartmentalization matches with the hypothesis of a multiprotein machinery that dynamically governs not only the EV sorting but also, conceivably, the intracellular retention of specific subsets of miRNAs.

At the present, the molecular players controlling the selective partition of miRNAs seems largely uncharacterized although recent evidence highlighted mechanisms of EV-loading depending on sequence-specific RNA-binding proteins (RBPs). On this regard, the first identified was the Heterogeneous Nuclear Ribonucleoprotein A2B1 (hnRNPA2B1), recognizing miRNAs containing the EXO motif (G/U,G,A/G/C,G/C) and guiding their inclusion in EVs secreted by human primary T-lymphocytes (*8*). A functional role was successively attributed to the Synaptotagmin-binding Cytoplasmic RNA-Interacting Protein (SYNCRIP or hnRNPQ) that, by binding to miRNAs embedding a short hEXO motif (G/A/U,G/U/A,G/A/U,C/A/G,U/A,G/C) through its NURR domain and its RRMs recognizing the 5' of the motif (*7, 9*), guides their inclusion in EVs secreted by hepatocytes. Notably, hEXO was shown to increase the EV/cell ratio of miRNAs as demonstrated by the insertion of this sequence in a cell-retained miRNA (*7, 9*) thus paving the way for further manipulative perspectives in the growing field of RNA-based therapies (*10-12*).

More recently, similar functional evidence gathered by the chimeric approach has been extended to newly identified motifs, causing miRNAs export or intracellular retention (*5*). While for these

export motifs two RNA-binding proteins (ALYREF and FUS) have been characterized (*5*) for the retention motifs, defined as "CELL" (*5*), as for the previously described "CL" motifs (*8*) the identification of interacting protein/s remains unaddressed.

We here aimed at the investigation of molecular players responsible for miRNAs intracellular retention. Overall, we gathered evidence on the interactions among several CELL-embedding miRNAs and two RBPs: SYNCRIP and the multifunctional RNA-binding protein PCBP2. Functionally, PCBP2 promotes cellular retention in SYNCRIP-bound miRNAs and the already described SYNCRIP loading activity appears impeded by PCBP2.

RESULTS

PCBP2 recognizes a CELL motif and has a functional role in intracellular retention of

miRNA-155-3p.

Aiming to identify RBPs involved in the intracellular retention, proteins from hepatocytes were

used in RNA-pull-down by using as specific bait the miRNA-155-3p, selected for the presence of

the CELL-motif identified in AML12 cells (*5*) (Figure 1A).

Label-free nLC-MS/MS proteomic analysis allowed to identify miR-155-3p-interacting proteins

(Supplementary Data File 1) and Label-free quantification intensities analysis identified 21 proteins

enriched in miR-155-3p pulldown with respect to miR-155-3p mutated in the CELL motif (miR-

155-3p no-CELL) (Figure 1A, Supplementary Data File 2 and Figure 1B). Twelve of them are

classified as RNA-binding proteins, with six containing at least one canonical RNA-binding domain

(*13-16*) (Supplementary Data File 2). For three of them, RNA-binding preferences are also known

(i.e. PCBP2 prefers CU-rich sequences (*15, 17*), LARP1 recognizes the CAP and the 5' top motif in

mRNAs (*18*) and STAU1 binds to double-stranded RNAs (*19*).

Among them, PCBP2 interaction with miR-155-3p was confirmed by RIP assay (Figure 1C, left

panel) and RNA pull-down followed by western blot analysis (Figure 1D). A second proof of

concept of the requirement of PCBP2/CELL motif interaction is provided by the observation on the

CELL motif-devoid miR-365-2-5p (Figure 1C, middle panel); RNA pull-down confirmed the

absence of interaction with PCBP2 (Figure 1E). At least *in vitro*, this binding appears direct and

sequence specific as demonstrated by the use of a recombinant protein in RNA pull-down (Figure

1F). The introduction of specific mutations allowed to test the requirement of the CELL motif since

its modification (miR-155-3p no-CELL) impairs PCBP2 binding (Figure 1D).

Notably, the inspection of the miR-155-3p sequence revealed the presence of the previously

identified SYNCRIP binding site and both MS/MS analysis (Supplementary Data File 1) and RIP

assay (Figure 1G) confirmed SYNCRIP binding to this miRNA independently of the CELL motif

mutation (Figure 1D).

Surprisingly, the permutation of the downstream two nucleotides removing SYNCRIP-binding

motif (miR-155-3p no-hEXO, Figure 1A) impairs PCPB2 binding despite the conservation of the

CELL retention motif (Figure 1D).

This suggests a possible SYNCRIP requirement for PCBP2 binding. To test this hypothesis, RIP

assay was performed on miR-31-3p, embedding the sole CELL motif, indicating the absence of

PCBP2 binding (Figure 1C, right panel).

Functionally, PCBP2 role in miRNA partition, was addressed by its silencing. Notably, PCBP2

interference (Supplementary Figure 1A and B) enhances miRNA-155-3p loading in EVs with

respect to control cell-derived EVs (Figure 1H, left panel). As a further control that miRNAs

without the CELL motif are not affected by PCBP2 silencing, the expression levels of miR-365-2-

5p (embedding the sole hEXO motif) were analyzed in EVs and cells, and resulted not differentially

exported (Figure 1H, middle panel). As expected SYNCRIP silencing (Supplementary Figure 1C-

D) reduces miR-155-3p export into EVs (Figure 1H, right panel); (for EVs characterization see

Supplementary Figure 2 A and B).

Overall, these data demonstrated that *i)* PCBP2 interacts with miRNA-155-3p, as proved by RIP analysis and RNA pull-down, *ii)* the interaction is CELL-motif-dependent while an unexpected role for the hEXO motif is also unveiled and *iii)* PCBP2 favors the intracellular localization of this miRNA.

PCBP2 binding to miR-155-3p is both sequence- and SYNCRIP-dependent.

The observation that loading (hEXO) and retention (CELL) motifs are both present in miR-155-3p sequence prompted us to investigate on the hypothesis of a sequence- and SYNCRIP-dependent PCBP2 binding ability.

First, we observed that the two proteins interact each other (Figure 2A) and more interestingly that both RBPs bind to miR-155-3p contemporarily as indicated by the ultra-shift obtained in EMSA assay (Figure 2B).

To challenge the hypothesis of a SYNCRIP-dependent PCBP2 binding, EMSA assay was performed in PCBP2-silenced and in SYNCRIP-silenced cells (Supplementary Figure 1A, B, C and D respectively). As shown in figure 2C while the PCBP2 silencing does not affect SYNCRIP binding, SYNCRIP silencing impairs also PCBP2 binding. Furthermore, RIP assay was performed on SYNCRIP-silenced cells; as shown in figure 2D, SYNCRIP silencing impairs PCBP2 binding to miR-155-3p.

This evidence supports the unpredictable mechanism where SYNCRIP binding appears a prerequisite for PCBP2 recruitment. To further confirm and extend this observation, a number of mutants were designed and tested by RNA pull-down for the binding capacity of these two proteins; results indicate that *i)* mutagenesis of the sole hEXO motif (miR26b-3p no-hEXO) on miR-26 backbone (bearing both hEXO and CELL motifs), impairs also PCBP2 binding (Figure 3A) and conversely *ii)* the *de novo* inclusion of a hEXO motif in miR-31-3p backbone (bearing only CELL-

48   motifs) (miR-31-3p +hEXO) confers a *de novo* PCBP2 binding ability to this mutant; furthermore,

49   mutation in the CELL motif (miR-31-3p +hEXO no-CELL) impairs PCBP2 binding (Figure 3B).

Overall, these data indicate that PCBP2 binding requires both the CELL motif and SYNCRIP

binding; in other word, SYNCRIP binding is epistatic to PCBP2 recruitment.

PCBP2 functionally dominates on SYNCRIP EV-loading activity on a repertoire of miRNAs embedding CELL and hEXO motifs.

To extend the evidence for the role of PCBP2 in miRNA compartmentalization and to confirm its mechanistic role *i)* PCBP2 and *ii)* SYNCRIP functional role in miRNA EVs/cell partition was evaluated, and *iii)* PCBP2, *iv)* SYNCRIP and *v)* SYNCRIP-dependent PCBP2 binding were assessed. First, NGS analysis of miRNAs exported in EVs produced by control and PCBP2-silenced murine hepatocytes allowed the selection of further miRNAs differentially loaded in EVs in correlation to PCBP2 (Supplementary Data File 3, Figure 4A, B).

Then, the functional role of PCBP2 was assessed by means of qRT-PCR performed on 9 miRNAs expressed in EVs and embedding both CELL and hEXO motifs in extra-seed position (see Methods section) in comparison to 2 miRNAs embedding either the CELL or the hEXO motif (Figure 4C).

In order to consider a possible impact of PCPB2 on miRNAs steady state level (resulting from variation in transcription and biogenesis), miRNA abundance in EVs and in the intracellular compartment was analyzed by qRT-PCR as EVs/cell ratio. Results indicate that PCBP2 silencing releases a SYNCRIP-dependent loading of miRNAs in the EVs (Figure 4D) thus highlighting a dominant PCBP2 cell-retention function on SYNCRIP-dependent export, evaluated on these miRNAs in Figure 4E.

With respect to the molecular mechanism, the analysis by RIP-qPCR assay demonstrates the PCBP2 binding to miRs-345-3p, 23a-5p, 214-3p, 155-5p, 181d-5p, 3084-5p, 122b-3p, 192-5p, 26b-3p that bear both the CELL and hEXO motifs; conversely, the presence of either the sole hEXO

(miR-365-2-5p) or the sole CELL (miR-31-3p) does not allow PCBP2 binding (Figure 5A). As expected, the presence of hEXO motif alone or in combination with the CELL motif is sufficient for SYNCRIP binding to all the analyzed miRNAs (Figure 5B).

Furthermore, in line with data obtained for miR-155-3p (see Figure 2D), results shown in Figure 6 demonstrate the SYNCRIP-dependent PCBP2 binding assessed in hepatocytes silenced for SYNCRIP.

Overall, these data indicate that the previously described SYNCRIP capacity to act as EV loader is functionally limited by the presence of a here-identified sequence- and SYNCRIP-dependent retention mechanism mediated by the RNA-binding protein PCBP2.

DISCUSSION

The main finding of this investigation is the identification of PCBP2 as a new regulator of miRNA partition between intracellular and EV compartments. Evidence here gathered indicates that: *i)* PCBP2 binding requires both the miRNA CELL sequence and the RBP SYNCRIP, which in turn recognizes its specific hEXO *consensus*; in other words, SYNCRIP binding is a prerequisite for PCBP2 recruitment; *ii)* PCBP2 cell retention function is dominant over the EV-loading SYNCRIP one; in other word PCPB2 impairs SYNCRIP-mediated miRNA export (Figure 7).

While SYNCRIP EV-loading activity has been previously well characterized by means of both
functional and structural analysis (*7, 9*), the role of PCPB2 as mediator of miRNA intracellular
retention is here disclosed for the first time.
Indeed, no data were previously reported on the RBPs involved in miRNA cell retention even if the
role of specific *consensus* sequences has been previously defined by means of functional assays
involving introduction/removal of the CELL motifs (*5*).
PCBP2 protein (similarly to SYNCRIP) displays a pleiotropic function; specifically, it is a well-
characterized member of the Poly-rC-binding proteins (PCBPs), a group of multifunctional RNA-

binding proteins that contain three highly conserved RNA binding KH domains and that may

shuttle between the nucleus and the cytoplasm (*20*). A large body of evidence, points to its role in

controlling multiple processes including RNA maturation and trafficking, RNA editing,

translational activation or repression and mRNA degradation (*21-26*).

Its potential impact on miRNA biogenesis suggested us to analyze miRNA partition as the EVs/cell

ratio thus circumventing variation deriving from the intracellular expression levels.

Here, PCBP2 was found to directly bind to miRNAs, sharing one of the CELL-motifs previously identified by *in silico* sequence analysis of the miRNAs that were retained in AML12 mouse hepatocytes (*5*). The use of specific insertion/removal mutants and knock-down cellular systems here highlighted the SYNCRIP recruitment as a prerequisite for PCBP2 binding, this verified on miRs-155-3p, 26b-3p and 31-3p. Furthermore, this conclusion was extended by RIP analysis to further 7 miRs selected on the basis of an NGS approach and of a bioinformatic analysis.

Of note, among them, miRs-155-3p, 23a-5p, 155-5p, 192-5p and 26b-3p display important EV-mediated functions in relation to pathophysiology (*27-30*).

The here proposed mechanism implies that the export activity of SYNCRIP is specifically impaired by PCBP2 and highlights that the miRNA partition is not only related to the presence of specific RBPs/export sequences interaction. The final functional compartmentalization output appears the result of an integrated system of RNA/proteins interactions, here only partially unveiled, whose dynamics may provide elements for the explanation of the EVs miRNA cargo specificity and for its variation coherently to cellular plasticity. Of note, recent research highlights a further level of complexity since miRNA epitranscriptomic modifications, while impairing miRNA intracellular function, appear instrumental to miRNA loading in EVs (*4*).

The described multiple RNA/proteins interactions provide a further step in the process of clarification of the mechanisms that may yield value in the control of cellular communication in pathophysiological processes. The knowledge of molecular players of miRNAs intracellular/EVs

partition could be soon instrumental for the development of RNA-based manipulations holding

therapeutic perspectives.

MATERIALS AND METHODS

Cell Culture Conditions

Nontumorigenic murine hepatocyte 3A cells (*3, 31*) were grown at 37°C, in a humidified

atmosphere with 5% CO_2, in RPMI 1640 medium supplemented with 10% FBS (Gibco Life

Technology), 50 ng/mL epidermal growth factor (EGF), 30 ng/mL insulin growth factor (IGF) II

(PeproTech), 10 mg/mL insulin (Roche), and penicillin/streptomycin, on dishes coated with

collagen I (Collagen I, Rat Tail; Gibco Life Technology).

Extracellular vesicle Purification

Extracellular vesicles were prepared according to International Society of Extracellular Vesicles

(ISEV) recommendations(*32*). Conditioned media (CM) from 150 mm plates each containing

250000 hepatocytes were collected after 72-hrs culture in complete medium containing EV-

depleted FBS. Cell-conditioned media were centrifuged at 2000 × g for 20 min at 4°C to remove

dead cells and then at 20000 × g for 30 min at 4°C. Cleared supernatants were passed through 0.22

mm filter membranes, ultracentrifuged in a SW32 Ti rotor at 100,000 rpm for 70 min at 4°C, and

finally resuspended in PBS. The EVs resuspension was analyzed by EXOID-V1-SC (IZON) for size

and concentration characterization.

Biotin miRNA Pull-Down

Biotin miRNA pull-down experiments were performed on cytoplasmic extracts. Briefly, cells were

lysed in hypotonic buffer (10 mM Tris-Cl [pH 7.5], 20 mM KCl, 1.5 mM $MgCl_2$, 5 mM DTT, 0.5

mM EGTA, 5% glycerol, 0.5% NP40, and 40 U/mL RNAsin [Promega]) supplemented with

protease inhibitors (Roche Applied Science). Lysates were incubated on a rotating platform for 30

min at 4°C and then centrifuged at 13000 rpm for 30 min at 4°C. Protein concentration was

determined with Protein Assay Dye Reagent (Bio-Rad) based on the Bradford assay.

Samples (2 mg of proteins) were incubated for 1 hr at 4°C with 10 nmol synthetic single strand

miRNA oligonucleotides containing a biotin modification attached to the 5' and via a spacer arm

(IDT, Intregrated DNA Technology) (Table 1).

Dynabeads™ M-280 Streptavidin (50 μl/sample, Invitrogen™), previously blocked with 1 mg/mL

yeast tRNA (Roche Applied Science), were added to reaction mixture for 90 min at 4°C, and then

the beads were washed three times with cold lysis buffer and once with PBS. Elution was

performed at room temperature for 5 min in Laemli Buffer (containing 2-β mercaptoethanol and

SDS).

Detection of miRNA/RBPs interaction was evaluated by WB on 10% of Input sample and 50% of

the pulled-down samples.

Pull down assay with the recombinant PCBP2 (PCBP2 (NM_001103165) mouse recombinant

protein TP522190, Origene) was performed with 4ug of protein.

Protein Digestion, Peptide Purification and nanoLC Analysis

Proteins obtained from the pull-down experiments with miR-155-3p or random scrambled miRNA

were separated on 4-12% gradient gels (Invitrogen) and stained by Simply Blue Safe Stain staining.

Fourteen sections of the gel lane were cut. Protein digestion of gel pieces and peptide purification

were performed as previously described in (*33*). Peptides resuspended in a suitable nanoLC

injection volume of 2.5% ACN/0.1% TFA and 0.1% formic acid were then analyzed by an UltiMate

3000 RSLCnano-LC system, (Thermo Fisher Scientific) connected on-line via a nano-ESI source to

an Q Exactive plus TM Hybrid Quadrupole-OrbitrapTM Mass Spectrometer (Thermo Fisher

Scientific) as in (*3*). Proteins were automatically identified by MaxQuant (v. 1.6.17.0) software.

Tandem mass spectra were searched against the Mus Musculus dataset of UniprotKB database.

Quantitative comparison among miR-155-3p WT and miR-155-3p no-CELL was performed using

the label-free quantification algorithm calculated by MaxQuant software.

SDS-PAGE and Western Blotting

Cells were lysed in Triton 1X Buffer, subsequently the proteins were analyzed as in (*34*). The

following primary antibodies were used for immunoblotting: α-PCBP2 (AV40568 – Sigma

Aldrich), α- SYNCRIP (MAB11004 – Merck-Millipore), α-HSP90 (sc-13119 - Santa Cruz

Biotech.), α-LAMP1 (Ab24170-Abcam), α-CD63 (sc5275-Santa Cruz Biotech.), α-synthenin

(Ab133267-Abcam), α-CALNEXIN (NB100-1965 – Novus Biologicals), α-GAPDH (MAB-374 -

Merck-Millipore) used as a loading control. The immune complexes were detected with horseradish

peroxidase-conjugated species-specific secondary antiserum: (α-Rabbit 172-1019 and α-Mouse

170-6516 Bio-Rad Laboratories), then by enhanced chemiluminescence reaction (Bio-Rad

Laboratories). Densitometric analysis of protein expression was performed by using the Fiji-Image

J image processing package.

RNA Extraction, RT-PCR and Real-Time qPCR

miRNAs were extracted by miRNeasy Mini Kit and RNeasy MinElute Cleanup Kit (QIAGEN) and

reverse transcribed with MystiCq® microRNA cDNA Synthesis Mix (Sigma-Aldrich). Quantitative

polymerase chain reaction (RT-qPCR) analyses were performed according to MIQE guidelines.

cDNAs were amplified by qPCR reaction using GoTaq qPCR Master Mix (Promega, Madison, WI,

USA). Relative amounts, obtained with $2^{\wedge}(-\Delta Ct)$ method, were normalized with respect to the cel-

miR-39 Spike-In (59000; NORGEN), previously added into miRNA samples.

Total RNA was extracted by ReliaPrep™ RNA Tissue Miniprep System (Promega, USA) and

reverse transcribed with iScriptTM c-DNA Synthesis Kit (Bio-Rad Laboratories Inc., USA).

Quantitative polymerase chain reaction (RT-qPCR) analyses were performed according to MIQE

guidelines. cDNAs were amplified by qPCR reaction using GoTaq qPCR Master Mix (Promega,

Madison, WI, USA). Relative amounts, obtained with $2^{\wedge}(-\Delta Ct)$ method, were normalized with

respect to the housekeeping gene 18S. Oligonucleotide sequences are reported in Table 3.

The results were analyzed with Manager Software (Bio-Rad) and calculated by the $\Delta C(t)$ method.

Co-Immunoprecipitation

Cells were lysed with IP Lysis Buffer (150 mM NaCl, 50 mM Tris-HCl pH 7.5, 5mM EGTA pH 8,

50 mM NaF pH 8, 1,5 mM $MgCl_2$, 1% TRITON-X100 and 10% glycerol) containing freshly added

cocktail protease inhibitors (complete EDTA-free Protease Inhibitor Cocktail; SigmaAldrich) and

phosphatase inhibitors (5 mM EGTA pH 8.0; 50 mM sodium fluoride; 5 mM sodium

orthovanadate). Lysates were incubated on a rotating platform for 2h at 4°C and then centrifuged at

13000 rpm for 30 min at 4°C. Protein concentration was determined with Protein Assay Dye

Reagent (Bio-Rad), based on the Bradford assay.

2 mg of proteins (one for the specific antibody and one for the corresponding aspecific IgG) were

precleared adding 40 µL of Protein A Sepharose or Protein G Sepharose (GE HealthCare) for 3 hrs

at 4°C in a total volume of 1 ml of IP Lysis Buffer in rotation. Then, Protein A or G Sepharose was

removed by centrifugation and the extracts were incubated with 5 µg of specific antibody α-PCBP2

(cod. RN025P - MBL), SYNCRIP (MAB11004, Merck-Millipore), Normal Rabbit IgG (12-370-

Millipore) or Normal Mouse IgG (12-371-Millipore), the last two used as negative controls, to

proceed with immunoprecipitation at 4°C overnight. Immuno-complexes were collected adding 50

µL of Protein A or G Sepharose for 3 hrs at 4°C in rotation. The immunoprecipitated proteins were

washed three times with Net Gel Buffer (150 mM NaCl, 50 mM Tris-HCl pH 7.5, 1mM EDTA,

0.1% NP40 and 0.25% gelatin) and once with RIPA Buffer (150 mM NaCl, 1% NP40, 0,5%

Sodium Deoxycholate, 0,1% SDS, 50 mM Tris HCl pH 8). Finally, immunoprecipitated proteins

were detached from Sepharose beads by adding 50 µL of Laemli Buffer 2X. Samples were boiled at

95°C for 5 min, beads were eliminated by centrifugation and 10% of input sample and 50% of each

immunoprecipitated sample were loaded on polyacrilammide gel and analyzed by Western Blotting.

EMSA

Cells were lysed in Triton Buffer at 4°C, for 30 min and 4 µg of protein extract were incubated with

0.5 pmol of biotinylated RNA oligonucleotides for 30 min at room temperature in REMSA Binding

Buffer, according to the manufacturer's protocol (Light Shift Chemiluminescent RNA EMSA Kit,

ThermoScientific 20158). 1 µg of each antibody was incubated with the protein-RNA complex:

anti-PCBP2 (RN025P; MBL), anti-SYNCRIP (MAB11004; Merck-Millipore) for supershift and

ultrashift analysis. The electrophoresis was performed in native 6% polyacrylamide gel in 0,5X

TBE. Transfer step was carried out at 25V, for 15 min in 0,5X TBE and the detection was

performed following manufacturer's instructions.

UV Cross-Linking RIP

CLIP was performed as reported in (*34*). Immunoprecipitated miRNAs were reverse transcribed and

analyzed by RT-qPCR amplifications. List of primers is reported in Table 2. Primary antibodies for

IP: anti-PCBP2 (RN025P; MBL), anti-SYNCRIP (MAB11004; Merck-Millipore) and as negative

controls Normal Rabbit IgG (12-370; Merck-Millipore) or Normal Mouse IgG (12-371; Merck-

Millipore).

shRNA Silencing

Stable PCBP2 knockdown was achieved through infection with shRNAs cloned in pSUPER retro

puro retroviral vector (Oligoengine). Viral supernatants were collected 48 hrs after transfection of

293gp packaging cells, filtered (0.45 mm), and added to hepatocytes. At 48 hrs post-infection,

selection was performed with 2 μg/mL puromycin for at least 1 week before analysis. The sequence

of shRNA scramble used as control was previously described(*35*). The sequences of shRNA oligos

used for cloning are reported in Table 4.

Motif Scanning Analysis

Murine mature miRNA sequences were retrieved from miRBase v22.1 database (*36*). The FIMO

tool (*37*) was used to scan these sequences for occurrences of hEXO, extended CELL (the bottom

motif identified in AML12 cells and reported in figure 2 by (*5*)) and core AUUA/G CELL motifs,

encoded as Position Probability Matrices, with parameters --bfile --motif-- --norc and setting the p-

value threshold to 0.1, 0.1 and 0.01, respectively. Motif instances falling in the seed regions

(nucleotides 2-7) were ignored.

Small RNA Sequencing

miRNA samples (two biological replicates per condition), to which the cel-miR-39 Spike-In

(59000; NORGEN) was previously added, were sequenced at Procomcure Biotech GmbH.

Sequencing libraries were prepared using the NEXTFLEX Small RNA-Seq Kit v4 (PerkinElmer).

The sequencing reaction was performed on an llumina NovaSeq 6000 instrument in 2x40bp paired-

end configuration, with a throughput of ~40 million read pairs per sample. FastqToolkit version

2.2.5 (available at https://www.illumina.com/products/by-type/informatics-products/basespace-

sequence-hub/apps/fastq-toolkit.html) was used to remove adapter sequences from the 3' end and to

filter out reads whose length and average quality after trimming were < 10 and < 30, respectively.

Only forward reads were kept for downstream analyses. The mirPRo software version 1.1.4 (*38*)

which utilizes NovoAlign (*39*) as its alignment engine, was used to align reads to a reference

composed of miRNA hairpin sequences downloaded from miRBase v22.1 database with the

addition of the spike-in, and to count reads mapping to mature miRNAs. A count matrix was

assembled, including only mature miRNAs with one or more reads in at least two cell and two EV

samples. Differential abundance analysis was performed using the DESeq2 R package (*40*). Size

factors were estimated directly from spike-in counts. For each mature miRNA, a likelihood ratio

test was conducted to assess differences between the EV/cell abundance ratios measured in the

shPCBP2 and shCTR conditions.

Statistical Analyses

For the qRT-PCR analysis, statistical differences were assessed with the one-tailed paired Student's

t-test using GraphPad Prism Version 9 (GraphPad Software). Data are presented as mean ± SEM,

and p values < 0.05 were considered statistically significant. For the statistical analysis of proteomic

studies, Perseus software (version 1.6.7.0) after log2 transformation of the intensity data was used.

Results were considered statistically significant at $p<0.05$.

Data availability

      The miRNA-seq data generated in this study have been deposited and are available in the GEO

      database under accession code GSE269709.

AKNOWLEDGEMENTS

We thank Prof. Andres Ramos for suggestions and critical reading of the manuscript, Andrea

Melito, Cristina Mordenti and Claudia Maldonado Torres for their help in molecular analyses and

data collection, Giovanna Sabarese for early stages experiments. S.G. is supported by the AIRC

Post-Doctoral Fellowship. This work was funded by Associazione Italiana per la Ricerca sul Cancro

(IG26290) to M.T. and by the Italian Ministry of Health "Ricerca Corrente - Linea 3 - Progetto 2-

INMI L. Spallanzani I.R.C.C.S." funding to C.M.; Istituto Pasteur-Fondazione Cenci-Bolognetti

(Anna Tramontano grant 2020) to C.C.; European Research Council (RIBOMYLOME_309545 and

ASTRA_855923) and the H2020 projects (IASIS_727658 and INFORE_825080) to G.G.T.;

Sapienza University of Rome (RM12218166AEFC72) to C.B.; SEED PNR-Finanziamento di

progetti di ricerca su temi di interesse trasversale per il PNR 2021 to C.B.; PNRR-Rome

Technopole to M.T. and PRIN: progetti di ricerca di rilevante interesse nazionale – Bando 2022

Prot. 2022ETPX42 to M.T.

REFERENCES

1. G. van Niel *et al.*, Challenges and directions in studying cell-cell communication by extracellular
vesicles. *Nat Rev Mol Cell Biol* **23**, 369-382 (2022).
2. M. Mathieu, L. Martin-Jaular, G. Lavieu, C. Thery, Specificities of secretion and uptake of exosomes
and other extracellular vesicles for cell-to-cell communication. *Nat Cell Biol* **21**, 9-17 (2019).
3. C. Montaldo *et al.*, Fibrogenic signals persist in DAA-treated HCV patients after sustained virological
response. *J Hepatol* **75**, 1301-1311 (2021).
4. S. Garbo *et al.*, m6A modification inhibits miRNAs' intracellular function, favoring their extracellular
export for intercellular communication. *Cell Rep* **43**, 114369 (2024).
5. R. Garcia-Martin *et al.*, MicroRNA sequence codes for small extracellular vesicle release and cellular
retention. *Nature* **601**, 446-451 (2022).
6. M. A. Mori, R. G. Ludwig, R. Garcia-Martin, B. B. Brandao, C. R. Kahn, Extracellular miRNAs: From
Biomarkers to Mediators of Physiology and Disease. *Cell Metab* **30**, 656-673 (2019).
7. L. Santangelo *et al.*, The RNA-Binding Protein SYNCRIP Is a Component of the Hepatocyte Exosomal
Machinery Controlling MicroRNA Sorting. *Cell Rep* **17**, 799-808 (2016).
8. C. Villarroya-Beltri *et al.*, Sumoylated hnRNPA2B1 controls the sorting of miRNAs into exosomes
through binding to specific motifs. *Nat Commun* **4**, 2980 (2013).
9. F. Hobor *et al.*, A cryptic RNA-binding domain mediates Syncrip recognition and exosomal
partitioning of miRNA targets. *Nat Commun* **9**, 831 (2018).
10. S. Garbo, R. Maione, M. Tripodi, C. Battistelli, Next RNA Therapeutics: The Mine of Non-Coding. *Int J
Mol Sci* **23**, (2022).
11. K. Paunovska, D. Loughrey, J. E. Dahlman, Drug delivery systems for RNA therapeutics. *Nat Rev
Genet* **23**, 265-280 (2022).
12. G. M. Traber, A. M. Yu, RNAi-Based Therapeutics and Novel RNA Bioengineering Technologies. *J
Pharmacol Exp Ther* **384**, 133-154 (2023).
13. K. B. Cook, H. Kazan, K. Zuberi, Q. Morris, T. R. Hughes, RBPDB: a database of RNA-binding
specificities. *Nucleic Acids Res* **39**, D301-308 (2011).
14. J. Y. Liao *et al.*, EuRBPDB: a comprehensive resource for annotation, functional and oncological
investigation of eukaryotic RNA binding proteins (RBPs). *Nucleic Acids Res* **48**, D307-D313 (2020).
15. D. Ray *et al.*, A compendium of RNA-binding motifs for decoding gene regulation. *Nature* **499**, 172-
177 (2013).
16. T. Treiber *et al.*, A Compendium of RNA-Binding Proteins that Regulate MicroRNA Biogenesis. *Mol
Cell* **66**, 270-284 e213 (2017).
17. E. L. Van Nostrand *et al.*, A large-scale binding and functional map of human RNA-binding proteins.
Nature **583**, 711-719 (2020).

18.   B. D. Fonseca *et al.*, La-related Protein 1 (LARP1) Represses Terminal Oligopyrimidine (TOP) mRNA
Translation Downstream of mTOR Complex 1 (mTORC1). *J Biol Chem* **290**, 15996-16020 (2015).
19.   L. Wickham, T. Duchaine, M. Luo, I. R. Nabi, L. DesGroseillers, Mammalian staufen is a double-
stranded-RNA- and tubulin-binding protein which localizes to the rough endoplasmic reticulum.
Mol Cell Biol **19**, 2220-2230 (1999).
20.   C. Yuan, M. Chen, X. Cai, Advances in poly(rC)-binding protein 2: Structure, molecular function, and
roles in cancer. *Biomed Pharmacother* **139**, 111719 (2021).
21.   A. Chaudhury, P. Chander, P. H. Howe, Heterogeneous nuclear ribonucleoproteins (hnRNPs) in
cellular processes: Focus on hnRNP E1's multifunctional regulatory roles. *RNA* **16**, 1449-1462
(2010).
22.   T. Ishii *et al.*, PCBP1 and PCBP2 both bind heavily oxidized RNA but cause opposing outcomes,
suppressing or increasing apoptosis under oxidative conditions. *J Biol Chem* **295**, 12247-12261
(2020).
23.   A. V. Makeyev, A. N. Chkheidze, S. A. Liebhaber, A set of highly conserved RNA-binding proteins,
alphaCP-1 and alphaCP-2, implicated in mRNA stabilization, are coexpressed from an intronless
gene and its intron-containing paralog. *J Biol Chem* **274**, 24849-24857 (1999).
24.   A. V. Makeyev, S. A. Liebhaber, The poly(C)-binding proteins: a multiplicity of functions and a search
for mechanisms. *RNA* **8**, 265-278 (2002).
25.   D. Perrotti *et al.*, BCR-ABL suppresses C/EBPalpha expression through inhibitory action of hnRNP
E2. *Nat Genet* **30**, 48-58 (2002).
26.   T. Yabe-Wada, C. C. Philpott, N. Onai, PCBP2 post-transcriptionally regulates sortilin expression by
binding to a C-rich element in its 3' UTR. *FEBS Open Bio* **10**, 407-413 (2020).
27.   A. Broermann *et al.*, Exosomal miRNAs as Potential Biomarkers to Monitor Phosphodiesterase 5
Inhibitor Induced Anti-Fibrotic Effects on CCl(4) Treated Rats. *Int J Mol Sci* **22**, (2020).
28.   S. G. Dosil *et al.*, Natural killer (NK) cell-derived extracellular-vesicle shuttled microRNAs control T
cell responses. *Elife* **11**, (2022).
29.   Y. Jiao, P. Xu, H. Shi, D. Chen, H. Shi, Advances on liver cell-derived exosomes in liver diseases. *J Cell
Mol Med* **25**, 15-26 (2021).
30.   Y. Zhao, J. Liu, Z. Xiong, S. Gu, X. Xia, Exosome-derived miR-23a-5p inhibits HCC proliferation and
angiogenesis by regulating PRDX2 expression: MiR-23a-5p/PRDX2 axis in HCC progression. *Heliyon*
10, e23168 (2024).
31.   A. Conigliaro, Amicone, L., Costa, V., De Santis Puzzonia, M., Mancone, C.,Sacchetti, B., Cicchini, C.,
Garibaldi, F., Brenner, D.A., Kisseleva, T., et al., in *(2013)*.
32.   W. K. Théry C, Aikawa E, Alcaraz MJ, Anderson JD, Andriantsitohaina R, Antoniou A, Arab T, Archer
F, Atkin-Smith GK, Ayre DC, Bach JM, Bachurski D, Baharvand H, Balaj L, Baldacchino S, Bauer NN et
al.. in *J Extracell Vesicles. 2018 Nov 23;7(1):1535750. doi: 10.1080/20013078.2018.1535750.
eCollection 2018.*
33.   C. Mancone *et al.*, Ferritin heavy chain is the host factor responsible for HCV-induced inhibition of
apoB-100 production and is required for efficient viral infection. *J Proteome Res* **11**, 2786-2797
(2012).
34.   C. Battistelli *et al.*, Design and Functional Validation of a Mutant Variant of the LncRNA HOTAIR to
Counteract Snail Function in Epithelial-to-Mesenchymal Transition. *Cancer Res* **81**, 103-113 (2021).
35.   G. A. Pasque V, Garrett N and Gurdon JB, in *EMBO J. 30, 2373–2387. (2011)*.
36.   A. Kozomara, M. Birgaoanu, S. Griffiths-Jones, miRBase: from microRNA sequences to function.
Nucleic Acids Res **47**, D155-D162 (2019).
37.   C. E. Grant, T. L. Bailey, W. S. Noble, FIMO: scanning for occurrences of a given motif.
Bioinformatics **27**, 1017-1018 (2011).
38.   J. Shi *et al.*, mirPRo-a novel standalone program for differential expression and variation analysis of
miRNAs. *Sci Rep* **5**, 14617 (2015).
39.   H. Li, N. Homer, A survey of sequence alignment algorithms for next-generation sequencing. *Brief
Bioinform* **11**, 473-483 (2010).

40.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biol* **15**, 550 (2014).

DECLARATION OF INTERESTS

The authors declare no competing interests.

SUPPLEMENTARY INFORMATION

Supplementary Information includes 2 figures and 3 data files.

331 **AUTHOR CONTRIBUTIONS**.

332 Molecular and biochemical analysis: F.M., S.G., L.Q., G.G. Proteomic analysis: F.M. and C.M.

333 Bioinformatics: A.C. and G.G.T. The project was designed by C.C., C.B. and M.T. The paper was

334 written by C.B. and M.T. and revised by all other authors.

FIGURES

A

miR-155-3p WT
CUCCUACCUGUUA GCAU UAAC

miR-155-3p no-CELL
CUCCUACCUGUUA GCAU GAUC

miR-155-3p no-hEXO
CUCCUACCUGUUA GCAU UAGU

Figure 1: PCBP2 recognizes the CELL motifs and has a functional role in intracellular retention of miRNA155 -3p

351 **Figure 1: PCBP2 recognizes the CELL motifs and has a functional role in intracellular**

352 **retention of miRNA-155-3p.**

A) Sequences of biotinylated oligos used as bait in pulldown experiment; CELL motifs (WT and mutated) are in grey, hEXO motifs (WT and mutated) are underlined. miRNA devoid of CELL motif (no-CELL), miRNA devoid of hEXO (no-hEXO).

B) Volcano plot comparing proteins bound to miRNA-155-3p no-CELL vs WT. Black curves represent the significant threshold at an a false-discovery rate (FDR) of 0.05 and S0 of 0.1. PCBP2 and SYNCRIP proteins are labeled in the plot.

C) CLIP of PCBP2 protein in murine hepatocytes. RT-qPCR analysis for miR-155-3p, miR-365-2-5p (CELL motif-devoid) and miR-31-3p (hEXO motif-devoid) is shown as IP/IgG. Data are the mean± SEM of three independent experiments.

D) RNA pull-down with the WT and mutated (no-CELL, no-hEXO) (sequences are reported in A) miR-155-3p followed by western blot for the indicated proteins (HSP90 is used as positive and GAPDH as negative controls respectively). Data are representative of three independent experiments.

E) RNA pull-down with the miR-365-2-5p followed by western blot for the indicated proteins. Data are representative of three independent experiments.

F) RNA pull-down by using the recombinant PCBP2 protein and with WT and mutated miR-155-3p (no-CELL) followed by western blot for PCBP2. Data are representative of three independent experiments.

G) CLIP of SYNCRIP protein in murine hepatocytes. RT-qPCR analysis for miR-155-3p is shown as IP/IgG. Data are the mean± SEM of three independent experiments.

H) (left and middle panels) EV miRNA-155-3p and miR-365-2-5p levels in shCTR and shPCBP2 cells analyzed by RT-qPCR. Data are expressed as ratio of miRNA expression in EVs with respect to the intracellular compartment (shCTR arbitrary value 1). Results are shown as the mean±S.E.M. of three independent experiments.

359 (right panel) EV miRNA-155-3p levels in shCTR and shSYNCRIP cells analyzed by RT-qPCR.

360 Data are expressed as ratio of miRNA expression in EVs with respect to the intracellular

361 compartment (shCTR arbitrary value 1). Results are shown as the mean±S.E.M. of three

362 independent experiments.

363 Data are considered statistically significant with $p < 0.05$ (Student's T test). *: $p < 0.05$; **: $p < 0.01$.

A

B

C **D**

miR-155-3p

Figure 2: PCBP2 binding to miR-155-3p is SYNCRIP-dependent

Figure 2: PCBP2 binding to miR-155-3p is SYNCRIP-dependent

A) Co-immunoprecipitation of PCBP2 and SYNCRIP. Immunoprecipitations with rabbit polyclonal anti-PCBP2, mouse monoclonal anti-SYNCRIP and the relative preimmune IgG were performed on protein extracts from hepatocytes. GAPDH is used as negative control. Immunoblots representative of three independent experiments are shown.

B) Electrophoretic mobility shift assay (EMSA): interactions of miR-155-3p with the indicated protein extracts (shifts) and Abs (anti-SYNCRIP and anti-PCBP2) (supershift) are shown. Ultrashift shown in lane 5 demonstrates concurrent binding of SYNCRIP and PCBP2 to miR-155-3p.

C) Electrophoretic mobility shift assay (EMSA): interactions of miR-155-3p with protein extracts from shCTR (1), shPCBP2 (2) and shSYNCRIP (3) cells (shifts) and Abs (anti-SYNCRIP and anti-PCBP2) (supershift) are shown.

D) CLIP of PCBP2 protein in murine hepatocytes both WT (shCTR) and silenced for SYNCRIP (shSYNCRIP). RT-qPCR analysis for the expression of miR-155-3p is shown as IP/IgG. Data are the mean± SEM of three independent experiments.

A

miR 26b-3p WT
CCUGUUCUCCAUUACUUGGCUC

miR 26b-3p no-hEXO
CCUGUUCUCCAUUACUUGGCGA

B

miR 31-3p WT
UGCUAUGCCAACAUAUUGCCAUC

miR 31-3p +hEXO
UGCUAUGCCAACAUAUUGGCUG

miR 31-3p +hEXO no-CELL
UGCUAUGCCAACAUAUUUGGCUG

397   **Figure 3: PCBP2 binding to miR-155-3p is sequence dependent**

Figure 3: PCBP2 binding to miR-155-3p is sequence dependent

A) RNA pull-down with the WT and mutated (sequences are reported above) miR-26b-3p followed by western blot for the indicated proteins. Data are representative of three independent experiments.

B) RNA pull-down with the WT and mutated (sequences are reported above) miR-31-3p followed by western blot for the indicated proteins. Data are representative of three independent experiments.

398   Data are considered statistically significant with p< 0.05 (Student's T test). *: p<0.05.

A-B) CELL motifs (WT and mutated) are in grey, hEXO motifs (WT and mutated) are underlined.

Figure 4: PCBP2 functionally dominates on SYNCRIP EV-loading activity on a repertoire of miRNAs embedding CELL and hEXO motifs

Figure 4: PCBP2 functionally dominates on SYNCRIP EV-loading activity on a repertoire of miRNAs embedding CELL and hEXO motifs

A) Volcano plot comparing miRNAs differently expressed from NGS data; miRNAs with Log2FC > 1 and Log2FC < -1 and p-value ≤ 0.10 were considered differently expressed. Downregulated miRNAs in shPCBP2 respect to shCTRL are represented as blue dots, upregulated miRNAs are represented as red dots.

B) Heatmap showing the Log2 fold enrichment (EV/CELL) of mature miRNAs in small extracellular vesicles derived from shCTRL cells versus shPCBP2 cells. miRNAs with Log2FE ≥ 1.0 and p-value ≤ 0.10 were considered to be differentially enriched.

C) List of selected miRNAs embedding CELL and/or hEXO motifs; consensus sequences are highlighted in grey or underlined respectively.

D) EV miRNA levels in shCTR and shPCBP2 cells analyzed by RT-qPCR. Data are expressed as
ratio of miRNA expression in EVs with respect to the intracellular compartment (shCTR arbitrary
value 1). Results are shown as the mean±S.E.M. of three independent experiments.

Data are considered statistically significant with p< 0.05 (Student's T test). *: p<0.05; **: p<0.01.

E) EV miRNA levels in shCTR and shSYNCRIP cells analyzed by RT-qPCR. Data are expressed
as ratio of miRNA expression in EVs with respect to the intracellular compartment
(shCTR arbitrary value 1). Results are shown as the mean±S.E.M. of three independent
experiments.

Data are considered statistically significant with p< 0.05 (Student's T test). *: p<0.05; **: p<0.01;
****: p<0.0001

Figure 5: PCBP2 and SYNCRIP bind to several miRNAs embedding CELL and hEXO motif sequences

Figure 5: PCBP2 and SYNCRIP bind to several miRNAs embedding CELL and hEXO motif sequences

A) CLIP of PCBP2 protein in murine hepatocytes. RT-qPCR analysis for the indicated miRNAs is shown as IP/IgG for each independent experiment (IgG arbitrary value 1). Data are the mean± SEM of three independent experiments.

Data are considered statistically significant with $p < 0.05$ (Student's T test). *: $p<0.05$; **: $p<0.01$.

B) CLIP of SYNCRIP protein in murine hepatocytes. RT-qPCR analysis for the indicated miRNAs is shown as IP/IgG for each independent experiment (IgG arbitrary value 1). Data are the mean± SEM of three independent experiments.

Data are considered statistically significant with $p < 0.05$ (Student's T test). *: $p<0.05$; **: $p<0.01$; ***: $p<0.001$.

Figure 6: PCBP2 binding to miRNAs requires SYNCRIP

Figure 6: PCBP2 binding to miRNAs requires SYNCRIP

CLIP of PCBP2 protein in murine hepatocytes both WT (shCTR) and silenced for SYNCRIP

(shSYNCRIP). RT-qPCR analysis for the indicated miRNAs is shown as IP/IgG. Data are the

mean± SEM of three independent experiments.

Data are considered statistically significant with $p < 0.05$ (Student's T test). *: $p<0.05$; ***:

$p<0.001$.

Figure 7: Schematic model of PCBP2/SYNCRIP dependent miRNA compartmentalization

Figure 7: Schematic model of PCBP2/SYNCRIP dependent miRNAs compartmentalization.

Left) hEXO-SYNCRIP interaction promotes miRNAs secretion into EVs.

Right) SYNCRIP-dependent PCBP2-CELL motif interaction promotes miRNAs intracellular

retention.

TABLES

Name	Oligonucleotides sequence
Biotin-miR-26b-3p WT	[Btn] 5' CCUGUUCUCCAUUACUUGGCUC 3'
Biotin-miR-26b-3p no-hEXO	[Btn] 5' CCUGUUCUCCAUUACUUGGCGA 3'
Biotin-miR-31-3p WT	[Btn] 5' UGCUAUGCCAACAUAUUGCCAUC 3'
Biotin-miR-31-3p+ hEXO	[Btn] 5' UGCUAUGCCAACAUAUUGGGCUG 3'
Biotin-miR-31-3p +hEXO no-CELL	[Btn] 5' UGCUAUGCCAACAUAUUUGGCUG 3'
Biotin-miR-155b-3p WT	[Btn] 5' CUCCUACCUGUUAGCAUUAAC 3'
Biotin-miR-155b-3p no-CELL	[Btn] 5' CUCCUACCUGUUAGCAUGAUC 3'
Biotin-miR-155b-3p no-hEXO	[Btn] 5' CUCCUACCUGUUAGCAUUAGU 3'
Biotin-miR-365-2-5p WT	[Btn] 5' AGGGACUUUCAGGGGCAGCUGUG 3'

495 **Table 1.** Biotinylated RNA oligonucleotides used in pull down experiments

miRNA	Primer sequence	Tm (ºC)
mmu-miR-23a-5p	GGGGTTCCTGGGGATGGGATTT	60
mmu-miR-26b-3p	CCTGTTCTCCATTACTTGGCTC	62
mmu-miR-31-3p	TGCTATGCCAACATATTGCCATC	61
mmu-miR-122b-3p	AAACACCATTGTCACACTCCAC	60
mmu-miR-155- 3p	CTCCTACCTGTTAGCATTAAC	57

mmu-miR-155-5p	TTAATGCTAATTGTGATAGGGGT	58
mmu-miR-181d-5p	AACATTCATTGTTGTCGGTGGGT	60
mmu-miR-192-5p	CTGACCTATGAATTGACAGCC	59
mmu-miR-214-3p	ACAGCAGGCACAGACAGGCAGT	60
mmu-miR-345-3p	CCCTGAACTAGGGGTCTGGAGAC	60
mmu-miR-365-2-5p	GACTTTCAGGGGCAGCTG	58
mmu-miR-3084-5p	GTTGAAGGTTAATTAGCAGAGT	60

497 **Table 2.** Primers for miRNA qPCR analysis.

Name	Primer sequence	Tm (°C)
PCBP2	*For* ACACCGGATTCAGTGGCA	58
	Rev TTGATTTTGGCGCCTTGACG	58
SYNCRIP	*For* ACCTTGCCAACACGTAACA	59
	Rev CCATAGCCTTGACACACCA	59
18s	*For* AGCACCCATTGCAACGTCTG	58
	Rev GCACGGCGACTACCATCG	58

499 **Table 3.** Primers for gene expression qPCR analysis.

Name	Sequence
PCBP2	*Sense* GATCCCCGAGCAGACCCATCCATAATTTCAAGAGAATTATGGATGGGT CTGCTCTTTTTA *Antisense* AGCTTAAAAAGAGCAGACCCATCCATAATTCTCTTGAAATTATGGATG

	GGTCTGCTCGGG
SYNCRIP	As reported in (*7*).
CTR	As reported in (*7*).

502   **Table 4.** Oligos for shRNA cloning in pSUPER.retro.puro vector.

542   **SUPPLEMENTARY INFORMATION**

Supplementary Figure 1: PCBP2 and SYNCRIP silencing

A

Particle diameter (nm)

Mean:	204 (Std Dev=56.5)	d50: 189	
Mode:	183	d10: 160	d90: 253
Maximum:	910	d90/d10: 1.58	
Minimum:	150	Span: 0.49	

Concentration [0-∞] (particles/ml)

Measured Concentration: 2.20e+09
Raw Concentration: 4.40e+10

EVs shCTR

Particle diameter (nm)

Mean:	219 (Std Dev=100.8)	d50: 182	
Mode:	166	d10: 152	d90: 325
Maximum:	972	d90/d10: 2.13	
Minimum:	146	Span: 0.94	

Concentration [0-∞] (particles/ml)

Measured Concentration: 6.11e+08
Raw Concentration: 1.22e+10

EVs shPCBP2

Particle diameter (nm)

Mean:	231 (Std Dev=83.7)	d50: 199	
Mode:	176	d10: 168	d90: 333
Maximum:	821	d90/d10: 1.98	
Minimum:	161	Span: 0.83	

Concentration [0-∞] (particles/ml)

Measured Concentration: 1.33e+09
Raw Concentration: 6.66e+09

EVs shSYNCRIP

B

WCE   EVs

LAMP1
CD63
SYNTHENIN
CALNEXIN

Supplementary Figure 2: shCTR, shPCBP2 and shSYNCRIP EV characterization

Supplementary Figure 1: PCBP2 and SYNCRIP silencing

a) Expression levels of PCBP2 in shCTR and shPCBP2 murine hepatocytes. Data are shown as the

mean±S.E.M. of three independent experiments.

b) (Left panel) Western-blot analysis for PCBP2 on protein extracts from hepatocytes silenced for

PCBP2 (3A shPCBP2) and relative control (3A shCTR). GAPDH has been used as loading control.

The figure is representative of three independent experiments. (Right panel) Densitometric analysis

of Western-blot signals. Data are shown as the mean±S.E.M. of three independent experiments.

c) Expression levels of SYNCRIP in shCTR and shSYNCRIP murine hepatocytes. Data are shown

as the mean±S.E.M. of three independent experiments.

d) (Left panel) Western-blot analysis for SYNCRIP on protein extracts from hepatocytes silenced

for SYNCRIP (3A shSYNCRIP) and relative control (3A shCTR). GAPDH has been used as

loading control. The figure is representative of three independent experiments. (Right panel)

Densitometric analysis of Western-blot signals. Data are shown as the mean±S.E.M. of three

independent experiments.

Supplementary Figure 2: shCTR, shPCBP2 and shSYNCRIP EV characterization

a) Particle diameter (nm) and concentration (particles/ml) of EVs evaluated by Exoid (IZON) (Top:

shCTR EVs, Middle: shPCBP2 EVs, Bottom: shSYNCRIP EVs).

d) Western-blot analysis for EV-specific (LAMP1, CD63, Synthenin) and intracellular (calnexin)

markers on protein extracts from hepatocytes (WCE, whole cell extract) and hepatocyte-derived

EVs (EVs).

Supplementary Data File 1:

Results of proteomic analysis of proteins bound to miRNA-155-3p no-CELL vs WT.

Supplementary Data File 2:

Results of proteomic analysis of proteins differentially bound to miRNA-155-3p no-CELL vs WT.

Supplementary Data File 3:

Results of the likelihood ratio test performed on small RNA Sequencing data to assess the

differential miRNA EV loading between shPCBP2 and shCTR conditions.