Title: Symbiosis collapses during development of asexual offspring in the absence of

heterotrophic feeding in a model cnidarian-algal symbiosis

Authors: Samuel A. Bedgood[1,3], Keyla Plichon[2,3], Virginia M. Weis[3]

Affiliations:

[1]New College of Florida, Division of Natural Sciences, Sarasota, FL, United States of America

[2]Université Côte d'Azur, CNRS, INSERM, Institute for Research on Cancer and Aging, Nice,

France

[3]Department of Integrative Biology, Oregon State University, Corvallis, OR, United States of

America

Corresponding author:

Samuel A. Bedgood

Email: sbedgood@ncf.edu

ORCID iD: 0000-0002-2510-0900

Acknowledgements

We thank Ruby Miller and Felix Jardini for their assistance in carrying out the experiment,

caring for the animals, and counting thousands of tentacles. We also thank Sofia Blaire for

contributing to the project and conducting a parallel experiment that informed the design of this

study. We appreciate the guidance of Dr. Nathan Kirk and Dr. Keith Garrison in sequencing.

Additionally, we thank the members of the Weis Lab for their valuable input on the experimental

design and analysis. Finally, we acknowledge the Center for Quantitative Life Sciences at

Oregon State University for their confocal microscopy and sequencing services, which were

essential to this work.

Funding

This research was supported by the National Science Foundation (NSF) through a grant awarded

to Virginia M. Weis (NSF Grant No. IOS2124119). The funding agency had no involvement in

the study design, data collection, analysis, or interpretation of the results, nor in the decision to

submit the manuscript for publication.

Abstract

A stable symbiosis between corals and dinoflagellate algae is crucial for coral reef health, and it

is driven by nutrient exchange and environmental interactions. Our understanding of the

homeostasis between host cnidarian and algal symbiont during the host adult stage is a longtime

area of focus, but little is known about the balance of partners during development and

regeneration. We investigated the role of symbiotic algae and heterotrophic feeding on

development in the sea anemone model organism commonly called Aiptasia. We focused on

asexually-produced offspring (G1), examining the effects of autotrophic and heterotrophic

nutrition on developmental rates. We found that the presence of symbionts enhanced growth in

fed conditions but impeded development and survival under starvation. The effect of symbiont

presence on starved offspring was dose-dependent where those offspring with more symbionts at

an earlier stage lost tentacles and mass faster than those with fewer symbionts. We hypothesize

that suppression of immunity during development explains the patterns we observe. Our results

provide insight into the metabolic costs and benefits of symbiosis under different nutritional

conditions during development and regeneration of symbiotic cnidarians.

Key words: Aipatsia, Host-symbiont dynamics, Development, Nutrient Exchange, Immunity

1. Introduction

Symbiosis between corals and dinoflagellate algae (Family Symbiodiniaceae) is foundational to a healthy coral reef ecosystem, and this relationship is based on nutrient exchange between partners [1–3]. The checks and balances to this relationship are complex and only partly understood. The homeostasis between host (coral) and symbiont (algae) keeps the biomass of each in check via four processes: 1) nutrient dynamics where the host controls nitrogen availability to the algae and the symbionts provide photosynthetically-derived sugars and lipids to the host [4–6], 2) host immune regulation of symbionts [7,8], 3) symbiont population dynamics and cell cycle regulation in host tissue [9], and 4) changing environmental factors such as light and temperature which affect the host's ability to maintain homeostasis and the symbiont's survival and growth [10–12]. Changes in one or more of these factors can result in a move from symbiosis to dysbiosis whereby algae are lost from the host [13]. Environmental perturbations, due to climate change, are occurring on a global scale causing the degradation of coral reefs through breakdown in symbiosis [14]. It is therefore critical to understand the functioning of this symbiosis under different stressors and life stages of the host.

The role of the host developmental stage on the establishment of host cnidarian-algae homeostasis has been overlooked. This is especially true in asexually-produced offspring, which are an important part of most cnidarian life history strategies [15,16]. Reef-building corals produce massive colonies via asexual polyp budding and colony fragmentation [17]. Homeostasis between host and symbiont must be maintained throughout the process with the parent polyp(s) passing algae to the asexual offspring.

Exaiptasia diaphana, a sea anemone commonly known as Aiptasia, is a well-established model for the study of coral-algal symbiosis [18]. It reproduces both sexually and asexually.

Sexual reproduction occurs through broadcast spawning of gametes, resulting in planula larvae

that settle and develop into polyps [19–21]. These F1 offspring produced by F0 parents acquire

their symbionts horizontally by ingesting algae during the planula or juvenile polyp stage (Fig. 1,

[21]). Asexual reproduction occurs through pedal laceration, where adult anemones, referred to

here as G0, leave behind small pieces of tissue from the pedal disk that develop into G1 offspring

[22,23]. Unlike F1 offspring, G1 offspring inherit their symbionts vertically, directly from the

G0 adult (Fig. 1; [22]). Each pedal lacerate is on average 0.5 mm in diameter (F [23], *pers. obs.*),

but contains hundreds of algae from the parent (*pers. obs.*, X.X.X.), which make up as much as

55-60% of the dry mass of a pedal lacerate [22]. Aposymbiotic pedal lacerates, like aposymbiotic

adults, can acquire symbionts from the environment, once they have developed a mouth.

Presnell et al. [23] showed that the presence of symbionts affects the developmental rate

of G1 offspring when starved; aposymbiotic pedal lacerates develop more rapidly than symbiotic

in the first few days of development. However, the growth rate in symbiotic offspring overtakes

that of aposymbiotic offspring 20 days after separation from G0 anemones [23]. The symbiosis

becomes beneficial only after an initial dampening of growth rate. The role of heterotrophic

feeding as an alternative or additive source of nutrition and the role of timing and onset of

symbiosis have not been explored in these early developmental stages or in later stages of

development beyond three weeks.

In this study, we extended the timeframe beyond the three week window to explore the

impact of nutrition, both from symbiotic photosynthate and heterotrophic feeding, on

developmental rates. We also introduced the variable of horizontal symbiont transmission

following laceration to determine effects from symbionts and nutrition during early- and late-

stage development. We hypothesized that starved symbiotic G1 juveniles would exhibit a more

rapid growth rate compared to their aposymbiotic counterparts, suggesting that the symbiotic

state confers a growth advantage under nutrient-limited conditions. Furthermore, we anticipated

that starved, freshly inoculated G1 juveniles would demonstrate an intermediate growth rate:

slower than symbiotic animals but faster than aposymbiotic animals. Lastly, we hypothesized

that heterotrophic feeding would have an additive effect on growth rate, where those that are fed

would grow faster regardless of symbiont presence. Surprisingly, our results strikingly refute

some of our hypotheses and bring up interesting new questions about the interacting effects of

immunity, host age and nutrition on symbiosis health and regulation.

2. Methods

a) Anemone and algal stocks

All sea anemone and algal cultures were kept in incubators (Percival Model: CU-36L5) at

$25°$ C under 20–30 µmol quanta m^{-2} s^{-1} using 5,000K fluorescent lights (Zoo Med Flora Sun) on

a 12/12h light/dark photoperiod. The algal cultures were maintained in 50 mL flasks using

autoclaved F/2 media (Bigelow NCMA, East Boothbay, ME, USA) without silicates. New

cultures were created once every two months by inoculating new F/2 media flasks with 5 mL of

the previous culture. Sea anemones for this study came from the Aiptasia clonal line H2,

originating from Hawaii [24]. To generate aposymbiotic (without algae) anemones, animals were

artificially bleached with menthol [25] once every three months. Aposymbiotic anemones used

in this study were menthol-bleached two months before the experiment began to minimize

effects from menthol on our results. Symbiotic parent anemones (G0s) were kept in clear 300 mL

tubs containing 1 µm filtered seawater (FSW) while aposymbiotic animals were kept under the

same conditions but in opaque containers to discourage re-inoculation with algae. G0 anemones

were fed *ad libitum* three times weekly with freshly-hatched *Artemia* nauplii and 100% water

changes with FSW were completed within 24 hours after feeding.

To verify the identity of algal species, algae from both G0 animals and algae in culture

were sequenced using Internal Transcribed Spacer 2 (ITS2). DNA was extracted from whole

anemones and algae pellets using the Qiagen DNeasy Plant Pro Kit. PCR amplification of the

target region was performed with ITS2-specific primers (primer sequence). We analyzed the

resulting PCR products through gel electrophoresis, followed by Sanger sequencing at the Center

for Quantitative Life Sciences, Oregon State University. Sequences were cleaned and

Symbiodiniaceae algae types identified using Geneious software and the SymPortal BLAST

database [26] to identify Symbiodiniaceae algae types. Algae from anemones and culture

matched type B1, *Breviolum minutum*, the native symbiont to the H2 clonal line [27]. The algae

within symbiotic parental sea anemones are assumed to be from the original clonal source while

the culture was strain SSB01, an axenic culture sourced from Aiptasia [24].

b) Experimental design and husbandry

We prepared artificially-produced pedal lacerates (G1s) following Presnell et al. [23]

with some modifications. We placed symbiotic and aposymbiotic G0 adults in separate

polypropylene petri dishes for two days so that they would fully attach. Once attached, we

excised two pieces of tissue from the pedal disk of each anemone using a scalpel [23], for a total

of 144 lacerates. The size of these artificially-produced lacerates was measured and the average

size was not different from naturally-produced lacerates (data not shown). Each lacerate was

placed in a well of a 12-well plate with 750 μL filtered seawater (FSW). The lacerates were left

to develop naturally, not removing the mucous film that develops on the tissue ball as described

in Presnell et al. [23]. We found no evidence for higher survival in lacerates that received this

treatment (see Supp. Fig. 1).

We produced symbiotic state treatments: aposymbiotic, inoculated and symbiotic as

illustrated in Figure 2. Symbiotic G1s were cut from symbiotic G0s, while inoculated and

aposymbiotic pedal lacerates G1s were cut from aposymbiotic G0s. The inoculated treatment

was colonized with algae (SSB01, *B. minutum*) in FSW at a final density of 3×10^5 cells/mL

within each well and left for 24 hours before a 100% water change. We inoculated using this

method twice to ensure a successful inoculation, at four and then seven days after laceration. By

day four, all anemones had developed a mouth, but only some had one tentacle (data not shown).

By day seven, most anemones had six or more tentacles, following the pattern reported by

Presnell et al [23]. On day eight, we removed anemones from wells, the wells were cleaned with

a cotton swab, and then the animals returned to the wells with fresh FSW. This removed any

residual symbionts and mucus left in the wells from the first seven days. All inoculated

anemones were confirmed to have algae by day 10 (data not shown).

Symbiotic state treatments were crossed with two feeding treatments, starved and fed, for

a total of 24 G1s in each treatment group at day zero. The starved treatments received no food

and fed treatments were given 4-5 live rotifers (source culture from K&P Aquaculture) three

times weekly beginning seven days after laceration. FSW changes were completed three times

weekly, 24 hours after feeding. We quantified tentacle number five to seven times weekly

beginning on day four. Tentacles begin as thick nubs and extend. We counted tentacles only if

they were longer than they were wide. The rotifers fed to anemones were kept at 26°C in 1L

bottles that received water changes once weekly. We sieved the rotifer culture with a 50 μm

nylon mesh, placed the sieved rotifers back in the bottle, and replaced the water with 900 mL

FSW and 100 mL *Tetraselmis* sp. microalgae once weekly. Live *Tetraselmis* sp. cultures were

maintained in 1L bottles with F/2 media in FSW refreshed weekly under 200 µmol quanta m^{-2} s^{-1}

LED lights (Barrina T5 Grow Lights, pinkish white).

c) Symbiont density and size

We imaged G1 anemones on a Zeiss Axio Observer.A1 inverted epifluorescence

microscope under brightfield illumination for pedal disk area and at 700 nm excitation for

symbiont imaging. All images were analyzed in ImageJ [28]. Pedal disk area was used as a non-

lethal measure of anemone size [29]. We measured symbiont density from images of tentacles by

selecting a region in the middle of each tentacle 2-8 mm in length and divided the proportion of

algae cell pixels (fluorescing) by the total number of pixels in the selected region, similar to

Maruyama et. al. [30]. We took the symbiont density of two tentacles per anemone. This allowed

for multiple non-destructive measurements from the same individuals.

d) Cell division

EdU (5-ethynyl 2´-deoxyuridine), a thymidine analog, gives a snapshot of host anemone

cell division rates *in vivo* (proportion of cells in S-phase) within an incubation period. We

sacrificed four anemones from each treatment for EdU labeling at three time points, after the

second, fourth, and tenth week. By the tenth week, there were no starved symbiotic or starved

inoculated G1s to sample. G1 anemones were incubated in 10 µM EdU in their wells for 24

hours before fixing in 4% formaldehyde, 0.2% glutaraldehyde, 0.1% Tween in PBS for 1.5

minutes then 4% formaldehyde and 0.1% Tween in PBS overnight. Each polyp was treated as

described in Tivey et al. [9] to label EdU+ cells (Click-iT EdU Alexa Fluor 555 imaging kit; Life

Technologies, Eugene, OR, USA) and host nuclei with Hoechst 33342. Briefly, tissue was

permeabilized in 0.2% PBST (0.2% Triton in PBS), washed in 0.02% PBST, blocked in 5% BSA

in PBS, labeled with EdU reaction, rinsed with 0.02% PBST, labeled with Hoechst 33342,

washed in 0.02% PBST, and mounted on a slide suspended in a final concentration of 87%

glycerol. Slides were imaged on a Zeiss LSM 780 NLO confocal microscope with three

excitation wavelengths: 555 nm (EdU-AF555), 405 nm (Hoechst), and 633 nm for algal

symbiont autofluorescence. Images were taken at multiple depths and combined into z-stacks for

better resolution across three dimensional structures. These z-stack images were analyzed in FIJI

[28] by selecting a region in the middle of a tentacle 2-8 mm in length and measuring the number

of pixels of EdU fluorescence standardized by the number of pixels of Hoechst fluorescence to

calculate a host cell division rate.

e) Statistical analyses

All analyses were completed in R v. 4.1.2 [31]. We used the package lme4 to create

models [32], car for ANOVA [33], and emmeans for posthoc analyses [34]. All models followed

the same design, which included a three-way interaction among symbiotic state treatment,

feeding treatment, and time, lm(data ~ symbiont * feed * day). We analyzed tentacle count data

with a generalized linear mixed model with a log link Poisson distribution. We analyzed

symbiont density, host size, and host cell division rate with linear models and checked

differences between observed and predicted values (residuals) for normality before analysis.

3. Results

a) Survival of starved anemones was reduced by the presence of symbionts

All anemones survived the initial development period before day 30, except for three in

the symbiotic treatments (Fig. 3). By day 30, the starved symbiotic anemones group had lost four

individuals while the fed symbiotic anemones group had lost one out of 24. After day 30, the

starved symbiotic and inoculated anemones rapidly died until there was only one G1 left in the

inoculated treatment. The rate of loss between starved symbiotic and inoculated anemones was

similar, but the inoculated group lagged approximately one week behind the symbiotic group.

Only two anemones died in the starved aposymbiotic treatment group by the end of the

experiment, and no fed anemones died after day one.

b) Development was affected by an interaction between feeding and the presence of symbionts

229   There were three distinct periods of development apparent in the data: Period 1, (days 0-

9), 2 (days 10-28), and 3 (days 29-70; Fig. 4). The analysis is divided into these three periods.

During Period 1, symbiotic state had an effect on host development (symbiont: χ^2 = 77.38, p <

0.001; symbiont*day: χ^2 = 78.08, p < 0.001; Fig. 4) similar to what was found during the first

few days of development by Presnell et al. [23]. Symbiotic G1s developed more slowly than all

other anemones in other treatments, including G1s that were inoculated with algae on days four

and seven. Symbiotic G1s had significantly fewer tentacles than all other groups on days 5-7

(Supp. Table 1). Feeding had a small effect on development during Period 1 where fed animals

grew tentacles at a faster rate (χ^2 = 4.93, p = 0.026). However, symbiotic G1s caught up to

animals in the rest of the treatments by day nine, the beginning of Period 2. There were no

significant differences between treatment groups on day nine.

240   During Period 2, there was a strong interaction between feeding and symbiotic state (χ^2

= 10.44, p = 0.005) where fed anemones developed faster than starved ones and the presence of

symbionts increased growth rates only when combined with feeding. (Fig. 4). Fed symbiotic

anemones, overtook animals in the rest of the treatments with an average of 16.3±0.6

(mean±SEM) tentacles by day 28. All starved treatments performed similarly, with G1s in the

starved symbiotic treatment having an average of 8.7±1.5 tentacles by the end of the period.

Tentacle numbers of fed symbiotic and fed inoculated animals were not significantly different (z

= -2.02, p = 0.329) and fell between these two groups at an average of 13.8±0.3 tentacles in the

fed inoculated treatment by day 28.

During Period 3, there was a strong interaction between feeding and symbiotic state

treatments ($\chi 2$ = 948.0, p < 0.001) where the presence of symbionts, both in the symbiotic and

inoculated treatments, had a positive effect on development if food was present but, crucially, a

deleterious effect when the host anemone was starved (Fig. 4). G1s in the fed inoculated

treatment caught up to those in the fed symbiotic treatment by the end of Period 3 (z = 0.03, p =

1.0). G1s in the fed aposymbiotic treatment had fewer tentacles than the fed symbiotic and

inoculated G1s throughout this period (Fig. 4), but no pairwise comparisons were significant

(Supp. Table 1). The starved aposymbiotic G1s had fewer tentacles than the fed aposymbiotic

G1s (day 70: z = 0.12, p < 0.001), but more tentacles than starved symbiotic and inoculated G1s

(day 64 starved symbiotic: z = 6.58, p < 0.001; day 64 starved inoculated: z = 7.76, p < 0.001).

The starved anemones that had algal symbionts notably lost tentacles during Period 3

(Fig. 4). G1s that were starved and had symbionts began losing tentacles on day 28 (Fig. 4). The

starved symbiotic G1s lost tentacles earlier than the starved inoculated G1s, but the starved

inoculated anemones caught up to the loss three weeks later (Fig. 4). Most starved anemones that

had symbionts lost all of their tentacles, eventually forming balls of tissue that were dead by day

70.

c) Symbiont density was higher in starved anemones

There was an effect of feeding (F = 56.34, p < 0.001), symbiotic state (F = 122.55, p <

0.001), and an interaction between feeding and symbiotic state (F = 18.38, p < 0.001) on

symbiont density in host tissues. The inoculated G1s began with no symbionts at day zero while

the natural-state symbiotic G1s started with hundreds of symbionts from G0s (*pers. obs.*,

X.X.X.). Symbiont density was first measured at day 10 when well-developed tentacles were

present. By this time, the starved symbiotic G1s had a higher symbiont density than the fed

symbiotic G1s (t = -6.31, p < 0.001), and this pattern persisted throughout the experiment (Fig.

5). The symbiont density within the inoculated anemones was lower than the symbiotic

anemones at day 10 (Fig. 5), and there was no difference between fed inoculated and starved

inoculated on day 10 (t = -0.05, p = 0.999) or day 25 (t = -1.70, p = 0.329). However, by day 25

the inoculated G1s had a similar density to the symbiotic G1s (Fig. 5). On day 66 there was only

one anemone left in the fed symbiotic group, but this anemone had 100% symbiont fluorescence

per tentacle area (Fig. 5). The starved inoculated G1s had a higher symbiont density than the G1s

in either of the fed treatments (fed symbiotic t = 6.27, p < 0.001; fed inoculated t = -6.55, p <

0.001), and there was no difference between the fed symbiotic and fed inoculated anemones (t =

-0.37, p = 0.982), which had the lowest density (Fig. 5). No algae was detected in the

aposymbiotic anemones throughout the experiment.

d) Growth was increased by symbionts only in the fed anemones

Size of anemones was affected by symbiotic state (F = 68.19, p < 0.001), feeding (F =

123.91, p < 0.001), and a three-way interaction between symbiotic state, feeding, and time (F =

7.15, p < 0.001). We visually size matched lacerates while cutting them for our treatments, but

by day two the symbiotic lacerates were significantly larger than all other treatments (Fig. 6;

Supp. Table 3). On days 10 and 24, the inoculated G1s had similar sizes to their respective

aposymbiotic treatments (Fig. 6; Supp. Table 3). On day 66 there was a clear interaction between

symbiotic state and feeding. There were no differences between any of the starved anemone

groups (Supp. Table 3) which lost 86.65±0.28% of their initial size, but there were large

differences among the three fed anemone groups. The fed symbiotic anemones were larger than

the fed inoculated anemones (t = -3.43, p = 0.009), and the fed inoculated anemones were larger

than the fed aposymbiotic anemones (t = -5.70, p < 0.001).

e) Host cell division was higher in fed anemones

Host cell division rate, as measured by EdU fluorescence was affected by feeding (F =

10.70, p = 0.002) but unaffected by symbiotic state (F = 18.46, p < 0.001). There were some

pairwise differences between symbiotic states on day 15 where fed symbiotic animals had higher

cell division than fed inoculated (t = -3.20, p = 0.030) and fed aposymbiotic anemones (t = -

3.22, p = 0.028), but the effect of symbiotic state in the fed anemones was gone by the next time

point (aposymbiotic vs. symbiotic: t = -0.722, p = 0.978). Fed anemones had higher cell division

than starved aposymbiotic anemones on days 29 and 70 (Supp. Fig. 2, Supp. Table 4).

4. Discussion

Our study demonstrated an interaction between symbiotic state and feeding on survival,

development, and growth of G1 (asexually-produced) offspring in a model cnidarian-algal

symbiosis. There was a strong interaction between symbiotic state and feeding where symbionts

were beneficial to the developing host anemone only when the host was fed. Fed anemones with

algae developed faster and grew larger than those without algae, and this effect was symbiont

dose-dependent. G1 anemones that started with symbionts grew faster than those that were

inoculated later and only started with a few cells that had to proliferate throughout the host. This

supported our initial hypothesis that the presence of symbionts would increase the growth rate of

G1 offspring.

The presence of symbionts in the starved treatments, however, did not increase the

growth rate of anemones, and, instead, slowed and eventually reversed tentacle growth after the

first three weeks, possibly after the offspring exhausted the resources from G0 adults. Starved

anemones shrank regardless of symbiotic state, but those with algae were more likely to lose all

tentacles and die. The effect was more pronounced in anemones with higher initial symbiont

densities: they died faster. This contradicted our hypothesis that the presence of symbionts would

boost development and growth. Instead, high symbiont densities in starved G1 anemones likely

exacerbated the adverse effects of starvation leading to their demise, as visualized by images of

EdU labeled dividing cells in fed compared to starved animals (Fig. 7).

Our findings expand on previous studies on sea anemones and corals. The symbiosis-by-

nutrition interaction has recently been a subject of studies performed in both adult corals [35]

and Aiptasia [36]. When corals were deprived of nutrients, for many months, they were shown to

'farm' their algae: digest a portion of the algal population to help them survive low nutrient

conditions. When anemones were starved for one year, both host biomass and symbiont densities

in host tissues decreased dramatically. However crucially, the symbionts continued to pass

carbon to the host: symbiosis homeostasis was maintained, the animals remained symbiotic and

the holobiont remained healthy. These results are in marked contrast to those found in our G1

juveniles where starvation caused profound symbiosis dysregulation and led to death.

Another study on Aiptasia by Peng et al. [36] found that aposymbiotic anemones

inoculated with algae and then starved for months demonstrated a fatal metabolic cost from

algae, leading to high mortality rates in the inoculated anemones as compared to the

aposymbiotic controls. Our results focusing on G1 offspring from pedal laceration, suggest that

developmental stage plays a crucial role in the regulation of homeostasis of the symbiosis in the

absence of heterotrophic feeding. The key differences between our study and Peng et. al. [36] is

that we included symbiotic G1s from the start while Peng's study included only aposymbiotic

and inoculated G1s, and we followed G1 anemones from initial laceration to adults while Peng's

study started with juveniles (6 tentacles, Stage 3 juveniles [23]). Our results also align with a

modeling study that used data from the Peng et al. [36] study to develop a food-dependent model

in Aiptasia [37]. The model developed in the study "demonstrates the role of the symbiont as an

amplifier of the host's state: a growing anemone grows better with symbionts, while a

malnourished anemone loses biomass faster with a symbiont than without." Our results

recapitulate and expand on these studies, demonstrating a stronger metabolic cost of algal

symbionts when algae are transmitted vertically at higher densities and at earlier developmental

stages. This indicates that earlier developmental stages may be less capable of regulating

symbiont populations under nutrient-limited conditions than adults, leading to symbiosis

dysregulation and host mortality.

Taken together, these studies suggest that host-symbiont biomass regulation and

homeostatic regulation of symbiosis differs between adult and developing animals. What

processes could explain this pattern? We suggest that the differences may be influenced by host

immunity. Asexual reproduction and regeneration in early-diverging metazoans are

evolutionarily tightly linked. Indeed pedal laceration can be considered a form of whole-body

regeneration (WBR), an area of evolutionary developmental biology that has been of interest to

biomedicine for many years [38]. In Aiptasia, regeneration, including pedal laceration, can

trigger immune responses that are linked with developmental pathways [39]. Studies on WBR in

a variety of animals including cnidarians suggest an inverse relationship between

regenerative/developmental processes and immune activation [40,41]. Transcriptomic studies on

Aiptasia regeneration have shown that innate immunity pathways are initially upregulated

following wounding but are downregulated as developmental programs take over [39]. This

could explain our results in G1 juveniles. After laceration, immunity is repressed to facilitate

WBR. However, when heterotrophic feeding is withheld and with immunity downregulated, the

symbiosis becomes dysregulated, leading to unchecked parasitic algal proliferation within the

host tissue and a fatal cost to the developing host. In contrast, when food is present,

dysregulation is avoided and normal development proceeds.

Our results also align with recent findings on the relationship between symbiont density

and host immunity in cnidarians. Studies on *Orbicella faveolata* and *Astrangia poculata* have

demonstrated that high symbiont densities in hosts can negatively impact host immunity,

particularly under stressful conditions such as nutrient limitation [42,43]. Increased symbiont

density in *O. faveolata* was associated with a significant suppression of immune gene expression,

potentially making the host more vulnerable to disease [42]. Similarly, in *A. poculata*, a positive

correlation between symbiont density and immune response was observed, but the immune

response varied with symbiont density, suggesting that under nutrient stress, higher symbiont

loads lead to an energetic burden that exacerbates host stress [43]. These findings suggest that

nutrition and symbiont density are critical factors that determine the balance between symbiosis

and immunity, and their dysregulation can lead to adverse outcomes, such as those observed in

our starved and symbiotic G1 anemones.

To further our understanding of the interplay between nutrition, development, and

symbionts, future studies could focus on algal cell cycle regulation during development using

methodologies similar to those employed by Tivey et. al. [9] and transcriptomic comparisons

across different feeding conditions, developmental stages, and symbiotic states. Examining

sexually-produced juvenile polyps, F1 offspring, could provide insights into whether the

outcomes observed in G1 offspring are consistent across different reproductive modes.

Maintaining homeostasis between partners in coral-algal symbiosis during development,

regeneration, and growth is essential for healthy coral reefs. A focus on nutrition during early

development and/or regeneration could be critical for successful conservation efforts.

Figures

Figure 1. Reproductive modes of Aiptasia showing both the F1 (sexual reproduction) and G1

generation (asexual reproduction). Brown indicates the presence of algal symbionts and gray

indicates no algal symbionts. Developmental stages are from Presnell et al. 2022. Created with

BioRender.

Figure 2. Illustration of the three algal symbiont experimental treatments. Brown indicates the

presence of algal symbionts and gray indicates no algal symbionts. Developmental stages are

from Presnell et al. 2022. Created with BioRender.

Figure 3. Proportion of pedal lacerates surviving over time (days post-laceration). Each

treatment began with 24 individuals. Vertical dashed lines separate Periods in the experiment.

Starved symbiotic offspring exhibited mortality starting in Period 2, while starved inoculated

anemones began to die at the beginning of Period 3. By day 70, only one starved anemone with

symbionts remained alive, with all others having died. Survival was highest in fed aposymbiotic

and inoculated anemones.

Figure 4. Pedal lacerates tentacle count over time (days post-laceration). Points are jittered to

differentiate error bars. Points are means ± standard error of the mean. Vertical dashed lines

separate Periods in the experiment. Sample sizes for each treatment at each time point were

reduced throughout the experiment because of lethal sampling for EdU labeling where days 0-15

had N = 24, days 16-29 had N = 20, and days 30-70 had N = 16. During Period 1, symbiotic

anemones exhibited a slower developmental rate compared to other groups. In Period 2, the

influence of food availability became apparent, as fed anemones showed faster development. By

Period 3, there was a clear interaction between symbiont presence and feeding; the presence of

symbionts was only beneficial to development when food was available.

Figure 5. Symbiont density of pedal lacerates over time (days post-laceration). The symbiont

density of symbiotic lacerates at time point zero was not measured. Points are means ± standard

error of the mean. Vertical dashed lines separate Periods in the experiment. Sample sizes for

each treatment at each time point were reduced throughout the experiment because of lethal

sampling for EdU labeling where days 0-15 had N = 24, days 16-29 had N = 20, and days 30-70

had N = 16. Symbiont density began higher in symbiotic anemones compared to inoculated

anemones. Throughout the experiment, fed anemones with symbionts maintained a stable

symbiont density, while starved anemones showed a notable increase in symbiont density,

reaching more than double the density observed in the fed treatment.

Figure 6. Size of pedal lacerates over time (days post laceration). Points are means ± standard

error of the mean. Vertical dashed lines separate Periods in the experiment. Sample sizes for

each treatment at each time point were reduced throughout the experiment because of lethal

sampling for EdU labeling where days 0-15 had N = 24, days 16-29 had N = 20, and days 30-70

had N = 16. Symbiotic lacerates were larger during Period 1, but those groups that were fed grew

larger during Periods 2 and 3. The presence of symbionts increased size only in the fed groups.

Figure 7. Representative z-stack images of anemones from day 29 (beginning of Period 3) of the

experiment incubated in EdU for 24 hours before fixation. Fluorescent channels mark host nuclei

in blue (Hoechst), host cell division in green (EdU), and algal symbionts in red (chlorophyll

autofluorescence).

References

1. Davy SK, Allemand D, Weis VM. 2012 Cell Biology of cnidarian-dinoflagellate symbiosis.

Microbiol. Mol. Biol. Rev. **76**, 229–261. (doi:10.1128/MMBR.05014-11)

2. Yellowlees D, Rees TAV, Leggat W. 2008 Metabolic interactions between algal symbionts

and invertebrate hosts. *Plant, Cell & Environment* **31**, 679–694. (doi:10.1111/j.1365-

3040.2008.01802.x)

3. Baker AC. 2003 Flexibility and specificity in coral-algal symbiosis: diversity, ecology, and

biogeography of Symbiodinium. *Annual Review of Ecology, Evolution, and Systematics* **34**,

661–689. (doi:10.1146/annurev.ecolsys.34.011802.132417)

4. McIlroy SE, terHorst CP, Teece M, Coffroth MA. 2022 Nutrient dynamics in coral

symbiosis depend on both the relative and absolute abundance of Symbiodiniaceae species.

Microbiome **10**, 192. (doi:10.1186/s40168-022-01382-0)

5. Cui G, Liew YJ, Li Y, Kharbatia N, Zahran NI, Emwas A-H, Eguiluz VM, Aranda M. 2019

Host-dependent nitrogen recycling as a mechanism of symbiont control in Aiptasia. *PLoS*

Genet. **15**, e1008189. (doi:10.1371/journal.pgen.1008189)

6. Rädecker N, Escrig S, Spangenberg JE, Voolstra CR, Meibom A. 2023 Coupled carbon and

nitrogen cycling regulates the cnidarian–algal symbiosis. *Nat. Commun.* **14**, 1–10.

(doi:10.1038/s41467-023-42579-7)

7. Mansfield KM, Gilmore TD. 2019 Innate immunity and cnidarian-Symbiodiniaceae

mutualism. *Developmental & Comparative Immunology* **90**, 199–209.

(doi:10.1016/j.dci.2018.09.020)

8. Matthews JL, Crowder CM, Oakley CA, Lutz A, Roessner U, Meyer E, Grossman AR, Weis

VM, Davy SK. 2017 Optimal nutrient exchange and immune responses operate in partner

specificity in the cnidarian-dinoflagellate symbiosis. *PNAS* **114**, 13194–13199.

(doi:10.1073/pnas.1710733114)

9. Tivey TR, Parkinson JE, Weis VM. 2020 Host and symbiont cell cycle coordination is

mediated by symbiotic state, nutrition, and partner identity in a model cnidarian-

dinoflagellate symbiosis. *mBio* **11**. (doi:10.1128/mBio.02626-19)

10. Downs CA *et al.* 2013 Heat-stress and light-stress induce different cellular pathologies in the

symbiotic dinoflagellate during coral bleaching. *PLOS ONE* **8**, e77173.

(doi:10.1371/journal.pone.0077173)

11. Rowan R, Knowlton N, Baker A, Jara J. 1997 Landscape ecology of algal symbionts creates

variation in episodes of coral bleaching. *Nature* **388**, 265–269. (doi:10.1038/40843)

12. Gong S, Li G, Liang J, Xu L, Tan Y, Jin X, Xia X, Yu K. 2023 Day-night cycle as a key

environmental factor affecting coral-Symbiodiniaceae symbiosis. *Ecological Indicators* **146**,

109890. (doi:10.1016/j.ecolind.2023.109890)

13. Helgoe J, Davy SK, Weis VM, Rodriguez-Lanetty M. 2024 Triggers, cascades, and

endpoints: connecting the dots of coral bleaching mechanisms. *Biological Reviews* **99**, 715–

752. (doi:10.1111/brv.13042)

14. Eddy TD, Lam VWY, Reygondeau G, Cisneros-Montemayor AM, Greer K, Palomares

MLD, Bruno JF, Ota Y, Cheung WWL. 2021 Global decline in capacity of coral reefs to

provide ecosystem services. *One Earth* **4**, 1278–1285. (doi:10.1016/j.oneear.2021.08.016)

15. Bocharova ES, Kozevich IA. 2011 Modes of reproduction in sea anemones (Cnidaria,

Anthozoa). *Biol Bull Russ Acad Sci* **38**, 849–860. (doi:10.1134/S1062359011090020)

16. Reitzel A, Stefanik D, Finnerty J. 2011 Asexual reproduction in Cnidaria: Comparative

processes and candidate mechanisms. pp. 101–113.

17. Caruso C *et al.* 2022 Genetic patterns in *Montipora capitata* across an environmental mosaic

in Kāne'ohe Bay, O'ahu, Hawai'i. *Molecular Ecology* **31**, 5201–5213.

(doi:10.1111/mec.16655)

18. Roberty S, Weis VM, Davy SK, Voolstra CR. 2024 Editorial: Aiptasia: a model system in

coral symbiosis research. *Front. Mar. Sci.* **11**. (doi:10.3389/fmars.2024.1370814)

19. Grawunder D, Hambleton EA, Bucher M, Wolfowicz I, Bechtoldt N, Guse A. 2015

Induction of gametogenesis in the cnidarian endosymbiosis model Aiptasia sp. *Scientific*

Reports **5**, 15677. (doi:10.1038/srep15677)

20. Clayton WS, Lasker HR. 1985 Individual and population growth in the asexually

reproducing anemone *Aiptasia pallida* Verrill. *Journal of Experimental Marine Biology and*

Ecology **90**, 249–258. (doi:10.1016/0022-0981(85)90170-4)

21. Maegele I, Rupp S, Özbek S, Guse A, Hambleton EA, Holstein TW. 2023 A predatory

gastrula leads to symbiosis-independent settlement in Aiptasia. *Proceedings of the National*

Academy of Sciences **120**, e2311872120. (doi:10.1073/pnas.2311872120)

22. Clayton W. 1985 Pedal laceration by the anemone *Aiptasia pallida*. *Mar. Ecol. Prog. Ser.*

21, 75–80. (doi:10.3354/meps021075)

23. Presnell JS, Wirsching E, Weis VM. 2022 Tentacle patterning during *Exaiptasia diaphana*

pedal lacerate development differs between symbiotic and aposymbiotic animals. *PeerJ* **10**,

e12770. (doi:10.7717/peerj.12770)

24. Xiang T, Hambleton EA, DeNofrio JC, Pringle JR, Grossman AR. 2013 Isolation of clonal

axenic strains of the symbiotic dinoflagellate Symbiodinium and their growth and host

specificity1. *Journal of Phycology* **49**, 447–458. (doi:10.1111/jpy.12055)

25. Matthews JL, Sproles AE, Oakley CA, Grossman AR, Weis VM, Davy SK. 2016 Menthol-

induced bleaching rapidly and effectively provides experimental aposymbiotic sea anemones

(Aiptasia sp.) for symbiosis investigations. *Journal of Experimental Biology* **219**, 306–310.

(doi:10.1242/jeb.128934)

26. Hume BCC, Smith EG, Ziegler M, Warrington HJM, Burt JA, LaJeunesse TC, Wiedenmann

J, Voolstra CR. 2019 SymPortal: A novel analytical framework and platform for coral algal

symbiont next-generation sequencing ITS2 profiling. *Molecular Ecology Resources* **19**,

1063–1080. (doi:https://doi.org/10.1111/1755-0998.13004)

27. Thornhill DJ, Xiang Y, Pettay DT, Zhong M, Santos SR. 2013 Population genetic data of a

model symbiotic cnidarian system reveal remarkable symbiotic specificity and vectored

introductions across ocean basins. *Molecular Ecology* **22**, 4499–4515.

(doi:10.1111/mec.12416)

28. Schindelin J *et al.* 2012 Fiji: an open-source platform for biological-image analysis. *Nature*

Methods **9**, 676–682. (doi:10.1038/nmeth.2019)

29. Bedgood SA, Bracken MES, Ryan WH, Levell ST, Wulff J. 2020 Nutritional drivers of adult

locomotion and asexual reproduction in a symbiont-hosting sea anemone Exaiptasia

diaphana. *Mar Biol* **167**, 39. (doi:10.1007/s00227-020-3649-3)

30. Maruyama S, Lab OW, Weis V. 2021 Symbiont density quantification in live Aiptasia.

31. R Core Team. 2021 *R: A Language and Environment for Statistical Computing*. Vienna,

Austria: R foundation for statistical computing. See https://www.R-project.org/.

32. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting linear mixed-effects models using

lme4. *Journal of Statistical Software* **67**, 1–48. (doi:10.18637/jss.v067.i01)

33. Fox J, Weisberg S. 2019 *An R Companion to Applied Regression*. Third. Thousand Oaks

CA: Sage. See https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

34. Lenth RV. 2021 *emmeans: Estimated Marginal Means, aka Least-Squares Means*. See

https://CRAN.R-project.org/package=emmeans.

35. Wiedenmann J *et al.* 2023 Reef-building corals farm and feed on their photosynthetic

symbionts. *Nature* , 1–7. (doi:10.1038/s41586-023-06442-5)

36. Peng S-E, Moret A, Chang C, Mayfield AB, Ren Y-T, Chen W-NU, Giordano M, Chen C-S.

2020 A shift away from mutualism under food-deprived conditions in an anemone-

dinoflagellate association. *PeerJ* **8**, e9745. (doi:10.7717/peerj.9745)

37. Kaare-Rasmussen JO, Moeller HV, Pfab F. 2023 Modeling food dependent symbiosis in

Exaiptasia pallida. *Ecological Modelling* **481**, 110325.

(doi:10.1016/j.ecolmodel.2023.110325)

38. Bideau L, Kerner P, Hui J, Vervoort M, Gazave E. 2021 Animal regeneration in the era of

transcriptomics. *Cell. Mol. Life Sci.* **78**, 3941–3956. (doi:10.1007/s00018-021-03760-7)

39. van der Burg CA, Pavasovic A, Gilding EK, Pelzer ES, Surm JM, Smith HL, Walsh TP,

Prentis PJ. 2020 The Rapid regenerative response of a model sea anemone species

Exaiptasia pallida is characterised by tissue plasticity and highly coordinated cell

communication. *Mar. Biotechnol.* **22**, 285–307. (doi:10.1007/s10126-020-09951-w)

40. Eming SA, Hammerschmidt M, Krieg T, Roers A. 2009 Interrelation of immunity and tissue

repair or regeneration. *Semin. Cell Dev. Biol.* **20**, 517–527.

(doi:10.1016/j.semcdb.2009.04.009)

41. van der Burg CA, Prentis PJ. 2021 The Tentacular spectacular: evolution of regeneration in

sea anemones. *Genes* **12**, 1072. (doi:10.3390/genes12071072)

42. Fuess LE, Palacio-Castro AM, Butler CC, Baker AC, Mydlarz LD. 2020 Increased algal

symbiont density reduces host immunity in a threatened caribbean coral species, Orbicella

faveolata. *Front. Ecol. Evol.* **8**. (doi:10.3389/fevo.2020.572942)

43. Changsut IV, Borbee EM, Womack HR, Shickle A, Sharp KH, Fuess LE. 2024

Photosymbiont density is correlated with constitutive and induced immunity in the

facultatively symbiotic coral, *Astrangia poculata. Integrative and Comparative Biology* ,

icae036. (doi:10.1093/icb/icae036)

563