A nucleic acid prodrug activates mitochondrial respiration and extends lifespan

Takahisa Anada*, Michiharu Kawahara, Taisei Shimada, Ryotaro Kuroda, Eriko Kage-Nakadai, Shingo Kobayashi, Masaru Tanaka*

T. Anada, S. Kobayashi, M. Tanaka

Institute for Materials Chemistry and Engineering, Kyushu university, 744 Motooka, Nishi-ku Fukuoka 819-0395, Japan

E-mail: takahisa_anada@ms.ifoc.kyushu-u.ac.jp, masaru_tanaka@ms.ifoc.kyushu-u.ac.jp

T. Anada, M. Kawahara, T. Shimada, R. Kuroda, M. Tanaka

Department of Applied Chemistry, Graduate School of Engineering, Kyushu University, 744 Motooka, Nishi-ku Fukuoka 819-0395, Japan

E. Kage-Nakadai

Laboratory of Environment and Health Science, Graduate School of Human Life and Ecology, Osaka Metropolitan University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan

E. Kage-Nakadai

Institute for Life and Medical Sciences, Kyoto University, 53 Shogoin Kawahara-cho, Sakyo-ku, Kyoto 606-8507, Japan

Keywords: nucleic acid prodrug, aging, mitochondria, ATP, lifespan

Abstract

Mitochondrial dysfunction caused by aging leads to decreased energy metabolism, resulting in functional decline and increased frailty in multiple tissues. Strategies for protecting and activating mitochondria under stressful conditions are required to suppress aging and age-related diseases. However, it is challenging to develop drugs capable of boosting mitochondrial respiration and compensating for the reduced intracellular adenosine triphosphate (ATP) levels. In this study, we developed a prodrug that stimulates the metabolism of intracellular adenine nucleotides (AXP: adenosine monophosphate (AMP), adenosine diphosphate (ADP), and ATP). It enhances AMP-activated protein kinase activity, fatty acid oxidation, oxidative stress resistance, and mitochondrial respiration, thereby increasing the intracellular ATP levels. Furthermore, this prodrug markedly extended the lifespan of *Caenorhabditis elegans*. The stimulation of the AXP energy metabolism (AXP stimulation) proposed in this study is expected to offer a new strategy against aging and pave the way for a novel approach in the bioenergetic molecules drug discovery.

1. Introduction

Aging and age-related diseases have been shown to be strongly associated with an imbalance between energy demand and supply[1]. Interventions such as calorie restriction, exercise, and molecules targeting longevity-related genes have been reported to be effective in addressing these issues[2]. Functions of mitochondria, the organelles responsible for energy production, decline with age, leading to age-

dependent reductions in intracellular adenosine triphosphate (ATP) levels. There are many reports that ATP levels decrease with age in human and various animals, such as in human fibroblasts[3], erythrocytes[4], skeletal muscles[5], rat heart tissues[6], mouse retinas and brains[7], *Drosophila*[8], and *Caenorhabditis elegans*[9] (*C. elegans*). ATP functions not only as an energy currency in biological systems, but also has diverse roles, including serving as a building block for ribonucleic acids, a donor in protein phosphorylation, and a solubilizing enhancer for maintaining proteostasis[10]. Therefore, its depletion might affect the maintenance of cellular functions. Mitochondrial dysfunction and decreased ATP levels impair energy metabolism and induce vulnerability in the entire body at the tissue level. Mitochondria play a crucial role not only in energy supply but also in the development of age-related diseases such as neurodegenerative and metabolic disorders[11–16]. Hence, there is a pressing need to develop techniques that protect against stress and activate the mitochondria. However, few drugs are capable of activating mitochondria and restoring reduced intracellular ATP levels, making the development of such mitochondria-activating drugs challenging. For this reason, this study hypothesized that by activating energy metabolism, particularly mitochondrial function, and increasing intracellular ATP concentration using pharmacological agents, it may be possible to slow the aging process.

ATP and its derivatives have been used as therapeutic agents to improve cardiovascular health, muscular performance, body composition, and fatigue recovery[17]. Despite the evidence of the therapeutic effects of ATP derivatives, their efficacy remains controversial. Some reports suggest that ATP administration is ineffective as a pharmaceutical agent[18] because of the limited blood stability[19] of adenosine derivatives and the lack of cell membrane permeability of the negatively charged phosphate groups, hindering entry into cells. Furthermore, cells recognize extracellular ATP as an inflammatory signal[20]. To minimize side effects, ATP administration techniques include drug delivery methods that encapsulate ATP in microparticles such as liposomes, micelles, and nanospheres[21]. Nevertheless, these approaches have limited utilization and applicability in the treatment of age-related diseases. Owing to the difficulty of increasing intracellular ATP concentrations, the long-term effects of ATP administration as a drug remain unclear.

Therefore, the present study was designed to develop the world's first prodrug with enhanced biological stability and cellular permeability of adenosine phosphate, capable of stimulating AXP (adenosine monophosphate (AMP), adenosine diphosphate (ADP), and ATP)-associated metabolic pathways, boosting mitochondrial respiration, ATP production, and reducing aging.

The strategy of the present study involved developing an AXP prodrug (proAX, with its chemical structure shown in Figure 1a and synthetic scheme in Figure S1) that can be converted to ATP via AMP

and ADP through intracellular enzymatic reactions and hydrolysis by phosphoramidating the phosphate moiety. Several phosphoramidate prodrugs (ProTides)[22] have been successfully applied to mononucleoside analogs as a class of nucleoside-based antivirals, such as Sofosbuvir and Remdesivir, targeting hepatitis C virus[23], and COVID-19[24]. Remdesivir is deemed safe for the treatment of hospitalized COVID-19 patients and approved by the Food and Drug Administration (FDA). These aryloxy phosphoramidate prodrugs are designed with the following proposed cellular activation pathway in the cell[25]: the prodrugs penetrate cell membranes, and the phosphoramidate moiety is metabolized stepwise by enzymatic and hydrolytic reactions to produce monophosphorylated nucleosides. These compounds are triphosphorylated by intracellular kinases and exert active medicinal effects. The potential intracellular activation mechanism of proAX developed in this study and based on this pathway is illustrated in Figure 1b. Moreover, we expected that, in addition to the direct conversion pathway of proAX to ATP, the transient increase in intracellular AMP levels might activate adenosine monophosphate-activated protein kinase (AMPK), which is involved in mitochondrial homeostasis regulation[26]. AMPK activation suppresses anabolic processes that consume ATP and promotes ATP synthesis through catabolic reactions.

Research on ProTide technology has primarily focused on its conversion into synthetic compounds that do not exist in nature. These compounds exhibited antiviral and anticancer properties. However, to the best of our knowledge, no instances of ProTide-type drugs exhibiting activity in a form that perfectly matches the structure of natural nucleic acids have been reported. To our knowledge, this is the first report of a prodrug that converts AXP into cells. In this study, we propose an innovative strategy for the "bioenergetic molecules drug discovery" to activate mitochondrial respiration and increase intracellular ATP levels, which decrease with aging or disease or both. We also demonstrated that proAX effectively extends the lifespan of *C. elegans*. We believe our findings open up new possibilities for nucleic acid prodrugs that achieve anti-aging through "stimulation of AXP-associated metabolic pathways (AXP stimulation)" distinct from existing anti-aging drugs.

Figure 1. The strategy of this study, and proAX enhances intracellular ATP levels without inducing cytotoxicity. a, The chemical structure of proAX. b, ProAX is internalized into a cell and undergoes cellular metabolic reactions. ProAX is metabolized by intracellular enzymes, such as esterases and phosphoamidases, and is then converted into AMP, ADP, and ATP. Transient increases in AMP levels could stimulate AMP-activated protein kinase (AMPK). The activation of AMPK could promote the ATP synthesis pathways through catabolic reactions. c, Comparison of intracellular ATP concentrations between HDF-29 and HDF-75, and the effects of varying proAX concentrations on intracellular ATP levels in HDF-75. d, Changes in intracellular ATP concentrations upon addition of adenosine and its derivatives (100 μM each) to HDF-75. e, Impacts of proAX addition on HDF-75 as determined by a LDH assay. For panels c and d, mean ± standard deviation shown, along with statistical significances for

relevant comparisons. Significance is assessed using a one-way ANOVA with Dunnett's multiple comparison test, with $n = 4$–19 replicates per condition. **$p < 0.01$, ****$p < 0.0001$; ns indicates not significant.

2. Results

2.1. Capability of ATP augmentation and cytotoxicity of the prodrug

The results of culturing normal human dermal fibroblasts from 29-year-old and 75-year-old donors (HDF-29 and HDF-75) are displayed in Figure 1c. In the absence of the prodrug, HDF-75 cells exhibited lower ATP levels than HDF-29 cells, a phenomenon previously reported in the literature [3]. We examined whether proAX increased the intracellular ATP levels in HDF-75 cells. After incubation with proAX for 24 h, the addition of the synthesized proAX to HDF-75 cells led to a concentration-dependent increase in intracellular ATP levels. With 100 µM proAX, a 2.5-fold increase in ATP levels was observed compared to the control (absence of proAX). We found that proAX effectively elevated the ATP levels in the cells of elderly individuals with reduced ATP levels. Similar results were obtained in HDF-29 cells (Figure S2a). These results demonstrate that the prodrug can increase intracellular ATP levels regardless of donor age. Addition of ATP analogs or proAX to compare intracellular ATP levels showed that adenosine, AMP, and ATP did not change the intracellular ATP concentration; however, addition of proAX led to a significant increase (Figure 1d). These findings confirm the effectiveness of the newly developed proAX in increasing intracellular ATP concentrations, unlike conventional ATP analogs, which are less effective because of their limited stability and cellular membrane permeability. We assessed the cytotoxicity of proAX in HDF-75 cells by live/dead staining. The results showed that proAX exhibited very low cytotoxicity (Figure S2b). Furthermore, the lactate dehydrogenase (LDH) assay indicated that proAX was not cytotoxic up to 256 µM in HDF-75 cells (Figure 1e), although its addition appeared to suppress cell proliferation (Figure S2c, d).

2.2. Evaluation of time-dependent changes of AMP, ADP, and ATP concentration in the cells

We analyzed changes in the intracellular concentrations of AMP, ADP, and ATP after proAX administration to HDF-75 cells using liquid chromatography mass spectrometry (LC/MS). AMP and ADP levels increased significantly within 3 h of proAX administration and continued to increase over 72 h (Figure 2a, b). After 3 h of prodrug administration, AMP and ADP levels in the cells were approximately 3-fold and 1.5-fold higher, respectively, than those observed without the prodrug. Conversely, an increase in ATP levels was observed later than that in AMP and ADP, starting after 6 h,

and remained significantly higher than that in the control condition without drug administration after 72 h (Figure 2c). These data indicated that when proAX was added at the start of the culture and incubated for three days without additional supplements, an increase in intracellular ATP levels was observed up to 72 h. Using these values, we calculated the adenylate energy charges in the cells with and without proAX administration at each time point. The energy charge was defined as (ATP +1/2ADP)/(ATP + ADP + AMP)[27]. Up to 24 h after drug administration, the energy charge in the prodrug group tended to be lower than that in the non-treated group; however, in both conditions, the value remained around 0.9, within maintaining a range under physiological conditions. Furthermore, we calculated changes in the AMP:ATP ratio induced by the addition of proAX. As illustrated in Figure 2e, proAX administration resulted in a 2.4-fold increase in the AMP: ATP ratio after 3 h. Over time, these ratios decreased. We found that proAX administration increased the adenine nucleotide pool (Figure 2f). These findings suggest that prodrug addition temporarily increased AMP, ADP, and ATP concentrations. Early after their addition, proAX enhanced the AMP:ATP ratio without deviating energy from the physiological norms. This indicates that proAX induces a pseudo-state of energy deficiency in the cells during the initial phase of addition.

Figure 2. Time-dependent changes in intracellular AXP levels and reactions in HDF-75 following the addition of proAX. a, Changes in AMP concentration. b, Changes in ADP concentration. c, Changes in ATP concentration. d, Changes in adenylate energy charge. e, Changes in AMP:ATP ratio. f, Changes in adenine nucleotide concentration (The amount of AMP is too low compared to ADP and ATP, making it barely visible on the graph). g, Time-dependent changes in phosphorylation of AMPK induced by proAX in HDF-75. Representative western blots with anti-P-AMPK (Thr172) and anti-AMPK antibodies, h, Time-dependent changes in phosphorylation of ACC induced by proAX in HDF-75. Representative Western blots with anti-P-ACC, and anti-ACC antibodies, i, Relative fatty acid oxidation (FAO) activity in HDF-75 treated with proAX for 3, 6, and 24 h. For all panels, HDF-75 were treated with 100 µM proAX, mean ± standard deviation shown, along with statistical significances between relevant comparisons. Significance is calculated using a one-way ANOVA with Sidak's

multiple comparisons test (a-d, f). Panel e uses an ANOVA followed by Dunnett's multiple comparisons test. ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, ns indicates not significant. For g, h, i panels, significance is calculated using a one-way ANOVA with Dunnett's multiple comparisons test (g, h), Sidak's multiple comparisons test (i). ****$p < 0.0001$, **$p < 0.01$.

2.3. AMPK activation by proAX

The AMPK monitors cellular energy status. AMPK activation is triggered by an increase in the cellular AMP: ATP ratio. Given the changes in these ratios, as indicated in Figure 2e, suggesting AMPK activation, we investigated the activation of AMPK by proAX and its time dependency in HDF-75 cells. Immunoblot analysis revealed that AMPK phosphorylation occurred within 3 h of treatment with 100 μM proAX (Figure 2g). AMPK activity increased approximately 4-fold 3 h after prodrug addition. AMPK activates fatty acid oxidation (FAO) by phosphorylating acetyl-CoA carboxylase (ACC), thereby reducing the levels of malonyl-CoA, an inhibitor of carnitine palmitoyltransferase (CPT). We assessed ACC phosphorylation after proAX administration (Figure 2h). ACC phosphorylation was observed within 3 h after the addition of proAX and was almost undetectable within 24 h. Following the confirmed phosphorylation of AMPK and ACC, we further explored fatty acid catabolism, specifically FAO, using a fatty acid β-oxidation-responsive fluorescent probe. The results indicated that at 3, 6, and 24 h post-proAX administration, FAO activity was significantly increased compared to that in the control group (no drug added) (Figure 2i).

2.4. Intracellular mechanisms of the ATP boosting

First, we explored the intracellular uptake pathway of proAX. Changes in the intracellular ATP levels following the addition of proAX and the nucleoside transporter inhibitor dipyridamole are shown in Figure 3a. The observation that higher concentrations of the inhibitor moderately reduced the increase in intracellular ATP levels induced by proAX suggests that proAX is taken up by cells via both passive diffusion and nucleoside transporters.

The increase in AMPK activity due to the addition of proAX was investigated to determine its effect on cellular ATP levels. This was evaluated by treating HDF-75 cells with dorsomorphin (Figure 3b), an AMPK inhibitor, and siRNA targeting AMPK (Figure S3a), followed by monitoring the changes in intracellular ATP concentration. ProAX-induced increase in cellular ATP levels was suppressed by dorsomorphin and siRNA. Furthermore, the phosphorylation of ACC by western blotting demonstrated that the treatment of proAX activated FAO processes. Collectively, these findings suggested that proAX

activates AMPK, which in turn stimulates FAO, leading to increased intracellular ATP concentrations. For comparison, the effect of 5-aminoimidazole-4-carboxamide-1-D-ribofuranoside (AICAR), an AMPK activator, on intracellular ATP levels was examined in HDF-75 cells. However, no increase in the ATP concentration was observed following AICAR treatment (Figure S3c). These results suggest that AMPK activity is necessary for the proAX-induced increase in ATP levels, and that AMPK activity alone is insufficient.

To investigate how proAX increases intracellular ATP levels, changes in ATP levels were examined in HDF-75 cells after the addition of oligomycin, an inhibitor of ATP synthase (Figure 3c). In the absence of proAX, the addition of oligomycin did not significantly change the intracellular ATP levels. In contrast, with proAX, the addition of oligomycin led to a dramatic decrease in intracellular ATP levels.

Next, we investigated the effects of proAX on mitochondrial respiration using an extracellular flux analyzer. At 3 and 24 h after proAX treatment, significant increases in basal respiration, maximal respiration, proton leakage, ATP production, and spare-respiratory capacity were observed (Figure 3d, e). These results demonstrated that proAX activates mitochondrial respiration-mediated energy metabolism.

Upon examining the changes in glucose consumption (Figure 3f) and lactate production (Figure 3g) before and after proAX administration, glucose consumption and lactate production were significantly increased 24 h after proAX administration.

Figure 3. Mechanisms of augmentation of intracellular ATP by proAX administration. a, Intracellular ATP changes in HDF-75 treated with 100 µM proAX in the presence or absence of nucleoside transport inhibitor, dipyridamole, b, Intracellular ATP changes in HDF-75 treated with 100 µM proAX in the

presence or absence of the AMPK-inhibitor compound-C (dorsomorphin), c, Intracellular ATP changes in HDF-75 treated with 100 μM proAX in the presence or absence of the ATP synthase inhibitor, oligomycin. The effects of proAX on mitochondrial respiration and their functions measured by an extracellular flux analyzer at 3 (d), and 24 (e) hours after treatment of 50 μM proAX. The effects of proAX on glucose consumption (f) and lactate production (g) at 3, 6, and 24 hours after treatment of 100 μM proAX. For all panels, mean ± sd shown, along with statistical significance between relevant comparisons. Significance is calculated using a one-way ANOVA with Dunnett's multiple comparisons test (a-c). For panel d-g, significance was calculated using a two-tailed unpaired t-test. ****$p < 0.0001$, ***$p < 0.001$. ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, ns indicates not significant.

2.5. The effects of proAX on oxidative stress resistance

Mitochondria generate cellular energy through oxidative phosphorylation (OXPHOS); however, this process also produces reactive oxygen species (ROS) as byproducts. ROS damage the mitochondrial DNA (mtDNA), proteins, and lipids. To further investigate the impact of proAX on cellular robustness, its effects on oxidative stress resistance were examined in both cultured cells and *C. elegans*. Upon exposure to oxidative stress induced by hydrogen peroxide (Figure 4a), a comparison of oxidative stress resistance between young and aged cells revealed that at all hydrogen peroxide concentrations, HDF-75 consistently had lower cell counts than HDF-29. This indicates a reduced resilience to oxidative stress in aged cells. Pretreatment of HDF-75 cells with proAX before adding hydrogen peroxide mitigated the decline in cell number, yielding a reduction in cell viability compared to that of HDF-29 cells. This suggests that proAX treatment enhances the resilience of aged cells to oxidative stress.

Paraquat, a mitochondrial toxicant, affects electron transport chain function through redox reactions, thereby increasing superoxide production[28]. To investigate whether proAX exhibits antioxidant stress effects at the organismal level, we selected *C. elegans* as the model organism. First, we examined whether proAX enhances the in vivo ATP levels at the organismal level in nematodes (Figure 4b). ProAX addition significantly increased the ATP levels per worm. These results clarify that proAX can increase ATP levels not only in cultured cells but also in individual organisms.

Here, the effects of proAX on oxidative stress resistance were examined by exposing 7-day-old *C. elegans* transferred to a NGM medium containing proAX to 200 mM paraquat. The results indicate that *C. elegans* cultured in the presence of proAX had a significantly extended lifespan (Figure 4c). Extension rate was 23%. Consequently, proAX enhances resilience to oxidative stress at the organismal level.

The effects of proAX on cells, obtained through experiments using cultured cells, are summarized in Figure 5.

Figure 4. Effect of proAX on oxidative stress resistance in fibroblast cells and *Caenorhabditis elegans*. a, Fibroblasts derived from donors of different ages were treated with indicated concentrations of hydrogen peroxide following pretreatment with or without proAX (100 μM). b, ATP levels in *C. elegans* on Day 7 when cultured with or without proAX (100 μM) (n = 25-35 worms). c, Effect of paraquat-induced stress on *C. elegans* pretreated with proAX (100 μM) for 7 days (n = 49 worms, each). For all panels, mean ± standard deviation shown, along with statistical significance between relevant comparisons. For the panel b, the center line of the box shows the median, box limits show interquartile ranges, and whiskers show the range of the data. Significance is calculated using one-way ANOVA with Tukey's multiple comparisons test (a), a two-tailed unpaired t-test (b), and log-rank test (c). ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, ns indicates not significant.

Figure 5. Summary of the effects of proAX on intracellular energy metabolism. ProAX penetrates cells, increases the AXP pool, and activates AMPK. The activation of AMPK promotes fatty acid oxidation and stimulates mitochondrial respiration. Furthermore, proAX is observed to enhance resistance to oxidative stress.

2.6. The effects of proAX on the lifespan of C. elegans

As *C. elegans* experiences a significant decrease in intracellular ATP levels with aging [9], it could be a suitable model for confirming the effects of ATP elevation. *C. elegans* were treated with various concentrations of proAX, ranging from 25 to 300 μM, and their lifespans were measured (Figure 6a, and Figure S4a-d). Surprisingly, proAX concentrations over 50 μM resulted in significantly increased longevity compared to that in the control group without proAX. Figure 6a shows the survival curves of *C. elegans* treated with 100 μM proAX and without the prodrug (control). In contrast, AMP, ATP, or AICAR did not significantly extend the lifespan of *C. elegans* (Figure 6b). Table 1 summarizes and compares the effects of proAX, AMP, ATP, and AICAR on lifespan extension. When varying the concentration of proAX, 100 μM proAX was found to have the highest efficacy in extending lifespan

(extension rate 24%, $p < 0.0001$). These results demonstrated the efficacy of proAX in extending the lifespan of *C. elegans*. To exclude the possibility of dietary restriction (DR) effects from proAX supplementation, worm body size was measured, as DR is known to increase longevity and reduce the growth of *C. elegans*. Furthermore, drugs that activate AMPK, such as metformin, have been reported to extend the lifespan of *C. elegans*, while inhibiting their growth[29]. There were no significant differences in body length between the treatments with or without proAX from Days 1 to 3 (Figure 6c). This was also the case for the body length differences between Days 4 and 20 (data not shown). Using dead bacterial lawns in lifespan experiments helps elucidate the impact of bacterial metabolism on the lifespan of *C. elegans*. Therefore, the bacteria were killed by heat, and lifespan analysis was performed in the same manner as for live bacteria. A significant lifespan extension effect of proAX was observed even when feeding on dead *Escherichia coli* (*E. coli*) (Figure S4e, and Table S1). This result confirms that the effect of proAX is not dependent on *E. coli* or its metabolism.

Figure 6. ProAX extends lifespan of *Caenorhabditis elegans*. a, Survival curves of *C. elegans* treated with proAX (100 µM). b, Survival curves of *C. elegans* treated with AMP, ATP, and AICAR (100 µM each). c, Effect of proAX on the growth of *C. elegans* over three days. For all panels, mean ± standard deviation shown, along with statistical significance between relevant comparisons. Significance is calculated using log-rank test (a, b), a two-tailed unpaired t-test (c), and ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, ns indicates not significant.

Table 1. Effect of proAX and other drugs on lifespan extension of *C. elegans*.

sample	Mean lifespan ± SE (days)	Extension rate (%)	N	*p* value (log-rank)
Control (no drug)	17.5 ± 0.3	-	326	-
25 µM proAX	18.1 ± 0.4	3.4	116	0.720
50 µM proAX	19.4 ± 0.5	10.9	111	0.002
100 µM proAX	21.7 ± 0.2	24	513	$< 10^{-4}$
200 µM proAX	19.3 ± 0.5	10.3	95	0.004
300 µM proAX	20.2 ± 0.6	15.4	69	0.0002
100 µM AMP	18.1 ± 0.4	3.4	145	0.823
100 µM ATP	17.7 ± 0.4	1.1	142	0.891
100 µM AICAR	18.0 ± 0.4	2.8	111	0.715

3. Discussion

The relationship between aging and energy metabolism has long been the subject of discussion. Among various factors that contribute to aging, mitochondrial dysfunction plays a pivotal role in the aging process. This dysfunction is also associated with the onset of age-related diseases. Small molecules essential for life activities such as NAD^+ and taurine[30,31], are known to slow the progression of aging in various animals. Reports suggest that supplementing these molecules from outside the body can extend the lifespan. Similarly, ATP levels decrease with age. The decline in ATP levels associated with reduced energy metabolism is thought to disrupt energy homeostasis and potentially accelerate aging. However, there are few reports on the effective supplementation of intracellular ATP levels with drugs. Various genetic mutations are known to extend the lifespan of *C. elegans*, with a significant increase in ATP levels observed in many mutant strains[14]. Kipreos et al. suggest that while genetic mutations increasing ATP levels benefit longevity, they propose the "energy maintenance theory of aging (EMTA)" as not extending longevity but as essential for survival maintenance in older animals[14]. Our results illuminate how AXP energy metabolism stimulation could provide a new paradigm for the role of energy metabolism in aging, particularly through its involvement in mitochondrial activity, the AXP pool, and the energy sensor AMPK.

We demonstrated that synthesized proAX increased intracellular ATP levels (Figure 1c). We found that the number of cells, 24 h after proAX administration, decreased in a concentration-dependent manner (Figure S2c, d). LDH and live/dead assays revealed no cytotoxicity of proAX (Figure 1e and Figure S2b), suggesting that the decrease in cell number was due to the suppression of cell proliferation. Numerous studies have shown that AMPK activators, such as metformin and AICAR, inhibit cell proliferation through AMPK activation[32,33]. AMPK conserves energy and enhances ATP utilization efficiency by inhibiting energy-intensive processes such as protein synthesis and cell proliferation. We confirmed that AICAR inhibited the proliferation of HDF-75 cells (Figure S3b). AMPK activation promotes catabolic pathways to restore energy homeostasis and enhance ATP production. However, addition of the conventional AMPK activator AICAR to HDF-75 cells did not increase intracellular ATP levels within 24 h (Figure S3c).

As expected from the molecular design, proAX was first converted into AMP in the cell, which significantly increased its concentration. Subsequently, the cellular levels of ADP and ATP increased (Figure 2a-c). Consequently, the intracellular reactions triggered by proAX increased the adenine nucleotide pool (Figure 2f). The energy charge stayed around 0.9 from 3 to 72-h post-proAX administration, indicating no abnormal cellular energy deficiency or excess (Figure 2d). AMPK activation is generally triggered by decreased energy status or stress. However, energy-charge fluctuations during proAX administration suggested that the cells did not experience significant energy stress, implying a temporary increase in AMP concentration (AMP/ATP) caused by proAX-induced AMPK activation (Figure 2g). The elevation of AMP and ADP levels by proAX may protect AMPK from dephosphorylation, potentially sustaining its activity longer[34]. This indicates that the prodrug may cause the cells to perceive pseudo-energy depletion.

The intracellular uptake pathway and the activation mechanism of the prodrug were examined. ProTide technology was developed to efficiently deliver nucleotide analogs by enhancing cell membrane permeability[35]. Investigations into the contribution of nucleoside transporters to the intracellular uptake of proAX suggested that, in addition to passive diffusion, some proportion of the prodrug was internalized via the transporters (Figure 3a), along with other nucleoside drugs for cancer chemotherapy[24]. Once inside the cell, proAX is thought to undergo cellular metabolic reactions similar to those of other ProTides. It is transformed into an alanine intermediate through esterase hydrolysis and converted into its monophosphate form (AMP) by phosphoamidases.

AMPK activation inhibits anabolic pathways and reduces ATP consumption. Simultaneously, it activates catabolic pathways that enhance ATP production via β-oxidation and glycolysis. Studies have shown that AMPK activation in the heart[36] and monocytes[37] leads to activation of the glycolytic

pathway. It has been reported that AMPK increases the flux of the glycolytic pathway[38]. In this study, glucose consumption and lactate production did not increase during short culture periods (3, and 6 h), but both increased after 24 h (Figure 3f, g). This suggests that the non-oxygen-dependent glycolytic pathway may be promoted by a sufficient increase in ATP levels. Kobayashi et al. demonstrated that AMPK activation increased fatty acid oxidation, leading to elevated mitochondrial respiration[39]. Our results suggest that AMPK activation enhances FAO (Figure 2i), thereby improving ATP production. However, as ATP production was not enhanced by AMPK activators such as AICAR (Figure S3c), it is conceivable that proAX not only activates AMPK, but also enhances ATP production through other mechanisms.

Several enzymes, including adenylate kinases, phosphorylate antiviral nucleotide analogs and natural AMP to ADP. Subsequently, these diphosphorylated drugs are further phosphorylated by multiple enzymes, including nucleoside diphosphate kinases, phosphoglycerate kinases, pyruvate kinases, and creatine kinase[40]. In the case of proAX, after being metabolized by AMP, these enzymes can convert AMP to ADP, and ADP to ATP. Although it is not yet possible to estimate the proportions of these reactions that contribute to ATP production accurately, experiments using oligomycin, an ATP synthase inhibitor, have shown that ATP production is significantly suppressed. This suggests that ATP production by proAX in the mitochondria is predominant. A close relationship exists between adenosine homeostasis, adenine nucleotide pool size, and mitochondrial bioenergetics[41–43]. When the concentration of ADP in the cell increases, it is transported into the matrix through the ADP/ATP exchange transport system, promoting ATP synthesis. ADP regulation of respiration leads to increased oxygen consumption in the mitochondria[44]. In other words, the prodrug likely boosted ADP levels, serving as a substrate for oxidative phosphorylation, thereby enhancing ATP production via mitochondrial respiration (Figure 3d, e). Kamatani et al. suggested that enhancing the purine salvage pathway, thereby reducing de novo purine synthesis, decreasing energy expenditure, and increasing the purine pool, leads to enhanced cellular ATP[45]. The critical role of mitochondria in cellular processes is well-established, with evidence linking mitochondrial dysfunction to aging and age-related diseases[46,47]. Strategies to enhance mitochondrial quality and function show the potential for widespread benefits, including aging and age-related disease prevention and treatment, as well as for improving overall cellular health and longevity. Our results demonstrate that the administration of proAX significantly enhances the energy metabolism of HDF cells via OXPHOS. This enhancement of energy metabolism may result from enhanced fatty acid oxidation due to AMPK activation and an increased adenine nucleotide pool available for ATP synthase.

Studies have shown that the intracellular adenine nucleotide pool correlates with the longevity of organisms such as yeast[48], *C. elegans*[49,50], and *Drosophila*[51]. In these organisms, lifespan extension is often linked to genetic mutations that increase the AMP/ATP ratio by affecting the adenine nucleotide pool, thereby activating AMPK.

Therapeutic approaches that activate AMPK, such as caloric restriction and metformin, have been shown to improve longevity and healthspan[52,53]. The effects of AMPK are elicited through mechanisms involving interactions with various factors. However, other studies have reported that AICAR does not extend *C. elegans*'s lifespan[54]. Our experiments confirmed that AICAR administration did not extend the lifespan of *C. elegans* (Figure 6b). Therefore, although AMPK activation influences the lifespan extension in *C. elegans*, its effect alone may be insufficient, indicating the involvement of other factors. This raises the question: What impact does enhancing intracellular ATP levels have on the lifespan of *C. elegans*? Insulin receptor mutants in C. elegans have been shown to promote mitochondrial fusion, increase ATP levels, and extend the lifespan[55]. Although few methods are available to pharmacologically enhance ATP levels, some studies have explored the relationship between increased ATP levels and the lifespan of *C. elegans*. Abe et al. reported that MA-5, which binds to ATP synthase and enhances ATP production, does not extend *C. elegans*'s lifespan[56]. Other reports suggest that a combination of febuxostat and vitamin C, an antioxidant, favors the purine salvage pathway, thereby aiding ATP production and extending the lifespan of *C. elegans* through antioxidant effects, which can be considered supportive of our study's findings[57]. In the present study, we demonstrated that proAX, without the addition of antioxidants, such as vitamin C, independently enhanced antioxidant resistance in both fibroblasts and *C. elegans* (Figure 4a, c). Decreased intracellular ATP levels lead to a reduced oxidative stress resistance[3]. Therefore, enhancing ATP production or activating the AMPK-FOXO pathway, which activates the antioxidant system via thioredoxin, may help lower ROS levels and improve oxidative stress resistance[58,59]. However, further detailed investigations are necessary to clarify the precise mechanisms underlying the enhanced resistance to oxidative stress by proAX. Although, activating mitochondrial respiration may potentially increase ROS production, the antioxidant effects of proAX could offset ROS generation caused by enhanced mitochondrial respiration.

4. Conclusion

Overall, proAX, which is converted into a natural energy currency, increases the intracellular adenine nucleotide pool. It exhibits various effects, including AMPK activation, enhanced FAO, and activation of mitochondrial respiration, leading to increased ATP production and improved resistance to oxidative stress. These combined actions are believed to significantly extend the average lifespan of *C. elegans*.

Therefore, the novel concept proposed by this technology, AXP energy metabolism stimulation (AXP stimulation), may inhibit the decline in energy metabolism and the biological defense mechanisms associated with aging, thereby preventing the onset of age-related diseases. Although further studies are necessary to understand how proAX extends the lifespan of *C. elegans*, the results of this study support the significant role of the adenine energy metabolism pathway in controlling lifespan, suggesting that drugs targeting these pathways may influence lifespan.

5. Experimental Section

Synthesis of proAX: ProAX was synthesized and its chemical identity was established using nuclear magnetic resonance (NMR) and fast atom bombardment mass spectrometry (FABMS) analyses, as described in the Supplementary Materials.

Cell cultures: Human dermal fibroblasts (HDFs) derived from 29- or 75-year-old donors were obtained from Cell Applications, Inc. (San Diego, CA, USA). Normal human dermal fibroblasts (NHDFs) were obtained from Lonza (Basel, Switzerland). HDF were grown at 37 ℃ and 5 % CO_2 in HDF growth medium (Cell Applications, Inc., San Diego, CA, USA). NHDFs were grown at 37 ℃ and 5 % CO_2 in DMEM/F12 supplemented with 10% FBS and 1% penicillin (all from Gibco, Carlsbad, CA, USA). ProAX was added at 3 h after seeding.

Reagents: Chemicals used in this study: Adenosine 5'-monophosphate (#017-20391, Fuji Film Wako Pure Chemicals, Osaka, Japan), Adenosine 5'-triphosphate (#019-09672, Fuji Film Wako Pure Chemicals, Osaka, Japan), AICAR (#015-22531, Fuji Film Wako Pure Chemicals, Osaka, Japan), Dipyridamole (#D2274, TCI, Tokyo, Japan), Dorsomorphin (#D5349, TCI, Tokyo, Japan), Etomoxir sodium salt hydrate (#E1905, sigma, St. Louis, MO, USA), Hydrogen Peroxide (#H1222, TCI, Tokyo, Japan), Oligomycin (#G270, Dojindo, Kumamoto, Japan), Paraquat (#D3685, TCI, Tokyo, Japan).

Antibodies: Antibodies used in this study: Anti-AMPKα (#2532; 1:1000), Anti-pThr172-AMPK (#2535; 1:1000), Anti-Acetyl-CoA Carboxylase (#3676; 1:1500), Anti-Phospho-CoA Carboxylase (#11818; 1:1500) from Cell Signaling Technology. Horseradish peroxidase-conjugated anti-rabbit IgG (#111-035-144; 1:10000) from Jackson ImmunoResearch.

Intracellular ATP Levels assay by Luciferin-luciferase: Intracellular ATP levels in HDFs incubated with different concentrations of proAX were measured using a luciferin–luciferase-based ATP assay kit (Dojindo, Kumamoto, Japan) according to the manufacturer's instructions. HDFs were seeded at 5,000 cells/well in 96-well plates. After 24 h of proAX treatment, cells were lysed in 0.2% Triton X-100 solution and sonicated in an ice bath. DNA concentration in cell lysate was measured

using a Quant-iT PicoGreen dsDNA kit (Thermo Fisher Scientific, Waltham, MA, USA). The luciferin-luciferase luminescence was measured using an Infinite 200PRO M Plex microplate reader (Tecan, Zürich, Switzerland). The ATP levels was normalized using the cell number, as determined by the DNA concentration.

Cytotoxicity Assay: The cytotoxicity of NHDF and HDF derived from a 75-year-old donor incubated with different concentrations of proAX was measured using a lactate dehydrogenase (LDH) assay kit (Dojindo, Kumamoto, Japan) or a Cellstain double staining kit (Dojindo, Kumamoto, Japan) according to the manufacturer's instructions. For the LDH assay, NHDFs were seeded at 5,000 cells/well in 96-well plates. After 24 h of treatment with proAX, the absorbance of the cells was measured at 490 nm using an Infinite 200PRO M Plex microplate reader (Tecan, Zürich, Switzerland). For Live/Dead cell staining, HDF were seeded at a density of 20,000 cells/well into 96-well plates. After 3 h of treatment with proAX, images were obtained using a Keyence BZ-X710 microscope (Keyence, Osaka, Japan).

Metabolite extraction and metabolome analysis[60]**:** After the culture medium was aspirated, plated HDF-75 cells were washed twice with PBS solution (10 mL and then 2 mL), treated with 740 µL of methanol, and left at rest for 30 s to inactivate enzymes. The cell extract was then treated with 500 µL of Milli-Q water containing internal standards (10 µM methionine sulfone) and left at rest for a further 30 s. Three hundred and fifty microliters of the extract was centrifugally filtered through a Millipore 5 kDa cutoff filter (UltrafreeMC-PLHCC, HMT) to remove macromolecules (9,100 × g, 4 ℃, 120 min). The filtrate was centrifugally concentrated and re-suspended in 30 µL of acetonitrile/ Milli-Q water (1:1 v/v) for metabolome analysis at Infinity Lab (Yamagata, Japan). HPLC instrument modules were obtained from the 1290 Infinity line (Agilent Technologies): binary pumps (Model G7120A), an autosampler (G7167B), and a temperature-controlled column compartment (Model G7116B). A stock solution of 100 mM ammonium acetate was prepared by dissolving ammonium acetate (0.1 mol) in water, adjusting to pH 9.0 with ammonium hydroxide, and correcting the final volume to 1 L with water. Hydrophilic interaction chromatography (HILIC) was performed using a 100 mm × 2.1 mm i.d. InfinityLab Poroshell 120 HILIC-Z column (Agilent Technologies). Solvent A was prepared by mixing 100 mL of the stock solution with 900 mL of water to yield a final ammonium acetate concentration of 10 mM (pH 9.0). Solvent B was prepared by mixing 100 mL of the stock solution with 900 mL of acetonitrile to yield a final concentration of 10 mM ammonium acetate (pH 9.0) in 80% acetonitrile. The deactivator additive (5 mM methylenediphosphonic acid) was spiked into the solvents for a final concentration of 5 µM for analysis. The flow rate was 0.3 mL/min, and the column temperature was set at 45 °C. A sample volume of 1 µL was injected into the column for each experiment. The gradient

began initially at 100% B, decreased linearly to 90% at 0.5 min, decreased linearly again to 70% at 10 min, and subsequently re-equilibrated at 95% B for a further 3 min. The sample temperature was maintained at 4 °C in the autosampler prior to analysis. Mass spectrometry analysis was performed using an Agilent 6470 B triple quadrupole mass spectrometer (Agilent Technologies) equipped with a Jet Stream ESI probe in positive-ion mode. A capillary voltage of 3,500 V, a source temperature of 200 °C, gas flow of 10 L/min, and nebulizer of 40 psi, sheath gas temperature of 300 °C, sheath gas flow of 12 L/min were used. Raw data were sampled at 667 ms/cycle in Dynamic Multiple Reaction Monitoring (MRM) mode. The concentrations of individual nucleotides were calculated from the peak area in the chromatogram detected using MRM relative to the internal standard methionine sulfone.

siRNA transfection: HDFs derived from the 75-year-old donor were transfected with AMPKα2 siRNA oligonucleotides (5' ACCGAGCUAUGAAGCAGCUGGAUUU 3') (Invitrogen, Waltham, MA, USA). Lipofectamine RNAiMAX (Thermo Fisher Scientific) was used for transfection according to the manufacturer's instructions. For each well of 96-well plate, 1 pmol of siRNA oligonucleotides and 0.3 μl lipofectamin RNAiMAX was added into 100 μl culture media of the cells.

Immunoblotting: Cells were lysed in RIPA buffer containing protease and phosphatase inhibitor cocktails (Nacalai Tesque, Kyoto, Japan). Cell lysates were sonicated for 10 s and heated at 95 °C for 5 min. Cell lysates were separated on 12 % or 4–20 % gels (Bio-Rad, Hercules, CA, United states) for SDS-PAGE. The proteins were transferred onto PDVF membranes (Bio-Rad) and blocked in blocking buffer (Nacalai Tesque) for 1 h. Next, the membranes were incubated overnight at 4 °C with the primary antibodies. The membranes were rinsed three times with TBS-T ((Tris-buffered saline)- 0.01% Tween), then incubated with secondary HRP-conjugated antibodies for 1 h at 25 °C. The membrane was washed with TBS-T and treated with ImmunoStar Zeta reagent (Wako Pure Chemical Industries). Chemiluminescent images of the membranes were acquired using LuminoGraph I (WSE-6100; ATTO).

Fatty acid oxidation (FAO) activity: FAO activity in HDFs derived from a 75-year-old donor was detected using FAOBlue (Funakoshi, Tokyo, Japan), according to the manufacturer's instructions. HDFs were seeded at 5,000 cells/well in 96-well plates and treated with proAX (final concentration 100 μM) for 3, 6, and 24 h. The cells were then treated with FAO blue (final concentration 5 μM) for 2 h. The fluorescence (excitation/emission = 405/460 nm) of the cells was measured using an Infinite 200PRO M Plex microplate reader (Tecan, Zürich, Switzerland).

Glucose consumption and lactate release: HDFs derived from 75-year-old donors were seeded at 5,000 cells/well in 96-well plates and treated with proAX (final concentration 100 μM) for 3, 6, and 24 h. The supernatant was collected and diluted 1:30 for the glucose assay or 1:10 for the lactate assay in ultrapure water. The diluted supernatant was measured using a glucose assay kit or

glycolysis/OXPHOS assay kit (Dojindo, Kumamoto, Japan). Absorbance was measured at 450 nm using an Infinite 200PRO M Plex microplate reader (Tecan, Zürich, Switzerland).

Extracellular flux analyzer: A Seahorse XF HS Mini Analyzer (Agilent Technologies) was used to measure OCR. HDFs derived from 75-year-old donors were seeded at 8,000 cells/well in XFp cell culture miniplates and treated with proAX (final concentration, 50 μM) for 3 and 24 h. Mitochondrial functions were examined through sequential injections of oligomycin (1.5 μM), Carbonyl cyanide-4 (trifluoromethoxy) phenylhydrazone (FCCP) (5 μM), and rotenone/antimycin A (Rot/AA) (0.5 μM). The obtained data were analyzed using the Wave software (version 10.0.1.72; Agilent Technologies).

Oxidative stress assay in HDFs: The viability of HDFs was measured using the cell counting kit-8 (CCK-8) (Dojindo, Kumamoto, Japan). HDFs were seeded at 5,000 cells/well in 96-well plates. Treatment with proAX (final concentration 100 μM), HDFs were incubated for 24 h. After washing out the pretreatment medium, cells were exposed to H_2O_2 (300–400 μM) in serum-free medium (DMEM/F12) at 37 ºC. After incubation for 2 h, the absorbance of the cells was measured at 450 nm using an Infinite 200PRO M Plex microplate reader (Tecan, Zürich, Switzerland).

C. elegans cultures: Wild-type N2 *C. elegans* were provided by the Caenorhabditis Genetics Center (CGC, University of Minnesota). The worms were grown and maintained on nematode growth medium (NGM) plates spread with live or dead *E. coli* OP50 as food. Dead OP50 was prepared by incubating in 65 ºC for 30 min. All the worms were cultured at 25 ºC. proAX was added to the NGM plate to a final concentration of 25–300 μM. The comparison samples were used in which 100 μM AMP, ATP, and AICAR were added to the NGM plate. All NGM plates were supplemented with DMSO at a final concentration of 1%.

Lifespan Assays: After the age-synchronized worms were cultured to the L4 stage, 35 worms were transferred to the desired NGM plates. The worms were transferred to fresh NGM plates with or without drugs every 1–2 days. Plates without any added drugs were used as the control. Live and dead worms were counted during transfer. Worms that displayed no movement after gentle contact with a platinum picker were considered dead. Kaplan–Meier curves were generated using Graph Pad Prism 10 (GraphPad Software, CA, USA) and calculated by Excel (Windows 365).

Body length of C. Elegans: After age synchronization, the worms were transferred to the desired NGM plates and cultured for 72 h. The body length of each live worm was measured using a VHX-900F instrument (KEYENCE, Osaka, Japan).

ATP levels of C. Elegans: After age synchronization, the worms were transferred to the NGM plates with or without proAX and cultured for 4 days. Bacteria attached to the bodies of 7-day-old worms were removed by allowing the worms to crawl onto an NGM agar plate with M9 buffer. Worms were collected in 60 µM M9 buffer, and endogenous ATP was extracted by sonication on ice for 1 min (Ultrasonic Homogenizer UP-21P; TOMY, Tokyo, Japan) and centrifugation (15,000 rpm, 10 min, 4 ºC. An ATP assay kit (Dojindo, Kumamoto, Japan) was used to measure the endogenous ATP levels. Luminescence was measured using an Infinite 200PRO M Plex microplate reader (Tecan).

Paraquat sensitivity of C. Elegans: After age synchronization, the worms were transferred to the NGM plates with or without proAX and cultured for 4 days. To assay paraquat sensitivity, the 7-day-old worms were collected and transferred to NGM plates containing 200 mM paraquat. Worms that displayed no movement after gentle contact with a platinum picker were considered dead.

Statistics and reproducibility: Data are presented as mean ± s.d. for the indicated number of biological replicates (n) from least three independent experiments. Statistical analyses were performed using GraphPad Prism v.10.2 (GraphPad Software, CA, USA). The relevant statistical tests are described in the figure legends. Time-to-event data were analyzed using Kaplan–Meier curves, and differences in survival were determined using the log-rank test by Excel (for Microsoft 365). P set at $p <$ 0.05 was significant.

Acknowledgements

General: We thank to Takamasa Ishikawa and Ryuhei Kudo (Infinity Lab, INC, Yamagata, Japan) for metabolome analysis and measurement of AXP.

Funding: This study was supported in part by Grants-in-Aid for Scientific Research (KAKENHI) Grant Number JP23K18580 (for T. A.), JP19H05720 (for M. T.), JP22K19759 (for E. N. K.) from the Japan Society for the Promotion of Science (JSPS). Additionally, this study was partially supported by the Dynamic Alliance Open Innovation Bridging Human, Environment and Materials from the Ministry of Education, Culture, Sports, and Technology of Japan (MEXT) (for M. T., and T. A.), CORE[2]-A lab (for T. A.). This study was partially supported by AMED-Hashiwatashi research program A (for T. A.), JST SPRING Japan Grand Number JPMJSP2136 (for M. K.) and Nagase Science and Technology Foundation (for E. N. K.).

Conflict of Interest
The authors declare no conflict of interest.

Author Contributions

T.A. conceived the idea with assistance from M.T. T.A and M.T. planned the project. T.A. designed experiments. M.K., T.S., R.K. S.K. and T.A. mainly collected experimental data and carried out the analyses. T.A., M.K., T.S., R.K., E.K.N. S.K. and M.T. interpreted the data through the discussion. T. A., E. K.N., and M. T. obtained founding for the project. The manuscript was written through contributions of T.A., and M. K. All the authors have approved the final version of the manuscript.

Author Information

Corresponding Authors

T. Anada, M. Tanaka

Institute for Materials Chemistry and Engineering, Kyushu university, 744 Motooka, Nishi-ku Fukuoka 819-0395, Japan

E-mail: takahisa_anada@ms.ifoc.kyushu-u.ac.jp, masaru_tanaka@ms.ifoc.kyushu-u.ac.jp

Authors

T. Anada, M. Kawahara, T. Shimada, R. Kuroda, M. Tanaka

Department of Applied Chemistry, Graduate School of Engineering, Kyushu University, 744 Motooka, Nishi-ku Fukuoka 819-0395, Japan

E. Kage-Nakadai

Laboratory of Environment and Health Science, Graduate School of Human Life and Ecology, Osaka Metropolitan University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan

E. Kage-Nakadai

Institute for Life and Medical Sciences, Kyoto University, 53 Shogoin Kawahara-cho, Sakyo-ku, Kyoto 606-8507, Japan

References

(1) Shaulson, E. D.; Cohen, A. A.; Picard, M. The Brain–Body Energy Conservation Model of Aging. *Nat Aging* **2024**, in press. https://doi.org/10.1038/s43587-024-00716-x.

(2) Amorim, J. A.; Coppotelli, G.; Rolo, A. P.; Palmeira, C. M.; Ross, J. M.; Sinclair, D. A. Mitochondrial and Metabolic Dysfunction in Ageing and Age-Related Diseases. *Nat Rev Endocrinol* **2022**, *18* (4), 243–258.

(3) Miyoshi, N.; Oubrahim, H.; Chock, P. B.; Stadtman, E. R. Age-Dependent Cell Death and the Role of ATP in Hydrogen Peroxide-Induced Apoptosis and Necrosis. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103* (6), 1727–1731.

(4) Rabini, R. A.; Petruzzi, E.; Staffolani, R.; Tesei, M.; Fumelli, P.; Pazzagli, M.; Mazzanti, L. Diabetes Mellitus and Subjects' Ageing: A Study on the ATP Content and ATP-related Enzyme

Activities in Human Erythrocytes. *European Journal of Clinical Investigation* **1997**, *27* (4), 327–332.

(5) Short, K. R.; Bigelow, M. L.; Kahl, J.; Singh, R.; Coenen-Schimke, J.; Raghavakaimal, S.; Nair, K. S. Decline in Skeletal Muscle Mitochondrial Function with Aging in Humans. *Proceedings of the National Academy of Sciences* **2005**, *102* (15), 5618–5623.

(6) Kim, J. H.; Shrago, E.; Elson, C. E. Age-Related Changes in Respiration Coupled to Phosphorylation. II. Cardiac Mitochondria. *Mechanisms of Ageing and Development* **1988**, *46* (1), 279–290.

(7) Gkotsi, D.; Begum, R.; Salt, T.; Lascaratos, G.; Hogg, C.; Chau, K.-Y.; Schapira, A. H. V.; Jeffery, G. Recharging Mitochondrial Batteries in Old Eyes. Near Infra-Red Increases ATP. *Experimental Eye Research* **2014**, *122*, 50–53.

(8) Vernace, V. A.; Arnaud, L.; Schmidt-Glenewinkel, T.; Figueiredo-Pereira, M. E. Aging Perturbs 26S Proteasome Assembly in Drosophila Melanogaster. *The FASEB Journal* **2007**, *21* (11), 2672–2682.

(9) Braeckman, B. P.; Houthoofd, K.; Vanfleteren, J. R. Assessing Metabolic Activity in Aging Caenorhabditis Elegans: Concepts and Controversies: Assessing Metabolic Rate in Aging C. Elegans, B. P. Braeckman et Al. *Aging Cell* **2002**, *1* (2), 82–88.

(10) Patel, A.; Malinovska, L.; Saha, S.; Wang, J.; Alberti, S.; Krishnan, Y.; Hyman, A. A. ATP as a Biological Hydrotrope. *Science* **2017**, *356* (6339), 753–756.

(11) Chistiakov, D. A.; Sobenin, I. A.; Revin, V. V.; Orekhov, A. N.; Bobryshev, Y. V. Mitochondrial Aging and Age-Related Dysfunction of Mitochondria. *BioMed Research International* **2014**, *2014*, e238463.

(12) Correia-Melo, C.; Passos, J. F. Mitochondria: Are They Causal Players in Cellular Senescence? *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2015**, *1847* (11), 1373–1379.

(13) Nicolson, G. L. Mitochondrial Dysfunction and Chronic Disease: Treatment With Natural Supplements. *Integr Med (Encinitas)* **2014**, *13* (4), 35–43.

(14) Chaudhari, S. N.; Kipreos, E. T. The Energy Maintenance Theory of Aging: Maintaining Energy Metabolism to Allow Longevity. *BioEssays* **2018**, *40* (8), 1800005.

(15) Grevendonk, L.; Connell, N. J.; McCrum, C.; Fealy, C. E.; Bilet, L.; Bruls, Y. M. H.; Mevenkamp, J.; Schrauwen-Hinderling, V. B.; Jörgensen, J. A.; Moonen-Kornips, E.; Schaart, G.; Havekes, B.; de Vogel-van den Bosch, J.; Bragt, M. C. E.; Meijer, K.; Schrauwen, P.; Hoeks, J. Impact of Aging and Exercise on Skeletal Muscle Mitochondrial Capacity, Energy Metabolism, and Physical Function. *Nat Commun* **2021**, *12* (1), 4773.

(16) Suomalainen, A.; Nunnari, J. Mitochondria at the Crossroads of Health and Disease. *Cell* **2024**, *187* (11), 2601–2627.

(17) Jäger, R.; Purpura, M.; Rathmacher, J. A.; Fuller, J. C.; Pitchford, L. M.; Rossi, F. E.; Kerksick, C. M. Health and Ergogenic Potential of Oral Adenosine-5′-Triphosphate (ATP) Supplementation. *Journal of Functional Foods* **2021**, *78*, 104357.

(18) Arts, I. C.; Coolen, E. J.; Bours, M. J.; Huyghebaert, N.; Stuart, M. A. C.; Bast, A.; Dagnelie, P. C. Adenosine 5′-Triphosphate (ATP) Supplements Are Not Orally Bioavailable: A Randomized, Placebo-Controlled Cross-over Trial in Healthy Humans. *J Int Soc Sports Nutr* **2012**, *9*, 16.

(19) Moser, G. H.; Schrader, J.; Deussen, A. Turnover of Adenosine in Plasma of Human and Dog Blood. *American Journal of Physiology-Cell Physiology* **1989**, *256* (4), C799–C806.

(20) Riteau, N.; Gasse, P.; Fauconnier, L.; Gombault, A.; Couegnat, M.; Fick, L.; Kanellopoulos, J.; Quesniaux, V. F. J.; Marchand-Adam, S.; Crestani, B.; Ryffel, B.; Couillin, I. Extracellular ATP Is a Danger Signal Activating P2X7 Receptor in Lung Inflammation and Fibrosis. *Am J Respir Crit Care Med* **2010**, *182* (6), 774–783.

(21) Kazemzadeh-Narbat, M.; Annabi, N.; Tamayol, A.; Oklu, R.; Ghanem, A.; Khademhosseini, A. Adenosine-Associated Delivery Systems. *Journal of Drug Targeting* **2015**, *23* (7–8), 580–596.

(22) Mehellou, Y.; Rattan, H. S.; Balzarini, J. The ProTide Prodrug Technology: From the Concept to the Clinic. *J. Med. Chem.* **2018**, *61* (6), 2211–2226.

(23) Sofia, M. J.; Bao, D.; Chang, W.; Du, J.; Nagarathnam, D.; Rachakonda, S.; Reddy, P. G.; Ross, B. S.; Wang, P.; Zhang, H.-R.; Bansal, S.; Espiritu, C.; Keilman, M.; Lam, A. M.; Steuer, H. M. M.; Niu, C.; Otto, M. J.; Furman, P. A. Discovery of a β- D -2′-Deoxy-2′-α-Fluoro-2′-β- *C* - Methyluridine Nucleotide Prodrug (PSI-7977) for the Treatment of Hepatitis C Virus. *J. Med. Chem.* **2010**, *53* (19), 7202–7218.

(24) Beigel, J. H.; Tomashek, K. M.; Dodd, L. E.; Mehta, A. K.; Zingman, B. S.; Kalil, A. C.; Hohmann, E.; Chu, H. Y.; Luetkemeyer, A.; Kline, S.; Lopez de Castilla, D.; Finberg, R. W.; Dierberg, K.; Tapson, V.; Hsieh, L.; Patterson, T. F.; Paredes, R.; Sweeney, D. A.; Short, W. R.; Touloumi, G.; Lye, D. C.; Ohmagari, N.; Oh, M.; Ruiz-Palacios, G. M.; Benfield, T.; Fätkenheuer, G.; Kortepeter, M. G.; Atmar, R. L.; Creech, C. B.; Lundgren, J.; Babiker, A. G.; Pett, S.; Neaton, J. D.; Burgess, T. H.; Bonnett, T.; Green, M.; Makowski, M.; Osinusi, A.; Nayak, S.; Lane, H. C. Remdesivir for the Treatment of Covid-19 — Final Report. *New England Journal of Medicine* **2020**, *383* (19), 1813–1826.

(25) Mackman, R. L. Phosphoramidate Prodrugs Continue to Deliver, The Journey of Remdesivir (GS-5734) from RSV to SARS-CoV-2. *ACS Med. Chem. Lett.* **2022**, *13* (3), 338–347.

(26) Herzig, S.; Shaw, R. J. AMPK: Guardian of Metabolism and Mitochondrial Homeostasis. *Nat Rev Mol Cell Biol* **2018**, *19* (2), 121–135.

(27) Atkinson, D. E.; Walton, G. M. Adenosine Triphosphate Conservation in Metabolic Regulation: RAT LIVER CITRATE CLEAVAGE ENZYME. *Journal of Biological Chemistry* **1967**, *242* (13), 3239–3241.

(28) Bora, S.; Vardhan, G. S. H.; Deka, N.; Khataniar, L.; Gogoi, D.; Baruah, A. Paraquat Exposure over Generation Affects Lifespan and Reproduction through Mitochondrial Disruption in *C. Elegans*. *Toxicology* **2021**, *447*, 152632.

(29) Wu, L.; Zhou, B.; Oshiro-Rapley, N.; Li, M.; Paulo, J. A.; Webster, C. M.; Mou, F.; Kacergis, M. C.; Talkowski, M. E.; Carr, C. E.; Gygi, S. P.; Zheng, B.; Soukas, A. A. An Ancient, Unified Mechanism for Metformin Growth Inhibition in C. Elegans and Cancer. *Cell* **2016**, *167* (7), 1705-1718.e13.

(30) Yoshino, J.; Baur, J. A.; Imai, S. NAD+ Intermediates: The Biology and Therapeutic Potential of NMN and NR. *Cell Metabolism* **2018**, *27* (3), 513–528.

(31) Singh, P.; Gollapalli, K.; Mangiola, S.; Schranner, D.; Yusuf, M. A.; Chamoli, M.; Shi, S. L.; Lopes Bastos, B.; Nair, T.; Riermeier, A.; Vayndorf, E. M.; Wu, J. Z.; Nilakhe, A.; Nguyen, C. Q.; Muir, M.; Kiflezghi, M. G.; Foulger, A.; Junker, A.; Devine, J.; Sharan, K.; Chinta, S. J.; Rajput, S.; Rane, A.; Baumert, P.; Schönfelder, M.; Iavarone, F.; di Lorenzo, G.; Kumari, S.; Gupta, A.; Sarkar, R.; Khyriem, C.; Chawla, A. S.; Sharma, A.; Sarper, N.; Chattopadhyay, N.; Biswal, B. K.; Settembre, C.; Nagarajan, P.; Targoff, K. L.; Picard, M.; Gupta, S.; Velagapudi, V.; Papenfuss, A. T.; Kaya, A.; Ferreira, M. G.; Kennedy, B. K.; Andersen, J. K.; Lithgow, G. J.; Ali, A. M.; Mukhopadhyay, A.; Palotie, A.; Kastenmüller, G.; Kaeberlein, M.; Wackerhage, H.; Pal, B.; Yadav, V. K. Taurine Deficiency as a Driver of Aging. *Science* **2023**, *380* (6649), eabn9257.

(32) Griss, T.; Vincent, E. E.; Egnatchik, R.; Chen, J.; Ma, E. H.; Faubert, B.; Viollet, B.; DeBerardinis, R. J.; Jones, R. G. Metformin Antagonizes Cancer Cell Proliferation by Suppressing Mitochondrial-Dependent Biosynthesis. *PLOS Biology* **2015**, *13* (12), e1002309.

(33) Rattan, R.; Giri, S.; Singh, A. K.; Singh, I. 5-Aminoimidazole-4-Carboxamide-1-β-D-Ribofuranoside Inhibits Cancer Cell Proliferation *in Vitro* and *in Vivo* via AMP-Activated Protein Kinase*. *Journal of Biological Chemistry* **2005**, *280* (47), 39582–39593.

(34) Xiao, B.; Sanders, M. J.; Underwood, E.; Heath, R.; Mayer, F. V.; Carmena, D.; Jing, C.; Walker, P. A.; Eccleston, J. F.; Haire, L. F.; Saiu, P.; Howell, S. A.; Aasland, R.; Martin, S. R.; Carling, D.; Gamblin, S. J. Structure of Mammalian AMPK and Its Regulation by ADP. *Nature* **2011**, *472* (7342), 230–233.

(35) Shelton, J.; Lu, X.; Hollenbaugh, J. A.; Cho, J. H.; Amblard, F.; Schinazi, R. F. Metabolism, Biochemical Actions, and Chemical Synthesis of Anticancer Nucleosides, Nucleotides, and Base Analogs. *Chem. Rev.* **2016**, *116* (23), 14379–14455.

(36) Marsin, A.-S.; Bertrand, L.; Rider, M. H.; Deprez, J.; Beauloye, C.; Vincent, M. F.; Van den Berghe, G.; Carling, D.; Hue, L. Phosphorylation and Activation of Heart PFK-2 by AMPK Has a Role in the Stimulation of Glycolysis during Ischaemia. *Current Biology* **2000**, *10* (20), 1247–1255.

(37) Marsin, A.-S.; Bouzin, C.; Bertrand, L.; Hue, L. The Stimulation of Glycolysis by Hypoxia in Activated Monocytes Is Mediated by AMP-Activated Protein Kinase and Inducible 6-Phosphofructo-2-Kinase *. *Journal of Biological Chemistry* **2002**, *277* (34), 30778–30783.

(38) Bando, H.; Atsumi, T.; Nishio, T.; Niwa, H.; Mishima, S.; Shimizu, C.; Yoshioka, N.; Bucala, R.; Koike, T. Phosphorylation of the 6-Phosphofructo-2-Kinase/Fructose 2,6-Bisphosphatase/PFKFB3 Family of Glycolytic Regulators in Human Cancer. *Clin Cancer Res* **2005**, *11* (16), 5784–5792.

(39) Kobayashi, H.; Hatakeyama, H.; Nishimura, H.; Yokota, M.; Suzuki, S.; Tomabechi, Y.; Shirouzu, M.; Osada, H.; Mimaki, M.; Goto, Y.; Yoshida, M. Chemical Reversal of Abnormalities in Cells Carrying Mitochondrial DNA Mutations. *Nat Chem Biol* **2021**, *17* (3), 335–343.

(40) Li, R.; Liclican, A.; Xu, Y.; Pitts, J.; Niu, C.; Zhang, J.; Kim, C.; Zhao, X.; Soohoo, D.; Babusis, D.; Yue, Q.; Ma, B.; Murray, B. P.; Subramanian, R.; Xie, X.; Zou, J.; Bilello, J. P.; Li, L.; Schultz, B. E.; Sakowicz, R.; Smith, B. J.; Shi, P.-Y.; Murakami, E.; Feng, J. Y. Key Metabolic Enzymes Involved in Remdesivir Activation in Human Lung Cells. *Antimicrobial Agents and Chemotherapy* **2021**, *65* (9), 10.1128/aac.00602-21.

(41) Schild, L.; Blair, P. V.; Davis, W. I.; Baugh, S. Effect of Adenine Nucleotide Pool Size in Mitochondria on Intramitochondrial ATP levels1Enzymes: Creatine Kinase (EC 2.7.3.2).1. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1999**, *1413* (1), 14–20.

(42) Aprille, J. R.; Austin, J. Regulation of the Mitochondrial Adenine Nucleotide Pool Size. *Archives of Biochemistry and Biophysics* **1981**, *212* (2), 689–699.

(43) Takaine, M.; Imamura, H.; Yoshida, S. High and Stable ATP Levels Prevent Aberrant Intracellular Protein Aggregation in Yeast. *eLife* **2022**, *11*, e67659.

(44) Klingenberg, M. The ADP and ATP Transport in Mitochondria and Its Carrier. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **2008**, *1778* (10), 1978–2021.

(45) Johnson, T. A.; Jinnah, H. A.; Kamatani, N. Shortage of Cellular ATP as a Cause of Diseases and Strategies to Enhance ATP. *Frontiers in Pharmacology* **2019**, *10*.

(46) Vafai, S. B.; Mootha, V. K. Mitochondrial Disorders as Windows into an Ancient Organelle. *Nature* **2012**, *491* (7424), 374–383.

(47) Andreux, P. A.; Houtkooper, R. H.; Auwerx, J. Pharmacological Approaches to Restore Mitochondrial Function. *Nat Rev Drug Discov* **2013**, *12* (6), 465–483.

(48) Matecic, M.; Jr, D. L. S.; Pan, X.; Maqani, N.; Bekiranov, S.; Boeke, J. D.; Smith, J. S. A Microarray-Based Genetic Screen for Yeast Chronological Aging Factors. *PLOS Genetics* **2010**, *6* (4), e1000921.

(49) Apfeld, J.; O'Connor, G.; McDonagh, T.; DiStefano, P. S.; Curtis, R. The AMP-Activated Protein Kinase AAK-2 Links Energy Levels and Insulin-like Signals to Lifespan in C. Elegans. *Genes Dev* **2004**, *18* (24), 3004–3009.

(50) Liu, Y.; Zhang, X.; Wang, Y.; Wang, J.; Wei, H.; Zhang, C.; Zhang, Q. Cyclocodon Lancifolius Fruit Prolongs the Lifespan of Caenorhabditis Elegans via Antioxidation and Regulation of Purine Metabolism. *Food Funct.* **2024**.

(51) Stenesen, D.; Suh, J. M.; Seo, J.; Yu, K.; Lee, K.-S.; Kim, J.-S.; Min, K.-J.; Graff, J. M. Adenosine Nucleotide Biosynthesis and AMPK Regulate Adult Life Span and Mediate the Longevity Benefit of Caloric Restriction in Flies. *Cell Metabolism* **2013**, *17* (1), 101–112.

(52) Weir, H. J.; Yao, P.; Huynh, F. K.; Escoubas, C. C.; Goncalves, R. L.; Burkewitz, K.; Laboy, R.; Hirschey, M. D.; Mair, W. B. Dietary Restriction and AMPK Increase Lifespan via Mitochondrial Network and Peroxisome Remodeling. *Cell Metabolism* **2017**, *26* (6), 884-896.e5.

(53) Martin-Montalvo, A.; Mercken, E. M.; Mitchell, S. J.; Palacios, H. H.; Mote, P. L.; Scheibye-Knudsen, M.; Gomes, A. P.; Ward, T. M.; Minor, R. K.; Blouin, M.-J.; Schwab, M.; Pollak, M.; Zhang, Y.; Yu, Y.; Becker, K. G.; Bohr, V. A.; Ingram, D. K.; Sinclair, D. A.; Wolf, N. S.;

Spindler, S. R.; Bernier, M.; de Cabo, R. Metformin Improves Healthspan and Lifespan in Mice. *Nat Commun* **2013**, *4* (1), 2192.

(54) Riedinger, C.; Mendler, M.; Schlotterer, A.; Fleming, T.; Okun, J.; Hammes, H.-P.; Herzig, S.; Nawroth, P. P. High-Glucose Toxicity Is Mediated by AICAR-Transformylase/IMP Cyclohydrolase and Mitigated by AMP-Activated Protein Kinase in Caenorhabditis Elegans. *J Biol Chem* **2018**, *293* (13), 4845–4859.

(55) Chaudhari, S. N.; Kipreos, E. T. Increased Mitochondrial Fusion Allows the Survival of Older Animals in Diverse C. Elegans Longevity Pathways. *Nat Commun* **2017**, *8* (1), 182.

(56) Wu, X.; Seida, M.; Abe, T.; Higashitani, A. Mitochonic Acid 5 Attenuates Age-Related Neuromuscular Dysfunction Associated with Mitochondrial Ca2+ Overload in Caenorhabditis Elegans. *npj Aging* **2023**, *9* (1), 20.

(57) Yoshina, S.; Izuhara, L.; Kamatani, N.; Mitani, S. Regulation of Aging by Balancing Mitochondrial Function and Antioxidant Levels. *J Physiol Sci* **2022**, *72* (1), 28.

(58) Greer, E. L.; Oskoui, P. R.; Banko, M. R.; Maniar, J. M.; Gygi, M. P.; Gygi, S. P.; Brunet, A. The Energy Sensor AMP-Activated Protein Kinase Directly Regulates the Mammalian FOXO3 Transcription Factor*. *Journal of Biological Chemistry* **2007**, *282* (41), 30107–30119.

(59) Li, X.-N.; Song, J.; Zhang, L.; LeMaire, S. A.; Hou, X.; Zhang, C.; Coselli, J. S.; Chen, L.; Wang, X. L.; Zhang, Y.; Shen, Y. H. Activation of the AMPK-FOXO3 Pathway Reduces Fatty Acid–Induced Increase in Intracellular Reactive Oxygen Species by Upregulating Thioredoxin. *Diabetes* **2009**, *58* (10), 2246–2257.

(60) Maruyama, A.; Sato, Y.; Nakayama, J.; Murai, J.; Ishikawa, T.; Soga, T.; Makinoshima, H. De Novo Deoxyribonucleotide Biosynthesis Regulates Cell Growth and Tumor Progression in Small-Cell Lung Carcinoma. *Sci Rep* **2021**, *11* (1), 13474.