Thermal Plasticity of Stage Specific Development Time in *Drosophila melanogaster* under Shifting Temperatures

Aradhya Chattopadhyay[1], Rishav Roy[1,2], Payel Biswas[1,3], Shampa M. Ghosh[1]*

1: School of Biotechnology, Kalinga Institute of Industrial Technology (KIIT) Deemed to be University, Bhubaneswar 751024, Odisha, India

2: Huck Institute of Life Sciences, Pennsylvania State University, USA

3: School of Biological Sciences, University of Edinburgh, UK

* **Correspondence:** Shampa M. Ghosh
Email: shampa.ghosh@kiitbiotech.ac.in

Abstract:

Thermal shifts and fluctuations are inherent aspects of the natural environment. How different developmental stages of the fly respond to thermal fluctuations and shifts are not properly understood. In view of this, we studied how durations of different developmental stages are affected by current and previous stage developmental temperatures when they experience upshifts and downshifts of temperature during developmental transitions. The study revealed a complex interaction of current and prior stage temperatures in shaping development time. The egg-stage temperature is found to exert its effect on both larval and pupal stage. However, the impacts of larva-pupa thermal shifts on pupal and overall development time are greater than that of egg-larva thermal shifts. As one would predict, a prior cold temperature extends subsequent pupal duration, while prior warm condition reduces it. Surprisingly, for larval duration, both prior warm and cold condition had the same effect- both led to significant reductions in subsequent larval stage duration. The warm egg temperature had an accelerating effect on larval development as expected. The other shift from cold egg to warm larval condition that also made development faster than constant warm rearing, was possibly caused by a shift from prior sub-optimal to optimal thermal condition used in the experiment. We also found warm larval stage followed by cold pupal stage makes development time shorter than cold larval followed by warm pupal rearing, suggesting a dominating role of the early-stage temperature in determining overall course of development. And lastly, we also observed two subsequent thermal shifts leave a decelerating effect on pupal development, possibly revealing a difficulty in developmental adjustments in the face of frequent thermal changes.

Keywords: Development time, *Drosophila*, thermal plasticity, thermal shift

Introduction

Temperature influences diverse aspects of biology such as development, physiology,

metabolism, behaviour, and life history. Variation in temperature is an inherent aspect of the

natural environment and some organisms are affected more by thermal changes than the

others. Thermal effects are more pronounced in ectotherms as their body closely tracks the

ambient temperature (Addo-Bediako et al., 2000; Angilletta, 2009). Effects of temperature on

organisms are manifested through thermal plasticity of different biological traits. Studies on

thermal plasticity have gained momentum in past few decades as increasing attention is now

being paid on mechanistic and adaptive aspects of plasticity (Trotta et al., 2006; Cooper et al.,

2012; McDonald et al., 2018; Lafuente et al., 2024). However, most studies exploring

thermal plasticity involve investigating organismal performance at a range of constant

temperatures in the laboratory (Trotta et al., 2006; Klepsatel et al., 2019; Lafuente et al.,

2024). While such studies offer insights into understanding plastic and adaptive responses of

organisms to different thermal environments, how temporal changes in temperature during

development and lifetime of an organism influence its biology, is not clearly understood. In

view of this, we were interested to study how shifting temperatures during pre-adult stages

influence temporal coordination of development. We used the ectotherm *Drosophila*

melanogaster for our study.

Drosophila life cycle consists of egg, larva, pupa and the adult. During its developmental

cycle, the four life stages of the fly might encounter dramatically varied surroundings. The

egg stage is immobile and occurs mostly on the surface of rotting fruits; the larval stage is

movable but normally restricted within the fruit; larvae generally wander outside the rotting

fruit and pupariate nearby; and the adult stage is not within a substrate but is completely

mobile (Sokolowski et al., 1986; Powell, 1997). Consequently, the temperature that each of

these four life stages experiences can be dramatically different due to daily temperature

fluctuations, heatwaves, and substrate differences. Previous studies reveal thermal tolerance

and reaction norms may change between life stages, indicating some stages may be more

sensitive to thermal changes than others (e.g. Kingsolver et al., 2011; Austin and Moehring,

2013; Blanckenhorn et al., 2014; van Heerwaarden et al., 2014;). Therefore, we speculate

thermal changes during the course of development may have dissimilar impacts on duration

of different pre-adult stages. Thermal plasticity of development time has been studied

extensively in *Drosophila*, and like majority of ectotherms, cold temperature extends overall

development time of the fly (Davidson and Davidson, 1944; Partridge et al., 1994; Al-Saffar et al., 1995; Ghosh et al., 2013). Developmental changes during embryonic/egg stage are found to be temporally robust across a wide range of temperatures in flies, when tested at various constant temperatures (Chong et al., 2018). Similarly, various stages of pre-adult development like duration of larval instars and pupal stage show proportionate increase with decreasing temperature (Ghosh et al., 2013). The optimal temperature for the model species *Drosophila melanogaster* is 25°C, while 17°C stretches the development by two-fold (Ghosh et al., 2013). All pre-adult developmental stages like egg hatching, larval duration, pupal stage are increased by two-fold at 17°C compared to 25°C (Ghosh et al., 2013). While there is ample data available for thermal plasticity of development time in constant warm *vs.* constant cold conditions, there is little understanding about what fluctuating temperatures do to *Drosophila* development. The question is relevant given in nature most organisms including flies are frequently subjected to changing temperatures (Angilletta, 2009; Garrity et al., 2010). Changes in temperature can manifest through regular diurnal or seasonal fluctuations, or it can also be evident as sudden rise or fall in temperature. Some studies have employed 12- or 24-hour thermal fluctuations to study how those affect flies (Petavy et al., 2001; Manenti et al., 2015, 2021; Tobler et al., 2015), but we know little about how development adjusts itself when there are instances of abrupt thermal shifts during the course of development. In view of this, we were interested to see how thermal shifts during developmental transitions affect subsequent development time of the fly.

We investigated how thermal changes across different developmental stages like egg, larva and pupa shape duration of subsequent developmental stages and overall development time of flies. What kind of carryover effects are observed when developmental stages of flies are shifted from warm to cold and vice versa? If the current rearing condition is cold but previous stage was warm, does the earlier stage temperature show a carryover effect by shortening the duration of the later stage? Conversely, if a fly was initially developing under cold and then it is brought to warm condition, does it take longer than normal to develop due to carryover effect of the previous cold rearing? To address these questions, we subjected individual *Drosophila melanogaster* flies to thermal shifts at developmental transitions, *e.g.* either upon hatching, or pupariation, or at both transitions. We used two alternating temperatures, 25°C and 17°C to upshift or downshift the flies immediately after larval hatching and/or pupariation.

Materials and Methods

Population ancestry

We used a large, outbred laboratory population of *Drosophila melanogaster* named CO-mix
for this study. It is a descendant of ancestral CO-populations (Rose et al., 1984; Chippindale
et al., 1993, 1997). 5 replicates of CO populations were mixed to create the CO-mix
population. This population as well as its ancestors are being maintained at 25˚C. CO-mix
flies are maintained on a 14 days discrete generation cycle, under 24 hours light, and ~70%
relative humidity on standard cornmeal medium. Rearing density is controlled at ~70 eggs
per 6 ml food in a vial. Twenty five such vials are reared every generation for growing the
flies, and upon eclosion, all flies are transferred to a population cage, and the eggs are
collected after 3 days and incubated in vials, to initiate the next generation.

Experimental protocol

Stage specific development time was assayed at different thermal combinations. *Drosophila*
development includes three major developmental transitions, namely egg to larva (hatching),
larva to pupa (pupariation), and pupa to adult (eclosion). For different thermal treatments,
thermal shifts between the two treatment temperatures, namely, 17˚C and 25˚C were
performed either at hatching, or at pupariation, or at both. Total eight thermal combinations
were used for the experiment, as shown in Table 1.

Set	Egg	Larva	Pupa
1	25˚C	25˚C	25˚C
2	17˚C	25˚C	25˚C
3	25˚C	25˚C	17˚C
4	17˚C	25˚C	17˚C
5	17˚C	17˚C	17˚C
6	25˚C	17˚C	17˚C
7	17˚C	17˚C	25˚C
8	25˚C	17˚C	25˚C

Table 1: Schematic of the 8 thermal treatments for stage-specific development time assay

For each thermal treatment, 80 microcentrifuge tubes (MCTs) were used, each containing 300

ml of cornmeal food. A small agar piece containing a single egg was placed in each MCT

within 2 hours of egg lay from the CO-mix populations. In *D. melanogaster*, eggs appear

smooth and swollen when larvae are inside, and after larval hatching, the eggs appear

crumpled. This way hatched vs. unhatched eggs can be distinguished clearly under

microscope. For treatments in which eggs were reared at 25˚C (Treatments 1, 3, 6, 8), eggs

were monitored for larval hatching 12 hours onwards post egg collection, at every 2-hour

intervals. The time of larval hatching was noted down for each MCT. Hatching check was

continued till 26 hours post egg collection, when no new eggs hatched for 3 consecutive

checks. Eggs that failed to hatch and became blackened, were discarded. For treatments 6 and

8, individual tubes were shifted from 25˚C upon hatching, and incubated at 17˚C. For

treatments 1 and 3, MCTs were maintained at 25˚C. Similar protocol was followed for eggs

reared at 17˚C (Treatments 2, 4, 5, 7), and larval hatching was monitored from 32 hours up to

42 hours post egg collection. For treatments 2 and 4 MCTs were shifted from 17˚C upon

hatching, and incubated at 25˚C while for treatments 5 and 7 MCTs were maintained at 17˚C

throughout egg and larval stages.

Hatched larvae were allowed to grow at specific temperatures and subsequently monitored

for pupariation after few days. MCTs were monitored for formation of pupae on inner wall of

the tube or lower side of the tube's lid every 4 hours. Upon pupariation, the time was noted

and tubes were further kept at 25˚C or 17˚C, for treatments that involved no shifts at

pupariation. Checks were continued till no new pupa was observed for 3 successive checks.

Larvae that failed to reach the wandering stage or pupariate were discarded. For treatments

involving shifts at pupariation (Treatments 3, 4, 7, 8), MCTs with pupae were shifted to

specified temperature after noting their pupariation time. MCTs containing pupae were

incubated at specific temperatures to allow metamorphosis to occur. As metamorphosis

ensued over next few days and pupae became dark indicating development of adult flies

within pupal cases, individual tubes were monitored for eclosion. As eclosion started, checks

were continued every 4 hours and time was noted. Eclosion checks were continued till no

new eclosion was observed for 3 successive checks.

Data analysis

For any given MCT, egg duration was calculated by subtracting the midpoint of the 2 hour-

long egg collection window from time of larval hatching. Larval duration in a given tube was

calculated by subtracting hatching time from the time of pupariation. Pupal duration was

calculated by subtracting pupariation time from the time of eclosion. Total egg to adult

development time was also noted for individual flies in all 8 thermal treatments.

ANOVA was performed to compare egg to adult development time across all 8 thermal

treatments with treatment taken as fixed factor. For comparing egg duration at 17˚C (N =

151) and 25˚C (N = 116), one way analysis of variance (ANOVA) was performed on

individual egg duration data. Thermal treatment was treated as fixed factor. Compared to

25˚C, 17˚C increases duration of developmental stages two-fold in *D. melanogaster*, and

therefore it is obvious that an ambient temperature of 25˚C *vs.* 17˚C would significantly

affect the duration of any given developmental stage (Ghosh et al., 2013). In this study we

rather investigated when the temperature is same for a given developmental stage (*e.g.*

larval), how it is affected by contrasting temperature in the preceding stage/s. Therefore, two

separate ANOVAs were performed for larval duration, one for comparing treatments in

which larval stage was reared at 25˚C (25-25 vs. 17-25), another for treatments in which

larvae were reared at 17˚C (17-17 vs. 25-17). Treatment was taken as a fixed factor, and

ANOVA was performed for replicate MCT data points. Like larval duration, for pupal

duration two separate ANOVAs were performed, one for comparing all thermal treatments in

which the ambient pupal rearing temperature was 25˚C, another for comparing duration of all

treatments in which pupae were reared at 17˚C. For one pupal duration ANOVA, four groups

were compared (25-25-25 vs. 17-25-25 vs. 25-17-25 vs. 17-17-25). In the other pupal

duration ANOVA, comparisons were for 17-17-17 vs. 25-17-17 vs.17-25-17 vs. 25-25-17.

Post hoc comparisons were done using Tukey's honest significant difference (HSD) test for

analysis of pupal duration and total development time. Data analyses were carried out on

JMP Pro 17. Homoscedasticity of variance among different groups was verified by Levene's

test performed on R.

Results and Discussion

Carryover effect of prior stage temperature on Development is more pronounced when

current temperature is 17˚C as opposed to 25˚C

The effect of thermal treatment on total or egg to adult development time was highly

significant (p < 0.0001). Post hoc comparisons revealed total development time was

significantly different across all 8 treatments (Fig. 1). Effect of egg and larval rearing

temperatures on duration of subsequent developmental stages revealed an array of interesting

trends, detailed in following paragraphs. It is important to note that the effect of previous

stages' temperature on subsequent development time was more pronounced if the later stage

temperature was 17°C instead of 25°C. Given development is longer at 17°C, any change

caused by thermal environments of preceding stages got magnified at this temperature, while

the effects were small at 25°C because of shorter duration of development at this temperature.

Shift at egg-larva transition impacts total development time but the effect is small

Total development time was longest at 17-17-17 (434 h), followed by 25-17-17 (384 h). At

25-25-25 it was the shortest (212 h), followed by 17-25-25 (224 h) (Fig. 1, H *vs.* G). This

suggests though a shift upon egg-larva transition (shift 1) significantly affects total egg to

adult development time, the effect is small such that overall development time in these

treatments remain closest to the development time observed at constant temperatures.

Figure 1: Mean egg to adult development time at different thermal treatments. The error bars

represent standard error across replicate MCTs.

Thermal plasticity of egg duration

Egg duration measured by egg to larva hatching time was significantly affected by rearing

temperature ($p < 0.0001$). At 25°C, mean egg hatching time was 21 hours, while at 17°C, eggs

took 37 hours on average to hatch (Fig. 2.1). This result corroborates findings reported by

earlier studies (Powsner, 1935; Chong et al., 2018).

Figure 2: 2.1: Mean egg duration (larval hatching time) at 25˚C and 17˚C. 2.2 Mean larval

duration at different thermal treatments. The error bars in both plots represent standard error

across replicate MCTs.

Both upshift and downshift of temperature at egg-larva transition accelerate subsequent

development

When eggs are reared at 25˚C, average duration of egg stage is 16 hours less compared to

17˚C (Fig. 2.1). The egg rearing temperature of 25˚C not only makes the egg stage develop

fast, it also accelerates successive development. This is evident from the finding that the

overall shortening of development time observed at 25-17-17 compared to 17-17-17 is 50

hours on average (Fig. 1, B *vs*. A). This clearly shows that the warm egg rearing temperature

has a spillover effect, and it accelerates subsequent development of the fly. ANOVA revealed

significant impact of egg temperature on larval duration ($p = 0.0017$) when larvae were

reared at 17˚C. Larval duration at 17-17 was 194 h, while at 25-17, it was 184 h (Fig. 2.2),

indicating a significant reduction caused by prior egg stage temperature.

Our expectation was in case of the upshift, *i. e*. at 17-25-25, larval and pupal developments

would get slower compared to constant developmental temperature of 25˚C, but the results

did not meet this expectation. 17˚C hatched larvae upon being shifted to 25˚C (17-25) instead

showed a small but significant reduction of larval duration (90 h) compared to 25-25 larvae
(94 h) (Fig. 2.2), and no change in pupal duration (98 h) at 17-25-25 compared to that at 25-
25-25 (97 h) (Fig. 3, b *vs*. a). We had speculated a prior warm temperature would leave an
accelerating after-effect while a prior cold temperature would do the opposite. However,
finding from this experiment suggests it could be more complex than that. In a variable
thermal environment, developmental pace of a pre-adult life stage may depend upon previous
stage temperature, that is, whether it experienced warmer or colder environment before. In
addition, the pace may also depend upon whether the shift was from a sub-optimal to optimal
temperature as development may overcome the carry-over decelerating effect of a previous
cold temperature and rather may gain some momentum because of sudden attainment of
optimal temperature after a sub-optimal one. It is possible that the enzymes driving
development at cold and sub-optimal temperature like 17˚C gains momentum upon abrupt
shift to the optimal temperature, *i. e.* 25˚C that results in speeding up of larval development.
Future research delving into the effect of thermal shifts on efficiency of developmental
enzymes may shed more light on this.

Larva-pupa thermal shift: cold larval-warm pupal condition make development longer than
warm larval -cold pupal condition

Development time at 17-17-25 was longer (333 h) than that at 17-25-17 (319 h) (Fig. 1, C *vs*.
D). Similarly, flies took significantly longer time to develop at 25-17-25 (304 h), compared to
that at 25-25-17 (288 h) (Fig. 1, E *vs*. F). This shows colder larval stage followed by warm
pupal stage makes overall development longer, compared to warm larval stage followed by
cold pupal stage. This is not surprising. Firstly, cold larval rearing makes larval stage long,
and it also leaves a carryover effect on pupal stage such that pupal development at warm
becomes significantly extended compared to constant warm rearing. The converse also holds
true as warm larval rearing not only makes larval development fast but also makes pupal
development at cold significantly faster compared to constant cold rearing. Therefore 17˚C
larval-25˚C pupal development is longer compared to 25˚C larval-17˚C pupal rearing.

Figure 3: Mean pupal duration at different thermal treatments. The error bars represent standard error across replicate MCTs.

Effects of egg and larval temperatures on pupal duration at 25˚C

When pupae were reared at 25˚C, pupal duration was impacted by the thermal combination experienced during egg and larval stages (p < 0.0001). However, at 25-25-25 and 17-25-25, pupal durations were similar, *i. e.* 97 h, and 98 h respectively (Fig. 3, a *vs*. b), indicating egg stage temperature did not have any carryover effect up to the pupal stage when larvae and pupae both experienced 25˚C. At 25-17-25, nonetheless, pupal duration was 100 h (Fig. 3, c), which was significantly longer than the same at 25-25-25, *i. e.* 97 h, showing cold larval temperature has carryover effect on pupal development. At 17-17-25, pupal duration was 104 h (Fig. 3, d), which was significantly longer than the same at three other treatments, indicating combined effect of cold temperature during egg and larval stages have stronger carryover effect in extending pupal duration at 25˚C.

Effects of egg and larval temperatures on pupal duration at 17˚C

Pupal duration at 17˚C too was significantly affected by the temperatures experienced during
egg and larval stages (p < 0.0001). Post hoc comparisons revealed pupal duration at 17˚C was
significantly different across all four different thermal pre-treatments. When eggs were reared
at 25˚C, and larvae were downshifted to 17˚C upon hatching and reared at the same
temperature till eclosion (25-17-17), both larval (184 h) and pupal developments (178 h) were
significantly accelerated compared to development at a constant temperature of 17˚C,
revealing a strong carryover effect of warm egg temperature on both larval and pupal stages
developing at 17˚C (Fig. 3, g). Moreover, pupae took 204 hours to develop into adults at 17-
17-17, which was the longest (Fig. 3, e). At 25-25-17, it was the shortest, as it took 172 hours
to complete pupal metamorphosis (Fig. 3, h). The combined effect of warm temperatures
during egg and larval stages in accelerating pupal development was clearly evident from this.

Two subsequent thermal shifts at egg-larva and larva-pupa make subsequent pupal
development slow

The effects of the double thermal shifts on development time were remarkable. At both 25-
17-25 and 17-25-17, pupal metamorphosis took substantially long. The lengths of pupal stage
in these treatments were second only to treatments in which egg-larva rearing temperature
was 17-17. Among treatments involving pupal development at 25˚C, pupal duration at 25-17-
25 (100 h) was faster only than 17-17-25 (104 h), and among treatments involving pupal
development at 17˚C pupal duration at 17-25-17 (194 h) was slower only compared to17-17-
17 (204 h) (Fig. 3, c *vs*. d, f *vs*. e). Pupal duration at 25-17-25 significantly increased to 100 h,
from 97 h at 25-25-25, revealing a carryover effect of colder larval rearing environment
(17˚C) on pupal development (Fig. 3, c *vs*. a).

It was a little perplexing that pupal duration in the other double shift treatment, the one for
17˚C pupae, *i. e.* 17-25-17 was 194 h, which was significantly slower than 178 h, observed at
25-17-17 (Fig. 3, f *vs*. g). Intuitively, one would expect pupal stage to have greater impact of
larval rearing temperature than that of egg temperature, because (a) larval stage immediately
precedes the pupal stage, and (b) larval stage is much longer than the egg stage. However, the
finding indicates warm temperature during egg rather than larval stage caused bigger
reduction of pupal duration in this case. In this regard, we speculate two abrupt thermal shifts
at subsequent developmental transitions possibly make developmental adjustments more
challenging such that metamorphosis remains slow, and therefore carryover effect of 25˚C
larval temperature on pupal development is not found to be as large as expected.

Pupal duration is affected the most when shift is done at larva-pupa transition

For pupae developing at 17˚C, metamorphosis took the shortest time (172 h) in those which

were reared at 25˚C during egg and larval stages (25-25-17) (Fig. 3, h). Similarly, for pupae

reared at 25˚C, metamorphosis was the longest (104 h) for those reared at 17˚C through egg

and larval stages (17-17-25) (Fig. 3, d). This indicates pupal duration at warm temperature is

extended the most when both egg and larval rearing were in cold, and conversely, it is

accelerated the most when both previous stages were warm.

Conclusion

To summarize, this study reveals several exciting findings about thermal plasticity of

development time in *Drosophila melanogaster* under variable thermal conditions. First, it

shows thermal effects of preceding developmental stages on later stages are more

conspicuous when the later stage temperature is cold. As duration of development is longer in

cold, it provides a finer resolution of developmental duration, revealing greater plastic

changes caused by previous thermal conditions. Second, thermal shift early during

development at egg-larva transition affects subsequent developmental pace, but it has

relatively smaller impact on total egg to adult development time compared to thermal shift

done at a later stage, *i. e.* larva-pupa transition. This is particularly evident in pupal duration

as it is affected more by larva-pupa than egg-larva thermal shift. Third, a downshift from

warm egg stage to colder temperature predictably accelerates subsequent larval and pupal

development compared to constant cold, due to carry-over effect of prior warm condition.

However, a striking finding from this study is an upshift from cold egg temperature to warm

conditions also speeds up larval development making it more rapid than that at constant

warm, instead of showing a carry-over effect of prior slow pace at cold. We speculate this is a

result of gain in momentum presumably due to increased efficiency of developmental

enzymes, caused by a shift from sub-optimal to optimal conditions. However, for pupal

duration, effect of previous stage temperatures was as expected. Prior stage cold temperature

led to lengthening of pupal duration while warm egg or larval temperature led to its

reduction. Fourth, cold larval-warm pupal rearing extends development time more than

warm-larval-cold pupal rearing, as thermal condition of the earlier stage, that is, larva, and

the carry-over effect of that stage on the later pupal stage, play dominant roles in shaping

overall development time. Last, but not the least, two subsequent thermal shifts at egg-larva

and larva-pupa transitions seem to make developmental adjustments challenging, such that

pupal development remains slow under such conditions. To conclude, this study demonstrates

that thermal plasticity of developmental durations in *Drosophila* under shifting temperatures

is a complex phenomenon influenced by interaction between previous and ongoing thermal

conditions. It seems to depend on whether it involves an upshift *vs.* downshift of temperature,

timing and number of thermal shifts, optimality of the temperatures involved and possibly

developmental adjustments. This study paves the way for more research delving into roles of

gene expression and enzyme activities in coordinating development in thermally variable

environments.

Author contributions

SG conceived the idea, designed the experiment, analysed and interpreted the data, wrote the

manuscript. AC designed and performed the experiments, analysed data, prepared figures and

co-wrote the manuscript. RR designed and performed the experiments and analysed data. PB

designed and performed the experiments and collected data. All authors reviewed the

manuscript.

Conflict of interest statement

We declare we have no competing interests

Acknowledgements

We thank Adarsh Tripathi for helping with the experiments, Ranjit Pradhan and Manas

Ranjan Mallik for general help in the laboratory.

Funding

This work was supported by a fellowship to SG under Department of Science & Technology

Government of India, DST Women Scientist A scheme, SR/WOS-A/LS-1179/2015(G), and a

grant from Science and Engineering Research Board (DST-SERB), Government of India,

under start up research grant, SRG/2020/001573.

References

Addo-Bediako, A., Chown, S. L., and Gaston, K. J. (2000). Thermal tolerance, climatic
variability and latitude. *Proc R Soc Lond B Biol Sci* 267, 739–745. doi:
10.1098/RSPB.2000.1065

Al-Saffar, Z. Y., Grainger, J. N. R., and Aldrich, J. (1995). Influence of constant and changing
temperature and humidity on the development and survival of the eggs and pupae of
Drosophila melanogaster (Meigen). *J Therm Biol* 20, 389–397. doi: 10.1016/0306-
4565(94)00075-T

Angilletta, M. J. (2009). Thermal Adaptation: A Theoretical and Empirical Synthesis.
Thermal Adaptation: A Theoretical and Empirical Synthesis, 1–302. doi:
10.1093/ACPROF:OSO/9780198570875.001.1

Austin, C. J., and Moehring, A. J. (2013). Optimal temperature range of a plastic species,
Drosophila simulans. *Journal of Animal Ecology* 82, 663–672. doi: 10.1111/1365-
2656.12041

Blanckenhorn, W. U., Gautier, R., Nick, M., Puniamoorthy, N., and Schäfer, M. A. (2014).
Stage- and sex-specific heat tolerance in the yellow dung fly Scathophaga stercoraria. *J
Therm Biol* 46, 1–9. doi: 10.1016/J.JTHERBIO.2014.09.007

Chippindale, A. K., Leroi, A. M., Kim, S. B., and Rose, M. R. (1993). Phenotypic plasticity
and selection in Drosophila life-history evolution. I. Nutrition and the cost of
reproduction. *J Evol Biol* 6, 171–193. doi: 10.1046/J.1420-9101.1993.6020171.X

Chippindale, A. K., Leroi, A. M., Saing, H., Borash, D. J., and Rose, M. R. (1997).
Phenotypic plasticity and selection in Drosophila life history evolution. 2. Diet, mates
and the cost of reproduction. *J Evol Biol* 10, 269–293. doi: 10.1046/J.1420-
9101.1997.10030269.X

Chong, J., Amourda, C., and Saunders, T. E. (2018). Temporal development of Drosophila
embryos is highly robust across a wide temperature range. *J R Soc Interface* 15. doi:
10.1098/RSIF.2018.0304

Cooper, B. S., Tharp, J. M., Jernberg, I. I., and Angilletta, M. J. (2012). Developmental
plasticity of thermal tolerances in temperate and subtropical populations of Drosophila
melanogaster. *J Therm Biol* 37, 211–216. doi: 10.1016/J.JTHERBIO.2012.01.001

Davidson, A. J., and Davidson, B. J. B. (1944). On the Relationship between Temperature and
Rate of Development of Insects at constant Temperatures. *Journal of Animal Ecology*
13, 26. doi: 10.2307/1326

Garrity, P. A., Goodman, M. B., Samuel, A. D., and Sengupta, P. (2010). Running hot and
cold: behavioral strategies, neural circuits, and the molecular machinery for thermotaxis
in C. elegans and Drosophila. *Genes Dev* 24, 2365–2382. doi: 10.1101/GAD.1953710

Ghosh, S. M., Testa, N. D., and Shingleton, A. W. (2013). Temperature-size rule is mediated
by thermal plasticity of critical size in Drosophila melanogaster. *Proceedings of the
Royal Society B: Biological Sciences* 280. doi: 10.1098/RSPB.2013.0174

Kingsolver, J. G., Arthur Woods, H., Buckley, L. B., Potter, K. A., MacLean, H. J., and
Higgins, J. K. (2011). Complex Life Cycles and the Responses of Insects to Climate
Change. *Integr Comp Biol* 51, 719–732. doi: 10.1093/ICB/ICR015

Klepsatel, P., Girish, T. N., Dircksen, H., and Gáliková, M. (2019). Reproductive fitness of
Drosophila is maximised by optimal developmental temperature. *J Exp Biol* 222. doi:
10.1242/JEB.202184

Lafuente, E., Duneau, D., and Beldade, P. (2024). Genetic basis of variation in thermal
developmental plasticity for Drosophila melanogaster body pigmentation. *Mol Ecol* 33.
doi: 10.1111/MEC.17294

Manenti, T., Kjærsgaard, A., Schou, T. M., Pertoldi, C., Loeschcke, V., and Moghadam, N. N.
(2021). Responses to developmental temperature fluctuation in life history traits of five
drosophila species (Diptera: Drosophilidae) from different thermal niches. *Insects* 12.
doi: 10.3390/INSECTS12100925/S1

Manenti, T., Loeschcke, V., Moghadam, N. N., and Sørensen, J. G. (2015). Phenotypic
plasticity is not affected by experimental evolution in constant, predictable or
unpredictable fluctuating thermal environments. *J Evol Biol* 28, 2078–2087. doi:
10.1111/JEB.12735

McDonald, J. M. C., Ghosh, S. M., Gascoigne, S. J. L., and Shingleton, A. W. (2018).
Plasticity through canalization: The contrasting effect of temperature on trait size and
growth in Drosophila. *Front Cell Dev Biol* 6, 423798. doi:
10.3389/FCELL.2018.00156/BIBTEX

Partridge, L., Barrie, B., Fowler, K., and French, V. (1994). Thermal evolution of pre-adult
life history traits in Drosophila melanogaster. *J Evol Biol* 7, 645–663. doi:
10.1046/J.1420-9101.1994.7060645.X

Petavy, G., David, J. R., Gibert, P., and Moreteau, B. (2001). Viability and rate of
development at different temperatures in Drosophila: a comparison of constant and
alternating thermal regimes. *J Therm Biol* 26, 29–39. doi: 10.1016/S0306-
4565(00)00022-X

Powell, J. R. (1997). Progress and Prospects in Evolutionary Biology: The Drosophila Model.
Progress and Prospects in Evolutionary Biology. doi:
10.1093/OSO/9780195076912.001.0001

Powsner, L. (1935). The Effects of Temperature on the Durations of the Developmental
Stages of Drosophila melanogaster. *https://doi.org/10.1086/physzool.8.4.30151263* 8,
474–520. doi: 10.1086/PHYSZOOL.8.4.30151263

Rose, M. R., Dorey, M. L., Coyle, A. M., and Service, P. M. (2011). The morphology of
postponed senescence in Drosophila melanogaster. *https://doi.org/10.1139/z84-230* 62,
1576–1580. doi: 10.1139/Z84-230

Tobler, R., Hermisson, J., and Schlötterer, C. (2015). Parallel trait adaptation across opposing
thermal environments in experimental Drosophila melanogaster populations. *Evolution
(N Y)* 69, 1745–1759. doi: 10.1111/EVO.12705

Trotta, V., Calboli, F. C. F., Ziosi, M., Guerra, D., Pezzoli, M. C., David, J. R., et al. (2006).
Thermal plasticity in Drosophila melanogaster: A comparison of geographic populations.
BMC Evol Biol 6, 1–13. doi: 10.1186/1471-2148-6-67/TABLES/4

van Heerwaarden, B., Lee, R. F. H., Overgaard, J., and Sgrò, C. M. (2014). No patterns in
thermal plasticity along a latitudinal gradient in Drosophila simulans from eastern
Australia. *J Evol Biol* 27, 2541–2553. doi: 10.1111/JEB.12510

406