Rapid riparian ecosystem recovery in low-latitudinal North China following the end-Permian mass extinction

Wenwei Guo[1, 2], Li Tian[2*], Daoliang Chu[2], Wenchao Shu[2], Michael J. Benton[3], Jun Liu[4], Jinnan Tong[2]

[1]College of Culture and Tourism, Zhangzhou Institute of Technology, Zhangzhou 363000, People's Republic of China; [2]State Key Laboratory of Biogeology and Environmental Geology, China University of Geosciences, Wuhan, 430074, People's Republic of China; [3]School of Earth Sciences, University of Bristol, BS8 1RJ, UK; [4]Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing 100044, People's Republic of China

*Author for Correspondence: Li Tian, Email: tianlibgeg@cug.edu.cn

Abstract

The greatest mass extinction at the end-Permian, 252 million years ago (Ma), led to a tropical dead zone on land and sea. The speed of recovery of life has been debated, whether fast or slow, and terrestrial ecosystems are much less understood than marine. Here, we show fast reestablished riparian ecosystems in low-latitude North China within as little as ~2 million years (Myr) after the end-Permian extinction. The initial ichnoassemblages in shallow lacustrine and fluvial facies of late Smithian age are monospecific, devoid of infaunalization, with apparent size reduction. In the following Spathian, newly identified medium-sized carnivores, plant stems, root traces, coupled with improved ichnological criteria and significantly increased infaunalization, suggesting a relatively complex, multi-level trophic structured riverain ecosystem had been rebuilt. Specifically, burrowing behavior had re-emerged as a key life strategy not only to minimize stressful climatic conditions, but possible to escape predation.

Introduction

The end-Permian mass extinction (EPME) is one of the most devastating bio-crises in the history of life, destroying both marine and terrestrial ecosystems (Wignall, 2015; Fan et al., 2020). A key feature of the EPME is the clearing of life from tropical ecosystems (e.g., Sun et al., 2012). The environmental extremes, especially lethal heat, have largely flattened the tropical biodiversity peak, and marine animals were forced to migrated poleward or into deeper water (Liu et al., 2020; Song et al., 2020). Such effects are more brutal on land. A tropical "tetrapod gap", spanning between 15°N and ~31°S, prevailed through the Early Triassic, or at least during particular interval of intense global warming (Bernardi et al., 2018; Allen et al., 2020; Romano et al., 2020; Liu et al., 2022).

It has been demonstrated that long-term environmental perturbations after the EPME resulted in the delayed recovery until to the Middle Triassic in the sea (Chen and Benton, 2012), although several fast evolvers would bounce back fast before being killed by further repeat hyperthermal events through the Early Triassic, as evidenced by the Guiyang Biota found ~1 Myr after the EPME (Dai et al., 2023). However, the post-extinction recovery on land remained largely unclear. The taxonomic diversity of vertebrates may have re-flourished soon after the extinction in European Russia (Tverdokhlebov et al., 2003), but it apparently took longer, until the latest Early Triassic, in the Central European Basin (e.g., Mujal et al., 2017). Model results of the tetrapod-dominated paleocommunities from the Karoo also displayed a short lived, unstable ecosystem during the Early Triassic *Lystrosaurus* Assemblage Zone, before being replaced by a globally stable ecological structure established in the Middle Triassic (Roopnarinev et al., 2019; Viglietti et al., 2022). It seemed that plants showed a quick return in Australia (Vajda and Kear, 2024), yet reorganization of floral communities were hindered by repeated climatic stressors, such as the Smithian–Spathian warming event (Mays et al., 2020; Vajda et al., 2020). Likewise, the initial construction of the mesophytic flora was in the earliest Middle Triassic in North China (Shu et al., 2022).

Other body fossils, especially non-marine invertebrates, are relatively scarce in the post-extinction interval. The earliest Middle Triassic riverine community, consisting of insects, rare fishes and trace fossils in equatorial western peri-Tethys (Baucon et al., 2014; Mujal et al., 2020; Matamales-Andreu et al., 2021) and the Anisian–Ladinian deep lake biota, comprising of diverse insects, fishes, fish coprolites and plants in tropical North China (Zhao et al., 2020), were thought representative of the recovered terrestrial ecosystems after the EPME. However, recently studies revealed that the biodiversity would be not as low as expected in North China from the ichnological

point of view, relatively diversified trace fossils have been found during the late Early Triassic with

moderate bioturbations (Guo et al., 2019; Xing et al., 2021; Zheng et al., 2021). Here, we report the

unexpectedly fast recovered ecosystem from the tropical dead zone after the EPME, based on

compiled data of vertebrates, invertebrate trace fossils, and plant remains from Lower Triassic

successions and outcrops in North China, to show how animal behaved to survive in harsh conditions

and finally recovered from the mass extinction event.

Materials and Methods

Abundant trace fossils have been identified from the uppermost Permian–Lower Triassic of the

Shichuanhe, Dayulin, Liulin sections and Hongyatou, Tuncun, Mafang outcrops in the Central North

China Basin (Fig. 1). During the early Mesozoic, the North China Block was located at about 20°N

based on paleomagnetic reconstruction (Huang et al., 2018; Guo et al., 2022). The Permo-Triassic

strata, typically terrestrial red-beds, comprising the Sunjiagou, Liujiagou, Heshanggou (HSG) and

Ermaying formations in ascending order, display varied depositional environments from fluvial to

lacustrine facies (See Figure S1 and Table S1 in the Supporting Information). The Sunjiagou

Formation mainly consists of red massive siltstone and intercalated sandstone of varied thickness,

and paleosols and trace fossils were locally developed, demonstrating floodplain to lakeshore facies

(Ji et al., 2022; Yu et al., 2022). The Liujiagou Formation is characterized by thick cross-bedded

sandstone and intraclast conglomerate. Lenticular sand bodies and erosive bases of sandstones were

common in the lower part, indicating fluvial channel facies under braided river systems (Zhu et al.,

2019; Ji et al., 2022). Interlayered thick siltstone of lakeshore facies increased upwards, accompanied

by weak biological activities. The HSG is dominated by massive siltstones, with multi-layered

paleosols, diverse trace fossils and some plant remains, belonging to alluvial plain and lakeshore facies, with periodic aerial exposure (Zhu et al., 2020; Ji et al., 2022).

Recently, a U-Pb CA-ID-TIMS age calibrated magnetostratigraphy from the Shichuanhe section provided a basic geochronological timescale for the fossil-poor strata in North China (Guo et al., 2022; Figure S2). Accordingly, the Permian–Triassic Boundary is constrained in the upper part of the Sunjiagou Formation based on the age of 252.21 $\pm$ 0.15 Ma, with the base of Smithian and Spathian being roughly located in the lower and upper Liujiagou Formation respectively. However, loss of several magnetozones due to regional hiatus made it difficult to precisely place the Lower–Middle Triassic Boundary, which was tentatively put at the lithological contact of HSG and the overlying Ermaying Formation (Guo et al., 2022). The 245–247 Ma ages from tuff layers at the base of the Ermaying Formation roughly supported this correlation (Zhu et al., 2022).

In order to discriminate the recovery stages, several ichno-ecological criteria were incorporated. Both bioturbation index (BI) and ichnofabric index (ii), which have been critically reviewed by Luo et al. (2020), are used to quantify bioturbation intensity. Values of ii and BI, ranging from 1–6 and 0–6, respectively, co-indicate the gradual increase of biotic disturbance from no bioturbation to total homogenization of sediments. Ichnodiversity represents the number of ichnotaxa, but it is not strictly equivalent to biodiversity, as a certain trace can be made by different animals and multiple trace types can originate from a single taxon (Luo et al., 2019). Ichnodisparity emphasizes the variability of architectural designs of trace fossils (Buatois et al., 2017). Therefore, both ichnodiversity and ichnodisparity are employed to assess the behavioural responses of animals and the stage of infaunal biotic recovery (Table S2–4). Size and penetration depth were also measured in place that can represent average level of each burrow. Tiering, referring to the life position of an animal vertically

in the sediment, is divided into surficial, semi-infaunal (0–0.5 cm), shallow (0.5–6 cm), intermediate

(6–12 cm) and deep infaunal tiers (> 12 cm), adopted from Minter et al. (2017).

Thin sections of plant stem specimens were prepared to examine vertical and cross-sectional

microstructures. Meanwhile, Micro-CT scanning (SkyScan 1172 X-ray; State Key Laboratory of

Biogeology and Environmental Geology) was employed to reconstruct the internal structures of

stems.

Results and Discussion

Infaunal crisis and living strategy after the EPME in North China

An infaunal crisis, marked by the disappearance of the moderately diversified pre-extinction

ichnofauna and the absence of biogenic structures, was identified during the late Changhsingian–

early Smithian. The latest Permian floodplain facies mainly occupied by shallow–intermediate tiers

of freely motile non-specialized deposit-feeding animals (Figure 1), akin to the Paleozoic suites

reported before (Minter et al., 2017). Both the ichnodiversity and ichnodisparity declined abruptly in

the middle Sunjiagou Formation (near the 252.21 Ma aged tuff layer), following a prolong

non-bioturbated interval in North China (Guo et al., 2019, 2022; Xing et al., 2021). The infaunal

crisis was contemporary with the extirpation of the pareiasaur fauna (*Shihtienfenia*) and deforestation

of the youngest Palaeozoic *Ullmannia-Pseudovoltzia-Germaropteris* assemblage in North China,

representing the EPME on land (Liu et al., 2022; Shu et al., 2022).

The early resurged ichnofaunal in the upper Liujiagou Formation, about late Smithian in age,

were short-lived. This lakeshore ichnofauna, including seven ichnogenera of six ichnodisparities,

were characterized by surficial and semi-infaunal tiered simple burrows or trails, with rare trackways

and root traces, which weakly disturbed the sediments (Figure S3). Dwarfism in trace makers is observed from all ichnogenera, with burrow sizes reduced from 4.06 mm before the EPME (n = 779) to 2.06 mm (n = 341) in the late Smithian (Figure 4 and Figure S7), or single ichnogenus, such as *Kouphichnim* (Shu et al., 2018). The depauperate ichnofauna of the late Smithian were monospecific, representing initial recolonization of empty niches by opportunists, but the coeval thrived microbial mats indicated harsh environments, which might have inhibited the recovery of freshwater ecosystems (Tu et al., 2016; Chu et al., 2017; Mays et al., 2021).

Abundant trace fossils were identified in the Spathian HSG Formation, comprising 16 ichnogenera of nine ichnodisparities, and two informally designated types (Figure 2 and Figure S4–5). The morphologically complex ichnofauna were dominated by actively filled burrows, with few arthropod trackways, which were probably produced by decapod crustaceous, myriapods and insects (Table S5). The lakeshore and alluvial plain facies were occupied multi-tiered traces, including surficial trackways (e.g., *Diplichnites*; *Kouphichnium*), semi-infaunal (e.g., *Helminthoidichnites*; *Gordia*), shallow (e.g., *Palaeophycus*; *Scoyenia*), intermediate (e.g., *Taenidium*) and deep tiers (*Camborygma*; *Skolithos*; Figure 1c). Trace producers colonized varied ecospace and shallower tiers are generally crosscut by deeper penetrative burrows (Figure 2a), resulting in moderately to substantially bioturbations, with ii 2–3 and BI 3–4 at most layers. In several horizons, the uppermost few centimeters of sediments are totally obliterated, mostly by the activity of deposit feeding animals. However, distal terminal fan facies are weakly reworked by simple traces such as *Skolithos* and *Palaeophycus* or root traces. Additionally, average burrow sizes of all ichnogenera also increased to 3.9 mm in the HSG (n = 2241, Figure 4 and Figure S7–8).

Several large burrows in the Spthian indicate the occurrence of advanced ecosystem engineers.

Camborygma litonomos were found in the basal HSG of the Shichuanhe section and outcrop in Hongyatou, co-occurring with *in situ* preserved *Neocalamites* plant fossils, and other traces, such as *Diplichnites*, *Monomorphichnus* and *Skolithos* (Figure S6). Surfaces of *C. litonomos* were occasionally intertwined with rhizotubules (Figure 2l), suggesting that those plants could be important constituents in the diet of crayfish or that the crayfishes may use the roots to hide among. In addition, another large burrow was identified at the Liulin section. The sub-horizontal unbranched burrow displayed probable longitudinal scratch marks on the external surface, but poor preservation and limited specimens hindered a definite designation (Figure S5). ?*Beaconites*, found in the lower part of the HSG at Mafang, are characterized by meniscate structures (Figure 2o–p and Figure S5C–E), akin to those of large *Beaconites* from breccia facies in the Devonian red sandstone of Britain (Brück, 1987). Although the biological nature of these large burrows cannot be confirmed, their activities increased biogenic reworking of sediment and soils, improved geochemical recycling and ecosystem complexity in the Spathian, implying key roles in ecosystem functioning.

Although climatic and environmental conditions in the early Spathian were still not cool and wet enough for thriving and abundant life, a fossorial strategy would have been useful for continental animals to avoid heat and aridity. Midday temperatures > 35°C, as occurred during peaks of global warming at the EPME and at points through the Early Triassic, cannot be tolerated for long by terrestrial (or aquatic) animals (Benton, 2018; Liu et al., 2022). The increase in tetrapod burrow abundance and complexity in the Lower Triassic suggest that a fossorial lifestyle allowed tetrapods to endure harsh post-extinction environmental conditions (Marchettti et al., 2024). Likewise, we envisaged that infaunalization could also have been a vital strategy for invertebrates to survive and thrive. More intensively occupied ecospace in the late Spathian, characterized by increased

burrowing and complicated crosscutting relationships among ichnogenera, may indicate an adaptive

response to heightened predation pressure or competition for available resources.

Fast recovered terrestrial ecosystem in low-latitudinal region

Newly discovered tetrapods from the HSG provide crucial insight into the post-EPME

ecosystem. Historically, vertebrates found in this lithological unit were exclusively from its middle–

upper portions (Li et al., 2008; Fig. 3). The presence of Archosauromorpha (e.g., *Fugusuchus*; Fig.

3c) and Procolophonomorpha (e.g., *Eumetabolodon*; Fig. 3d) could reflect their tolerance to hot and

arid condition, while the initial diversification of archosauromorphs in the Olenekian was interpreted

as a response to empty ecological space after the EPME (McLoughlin et al., 2020; Romano et al.,

2020). Herein, a cluster of tetrapod skeletons, includes a few articulated bones, were found near the

base of HSG (early Spathian; Figure 3i). The body trunk lengths are estimated at 30–40 cm for the

well exposed vertebrates, and the postcranial skeleton suggests a carnivorous feeding strategy.

Although the specimens are not yet fully prepared for taxonomic description, they clearly show the

existence of tetrapod at this level.

Plants, as key components of the ecosystem, were patchy distributed in the Early Triassic (Shu

et al., 2022). However, root traces (rhizoliths) are quite abundant, especially at the Dayulin section

(Figure 2d and Figure S6). The different styles of preservation and various morphologies of

rhizoliths provide information about the moderately to relatively well-drained red paleosols in

alluvial plain facies in North China (Kraus and Hasiotis, 2006). Specifically, *in situ* preserved

vertical *Neocalamites* stems and *Pleuromeia* found in the lower HSG of the Shichuanhe and

Hongyatou sections, both emerged in tandem with crayfish trace fossils and other burrows.

Delayed terrestrial recovery was proposed based on the Tongchuan fauna from North China,

which consisted of diverse insects, ostracods, fishes, etc, signalling a fully recovered deep lake ecosystem in the middle Triassic, i.e., ~8–12 Myr after the EPME (Zheng et al., 2018; Zhao et al., 2020). However, compiled paleontological data herein show the reconstruction of ecosystem on land in tropical region was as early as in the Spathian, ~2 Myr after the EPME (Figure S9). The enhanced floral coverages, attributed to different types of roots and plant fossils, have great impacts to the initiation of post-EPME ecosystem, especially cooccurred stems and trace fossils, which suggest that riverain realms may be refugia for the survival and evolution of the Spahtian biota. Increased abundance and quantity of faunal communities can be inferred from ichnodiversity, with possible candidate of trace-producers including limuloids, crayfishes, spinicaudatan, insects and even small-sized vertebrates (Table S5), and moderate bioturbations made by high-density resting traces, respectively. Coeval tetrapods, despite rare, do show the existence of carnivores and further improved the local ecological structures. Reconstruction of freshwater ecosystems in the Spathian was also facilitated by amelioration of the climate (Fig. 4). Paleosol-based paleoclimatic reconstructions suggest that precipitation was ~520–680 mm/y in the late Spathian, with pCO_2 estimated from paleosols at 1523 ± 417 ppm (Joachimski et al., 2022; Yu et al., 2022), indicating mitigation of hyperthermal conditions. Geochemical proxies of weathering intensity, salinity and clayiness, along with increased hygrophyte/xerophyte ratio, also demonstrate the transition to wetter conditions (Shu et al., 2022; Zhu et al., 2022).

Conclusions

The diverse community of the HSG Formation, consisting of tetrapods, plant stems, rhizoliths, and diverse ichnofossils suggests a rapid recovery of life in low-latitude terrestrial environments in

the Spathian, as little as ~2 Myr after the end-Permian biotic crisis. The newly discovered vertebrate fossils are medium-sized carnivores of approximately early Spathian age, representing the earliest tetrapod found in North China. High ichnodiversity and ichnodisparity, and three types of large burrows, not only results in intensified bioturbation, but provide additional information about the enriched local biota. Furthermore, enhanced burrowing behavior is considered a key survival-recovery strategy to adapt to harsh climatic and environmental conditions on land. The cooccurred trace fossils and stems, and coeval tetrapod in alluvial plain facies, suggesting that the riverain regions could be refugia for the reorganization of post-EPME ecosystem.

Data Availability Statement

All supplementary data related to this paper is available in the end of this manuscript.

Article and author information

Author details

Wenwei Guo

● College of Culture and Tourism, Zhangzhou Institute of Technology, Zhangzhou, People's Republic of China

● State Key Laboratory of Biogeology and Environmental Geology, School of Earth Science, China University of Geosciences, Wuhan, People's Republic of China

Contribution: Conceptualization; Methodology; Validation; Formal analysis; Investigation; Data curation; Writing–original draft preparation; Writing–review & editing; Visualization

Competing interests: No competing interests declared

Li Tian

State Key Laboratory of Biogeology and Environmental Geology, School of Earth Science, China

University of Geosciences, Wuhan, People's Republic of China

Contribution: Conceptualization; Methodology; Validation; Formal analysis; Investigation; Data

curation; Writing–original draft preparation; Writing–review & editing; Visualisation; Project

administration; Funding acquisition

For correspondence: tianlibgeg@cug.edu.cn

Competing interests: No competing interests declared

Daoliang Chu

State Key Laboratory of Biogeology and Environmental Geology, School of Earth Science, China

University of Geosciences, Wuhan, People's Republic of China

Contribution: Conceptualization; Methodology; Validation; Investigation; Data curation; Writing–

original draft preparation; Writing–review & editing; Visualization; Project administration; Funding

acquisition

Competing interests: No competing interests declared

Wenchao Shu

State Key Laboratory of Biogeology and Environmental Geology, School of Earth Science, China

University of Geosciences, Wuhan, People's Republic of China

Contribution: Methodology; Validation; Formal analysis; Investigation; Writing–original draft

preparation; Writing–review & editing; visualization

Competing interests: No competing interests declared

Michael J. Benton

School of Earth Sciences, University of Bristol, UK

Contribution: Conceptualization; Methodology; Validation; Formal analysis; Investigation;

Writing–review & editing; Visualization; Funding acquisition

Competing interests: No competing interests declared

Jun Liu

Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing,

People's Republic of China

Contribution: Conceptualization; Methodology; Validation; Formal analysis; Investigation;

Writing–review & editing; Visualization; Funding acquisition

Competing interests: No competing interests declared

Jinnan Tong

State Key Laboratory of Biogeology and Environmental Geology, School of Earth Science, China

University of Geosciences, Wuhan, People's Republic of China

Contribution: Conceptualization; Validation; Formal analysis; Investigation; Writing–review &

editing; Visualization; Supervision; Project administration; Funding acquisition

Competing interests: No competing interests declared

Acknowledgments

We thank Kaixuan Ji, Gan Liu and Yuyang Wu for assistance in the field. We also thank anonymous reviewers for their comments and constructive suggestions. This work was supported by the National Natural Science Foundation of China (grant nos. 42030513) to L. T., D. C. and J. T., the Strategic Priority Research Program of the Chinese Academy of Sciences (grant no. XDB26000000) to J. L. and the Natural Environment Research Council (UK) of grant no. NE/P013724/1 to M. J. B.

References

Allen BJ, Wignall PB, Hill DJ, Saupe EE, Dunhill AM. 2020. The latitudinal diversity gradient of tetrapods across the Permo-Triassic mass extinction and recovery interval. *Proceedings of the Royal Society B*, **287**: 20201125. https://doi.org/10.1098/rspb.2020.1125

Baucon A, Ronchi A, Felletti F, Neto de Carvalho C. 2014. Evolution of Crustaceans at the edge of the end-Permian crisis: Ichnonetwork analysis of the fluvial succession of Nurra (Permian–Triassic, Sardinia, Italy). *Palaeogeography*, *Palaeoclimatology*, *Palaeoecology*, **410**: 74–103. https://doi.org/10.1016/j.palaeo.2014.05.034

Benton MJ. 2018. Hyperthermal-driven mass extinctions: killing models during the Permian–Triassic mass extinction. *Philosophical Transactions of the Royal Society A*, **376**: 20170076. https://doi.org/10.1098/rsta.2017.0076

Bernardi M, Petti FM, Benton MJ. 2018. Tetrapod distribution and temperature rise during the Permian-Triassic mass extinction. *Proceedings of the Royal Society B*, **285**: 20172331. https://doi.org/10.1098/rspb.2017.2331

Brück PM. 1987. A note on the trace fossil *Beaconites barretti* in the Old Red Sandstone of County

Dublin, Ireland. *Proceedings of the Geologists' Association*, **98**: 259–263.

https://doi.org/10.1016/S0016-7878(87)80043-3

Buatois LA, Wisshak M, Wilson MA, Mángano MG, 2017. Categories of architectural designs in

trace fossils: a measure of ichnodisparity. *Earth-Science Reviews*, **164**: 102–181.

https://doi.org/10.1016/j.earscirev.2016.08.009

Buatois LA, Borruel-Abadía V, De la Horra R, Galán-Abellán AB, López-Gómez J, Barrenechea JF,

Arche A. 2021. Impact of Permian mass extinctions on continental invertebrate infauna. *Terra*

Nova, **33**: 455–464. https://doi.org/10.1111/ter.12530.

Chen ZQ, Benton MJ. 2012. The timing and pattern of biotic recovery following the end-Permian

mass extinction. *Nature Geoscience*, **5**: 375–383. https://doi.org/10.1038/ngeo1475

Chu DL, Tong JN, Bottjer DJ, Song HJ, Song HY, Benton MJ, Tian Li, Guo WW. 2017. Microbial

mats in the terrestrial Lower Triassic of North China and implications for the Permian–Triassic

mass extinction. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **474**: 214–231.

https://doi.org/10.1016/j.palaeo.2016.06.013

Dai X, Davies JHFL, Yuan Z, Brayard A, Ovtcharova M, Xu GH, Liu XK, Smith CPA, Schweitzer

CE, Li MT, Perrot MG, Jiang SY, Miao LY, Cao YR, Yan J, Bai RY, Wang FY, Guo W, Song HY,

Tian L, Dal Corso J, Liu YT, Chu DL, Song HJ. 2023. A Mesozoic fossil lagerstätte from 250.8

million years ago shows a modern-type marine ecosystem. *Science*, **379**: 567–572. DOI:

10.1126/science.adf1622

Fan JX, Shen SZ, Erwin DH, Sadler PM, MacLeod N, Cheng QM, Hou XD, Yang J, Wang XD,

Wang Y, Zhang H, Chen X, Li GX, Zhang YC, Shi YK, Yuan DX, Chen Q, Zhang LN, Li C,

Zhao YY. 2020. A high-resolution summary of Cambrian to early Triassic marine invertebrate

biodiversity. *Science*, **367**: 272–277. DOI: 10.1126/science.aax4953

Guo WW, Tong JN, Tian L, Chu DL, Bottjer DJ, Shu WC, Ji KX. 2019. Secular variations of

ichnofossils from the terrestrial Late Permian-Middle Triassic succession at the Shichuanhe

section in Shaanxi Province, North China. *Global and Planetary Change*, **181**: 102978.

https://doi.org/10.1016/j.gloplacha.2019.102978

Guo WW, Tong JN, He Q, Hounslow MW, Song HY, Dal Corso J, Wignall PB, Ramezani J, Tian L,

Chu DL. 2022. Late Permian–Middle Triassic magnetostratigraphy in North China and its

implications for terrestrial-marine correlations. *Earth and Planetary Science Letters*, **585**:

117519. https://doi.org/10.1016/j.epsl.2022.117519

Huang BC, Yan Y, Piper JDA, Zhang D, Yi Z, Yu S, Zhou T. 2018. Paleomagnetic constraints on the

paleogeography of the East Asian blocks during Late Paleozoic and Early Mesozoic times.

Earth-Science Reviews, **186**: 8–36. https://doi.org/10.1016/j.earscirev.2018.02.004

Ji K, Wignall PB, Tong JN, Yu YY, Guo WW, Shu WC, Chu DL. 2022. Sedimentology of the latest

Permian to Early Triassic in the terrestrial settings of the North China Basin: Low-latitude

climate change during a warming-driven crisis. *Geological Society of America Bulletin*, **135**:

481-503. https://doi.org/10.1130/b36260.1

Joachimski MM, Müller J, Gallagher TM, Mathes G, Chu DL, Mouraviev F, Silantiev V, Sun YD,

Tong JN. 2022. Five million years of high atmospheric CO_2 in the aftermath of the

Permian-Triassic mass extinction. *Geology*, **50**: 650–654. https://doi.org/10.1130/G49714.1

Kraus MJ, Hasiotis ST. 2006. Significance of Different Modes of Rhizolith Preservation to

Interpreting Paleoenvironmental and Paleohydrologic Settings: Examples from Paleogene

344 Paleosols, Bighorn Basin, Wyoming, U.S.A. *Journal of Sedimentary Research*, **76**: 633–646.

345 https://doi.org/10.2110/jsr.2006.052

346 **Li JL**, Wu X, Zhang F. 2008. *The Chinese Fossil Reptiles and Their Kin*. Science Press, Beijing.

347 **Liu J**, Abdala F, Angielczyk KD, Sidor CA. 2022. Tetrapod turnover during the Permo-Triassic

348 transition explained by temperature change. *Earth-Science Reviews*, **224**: 103886.

349 https://doi.org/10.1016/j.earscirev.2021.103886.

350 **Liu XK**, Song HJ, Bond DPG, Tong JN, Benton MJ. 2020. Migration controls extinction and

351 survival patterns of foraminifers during the Permian-Triassic crisis in South China.

352 *Earth-Science Reviews*, **209**: 103329. https://doi.org/10.1016/j.earscirev.2020.103329

353 **Luo M**, Shi GR, Buatois LA, Chen ZQ. 2020. Trace fossils as proxy for biotic recovery after the

354 end-Permian mass extinction. A critical review. *Earth-Science Reviews*, **203**: 103059.

355 https://doi.org/10.1016/j.earscirev.2019.103059

356 **Matamales-Andreu R**, Penalver E, Mujal E, Oms O, Scholze F, Juarez J, Galobart A, Fortuny J.

357 2021. Early-Middle Triassic fluvial ecosystems of Mallorca (Balearic Islands): Biotic

358 communities and environmental evolution in the equatorial western peri-Tethys. *Earth-Science*

359 *Reviews*, **222**: 103783. https://doi.org/10.1016/j.earscirev.2021.103783

360 **Marchetti L**, MacDougall MJ, Buchwitz M, Canoville A, Herde M, Kammerer CF, Fröbisch J. 2024.

361 Origin and early evolution of vertebrate burrowing behaviour. *Earth-Science Reviews*, **250**:

362 104702. https://doi.org/10.1016/j.earscirev.2024.104702

363 **Mays C**, Vajda V, Frank TD, Fielding CR, Nicoll RS, Tevyaw AP, McLoughlin S. 2020. Refined

364 Permian–Triassic floristic timeline reveals early collapse and delayed recovery of south polar

365 terrestrial ecosystems. *Geological Society of America Bulletin*, **132**: 1489–1513.

366 https://doi.org/10.1130/B35355.1

Mays C, McLoughlin S, Frank TD, Fielding CR, Slater SM, Vajda V. 2021. Lethal microbial blooms delayed freshwater ecosystem recovery following the end-Permian extinction. *Nature Communications*, **12**: 5511. https://doi.org/10.1038/s41467-021-25711-3

McLoughlin S, Mays C, Vajda V, Bocking M, Frank TD, Fielding CR. 2020. Dwelling in the Dead Zone—Vertebrate Burrows Immediately Succeeding the End-Permian Extinction Event in Australia. *Palaios*, **35**: 342-357. https://doi.org/10.2110/palo.2020.007

Minter NJ, Buatois LA, Mángano MG, Davies NS, Gibling MR, MacNaughton RB, Labandeira CC. 2017. Early bursts of diversification defined the faunal colonization of land. *Nature Ecology & Evolution*, **1**: https://doi.org/10.1038/s41559-017-0175

Mujal E, Fortuny J, Pérez-Cano J, Dinarès-Turell J, Ibáñez-Insa J, Oms O, Vila I, Bolet A, Anadón P. 2017. Integrated multi-stratigraphic study of the Coll de Terrers late Permian–Early Triassic continental succession from the Catalan Pyrenees (NE Iberian Peninsula): A geologic reference record for equatorial Pangaea. *Global and Planetary Change*, **159**: 46–60. https://doi.org/10.1016/j.gloplacha.2017.10.004

Mujal E, Marchetti L, Schoch RR, Fortuny J. 2020. Upper Paleozoic to Lower Mesozoic tetrapod ichnology revisited: photogrammetry and relative depth pattern inferences on functional prevalence of Autopodia. *Frontiers in Earth Science*, **8**: 248. https://doi.org/10.3389/feart.2020.00248

Romano M, Bernardi M, Petti FM, Rubidge B, Hancox J, Benton MJ. 2020. Early Triassic terrestrial tetrapod fauna: a review. *Earth-Science Reviews*, **210**: 103331. https://doi.org/10.1016/j.earscirev.2020.103331

Roopnarine PD, Angielczyk KD, Weik A, Dineen A. 2019. Ecological persistence, incumbency and

reorganization in the Karoo Basin during the Permian-Triassic transition. *Earth-Science*

Reviews, **189**: 244−263. https://doi.org/10.1016/j.earscirev.2018.10.014

Shu WC, Tong JN, Tian L, Benton MJ, Chu DL, Yu JX, Guo WW. 2018. Limuloid trackways from

Permian–Triassic continental successions of North China. *Palaeogeography*, *Palaeoclimatology*,

Palaeoecology, **508**: 71–90. https://doi.org/10.1016/j.palaeo.2018.07.022

Shu WC, Tong JN, Yu JX, Hilton J, Benton MJ, Shi X, Diez JB, Wignall PB, Chu DL, Tian L, Yi,

ZX, Mao YD. 2022. Permian−Middle Triassic floral succession in North China and implications

for the great transition of continental ecosystems. *Geological Society of America Bulletin*, **135**:

1747–1767. https://doi.org/10.1130/B36316.1

Song HJ, Huang S, Jia EH, Dai X, Wignall PB, Dunhill AM. 2020. Flat latitudinal diversity gradient

caused by the Permian-Triassic mass extinction. *Proceedings of the National Academy of*

Sciences of the United States of America, **117**: 17578–17583.

https://doi.org/10.1073/pnas.1918953117

Sun YD, Joachimski MM, Wignall PB, Yan CB, Chen YL, Jiang HS, Wang LN, Lai XL. 2012.

Lethally hot temperatures during the Early Triassic greenhouse. *Science*, **338**: 366–370.

DOI:10.1126/science.1224126

Tu CY, Chen ZQ, Retallack GJ, Huang Y, Fang Y. 2016. Proliferation of MISS-related microbial

mats following the end-Permian mass extinction in terrestrial ecosystems: Evidence from the

Lower Triassic of the Yiyang area, Henan Province, North China. *Sedimentary Geology*, **333**:

50−69. https://doi.org/10.1016/j.sedgeo.2015.12.006

Tverdokhlebov VP, Tverdokhlebova GI, Surkov MV, Benton MJ. 2003. Tetrapod localities from the

Triassic of the SE of European Russia. *Earth-Science Reviews*, **60**: 1–66.

https://doi.org/10.1016/S0012-8252(02)00076-4

Vajda V, McLoughlin S, Mays C, Frank TD, Fielding CR, Tevyaw A, Lehsten V, Bocking M, Nicoll

RS. 2020. End-Permian (252 Mya) deforestation, wildfires and flooding – an ancient biotic

crisis with lessons for the present. *Earth and Planetary Science Letters*, **529**: 115875.

https://doi.org/10.1016/j.epsl.2019.115875

Vajda V, Kear BP. 2024. An earliest Triassic riparian ecosystem from the Bulgo Sandstone (Sydney

Basin), Australia: palynofloral evidence of a high-latitude terrestrial vertebrate habitat after the

end-Permian mass extinction. *Alcheringa: An Austalasian Journal of Palaeontology*, 1−12.

https://doi.org/10.1080/03115518.2024.2392489

Viglietti PA, Rojas A, Rosvall M, Klimes B, Angielczyk KD. 2022. Network-based biostratigraphy

for the late Permian to mid-Triassic Beaufort Group (Karoo Supergroup) in South Africa

enhances biozone applicability and stratigraphic correlation. *Palaeontology*, **65**: e12622.

https://doi.org/10.1111/pala.12622

Wignall PB. 2015. *The Worst of Times: How Life on Earth Survived Eighty Million Years of*

Extinction. Princeton University Press, NJ: Princeton.

Xing ZF, Lin J, Fu YX, Zheng W, Liu YL, Qi YA. 2021. Trace fossils from the Lower Triassic of

North China — a potential signature of the gradual recovery of a terrestrial ecosystem.

Palaeoworld, **30**: 95–105. https://doi.org/10.1016/j.palwor.2020.06.002

Yu YY, Tian L, Chu DL, Song HY, Guo WW, Tong JN. 2022. Latest Permian–Early Triassic

paleoclimatic reconstruction by sedimentary and isotopic analyses of paleosols from the

Shichuanhe section in central North China Basin. *Palaeogeography, Palaeoclimatology,*

Palaeoecology, **585**: 110726. https://doi.org/10.1016/j.palaeo.2021.110726

Zhao XD, Zheng DR, Xie GW, Jenkyns HC, Guan CG, Fang Y, He J, Yuan XQ, Xue NH, Wang H,

Li S, Jarzembowski, EA, Zhang HC, Wang B. 2020. Recovery of lacustrine ecosystems after the

end-Permian mass extinction. *Geology*, **48**: 609–613. https://doi.org/10.1130/G47502.1

Zheng DR, Chang SC, Wang H, Fang Y, Wang J, Feng CQ, Xie GW, Jarzembowski EA, Zhang HC,

Wang B. 2018. Middle-Late Triassic insect radiation revealed by diverse fossils and isotopic

ages from China. *Science Advances*, **4**: eaat1380. https://doi.org/10.1126/sciadv.aat1380

Zheng W, Wan EZ, Xu X, Li D, Dai MY, Qi YA, Xing ZF, Liu YL. 2021. The variations of

terrestrial trace fossils and sedimentary substrates after the end-Permian extinction in the

Dengfeng area, North China. *Geological Journal*, **58**: 1223–1238.

https://doi.org/10.1002/gj.4657

Zhu ZC, Liu YQ, Kuang HW, Benton MJ, Newell AJ, Xu H, An W, Ji SA, Xu SC, Peng N, Zhai QG.

2019. Altered fluvial patterns in North China indicate rapid climate change linked to the

Permian-Triassic mass extinction. *Scientific Reports*, **9**: 16818.

https://doi.org/10.1038/s41598-019-53321-z

Zhu ZC, Kuang HW, Liu YQ, Benton MJ, Newell AJ, Xu H, An W, Ji SA, Xu SC, Peng N, Zhai QG.

2020. Intensifying aeolian activity following the end-Permian mass extinction: Evidence from

the Late Permian–Early Triassic terrestrial sedimentary record of the Ordos Basin, North China.

Sedimentology, **67**: 2691-2720. https://doi.org/10.1111/sed.12716

Zhu ZC, Liu YQ, Kuang HW, Newell AJ, Peng N, Cui MM, Benton MJ. 2022. Improving

paleoenvironment in North China aided Triassic biotic recovery on land following the

end-Permian mass extinction. *Global and Planetary Change*, **216**: 103914.

Figure 1. Location of the studied regions and lithological columns. (a–b) Permian-Triassic paleogeographic map of North China and the studied successions (stars) and outcrops (points). Base map of a is modified from Sun et al. (2012). (c) Depositional facies and detailed distribution of trace fossils in three main successions.

Figure 2. Ichnofossils from the Heshanggou Formation of North China. (a) Shallow tiers *Kouphichnium* and *Helminthoidichnites* (He) are crosscut by immediate tier *Palaeophycus* (Pa). (b) *Skolithos* cf. *serratus* with faint oblique striations. (c) Y-shaped *Psilonichnus* isp. (d) Downward unbranched and tapered rhizocretion. (e) Shallow tiers *Beaconites coronus*, arrows show tightly stacked arcuate meniscus. (f) *Camborygma* isp. shows enlarged terminal chamber and possible transverse scratches (arrows). (g) *Gordia* isp. with darker and finer infills. (h) High density *Palaeophycus tubularis* preserved on the sole of thick sandstone. (i) Inclined *Planolites beverleyensis* within siltstone. (j) *Taenidium barretti* (arrows) pass through the rippled surface. (k)

Horizontal *Camborygma*, the outer surface is intertwined with tinny root traces (arrows). (l) Biserial

Diplichnites gouldi. (m-n) Internode cross-section of *Neocalamites* stem and micro-CT structure,

showing the clear ribs and grooves. (o-p) Large ?*Beaconites* on top of rippled sandstone, rectangle in

(p) shows meniscus-like portions comprising of gritty infillings. Scale bar of (d, f, h, o-p) are 40 mm,

the rest are all 20 mm.

Figure 3. Vertebrates from the Heshanggou Formation of North China. (a–b) Skull elements of

Xilousuchus sapingensis and drawings. (c) *Fugusuchus hejiapanensis*. (d) *Eumetabolodon*

bathycephalus. (e–f) *Shaanbeikannemeyeria xilouensis* and drawing. (g) *Pentaedrusaurus*

ordosianus. (h) *Hazhenia concava*, (i) Dashed line shows the boundary between the Liujiagou and

Heshanggou formations. Arrow displays fossil horizon of the inserted picture at the base of the

Heshanggou Formation. (b, c, f) are from Li et al. (2008). Scale bars are 40 mm.

Figure 4. Ichnofossil criteria in North China and global terrestrial ecosystem changes from latest Permian to earliest Middle Triassic. Geochronological timescale is based on the latest version of the International Chronostratigraphic Chart (https://stratigraphy.org/). Plant richness from Shu et al. (2022). Numbers in Burrow size column represent the mean trace fossil sizes from the investigated interval. Drainage conditions of paleosols are inferred from preservation of rhizoliths and their relative depths. Ranges of microbially induced sedimentary structures (MISS) are from Chu et al. (2017). Atmospheric CO_2 curves are modified from Joachimski et al. (2022). Abbreviations: Ind. = Induan; G. = Griesbachian; D. = Dienerian.

Supplementary files for

Rapid riparian ecosystem recovery in low-latitudinal North China following the end-Permian mass extinction

Wenwei Guo[1, 2], Li Tian[2*], Daoliang Chu[2], Wenchao Shu[2], Michael J. Benton[3], Jun Liu[4], and Jinnan Tong[2]

[1]College of Culture and Tourism, Zhangzhou Institute of Technology, Zhangzhou 363000, People's Republic of China.

[2]State Key Laboratory of Biogeology and Environmental Geology, China University of Geosciences, Wuhan, 430074, People's Republic of China.

[3]School of Earth Sciences, University of Bristol, BS8 1RJ, UK.

[4]Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing 100044, People's Republic of China

Introduction

The Supplementary files include: (1) file 1: Early Triassic chronostratigraphic timescale in North China (Figures S1-2) and brief description of facies associations of the studied sections (Table S1); file 2: detailed description of traces fossils (Fig S3-6), fossil size variations (Figures S7-8), reconstruction of the late Early Triassic (Heshanggou Formation) coastal mudplain to alluvial ecosystem (Figure S9), designations of ichnodiversity and tiering level (Tables S2-4) and possible trace producers on land   (Table S5).

File 1. Integrated stratigraphic timescale and depositional environment

Detailed sedimentological investigations at the three studied sections have been undertaken by Ji et al. (2022) and Yu et al., (2022). Therefore, we primarily summarize the lithofacies and facies associations, and add more information about trace fossils (Figure S1 and Table S1).

Body fossil records in the red bed predominated Permian–Triassic transitional successions were too limited to establish a high-resolution timescale for global correlation. Based on the U-Pb CA-ID-TIMS age calibrated magnetostratigraphy from the Shichuanhe section (Guo et al., 2022), we complied other detrital zircon U-Pb ages during the Early Triassic, to provide a better constrained chronostratigraphy in North China (Zhu et al., 2019; 2022; Lu et al., 2022).

Figure S1. Lithology, sedimentology and ichnology of the studied sections.

Figure S2. Early Triassic geochronological scale in North China. The Geo-magnetic polarity timescale (GPTS) is from Hounslow and Muttoni (2010) and Hounslow and Balabanov (2018). Magnetostratigraphy and the CA-ID-TIMS age of [1] are from Guo et al. (2022). Detrital zircon U-Pb ages are based on: [2] = tuffaceous claystones sampled from the uppermost part of the Sunjiagou Formation (Lu et al., 2022), [2] = around middle part of the Liujiagou Formation (Zhu et al., 2019), [3] = sandstone from the base of the Heshanggou Formation and tuffs from the bottom Ermaying Formation (EMY; Zhu et al., 2022). G. = Griesbachian, D. = Dienerian.

 Table S1. Brief description of lithofacies, facies associations (FA) and trace fossils at the studied sections.

FA	Lithology	Trace fossils	Depositional environments
FA1	• Dominated by red mud-siltstone with paleosols (Msp) and interbedded with cross-bedded sandstone (Scb). • Well-developed paleosols in Msp, of moderate to very high maturity, contained layered calcareous nodules up to 10 cm in diameter, which were assigned to Aridisols and Vertisols.	• Root traces (Rh) and burrows (e.g., *Planolites* (*Pl*, *Taenidium* (*Ta*)) are common in Msp. • Scb is also bioturbated, especially at the Shichuanhe section. Diverse traces, including *Lockeia* (*Lo*), *Protovirgularis*, *Pa*, *Ta*, *Spongeliomorpha*, and several informally designated types have been identified. • Generally, animal traces are crosscut by roots.	• Distributed in the middle and upper Sunjiagou Formation. • **Arid floodplain.**
FA2	• Comprised of grayish-green silty mudstone (Ms), marine marls and calcareous sandstone. • A mixed continental-marine fauna, including conchostracans, bivalves, brachiopods and plant debris, have been identified, indicative of **terrestrial facies with periodical marine flooding events** (Chu et al., 2019).	No trace fossil found.	• Exclusively distributed in the lower Sunjiagou Formation of Shichuanhe section.
FA3	• Comprise red Ms and Sandstone (S). • In upper Sunjiagou Formation, gypsum crystals, blocky calcite cements, Fe-oxide hypocoating occasionally occurred.	• Bioturbation is absent. • Strata in the lower Heshanggou Formation are moderately reworked, several ichnogenera (e.g., *Pl*, *Palaeophycus* (*Pa*), *Beaconites* (*Be*), *Ta*), plant debris, root traces, and vertebrate remains, have been identified.	• Distributed in upper Sunjiagou and lower Heshanggou formations. • **Costal mudplain facies.**
FA4	• Dominated by Pale red to grey purple Scb, planar-laminated sandstone (Sp), interbedded with thin dark	• *Kouphichnium* (*Kou*) (Shu et al., 2018) and tiny *Helminthoidichnites* have been found in Si.	• Distributed in the Liujiagou and

	red siltstone (Si), where Inceptisols are weakly- developed.	• Vertical burrows of *Skolithos* (*Sk*) and Rh, occurred occasionally in Scb, but the sediments were weakly bioturbated.	Ermaying formations. • **Low sinuosity fluvial system.**
FA5	• Consists of Scb, planar-laminated sandstone (Sp), hummocky and swaley cross-stratified sandstone (Sh), current and wave-rippled sandstone (Scr, Swr), interbedded with Intraformational conglomerate (Ci), which • Microbially induced sedimentary structures (MISS) are quite common, especially at Liulin Section (Chu et al., 2015, 2017).	• Depauperated ichnofauna, dominated by monospecific ichnoassemblage, i.e., *Sk*, *Lo*, *Gordia*. • Bioturbation in this interval is weak.	• Distributed in the middle-upper Liujiagou Formation. • **Shallow lake and lake margin facies, with influence of major storm events.**
FA6	• Consists of dark red massive Msp, Scb, and Sp. • Sheetlike sandy bodies display lightly erosive bases.	• Msp contain diverse trace fossils, including *Kou*, *Diplichnites*, *Be*, *Ta*, *Camborygma*, *Psinolichnus* and various types of root traces, which moderately disturbed the sediments. • Trace fossils in sandstones are mainly *Sk*, *Pl* and Rh. • In several horizons, the top of several centimeters of the strata is strongly reworked.	• Heshanggou Formation. • **Alluvial plain with shallow lakes facies.**
FA7	• Dominated by massive sandstone with sharp base and trough cross-bedded or planar lamination, interlayered with dark red Msp and intraclast conglomerates, constituting several fining-upward sequences. • Vertisols and calcic Inceptisols, of low maturity, are well-developed in Msp.	• Relatively simple ichnoassemblage, dominated by *Sk*, *Pa*, *Ta* and Rh • Root traces generally crosscut animal traces.	• In the Heshanggou Formation of Dayulin section. • **Distal terminal fan facies**

File 2. Ichnological information

In this section, all trace fossils encountered (Figures S2–5) and their size variations Figures S6–7) are provided herein. Ichnodisparity designation (Tables S2–4) follows Buatois et al. (2017). Traces from the Shichuanhe section have been documented by Guo et al. (2019), but several new ichnogenera have been found during the ongoing field seasons and are described here (Figure S3). Additionally, possible trace maker(s) on land are also discussed, based on paleontological and neoichnological evidence (Table S5). Accordingly, an artistic illustration of the early recovered ecosystem on land in North China have been reconstructed (Figure S8).

Figure S3. Ichnofauna from the Liujiagou Formation at Liulin section. (a-b) Ovate to elongated almond-shaped *Lockeia siliquaria* roughly arranged along a similar direction, fossils were preserved as convex hypichia in sandstone with desiccation cracks (arrow in a). (c-dd) Simple unbranched surficial burrows with sharp turns (arrows in d), self-overcrossing occasionally seen, traces are identified as *Gordia indianaensis* and preserved in rippled sandstone (arrow in c). (e) Vertical *Skolithos linearis* in sandstone. (f) Vertical root traces, showing inconsistent burrow sizes and tiny bifurcations (arrows). (g) *Kouphichnium didactylum* with simple foot imprints and a pusher imprint.

Figure S4. Ichnofauna from the Heshanggou Formation. (a) Y-shaped *Psilonichnus* cf. *upsilon*, the branch portion displays similar size with the main tube. FA3 costal mudplain facies at Shichuanhe section. (b) *Diplichnites gouldi*, evenly spaced imprints are comma or elongated and incline to the midline. Fossils preserved in FA3 at Shichuanhe section. (c) High-density simple vertical *Skolithos linearis* at Dayunlin section. (d) Actively filled *Planolites beverleyensis* (*Pl*) pass through surficial Helminthoidichnites tenuis (*He*), which in turn, crosscut by *Skolithos* (*Sk*). Dayunlin section. (e) Horizontal *Taenidium serpentinum*, distance between menisci (arrow) is roughly equal to the burrow width. Dayunlin section. (f) Crowded *Palaeophycus tubularis* preserved on the sole of thick sandstone. Liulin section. (g) Horizontal to inclined, straight to curve, lined *Beaconites coronus* with moderately arcuate menisci. Liulin section.

Figure S5. Large burrows from the Heshanggou Formation. (a-b) Horizontal but inclined unlined burrow with
elliptical cross sections, branching and terminal chamber were no found. Possible lateral scratches could be seen on
the side (Arrows in b). trace fossil is informally assigned to undesigned burrow. Liuin section. (c-e) Large ?
Beaconites and other relatively small traces (e.g., *Palaeophycus*) on the rippled siltstone (ripple marks can be seen
at the top-right of c). Burrows display meniscus-like portions, comprising of gritty infillings (Arrow in e), akin to
Beaconites reported from breccia facies in the Devonian (Brück, 1987). Mafang outcrops. (f-h) Vertical to
subhorizontal burrows with circular cross sections, filled with calcareous sand. Trace fossils are similar to crayfish
burrows (*Camborygma*), although branches, terminal chamber or scratch marks were no found. Hongyatou outcrop.
(i-j) In situ preserved *Neocalamites* stems in the lower part of the Heshanggou Formation at Shichuanhe section and
Hongyatou outcrop, respectively. Typically, all the stems are associated with *Camborygma* and other burrows and
trackways.

Figure S6. Root traces from the Heshanggou Formation in studied sections and outcrops. (a-h) Vertical branched rhizolith with green or pale purple haloes, showing irregular burrow sizes and lateral small rootlets (d, f), Yiyang section. (i) Herringbone-like branched rhizotubules, Yiyang section. (j-k) Vertical tapered unbranched or bifurcated rhiozocretions, Yiyang section. (l) Simple vertical branched roots, Tuncun outcrops. (m) Horizontal small rootlets, Yiyang section. Classification of root traces are based on Klappa (1980) and Kraus and Hasiotis (2006). All scale bars are 20 mm.

584
585 **Figure S7. Size variation of all ichnogenera from there sections.** EMY = Ermaying Formation.

587 **Figure S8. Size variation of sing ichnogenus from Liulin and Dayulin sections during the Spathian.** EMY =

588 Ermaying Formation.

Figure S9. Reconstruction of the Spathian (Heshanggou Formation) coastal mudplain to alluvial ecosystem in North China. Plant communities in the coastal mudplain and alluvial facies are depauperated, dominated by *Neocalamites* and *Pleuromeia*, and only diversified at the top of the Heshanggou Formation (late Spathian; Shu et al., 2022). Reoccurrence of plants and tetrapods, coupled with diverse invertebrate, including limuloid, crayfish, spinicaudatan, and insect, etc., reveal reorganization of a relatively complex ecosystem in riverain regions during the Spathian.

Table S2. List of ichnogenera from the Shichuanhe section, Ichnodisparity designations and Tiering level.

	Ichnodisparity	Ichnogenus	Tier
Sunjiagou Formation	10-Passively filled horizontal burrows	*Palaeophycus*	Shallow-infaunal
	35-Vertical unbranched burrows	*Skolithos*	Deep -infaunal
	50-Maze and boxwork burrows	*Spongeliomorpha*	Shallow-infaunal
	12-Simple actively filled (meniscate) horizontal to oblique	*Taenidium*	Intermediate-infaunal
	1-Simple horizontal trails	*Helminthoidichnites*	Semi-infaunal
	3-Chevronate trails	*Protovirgularia*	Shallow-infaunal
	30-Isolated and serial oval to almond-shaped burrows	*Lockeia*	Semi-infaunal
	Rhizoliths	Rhizoliths	Deep-infaunal
Liujiagou Formation	10-Passively filled horizontal burrows	*Palaeophycus*	Shallow-infaunal
	35-Vertical unbranched burrows	*Skolithos*	Deep-infaunal
	36-Vertical single U- and Y-shaped burrows	*Arenicolites*	Deep-infaunal
		Psilonichnus	Intermediate-infaunal
	52-Simple to complex burrows with terminal chambers	*Camborygma*	Deep-infaunal
		Macanopsis	Intermediate-infaunal
	1-Simple horizontal trails	*Helminthoidichnites*	Semi-infaunal

Heshanggou Formation		Diplichnites	Surficial
	6-Trackways and scratch imprints	Monomorphichnus	Surficial
		Kouphichnium	Surficial
	10-Passively filled horizontal burrows	Palaeophycus	Shallow to Intermediate-infaunal
	11-Simple actively filled (massive) horizontal to oblique	Planolites	Intermediate-infaunal
	12-Simple actively filled (meniscate) horizontal to oblique	Scoyenia	Intermediate-infaunal
		Taenidium	Shallow-infaunal
	35-Vertical unbranched burrows	Skolithos	Deep-infaunal
	Rhizoliths	Rhizoliths	Deep-infaunal

605 **Table S3. List of ichnogenera from the Heshanggou Formation of the Dayulin section, Ichnodisparity**
606 **designations and tiering level.**

	Ichnodisparity	Ichnogenus	Tier
Heshanggou Formation	1-Simple horizontal trails	Helminthoidichnites	Semi-infaunal
	10-Passively filled horizontal burrows	Palaeophycus	Shallow to Intermediate-infaunal
	11-Simple actively filled (massive) horizontal to oblique	Planolites	Intermediate to deep- infaunal
	12-Simple actively filled (meniscate) horizontal to oblique	Taenidium	Intermediate-infaunal
		Beaconites	Intermediate-infaunal
		Scoyenia	Intermediate-infaunal
	35-Vertical unbranched burrows	Skolithos	Deep-infaunal
	36-Vertical single U- and Y-shaped burrows	Arenicolites	Deep-infaunal
		Psilonichnus	Intermediate to deep- infaunal
	Rhizoliths	Rhizoliths	Deep-infaunal

609 **Table S4. List of ichnogenera from the Liulin section, Ichnodisparity designations and tiering level.**

	Ichnodisparity	Ichnogenus	Tier
Sunjiagou Formation	10-Passively filled horizontal burrows	Palaeophycus	semi-infaunal
	11-Simple actively filled (massive) horizontal to oblique	Planolites	Intermediate-infaunal
	12-Simple actively filled (meniscate) horizontal to oblique	Taenidium	semi-infaunal
	Rhizoliths	Rhizoliths	Deep-infaunal
Liujiagou Formation	1-Simple horizontal trails	Helminthoidichnites	Semi-infaunal
		Gordia	Semi-infaunal
	6-Trackways and scratch imprints	Kouphichnium	Surficial
	30-Isolated and serial oval to almond-shaped burrows	Lockeia	Shallow to Intermediate-infaunal
	Rhizoliths	Rhizoliths	Deep-infaunal
Heshanggou	6-Trackways and scratch imprints	Kouphichnium	Surficial

Formation		Diplichnites	Surficial
	10-Passively filled horizontal burrows	Palaeophycus	Shallow to Intermediate-infaunal
	11-Simple actively filled (massive) horizontal to oblique	Planolites	Intermediate-infaunal
	35-Vertical unbranched burrows	Skolithos	Deep-infaunal
	12-Simple actively filled (meniscate) horizontal to oblique	Taenidium	Intermediate-infaunal
		Beaconites	Intermediate-infaunal
	1-Simple horizontal trails	Helminthoidichnites	Semi-infaunal
		Gordia	Shallow-infaunal
	Rhizoliths	Rhizoliths	Deep-infaunal

612 **Table S5. Possible trace producers on land.**

Morphology	Ichnogenus	Intepretation
Simple horizontal trails	*Helminthoidichnites tenuis*	• Could be made by arthropods, vermiform animals, or insect larvae (Buatois et al., 1996, 1998).
Passively or actively filled burrow	*Palaeophycus*, *Planolites*	• Probably produced by insects or arthropods (Pemberton and Frey, 1982; Buatois and Mángano, 1993; Keighley and Pickerill, 1995).
Burrows with terminal chambers	*Camborygma*	• Freshwater crayfishes (Hasiotis and Mitchell, 1993; Hasiotis et al., 1993).
Vertical unbranched, single U- or Y-shaped burrows	*Skolithos,*	• Most of the *Skolithos* are smooth, but several specimens display distinct subvertical scratches. • arachnids, coleopterans, crickets, cicadas, bees or annelids can all build such burrows (Schlirf et al., 2001; do Nascimento and Netto, 2019).
	Arenicolites,	• Deeply extended with tiny tubes and wide limb distance. • Burrowed by polychaeta worms, crustaceans or may fly larvaes (Bromley, 1996; Rindsberg and Kopaska-Merkel, 2005).
	Psilonichnus	• Display similar sizes with branches, with bifurcation angle at 15°-45°. • Generally occurred in the Cenozoic marine facies, and constructed by decapod (Frey et al., 1984; de Carvalho, 2016). • Neoichnological investigations show that terrestrial arthropod, such as scorpion, can also produce similar Y-shaped burrows as temporary shelters or permanent dwellings (Hembree, 2016).
Backfilled burrows	*Beaconites* *Scoyenia* *Taenidium*	• Could be made by insects (cicadas, beetle larvaes), millipedes, oligochetes (earthworms) or vermiform on land, based on modern observations (Keighley and

		Pickerill, 1994; Rodríguez-Tovar et al., 2016; Boyd and McIlroy, 2017).
	? *Beaconites*	• Could be produced by either invertebrate, like arthropods and polychaete worms, or vertebrate, such as reptiles, amphibians (Brück, 1987; Allen and Williams, 1981; Morrissey and Braddy, 2004).
Trackways and scratch imprints	*Diplichnites* *Monomorphichnus*	• Could be left by myriapods or morphological similar arthropods (Bradshaw, 1981).
	Kouphichnium	• Limuloid (i.e., Shu et al., 2018).

References

Allen JRL, Williams BPJ. 1981. *Beaconites antarcticus*: a giant channel-associated trace fossil from the Lower Old Red Sandstone of South Wales and the Welsh Borders. *Geological Journal*, **16**: 255–269. https://doi.org/10.1002/gj.3350160405

Boyd C, McIlroy D. 2017. Three-Dimensional Morphology of *Beaconites capronus* from Northeast England. *Ichnos*, **24**: 250–258. doi: 10.1080/10420940.2017.1282862

Bradshaw MA. 1981. Paleoenvironmental interpretations and systematics of Devonian trace fossils from the Taylor Group (lower Beacon Supergroup), Antarctica. *New Zealand Journal of Geology and Geophysics*, **24**: 615–652. https://doi.org/10.1080/00288306.1981.10421537

Bromley RG. 1996. *Trace Fossils: Biology, Taxonomy and Applications*. London: Chapman & Hall.

Brück PM. 1987. A note on the trace fossil Beaconites barretti in the Old Red Sandstone of County Dublin, Ireland. *Proceedings of the Geologists' Association*, **98**: 259–263. https://doi.org/10.1016/S0016-7878(87)80043-3

Buatois LA, Mángano MG. 1993. Trace fossils from a Carboniferous turbiditic lake: implications for the recognition of additional nonmarine ichnofacies. *Ichnos*, **2**: 237–258. doi: 10.1080/10420949309380098

Buatois LA, Mángano MG, Wu XT, Zhang GC. 1996. Trace fossils from Jurassic lacustrine turbidites of the Anyao Formation (central China) and their environmental and evolutionary significance. *Ichnos*, **4**: 287–303. doi: 10.1080/10420949609380137

Buatois LA, Mángano MG, Maples G, Lanier WP. 1998. Ichnology of an upper Carboniferous fluvio-estuarine paleovalley: The Tonganoxie sandstone, Buildex quarry, eastern Kansas, USA. *Journal of Paleontology*, **72**: 152–180. doi: 10.1017/S0022336000024094

Buatois LA, Wisshak M, Wilson MA, Mángano MG. 2017. Categories of architectural designs in trace fossils: a measure of ichnodisparity. *Earth-Science Reviews*, **164**: 102–181. https://doi.org/10.1016/j.earscirev.2016.08.009

Chu DL, Tong JN, Song HJ, Benton MJ, Bottjer DJ, Song HY, Tian L. 2015. Early Triassic wrinkle structures on land: stressed environments and oases for life. *Scientific Reports*, **5**: 10109, https://doi.org/10.1038/srep10109.

Chu DL, Tong JN, Bottjer DJ, Song HJ, Song HY, Benton MJ, Tian L, Guo WW. 2017. Microbial mats in the terrestrial Lower Triassic of North China and implications for the Permian–Triassic mass extinction. *Palaeogeography*, *Palaeoclimatology*, *Palaeoecology*, **474**: 214–231. https://doi.org/10.1016/j.palaeo.2016.06.013

Chu DL, Tong JN, Benton MJ, Yu JX, Huang YF. 2019. Mixed continental-marine biotas following the Permian-Triassic mass extinction in South and North China. *Palaeogeography*, *Palaeoclimatology*,

Palaeoecology, 519: 95–107. https://doi.org/10.1016/j.palaeo.2017.10.028

de Carvalho CN. 2016. *Psilonichnus* Fürsich, 1981 in its type-locality (Praia do Salgado, western Portugal). *Comunicaçõe Geológicas*, **103**: 13–21.

do Nascimento DL, Netto RG. 2019. *Skolithos serratus* in paleosols: Paleobiological, paleoecological, and paleobiogeographical insights. *Palaeogeography*, *Palaeoclimatology*, *Palaeoecology*, **530**: 152–162. https://doi.org/10.1016/j.palaeo.2019.05.044

Frey RW, Curran HA, Pemberton SG. 1984. Tracemaking activities of crabs and their environmental significance: the ichnogenus *Psilonichnus*. *Journal of Paleontology*, **58**: 333–350.

Guo WW, Tong JN, Tian L, Chu DL, Bottjer DJ, Shu WC, Ji KX. 2019. Secular variations of ichnofossils from the terrestrial Late Permian-Middle Triassic succession at the Shichuanhe section in Shaanxi Province, North China. Global and Planetary Change, 181: 102978. https://doi.org/10.1016/j.gloplacha.2019.102978

Guo WW, Tong JN, He Q, Hounslow MW, Song HY, Dal Corso J, Wignall PB, Ramezani J, Tian L, Chu DL. 2022. Late Permian–Middle Triassic magnetostratigraphy in North China and its implications for terrestrial-marine correlations. Earth and Planetary Science Letters, 585: 117519. https://doi.org/10.1016/j.epsl.2022.117519

Hasiotis ST, Mitchell CE. 1993. A comparison of crayfish burrow morphologies: Triassic and Holocene fossil, paleo-and neo-ichnological evidence, and the identification of their burrowing signatures. *Ichnos*, **2**: 291–314. https://doi.org/10.1080/10420949309380104

Hasiotis ST, Mitchell CE, Dubiel RF. 1993. Application of morphologic burrow interpretations to discern continental burrow architects: lungfish or crayfish? *Ichnos*, **2**: 315–333. https://doi.org/10.1080/10420949309380105

Hembree D. 2016. Using experimental neoichnology and quantitative analyses to improve the interpretation of continental trace fossils. *Ichnos*, **23**: 262–297. https://doi.org/10.1080/10420940.2016.1202684.

Hounslow MW, Balabanov YP. 2018. A geomagnetic polarity timescale for the Permian, calibrated to stage boundaries. *Geological Society, London, Special Publications*, **450**: 61–103. https://doi.org/10.1144/SP450.8

Hounslow MW, Muttoni G. 2010. The geomagnetic polarity timescale for the Triassic: linkage to stage boundary definitions. *Geological Society, London, Special Publications*, **334**: 61–102. https://doi.org/10.1144/SP334.4

Ji KX, Wignall PB, Tong JN, Yu YY, Guo WW, Shu WC, Chu DL. 2022. Sedimentology of the latest Permian to Early Triassic in the terrestrial settings of the North China Basin: Low-latitude climate change during a warming-driven crisis. *Geological Society of America Bulletin*, **135**: 481-503. https://doi.org/10.1130/b36260.1

Keighley DG, Pickerill RK. 1994. The ichnogenus *Beaconites* and its distinction from *Ancorichnus* and *Taenidium*. *Palaeontology*, **37**: 305–338.

Keighley DG, Pickerill RK. 1995. Commentary: the ichnotaxa *Palaeophycus* and *Planolites*: historical perspectives and recommendations. *Ichnos*, **3**: 301–309. doi: 10.1080/10420949509386400

Klappa CF. 1980. Rhizoliths in terrestrial carbonates: classification, recognition, genesis and significance. *Sedimentology*, **27**: 613–629. https://doi.org/10.1111/j.1365-3091.1980.tb01651.x.

Kraus MJ, Hasiotis ST. 2006. Significance of Different Modes of Rhizolith Preservation to Interpreting Paleoenvironmental and Paleohydrologic Settings: Examples from Paleogene Paleosols, Bighorn Basin, Wyoming, U.S.A. *Journal of Sedimentary Research*, **76**: 633–646. https://doi.org/10.2110/jsr.2006.052

Lu J, Wang Y, Yang M, Zhang P, Bond DPG, Shao L, Hilton J. 2022. Diachronous end-Permian terrestrial ecosystem collapse with its origin in wildfires. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **594**: 110960. https://doi.org/10.1016/j.palaeo.2022.110960

Morrissey LB, Braddy SJ. 2004. Terrestrial trace fossils from the Lower Old Red Sandstone, southwest Wales. *Geological Journal*, **39**: 315–336. https://doi.org/10.1002/gj.991

Pemberton SG, Frey RW. 1982. Trace fossil nomenclature and the *Planolites-Palaeophycus* dilemma. *Journal of Paleontology*, **56**: 843–881.

Rindsberg AK, Kopaska-Merkel DC. 2005. *Treptichnus* and *Arenicolites* from the Steven C. Minkin paleozoic footprint site (Langsettian, Alabama, USA). *Pennsylvanian Footprints in the Black Warrior Basin of Alabama*, **1**: 121–141.

Rodríguez-Tovar FJ, Alcalá L, Cobos A. 2016. *Taenidium* at the lower Barremian El Hoyo dinosaur tracksite (Teruel, Spain): Assessing palaeoenvironmental conditions for the invertebrate community. *Cretaceous Research*, **65**: 48–58. https://doi.org/10.1016/j.cretres.2016.04.013.

Schlirf M, Uchman A, Kümmel M. 2001. Upper Triassic (Keuper) non-marine trace fossils from the Haßberge area (Franconia, south-eastern Germany). *Paläontologische Zeitschrift*, **75**: 71–96. https://doi.org/10.1007/BF03022599

Shu WC, Tong JN, Tian L, Benton MJ, Chu DL, Yu JX, Guo WW. 2018. Limuloid trackways from Permian–Triassic continental successions of North China. Palaeogeography, Palaeoclimatology, Palaeoecology, 508: 71–90. https://doi.org/10.1016/j.palaeo.2018.07.022

Shu WC, Tong JN, Yu JX, Hilton J, Benton MJ, Shi X, Diez JB, Wignall PB, Chu DL, Tian L, Yi, ZX, Mao YD. 2022. Permian−Middle Triassic floral succession in North China and implications for the great transition of continental ecosystems. Geological Society of America Bulletin, 135: 1747–1767. https://doi.org/10.1130/B36316.1

Yu YY, Tian L, Chu DL, Song HY, Guo WW, Tong JN. 2022. Latest Permian–Early Triassic paleoclimatic reconstruction by sedimentary and isotopic analyses of paleosols from the Shichuanhe section in central North China Basin. Palaeogeography, Palaeoclimatology, Palaeoecology, 585: 110726. https://doi.org/10.1016/j.palaeo.2021.110726

Zhu ZC, Liu YQ, Kuang HW, Benton MJ, Newell AJ, Xu H, An W, Ji SA, Xu SC, Peng N, Zhai QG. 2019. Altered fluvial patterns in North China indicate rapid climate change linked to the Permian-Triassic mass extinction. Scientific Reports, 9: 16818. https://doi.org/10.1038/s41598-019-53321-z

Zhu ZC, Liu YQ, Kuang HW, Newell AJ, Peng N, Cui MM, Benton MJ. 2022. Improving paleoenvironment in North China aided Triassic biotic recovery on land following the end-Permian mass extinction. Global and Planetary Change, 216: 103914. https://doi.org/10.1016/j.gloplacha.2022.103914