Structural evolution of nitrogenase enzymes over geologic time

Bruno Cuevas-Zuviría[1,2†], Franka Detemple[3], Kaustubh Amritkar[1], Amanda K. Garcia[1], Lance C. Seefeldt[4], Oliver Einsle[3] and Betül Kaçar[1*]

1. Department of Bacteriology, University of Wisconsin-Madison, Madison, Wisconsin, United States
2. Centro de Biotecnología y Genómica de Plantas, Universidad Politécnica de Madrid (UPM)—Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria-CSIC (INIA/CSIC), Campus de Montegancedo, Madrid, Spain
3. Institute of Biochemistry, University of Freiburg, Freiburg, Germany
4. Department of Chemistry and Biochemistry, Utah State University, Logan, Utah, United States

[†]Affiliation where research was conducted: Department of Bacteriology, University of Wisconsin-Madison, Madison, United States

[*]To whom correspondence should be addressed: bkacar@wisc.edu

Short title: Three billion years of nitrogenase diversity

ABSTRACT

Life on Earth is about 4 billion years old—nearly as old as the planet itself. Over this

immense timespan, living systems and their biomolecules have both adapted to and

driven profound changes in the Earth's environment. Among these, certain critical

enzymes emerged early and have persisted through planetary extremes. Here, we

implement an integrated approach to investigate the structural evolution of nitrogenase,

an ancient and globally essential enzyme responsible for biological nitrogen fixation.

Despite the ecological diversity of its host microbes, nitrogenase retains strict functional

constraints, including extreme oxygen sensitivity, high energy demands, and substrate

availability. We combined phylogenetics, ancestral sequence reconstruction, protein

crystallography and deep-learning based structural prediction to resurrect nearly three

billion years of nitrogenase structural history. This effort represents the first effort to

predict the full set of extant and ancestral structures along the evolutionary tree of a

single enzyme, yielding over 5000 structural models. Our framework lays a foundation

for reconstructing key structural constraints that shape protein evolution and examining

ancient enzymes within the broader context of phylogenetic relationships and

environmental transitions across geological timescales.

INTRODUCTION

For billion years, life and Earth had been intimately linked through globally significant biogeochemical processes. Understanding how these processes evolved requires insight into the molecular machinery that drives them, enzymes, and how their activities have adapted over geological timescales. Tracing the deep evolutionary history of life requires looking beyond the fossil record, which is sparse and incomplete for Earth's earliest eras (Knoll 2003). Genetic sequences, especially those encoding enzymes central to biogeochemical processes, offer an alternative archive of molecular evolution. Advances in evolutionary modeling, ancestral reconstruction, and high-throughput structure prediction (e.g., AlphaFold; Jumper et al., 2021) now make it possible to reconstruct and examine extinct enzymes with unprecedented resolution.

A prime example of an ancient, globally influential enzyme is nitrogenase. Nitrogenase provides access to bioessential nitrogen via the reduction of highly inert atmospheric dinitrogen (N_2) to ammonia (NH_3) (Rucker and Kaçar, 2024). In select microbes called diazotrophs, this pathway is exclusively performed by members of the nitrogenase metalloenzyme family that catalyze the cleavage of the strong $N{\equiv}N$ bond. Age-calibrated, phylogenomic analyses of nitrogenase genes (Parsons et al., 2021) and nitrogen isotopic signatures preserved in the geologic record (Stüeken et al., 2015) both provide evidence for the existence of biological nitrogen fixation more than three billion years ago. Thus, as the sole biological means for fixing essential nitrogen, nitrogenases have underlaid the productivity of the biosphere for most of Earth's history (Falkowski, 1997; Navarro-González et al., 2001; Sánchez-Baracaldo et al., 2014).

Despite the antiquity and ecological importance of nitrogenases, nearly nothing is known about how their enzymatic properties have varied in the past (Rucker and Kaçar, 2024), particularly in response to ancient, global environmental transitions and the distribution of nitrogenase genes across ecologically diverse diazotrophs (Sobol et al., 2025). Notably, nitrogenases are extremely sensitive to oxygen, which degrades the bound metalloclusters required for their enzymatic activity and multimeric structure

(Robson and Postgate, 1980). Nevertheless, these enzymes, which originated in
anaerobic organisms inhabiting an anoxic planet >3.2 billion years ago (Stüeken et al.,
2015), evolved through the progressive oxygenation of Earth's surface environment
(beginning ~2.4 to 3 billion years ago (Lyons et al., 2024, 2014) and genetic acquisition
by aerobic organisms (Boyd et al., 2015). Though several regulatory mechanisms and
accessory proteins that today shield nitrogenase from oxygen have been identified
(Dixon and Kahn, 2004; Robson and Postgate, 1980; Schlesier et al., 2016; Takimoto et
al., 2022), molecular features associated with oxygen protection adaptation have not
been investigated within the nitrogenase enzyme itself. Another outcome of Earth
surface oxygenation is the spaciotemporal shift in the environmental bioavailability of
redox-sensitive trace metals used by nitrogenases. Specifically, molybdenum (Mo),
vanadium (V), and iron (Fe) (Johnson et al., 2021; Moore et al., 2017) are proposed to
have shaped the diversification of the variably metal-dependent Mo-, V-, and Fe-
nitrogenase isozymes (Anbar and Knoll, 2002). The emergence of novel features within
the nitrogenase enzyme has been suggested as necessary for maintaining biological
nitrogen fixation amid global marine geochemical shifts (Cuevas-Zuviría et al., 2024),
yet little is known about the molecular changes and constraints that accompanied their
diversification and evolution across time. This knowledge gap limits our ability to link
molecular innovations to environmental transitions in nitrogenase evolution, obscuring
the principles behind an enzyme capable of catalyzing one of nature's most challenging
reactions.
A window into the evolution of nitrogenase function can be generated by probing the
historical protein structure space representative of this ancient enzyme family. In this
study, we present a combinatorial approach for reconstructing ancient nitrogenase
structures and tracking their evolution across Earth history. We draw from a
phylogenetic dataset of extant and inferred ancient nitrogenase proteins to reconstruct
more than three billion years of nitrogenase structural evolution. Our study combines
ancestral sequence reconstruction with massive machine-learning-enabled structure
prediction of present-day and ancestral nitrogenases, as well as protein crystallographic
investigations of select, functionally characterized ancestors. We thus present the first
effort to predict all extant (770) and ancestral (4,608) enzyme structures across the

nitrogenase evolutionary tree, and interpret key observations across an enzyme's
structural history within the context of planetary environmental change.

RESULTS

Structural prediction of ancient nitrogenase enzymes across time
We began our structural investigation of nitrogenase evolutionary history by conducting
a large-scale structure prediction analysis, encompassing 5378 protein structures, a
more than threefold increase compared to available nitrogenase structures in the PDB.
We then analyzed our phylogenetic dataset to identify notable structural changes. Our
dataset includes representatives of all three known nitrogenase isozymes, which are
differentiated by the metal composition of their active-site metallocluster: Nif (encoded
by *nifHDK* genes; incorporates an iron-molybdenum cofactor, "FeMoco"), Vnf (encoded
by *vnfHDGK* genes; incorporates an iron-vanadium cofactor, "FeVco"), and Anf
(encoded by *anfHDGK* genes; incorporates an iron-only cofactor, "FeFeco"). H-, D-, and
K-subunit sequences were analyzed for each nitrogenase homolog (G-subunit
sequences unique to Vnf and Anf were not included in the analysis; see **Materials and**
Methods). In all nitrogenase isozymes, H-subunits form a homodimeric reductase
component ("HH") that transiently associates with and transfers electrons to a catalytic
core (Einsle and Rees, 2020). This catalytic component is a heterotetramer composed
of two D- and two K-subunits ("DDKK" or $\alpha_2\beta_2$) and contains the active site
metallocluster where N_2 is bound and reduced to NH_3.

The topology of the nitrogenase phylogenetic dataset (Garcia et al., 2023) used in the
present study reveals five major clades: Nif-I, Nif-II, Nif-III, Vnf, and Anf (**Figure 1A**).
Both Vnf and Anf homologs are nested within the Nif-III clade. Aside from sequence-
level homology of their constituent proteins, clades are also distinguished by associated
features including metal dependence of the nitrogenase enzyme, taxonomy, host
ecology, and complexity of the nitrogenase-associated gene network (Cuevas-Zuviría et
al., 2024). In extant diazotrophs, nitrogenases require a suite of associated genes for
their regulation, assembly, and maintenance (Campo et al., 2022). The content and

complexity of this cellular network for nitrogen fixation tends to be characteristic of the
major clades and host ecologies of nitrogenases. For example, Nif-I nitrogenases are
primarily hosted by aerobic or facultatively anaerobic diazotrophs that have among the
largest number of nitrogenase-associated genes (Boyd et al., 2015). By comparison,
Nif-II and Nif-III nitrogenases are hosted nearly exclusively by anaerobic diazotrophs
and have comparatively smaller nitrogen fixation gene networks (Cuevas-Zuviría et al.,
2024; Garcia et al., 2020). Vnf and Anf isozymes, though sharing certain associated
genes with Nif, are also supported by distinct, dedicated gene clusters (Garcia et al.,
2022, 2020). We hypothesized that these differences would manifest in correlated
structural differences across the enzyme family.

Figure 1. Overview of the methodological pipeline for massive protein structure
prediction and experimental crystallographic analysis of extant and ancestral
nitrogenase enzymes.   **A)** Nitrogenase phylogeny built from concatenated
Nif/Vnf/AnfHDK protein sequences. Clades are labeled according to the nomenclature
used by Raymond et al. 2004. (Nif-I: Group I Nif, Nif-II: Group II Nif, Nif-III: Group III Nif).
Nif, Vnf, or Anf homologs from select model organisms are labeled with dashed lines
(*Avin*: *Azotobacter vinelandii*, *Cpas*: *Clostridium pasteurianum*, *Kpne*: *Klebsiella*

pneumoniae, *Mace*: *Methanosarcina acetivorans*, *Rpal*: *Rhodopseudomonas palustris*).

Anc1A/B and Anc2 ancestors targeted for crystallographic analysis are labeled with

stars. **B)** Graphical overview of the pipeline for nitrogenase protein structure prediction

and crystallization (see Materials and Methods for further details). Colored rectangles

correspond to protein sequences for H, D, and K subunits. For each ancestral node,

protein structures were predicted for the most likely ancestral sequence ("ML") and five

alternative sequences ("Alt") reconstructed based on the site-wise posterior probability

distributions in the ancestral sequence. Ancestors Anc1A and Anc2 hybrid enzymes

were crystallized containing an ancestral NifD subunit and WT NifH and NifK subunits

(WT subunits indicated by lighter color). All predicted structures are publicly available at

https://nsdb.bact.wisc.edu.

We employed a high-throughput, AlphaFold-based computational pipeline to generate

multimeric HH and DDKK structural predictions for each nitrogenase homolog in our

dataset (**Figure 1B**). For each of the 385 extant targets, a single structure was

predicted. For each of the 384 ancestral targets, six structures were predicted, sourced

from the single most likely ancestral protein sequence and five alternative ancestral

sequences generated by substitutions at ambiguously reconstructed sites (see

Materials and Methods). In total, 2,689 unique extant and ancestral nitrogenase

variants were targeted.   All structures were generated in approximately 805 hours,

including GPU computations and MMseqs2 alignments performed using two different

protocols: one for extant or most likely ancestral sequences, and another for ancestral

variants. HH and DDKK structures were predicted independently, resulting in a total of

5378 individual nitrogenase protein structures generated from our pipeline. For

comparison, the Protein Data Bank currently contains structures of nitrogenase

homologs (from seven different organisms, spanning 15 HH and 24 DDKK 104

structures) (**Table S1**).

We found that all nitrogenase structures were predicted with high confidence, assessed

by the predicted Local Distance Difference Test (pLDDT) (Jumper et al., 2021; Mariani

et al., 2013). pLDDT scores ranged between ~82 and 98 (**Figure S1**). Within this range,

confidence was lowest for H-subunit structures (median pLDDT ≈ 88) and highest for K-
subunit structures (median pLDDT ≈ 98). Furthermore, we built a public web database,
"Nitrogenase Structural DB," (accessible at https://nsdb.bact.wisc.edu), to host all
nitrogenase structures predicted in this study. Structures are visualizable, downloadable
(PDB format files), and can be searched by sequence or node ID (annotated on the
nitrogenase phylogeny in **Figure S2**), NCBI taxon ID, and species name.
199

Ancient structural insertion events distinguish major nitrogenase clades
We observe that predicted nitrogenase structures have sufficient variations to resolve
phylogenetic relationships. To assess structural variation, we calculated the Root Mean
Squared Deviation (RMSD) using US-Align (Zhang et al., 2022). Although in this case,
this metric indicates overall small differences (< 3 Å) (**Figure 2A**), it still strongly
correlates with sequence identity (**Figure 2B**), indicating that even minor structural
variations can recapitulate sequence-based phylogenetic distinctions. Hierarchical
clustering of nitrogenase structures by pairwise RMSD largely reproduces the major
clades (Nif-I, Nif-II, Nif-III, Vnf/Anf) (**Figure 2C**). To confirm that this correlation between
sequence identity and structure is not simply an artifact resulting from alignment length
bias, we applied two complementary metrics: $RMSD_{100}$ (Carugo and Pogor 2001) and
TM-score (Zhang and Skolnick, 2004). $RMSD_{100}$ is a normalized variant of RMSD
designed specifically to correct structural alignment length bias. The results obtained
with these alternative metrics (**Figure S3**) were similar to the results of **Figure 2**,
differing primarily by a scaling factor of approximately one-third.
215

TM-score enables a global comparison of structural similarity ranging between 1
(identical) and 0 (no similarity). All DDKK structures have relatively high pairwise TM-
scores (> .84); TM-score > 0.5 indicates comparable folding (Xu and Zhang, 2010) and
hierarchical clustering of nitrogenase structures by pairwise TM-scores largely
reproduces the major clades (Nif-I, Nif-II, Nif-III, Vnf/Anf) observed in the protein
sequence phylogeny (**Figure S4**).
222

Upon finding a correlation between structure similarity and phylogenetic relationships,
we developed a method to sample the inherent uncertainty of ASR outputs by
generating sequences with alternative residues at ambiguous positions (see **Materials**
and Methods). We predicted the resulting structures and compared the variability of
these outputs against the maximum-likelihood sequence. **Figure S5A** shows the overall
distribution of RMSDs for randomized variants against the maximum likelihood variant.
RMSD for most divergent variants fall within two angstroms, indicating very high
structural similarity. As shown, the RMSD does not correlate with the distance to the
tree's root, indicating no systemic relationship between structural similarity and tree
depth (**Figure S5B**). Inferred structures, even those derived from deep nodes within the
phylogenetic tree, are robust against the statistical uncertainty of ancestral sequence
inference. Finally, we computed the all-vs-all RMSD at the DDKK active sites (FeMoco
and 8Fe7S; **Figures S6A** and **S6B**). The results indicate that the active sites have high
structural conservation, especially around the 8Fe7S cluster (RMSD < 0.5Å, **Figure**
S6C), which could be expected given the tight coordination of the metal cofactor with its
surrounding cysteines. It is worth noting that AlphaFold2 does not use information about
metal cofactors when building the model.
240

Figure 2. Global analyses of nitrogenase DDKK sequence and structural diversity.
A) Root Mean Squared Deviation (RMSD) distribution on paired nitrogenase alignments across extant and ML-ancestor pairs. **B)** Sequence identity and structural similarity (quantified by root mean square deviation (RMSD) of aligned predicted structures) distribution of paired nitrogenase alignments. **C)** Hierarchical clustering of predicted nitrogenase structures based on structural similarity (RMSD). Each tile in the heatmap corresponds to the RMSD between two nitrogenase structures.

Our global comparisons of predicted nitrogenase structures suggest that, despite relatively high conservation, certain structural distinctions between major clades evolved early and persisted in the extant descendants. We examined our dataset to identify these distinctions and map them to the evolutionary trajectory on the nitrogenase family history.

The structural predictions identified three insertion events that distinguish nitrogenase

variants. These insertions likely contribute significantly to the phylogenetic signal

encoded within structural features (**Figure 3A**): a D-subunit C-terminus extension (~60

residues) unique to Anf (**Figure 3B**), a K-subunit N-terminus extension (~8 to 40

residues) unique to Nif-I (**Figure 3C**), and a D-subunit insertion (~55 residues) unique to

Nif-II (**Figure 3D**). The relationships between these insertion events and the major

nitrogenase clades are also associated with the significant differences in host ecology

and nitrogen fixation genetics across clades.

We found that the K-subunit N-terminus is predominantly found in oxygen-tolerant,

aerobic or facultatively anaerobic diazotrophs, whereas the D-subunit insertion of Nif-II

is mostly found in anaerobic diazotrophs. Our ancestral reconstructions constrain the

initial appearance of each insertion to the branch leading to the last common ancestor

of the clade. Both D-subunit insertions appear within a single branch and thereafter

remain relatively conserved in length. By contrast, we observed that the K-subunit N-

terminus insertion evolved progressively in length (**Figure 3B, 3D**) through several

deep-time ancestors of the Nif-I clade (**Figure 3C).** The elongation of this insertion

recapitulates trends in the K-subunit N-terminus lengths of extant Nif-I homologs, where

early-diverged lineages have short (~8 residue) extensions and later-diverged lineages

have longer (~40 residue) extensions. The extent to which the N-terminus of the K-

subunit affects functional importance in relation to oxygen demand warrants further

empirical investigation. As a side observation, our analysis detected no major deletion

events in the evolutionary history of nitrogenase. Oldest ancestors (including the last

common ancestor of the entire nitrogenase family) and Group III nitrogenases are

among the shortest in sequence length and do not contain any of the insertions

described above.

284
Figure 3. Nitrogenase structure variation in a phylogenetic context. A) Nitrogenase
protein phylogeny. Branches and ancestral nodes corresponding to structural insertion
events, as well as representative extant variants conserving those insertions, are
highlighted and/or labeled. Clade and node colors correspond to the subunit for which
an insertion is observed (i.e., blue for the D subunit, red for the K subunit). **B)**
Elongation of the NifD C-terminus coincident with the origin of the Anf clade. **C)**
Progressive elongation of the NifK N-terminus through the early evolution of the Nif-I
clade. **D)** Insertion within NifD coincident with the origin of the Nif-II clade. **B-D)** All
visualized structures are predicted unless otherwise specified with the corresponding
Protein Data Bank identifier. Bound G- and H-subunit structures were not predicted
together with the NifDK structures and are thus indicated with an asterisk. The binding
positions of the G- and H-subunit structures are inferred based on alignment with PDB
8BOQ (Trncik et al., 2023) and PDB 1M34 (Schmid et al., 2002), respectively.

The locations of these insertions within the nitrogenase structure and their encoding
genes provide insight into their possible functional contributions. First, all insertions are
exposed at the surface of the DDKK protein complex (**Figures 3B-D**), and all comprise
varying proportions of helix or loop regions in both extant and ancestral structures. The
position of the D-subunit C-terminus extension (**Figure 3B**) and the K-subunit N-

terminus extension (**Figure 3C**) is similar. Both line the edges of the interface between D- and K- subunit proteins, possibly stabilizing the complex. It is possible that the N-terminus of NifK wraps around the entry point of small-molecules (Barney et al., 2009; Morrison et al., 2015), and help in reducing access for oxygen in Nif-I hosted by aerobes. The D-subunit insertion in Nif-II instead protrudes near the inferred HH binding site. Though no crystallographic structures of a bound HH-DDKK nitrogenase complex are available for Nif-II homologs (and we did not predict bound structures here), the Nif-II D-subunit insertion is likely capable of H-subunit interactions based on alignment with the binding site in Nif-I nitrogenases (**Figure 3D**).

Phylogenetic trends of nitrogenase DDKK complex structures

Following the identification of structural innovations in the history of nitrogenases, we investigated evolutionary trends in calculated structural attributes for DDKK structures, including surface properties and intermolecular contacts (**Figure S7A**). Broadly, we found that many structural features carry clade-specific trends and/or are associated with the presence of the insertions described above. For example, total surface area (summed over each protein subunit in the DDKK complex) is distinct between Nif-I, Nif-II, and Nif-III/Vnf/Anf, and greatest in the Nif-II clade (**Figure S7B**). Trends in surface area are likely driven in part by the presence and conformation of the surface-exposed insertions. We also examined attributes related to protein-protein interactions within DDKK. In all cases, we found that trends in these attributes were influenced most strongly by the presence of the K-subunit N-terminus extension in the Nif-I clade. Nif-I homologs (and particularly those with longer K-subunit extensions) have the largest number of intersubunit molecular contacts, and intersubunit binding affinities, as calculated with Prodigy (Vangone and Bonvin 2015) (**Figure S7C, Figure 4B**). These patterns support our hypothesis that the K-subunit N-terminal extension (**Figure 3C** and **4C**) contributes positively to DDKK complex interaction strength and stability. By contrast, homologs with the least number of intersubunit contacts and weakest binding affinities are Nif-III nitrogenases, which notably contain none of the described insertions. We also observed an increase in the total surface area of the nitrogenase complex

within the Nif-II group (**Figure S7B**). The only structural attribute that we found correlated with ancestor age was the proportion of charged residues across DDKK proteins (**Figure S7E**). Accordingly, we observed a weak negative correlation between phylogenetic distance from the root and proportion of charged residues (Spearman R ≈ -0.261). We did not detect similar trends for other attributes. Rather, the attributes of nitrogenase ancestors largely resemble those of their descendants in a clade-specific manner. These results reinforce the idea that major structural distinctions were set early in nitrogenase history and conserved in respective clades.

Figure 4. Phylogenetic patterns of nitrogenase structural attributes calculated across DDKK proteins. A) Node taxonomy overview. **B)** Binding affinity prediction for extant and ancestral nitrogenase nodes mapped to the nitrogenase phylogeny. **C)** Structure overview of the D and K subunit interactions around the N-terminal insertion in Nif-I nitrogenases. Structure corresponding to *Azotobacter vinelandii* NifDK (PDB code: 3U7Q). Note: We reduced the number of displayed phylogenetic nodes to mitigate visual overcrowding, refer to **Supplementary Figure S7** for a complete visualization of all nodes.

Crystalized nitrogenase ancestors confirm evolutionary substitutions at the HH-DDKK interface

In this work, we created a large dataset of Nitrogenase structures, expanding upon the limited structural reference data that the prediction methods rely on. Therefore, we continued our structural investigation of nitrogenase evolutionary history by crystallographic characterization of select nitrogenase ancestors which helped us assess the accuracy of the structure prediction methods. Three previously characterized ancestral variants were targeted from phylogenetic nodes within the evolutionary lineage of *A. vinelandii* Nif (wild-type, "WT") (**Figure 5A**) (Garcia et al., 2023): Anc1A, Anc1B and Anc2. Anc1A and Anc1B are the same age, whereas Anc2 is reconstructed from an older phylogenetic node. WT, Anc1, and Anc2 belong to a Mo-nitrogenase clade, formerly designated as 'Group I' by (Raymond et al., 2004). This clade includes homologs from various aerobic and facultatively anaerobic organisms, such as proteobacteria and cyanobacteria (Garcia et al., 2023). An estimated maximum age of approximately 2.5 billion years for Group I nitrogenases —and consequently for Anc1 and Anc2— is inferred from the timing of the GOE (Garcia et al., 2023; Lyons et al., 2014). As an additional distinction, Anc1A and Anc2 contain an ancestral D-subunit protein complexed with WT H- and K-subunit proteins. Anc1B, by contrast, is composed of ancestral H-, D-, and K-subunit proteins. Ancestral variants preferentially bind to the ATP cofactor (Harris et al., 2024) and exhibit the same general mechanism for N_2 binding and reduction conserved across all known nitrogenases to date, but with

decreased activity for reduction of N_2 or other substrates (Garcia et al., 2023). Thus, the

sequence differences between ancestors and WT are sufficient to generate phenotypic

changes.

Ancestral nitrogenase DDKK proteins were isolated from genomically modified strains of

A. vinelandii (Garcia et al., 2023), purified by chromatographic methods, and crystallized

under anoxic conditions (**Figure S8**) (see **Materials and Methods**). We found that

ancestral nitrogenases required different conditions for crystallization relative to that

required for modern proteins. Specifically, all ancestors crystallized at pH 6.5, which is

unusually acidic for nitrogenase (Peters et al., 1997; Spatzal et al., 2011). In addition, all

ancestral proteins crystallized in space group $P\,2_12_12_1$, which differs from the space

groups for the same proteins in *A. vinelandii* NifDK (i.e., WT; space group $P\,2_1$),

Clostridium pasteurianum Nif (space group $P\,2_1$), and *Klebsiella pneumoniae* NifDK

(space group $C\,2$). We observed one ancestral DDKK protein per asymmetric unit. All

three ancestors showed anisotropic diffraction: 2.65 to 2.93 Å for Anc1A, 1.82 to 2.8 Å

for Anc1B, and 1.82 to 2.42 Å for Anc2. All structures were solved by molecular

replacement using WT *A. vinelandii* (PDB 3U7Q).

As in extant nitrogenase proteins, each ancestral D- or K-subunit structure exhibits

three Rossmann-fold domains (**Figure 5B**). The P-cluster, an 8Fe7S metallocluster that

mediates electron flow to the active site, is located on the pseudo-twofold axis relating

the structurally similar D- and K-subunits and was modeled in the all-ferrous P^N-state.

Both the P-cluster and the active-site cofactor, FeMoco, were modeled with full

occupancy. Global structural differences between the ancestors and WT *A. vinelandii*

DDKK are relatively minor, and within the range of variation for available extant

structures. Among all ancestors, root mean square deviations (RMSD) for all atom

positions are largest between Anc1B and WT DDKK (0.36 Å; compared to RMSD of

0.50 Å between *A. vinelandii* and *K. pneumoniae* DDKK). This observation may be

explained by the larger number of amino acid substitutions between the fully ancestral

Anc1B complex and WT, whereas only the D-subunit contains substitutions in the Anc1A

and Anc2 hybrid complexes (**Figure 5C**). Among the two hybrid ancestors, the RMSD

between Anc2 and WT (0.24 Å) is smaller than that between Anc1A and WT (0.31 Å). Furthermore, we observe that the position of the twofold symmetry axis relating the two DK heterodimers of a single MoFe protein shifts slightly between Anc1B and WT (**Figure S9**). The same was previously observed in VFe protein (Sippel and Einsle, 2017) and FeFe protein (Trncik et al., 2023) and may reflect slight differences in the interface between the two copies of NifK.

Figure 5. Crystal structures of targeted ancestral nitrogenases. A) Nitrogenase protein phylogeny from which nitrogenase ancestral proteins were reconstructed and crystallized for structural characterization. Major clades are labeled following (Garcia et al., 2023). **B)** Residue-level RMSD between the crystallized ancestral and wild-type (*A. vinelandii*) NifDK/NifH structures. **C)** Spatial distribution of ancestral amino acid substitutions relative to WT. Crystallized protein complexes for Anc1A and Anc2 contain ancestral NifD and WT NifH and NifK. Therefore, ancestral substitutions are only in the NifD subunit for these structures. **D)** Ancestral amino acid substitutions within the NifH-NifDK interface of Anc1B. Bound positions of NifH are inferred by alignment with either the nucleotide-free (PDB 2AFH (Tezcan et al., 2005)) or MgATP-bound (PDB 7UT8 (Rutledge et al., 2022)) structures. **E-G)** Close views of specific, ancestral amino acid substitutions that are inferred to impact NifH-NifDK interactions.

Figure 6. Comparison between crystallographic/cryo-EM and AlphaFold-predicted nitrogenase structures. A) RMSD between crystallographic/cryo-EM structures available in the Protein Data Bank (PDB) and AlphaFold predictions. Each data point represents a comparison between one of five AlphaFold replicates and each PDB structure. **B)** RMSD between experimental structures obtained in this work and their AlphaFold predictions. Each data point represents a comparison between one AlphaFold replicate and the experimental structure **C**) Cα- Cα distance between the Anc1A structural prediction and its crystallographic reference, mapped on the

nitrogenase structure through a colormap. Blue residues denote larger distances C□-

C□ distance, which indicate lower prediction fidelity on these regions.

In our previous study, we found that some of the ancestral nitrogenases had decreased

N_2-reduction activity *in vitro* relative to WT (Garcia et al., 2023). The generation of

crystallographic structures for these ancestors provide an opportunity to examine any

structural features that might be responsible for the decrease in activity in these

variants. Given that most ancestral DDKK substitutions were not close to

metalloclusters, we hypothesized that those situated at the DDKK-HH interface might

have modified the activity of the ancestors by impeding protein-protein interactions that

are necessary for electron transfer to the nitrogenase active site, as well as cofactor

loading and complex maturation (Burén et al., 2020). Snapshots of crystal structure

interface, highlighting mutations and residue interactions are shown in **Figure 5E-G**.

Notably, a non-conserved Nif-D α-helix (positions 165-174 in Anc1B) lies within

interacting distance of HH. This helix contains several substitutions across all ancestors,

albeit the positioning of polar amino acids differs among Anc1A, Anc1B, and Anc2.

Herein, the D-subunit S165A substitution is consistently present across all ancestors

and may influence interactions with HH S175 in the ATP-bound complex, potentially

reducing stability (**Figure 5E)**. Additionally, two unique single-point substitutions in the

ancestral K-subunit—K211N and G307N, specific to Anc1B—are likely to impact

interactions with HH E111 in the ATP-bound state (**Figure 5G**). Positioned centrally

within the DDKK-HH interface, these substitutions alter the electrostatic surface of HH.

In Anc1B, the substitution HH S177G is located near the interface with NifDK, with

G307N in Anc1B NifK identified as a potential interaction partner (**Figure 5F**).

Predictions for nitrogenase variants had high similarity (average RMSD < 1.4 Å) to

experimentally determined structures (**Figure 6A**). This was also true for the ancestral

nodes selected for crystallographic analysis in this study (**Figure 6B, Figure 6C**),

intriguingly, none of which were included in the structural database used for AlphaFold

training. Beyond the limited number of substitutions within the interfaces of Anc1A/B and
Anc2, the changes in the ancestral enzymes were distributed quite evenly across the
protein subunits.

DISCUSSION

In this study, we combined targeted crystallization experiments of ancestral nitrogenase
enzymes with large-scale 3D structure predictions spanning both present-day and
ancestral nitrogenase diversity. Building on the recent expansion of predicted structures
for extant proteins, this work marks a pioneering effort to apply these methods to
explore the ancestry of an entire enzyme family. The three crystallographic and 5,378
predicted structures generated in this study represent approximately a 50-fold increase
in available nitrogenase complex structures, and importantly includes an extensive
sampling of historical structural variation.

Both crystallographic and predicted structure of nitrogenase ancestors highlight the
long-term evolutionary conservation of core structural features, consistent with
previously characterized preservation of mechanistic aspects in nitrogenase catalysis
(Garcia et al., 2023, Harris et al. 2024). Nevertheless, historical structural variation
sufficiently distinguishes major nitrogenase clades independent of protein sequence
information and this phylogenetic variation is driven primarily by different insertion
events that occurred early in nitrogenase evolutionary history. For example, we show
that the extension of the K-subunit N-terminus emerged during the early diversification
of Group I nitrogenases coinciding with the proliferation of nitrogenases into oxygen-
tolerant bacterial lineages and the planetary rise in atmospheric oxygen.

Nitrogenase structural features that might be associated with increased compatibility
between the oxygen-sensitive nitrogenase enzyme and oxygen-tolerant host organisms
are of significant interest given efforts to transfer N-fixation genes to crops (Vicente and
Dean, 2017). Furthermore, if such structural features are in fact required for
compatibility with oxygen tolerant hosts, their emergence would represent a pivotal

transition in Earth history given the importance of aerobic N-fixers for present-day global nitrogen cycling. For example, *Trichodesmium*, a single genus of oxygenic phototrophic cyanobacteria, today contributes 60-80 Tg of the total 100-200 Tg fixed N produced annually in modern oceans (Bergman et al., 2013). It was also proposed that the O_2 sensitivity of nitrogenase limited the extent of oxygen production by primary producers (Mrnjavac et al., 2024).

The occurrence of the NifK N-terminus extension in oxygen tolerant diazotrophs, combined with increased inter-subunit interactions and affinity predicted by our *in-silico* analyses, points to a possible functional role in response to the planetary rise of oxygen. Aerobic diazotrophs today utilize a variety of oxygen protection strategies for N-fixation. These include physical separation of nitrogenases into anaerobic, differentiated cells, temporal decoupling of N-fixation from oxygen production in cyanobacteria, and increased rates of aerobic respiration (Mus et al., 2019; Robson and Postgate, 1980). Further, the Shethna protein II was found to shield nitrogenase from temporary oxygen exposure by forming a complex with nitrogenase proteins and rendering the enzyme inactive yet oxygen-tolerant (Schlesier et al., 2016). Nevertheless, evolutionary modifications to the nitrogenase enzyme itself that improve its compatibility in aerobic diazotrophs have not yet been identified. The NifK N-terminus extension discussed here may play a direct role, by stabilizing oxygen-sensitive, intersubunit protein interactions within the nitrogenase enzyme. Alternatively, it might play an indirect role by shaping potential interaction sites with accessory proteins (like the Shethna protein II) that improve oxygen protection. Notably, while the NifK extension may contribute to oxygen protection, it is absent in Vnf and Anf nitrogenases, which are nevertheless present in aerobic and facultative anaerobic diazotrophs. Given the dramatic environmental shifts brought by the GOE, it remains uncertain to what extent these emergent changes were beneficial at the time of their appearance. For example, within the framework of constructively neutral evolution, these changes may have initially been neutral, only later acquiring adaptive value as environmental conditions or biological contexts shifted (Brunet and Doolittle, 2018).

Collectively, we identify major structural transitions in the history of nitrogenase
enzymes that can serve as a foundation for future experimental studies examining the
interplay between cellular and environmental factors that impact enzyme evolution.
Nonetheless, the insights gained by these structure prediction methods carry inherent
uncertainty. For example, accurate active-site side chain prediction would require
incorporation of metal-cofactors into the active site, which is currently missing in our
AlphaFold2-based pipeline. Future versions of AlphaFold, with enhanced capacity to
capture finer-scale features (e.g., side chain structures), may enable more detailed
estimates and reveal subtle structural and historical variations.
546
Our study shows that, while nitrogenase retains a conserved multimeric core structure,
it has also evolved novel structural features aligned with key environmental transitions
in Earth's history. Subtle, modular changes, particularly those distal to the active site,
likely played a crucial role in the enzyme's adaptation and long-term persistence of over
geologic time. These findings suggest that transient regulatory adaptations in response
to global shifts may have significantly influenced the trajectory of protein evolution.
Reconstructing the lost histories of ancient metabolisms (Kaçar 2024), including their
structural transformations, opens new avenues for exploring how climatological and
ecological forces shape molecular evolution.
556
557

MATERIALS AND METHODS

Nitrogenase ancestral sequence reconstruction, selection, and resurrection in *A. vinelandii*

All extant and ancestral nitrogenase protein sequences were drawn from a previous nitrogenase sequence dataset (Garcia et al., 2023). Briefly, 385 nitrogenase sequences and 385 outgroup sequences were curated from the NCBI nr database by BLASTp for phylogenetic tree building and ancestral sequence reconstruction. For ancestors Anc1A and Anc2, aligned sequences (MAFFT v7.450 (Katoh and Standley, 2013) were trimmed (trimAl v1.2 (Capella-Gutiérrez et al., 2009)) and both tree reconstruction and ancestral sequence inference were performed by RAxML v8.2.10 (Stamatakis, 2014) with the LG + G + F evolutionary model. Model selection was performed by ModelFinder (Kalyaanamoorthy et al., 2017) in IQ-TREE v.1.6.12 (Nguyen et al., 2015). For ancestor Anc1B, the untrimmed sequence alignment was used for tree reconstruction by RAxML and ancestral sequence inference by PAML v4.9j (Yang, 2007) using the LG + G + F model. This second analysis was conducted due to concerns that RAxML v8.2 does not perform full, marginal ancestral sequence reconstructed as described by (Yang et al., 1995). Anc1A and Anc1B are equivalent in their set of descendants and, despite the difference in reconstruction methods, have NifD proteins with 95% identity (**Figure S10**).

In addition to the ancestral sequences described above, we calculated the statistical uncertainty of ASR by generating five alternative variants for each ancestor in the nitrogenase phylogeny. These include: one variant that contains the second most probable residue at each ambiguous position (probability of most probable ancestral state < 0.7) ("*altall*") and another four variants selected by randomly sampling the ASR posterior probability distribution at each ambiguous site ("*alt2*," "*alt3*," "*alt4*," "*alt5*"). The *altall* and *alt2-5* sequences correspond to the alternative sequences considered in **Figure 1**.

Engineering of *A. vinelandii* strains was performed as previously described, replacing
the native WT *nifD* gene with an ancestral variant (Garcia et al., 2023). A. *vinelandi*i WT
('DJ'), DJ2278 (ΔnifD::KanR), and DJ2102 (Strep-II-tagged WT NifD were generously
provided by Dennis Dean (Virginia Tech) for strain engineering (Santos, 2018).
Following genomic integration of ancestral nitrogenase genes into *A. vinelandii*,
transformants were passaged at least three times to ensure phenotypic stability prior to
storage at –80 °C in phosphate buffer containing 7% DMSO. Verification of the
engineered strains was performed by Sanger sequencing of the *nifD* region as well as
whole genome sequencing.

Expression and purification of nitrogenase proteins *A. vinelandii*

Cells from DMSO stock were directly grown in a pre-culture using liquid Burk-medium
containing ammonium sulfate at 37 °C and 180 rpm until an optical density at 600 nm of
at least 1.5 was reached. Further a second pre-culture without ammonium-sulfate was
grown under the same conditions and used to inoculated 500 mL main cultures at 30 °C
and 180 rpm. Cells were harvested when an optical density at 600 nm of 1.7-2.5 was
reached.

Anc1A: Cells were grown in a 10 L Eppendorf BF120 bioreactor at 30 °C in Burk-
medium containing urea as the fixed N source. Dissolved oxygen (DO) was held at 20%
using lab air at a constant 10 SLPM and cascade control of agitation (100 – 500 RPM)
and supplemental O_2 (0 – 30%). At an optical density at 600 nm of ~8 the cells were
pelleted by centrifugation and resuspended in Burk-medium without a fixed N source to
induce expression of nitrogenase. Cells were allowed to express for 4 hours with DO
held at 10% with the same air flow and cascade control before harvesting.

Isolation of MoFe proteins: MoFe protein is oxygen sensitive therefore all consecutive
steps were conducted either in an anoxic chamber (Coy Laboratories) under 95% N_2
and 5% H_2 atmosphere or using modified Schlenk techniques under N_2 flow. Anc1
contains two additional crystallographic forms, Anc1A and Anc1B; the two are the same

age, and Anc1A contains an ancestral D-subunit protein complexed with WT H- and K-
subunit proteins, Anc1B is by contrast is composed of ancestral H-, D-, and K-subunit
proteins. They were isolated as follows:

Anc1B (MoFe and Fe protein): Cell pellets were resuspended in lysis buffer (50 mM
Tris/HCl at pH 7.4, 2.5 mM $Na_2S_2O_4$) and opened at 1500 bar in a SUP Maximator HPL-
6 at 2 °C under anoxic conditions. The cell free extract was centrifuged at 87,000 x g at
4 °C for 60 minutes, filtered through a 0.45 μM syringe filter (Filtropur) and were loaded
onto two equilibrated 5 mL HiTrap Q HP columns. After washing with 12.5 % elution
buffer (50 mM Tris/HCl at pH 7.4, 1000 mM NaCl, 2.5 mM Na2S2O4), the amount of
elution buffer was increased in 2.5 % increments up to 35 %, followed by steps at 40
and 50 %. MoFe and Fe protein eluted after 300 mM NaCl and 325 mM NaCl
respectively. Step elution with 40 % elution buffer was used to concentrate. The proteins
were loaded separately onto a size exclusion chromatography (Superdex S200, GE
Healthcare) equilibrated with 20 mM Tris/HCl at pH 7.4, 200 mM NaCl, 2.5 mM
$Na_2S_2O_4$, concentrated (Vivaspin 20, 50 or 30 kDa MWCO, Sartorius) and flash frozen
in liquid nitrogen. A second size exclusion chromatography step (Superdex 200
Increase 10/300 GL) was carried out with MoFe protein, which was subsequently
concentrated, and flash frozen in liquid nitrogen.

Anc1A: Cell pellets were resuspended in lysis buffer (50 mM Tris/HCl pH 7.9, 20%
glycerol (v/v), 500 mM NaCl, and 2 mM $Na_2S_2O_4$). Cells were lysed with a Nano DeBee
45-2 High Pressure Homogenizer and then ultra-centrifugated to produce cell free
lysate. Cell free lysate was loaded onto an equilibrated (50 mM Tris/HCl pH 7.9, 500
mM NaCl, and 2 mM $Na_2S_2O_4$) 20 mL StrepTactinXT 4Flow column. The column was
washed with 2 column volumes of equilibration buffer, followed by elution of the protein
in a single fraction with elution buffer (50 mM Tris/HCl pH 8, 150 mM NaCl, 50 mM
biotin, and 2 mM $Na_2S_2O_4$). Protein was then loaded on an equilibrated (20 mM Tris/HCl
at pH 7.4, 200 mM NaCl, 2.5 mM $Na_2S_2O_4$) size exclusion chromatography (Superdex
200 Increase 10/300 GL). Subsequently it was concentrated, and flash frozen in liquid
nitrogen.

Anc2: Cell pellets were resuspended in lysis buffer (20 mM Tris/HCl at pH 7.4, 200 mM
NaCl, 2.5 mM $Na_2S_2O_4$) and opened, centrifuged and filtered following the Anc1B
protocol. The cell free extract was loaded onto a with lysis buffer equilibrated
Streptavidin XT 4 Flow column. Through addition of 50 mM Biotin to the lysis buffer
Anc2 eluted. Size exclusion chromatography (Superdex S200, GE Healthcare) was
equilibrated using lysis buffer and the protein loaded. Protein was concentrated
(Vivaspin 20, 50 kDa MWCO, Sartorius), and flash frozen in liquid nitrogen until further
use.

Crystallization and data collection

Protein crystallization was carried out in an anoxic chamber with less than 5 ppm of O_2.
Solutions used for crystallization were degassed using modified Schlenck techniques
and crystallization was carried out using the sitting-drop vapor diffusion technique in 96
well plates (Anc1A and Anc1B) (Swissci 96-well two-drop plate, Hampton Research) or
24 well plates (Anc2) (24-well crystallization Cryschem M plates, Hampton Research).
Crystals which were used for microseeding were added to 50 μl reservoir solution and a
PTFE seed bead (Hampton Research). The crystals were crushed by 6 times vortexing
for 30 s and then diluted with reservoir solution. 7.5 mM of $Na_2S_2O_4$ were added to the
protein prior to crystallization. 2 μl of 10 mg/ml protein solution (Anc2) were mixed with
the same volume of reservoir solution containing 4.5% (w/v) of polyethylene glycol
2000, 3350, 4000 and polyethylene glycol mono methylether 5000, 0.1 M MES/NaOH at
pH 6.5, 10% (v/v) of ethylene glycol and 0.15 M Mg acetate. 0.2 μl of 1:100 diluted seed
were added.

For Anc1A, 0.3 μl of 9 mg/ml protein solution were mixed with 0.4 μl of reservoir solution
containing 0.2 M $LiSO_4$ x 1 H_2O, 0.1 M bis-Tris/HCl at pH 6.5, 25%(w/v) of polyethylene
glycol 3350. 0.1 μl of 1:400 diluted seed was added. 0.4 μl of 9 mg/ml protein solution
(Anc1B) was mixed with 0.4 μl of reservoir solution containing 0.1 M MES/imidazole at
pH 6.5, 12.5% of (w/v) of polyethylene glycol 1000, 3350 and (v/v) 2-methyl-2,4-
pentandiol each and 0.02 M of 1,6-hexanediol, 1-butanol, 1,4-butanediol, (RS)-1,2-

propanediol, 2-propanol, and 1,3-propanediol, each with addition of 0.1 µl of 1:400

diluted seed. 0.8 µl of 8 mg/ml Anc1B-H was mixed with 0.8 µl of reservoir containing

0.1 M MES/NaOH at pH 6.5, 7.5% of (w/v) polyethylene glycol 6000, 8000, 10000 each,

10%(vol/vol) isopropanol. 0.2 µl of 1:400 diluted seed stock was added.

Crystals were harvested into nylon loops and flash-frozen in liquid nitrogen. Diffraction

data for Anc2 were collected at the Swiss Light Source (Paul Scherrer Institute) on

beamline X06SA using an EIGER 16M X detector at an X-ray wavelength of 1.0000 Å.

Diffraction data for Anc1A and B were collected at the ESRF on beamline ID30B using

an EIGER2 X 9M at an X-ray wavelength of 0.8856 A. Diffraction date for Anc1B-H was

collected at the ESRF on beamline ID23-1 using an EIGER2 16M CdTe detector at an

X-ray wavelength of 0.8856 A.

Structure solution and refinement

The crystallographic phase problem was solved using the WT structure of MoFe or Fe

protein, respectively, from *A. vinelandii* (PDB:3U7Q, 1G1M). A single solution was

obtained for the mutants using MOLREP (Anc1B, Anc1B-NifH) (Vagin and Teplyakov,

2010) or PhaserMR (Anc1A, Anc2) (McCoy et al., 2007). The obtained model was

refined using iterative rounds of BUSTER (Anc1A, Anc2) (Blanc et al., 2004) or

REFMAC5 (all) (Murshudov et al., 2011) and model building in COOT (Emsley et al.,

2010). **Tables S2 to S5** show the statistics of the refinement process.

Ancient and extant protein structure prediction

Structure predictions based on Alphafold-derived methods require two steps: generating

deep sequence alignments and predicting structures based on those alignments

(Jumper et al., 2021; Senior et al., 2020; Tunyasuvunakool et al., 2021). Additionally,

structural templates can be added to the process, but they are not mandatory. The first

step depends on searching large sequence databases. The optimization of this step

requires significant memory resources which are limiting in many cases. Colabfold

provides a public MMSeqs2 server (Steinegger and Söding, 2017) that generates
multiple-sequence alignments tailored to Alphafold requirements to solve this issue
(Mirdita et al., 2022). However, the public nature of this server implies that users can
only perform a limited number of queries per unit of time. The second step consists of
running forward predictions on a deep neural network. This step is usually performed in
high-performance graphical processing units (GPU). To predict nitrogenase structures,
we have considered a *bonafide* protocol and a recycling protocol.

Bonafide protocol – In this protocol, we rely on the Colabfold methodology, which
consists of using MMSeqs2 to generate a deep sequence alignment covering the whole
sequence of the query and running forward predictions with 3 recycles using the
Alphafold neural network. In this step, we avoided using structural templates and
optimizing side chains using the AMBER force field.

Recycling protocol – To predict variant structures with a high level of sequence identity
to another reference sequence, we implemented a different pipeline based on recycling
the MSA generated by MMSeqs2. The main reason for this additional step is to avoid
blocking a public resource with thousands of similar requests. We re-aligned the
corresponding concatenation of sequences to the MSA generated on the structure
prediction of the maximum-likelihood ancestral sequence using MAFFT with its options
"--keepLength" and "--addResidue" (Katoh and Standley, 2013). Previous conversions
from A3M to FASTA format were carried out using HH-suite convert.pl script (Steinegger
et al., 2019). The result of this re-alignment was converted to A3M again, and the files
were modified to include the headers specifying the number of chains for each
sequence and the length of each sequence in the concatenated dataset. The rest of the
protocol proceeded as the *bonafide* protocol, using 3 recycles without templates nor
side-chain optimization.

Protein structure analysis pipeline

The final dataset includes a total of 5378 structures corresponding to extant and

ancestral structures. We utilized semi-automatic methods to analyze the high number of

structures. Our pipeline for the analysis of structures includes the following steps. (1)

Aligning every structure against a template using US-Align (Zhang et al., 2022). For the

component-1 complexes (chains DDKK), Vnf nitrogenase (PDB: 5N6Y) was used as a

template. For the component-2, the NifH dimer was used as a template (PDB: 1G1M),

both templates from A. *vinelandii.* (2) Modifying the name of chains to match the

reference structure, and (3) Extracting the pLDDT, predicted local distance difference

test, which is the standard uncertainty predictor of Alphafold (Jumper et al., 2021) from

the B-factor column of PDB files and trimming regions of the structure with low

prediction confidence.

The similarity of two protein structures is computed using US-Align (Zhang et al., 2022),

which extends the structural alignment capabilities of the TM-align (Zhang and Skolnick,

2005) method to proteins containing multiple chains. US-Align provides two metrics that

provide an assessment of the similarity between proteins: first, the Root Mean Square

Deviation (RMSD), which measures the mean distance in which a set of atoms deviates

from another; and the TM-score, which is a metric bound between 0 (no similarity) and 1

(absolute similarity).

[FORMULA 1: RMSD]

$$RMSD = \sqrt{\frac{1}{N}\sum_{i}^{N}\left(x_i^{ref} - x_i\right)^2}$$

765   Where N is the number of atoms, x_i are the 3D coordinates of the aligned atoms, and

766   x_i^{ref} are the 3D coordinates of the reference atoms. In this work, we compute this

767   function solely based on the coordinates of α-carbons.

768   [FORMULA 2: TM-SCORE]

$$TMScore = \max\left(\frac{1}{L^{ref}}\sum_{i}^{L}\frac{1}{1+\left(\frac{d_i}{d_0(L^{ref})}\right)^2}\right)$$

Where L^{ref} is the length of the reference protein, L is the number of aligned residues, d_i
is the distance between a par of aligned residues, and $d_0(L^{ref})$ is a normalization term to
account for the size of the proteins under alignment. We additionally attempted to
correct possible bias connected to the structural alignment size using the RMSD100
metric (Carugo and Pongor 2001), defined as:
[FORMULA 3: RMSD100]
776

$$RMSD_{100} = RMSD \cdot \frac{1}{1+\ln\sqrt{\frac{L}{100}}}$$

777   Where L is the length of the structural alignment.
778

We have combined different programs' outputs to study the structural features among
the different nitrogenases. The list of studied features and the respective programs are
summarized in **Table S6**. For normal-mode derived properties and geometric properties
(e.g., gyration radius), we used ProDy (Bakan et al. 2011). For affinity predictions, we
used PRODIGY (Vangone and Bonvin 2015). For surface solvent-accessible areas, we
used FreeSASA (Mitternacht 2016). Finally, for residue-interaction network properties,
we used RING (Clementel et al. 2022).
786

787   **Nitrogenase structure repository**
788

789   We implemented a lightweight server using Streamlit framework in Python providing
790   access and the ability to download to all reconstructed structures using taxon IDs, gene
791   names, or species names available at https://nsdb.bact.wisc.edu.
792   **Visualization**
793

To visualize quantitative data, we employed the Seaborn Python library. To visualize
protein structures, we used ChimeraX-v1.5 (Goddard et al., 2018).

DATA AVAILABILITY

All codes are available at Github: http://github.com/kacarlab/nif-structures

All crystal structures will be available at the Protein Database upon publication: Anc1A
DDKK: 9HB9, Anc 1B HH: 9HAZ, Anc1B DDKK: 9HBN and Anc2 DDKK: 9HBC.

All structures are available at the nitrogenase structure repository at the UW-Madison
Department of Bacteriology hosted https://nsdb.bact.wisc.edu and at Zenodo
10.5281/zenodo.13351063.

AUTHOR CONTRIBUTIONS
Conceptualization and study design: BCZ and BK
Data collection: BCZ, FD, KA, AKG
Writing (first draft): BCZ and BK
Writing (reviewing/editing): All authors

ACKNOWLEDGEMENTS

B.C.Z acknowledges the Margarita Salas Postdoctoral Fellowship, founded by the
Unión Europea - Next Generation EU (B.C.Z.; UP2021-035). This work was supported
by the National Aeronautics and Space Administration (NASA) Interdisciplinary
Consortium for Astrobiology Research: Metal Utilization and Selection Across Eons,
MUSE [*80NSSC17K0296*] with additional support from the Human Frontier Science
Program (HFSP) [*RGY0072/2021*] (KA), W. M. Keck Foundation, the Hypothesis Fund
and the NASA Exobiology Program [*NNH23ZDA001N*]. O.E. acknowledges support
from the European Research Council (Horizon Europe, AdG 101141673). We thank the
Center for High Throughput Computing (CHTC), Dr. Derek Harris for assistance in
nitrogenase protein purification, Dr. Janet Newlands for the IT support, the Department

REFERENCES

Anbar AD, Knoll AH. 2002. Proterozoic Ocean Chemistry and Evolution: A Bioinorganic Bridge? *Science* **297**:1137–1142. doi:10.1126/science.1069651

Bakan A, Meireles LM, Bahar I. 2011. ProDy: Protein Dynamics Inferred from Theory and Experiments. *Bioinformatics* **27**:1575–1577. doi:10.1093/bioinformatics/btr168

Barney BM, Yurth MG, Santos PCD, Dean DR, Seefeldt LC. 2009. A substrate channel in the nitrogenase MoFe protein. *JBIC J Biol Inorg Chem* **14**:1015–1022. doi:10.1007/s00775-009-0544-2

Bergman B, Sandh G, Lin S, Larsson J, Carpenter EJ. 2013. Trichodesmium – a widespread marine cyanobacterium with unusual nitrogen fixation properties. *FEMS Microbiol Rev* **37**:286–302. doi:10.1111/j.1574-6976.2012.00352.x

Blanc E, Roversi P, Vonrhein C, Flensburg C, Lea SM, Bricogne G. 2004. Refinement of severely incomplete structures with maximum likelihood in BUSTER–TNT. *Acta Crystallogr Sect D: Biol Crystallogr* **60**:2210–2221. doi:10.1107/s0907444904016427

Boyd ES, Costas AMG, Hamilton TL, Mus F, Peters JW. 2015. Evolution of Molybdenum Nitrogenase during the Transition from Anaerobic to Aerobic Metabolism. *J Bacteriol* **197**:1690–1699. doi:10.1128/jb.02611-14

Brunet TDP, Doolittle WF. 2018. The generality of Constructive Neutral Evolution. *Biol Philos* **33**:2. doi:10.1007/s10539-018-9614-6

Burén S, Jiménez-Vicente E, Echavarri-Erasun C, Rubio LM. 2020. Biosynthesis of Nitrogenase Cofactors. *Chem Rev* **120**:4921–4968. doi:10.1021/acs.chemrev.9b00489

Campo JSM del, Rigsbee J, Batista MB, Mus F, Rubio LM, Einsle O, Peters JW, Dixon R, Dean DR, Santos PCD. 2022. Overview of physiological, biochemical, and regulatory aspects of nitrogen fixation in Azotobacter vinelandii. *Crit Rev Biochem Mol Biol* **57**:492–538. doi:10.1080/10409238.2023.2181309

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**:1972–1973. doi:10.1093/bioinformatics/btp348

Carugo O, Pongor S. 2001. A normalized root-mean-spuare distance for comparing protein three-dimensional structures. *Protein Sci* **10**:1470–1473. doi:10.1110/ps.690101

Clementel D, Conte AD, Monzon AM, Camagni GF, Minervini G, Piovesan D, Tosatto SCE. 2022. RING 3.0: fast generation of probabilistic residue interaction networks from structural ensembles. *Nucleic Acids Res* **50**:W651–W656. doi:10.1093/nar/gkac365

Cuevas-Zuviría B, Garcia AK, Rivier AJ, Rucker HR, Carruthers BM, Kaçar B. 2024.
Emergence of an orphan nitrogenase protein following atmospheric oxygenation. *Mol*
Biol Evol msae067. doi:10.1093/molbev/msae067

Dixon R, Kahn D. 2004. Genetic regulation of biological nitrogen fixation. *Nat Rev*
Microbiol **2**:621–631. doi:10.1038/nrmicro954

Einsle O, Rees DC. 2020. Structural Enzymology of Nitrogenase Enzymes. *Chem Rev*
120:4969–5004. doi:10.1021/acs.chemrev.0c00067

Emsley P, Lohkamp B, Scott WG, Cowtan K. 2010. Features and development of Coot.
Acta Crystallogr Sect D: Biol Crystallogr **66**:486–501. doi:10.1107/s0907444910007493

Falkowski PG. 1997. Evolution of the nitrogen cycle and its influence on the biological
sequestration of CO2 in the ocean. *Nature* **387**:272–275. doi:10.1038/387272a0

Garcia AK, Harris DF, Rivier AJ, Carruthers BM, Pinochet-Barros A, Seefeldt LC, Kaçar
B. 2023. Nitrogenase resurrection and the evolution of a singular enzymatic
mechanism. *eLife* **12**:e85003. doi:10.7554/elife.85003

Garcia AK, Kaçar B. 2019. How to resurrect ancestral proteins as proxies for ancient
biogeochemistry. *Free Radic Biol Med* **140**:260–269.
doi:10.1016/j.freeradbiomed.2019.03.033

Garcia AK, Kolaczkowski B, Kaçar B. 2022. Reconstruction of Nitrogenase
Predecessors Suggests Origin from Maturase-Like Proteins. *Genome Biol Evol*
14:evac031. doi:10.1093/gbe/evac031

Garcia AK, McShea H, Kolaczkowski B, Kaçar B. 2020. Reconstructing the evolutionary
history of nitrogenases: Evidence for ancestral molybdenum-cofactor utilization.
Geobiology **18**:394–411. doi:10.1111/gbi.12381

Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, Ferrin TE.
2018. UCSF ChimeraX: Meeting modern challenges in visualization and analysis.
Protein Sci **27**:14–25. doi:10.1002/pro.3235

Harris DF, Rucker HR, Garcia AK, Yang Z-Y, Chang SD, Feinsilber H, Kaçar B, Seefeldt
LC. 2024. Ancient nitrogenases are ATP dependent. *mBio* **15**:e01271-24.
doi:10.1128/mbio.01271-24

Johnson AC, Ostrander CM, Romaniello SJ, Reinhard CT, Greaney AT, Lyons TW,
Anbar AD. 2021. Reconciling evidence of oxidative weathering and atmospheric anoxia
on Archean Earth. *Sci Adv* **7**:eabj0108. doi:10.1126/sciadv.abj0108

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool
K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie

A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D,
Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver
D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate
protein structure prediction with AlphaFold. *Nature* **596**:583–589. doi:10.1038/s41586-
021-03819-2

Kaçar B. 2024. Reconstructing Early Microbial Life. *Annu Rev Microbiol*.
doi:10.1146/annurev-micro-041522-103400

Kalyaanamoorthy S, Minh BQ, Wong TKF, Haeseler A von, Jermiin LS. 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods*
14:587–589. doi:10.1038/nmeth.4285

Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Mol Biol Evol* **30**:772–780.
doi:10.1093/molbev/mst010

Knoll, A.H. (2003). Life on a Young Planet: The First Three Billion Years of Evolution on
Earth. Princeton University Press.

Lyons TW, Reinhard CT, Planavsky NJ. 2014. The rise of oxygen in Earth's early ocean
and atmosphere. *Nature* **506**:307–315. doi:10.1038/nature13068

Lyons TW, Tino CJ, Fournier GP, Anderson RE, Leavitt WD, Konhauser KO, Stüeken
EE. 2024. Co-evolution of early Earth environments and microbial life. *Nat Rev*
Microbiol **22**:572–586. doi:10.1038/s41579-024-01044-y

Mariani V, Biasini M, Barbato A, Schwede T. 2013. lDDT: a local superposition-free
score for comparing protein structures and models using distance difference tests.
Bioinformatics **29**:2722–2728. doi:10.1093/bioinformatics/btt473

McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ. 2007.
Phaser crystallographic software. *J Appl Crystallogr* **40**:658–674.
doi:10.1107/s0021889807021206

Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. 2022.
ColabFold: making protein folding accessible to all. *Nat Methods* **19**:679–682.
doi:10.1038/s41592-022-01488-1

Mitternacht S. 2016. FreeSASA: An open source C library for solvent accessible surface
area calculations. *F1000Research* **5**:189. doi:10.12688/f1000research.7931.1

Moore EK, Jelen BI, Giovannelli D, Raanan H, Falkowski PG. 2017. Metal availability
and the expanding network of microbial metabolisms in the Archaean eon. *Nat Geosci*
10:629–636. doi:10.1038/ngeo3006

Morrison CN, Hoy JA, Zhang L, Einsle O, Rees DC. 2015. Substrate Pathways in the
Nitrogenase MoFe Protein by Experimental Identification of Small Molecule Binding
Sites. *Biochemistry* **54**:2052–2060. doi:10.1021/bi501313k

Mrnjavac N, Esposti MD, Mizrahi I, Martin WF, Allen JF. 2024. Three enzymes governed
the rise of O2 on Earth. *Biochim Biophys Acta (BBA) - Bioenerg* **1865**:149495.
doi:10.1016/j.bbabio.2024.149495

Murshudov GN, Skubák P, Lebedev AA, Pannu NS, Steiner RA, Nicholls RA, Winn MD,
Long F, Vagin AA. 2011. REFMAC5 for the refinement of macromolecular crystal
structures. *Acta Crystallogr Sect D* **67**:355–367. doi:10.1107/s0907444911001314

Mus F, Colman DR, Peters JW, Boyd ES. 2019. Geobiological feedbacks, oxygen, and
the evolution of nitrogenase. *Free Radic Biol Med* **140**:250–259.
doi:10.1016/j.freeradbiomed.2019.01.050

Navarro-González R, McKay CP, Mvondo DN. 2001. A possible nitrogen crisis for
Archaean life due to reduced nitrogen fixation by lightning. *Nature* **412**:61–64.
doi:10.1038/35083537

Nguyen L-T, Schmidt HA, Haeseler A von, Minh BQ. 2015. IQ-TREE: A Fast and
Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol*
Evol **32**:268–274. doi:10.1093/molbev/msu300

Parsons C, Stüeken EE, Rosen CJ, Mateos K, Anderson RE. 2021. Radiation of
nitrogen-metabolizing enzymes across the tree of life tracks environmental transitions in
Earth history. *Geobiology* **19**:18–34. doi:10.1111/gbi.12419

Peters JW, Stowell MHB, Soltis SM, Finnegan MG, Johnson MK, Rees DC. 1997.
Redox-Dependent Structural Changes in the Nitrogenase P-Cluster † , ‡. *Biochemistry*
36:1181–1187. doi:10.1021/bi9626665

Raymond J, Siefert JL, Staples CR, Blankenship RE. 2004. The Natural History of
Nitrogen Fixation. *Mol Biol Evol* **21**:541–554. doi:10.1093/molbev/msh047

Robson RL, Postgate JR. 1980. Oxygen and Hydrogen in Biological Nitrogen Fixation.
Annu Rev Microbiol **34**:183–207. doi:10.1146/annurev.mi.34.100180.001151

Rucker HR, Kaçar B. 2024. Enigmatic evolution of microbial nitrogen fixation: insights
from Earth's past. *Trends Microbiol* **32**:554–564. doi:10.1016/j.tim.2023.03.011

Rutledge HL, Cook BD, Nguyen HPM, HerzikJr. MA, Tezcan FA. 2022. Structures of the
nitrogenase complex prepared under catalytic turnover conditions. *Science* **377**:865–
869. doi:10.1126/science.abq7641

Sánchez-Baracaldo P, Ridgwell A, Raven JA. 2014. A Neoproterozoic Transition in the Marine Nitrogen Cycle. *Curr Biol* **24**:652–657. doi:10.1016/j.cub.2014.01.041

Santos PCD. 2018. Metalloproteins, Methods and Protocols. *Methods Mol Biol* **1876**:91–109. doi:10.1007/978-1-4939-8864-8_6

Schlesier J, Rohde M, Gerhardt S, Einsle O. 2016. A Conformational Switch Triggers Nitrogenase Protection from Oxygen Damage by Shethna Protein II (FeSII). *J Am Chem Soc* **138**:239–247. doi:10.1021/jacs.5b10341

Schmid B, Einsle O, Chiu H-J, Willing A, Yoshida M, Howard JB, Rees DC. 2002. Biochemical and Structural Characterization of the Cross-Linked Complex of Nitrogenase: Comparison to the ADP-AlF4 --Stabilized Structure † , ‡. *Biochemistry* **41**:15557–15565. doi:10.1021/bi026642b

Senior AW, Evans R, Jumper J, Kirkpatrick J, Sifre L, Green T, Qin C, Žídek A, Nelson AWR, Bridgland A, Penedones H, Petersen S, Simonyan K, Crossan S, Kohli P, Jones DT, Silver D, Kavukcuoglu K, Hassabis D. 2020. Improved protein structure prediction using potentials from deep learning. *Nature* **577**:706–710. doi:10.1038/s41586-019-1923-7

Sippel D, Einsle O. 2017. The structure of vanadium nitrogenase reveals an unusual bridging ligand. *Nat Chem Biol* **13**:956–960. doi:10.1038/nchembio.2428

Sobol, M.S., Ané, C., McMahon, K.D. and Kaçar, B., 2025. Ecological constraints and evolutionary trade-offs shape nitrogen fixation across habitats. bioRxiv, pp.2025-05.

Spatzal T, Aksoyoglu M, Zhang L, Andrade SLA, Schleicher E, Weber S, Rees DC, Einsle O. 2011. Evidence for Interstitial Carbon in Nitrogenase FeMo Cofactor. *Science* **334**:940–940. doi:10.1126/science.1214025

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**:1312–1313. doi:10.1093/bioinformatics/btu033

Steinegger M, Meier M, Mirdita M, Vöhringer H, Haunsberger SJ, Söding J. 2019. HH-suite3 for fast remote homology detection and deep protein annotation. *BMC Bioinform* **20**:473. doi:10.1186/s12859-019-3019-7

Steinegger M, Söding J. 2017. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nat Biotechnol* **35**:1026–1028. doi:10.1038/nbt.3988

Stüeken EE, Buick R, Guy BM, Koehler MC. 2015. Isotopic evidence for biological nitrogen fixation by molybdenum-nitrogenase from 3.2 Gyr. *Nature* **520**:666–669. doi:10.1038/nature14180

Takimoto R, Tatemichi Y, Aoki W, Kosaka Y, Minakuchi H, Ueda M, Kuroda K. 2022. A
critical role of an oxygen-responsive gene for aerobic nitrogenase activity in Azotobacter
vinelandii and its application to Escherichia coli. *Sci Rep-uk* **12**:4182.
doi:10.1038/s41598-022-08007-4

Tezcan FA, Kaiser JT, Mustafi D, Walton MY, Howard JB, Rees DC. 2005. Nitrogenase
Complexes: Multiple Docking Sites for a Nucleotide Switch Protein. *Science* **309**:1377–
1380. doi:10.1126/science.1115653

Trncik C, Detemple F, Einsle O. 2023. Iron-only Fe-nitrogenase underscores common
catalytic principles in biological nitrogen fixation. *Nat Catal* 1–10. doi:10.1038/s41929-
023-00952-1

Tunyasuvunakool K, Adler J, Wu Z, Green T, Zielinski M, Žídek A, Bridgland A, Cowie A,
Meyer C, Laydon A, Velankar S, Kleywegt GJ, Bateman A, Evans R, Pritzel A, Figurnov
M, Ronneberger O, Bates R, Kohl SAA, Potapenko A, Ballard AJ, Romera-Paredes B,
Nikolov S, Jain R, Clancy E, Reiman D, Petersen S, Senior AW, Kavukcuoglu K, Birney
E, Kohli P, Jumper J, Hassabis D. 2021. Highly accurate protein structure prediction for
the human proteome. *Nature* **596**:590–596. doi:10.1038/s41586-021-03828-1

Vagin A, Teplyakov A. 2010. Molecular replacement with MOLREP. *Acta Crystallogr
Sect D: Biol Crystallogr* **66**:22–25. doi:10.1107/s0907444909042589

Vangone A, Bonvin AM. 2015. Contacts-based prediction of binding affinity in protein–
protein complexes. *Elife* **4**:e07454. doi:10.7554/elife.07454

Vicente EJ, Dean DR. 2017. Keeping the nitrogen-fixation dream alive. *Proc Natl Acad
Sci* **114**:3009–3011. doi:10.1073/pnas.1701560114

Xu J, Zhang Y. 2010. How significant is a protein structure similarity with TM-score =
0.5? *Bioinformatics* **26**:889–895. doi:10.1093/bioinformatics/btq066

Yang Z. 2007. PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Mol Biol Evol*
24:1586–1591. doi:10.1093/molbev/msm088

Yang Z, Kumar S, Nei M. 1995. A new method of inference of ancestral nucleotide and
amino acid sequences. *Genetics* **141**:1641–1650. doi:10.1093/genetics/141.4.1641

Zhang C, Shine M, Pyle AM, Zhang Y. 2022. US-align: universal structure alignments of
proteins, nucleic acids, and macromolecular complexes. *Nat Methods* **19**:1109–1115.
doi:10.1038/s41592-022-01585-1

Zhang S, Krieger JM, Zhang Y, Kaya C, Kaynak B, Mikulska-Ruminska K, Doruker P, Li
H, Bahar I. 2021. ProDy 2.0: increased scale and scope after 10 years of protein
dynamics modelling with Python. *Bioinformatics* **37**:3657–3659.
doi:10.1093/bioinformatics/btab187

Zhang Y, Skolnick J. 2005. TM-align: a protein structure alignment algorithm based on
the TM-score. *Nucleic Acids Res* **33**:2302–2309. doi:10.1093/nar/gki524

Zhang Y, Skolnick J. 2004. Scoring function for automated assessment of protein
structure template quality. *Proteins: Struct, Funct, Bioinform* **57**:702–710.
doi:10.1002/prot.20264

1033

1034