Title

Insulin/insulin-like growth factor signaling pathway promotes higher fat storage in

Drosophila females

Authors

Puja Biswas[1], Jennifer A. Bako[1§], J. Beatrice Liston[1§], Huaxu Yu[2,3], Lianna W. Wat[1,4],

Colin J. Miller[1], Michael D. Gordon[5], Tao Huan[2], Molly Stanley[6], Elizabeth J. Rideout[1,7, *]

Affiliations

[1] Department of Cellular and Physiological Sciences, Life Sciences Institute, The

University of British Columbia, Vancouver, Canada.

[2] Department of Chemistry, Faculty of Science, The University of British Columbia,

Vancouver, Canada.

[3] Present address: West Coast Metabolomics Center, University of California, Davis,

California 95616, USA.

[4] Present address: Department of Pathology & Department of Neurobiology, Stanford

Diabetes Research Center, Stanford Cardiovascular Institute, Stanford University School

of Medicine, Stanford, California, USA.

[5] Department of Zoology, Life Sciences Institute, and Djavad Mowafaghian Centre for

Brain Health, The University of British Columbia, Vancouver, Canada.

[6] Department of Biology, The University of Vermont, Burlington, Vermont, USA.

[7] Lead contact

[§] Equal contribution

* Correspondence: elizabeth.rideout@ubc.ca

Summary

In *Drosophila*, adult females store more fat than males. While the mechanisms that restrict

body fat in males are becoming clearer, less is known about how females achieve higher

fat storage. Here, we perform a detailed investigation of the mechanisms that promote

higher fat storage in females. We show greater intake of dietary sugar supports higher fat

storage due to female-biased remodeling of the fat body lipidome. Dietary sugar stimulates

a female-specific increase in *Drosophila* insulin-like peptide 3 (Dilp3), which acts together

with greater peripheral insulin sensitivity to augment insulin/insulin-like growth factor

signaling pathway (IIS) activity in adult females. Indeed, Dilp3 overexpression prevented

the female-biased decrease in body fat after removal of dietary sugar. Given that adult-

specific IIS inhibition caused a female-biased decrease in body fat, our data reveal IIS as a

key determinant of female fat storage.

Keywords

Sex difference, body fat, triglyceride, insulin/insulin-like growth factor signaling pathway,

sugar, insulin sensitivity

Introduction

In *Drosophila*, as in other animals, triglyceride is the main stored form of fat[1]. The primary depot for stored fat in insects is called the fat body[1]; however, other cell types also synthesize and store triglyceride (*e.g.*, gut, oenocytes, neurons, glia, muscle, testis, ovary)[2-9]. While many factors influence the total amount of stored fat, one important determinant of body fat in flies is biological sex[5,9,10]. For example, *Drosophila* virgin females store more fat than males[5], a difference that is amplified post-mating[9,11]. Virgin females also break down fat more slowly than males after nutrient withdrawal[5]. These sex differences in body fat regulation are physiologically significant: females use their greater fat stores to support the energetic demands of reproduction[9,12] and to survive long bouts of nutrient deprivation[5], whereas males need to restrict excess triglyceride accumulation to maintain fertility[8].

Over the past two decades, studies in *Drosophila* have played a key role in identifying many genes and pathways that contribute to whole-body fat storage. For example, loss-of-function experiments with genes predicted to promote triglyceride synthesis lead to reduced fat storage in both individual cells and in the whole body[12-16]. In contrast, genetically inhibiting genes that facilitate triglyceride breakdown significantly augment fat storage at the cellular and organismal levels[17-19]. Gain- and loss-of-function experiments with genes predicted to regulate lipid droplet dynamics, a specialized organelle dedicated to triglyceride storage, similarly influence cellular and organismal fat stores[16,19-24]. Beyond these enzymes directly involved in triglyceride metabolism and lipid droplet biology, large-scale screening efforts have revealed many additional genes that influence *Drosophila* body fat[4,18,25-39].

In addition to the action of individual genes, a large body of literature has identified hormones and peptides that influence whole-body fat storage[9,40-57]. Often, these hormones modulate whole-body fat storage via effects on cells that produce adipokinetic hormone

(Akh)[43-46,58] and *Drosophila* insulin-like peptides (Dilps)[30,43,45,59-62], where Akh and most

Dilps have opposing effects on whole-body fat storage[63]. For example, when nutrients are

plentiful, Dilps 2, 3, and 5 are released from the insulin-producing cells (IPC) in the

Drosophila brain into the hemolymph[43,59]. Binding of the Dilps to the insulin receptor (InR)

on the fat body stimulates the activity of the insulin/insulin-like growth factor signaling

pathway (IIS) to promote fat storage[53]. When nutrients are scarce, Akh is released into the

hemolymph from the Akh-producing cells (APC) in the *corpora cardiaca*[46,64-67]. Akh binds

to the Akh receptor located in the fat body where it promotes fat breakdown[55-57]. Despite

this advanced knowledge of factors that regulate fat storage, most studies used single- or

mixed-sex animal groups. As a result, less is known about the mechanisms underlying sex

differences in fat storage.

Recent studies have begun to close this knowledge gap. Both large-scale RNAseq

data and targeted gene expression studies reveal profound sex differences in expression

of genes that regulate cellular and whole-body fat storage[5,68,69]. For example, male flies

have higher mRNA levels of the *Drosophila* homolog of adipose triglyceride lipase called

brummer (*bmm*), an enzyme that limits triglyceride accumulation across most cell types[17].

This male-biased *bmm* expression is physiologically significant, as whole-body loss of

bmm abolishes the sex difference in fat storage via a male-biased increase in body fat[5]. A

similar male bias has also been reported in the production and secretion of Akh[10], which

prevents fat accumulation to restrict body fat in males[10]. A male bias in two central lipolytic

pathways therefore plays a key role in establishing the sex difference in body fat. Yet, the

mechanisms that allow increased fat storage in females are less clear, as females are

largely unaffected by loss of *bmm* and Akh[5,10].

In mated females, higher levels of steroid hormone ecdysone enhance body fat by

stimulating food intake[9]. Mated females also show higher mRNA levels of *dilp3*[70], where

dilp3 has been implicated in promoting fat storage[43,62]. Whether these mechanisms also

influence the sex difference in fat storage remains unclear. We therefore performed a

detailed analysis of adult virgin males and females to explore how females achieve a

higher level of fat storage. We found that virgin female flies normally consume more food

than males, and show that female fat storage is sensitive to changes in food quantity. In

particular, consumption of dietary sugar and not simply a greater number of calories

explained this higher fat storage in females. Unbiased lipidomic and metabolomic analysis

of abdominal carcasses from adult flies revealed dietary sugar caused a profoundly

female-biased metabolic remodeling of energy-storing cells including the fat body. Given

that ecdysone did not promote increased body fat in virgin females, we characterized sex

differences in IIS, the other key pathway that promotes fat storage. We identified

significant female biases in Dilp3 levels and fat body insulin sensitivity, leading to higher

IIS activity in the female fat body. These sex biases were significant: Dilp3 overexpression

in females bypassed the reduction in body fat caused by a low-sugar diet, and ablating the

IPC in adults caused a female-specific reduction in body fat. Together, these data identify

IIS as a key pathway that promotes higher fat storage in adult females.

Results

Dietary sugar promotes increased fat storage in adult females

To determine whether unmated females have higher food intake than males we used the

capillary feeder (CAFE) assay[71] to assess food consumption in 5-day-old *Drosophila* virgin

males and females. Five-day-old adult *Canton-S* (*CS*) females consumed significantly

more food per body weight over 24 hr than genotype-matched males (Figure 1A), a

phenotype we reproduced in 5-day-old *w*^{1118} adults (Figure S1A). Considering changes in

food intake affect body fat[26,47,72], this suggests higher food consumption may explain this

increased fat storage in females. To monitor body fat, we used a coupled colorimetric

assay to measure whole-body triglyceride in male and female flies[4,35,73], where triglyceride

content was normalized to mass to account for sex differences in body size as described

previously[5,10]. All non-normalized triglyceride data are available in Supplemental Table S1.

Importantly, coupled colorimetric assays have been shown to accurately quantify stored

triglyceride in *Drosophila*[74]. Reducing nutrient content in the diet significantly reduced body

fat in *CS* females (Figure 1B). Because body fat in *CS* males was unaffected by an

identical food dilution (Figure 1B), the sex difference in body fat was reduced (Figure 1B').

We next tested the contribution of specific diet components to higher fat storage in

females. Given that dietary sugar affects body fat in a mixed-sex population of

Drosophila[75-79], we transferred newly-eclosed *CS* and w^{1118} flies to a diet lacking added

sugar (0S) for 5 days[80]. In both w^{1118} and *CS* strains, adult females transferred to the 0S

diet had significantly less body fat compared with genotype-matched flies kept on a widely-

used lab diet with sugar (1S; Figure 1C and S1B)[81]. Because the magnitude of the

reduction in body fat was smaller in w^{1118} males (Figure 1C; sex:genotype interaction

p<0.0001), with the same trend in *CS* males (Figure S1B), the sex difference in fat storage

was reduced (Figure 1C').

The female-biased reduction in fat storage on 0S cannot be attributed to altered

food intake on the low-sugar diet, as food consumption was not different between flies kept

on the 0S diet compared with flies maintained on the 1S diet (Figure S1C). A decrease in

calories similarly cannot explain the effect of reduced sugar on body fat, as female flies

transferred to a reduced-calorie diet with sugar still had higher body fat than females kept

on the 0S diet (Figure 1D and 1D'). Because adding the quantity of either sucrose (20.5

g/L) or D-glucose (70.9 g/L) that we usually put in 1S food to the 0S diet was not sufficient

to restore fat storage to the same level as females maintained on the 1S diet (Figure 1E),

our data suggests it is the total amount of sugar rather than a specific sugar in the diet that

promotes increased fat storage in female flies. Indeed, providing females maintained on

1S with additional sugar was sufficient to further increase body fat (Figure 1F). Because

the magnitude of these sugar-dependent changes in body fat (-1% or +1% body fat) are

similar to the change in body fat we observe in females reared on a high-fat diet (+1%)[5],

where a high-fat diet is associated with a number of adverse physiological effects[82-84] and

changes in fertility and lifespan[85,86], the change in body fat we observed on the 0S diet is

physiologically relevant. Finally, supporting a model in which the nutritive value of sugar

rather than sweet taste promotes increased female fat storage, body fat was not higher in

either male or female flies kept on the 0S diet containing a non-nutritive but sweet-tasting

sugar[87] (Figure S1D). Thus, our data suggest that higher food intake, specifically

increased consumption of dietary sugar, promotes higher fat storage in females.

Female-biased effect of dietary sugar on metabolism

To assess how sex and dietary sugar affect the lipidome and metabolome of fat-storing

organs in *Drosophila*, we performed untargeted mass spectrometry (MS)-based lipidomic

profiling on abdominal carcasses isolated from adult male and female flies transferred to

either a 1S or 0S diet as adults. We detected 464 lipid species in total (Supplemental

Table S1). We first compared lipid abundance between males and females. We identified

sex differences in the abundance of 163 lipids in flies maintained on the 1S diet (p<0.01):

117 lipid species were significantly higher in females and 46 lipid species were significantly

higher in males (Figure 2A). When we examined the sex bias of lipids within each lipid

subclass, we found that triglyceride was the subclass with the most species showing a

female bias. Indeed, 48/57 (86%) of sex-biased triglyceride species showed a female bias

in abundance (Figure 2B). This validates data from our coupled colorimetric assays,

strongly supporting increased stored fat in females.

When we examined specific fatty acids within triglyceride species that show a sex-

bias, we found that triglycerides with a female bias contained a greater number of

monounsaturated fatty acids such as myristoleic acid (C14:1), palmitoleic acid (C16:1),

and oleic acid (C18:1) (Figure S2A). Triglycerides with a female bias also contained a

greater number of odd-chain fatty acids (Figure S2A) and C28:2 (7/7). This female bias in

odd-chain fatty acids aligns with published data showing odd-chain fatty acids are derived

from the gut microbiome, where the microbe load is higher in adult female intestines than

in males[88]. A female bias in C28:2 also aligns with the fact that it is a fatty acid precursor

used for synthesis of female-specific pheromones. Triglycerides with a male bias

contained a greater number of C24:1 (4/4) and myristic acid (C14:0) (Figure S2A). Beyond

triglyceride, we also noted a significant female bias in lipid species within subclasses such

as acyl-carnitine (acyl-Car) (4/5), ceramide (Cer) (4/7), diacylglycerol (DG) (6/9),

phosphatidylcholine (PC) (16/25), ether-linked phosphatidylcholine (PC-O) (4/6),

phosphatidylethanolamine (PE) (17/20), phosphatidylglycerol (PG) (3/3),

phosphatidylinositol (PI) (3/3), sterol-ester (SE) (4/4) and sterol (ST) (4/4) (Figure 2B).

Lipid species with a significant male bias included lipids within subclasses such as

lysophosphatidylcholine (LPC) (2/3) and ether-linked triacylglycerol (TG-O) (12/13) (Figure

2B). As with lipids, there was a significant female bias in many metabolites ($p<0.05$)

(Figure 2C), including asparagine, glutamine, glutamic acid, glucose-6-phosphate, ADP,

propionylcarnitine, pantothenic acid, which together suggest a female bias in glycolysis

and the tricarboxylic acid (TCA) cycle. Supporting this, females showed a trend toward

higher levels of citrate, a TCA cycle metabolite, than age- and genotype-matched males

($p=0.06$; Figure S2B). Thus, our data suggest that biological sex is a key regulator of the

lipidome and metabolome.

To determine how dietary sugar affects the lipidome and metabolome in each sex,

we compared these parameters in abdominal carcasses of flies transferred to the 1S or 0S

diets as adults. In females, we observed significant differences in levels of 218 lipids

between flies transferred to 1S compared with flies kept on 0S ($p<0.01$). Specifically,

dietary sugar was associated with an increased abundance of 154 lipids and reduced

abundance of 64 lipids (Figure 2D). Triglyceride showed the largest sugar-dependent

change in abundance, where females maintained on 0S had significantly lower triglyceride

abundance than females kept on 1S (61/81) (Figure S2C), consistent with our triglyceride

assay data. When we examined specific fatty acids within triglyceride species affected by

dietary sugar, we found that triglycerides with a bias toward females maintained on 0S had

fewer C16:1 and C18:1 fatty acids (Figure S2D). Triglycerides with a bias toward females

maintained on 0S similarly had fewer C28:2 fatty acids (Figure S2D). Broader effects of

reduced dietary sugar were also observed across many lipid subclasses in females: Acyl-

Car (3/4), Cer (5/9), DG (11/17), PC (29/33), PC-O (4/7), PE (25/27), PG (5/5), SE (1/1),

ST (1/1), Hex-Cer (1/1), PE-O (1/1) (Figure S2C). Lipids that were significantly higher in

abundance in females maintained on the 0S diet included NAE (1/1), PI (4/6), PS (1/1) and

TG-O (17/21) (Figure S2C). Because body size is fixed before the adult stage of

development[89], and the females are only fed 1S or 0S after eclosion, these diet-induced

alterations in lipid abundance cannot be explained by a sugar-dependent change in body

size. In males, only 27 lipid species were differentially regulated by dietary sugar: 17 lipids

were higher in abundance in flies maintained on a 1S diet whereas 10 lipids were lower in

abundance (Figure 2E). Lipids that were lower in abundance in males on the 0S diet

included PC (7/7), PC-O (2/3), triglyceride (4/6), Cer (1/1), Hex-Cer (1/1) (Figure S2E).

This female bias in sugar-induced changes in the fat body was reproduced in our

metabolomic data (Supplemental Table S1, Figure 2F): 38 metabolites showed a

difference in abundance in females maintained on the 0S diet whereas only 13 metabolites

were altered in males kept on 0S ($p<0.05$ threshold for both sexes) (Figure 2G).

Metabolites that were altered in females on the 0S diet indicate reduced glycolysis and

TCA cycle (asparagine, glutamine, glutamic acid, glucose-6-phosphate, ADP,

nicotinamide, propionylcarnitine, pantothenic acid) and lower oxidative stress (oxidized

glutathione, tans-4-hydroxy-L-proline, methioninesulfoxide, L-5-oxoproline (also known as

pyroglutamic acid), kynurenine) (Figure 2F). In males, metabolites that were altered

suggest reduced glycolysis on the 0S diet (nicotinamide, NAD, AMP) (Figure 2G). Using

an independent assay to monitor sugar-dependent changes in TCA cycle activity, we

found that both sexes had reduced citrate levels when maintained on 0S (Figure S2F).

While metabolic flux analysis will be an important future direction to extend knowledge of

sex differences in the metabolic fate of dietary sugar, our data suggest sugar caused a

strongly female-biased remodeling of the lipidome and metabolome in abdominal

carcasses. While the reason for this profound female bias in changes to the lipidome and

metabolome in response to dietary sugar is unclear, males showed higher levels of sugar

excretion on the 1S diet (Figure 2H).

Dietary sugar promotes sex differences in IIS regulation

Considering dietary sugar stimulates the production and secretion of Dilps from the IPCs in

the brain[90-93], we tested if Dilp production, Dilp secretion, and IIS activity differed between

males and females maintained on 1S. We first measured mRNA levels of IPC-derived

dilp2, *dilp3* and *dilp5* in adult heads using quantitative real-time polymerase chain reaction

(qPCR). Female heads had higher mRNA levels of *dilp3* than male heads (Figure 3A); no

sex differences in mRNA levels of *dilp2* and *dilp5* were found (Figure 3B and 3C). To test

whether dietary sugar caused the sex difference in *dilp3* mRNA levels, we compared *dilp3*

mRNA levels in males and females transferred to either 1S or 0S diets as adults. Female

flies transferred to 0S had significantly lower *dilp3* mRNA levels than females kept on 1S

(Figure 3D). Because there was no sugar-dependent change in male *dilp3* levels, the sex

difference in *dilp3* levels was eliminated (Figure 3D; sex:diet interaction p=0.0128),

showing dietary sugar plays a role in the female bias in *dilp3* mRNA levels. We next asked

whether this bias in *dilp3* mRNA levels extended to Dilp3 protein using a published anti-

Dilp3 antibody[94]. We observed a female bias in Dilp3 levels in the IPC of 5-day-old adults

in both the 1S and 0S diets (Figure S3A-S3E), confirming Dilp3 levels are higher in

females than in males.

Given the sex bias in *dilp3* mRNA and protein, we next monitored Dilp secretion

from the IPCs. We first expressed a calcium-responsive chimeric transcription factor LexA-

VP16-NFAT (*UAS-LexA-VP16-NFAT* [called *UAS-CaLexA*]) in the IPC using *dilp2-GAL4*

(full genotype *dilp2-GAL4>UAS-LexA-VP16-NFAT*). Because the chimeric transcription

factor drives expression of GFP in response to changes in cellular calcium levels, this

method provides an indirect readout of neuronal activity and Dilp release[95]. On the 1S diet,

there was no sex difference in GFP expression within the IPC (Figure S3F-S3H). To

confirm this, we directly monitored IPC activity using cell-attached electrophysiological

recordings. When we examined IPC spike rates, we found no significant differences

between males and females after a period of fasting (Figure 3E and 3F), or when the

fasting period was followed with refeeding on either whole food or D-glucose (Figure 3E

and 3F). These data suggest IPC activity is equivalent between the sexes, confirming

results from another recent study[96]. Given that Dilp3 levels in the IPC show a female bias,

these data suggest females have a greater quantity of Dilp3 available for secretion each

time the IPC are active.

To test whether the female bias in *dilp3* mRNA caused by dietary sugar was due to

an intrinsic difference in IPC, we expressed sex determination gene *transformer* (*tra*) in

male IPC and monitored *dilp3* mRNA levels. Normally, a functional Tra protein is

expressed only in females, where it directs most aspects of female sexual development[97-

100]. Because Tra expression in males specifies female development across multiple

tissues[101-107], this allows a female sexual identity to be conferred upon male IPC. In males

with IPC-specific expression of Tra, we observed no significant change in *dilp3* mRNA

levels (Figure S3I). This suggests the sexual identity of the IPC does not determine the

sex difference in *dilp3* mRNA levels. While the APCs have also been shown to influence

dilp3 mRNA levels in larvae[108], Tra expression in these cells similarly had no effect on the

sex difference in *dilp3* mRNA in adults (Figure S3J). Given that many groups of neurons

and neuropeptide-producing cells have been shown to influence *dilp3* mRNA levels in a

non-cell-autonomous manner[43,62, 109], we next used *elav-GAL4* to drive Tra expression

broadly across these cells. We found that Tra expression in all *elav-GAL4*-positive cells

caused a significant increase in male *dilp3* mRNA levels (Figure S3K), suggesting that Tra

acts in neurons or neuropeptide-producing cells other than the IPC to regulate the sex

difference in *dilp3* mRNA levels.

Given the female bias in *dilp3* mRNA and protein, and the known role of *dilp3* in

promoting IIS activity[108,110,111], we next compared IIS activity in males and females. To

assess IIS activity, we measured mRNA from known transcriptional targets of Forkhead

box, sub-group O (Foxo; FBgn0038197) in male and female abdominal carcasses. When

IIS activity is high, Foxo is phosphorylated and repressed, leading to lower mRNA levels of

known target genes *InR* and *bmm*[112-115]. In adult males and females maintained on 1S,

mRNA levels of Foxo target genes were significantly lower in females than in males

(Figure 3G and 3H). This suggests IIS activity is normally higher in females than in males.

Given that *Drosophila* female larvae are more insulin sensitive than male larvae[80], a sex

difference that also exists across many tissues in mammals[116,117] , we also compared

insulin sensitivity between adult virgin males and females kept on 1S. Considering female

abdominal carcasses showed significant insulin-induced repression of Foxo target genes

(Figure 3I and 3J), a response we did not observe in males (Figure 3K and 3L), our data

suggest female abdominal carcasses are more insulin sensitive than males. Taken

together, our data show that dietary sugar promotes higher expression of *dilp3* in females.

Increased *dilp3* mRNA and protein, combined with higher insulin sensitivity in females,

likely contribute to elevated IIS activity in females maintained on 1S.

Females require IIS to maintain fat storage

Insulin is a potent lipogenic hormone in multiple animals including *Drosophila*[53,118-122].

Increased IIS activity in female abdominal carcasses may therefore explain their higher fat

storage. Because reduced IIS activity during development significantly impairs larval

growth and survival to adulthood[54,103,110,111,123-125], and increased IIS activity during

development augments body size[103,125-127], we needed a system to modulate IIS activity

specifically in adults. Unfortunately, existing systems for inducible gene expression have

undesirable limitations for studying sex differences in fat metabolism. For example, the

Gene Switch system[128,129] induces transgene expression by feeding flies mifepristone,

which has sex-specific effects on adult physiology[130-132]. The AGES (Auxin-inducible Gene

Expression System) method uses auxin feeding to induce transgene expression[133], which

we and others recently showed affects feeding and triglyceride metabolism in a sex-biased

manner[134].

Another inducible system for gene expression, the temporal and regional gene

expression targeting (TARGET) system[135], normally requires long-term temperature shifts

at 29°C to inactivate temperature-sensitive repressor GAL80. Given that inactive GAL80

cannot repress GAL4, UAS transgene-mediated expression is initiated as long as the

animals are kept at 29°C. Unfortunately, 29°C is a temperature known to disrupt lipid

metabolism[136,137], making a long-term heat shock to induce transgene expression or

RNAi-mediated knockdown undesirable. To overcome this limitation, we reasoned that we

could deliver a brief 3 hr heat shock to newly-eclosed male and female flies to transiently

express proapoptotic gene *reaper* (*rpr*) specifically in the IPC (genotype: *dilp2-*

GAL4>UAS-rpr,tub-GAL80^ts). To validate expression of *dilp2-GAL4* in each sex, we first

confirmed that *dilp2-GAL4*-positive cells were morphologically similar to published

data[45,138-140], with no obvious differences between males and females (Figure S4A and

S4B). We also confirmed *dilp2-GAL4*-positive cells express Dilp3 (Figure S4C, S4C□,

S4C☐☐, S4D, S4D☐, S4D☐☐), in line with published data[43,138]. We next assessed whether

dilp2-GAL4 could be used to ablate the IPC. Importantly, mediated *rpr* overexpression is

an established way to ablate the IPC[123,141,142]. Based on published literature[110,111,123,142,143],

in adult heads the IPC are the only place that *dilp2* and *dilp3* mRNA are produced.

Therefore, we measured *dilp2* and *dilp3* mRNA levels as a direct way to test IPC ablation.

Supporting successful IPC ablation, mRNA levels of *dilp3* and *dilp2* were reduced in

adult heads from female and male flies after a brief heat shock compared with genotype-

matched controls without heat shock (Figure 4A, 4B, S4E, S4F). When we used anti-Dilp3

to check IPC ablation, many flies lacked Dilp3-positive cells in the *pars intercerebralis*

region of the brain where the IPC are located, indicating a complete IPC ablation in these

individuals (Figure S4G-S4J). In line with prior reports[141,142], some individuals had residual

Dilp3-producing cells, suggesting a partial IPC ablation (Figure S4K and S4L). Given that

our approach to fully/partially ablate the IPC caused phenotypes associated with IPC loss

and reduced insulin signaling such as an extremely long lifespan in both sexes (Figure 4C

and 4D)[141,144-148] and transiently reduced fertility in female but not male flies (Figure 4E

and 4F)[142], we conclude that our approach to IPC ablation is a valid method for assessing

how adult-specific reduction in IIS affects fat storage. While the reason for the transient

effect on female fertility remains unclear, previous studies show that changes to insulin

affect female pheromones[149] which could introduce a minor delay in egg-laying.

Using this system to ablate the IPC, we tested whether increased IIS was required

for females to store more body fat than males. We found that triglyceride levels were

significantly lower in 5-day-old females with ablated IPCs (genotype *dilp2-GAL4>UAS-*

rpr,tub-GAL80^{ts}) than in control females (genotypes *dilp2-GAL4>+,tub-GAL80^{ts}* and

+>UAS-rpr) (Figure 4G and S4M). In contrast, IPC ablation in males had no significant

effect on body fat (Figure 4G and S4M). We also found that females (genotype *dilp2-*

GAL4>UAS-rpr,tub-GAL80^{ts}) subjected to a brief heat shock had lower body fat compared

with genotype-matched females that were not heat-shocked (Figure 4H). While heat-

shocked *dilp2-GAL4>UAS-rpr,tub-GAL80*^{ts} males showed a decrease in body fat (Figure

4H), the magnitude of the decrease was greater in females (sex:genotype interaction

p=0.0071) (Figure S4N). The reduction in female body fat upon adult-specific loss of IPC

was not explained by the brief heat shocks, as control females (genotype *dilp2-*

GAL4>+,tub-GAL80^{ts}) subjected to an identical heat shock showed no significant effect on

body fat (Figure 4I).

The decrease in body fat was also not due to altered feeding in animals with IPC

ablation, as no significant differences in food consumption were present in either females

or males with IPC ablation compared with controls (Figure 4J and 4K). Instead, we identify

dilp3 as a key factor that contributes to higher female fat storage when dietary sugar is

present: Dilp3 overexpression in the IPC blocked the decrease in body fat normally

observed in adult females maintained on 0S (Figure S4O; diet:genotype interaction

p<0.0001). Dilp3 overexpression had no effect on sugar-induced changes to body fat in

males (diet:genotype interaction *p*=0.5588), possibly due to insulin resistance (Figure

S4P). Supporting a specific role for IIS in maintaining higher fat storage in females,

supplementing the diet of virgin females with an ecdysteroid called 20-hydroxyecdysone

(20HE) did not promote fat storage in females (Figure S4Q). Our data therefore reveal sex

differences in IIS regulation in adult flies, and demonstrate the importance of this

regulation to the male-female difference in fat storage.

Discussion

Our findings provide significant insight into the mechanisms by which adult females store

more fat than males. This information is critical to our understanding of how sex

differences in body fat are established and maintained, as past studies identified

mechanisms that operate in males with little effect in females[5,10]. Based on our data, we

propose a model in which higher consumption of dietary sugar in females leads to

elevated *dilp3* mRNA and protein levels, which act together with greater insulin sensitivity

to promote higher IIS activity and body fat in females. Supporting this model, reduced

sugar consumption or IPC ablation lessened the sex difference in fat storage by lowering

body fat in females with little effect in males, and Dilp3 overexpression prevented the

reduction in fat storage normally observed in females maintained on 0S. This reveals

previously unrecognized sex-specific IIS regulation in adult flies, and identifies Dilp3 and

IIS as key factors that support increased fat storage in adult females.

One important advance in our study was to extend knowledge of how food intake

affects physiology in unmated females. Past studies used mated females to show that

greater food consumption supports the quality and quantity of eggs produced[9,150-157]. While

greater total nutrient intake could explain these effects, specific macronutrients such as

dietary protein play individual roles in supporting egg production, egg-laying, and

postcopulatory behaviors in mated females[158-160]. Indeed, dietary protein is limiting for

Drosophila egg production[161], where its biological importance is shown by the fact that

mated females will adjust their food preferences to ensure adequate protein

intake[159,162,163]. Dietary protein does not hold the same physiological value for unmated

females, however, as virgin females do not adjust their preferences after protein

restriction[163]. Instead, our data suggest that dietary sugar is a key macronutrient for virgin

female physiology due to its effects on IIS activity and fat accumulation. Because IIS

activity supports normal germline stem cell proliferation and maintenance in the

ovary[54,142,161,164-167], and high levels of body fat are required for successful egg

production[12], dietary sugar consumption likely helps to establish a supportive metabolic

state prior to reproduction. Once mated, this role for dietary sugar becomes less important,

as reflected by the limited consumption of a sugar-rich diet in mated females compared

with males[168]. Thus, our data suggests that higher food intake in both virgin and mated

females provide key macronutrients to fuel the specific needs that accompany particular

life history stages (*e.g.* prior to mating or during egg production).

Another advance of our study was to reveal a profound sex difference in how

dietary sugar affects metabolism in the fat body. In females, sugar caused profound

remodeling of the lipidome and many changes in the metabolome. This sex difference was

most apparent when examining triglyceride, the lipid sub-class with the most obvious

sugar-dependent changes in female lipid abundance. However, females showed a sugar-

dependent increase across many lipid sub-classes (*e.g.*, DG, PG, PE, Acyl-Car). Given

that females have lower lipase activity than males[5], the presence of high diglyceride and

triglyceride levels suggest increased lipid synthesis in females. Diglyceride is also a

precursor for phospholipids such as PE and PC synthesis[1,169]. High levels of PE and PC in

females maintained on 1S further support higher lipid synthesis in females, a possibility

that will be important to test in future studies.

Beyond this broad lipid accumulation in females kept on 1S, other interesting

findings were that dietary sugar enhanced levels of a fatty acid precursor used for

synthesis of female-specific pheromones (28:2) and odd-chain fatty acids in the

triglyceride pool. Fatty acid 28:2 is the direct precursor for the synthesis of a cuticular

hydrocarbon called 7,11-heptacosadiene, the major female pheromone that mediates

female sexual attractiveness[170-172]. Thus, in addition to supporting IIS activity and fat

accumulation in females, dietary sugar consumption may also ensure females emit an

optimal pheromone profile. Indeed, sexual attractiveness is altered in females with

abnormal levels of cuticular hydrocarbons[173]. Given that the gut microbiota help produce

odd-chain fatty acids[174] and the microbe load is higher in adult female intestines than in

males[88], our data suggest that the intestinal microbiome may provide precursors used for

fat storage in females exposed to dietary sugar. Thus, dietary sugar provides an array of

physiological benefits for females to optimize reproduction.

In males, dietary sugar caused few changes in either lipids or metabolites. One

notable finding was that only one lipid subclass, TG-O, showed higher abundance in males

than in females. Ether-linked lipids such as TG-O are synthesized by a multi-step process

mediated by enzymes located both in the peroxisome and endoplasmic reticulum[175-177].

Because a recent study found high levels of TG-O in male flies chronically fed a high-

sugar diet[178], higher TG-O levels in males may indicate sugar-induced lipotoxicity. Indeed,

elevated TG-O levels were associated with metabolic dysfunction, as genetically reducing

production of ether-linked lipids in male flies reversed defects in glucose levels, cardiac

function, and insulin signaling caused by the high-sugar diet[178,179]. Future studies will

therefore need to examine ether-linked lipid synthesis in both sexes to determine whether

male-female differences exist in this process and its relationship to metabolic dysfunction.

Follow-up studies will also need to determine the ultimate reason that males largely fail to

adjust the lipidome in response to dietary sugar. Potential explanations suggested by our

data include insulin resistance, increased sugar excretion, and high lipase and lipolytic

pathway activity[5,10]. Published data suggest that greater carbohydrate breakdown within

the male gut to support spermatogenesis might also explain the female-biased lipidomic

response to dietary sugar[105].

Other key findings of our study include showing that dietary sugar promotes a

female bias in *dilp3* levels, and revealing a female bias in IIS activity within the adult fat

body. This extends knowledge from past studies on sex differences in IIS activity in third-

instar larvae[103] by showing this female bias in IIS activity is present in adults and is

physiologically significant. Given that elevated *dilp3* levels are known to promote body

fat[43,62,180] and prolong starvation resistance[61,181,182] , and that Dilp3 overexpression blocks

the reduction in body fat normally observed on the 0S diet, this female bias in *dilp3* is

significant for higher fat accumulation in females. However, a number of important future

directions remain. First, it will be important to determine whether Dilp3 signaling via the

insulin receptor drives all the lipidomic changes we observed due to dietary sugar

consumption, or whether other Dilps play a role. Second, it will be important in future

studies to reveal the mechanisms underlying the sex bias in *dilp3* levels. We identified sex

determination gene Tra as a key regulator of the sex difference in *dilp3* mRNA levels.

While it is clear that Tra does not act directly on the IPC or in the Akh-producing cells to

influence the sex difference in *dilp3* mRNA levels, the cell type in which Tra acts to specify

this difference remains unclear. Future studies will need to determine whether Tra acts in

cells that produce *dilp3*-regulatory factors such as gut-derived neuropeptide F[43],

octopamine[181], tachykinin[109], leucokinin[61,183], adiponectin receptor[184], Allatostatin A[45],

bursicon[62], drosulfakinin[182], and potentially Akh[108] and Limostatin[180,185] to regulate the sex

difference in *dilp3* mRNA levels, and whether this involves changes in food intake. Indeed,

sex differences in expression of select *dilp3*-regulatory factors have been reported[10,104,186],

where these factors have sex-specific effects on physiology[10,42-46,61,62,109,180,181] and

influence mating and/or reproductive behaviors[187-195].

Overall, our findings identify IIS as a major determinant of female fat storage, with

only minor effects on male body fat. This contrasts with males, where a set point of stored

fat is maintained by *bmm* and Akh, pathways with little effect on female body fat[5,10]. The

primary mechanisms that establish fat storage are therefore not identical between the

sexes: females sustain body fat primarily via the anabolic effects of insulin whereas body

fat in males is maintained by the lipolytic effects of *bmm* and Akh. Future studies will need

to determine whether the sex-biased effects of changing IIS and Akh on body fat are

reproduced in other aspects of physiology regulated by these pathways (*e.g.* locomotion,

lifespan, aggression, feeding, sleep)[64,141,142,149,152,196-199]. Future studies will also need to

determine whether the extensive crosstalk between IIS and Akh identified in single- and

mixed-sex studies in adults and larvae is shared between males and females[45,57,108,123,200].

Because IIS and Akh mediate responses to a variety of extrinsic and intrinsic cues, gaining

a better understanding of the interplay between these two major hormone pathways in

males and females will provide critical insights into how diverse cues are integrated to

maintain whole-body triglyceride homeostasis in each sex. Given that sex differences in

sucrose-induced changes to mammalian adipose tissue lipolysis, hepatic lipogenesis, and

adipose tissue gene expression broadly reproduce the effects we report in flies[201], the sex-

biased effects of IIS and Akh on body fat we observed have implications for our

understanding of sex differences in parallel hormone systems in mammals (*e.g.* insulin,

glucagon) and the sex-biased risk of diseases associated with these hormones[202-206].

Limitations of the study. Our study has several limitations. First, the limitations of current

Drosophila inducible gene expression systems mean that we overexpressed *dilp3*

throughout development instead of just in adults. We therefore cannot rule out the

possibility that Dilp3 acts in development to influence adult fat storage. Second, while we

defined a role for Dilp3 in regulating the sex difference in body fat, we cannot rule out a

role for other Dilps.

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to, and

will be fulfilled by, lead contact Elizabeth J. Rideout (elizabeth.rideout@ubc.ca).

Materials availability

This study did not generate new unique reagents.

Data and code availability

• Fly lipidomic and metabolomic data are available in Supplemental Table S1.

• This paper does not report original code.

• Any additional information required to reanalyze the data reported in this paper is

available from the lead contact upon request.

Acknowledgments. This study was supported by operating grants to EJR from the

Canadian Institutes for Health Research (PJT-153072 and PJT-183786), CIHR Sex and

Gender Science chair program (GS4-171365), Michael Smith Foundation for Health

Research (16876), and the Canadian Foundation for Innovation (JELF-34879). PB was

supported by a 4-year CELL fellowship from UBC. LWW was supported by a British

Columbia Graduate Scholarship Award. TH/HY were supported by Natural Sciences and

Engineering Research Council of Canada (NSERC; 2020-04895). MDG and MS were

supported by NSERC (RGPIN-2016-03857). We used FlyBase to find information, stocks,

and literature for our project. FlyBase is supported by a grant from the National Human

Genome Research Institute at the U.S. National Institutes of Health (U41 HG000739) and

by the British Medical Research Council (MR/N030117/1). Stocks obtained from the

Bloomington *Drosophila* Stock Center (NIHP40OD018537) were used in this study. The

authors thank the TRiP at Harvard Medical School (NIH/NIGMS R01-GM084947) for

providing transgenic RNAi fly stocks and/or plasmid vectors used in this study. We also

thank Dr. Jan Veenstra (University of Bordeaux) and Dr. Patrick Callaerts (University of

Leuven) for providing anti-Dilp3 antibody, and Dr. Gaiti Hasan (National Centre for

Biological Science, India) for providing *UAS-dilp3* flies. The graphical abstract was

prepared using BioRender (https://BioRender.com). The funders had no role in study

design, data collection and analysis, decision to publish, or preparation of the manuscript.

We thank members of the Rideout lab for valuable feedback. We acknowledge that our

research takes place on the traditional, ancestral, and unceded territory of the Musqueam

people; a privilege for which we are grateful.

Author contributions

P. B. conceived studies, collected samples, conducted experiments, interpreted

experiments, wrote and edited the manuscript.

J. A. B. and J. B. L. performed dissections and collected data related to brain anti-Dilp3

immunofluorescence.

H. Y. performed lipidomic and metabolomic experiments.

L. W. W. performed ecdysone experiment.

C. J. M. performed brain dissections and monitored IPC activity using the CaLexA system.

M. D. G. edited the manuscript and acquired funding for electrophysiological recordings of

IPC.

T. H. edited the manuscript and acquired funding for lipidomic and metabolomic analyses.

M. S. conceived studies, analyzed data, interpreted experiments, and contributed to writing

and editing of the manuscript related to electrophysiological recordings of IPC.

E. J. R. conceived studies, interpreted experiments, wrote and edited the manuscript,

acquired funding, and is the guarantor of this work.

Declaration of interests

The authors declare no competing interests.

Supplemental information

Document S1. Supplemental Figures S1-S4

Table S1. Excel file containing raw data with calculations

Table S2. Excel file containing statistics for all data

Figure Legends

Figure 1. Dietary sugar promotes increased fat storage in adult females.

(A) Food consumption per unit of body weight was significantly higher in 5-day-old adult

Canton-S (CS) females than males (p=0.0072; Student's *t*-test); n=8 biological replicates.

(B) Whole-body triglyceride levels were significantly lower in 5-day-old *CS* females

maintained on a diet with 20% of normal nutrient content compared with females kept on

our normal diet (100% diet; $p<0.0001$). No change in whole-body triglyceride was

observed in male flies ($p>0.9999$; sex:diet interaction $p<0.0001$). Two-way ANOVA

followed by Bonferroni post-hoc test; n=16 biological replicates.

(B') Sex difference in body fat represented as a percent of the male-female difference in

our normal diet.

(C) Whole-body triglyceride levels were significantly higher in 5-day-old adult w^{1118} females

and males maintained on the 1S diet compared with sex-matched flies kept on the 0S diet

(Females: $p<0.0001$; Males: $p=0.0068$). The magnitude of the decrease in body fat was

greater in females than in males (sex:diet interaction $p<0.0001$). Two-way ANOVA

followed by Tukey HSD on data processed using the aligned rank transform for non-

parametric data; n=16 biological replicates.

(C') Sex difference in body fat represented as a percent of the male-female difference in

the 1S diet.

(D) Whole-body triglyceride levels were significantly higher in 5-day-old adult w^{1118} females

maintained on sugar-containing diet with reduced calories (1S-LC) compared with females

kept on the 0S diet (same calorie content as 1S-LC) ($p=0.0032$); no significant change in

body fat was observed in males ($p=0.3715$; sex:diet interaction $p=0.0019$). Two-way

ANOVA followed by Bonferroni post-hoc test; n=8 biological replicates.

(D') Sex difference in body fat represented as a percent of the male-female difference in

the lower-calorie 1S diet.

(E) Whole-body triglyceride levels were not significantly different between 5-day-old adult

w^{1118} females maintained on the 0S diet and sex-matched flies kept on the 0S diet with

either added 20.5 g/L of sucrose or 70.9 g/L of D-glucose diet (females: $p=0.9998$

(sucrose), p=0.7532 (D-glucose). One-way ANOVA followed by Tukey HSD test; n=8

biological replicates.

(F) Whole-body triglyceride levels were significantly higher in 5-day-old adult w^{1118} females

maintained on our lab diet with 30% added sugar (HS) compared with females kept on the

1S diet (p=0.0036; Student's t-test). n=8 biological replicates.

All data plotted as mean ± SEM. ns indicates not significant with p>0.05; * p<0.05, **

p<0.01, *** p<0.001, **** p<0.0001.

See also Figure S1.

Figure 2. Female-biased effect of dietary sugar on metabolism.

(A) Volcano plot of lipid species that are differentially regulated between 5-day-old w^{1118}

adult female and male fat bodies in 1S diet (colored dots represent p<0.01; Student's t-

test).

(B) Number of lipid species within each sub-class that show higher abundance in either

female (purple) or male (green) 5-day-old adult fat bodies in 1S diet (p<0.01; Student's t-

test). For example, 48/57 TG species are more abundant in females than in males,

whereas 12/13 TG-O species are more abundant in males than in females.

(C) Sex bias of all metabolites with differential abundance between female and male fat

bodies in 1S diet (p<0.05; Student's t-test).

(D) Volcano plot of lipid species that are differentially regulated when females are

maintained on a diet without sugar (colored dots represent p<0.01; Student's t-test).

(E) Volcano plot of lipid species that are differentially regulated when males are

maintained on a diet without sugar (colored dots represent p<0.01; Student's t-test).

(F) Volcano plot of metabolites that are differentially regulated when females are

maintained on a diet without sugar (colored dots represent p<0.05; Student's t-test).

(G) Volcano plot of metabolites that are differentially regulated when males are maintained

on a diet without sugar (colored dots represent $p<0.05$; Student's t-test).

(H) Glucose levels in the excreta of 6-day-old adult w^{1118} males were higher than females

($p=0.0461$, Student's t-test); n=4-6 biological replicates. Data plotted as mean ± SEM. *

$p<0.05$

TG: triglyceride; TG-O: ether-linked TG; DG: diglyceride; PI: phosphatidylinositol; PC:

phosphatidylcholine; PC-O: ether-linked PC; ST: sterol; LPC: lyso-PC; PG:

phosphatidylglycerol; PS: phosphatidylserine; PE: phosphatidylethanolamine; PE-O or PE-

P: ether-linked PE; NAE: N-acetylethanolamine; Cer: Ceramide; Acyl-Car: acyl-carnitine;

SE: sterol ester; Hex-Cer: hexosyl-ceramide.

See also Figure S2.

Figure 3. Dietary sugar promotes sex differences in IIS regulation.

(A) mRNA levels of IPC-derived *dilp3* in heads were significantly higher in 5-day-old w^{1118}

females compared with males ($p<0.0001$; Student's t-test); n=7 biological replicates.

(B, C) mRNA levels of IPC-derived *dilp2* (B) and *dilp5* (C) in heads were not significantly

different between 5-day-old w^{1118} males and females ($p=0.1282$, Mann-Whitney test

(*dilp2*); $p=0.5753$, Student's t-test (*dilp5*)); n=7 biological replicates.

(D) mRNA levels of IPC-derived *dilp3* in heads were significantly lower in females

maintained on the 0S diet compared to females kept on the 1S diet ($p<0.0001$). There was

no sugar-dependent change in *dilp3* mRNA in males ($p=0.4809$; sex:diet interaction

$p=0.0128$). Two-way ANOVA followed by Bonferroni post-hoc test; n=7 biological

replicates. Note: *dilp3* mRNA levels in 1S diet were replotted from Figure 3A as these data

were collected in the same experiment.

(E) Representative cell-attached recordings of IPC activity in 4-8-day-old female (left) and

male (right) flies that were either starved for 24 hr (top), re-fed with D-glucose (middle), or

re-fed with whole food (bottom).

(F) Quantification of cell-attached recordings of IPC activity in 4-8-day-old male and female

flies that were either starved for 24 hr, re-fed with D-glucose, or re-fed with whole food.

n=10-15 individuals per sex and per condition; error bars indicate SEM.

(G, H) mRNA levels of Foxo target genes-*InR* (G) *and bmm* (H) in fat bodies were

significantly higher in 5-day-old w^{1118} males compared to females in 1S diet (p=0.0077

(*InR*); p=0.0022 (*bmm*) Student's *t*-test); n=7 biological replicates.

(I, J) mRNA levels of Foxo target genes-*InR* (I) and *bmm* (J) in 5-day-old w^{1118} female fat

bodies were significantly reduced after stimulation with 1 µM insulin compared with fat

bodies lacking insulin stimulation (p=0.0208 (*InR*); p=0.0100 (*bmm*); Student's *t*-test); n=6

biological replicates.

(K, L) mRNA levels of Foxo target genes-*InR* (K) and *bmm* (L) in 5-day-old w^{1118} male fat

bodies were not significantly different between unstimulated and insulin-stimulated

conditions (p=0.2194 (*InR*); p=0.7649 (*bmm*); Student's *t*-test); n=6 biological replicates.

Data plotted as mean ± SEM. ns indicates not significant with p>0.05; * p<0.05, ** p<0.01,

*** p<0.001, **** p<0.0001.

See also Figure S3.

Figure 4. Females require IIS to maintain fat storage.

(A, B) mRNA levels of IPC-derived *dilp3* in heads were significantly lower in both female

(A; p=0.0005, Student's *t*-test) and male (B; p=0.0092, Student's *t*-test) *dilp2-GAL4>UAS-*

rpr,tub-GAL80^{ts} flies with heat shock at 29°C compared to sex-matched flies reared in

room temperature; n=4-6 biological replicates.

(C) Lifespan was significantly longer in *dilp2-GAL4>UAS-rpr,tub-GAL80^{ts}* females

compared with *dilp2-GAL4>+,tub-GAL80^{ts}* and *+>UAS-rpr* controls ($p<0.0001$ and

$p<0.0001$, respectively; Log-rank test); n=100-213 females.

(D) Lifespan was significantly longer in *dilp2-GAL4>UAS-rpr,tub-GAL80^{ts}* males compared

with *dilp2-GAL4>+,tub-GAL80^{ts}* and *+>UAS-rpr* controls ($p<0.0001$ and $p<0.0001$,

respectively; Log-rank test); n=100-148 males.

(E) The number of eggs laid by females in the first 24-hour period post-mating that

developed into pupae was significantly lower in *dilp2-GAL4>UAS-rpr,tub-GAL80^{ts}* females

compared with both *dilp2-GAL4>+,tub-GAL80^{ts}* and *+>UAS-rpr* controls ($p<0.0001$ and

$p<0.0001$, respectively; one-way ANOVA followed by Tukey HSD). n=30 females.

(F) The number of eggs laid by females mated to *dilp2-GAL4>UAS-rpr; tub-GAL80^{ts}* males

were significantly lower compared to control *dilp2-GAL4>+; tub-GAL80^{ts}* but no change

compared to *+>UAS-rpr* control males that developed into pupae over the first 24 hr of

after 24 hr mating ($p=0.0032$ and $p=0.7420$, respectively; one-way ANOVA followed by

Tukey HSD). n=22-30 males.

(G) Whole-body triglyceride levels were significantly lower in *dilp2-GAL4>UAS-rpr,tub-*

GAL80^{ts} females compared with both *dilp2-GAL4>+,tub-GAL80^{ts}* and *+>UAS-rpr* control

females ($p<0.0001$ and $p<0.0001$, respectively). Whole-body triglyceride levels were not

significantly different in *dilp2-GAL4>UAS-rpr,tub-GAL80^{ts}* males compared with *dilp2-*

GAL4>+,tub-GAL80^{ts} control males but significantly different from *+>UAS-rpr* control males

($p=0.9397$ and $p=0.0022$, respectively) (sex:genotype interaction $p=0.0002$). Two-way

ANOVA followed by Bonferroni post-hoc test; n=8 biological replicates.

(H) Whole-body triglyceride levels were significantly lower in female *dilp2-GAL4>UAS-*

rpr,tub-GAL80^{ts} after heat shock compared with genotype-matched females without heat

shock ($p<0.0001$). A reduction in whole-body triglyceride levels was also observed in

males ($p=0.0012$); however, the magnitude of the decrease was smaller than in females

(sex:genotype interaction p=0.0071). Two-way ANOVA followed by Bonferroni post-hoc

test; n=16 biological replicates.

(I) Whole-body triglyceride levels were not significantly different between *dilp2-*

GAL4>+,tub-GAL80^{ts} females that received a brief heat shock compared with *dilp2-*

GAL4>+,tub-GAL80^{ts} females that received no heat shock (p=0.8107; Student's *t*-test);

n=8 biological replicates.

(J, K) Food consumption was not significantly different in either females (J) or males (K)

between *dilp2-GAL4>UAS-rpr,tub-GAL80*^{ts} flies and *dilp2-GAL4>+,tub-GAL80*^{ts} or *+>UAS-*

rpr controls (females: p=0.9999 [GAL4 control] and p=0.9318 [UAS control]; males:

p=0.6516 [GAL4 control] and p=0.0628 [UAS control]). One-way ANOVA followed by

Tukey HSD; n=8-9 biological replicates.

All data in (A, B, E-K) plotted as mean ± SEM. ns indicates not significant with p>0.05, *

p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001.

See also Figure S4.

STAR⬚Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-Dilp3	Dr. Patrick Callaerts; Dr. Jan Veenstra	[94,207]
Goat anti-Rabbit Alexa Fluor 488	Invitrogen	Cat#A11008
Chemicals, peptides, and recombinant proteins		
Water-soluble yeast extract	Sigma-Aldrich	Cat#Y1625
Tween 20	Amresco	Cat#0777
1xPBS	Sigma-Aldrich	Cat#P5493
5% blue dye	Ward's Science	Cat#470300-996
Trizol	Thermo Fisher Scientific	Cat#15596018
Molecular Biology Grade Water	Corning	Cat#46-000-CV
16% paraformaldehyde	Electron Microscopy Sciences	Cat#15710
Hoechst	Sigma-Aldrich	Cat#33342
Schneider's *Drosophila* solution	Thermo Scientific	Cat#21720024
Recombinant human insulin	Sigma-Aldrich	Cat#I2643
Sodium bicarbonate	Sigma-Aldrich	Cat#S6014
NaCl	Sigma-Aldrich	Cat#S7653
Sodium phosphate	Millipore Sigma	Cat#567545
HEPES	Fisher Scientific	Cat#BP310-100
Ribose	Sigma-Aldrich	Cat#1603108
$CaCl_2$	Sigma-Aldrich	Cat#C7902
$MgCl_2$	Sigma-Aldrich	Cat#M9272
KCl	Sigma-Aldrich	Cat#P9541
L-glucose	Biosynth	Cat#MG05247
20-hydroxyecdysone	abcam	Cat#ab142425
Goat normal serum	Sigma-Aldrich	Cat#S26
Triton X-100	Amresco	Cat#M143
SlowFade Diamond Antifade Mountant	Invitrogen	Cat#S36972
Critical commercial assays		
Free glycerol reagent	Sigma-Aldrich	Cat#F6428
Triglyceride reagent	Sigma-Aldrich	Cat#T2449
Glycerol standard	Sigma-Aldrich	Cat#G7793
Triglyceride standard	Stanbio	Cat#FT7610
Enzymatic Triglyceride Reagent	Stanbio	Cat#BD386a/d
Triglyceride Activator	Stanbio	Cat#BD386b
Glucose reagent	Sigma-Aldrich	Cat#GAHK20
QuantiTect Reverse Transcription Kit	Qiagen	Cat#205314
Citrate assay kit	Sigma-Aldrich	Cat#MAK333
Bradford protein assay kit	Bio-Rad	Cat#5000204
Experimental models: organisms/strains		

D. melanogaster: Canton-S	Bloomington *Drosophila* Stock Center (BDSC)	RRID: BDSC_64349
D. melanogaster: w[1118]	BDSC	RRID: BDSC_3605
D. melanogaster: dilp2-GAL4	BDSC	RRID: BDSC_37516
D. melanogaster: UAS-reaper	BDSC	RRID: BDSC_5823
D. melanogaster: UAS-CaLexA	BDSC	RRID: BDSC_66542
D. melanogaster: tubP-GAL80[ts]	BDSC	RRID: BDSC_7018
D. melanogaster: UAS-mRFP	Dr. Douglas Allan	N/A
D. melanogaster: UAS-mCD8::GFP	BDSC	RRID: BDSC_5137
D. melanogaster: elav-GAL4	BDSC	RRID: BDSC_458
D. melanogaster: UAS-tra[F]	BDSC	RRID: BDSC_4590
D. melanogaster: UAS-dilp3	Dr. Gaiti Hasan	[111,208]
D. melanogaster: Akh-GAL4	Dr. Michael Gordon	N/A
D. melanogaster: UAS-nRFP	BDSC	RRID: BDSC_30556
Oligonucleotides		
Primer: *β tubulin*-Forward: ATCATCACACACGGACAGGA	This study	N/A
Primer: *β tubulin*-Reverse: GAGCTGGATGATGGGGAGTA	This study	N/A
Primer: *Actin5C*-Forward: TTGTCTGGGCAAGAGGATCAG	This study	N/A
Primer: *Actin5C*-Reverse: ACCACTCGCACTTGCACTTTC	This study	N/A
Primer: *InR*-Forward: GCTGCATCTCCTGTCGAAAT	This study	N/A
Primer: *InR*-Reverse: CGTTGGACAGTGGGTGATAC	This study	N/A
Primer: *bmm*-Forward: GTCCCTTCAGTCCCTCCTTC	This study	N/A
Primer: *bmm*-Reverse: TATGAAGCACGCACACAACA	This study	N/A
Primer: *dilp2*-Forward: CCGGCTCGACCCAACTTAAT	This study	N/A
Primer: *dilp2*-Reverse: GCACACCATACTCAGCACCT	This study	N/A
Primer: *dilp3*-Forward: AGAGAACTTTGGACCCCGTGAA	This study	N/A
Primer: *dilp3*-Reverse: TGAACCGAACTATCACTCAACAGTCT	This study	N/A

Primer: *dilp5*-Forward: GAGGCACCTTGGGCCTATTC	This study	N/A
Primer: *dilp5*-Reverse: CATGTGGTGAGATTCGGAGCTA	This study	N/A
Software and algorithms		
ImageJ	NIH Image[209]	https://imagej.net/ImageJ
Prism 10	Graphpad	https://www.graphpad.com/
MAFFIN R package[210]		
pCLAMP Software Suite	Molecular Devices	https://www.moleculardevices.com

Experimental model and study participant details

Fly husbandry

Detailed lists of strains are provided in the key resources table. The recipe for our regular lab diet (1S) is as follows: 20.5 g/L sucrose, 70.9 g/L D-glucose, 48.5 g/L cornmeal, 45.3 g/L yeast, 4.55 g/L agar, 0.5g CaCl2•2H2O, 0.5 g MgSO4•7H2O, 11.77 mL/L acid mix (propionic acid/phosphoric acid). To make 1 L of 20% food, 800 mL 0.7% agar was mixed with 200 mL of 1S food. The low-sugar diet (0S) was made by leaving out sucrose and D-glucose from the 1S diet. The 1S diet was calorie-matched to 0S (1S-low calorie) by removing 8.6 g/L cornmeal from the 1S diet. For diets with sugar added back, either 20.5 g/L sucrose (0S+sucrose) or 70.9 g/L D-glucose (0S+D-glucose) were added to 0S, as this reflects the usual quantity of each sugar in the 1S diet. To monitor the effect of non-nutritive sugar on body fat, we added 62.5 g/L L-glucose (0S+L-glucose; Biosynth, MG05247) to the 0S diet[87]. To make a 30% high sugar diet, 300 g/L extra sucrose was added to the 1S diet, in line with published work[211,212]. For ecdysone-supplemented food, 0.5 mg/mL of 20-hydroxyecdysone (20-HE) in ethanol was added to the food (abcam, ab142425). For control food, the same ethanol volume without 20-HE was added to the food. For all experiments, we allowed female flies to lay eggs on grape juice agar plates for 12 hr. At 24 hr after egg laying, 50 larvae were picked into vials containing 10 mL food

and reared at 22°C. Males and females were distinguished by the presence of sex combs

late in the pupal period and placed into single-sex vials to eclose. After eclosion, adult flies

were maintained at a density of twenty flies per vial in single-sex groups. Unless otherwise

stated, all experiments used 5- to 7-day-old virgin flies.

Adult weight

Groups of 10 flies were placed in preweighed 1.5 ml tubes (Diamed Lab Supplies,

DIATEC610-2550) and weighed on an analytical balance (Mettler-Toledo, ME104).

Method details

Capillary feeder (CAFE) assay

One biological replicate consisted of measurements from one group of 10 flies. Flies were

placed into a specialized 15 ml conical vial with access to two capillary tubes (A-M

Systems; 626000), as previously described[87]. Briefly, capillary tubes were filled with fly

food media containing nutrients equivalent to the 1S diet (20.5 g/L sucrose, 70.9 g/L D-

glucose, 45.3 g/L yeast extract (Sigma-Aldrich; Y1625) in water) or 0S diet (45.3 g/L yeast

extract in water). To prevent evaporation, approximately 0.5 µl of mineral oil was added to

the top of each capillary tube. The vials were placed into fitted holes in the lid of a large

plastic container. To maintain high humidity throughout the experiment, a shallow layer of

water was poured into the base of the container. The meniscus of the fly food media was

marked before the start of the experiment and again after 24 hr. The distance between the

marks was used to quantify the volume of fly food media that was consumed by the 10

flies (1 mm=0.15 µl). To account for sex differences in body size, the volume of fly food

consumed was normalized to the weight of individual flies.

Whole-body triglyceride measurements

One biological replicate consisted of five flies. Flies were collected in a 1.5 ml tube and

homogenized in 200-350 µl of 0.1% Tween (Amresco, 0777-1L) in 1X phosphate-buffered

saline (PBS; Sigma-Aldrich, P5493) using 50 µl of glass beads (Sigma-Aldrich, Z250473)

agitated at 8 m/s for 5 s (OMNI International BeadRuptor 24). Triglyceride concentration

was measured using either Sigma kit (G7793, T2449, F6428) or Stanbio kit (FT7610,

BD386a/d, BD386b) according to the manufacturer's instructions and as described

previously[4,5] with the following modifications. For Sigma kit, 30 µl of homogenate was

transferred in duplicate to a 96-well plate and heat treated at 70°C for 10 min (Bio-Rad

T100 Thermal Cycler). Standard curve was prepared using glycerol standard (Sigma,

G7793) for concentrations of 0 mg/ml, 0.125 mg/ml, 0.25 mg/ml, 0.5 mg/ml, 1 mg/ml, and

2 mg/ml. After heat treatment, 30 µl of each concentration of standard solution was

transferred to the same 96-well plate in duplicate. 30 µl of triglyceride reagent (Sigma,

T2449) was added to 1 set of samples/standard solutions and 0.1% Tween in 1x PBS was

added to the other set of samples/standard solutions. After 40 min incubation at 37°C, the

96-well plate was centrifuged at 3800 rpm for 3 min at 4°C (VWR® Benchtop General

Purpose Centrifuges, 10830-744) to precipitate cellular debris. 7.5 µl of supernatant was

transferred to a 384-well plate-reader plate and 25 µl of free glycerol reagent (Sigma,

F6428) was added to each well. Plate was incubated at 37°C for 5 min, then absorbance

was read at 540 nm (Thermo Scientific – Multiskan FC Microplate Photometer). For the

Stanbio kit, 10 µl of either homogenate or triglyceride standard (FT7610) was added to 190

µl of activated triglyceride reagent (Enzymatic Triglyceride Reagent, BD386a/d;

Triglyceride Activator, Cat. No. BD386b) in the 96-well plate. After 15 min incubation at

room temperature, the absorbance was read at 540 nm. Triglyceride concentration was

used to calculate percent body fat.

Lipidomics and metabolomics

Sample preparation

Abdominal carcasses were isolated from 5-day-old male and female w^{1118} flies maintained

on the 1S or 0S diet after eclosion. One biological replicate consisted of 30 abdominal

carcasses dissected from 5- to 7-day-old adult males and females; a total of 9 biological

replicates were collected for each sex and condition. After dissection, each carcass was

snap-frozen on dry ice in a 2 mL screw-cap microcentrifuge tube (Thermo Fisher Scientific,

Waltham, MA) and stored at –80˚C until further processing. For a dual-phase extraction,

samples were homogenized with a Mini-Beadbeater-24 (Biospec Products, Bartlesville,

United States) at 3800 rpm. A total of 333 µl 90% MeOH was added and two 15 sec

homogenization cycles were performed with a 1 min rest on dry ice in between. The

homogenized solution was mixed with 1000 µl of methyl tert-butyl ether and shaken for 5

min. To induce the phase separation, 317 µl of water was added. After mixing, the solution

rested at room temperature for 10 min and then the two-layer solution was centrifuged at

14,000 rpm for 15 min at 4°C. For LC-MS analysis, the clear upper and lower layers were

separated into new vials, dried in a SpeedVac at 20°C, and reconstituted in 70 µl of

acetonitrile and isopropanol (1:1, v/v) and 70 µl of acetonitrile and water (1:1, v/v) mixed

solvent, respectively. A method blank sample was also prepared following the same

protocol but without *Drosophila* abdomens.

LC-MS/MS experiment

For metabolomics, hydrophilic interaction chromatography (HILIC) separation was

performed on a SeQuant ZIC-HILIC column (50 mm × 2.1 mm, 5 µm, 200 Å) with an

elution time of 8 min. Mobile phase A was 5% acetonitrile in water with 10 mM ammonium

acetate (Thermo Fisher Scientific) with pH = 9.8, adjusted by ammonium hydroxide

(MilliporeSigma), and mobile phase B was 5% water in acetonitrile with no buffer. The LC

elution gradient was optimized using the BAGO software[189] as follows: 0 min, 10% A; 1

min, 10% A; 2 min, 20% A; 5 min, 20% A; 6 min, 45% A; 7 min, 95% A; 8 min, 95% A. For

lipidomics, reversed phase (RP) separation was achieved on a Waters UPLC Acquity BEH

C18 Column (1.0 mm ×100 mm, 1.7 µm, 130 Å, Milford, MA, USA) with an elution time of

25 min. Mobile phase A was acetonitrile and water (6:4, v/v) with 2 mM ammonium

formate (pH = 4.8, adjusted by formic acid), and mobile phase B was isopropanol and

acetonitrile (9:1, v/v). LC elution gradient was optimized using the BAGO software as

follows: 0 min, 5% B; 1 min, 5% B; 4 min, 35% B; 7 min, 40% B; 10 min, 50% B; 13 min,

55% B; 16 min, 55% B; 19 min, 75% B; 22 min, 75% B; 25 min, 95% B.

The MS was operated in electrospray ionization (ESI) positive mode. The ESI

source conditions were set as follows: dry gas temperature, 220°C; dry gas flow, 7 L/min;

nebulizer gas pressure, 1.6 bar; capillary voltage, 4500 V for positive mode. The MS1

analysis was conducted using following parameters: mass range, 70-1000 *m/z*; spectrum

type: centroid, calculated using maximum intensity; absolute intensity threshold: 250.

Data-dependent MS/MS analysis parameters: collision energy: 16-30 eV; cycle time: 3 s;

spectra rate: 4 Hz when intensity $< 10^4$ and 12 Hz when intensity $> 10^5$, linearly increased

from 10^4 to 10^5. External calibration was applied using sodium formate to ensure the *m/z*

accuracy before sample analysis.

Data processing

The raw MS data were converted to ABF format in Reifycs Abf Converter (ver. 4.0.0). The

converted data were processed in MS-DIAL (ver. 4.90) for feature extraction, feature

alignment, and compound identification. The data processing parameters in MS-DIAL were

set as follows: MS1 tolerance, 0.01 Da; MS/MS tolerance, 0.05 Da; mass slice width, 0.05

Da; smoothing method, linear weighted moving average; smoothing level, 3 scans;

minimum peak width, 5 scans. Compound identification was achieved using NIST 20 for

metabolomics and the internal library within MS-DIAL for lipidomics. Post-acquisition

sample normalization was performed using the *MAFFIN* R package[210] to ensure fair

quantitative comparison.

Citrate assay

For females, one biological replicate consisted of ten fat bodies; for males one biological

replicate consisted of fifteen fat bodies. Tissues were collected in a 1.5 ml tube and

homogenized in 100 µl of 1XPBS (PBS; Sigma-Aldrich, P5493) using 50 µl of glass beads

(Sigma-Aldrich, Z250473) agitated at 8 m/s for 5 s (OMNI International BeadRuptor 24).

Citrate concentration was measured using citrate assay kit (Sigma-Aldrich MAK333)

according to the manufacturer's instructions. Briefly, the homogenized solution was

centrifuged at 14,000 rpm for 5 min at 4°C (VWR® Benchtop General Purpose

Centrifuges, 10830-744) to precipitate cellular debris. 20 µl of supernatant was transferred

in duplicate to a clear flat bottom 96-well plate, to which either 80 µl of reaction mix (85 µl

of developer (#MAK333A) + 1 µl of CL enzyme solution (#MAK333B) + 1 µl of dye reagent

(#MAK333C) + 1 µl of ODC enzyme (#MAK333D)) or blank reaction mix (85 µl of

developer (#MAK333A) + 1 µl of dye reagent (#MAK333C) + 1 µl of ODC enzyme

(#MAK333D)) were added. Standard curve was prepared using citrate standard

(#MAK333E) for concentrations of 0 µM, 30 µM, 60 µM, 120 µM, 240 µM, and 400 µM. 20

µL of each standard solution was added to the 96 well plate, to which 80 µl of reaction mix

was added. The whole plate was incubated at room temperature for 15 min and the

absorbance was measured at 570 nm. The citrate levels were normalized to the protein

concentration of each sample. Protein concentration was determined by Bradford assay

(Bio-Rad #5000204), as we described previously[5].

Excrement glucose measurements

One biological replicate consisted of 30 female flies or 60 male flies. Flies were fed with 1S

diet containing 5% blue dye (FD&C Blue no.1, Ward's Science, 470300-996) for 3.5 hr,

and subsequently transferred into a 1.5 mL tube for excrement collection at room

temperature. After 6 hr in the 1.5 mL tube, flies were discarded and 100 µL of 1x PBS was

added to each tube; tubes were vortexed to dissolve and pool all the blue-stained

excrement deposited in the tube. For blue dye measurement, 10 µL of the excrement was

added to 10 µL of 1x PBS in 384-well plate and absorbance was measured at 630 nm

(Thermo Scientific – Multiskan FC Microplate Photometer). The rest of the excreta (90 µL)

was added to 110 µL of Glucose Reagent (Sigma-Aldrich, GAHK20) in 96-well plate and

incubated overnight at 37 °C for measurement of glucose levels at 340 nm (Tecan Austria

GmbH-Tecan Spark). Glucose content in the excrement was normalized with blue dye to

account for sex differences in excreta volume, and to account for the number of flies that

deposited the excreta[213].

RNA extraction, cDNA synthesis, and Quantitative real-time PCR (qPCR)

One biological replicate consisted of 10 adult heads or 15 adult fat bodies. Samples were

homogenized in 500 µl of Trizol (Thermo Fisher Scientific; 15596018) and precipitated

using isopropanol and 75% ethanol to extract RNA. Pelleted RNA was resuspended using

20-30 µl of molecular biology grade water (Corning, 46-000-CV) and stored at -80°C until

use. Each experiment contained 4-7 biological replicates per sex, per genotype, and per

diet; each experiment was repeated twice.

For cDNA synthesis, an equal amount of RNA per reaction was DNase-treated and

reverse transcribed according to manufacturer's instructions using the QuantiTect Reverse

Transcription Kit (Qiagen, 205314). Relative mRNA transcript levels were quantified using

qPCR as described previously[5]. Data were normalized to the average fold change of

Actin5C and *β-tubulin*.

Immunostaining

Brains were dissected in cold 1× PBS and fixed in fresh 4% paraformaldehyde at room

temperature for 40 min. The brains were washed with 1× PBS three times and washed in

PBST (PBS + 0.1% Triton X-100) for 10 min. The samples were blocked for 1 hr at 4°C in

PBST + 3% goat normal serum (Sigma-Aldrich; S26). Samples were incubated with rabbit

anti-Dilp3 (1:200)[208] in blocking solution overnight at 4°C with gentle agitation. Samples

were washed three times with PBST and then incubated with secondary antibody (goat

anti-rabbit Alexa Fluor 488; 1:500) overnight at 4°C with gentle agitation. Samples were

washed with PBST three times, with Hoescht added to the final PBST wash for 30 min

(Sigma-Aldrich 33342, 1:1000). Brains were washed for 10 min in 1× PBS, mounted using

SlowFade Diamond Antifade mountant (Invitrogen; S36972), and imaged using a Leica

SP5 laser scanning confocal microscope. To quantify signal, we manually segmented IPC

clusters and used Z-Project "sum slices" method, measured RawIntDen of IPC clusters

and background after subtracting background RawIntDen. Signal was normalized to IPC

size and to number of slices. To quantify anti-Dilp3 levels 16-21 brain were analyzed per

sex and per condition.

IPC morphology

Brains were dissected in cold 1× PBS and fixed in fresh 4% paraformaldehyde at room

temperature for 40 min. Brains were washed with 1× PBS for 5 min, to which Hoescht was

added to a final dilution of 1:1000 for a 10 min incubation, followed by a 5 min wash in 1×

PBS. Samples were mounted in a 100% glucose solution. Images were acquired using a

Leica SP5 laser scanning confocal microscope; we dissected five brains per sex.

IPC calcium level measurements

To isolate the IPCs, individual flies were removed from food and anesthetized briefly with

CO_2. Brains were dissected in cold 1× PBS. Samples were fixed in 4% paraformaldehyde

(Electron Microscopy Science 15710) for 40 min, followed by two washes in cold 1× PBS.

Samples were incubated with Hoechst (Sigma-Aldrich, 33342) at a concentration of 1:1000

for 30 min and mounted in 75% (w/v) sucrose solution. Images were captured using a

LSI3-Leica SP5 laser scanning confocal microscope and processed using Fiji (ImageJ;[192]).

To quantify IPC activity we used IPC-specific GAL4 driver *dilp2-GAL4*[123] to drive

expression of both *UAS-mRFP* and *UAS-CaLexA* (*UAS-LexA-VP16-NFAT*)[95]. GFP

intensity in each IPC cluster was quantified by measuring the sum of pixel intensity within

the region of interest for both GFP and RFP. GFP was normalized to RFP that was co-

expressed in the IPC to account for inter-individual and/or sex differences in GAL4-driven

transgene expression (ImageJ;[209]).

IPC electrophysiology

Loose, whole-cell cell-attached patch clamp recordings were performed *in vivo* as

previously described[214,215]. Adult flies 4-8 days old were food deprived on 1% agar for 24

hr. Flies were either kept on agar (starved), moved to 500 mM D-glucose for 30 min, or

moved to whole food for 2 hr prior to mounting. Flies were briefly anesthetized with CO_2

and mounted by the cervix into a custom chamber. Nail polish was applied around the

eyes and over the labellum, adhering the mouthparts to the chamber to prevent excessive

movement. After recovering in humidity chamber for 45 min, the cuticle and air sacs

covering the *pars intercerebralis* were removed and the head and exposed brain were

immersed in Adult Hemolymph-Like (AHL) solution (108 mM NaCl, 5 mM KCl, 4 mM

$NaHCO_3$, 1 mM NaH_2PO_4, 5 mM HEPES, 15 mM ribose, 2 mM Ca^{2+}, 8.2 mM Mg^{2+}, pH

7.5). IPCs were identified by expression of *UAS-mCD8::GFP* and glass electrodes (1.5-3.5

MΩ) filled with AHL were used for all recordings, with patch resistances ranging from 50-

500 MΩ. Spikes were recorded in voltage-clamp mode in Multiclamp 700B at 20 kHz.

Recordings were passed through a low-pass filter at 2kHz and high-pass filter of 5 Hz, and

then band-pass filtered between 2 and 2000 Hz with a Bessel (8-pole) filter. Event

detection was performed in Clampfit for spike analysis, with a threshold set between -6 to -

12 pA. The number of spikes in a 30-60 second window for a given cell were counted and

converted to spikes/sec. Data from 6-8 flies were collected per group, with recordings from

1-3 IPCs per animal.

Insulin sensitivity test

To measure insulin sensitivity, we adapted a published protocol[79]. Briefly, abdominal

carcasses were isolated from 5-day-old adult flies; one biological replicate consisted of 15

abdominal carcasses. The carcasses were placed in a dish with ice-cold Schneider's

Drosophila solution (21720024, Thermo Scientific) for 30 min. After this initial incubation,

Schneider's media was replaced with 2 mL of fresh Schneider's *Drosophila* solution or

fresh Schneider's *Drosophila* solution with recombinant human insulin (final concentration

1 µM; I2643, Sigma-Aldrich) and incubated at room temperature for 30 min. After

stimulation, carcasses were collected in a tube, snap-frozen, and used for RNA isolation

and qPCR analysis.

Lifespan

After eclosion, unmated male and female flies were transferred to new vials of fresh food

every 2–3 days until no living flies remained in the vial. Deaths were recorded when the

flies were transferred to fresh food.

Female fecundity

One 5-day-old virgin female was placed with a group of three 5-day-old virgin *CS* males for

1 hr. After a 1 hr mating period, the males were removed and the females were allowed to

lay eggs. Females were transferred into new vials for egg-laying every 24 hr; the total

number of pupae in each vial was counted several days later.

Male fertility

One singly-housed 5-day-old virgin male was placed with a group of three 5-day-old virgin

CS females and allowed to interact for 1 hr. After the 1 hr mating period, the male was

removed from the vial and the females were allowed to lay eggs. Females were

transferred into new vials for egg-laying every 24 hr; the total number of pupae in each vial

was counted several days later.

Quantification and statistical analysis

Statistical analyses and data presentation were completed using GraphPad Prism 10

(GraphPad Software, San Diego, CA, USA) (Supplemental Table S2). All data were tested

for normality using the Shapiro-Wilk test. Normally-distributed data was subjected to

parametric tests, including Student's *t*-test, one-way ANOVA followed by Tukey's HSD,

and two-way ANOVA followed by Bonferroni post-hoc test as appropriate. For non-

normally-distributed data we used Mann-Whitney test and Kruskal-Wallis test followed by

Dunn's test. For two-way ANOVA involving data that do not satisfy the normality

assumption, aligned rank transformation was first applied using the art() function from the

ARTool R package[216]. Then, ANOVA was performed on the transformed data with the

base R anova() function. Finally, the art.con() function from the ARTool package was used

to extract the main as well as the interaction effects. Default parameters were used in each

step of the analysis. For all statistical analyses, differences were considered significant if

$p < 0.05$.

    Supplemental Figure Legends

    Figure S1. Dietary sugar levels change body fat but not food consumption in adult

    flies. Related to Figure 1.

(A) Food consumption per unit of body weight was significantly higher in 5-day-old adult

w^{1118} females than males (p=0.0483; Student's t-test); n=19 biological replicates.

(B) Whole-body triglyceride levels were significantly lower in 5-day-old CS females

maintained on the 0S diet compared with females kept on the 1S diet (p=0.0181); no

significant change in body fat was observed in males (p=0.0582; sex:diet interaction

p=0.7239). Two-way ANOVA followed by Bonferroni post-hoc test; n=8 biological

replicates.

(C) Food consumption was not significantly affected by diet in either 5-day-old adult w^{1118}

females or males (females: p>0.9999; males: p>0.9999; sex:diet interaction p=0.8450).

Two-way ANOVA followed by Bonferroni post-hoc test; n=10-19 biological replicates.

Note: food consumption data in 1S diet was replotted from panel S1A as food intake data

on 1S and 0S were collected in the same series of experiments.

(D) Whole-body triglyceride levels were significantly lower in 5-day-old w^{1118} adult females

and males maintained on 0S with L-glucose diet compared with sex-matched flies kept on

the 1S diet (females: p<0.0001; males: p<0.0001); the magnitude of the decrease in body

fat was greater in females than in males (sex:diet interaction p=0.0029). Two-way ANOVA

followed by Bonferroni post-hoc test; n=8 biological replicates. Note: the 1S triglyceride

data was replotted from Figure 1E as they were part of the same large experiment.

All data plotted as mean ± SEM. ns indicates not significant with p>0.05; * p<0.05, ****

p<0.0001.

1003

1004   **Figure S2. Sugar-induced changes in lipid species in the adult fat body. Related to**

1005   **Figure 2.**

(A) Specific fatty acids are enriched within triglyceride species that show a sex bias in abundance in 5-day-old adult w^{1118} female (purple) and male (green) flies maintained on 1S (p<0.01; Student's t-test). For example, all triglycerides carrying fatty acid C28:2 showed a female bias in abundance.

(B) Citrate levels in the fat body of 5-day-old adult w^{1118} males showed a trend toward lower levels of this metabolite compared with females (p=0.0630, Student's t-test); n=6-7 biological replicates. Data plotted as mean ± SEM.

(C) Number of lipid species within each subclass that show higher abundance in either 1S (purple) or 0S (pink) in 5-day-old adult w^{1118} females (p<0.01; Student's t-test). For example, 17/21 TG-O species are more abundant in 0S than in 1S, whereas 25/27 PE species are more abundant in 1S than 0S.

(D) Specific fatty acids are enriched within triglyceride species that show sugar-dependent regulation in 5-day-old adult w^{1118} females (p<0.01; Student's t-test); 1S (purple), 0S (pink). For example, 5/5 triglycerides that contain C28:2 are higher in the 1S diet, whereas 22/29 C14:0 are higher in 0S.

(E) Number of lipid species within each subclass that show higher abundance in either 1S (green) or 0S (blue) in 5-day-old adult w^{1118} males (p<0.01; Student's t-test). For example, 4/6 TG species are more abundant in 1S than in 0S.

(F) Citrate levels in the fat body were significantly lower in 5-day-old w^{1118} adult females and males maintained on 0S diet compared with sex-matched flies kept on the 1S diet (females: p=0.0193; males: p=0.0409; sex:diet interaction p=0.8806). Two-way ANOVA followed by Bonferroni post-hoc test; n=6-8 biological replicates. Data plotted as mean ± SEM. Note: the 1S citrate data was replotted from Figure S2B as they were part of the same large experiment.

1031   TG: triglyceride; TG-O: ether-linked TG; DG: diglyceride; PC: phosphatidylcholine; PC-O:

1032   ether-linked PC; PE: phosphatidylethanolamine; PG: phosphatidylglycerol; PI:

1033   phosphatidylinositol;. Acyl-Car: acyl-carnitine; Cer: Ceramide; ST: sterol; PS:

1034   phosphatidylserine; PE-O or PE-P: ether-linked PE; NAE: N-acetylethanolamine; SE:

1035   sterol ester; Hex-Cer: hexosyl-ceramide.

1036

Figure S3. Sex-specific regulation of *dilp3* mRNA and protein. Related to Figure 3.

(A-D) Representative images of maximum Z-projections from brains of 5-day-old w^{1118} flies

labelled with anti-Dilp3. Females maintained on 1S (A) and 0S (B); males maintained on

1S (C) and 0S (D). Scale bar represents 10 µm.

(E) Quantification of anti-Dilp3 levels in brains 5-day-old w^{1118} males and females

maintained on either 1S or 0S.

(F, G) Representative images of maximum Z-projections from brains of 5-day-old *dilp2-*

GAL4>UAS-CaLexA,UAS-mRFP females (F) and males (G) maintained on 1S showing

GFP levels as a readout for IPC Ca^{2+} activity. Scale bar represents 50 µm.

(H) Quantification of GFP levels in brains of 5-day-old *dilp2-GAL4>UAS-CaLexA,UAS-*

mRFP males and females maintained on 1S. Quantification of GFP levels due to IPC-

specific expression of *UAS-CaLexA* in the IPCs, normalized to RFP levels. Normalized

GFP intensity was not significantly different between females and males in 1S diet

(p=0.4782; Mann-Whitney test); n=15-23 animals.

(I) Changing the sexual identity of the IPC by expressing Tra^F (*dilp2-GAL4>UAS-tra^F*) did

not significantly alter mRNA levels of IPC-derived *dilp3* in head (p=0.7859 and p=0.0460,

respectively). One-way ANOVA followed by Tukey HSD; n=4 biological replicates.

(J) Changing the sexual identity of the APC by expressing Tra^F (*Akh-GAL4>UAS-tra^F*) did

not significantly alter mRNA levels of IPC-derived *dilp3* in heads (p=0.7143 and p=0.0009,

respectively). One-way ANOVA followed by Tukey HSD; n=6 biological replicates.

(K) Changing the sexual identity of neurons and neuropeptide-expressing cells using Tra^F

(*elav-GAL4>UAS-tra*^{F}) significantly increased mRNA levels of IPC-derived *dilp3* in heads

($p<0.0001$ and $p<0.0001$, respectively). One-way ANOVA followed by Tukey HSD; n=6

biological replicates.

Data plotted as mean ± SEM. ns indicates not significant with $p>0.05$, * $p<0.05$, ** $p<0.01$,

*** $p<0.001$, **** $p<0.0001$.

1063

Figure S4. Validating IPC ablation using *dilp2-GAL4*. Related to Figure 4.

(A, B) Representative images of maximum Z-projections in 5-day-old flies where *dilp2-*

GAL4 was used to drive expression of membrane-bound GFP (*UAS-mCD8::GFP*) to

visualize IPC morphology; female (A) and male (B) flies are shown. Scale bar represents

50 µm.

(C, D) Representative images of maximum Z-projections in 5-day-old flies where *dilp2-*

GAL4 was used to drive expression of nuclear RFP (*UAS-nRFP*) to assess colocalization

with anti-Dilp3; female (C, C□, C□□) and male (D, D□, D□□) flies. Scale bar represents

100 µm.

(E, F) mRNA levels of IPC-derived *dilp2* in heads were significantly lower in female (E;

$p=0.0001$, Student's *t*-test) and showed a trend of lower *dilp2* in male (F; $p=0.06$, Mann-

Whitney test) *dilp2-GAL4>UAS-rpr; tub-GAL80*^{ts} flies with heat shock at 29°C compared to

sex-matched flies reared in room temperature; n=4-6 biological replicates.

(G-L) Representative images of maximum Z-projections in 5-day-old flies with (I-L) and

without (G, H) *dilp2-GAL4-*mediated IPC ablation at the adult stage via heat shock-induced

rpr overexpression. Brains were labelled with anti-Dilp3 to assess IPC ablation. As

expected, IPC are present in animals without heat shock; IPC are fully (I, J) or partially (K,

L) absent in animals with heat shock. Scale bar represents 100 µm.

(M) Sex difference in body fat represented as a percent of the male-female difference in

the control *dilp2-GAL4>+,tub-GAL80^{ts}* genotype.

(N) Sex difference in body fat represented as a percent of the male-female difference in

the *dilp2-GAL4>rpr,tub-GAL80^{ts}* genotype without heat shock.

(O) *dilp2-GAL4>+ and +>UAS-dilp3* females maintained on 0S had significantly lower body

fat than genotype-matched females kept on 1S, a sugar-dependent reduction in body fat

that was blocked in females with IPC-specific overexpression of Dilp3 (*dilp2-GAL4>UAS-*

dilp3; 1S to 0S $p<0.0001$ [GAL4], $p<0.0001$ [UAS] and $p=0.5298$ [Dilp3 overexpression]).

Two-way ANOVA followed by Bonferroni post-hoc test; n=8 biological replicates.

(P) *dilp2-GAL4>+ and +>UAS-dilp3* males maintained on 0S had significantly lower body

fat than genotype-matched females kept on 1S, a sugar-dependent reduction in body fat

that was blocked in males with IPC-specific overexpression of Dilp3 (*dilp2-GAL4>UAS-*

dilp3; $p=0.0029$, $p=0.0452$ and $p=0.1426$, respectively). Two-way ANOVA followed by

Bonferroni post-hoc test; n=8 biological replicates.

(Q) Whole-body triglyceride levels were significantly lower in 5-day-old adult w^{1118} females

maintained on a diet supplemented with 0.5 mg/mL 20-HE compared with control females

($p=0.0067$, Student's t-test); n=8 biological replicates.

Data plotted as mean ± SEM. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$.

1100

References:

1. Kühnlein, R.P. (2012). Lipid droplet-based storage fat metabolism in Drosophila:

Thematic Review Series: Lipid Droplet Synthesis and Metabolism: from Yeast to Man. J.

Lipid Res. *53*, 1430–1436. https://doi.org/10.1194/jlr.R024299.

2. Kühnlein, R.P. (2011). The contribution of the *Drosophila* model to lipid droplet

research. Prog. Lipid Res. *50*, 348–356. https://doi.org/10.1016/j.plipres.2011.04.001.

3. Gutierrez, E., Wiggins, D., Fielding, B., and Gould, A.P. (2007). Specialized hepatocyte-

like cells regulate Drosophila lipid metabolism. Nature *445*, 275–280.

https://doi.org/10.1038/nature05382.

4. Sieber, M.H., and Thummel, C.S. (2009). The DHR96 Nuclear Receptor Controls

Triacylglycerol Homeostasis in Drosophila. Cell Metab. *10*, 481–490.

https://doi.org/10.1016/j.cmet.2009.10.010.

5. Wat, L.W., Chao, C., Bartlett, R., Buchanan, J.L., Millington, J.W., Chih, H.J.,

Chowdhury, Z.S., Biswas, P., Huang, V., Shin, L.J., et al. (2020). A role for triglyceride

lipase brummer in the regulation of sex differences in Drosophila fat storage and

breakdown. PLOS Biol. *18*, e3000595. https://doi.org/10.1371/journal.pbio.3000595.

6. Kis, V., Barti, B., Lippai, M., and Sass, M. (2015). Specialized Cortex Glial Cells

Accumulate Lipid Droplets in Drosophila melanogaster. PloS One *10*, e0131250.

https://doi.org/10.1371/journal.pone.0131250.

7. Villanueva, J.E., Livelo, C., Trujillo, A.S., Chandran, S., Woodworth, B., Andrade, L., Le,

H.D., Manor, U., Panda, S., and Melkani, G.C. (2019). Time-restricted feeding restores

muscle function in Drosophila models of obesity and circadian-rhythm disruption. Nat.

Commun. *10*, 2700. https://doi.org/10.1038/s41467-019-10563-9.

8. Chao, C.F., Pesch, Y.-Y., Yu, H., Wang, C., Aristizabal, M.J., Huan, T., Tanentzapf, G.,

and Rideout, E.J. (2024). An important role for triglyceride in regulating spermatogenesis.

eLife *12*. https://doi.org/10.7554/eLife.87523.3.

9. Sieber, M.H., and Spradling, A.C. (2015). Steroid Signaling Establishes a Female

Metabolic State and Regulates SREBP to Control Oocyte Lipid Accumulation. Curr. Biol.

25, 993–1004. https://doi.org/10.1016/j.cub.2015.02.019.

10. Wat, L.W., Chowdhury, Z.S., Millington, J.W., Biswas, P., and Rideout, E.J. (2021).

Sex determination gene transformer regulates the male-female difference in Drosophila fat

storage via the adipokinetic hormone pathway. eLife *10*, e72350.

https://doi.org/10.7554/eLife.72350.

11. Schwasinger-Schmidt, T.E., Kachman, S.D., and Harshman, L.G. (2012). Evolution of

starvation resistance in Drosophila melanogaster: measurement of direct and correlated

responses to artificial selection. J. Evol. Biol. *25*, 378–387. https://doi.org/10.1111/j.1420-

9101.2011.02428.x.

12. Buszczak, M., Lu, X., Segraves, W.A., Chang, T.Y., and Cooley, L. (2002). Mutations

in the midway Gene Disrupt a Drosophila Acyl Coenzyme A: Diacylglycerol

Acyltransferase. Genetics *160*, 1511–1518. https://doi.org/10.1093/genetics/160.4.1511.

13. Wilfling, F., Wang, H., Haas, J.T., Krahmer, N., Gould, T.J., Uchida, A., Cheng, J.-X.,

Graham, M., Christiano, R., Fröhlich, F., et al. (2013). Triacylglycerol synthesis enzymes

mediate lipid droplet growth by relocalizing from the ER to lipid droplets. Dev. Cell *24*,

384–399. https://doi.org/10.1016/j.devcel.2013.01.013.

14. Bailey, A.P., Koster, G., Guillermier, C., Hirst, E.M.A., MacRae, J.I., Lechene, C.P.,

Postle, A.D., and Gould, A.P. (2015). Antioxidant Role for Lipid Droplets in a Stem Cell

Niche of Drosophila. Cell *163*, 340–353. https://doi.org/10.1016/j.cell.2015.09.020.

15. Ugrankar, R., Liu, Y., Provaznik, J., Schmitt, S., and Lehmann, M. (2011). Lipin is a

central regulator of adipose tissue development and function in Drosophila melanogaster.

Mol. Cell. Biol. *31*, 1646–1656. https://doi.org/10.1128/MCB.01335-10.

16. Beller, M., Bulankina, A.V., Hsiao, H.-H., Urlaub, H., Jäckle, H., and Kühnlein, R.P.

(2010). PERILIPIN-dependent control of lipid droplet structure and fat storage in

Drosophila. Cell Metab. *12*, 521–532. https://doi.org/10.1016/j.cmet.2010.10.001.

17. Grönke, S., Mildner, A., Fellert, S., Tennagels, N., Petry, S., Müller, G., Jäckle, H., and

Kühnlein, R.P. (2005). Brummer lipase is an evolutionary conserved fat storage regulator

in Drosophila. Cell Metab. *1*, 323–330. https://doi.org/10.1016/j.cmet.2005.04.003.

18. Palm, W., Sampaio, J.L., Brankatschk, M., Carvalho, M., Mahmoud, A., Shevchenko,

A., and Eaton, S. (2012). Lipoproteins in Drosophila melanogaster--assembly, function,

and influence on tissue lipid composition. PLoS Genet. *8*, e1002828.

https://doi.org/10.1371/journal.pgen.1002828.

19. Bi, J., Xiang, Y., Chen, H., Liu, Z., Grönke, S., Kühnlein, R.P., and Huang, X. (2012).

Opposite and redundant roles of the two Drosophila perilipins in lipid mobilization. J. Cell

Sci. *125*, 3568–3577. https://doi.org/10.1242/jcs.101329.

20. Tian, Y., Bi, J., Shui, G., Liu, Z., Xiang, Y., Liu, Y., Wenk, M.R., Yang, H., and Huang,

X. (2011). Tissue-autonomous function of Drosophila seipin in preventing ectopic lipid

droplet formation. PLoS Genet. *7*, e1001364.

https://doi.org/10.1371/journal.pgen.1001364.

21. Ding, L., Yang, X., Tian, H., Liang, J., Zhang, F., Wang, G., Wang, Y., Ding, M., Shui,

G., and Huang, X. (2018). Seipin regulates lipid homeostasis by ensuring calcium-

dependent mitochondrial metabolism. EMBO J. *37*, e97572.

https://doi.org/10.15252/embj.201797572.

22. Teixeira, L., Rabouille, C., Rørth, P., Ephrussi, A., and Vanzo, N.F. (2003). Drosophila

Perilipin/ADRP homologue Lsd2 regulates lipid metabolism. Mech. Dev. *120*, 1071–1081.

https://doi.org/10.1016/s0925-4773(03)00158-8.

23. Grönke, S., Beller, M., Fellert, S., Ramakrishnan, H., Jäckle, H., and Kühnlein, R.P.

(2003). Control of fat storage by a Drosophila PAT domain protein. Curr. Biol. CB *13*, 603–

606. https://doi.org/10.1016/s0960-9822(03)00175-1.

24. Fauny, J.D., Silber, J., and Zider, A. (2005). Drosophila Lipid Storage Droplet 2 gene

(Lsd-2) is expressed and controls lipid storage in wing imaginal discs. Dev. Dyn. Off. Publ.

Am. Assoc. Anat. *232*, 725–732. https://doi.org/10.1002/dvdy.20277.

25. Pospisilik, J.A., Schramek, D., Schnidar, H., Cronin, S.J.F., Nehme, N.T., Zhang, X.,

Knauf, C., Cani, P.D., Aumayr, K., Todoric, J., et al. (2010). Drosophila Genome-wide

Obesity Screen Reveals Hedgehog as a Determinant of Brown versus White Adipose Cell

Fate. Cell *140*, 148–160. https://doi.org/10.1016/j.cell.2009.12.027.

26. Baumbach, J., Hummel, P., Bickmeyer, I., Kowalczyk, K.M., Frank, M., Knorr, K.,

Hildebrandt, A., Riedel, D., Jäckle, H., and Kühnlein, R.P. (2014). A Drosophila in vivo

screen identifies store-operated calcium entry as a key regulator of adiposity. Cell Metab.

19, 331–343. https://doi.org/10.1016/j.cmet.2013.12.004.

27. Baranski, T.J., Kraja, A.T., Fink, J.L., Feitosa, M., Lenzini, P.A., Borecki, I.B., Liu, C.-

T., Cupples, L.A., North, K.E., and Province, M.A. (2018). A high throughput, functional

screen of human Body Mass Index GWAS loci using tissue-specific RNAi Drosophila

melanogaster crosses. PLoS Genet. *14*, e1007222.

https://doi.org/10.1371/journal.pgen.1007222.

28. Kunte, A.S., Matthews, K.A., and Rawson, R.B. (2006). Fatty acid auxotrophy in

Drosophila larvae lacking SREBP. Cell Metab. *3*, 439–448.

https://doi.org/10.1016/j.cmet.2006.04.011.

29. Teleman, A.A., Chen, Y.-W., and Cohen, S.M. (2005). 4E-BP functions as a metabolic

brake used under stress conditions but not during normal growth. Genes Dev. *19*, 1844–

1848. https://doi.org/10.1101/gad.341505.

30. Rajan, A., and Perrimon, N. (2012). Drosophila cytokine unpaired 2 regulates

physiological homeostasis by remotely controlling insulin secretion. Cell *151*, 123–137.

https://doi.org/10.1016/j.cell.2012.08.019.

31. Reis, T., Gilst, M.R.V., and Hariharan, I.K. (2010). A Buoyancy-Based Screen of

Drosophila Larvae for Fat-Storage Mutants Reveals a Role for Sir2 in Coupling Fat

Storage to Nutrient Availability. PLOS Genet. *6*, e1001206.

https://doi.org/10.1371/journal.pgen.1001206.

32. Tsuda-Sakurai, K., Seong, K.-H., Horiuchi, J., Aigaki, T., and Tsuda, M. (2015).

Identification of a novel role for Drosophila MESR4 in lipid metabolism. Genes Cells *20*,

358–365. https://doi.org/10.1111/gtc.12221.

33. Bennick, R.A., Nagengast, A.A., and DiAngelo, J.R. (2019). The SR proteins SF2 and

RBP1 regulate triglyceride storage in the fat body of *Drosophila*. Biochem. Biophys. Res.

Commun. *516*, 928–933. https://doi.org/10.1016/j.bbrc.2019.06.151.

34. Kishita, Y., Tsuda, M., and Aigaki, T. (2012). Impaired fatty acid oxidation in a

Drosophila model of mitochondrial trifunctional protein (MTP) deficiency. Biochem.

Biophys. Res. Commun. *419*, 344–349. https://doi.org/10.1016/j.bbrc.2012.02.026.

35. Palanker, L., Tennessen, J.M., Lam, G., and Thummel, C.S. (2009). Drosophila HNF4

Regulates Lipid Mobilization and β-Oxidation. Cell Metab. *9*, 228–239.

https://doi.org/10.1016/j.cmet.2009.01.009.

36. Lee, S., Bao, H., Ishikawa, Z., Wang, W., and Lim, H.-Y. (2017). Cardiomyocyte

Regulation of Systemic Lipid Metabolism by the Apolipoprotein B-Containing Lipoproteins

in Drosophila. PLoS Genet. *13*, e1006555. https://doi.org/10.1371/journal.pgen.1006555.

37. Parra-Peralbo, E., and Culi, J. (2011). Drosophila lipophorin receptors mediate the

uptake of neutral lipids in oocytes and imaginal disc cells by an endocytosis-independent

mechanism. PLoS Genet. *7*, e1001297. https://doi.org/10.1371/journal.pgen.1001297.

38. Lim, H.-Y., Wang, W., Wessells, R.J., Ocorr, K., and Bodmer, R. (2011). Phospholipid

homeostasis regulates lipid metabolism and cardiac function through SREBP signaling in

Drosophila. Genes Dev. *25*, 189–200. https://doi.org/10.1101/gad.1992411.

39. Bauer, R., Voelzmann, A., Breiden, B., Schepers, U., Farwanah, H., Hahn, I., Eckardt,

F., Sandhoff, K., and Hoch, M. (2009). Schlank, a member of the ceramide synthase family

controls growth and body fat in Drosophila. EMBO J. *28*, 3706–3716.

https://doi.org/10.1038/emboj.2009.305.

40. Li, Y., Hoffmann, J., Li, Y., Stephano, F., Bruchhaus, I., Fink, C., and Roeder, T.

(2016). Octopamine controls starvation resistance, life span and metabolic traits in

Drosophila. Sci. Rep. *6*, 35359. https://doi.org/10.1038/srep35359.

41. Neckameyer, W.S., Coleman, C.M., Eadie, S., and Goodwin, S.F. (2007).

Compartmentalization of neuronal and peripheral serotonin synthesis in Drosophila

melanogaster. Genes Brain Behav. *6*, 756–769. https://doi.org/10.1111/j.1601-

183X.2007.00307.x.

42. Malita, A., Kubrak, O., Koyama, T., Ahrentløv, N., Texada, M.J., Nagy, S., Halberg,

K.V., and Rewitz, K. (2022). A gut-derived hormone suppresses sugar appetite and

regulates food choice in Drosophila. Nat. Metab. *4*, 1532–1550.

https://doi.org/10.1038/s42255-022-00672-z.

43. Yoshinari, Y., Kosakamoto, H., Kamiyama, T., Hoshino, R., Matsuoka, R., Kondo, S.,

Tanimoto, H., Nakamura, A., Obata, F., and Niwa, R. (2021). The sugar-responsive

enteroendocrine neuropeptide F regulates lipid metabolism through glucagon-like and

insulin-like hormones in Drosophila melanogaster. Nat. Commun. *12*, 4818.

https://doi.org/10.1038/s41467-021-25146-w.

44. Scopelliti, A., Bauer, C., Yu, Y., Zhang, T., Kruspig, B., Murphy, D.J., Vidal, M.,

Maddocks, O.D.K., and Cordero, J.B. (2019). A Neuronal Relay Mediates a Nutrient

Responsive Gut/Fat Body Axis Regulating Energy Homeostasis in Adult Drosophila. Cell

Metab. *29*, 269-284.e10. https://doi.org/10.1016/j.cmet.2018.09.021.

45. Hentze, J.L., Carlsson, M.A., Kondo, S., Nässel, D.R., and Rewitz, K.F. (2015). The

Neuropeptide Allatostatin A Regulates Metabolism and Feeding Decisions in Drosophila.

Sci. Rep. *5*, 11680. https://doi.org/10.1038/srep11680.

46. Kubrak, O., Koyama, T., Ahrentløv, N., Jensen, L., Malita, A., Naseem, M.T., Lassen,

M., Nagy, S., Texada, M.J., Halberg, K.V., et al. (2022). The gut hormone Allatostatin

C/Somatostatin regulates food intake and metabolic homeostasis under nutrient stress.

Nat. Commun. *13*, 692. https://doi.org/10.1038/s41467-022-28268-x.

47. Beshel, J., Dubnau, J., and Zhong, Y. (2017). A Leptin analog locally produced in the

brain acts via a conserved neural circuit to modulate obesity-linked behaviors in

Drosophila. Cell Metab. *25*, 208–217. https://doi.org/10.1016/j.cmet.2016.12.013.

48. Yamamoto, R., Bai, H., Dolezal, A.G., Amdam, G., and Tatar, M. (2013). Juvenile

hormone regulation of Drosophila aging. BMC Biol. *11*, 85. https://doi.org/10.1186/1741-

7007-11-85.

49. Kamoshida, Y., Fujiyama-Nakamura, S., Kimura, S., Suzuki, E., Lim, J., Shiozaki-Sato,

Y., Kato, S., and Takeyama, K.-I. (2012). Ecdysone receptor (EcR) suppresses lipid

accumulation in the Drosophila fat body via transcription control. Biochem. Biophys. Res.

Commun. *421*, 203–207. https://doi.org/10.1016/j.bbrc.2012.03.135.

50. Song, W., Veenstra, J.A., and Perrimon, N. (2014). Control of lipid metabolism by

tachykinin in Drosophila. Cell Rep. *9*, 40–47. https://doi.org/10.1016/j.celrep.2014.08.060.

51. Gera, J., Agard, M., Nave, H., Sajadi, F., Thorat, L., Kondo, S., Nässel, D.R., Paluzzi,

J.-P.V., and Zandawala, M. (2024). Anti-diuretic hormone ITP signals via a guanylate

cyclase receptor to modulate systemic homeostasis in Drosophila. eLife *13*.

https://doi.org/10.7554/eLife.97043.1.

52. Gáliková, M., and Klepsatel, P. (2022). Ion transport peptide regulates energy intake,

expenditure, and metabolic homeostasis in Drosophila. Genetics *222*, iyac150.

https://doi.org/10.1093/genetics/iyac150.

53. DiAngelo, J.R., and Birnbaum, M.J. (2009). Regulation of Fat Cell Mass by Insulin in

Drosophila melanogaster. Mol. Cell. Biol. *29*, 6341–6352.

https://doi.org/10.1128/MCB.00675-09.

54. Grönke, S., Clarke, D.-F., Broughton, S., Andrews, T.D., and Partridge, L. (2010).

Molecular evolution and functional characterization of Drosophila insulin-like peptides.

PLoS Genet. *6*, e1000857. https://doi.org/10.1371/journal.pgen.1000857.

55. Grönke, S., Müller, G., Hirsch, J., Fellert, S., Andreou, A., Haase, T., Jäckle, H., and

Kühnlein, R.P. (2007). Dual lipolytic control of body fat storage and mobilization in

Drosophila. PLoS Biol. *5*, e137. https://doi.org/10.1371/journal.pbio.0050137.

56. Bharucha, K.N., Tarr, P., and Zipursky, S.L. (2008). A glucagon-like endocrine

pathway in Drosophila modulates both lipid and carbohydrate homeostasis. J. Exp. Biol.

211, 3103–3110. https://doi.org/10.1242/jeb.016451.

57. Gáliková, M., Diesner, M., Klepsatel, P., Hehlert, P., Xu, Y., Bickmeyer, I., Predel, R.,

and Kühnlein, R.P. (2015). Energy Homeostasis Control in Drosophila Adipokinetic

Hormone Mutants. Genetics *201*, 665–683. https://doi.org/10.1534/genetics.115.178897.

58. Zhao, X., and Karpac, J. (2017). Muscle Directs Diurnal Energy Homeostasis through

a Myokine-Dependent Hormone Module in Drosophila. Curr. Biol. CB *27*, 1941-1955.e6.

https://doi.org/10.1016/j.cub.2017.06.004.

59. Kapan, N., Lushchak, O.V., Luo, J., and Nässel, D.R. (2012). Identified peptidergic

neurons in the Drosophila brain regulate insulin-producing cells, stress responses and

metabolism by coexpressed short neuropeptide F and corazonin. Cell. Mol. Life Sci. CMLS

69, 4051–4066. https://doi.org/10.1007/s00018-012-1097-z.

60. Luo, J., Becnel, J., Nichols, C.D., and Nässel, D.R. (2012). Insulin-producing cells in

the brain of adult Drosophila are regulated by the serotonin 5-HT1A receptor. Cell. Mol.

Life Sci. CMLS *69*, 471–484. https://doi.org/10.1007/s00018-011-0789-0.

61. Zandawala, M., Yurgel, M.E., Texada, M.J., Liao, S., Rewitz, K.F., Keene, A.C., and

Nässel, D.R. (2018). Modulation of Drosophila post-feeding physiology and behavior by

the neuropeptide leucokinin. PLoS Genet. *14*, e1007767.

https://doi.org/10.1371/journal.pgen.1007767.

62. Kubrak, O., Jørgensen, A.F., Koyama, T., Lassen, M., Nagy, S., Hald, J., Mazzoni, G.,

Madsen, D., Hansen, J.B., Larsen, M.R., et al. (2024). LGR signaling mediates muscle-

adipose tissue crosstalk and protects against diet-induced insulin resistance. Nat.

Commun. *15*, 6126. https://doi.org/10.1038/s41467-024-50468-w.

63. Heier, C., and Kühnlein, R.P. (2018). Triacylglycerol Metabolism in Drosophila

melanogaster. Genetics *210*, 1163–1184. https://doi.org/10.1534/genetics.118.301583.

64. Lee, G., and Park, J.H. (2004). Hemolymph sugar homeostasis and starvation-induced

hyperactivity affected by genetic manipulations of the adipokinetic hormone-encoding gene

in Drosophila melanogaster. Genetics *167*, 311–323.

https://doi.org/10.1534/genetics.167.1.311.

65. Noyes, B.E., Katz, F.N., and Schaffer, M.H. (1995). Identification and expression of the

Drosophila adipokinetic hormone gene. Mol. Cell. Endocrinol. *109*, 133–141.

https://doi.org/10.1016/0303-7207(95)03492-p.

66. Mochanová, M., Tomčala, A., Svobodová, Z., and Kodrík, D. (2018). Role of

adipokinetic hormone during starvation in Drosophila. Comp. Biochem. Physiol. B

Biochem. Mol. Biol. *226*, 26–35. https://doi.org/10.1016/j.cbpb.2018.08.004.

67. He, Q., Du, J., Wei, L., and Zhao, Z. (2020). AKH-FOXO pathway regulates starvation-

induced sleep loss through remodeling of the small ventral lateral neuron dorsal

projections. PLoS Genet. *16*, e1009181. https://doi.org/10.1371/journal.pgen.1009181.

68. Li, H., Janssens, J., De Waegeneer, M., Kolluru, S.S., Davie, K., Gardeux, V., Saelens,

W., David, F., Brbić, M., Spanier, K., et al. (2022). Fly Cell Atlas: a single-nucleus

transcriptomic atlas of the adult fruit fly. Science *375*, eabk2432.

https://doi.org/10.1126/science.abk2432.

69. Leader, D.P., Krause, S.A., Pandit, A., Davies, S.A., and Dow, J.A.T. (2018). FlyAtlas

2: a new version of the Drosophila melanogaster expression atlas with RNA-Seq, miRNA-

Seq and sex-specific data. Nucleic Acids Res. *46*, D809–D815.

https://doi.org/10.1093/nar/gkx976.

70. De Groef, S., Ribeiro Lopes, M., Winant, M., Rosschaert, E., Wilms, T., Bolckmans, L.,

Calevro, F., and Callaerts, P. (2024). Reference genes to study the sex-biased expression

of genes regulating Drosophila metabolism. Sci. Rep. *14*, 9518.

https://doi.org/10.1038/s41598-024-58863-5.

71. Ja, W.W., Carvalho, G.B., Mak, E.M., de la Rosa, N.N., Fang, A.Y., Liong, J.C.,

Brummel, T., and Benzer, S. (2007). Prandiology of Drosophila and the CAFE assay. Proc.

Natl. Acad. Sci. U. S. A. *104*, 8253–8256. https://doi.org/10.1073/pnas.0702726104.

72. Al-Anzi, B., Sapin, V., Waters, C., Zinn, K., Wyman, R.J., and Benzer, S. (2009).

Obesity-Blocking Neurons in Drosophila. Neuron *63*, 329–341.

https://doi.org/10.1016/j.neuron.2009.07.021.

73. Tennessen, J.M., Barry, W.E., Cox, J., and Thummel, C.S. (2014). Methods for

studying metabolism in Drosophila. Methods San Diego Calif *68*, 105–115.

https://doi.org/10.1016/j.ymeth.2014.02.034.

74. Hildebrandt, A., Bickmeyer, I., and Kühnlein, R.P. (2011). Reliable Drosophila body fat

quantification by a coupled colorimetric assay. PloS One *6*, e23796.

https://doi.org/10.1371/journal.pone.0023796.

75. De Groef, S., Wilms, T., Balmand, S., Calevro, F., and Callaerts, P. (2021). Sexual

Dimorphism in Metabolic Responses to Western Diet in Drosophila melanogaster.

Biomolecules *12*, 33. https://doi.org/10.3390/biom12010033.

76. Musselman, L.P., Fink, J.L., and Baranski, T.J. (2019). Similar effects of high-fructose

and high-glucose feeding in a Drosophila model of obesity and diabetes. PLoS ONE *14*,

e0217096. https://doi.org/10.1371/journal.pone.0217096.

77. Rovenko, B.M., Perkhulyn, N.V., Gospodaryov, D.V., Sanz, A., Lushchak, O.V., and

Lushchak, V.I. (2015). High consumption of fructose rather than glucose promotes a diet-

induced obese phenotype in *Drosophila melanogaster*. Comp. Biochem. Physiol. A. Mol.

Integr. Physiol. *180*, 75–85. https://doi.org/10.1016/j.cbpa.2014.11.008.

78. Dam, E. van, Leeuwen, L.A.G. van, Santos, E. dos, James, J., Best, L., Lennicke, C.,

Vincent, A.J., Marinos, G., Foley, A., Buricova, M., et al. (2020). Sugar-Induced Obesity

and Insulin Resistance Are Uncoupled from Shortened Survival in Drosophila. Cell Metab.

31, 710-725.e7. https://doi.org/10.1016/j.cmet.2020.02.016.

79. Musselman, L.P., Fink, J.L., Narzinski, K., Ramachandran, P.V., Sukumar

Hathiramani, S., Cagan, R.L., and Baranski, T.J. (2011). A high-sugar diet produces

obesity and insulin resistance in wild-type Drosophila. Dis. Model. Mech. *4*, 842–849.

https://doi.org/10.1242/dmm.007948.

80. Millington, J.W., Biswas, P., Chao, C., Xia, Y.H., Wat, L.W., Brownrigg, G.P., Sun, Z.,

Basner-Collins, P.J., Klein Geltink, R.I., and Rideout, E.J. (2022). A low-sugar diet

enhances Drosophila body size in males and females via sex-specific mechanisms. Dev.

Camb. Engl. *149*, dev200491. https://doi.org/10.1242/dev.200491.

81. Lewis, E.B. (1960). A new standard food medium. Drosoph. Inf. Serv. *34*, 1–55.

82. Nayak, N., and Mishra, M. (2021). High fat diet induced abnormalities in metabolism,

growth, behavior, and circadian clock in *Drosophila melanogaster*. Life Sci. *281*, 119758.

https://doi.org/10.1016/j.lfs.2021.119758.

83. Birse, R.T., Choi, J., Reardon, K., Rodriguez, J., Graham, S., Diop, S., Ocorr, K.,

Bodmer, R., and Oldham, S. (2010). High-fat-diet-induced obesity and heart dysfunction

are regulated by the TOR pathway in Drosophila. Cell Metab. *12*, 533–544.

https://doi.org/10.1016/j.cmet.2010.09.014.

84. Heinrichsen, E.T., and Haddad, G.G. (2012). Role of High-Fat Diet in Stress Response

of Drosophila. PLOS ONE *7*, e42587. https://doi.org/10.1371/journal.pone.0042587.

85. Woodcock, K.J., Kierdorf, K., Pouchelon, C.A., Vivancos, V., Dionne, M.S., and

Geissmann, F. (2015). Macrophage-Derived upd3 Cytokine Causes Impaired Glucose

Homeostasis and Reduced Lifespan in Drosophila Fed a Lipid-Rich Diet. Immunity *42*,

133–144. https://doi.org/10.1016/j.immuni.2014.12.023.

86. Liao, S., Amcoff, M., and Nässel, D.R. (2021). Impact of high-fat diet on lifespan,

metabolism, fecundity and behavioral senescence in *Drosophila*. Insect Biochem. Mol.

Biol. *133*, 103495. https://doi.org/10.1016/j.ibmb.2020.103495.

87. Stafford, J.W., Lynd, K.M., Jung, A.Y., and Gordon, M.D. (2012). Integration of Taste

and Calorie Sensing in Drosophila. J. Neurosci. *32*, 14767–14774.

https://doi.org/10.1523/JNEUROSCI.1887-12.2012.

88. Regan, J.C., Khericha, M., Dobson, A.J., Bolukbasi, E., Rattanavirotkul, N., and

Partridge, L. (2016). Sex difference in pathology of the ageing gut mediates the greater

response of female lifespan to dietary restriction. eLife *5*, e10956.

https://doi.org/10.7554/eLife.10956.

89. Mirth, C.K., and Riddiford, L.M. (2007). Size assessment and growth control: how adult

size is determined in insects. BioEssays News Rev. Mol. Cell. Dev. Biol. *29*, 344–355.

https://doi.org/10.1002/bies.20552.

90. Kréneisz, O., Chen, X., Fridell, Y.-W.C., and Mulkey, D.K. (2010). Glucose increases

activity and Ca2+ in insulin-producing cells of adult Drosophila. Neuroreport *21*, 1116–

1120. https://doi.org/10.1097/WNR.0b013e3283409200.

91. Nässel, D.R., Liu, Y., and Luo, J. (2015). Insulin/IGF signaling and its regulation in

Drosophila. Gen. Comp. Endocrinol. *221*, 255–266.

https://doi.org/10.1016/j.ygcen.2014.11.021.

92. Park, S., Alfa, R.W., Topper, S.M., Kim, G.E.S., Kockel, L., and Kim, S.K. (2014). A

Genetic Strategy to Measure Circulating Drosophila Insulin Reveals Genes Regulating

Insulin Production and Secretion. PLOS Genet. *10*, e1004555.

https://doi.org/10.1371/journal.pgen.1004555.

93. Oh, Y., Lai, J.S.-Y., Mills, H.J., Erdjument-Bromage, H., Giammarinaro, B., Saadipour,

K., Wang, J.G., Abu, F., Neubert, T.A., and Suh, G.S.B. (2019). A glucose-sensing neuron

pair regulates insulin and glucagon in Drosophila. Nature *574*, 559–564.

https://doi.org/10.1038/s41586-019-1675-4.

94. Buhler, K., Clements, J., Winant, M., Bolckmans, L., Vulsteke, V., and Callaerts, P.

(2018). Growth control through regulation of insulin signalling by nutrition-activated steroid

hormone in Drosophila. Development *145*, dev165654.

https://doi.org/10.1242/dev.165654.

95. Masuyama, K., Zhang, Y., Rao, Y., and Wang, J.W. (2012). Mapping Neural Circuits

with Activity-Dependent Nuclear Import of a Transcription Factor. J. Neurogenet. *26*, 89–

102. https://doi.org/10.3109/01677063.2011.642910.

96. Bisen, R.S., Iqbal, F.M., Cascino-Milani, F., Bockemühl, T., and Ache, J.M. (2024).

Nutritional state-dependent modulation of Insulin-Producing Cells in Drosophila. eLife *13*.

https://doi.org/10.7554/eLife.98514.1.

97. Sturtevant, A.H. (1945). A Gene in Drosophila Melanogaster That Transforms Females

into Males. Genetics *30*, 297–299. https://doi.org/10.1093/genetics/30.3.297.

98. McKeown, M., Belote, J.M., and Baker, B.S. (1987). A molecular analysis of

transformer, a gene in drosophila melanogaster that controls female sexual differentiation.

Cell *48*, 489–499. https://doi.org/10.1016/0092-8674(87)90199-1.

99. Camara, N., Whitworth, C., and Van Doren, M. (2008). The creation of sexual

dimorphism in the Drosophila soma. Curr. Top. Dev. Biol. *83*, 65–107.

https://doi.org/10.1016/S0070-2153(08)00403-1.

100. Oliver, B. (2002). Genetic control of germline sexual dimorphism in *Drosophila*. In

International Review of Cytology, K. W. Jeon, ed. (Academic Press), pp. 1–60.

https://doi.org/10.1016/S0074-7696(02)19010-3.

101. Burtis, K.C. (1993). The regulation of sex determination and sexually dimorphic

differentiation in Drosophila. Curr. Opin. Cell Biol. *5*, 1006–1014.

https://doi.org/10.1016/0955-0674(93)90085-5.

102. Rideout, E.J., Dornan, A.J., Neville, M.C., Eadie, S., and Goodwin, S.F. (2010).

Control of sexual differentiation and behavior by the doublesex gene in Drosophila

melanogaster. Nat. Neurosci. *13*, 458–466. https://doi.org/10.1038/nn.2515.

103. Rideout, E.J., Narsaiya, M.S., and Grewal, S.S. (2015). The Sex Determination Gene

transformer Regulates Male-Female Differences in Drosophila Body Size. PLOS Genet.

11, e1005683. https://doi.org/10.1371/journal.pgen.1005683.

104. Hudry, B., Khadayate, S., and Miguel-Aliaga, I. (2016). The sexual identity of adult

intestinal stem cells controls organ size and plasticity. Nature *530*, 344–348.

https://doi.org/10.1038/nature16953.

105. Hudry, B., de Goeij, E., Mineo, A., Gaspar, P., Hadjieconomou, D., Studd, C.,

Mokochinski, J.B., Kramer, H.B., Plaçais, P.-Y., Preat, T., et al. (2019). Sex Differences in

Intestinal Carbohydrate Metabolism Promote Food Intake and Sperm Maturation. Cell *178*,

901-918.e16. https://doi.org/10.1016/j.cell.2019.07.029.

106. Evans, D.S., and Cline, T.W. (2007). Drosophila melanogaster Male Somatic Cells

Feminized Solely by TraF Can Collaborate With Female Germ Cells to Make Functional

Eggs. Genetics *175*, 631–642. https://doi.org/10.1534/genetics.106.066332.

107. Regan, J.C., Lu, Y.-X., Ureña, E., Meilenbrock, R.L., Catterson, J.H., Kißler, D.,

Fröhlich, J., Funk, E., and Partridge, L. (2022). Sexual identity of enterocytes regulates

autophagy to determine intestinal health, lifespan and responses to rapamycin. Nat. Aging

2, 1145–1158. https://doi.org/10.1038/s43587-022-00308-7.

108. Kim, J., and Neufeld, T.P. (2015). Dietary sugar promotes systemic TOR activation in

Drosophila through AKH-dependent selective secretion of Dilp3. Nat. Commun. *6*, 6846.

https://doi.org/10.1038/ncomms7846.

109. Birse, R.T., Söderberg, J.A.E., Luo, J., Winther, Å.M.E., and Nässel, D.R. (2011).

Regulation of insulin-producing cells in the adult Drosophila brain via the tachykinin

peptide receptor DTKR. J. Exp. Biol. *214*, 4201–4208. https://doi.org/10.1242/jeb.062091.

110. Brogiolo, W., Stocker, H., Ikeya, T., Rintelen, F., Fernandez, R., and Hafen, E.

(2001). An evolutionarily conserved function of the *Drosophila* insulin receptor and insulin-

like peptides in growth control. Curr. Biol. *11*, 213–221. https://doi.org/10.1016/S0960-

9822(01)00068-9.

111. Ikeya, T., Galic, M., Belawat, P., Nairz, K., and Hafen, E. (2002). Nutrient-dependent

expression of insulin-like peptides from neuroendocrine cells in the CNS contributes to

growth regulation in Drosophila. Curr. Biol. CB *12*, 1293–1300.

https://doi.org/10.1016/s0960-9822(02)01043-6.

112. Alic, N., Andrews, T.D., Giannakou, M.E., Papatheodorou, I., Slack, C., Hoddinott,

M.P., Cochemé, H.M., Schuster, E.F., Thornton, J.M., and Partridge, L. (2011).

Genome-wide dFOXO targets and topology of the transcriptomic response to stress and

insulin signalling. Mol. Syst. Biol. *7*, 502. https://doi.org/10.1038/msb.2011.36.

113. Puig, O., and Tjian, R. (2005). Transcriptional feedback control of insulin receptor by

dFOXO/FOXO1. Genes Dev. *19*, 2435–2446. https://doi.org/10.1101/gad.1340505.

114. Puig, O., Marr, M.T., Ruhf, M.L., and Tjian, R. (2003). Control of cell number by

Drosophila FOXO: downstream and feedback regulation of the insulin receptor pathway.

Genes Dev. *17*, 2006–2020. https://doi.org/10.1101/gad.1098703.

115. Zinke, I., Schütz, C.S., Katzenberger, J.D., Bauer, M., and Pankratz, M.J. (2002).

Nutrient control of gene expression in Drosophila: microarray analysis of starvation and

sugar-dependent response. EMBO J. *21*, 6162–6173.

https://doi.org/10.1093/emboj/cdf600.

116. Macotela, Y., Boucher, J., Tran, T.T., and Kahn, C.R. (2009). Sex and depot

differences in adipocyte insulin sensitivity and glucose metabolism. Diabetes *58*, 803–812.

https://doi.org/10.2337/db08-1054.

117. Geer, E.B., and Shen, W. (2009). Gender differences in insulin resistance, body

composition, and energy balance. Gend. Med. *6 Suppl 1*, 60–75.

https://doi.org/10.1016/j.genm.2009.02.002.

118. Engel, F.L., and White, J.E. (1960). Some Hormonal Influences on Fat Mobilization

from Adipose Tissue. Am. J. Clin. Nutr. *8*, 691–704. https://doi.org/10.1093/ajcn/8.5.691.

119. Zhao, J., Wu, Y., Rong, X., Zheng, C., and Guo, J. (2020). Anti-Lipolysis Induced by

Insulin in Diverse Pathophysiologic Conditions of Adipose Tissue. Diabetes Metab. Syndr.

Obes. Targets Ther. *13*, 1575–1585. https://doi.org/10.2147/DMSO.S250699.

120. Chakrabarti, P., Kim, J.Y., Singh, M., Shin, Y.-K., Kim, J., Kumbrink, J., Wu, Y., Lee,

M.-J., Kirsch, K.H., Fried, S.K., et al. (2013). Insulin Inhibits Lipolysis in Adipocytes via the

Evolutionarily Conserved mTORC1-Egr1-ATGL-Mediated Pathway. Mol. Cell. Biol. *33*,

3659–3666. https://doi.org/10.1128/MCB.01584-12.

121. Scherer, T., O'Hare, J., Diggs-Andrews, K., Schweiger, M., Cheng, B., Lindtner, C.,

Zielinski, E., Vempati, P., Su, K., Dighe, S., et al. (2011). Brain Insulin Controls Adipose

Tissue Lipolysis and Lipogenesis. Cell Metab. *13*, 183–194.

https://doi.org/10.1016/j.cmet.2011.01.008.

122. Krycer, J.R., Quek, L.-E., Francis, D., Zadoorian, A., Weiss, F.C., Cooke, K.C.,

Nelson, M.E., Diaz-Vegas, A., Humphrey, S.J., Scalzo, R., et al. (2020). Insulin signaling

requires glucose to promote lipid anabolism in adipocytes. J. Biol. Chem. *295*, 13250–

13266. https://doi.org/10.1074/jbc.RA120.014907.

123. Rulifson, E.J., Kim, S.K., and Nusse, R. (2002). Ablation of Insulin-Producing

Neurons in Flies: Growth and Diabetic Phenotypes. Science *296*, 1118–1120.

https://doi.org/10.1126/science.1070058.

124. Clancy, D.J., Gems, D., Harshman, L.G., Oldham, S., Stocker, H., Hafen, E.,

Leevers, S.J., and Partridge, L. (2001). Extension of life-span by loss of CHICO, a

Drosophila insulin receptor substrate protein. Science *292*, 104–106.

https://doi.org/10.1126/science.1057991.

125. Millington, J.W., Brownrigg, G.P., Basner-Collins, P.J., Sun, Z., and Rideout, E.J.

(2021). Genetic manipulation of insulin/insulin-like growth factor signaling pathway activity

has sex-biased effects on Drosophila body size. G3 GenesGenomesGenetics *11*, jkaa067.

https://doi.org/10.1093/g3journal/jkaa067.

126. Grewal, S.S. (2009). Insulin/TOR signaling in growth and homeostasis: a view from

the fly world. Int. J. Biochem. Cell Biol. *41*, 1006–1010.

https://doi.org/10.1016/j.biocel.2008.10.010.

127. Teleman, A.A. (2009). Molecular mechanisms of metabolic regulation by insulin in

Drosophila. Biochem. J. *425*, 13–26. https://doi.org/10.1042/BJ20091181.

128. Osterwalder, T., Yoon, K.S., White, B.H., and Keshishian, H. (2001). A conditional

tissue-specific transgene expression system using inducible GAL4. Proc. Natl. Acad. Sci.

U. S. A. *98*, 12596–12601. https://doi.org/10.1073/pnas.221303298.

129. Roman, G., Endo, K., Zong, L., and Davis, R.L. (2001). P{Switch}, a system for

spatial and temporal control of gene expression in Drosophila melanogaster. Proc. Natl.

Acad. Sci. *98*, 12602–12607. https://doi.org/10.1073/pnas.221303998.

130. Poirier, L., Shane, A., Zheng, J., and Seroude, L. (2008). Characterization of the

Drosophila Gene-Switch system in aging studies: a cautionary tale. Aging Cell *7*, 758–770.

https://doi.org/10.1111/j.1474-9726.2008.00421.x.

131. Yamada, R., Deshpande, S.A., Keebaugh, E.S., Ehrlich, M.R., Soto Obando, A., and

Ja, W.W. (2017). Mifepristone Reduces Food Palatability and Affects Drosophila Feeding

and Lifespan. J. Gerontol. A. Biol. Sci. Med. Sci. *72*, 173–180.

https://doi.org/10.1093/gerona/glw072.

132. Landis, G.N., Salomon, M.P., Keroles, D., Brookes, N., Sekimura, T., and Tower, J.

(2015). The progesterone antagonist mifepristone/RU486 blocks the negative effect on life

span caused by mating in female Drosophila. Aging *7*, 53–69.

https://doi.org/10.18632/aging.100721.

133. McClure, C.D., Hassan, A., Aughey, G.N., Butt, K., Estacio-Gómez, A., Duggal, A.,

Ying Sia, C., Barber, A.F., and Southall, T.D. (2022). An auxin-inducible, GAL4-

compatible, gene expression system for Drosophila. eLife *11*, e67598.

https://doi.org/10.7554/eLife.67598.

134. Fleck, S.A., Biswas, P., DeWitt, E.D., Knuteson, R.L., Eisman, R.C., Nemkov, T.,

D'Alessandro, A., Tennessen, J.M., Rideout, E., and Weaver, L.N. (2024). Auxin exposure

disrupts feeding behavior and fatty acid metabolism in adult Drosophila. eLife *12*,

RP91953. https://doi.org/10.7554/eLife.91953.

135. McGuire, S.E., Le, P.T., Osborn, A.J., Matsumoto, K., and Davis, R.L. (2003).

Spatiotemporal rescue of memory dysfunction in Drosophila. Science *302*, 1765–1768.

https://doi.org/10.1126/science.1089035.

136. Klepsatel, P., Gáliková, M., Xu, Y., and Kühnlein, R.P. (2016). Thermal stress

depletes energy reserves in Drosophila. Sci. Rep. *6*, 33667.

https://doi.org/10.1038/srep33667.

137. Klepsatel, P., Wildridge, D., and Gáliková, M. (2019). Temperature induces changes

in Drosophila energy stores. Sci. Rep. *9*, 5239. https://doi.org/10.1038/s41598-019-41754-

5.

138. Tanabe, K., Itoh, M., and Tonoki, A. (2017). Age-Related Changes in Insulin-like

Signaling Lead to Intermediate-Term Memory Impairment in *Drosophila*. Cell Rep. *18*,

1598–1605. https://doi.org/10.1016/j.celrep.2017.01.053.

139. Enell, L.E., Kapan, N., Söderberg, J.A.E., Kahsai, L., and Nässel, D.R. (2010). Insulin

Signaling, Lifespan and Stress Resistance Are Modulated by Metabotropic GABA

Receptors on Insulin Producing Cells in the Brain of Drosophila. PLOS ONE *5*, e15780.

https://doi.org/10.1371/journal.pone.0015780.

140. Corl, A.B., Rodan, A.R., and Heberlein, U. (2005). Insulin signaling in the nervous

system regulates ethanol intoxication in Drosophila melanogaster. Nat. Neurosci. *8*, 18–

19. https://doi.org/10.1038/nn1363.

141. Haselton, A., Sharmin, E., Schrader, J., Sah, M., Poon, P., and Fridell, Y.-W.C.

(2010). Partial ablation of adult Drosophila insulin-producing neurons modulates glucose

homeostasis and extends life span without insulin resistance. Cell Cycle *9*, 3063–3071.

https://doi.org/10.4161/cc.9.15.12458.

142. Broughton, S.J., Piper, M.D.W., Ikeya, T., Bass, T.M., Jacobson, J., Driege, Y.,

Martinez, P., Hafen, E., Withers, D.J., Leevers, S.J., et al. (2005). Longer lifespan, altered

metabolism, and stress resistance in Drosophila from ablation of cells making insulin-like

ligands. Proc. Natl. Acad. Sci. U. S. A. *102*, 3105–3110.

https://doi.org/10.1073/pnas.0405775102.

143. Nässel, D.R., Kubrak, O.I., Liu, Y., Luo, J., and Lushchak, O.V. (2013). Factors that

regulate insulin producing cells and their output in Drosophila. Front. Physiol. *4*, 252.

https://doi.org/10.3389/fphys.2013.00252.

144. Kimura, K.D., Tissenbaum, H.A., Liu, Y., and Ruvkun, G. (1997). daf-2, an insulin

receptor-like gene that regulates longevity and diapause in Caenorhabditis elegans.

Science *277*, 942–946. https://doi.org/10.1126/science.277.5328.942.

145. Kenyon, C., Chang, J., Gensch, E., Rudner, A., and Tabtiang, R. (1993). A C.

elegans mutant that lives twice as long as wild type. Nature *366*, 461–464.

https://doi.org/10.1038/366461a0.

146. Partridge, L., Alic, N., Bjedov, I., and Piper, M.D.W. (2011). Ageing in Drosophila: The

role of the insulin/Igf and TOR signalling network. Exp. Gerontol. *46*, 376–381.

https://doi.org/10.1016/j.exger.2010.09.003.

147. Blüher, M., Kahn, B.B., and Kahn, C.R. (2003). Extended longevity in mice lacking

the insulin receptor in adipose tissue. Science *299*, 572–574.

https://doi.org/10.1126/science.1078223.

148. Fontana, L., Partridge, L., and Longo, V.D. (2010). Extending healthy life span--from

yeast to humans. Science *328*, 321–326. https://doi.org/10.1126/science.1172539.

149. Kuo, T.-H., Fedina, T.Y., Hansen, I., Dreisewerd, K., Dierick, H.A., Yew, J.Y., and

Pletcher, S.D. (2012). Insulin Signaling Mediates Sexual Attractiveness in Drosophila.

PLoS Genet. *8*, e1002684. https://doi.org/10.1371/journal.pgen.1002684.

150. Carvalho, G.B., Kapahi, P., Anderson, D.J., and Benzer, S. (2006). Allocrine

Modulation of Feeding Behavior by the Sex Peptide of *Drosophila*. Curr. Biol. *16*, 692–696.

https://doi.org/10.1016/j.cub.2006.02.064.

151. Hadjieconomou, D., King, G., Gaspar, P., Mineo, A., Blackie, L., Ameku, T., Studd,

C., de Mendoza, A., Diao, F., White, B.H., et al. (2020). Enteric neurons increase maternal

food intake during reproduction. Nature *587*, 455–459. https://doi.org/10.1038/s41586-020-

2866-8.

152. Cognigni, P., Bailey, A.P., and Miguel-Aliaga, I. (2011). Enteric Neurons and

Systemic Signals Couple Nutritional and Reproductive Status with Intestinal Homeostasis.

Cell Metab. *13*, 92–104. https://doi.org/10.1016/j.cmet.2010.12.010.

153. Ja, W.W., Carvalho, G.B., Zid, B.M., Mak, E.M., Brummel, T., and Benzer, S. (2009).

Water- and nutrient-dependent effects of dietary restriction on Drosophila lifespan. Proc.

Natl. Acad. Sci. U. S. A. *106*, 18633–18637. https://doi.org/10.1073/pnas.0908016106.

154. Lee, K.P., Simpson, S.J., Clissold, F.J., Brooks, R., Ballard, J.W.O., Taylor, P.W.,

Soran, N., and Raubenheimer, D. (2008). Lifespan and reproduction in Drosophila: New

insights from nutritional geometry. Proc. Natl. Acad. Sci. U. S. A. *105*, 2498–2503.

https://doi.org/10.1073/pnas.0710787105.

155. Chapman, T., and Partridge, L. (1996). Female fitness in Drosophila melanogaster:

an interaction between the effect of nutrition and of encounter rate with males. Proc. Biol.

Sci. *263*, 755–759. https://doi.org/10.1098/rspb.1996.0113.

156. Barnes, A.I., Wigby, S., Boone, J.M., Partridge, L., and Chapman, T. (2008). Feeding,

fecundity and lifespan in female Drosophila melanogaster. Proc. Biol. Sci. *275*, 1675–

1683. https://doi.org/10.1098/rspb.2008.0139.

157. Grandison, R.C., Piper, M.D.W., and Partridge, L. (2009). Amino-acid imbalance

explains extension of lifespan by dietary restriction in Drosophila. Nature *462*, 1061–1064.

https://doi.org/10.1038/nature08619.

158. Laturney, M., Sterne, G.R., and Scott, K. (2023). Mating activates neuroendocrine

pathways signaling hunger in Drosophila females. eLife *12*, e85117.

https://doi.org/10.7554/eLife.85117.

159. Strilbytska, O., Yurkevych, I., Semaniuk, U., Gospodaryov, D., Simpson, S.J., and

Lushchak, O. (2024). Life-History Trade-Offs in Drosophila: Flies Select a Diet to Maximize

Reproduction at the Expense of Lifespan. J. Gerontol. Ser. A *79*, glae057.

https://doi.org/10.1093/gerona/glae057.

160. Lee, K.P., Kim, J.-S., and Min, K.-J. (2013). Sexual dimorphism in nutrient intake and

life span is mediated by mating in *Drosophila melanogaster*. Anim. Behav. *86*, 987–992.

https://doi.org/10.1016/j.anbehav.2013.08.018.

161. Drummond-Barbosa, D., and Spradling, A.C. (2001). Stem cells and their progeny

respond to nutritional changes during Drosophila oogenesis. Dev. Biol. *231*, 265–278.

https://doi.org/10.1006/dbio.2000.0135.

162. Camus, M.F., Huang, C.-C., Reuter, M., and Fowler, K. (2018). Dietary choices are

influenced by genotype, mating status, and sex in Drosophila melanogaster. Ecol. Evol. *8,*

5385–5393. https://doi.org/10.1002/ece3.4055.

163. Ribeiro, C., and Dickson, B.J. (2010). Sex peptide receptor and neuronal TOR/S6K

signaling modulate nutrient balancing in Drosophila. Curr. Biol. CB *20*, 1000–1005.

https://doi.org/10.1016/j.cub.2010.03.061.

164. Hsu, H.-J., and Drummond-Barbosa, D. (2009). Insulin levels control female germline

stem cell maintenance via the niche in Drosophila. Proc. Natl. Acad. Sci. *106*, 1117–1121.

https://doi.org/10.1073/pnas.0809144106.

165. Burn, K.M., Shimada, Y., Ayers, K., Lu, F., Hudson, A.M., and Cooley, L. (2015).

Somatic insulin signaling regulates a germline starvation response in Drosophila egg

chambers. Dev. Biol. *398*, 206–217. https://doi.org/10.1016/j.ydbio.2014.11.021.

166. LaFever, L., and Drummond-Barbosa, D. (2005). Direct control of germline stem cell

division and cyst growth by neural insulin in Drosophila. Science *309*, 1071–1073.

https://doi.org/10.1126/science.1111410.

167. Richard, D.S., Rybczynski, R., Wilson, T.G., Wang, Y., Wayne, M.L., Zhou, Y.,

Partridge, L., and Harshman, L.G. (2005). Insulin signaling is necessary for vitellogenesis

in Drosophila melanogaster independent of the roles of juvenile hormone and

ecdysteroids: female sterility of the chico1 insulin signaling mutation is autonomous to the

ovary. J. Insect Physiol. *51*, 455–464. https://doi.org/10.1016/j.jinsphys.2004.12.013.

168. Chandegra, B., Tang, J.L.Y., Chi, H., and Alic, N. (2017). Sexually dimorphic effects

of dietary sugar on lifespan, feeding and starvation resistance in Drosophila. Aging *9*,

2521–2528. https://doi.org/10.18632/aging.101335.

169. Eichmann, T.O., and Lass, A. (2015). DAG tales: the multiple faces of

diacylglycerol—stereochemistry, metabolism, and signaling. Cell. Mol. Life Sci. CMLS *72*,

3931–3952. https://doi.org/10.1007/s00018-015-1982-3.

170. Antony, C., and Jallon, J.-M. (1982). The chemical basis for sex recognition in

Drosophila melanogaster. J. Insect Physiol. *28*, 873–880. https://doi.org/10.1016/0022-

1910(82)90101-9.

171. Jallon, J.M. (1984). A few chemical words exchanged by Drosophila during courtship

and mating. Behav. Genet. *14*, 441–478. https://doi.org/10.1007/BF01065444.

172. Fedina, T.Y., Kuo, T.-H., Dreisewerd, K., Dierick, H.A., Yew, J.Y., and Pletcher, S.D.

(2012). Dietary Effects on Cuticular Hydrocarbons and Sexual Attractiveness in

Drosophila. PLoS ONE *7*, e49799. https://doi.org/10.1371/journal.pone.0049799.

173. Wicker-Thomas, C., Garrido, D., Bontonou, G., Napal, L., Mazuras, N., Denis, B.,

Rubin, T., Parvy, J.-P., and Montagne, J. (2015). Flexible origin of

hydrocarbon/pheromone precursors in Drosophila melanogaster. J. Lipid Res. *56*, 2094–

2101. https://doi.org/10.1194/jlr.M060368.

174. Scheitz, C.J.F., Guo, Y., Early, A.M., Harshman, L.G., and Clark, A.G. (2013).

Heritability and Inter-Population Differences in Lipid Profiles of Drosophila melanogaster.

PLOS ONE *8*, e72726. https://doi.org/10.1371/journal.pone.0072726.

175. Faust, J.E., Manisundaram, A., Ivanova, P.T., Milne, S.B., Summerville, J.B., Brown,

H.A., Wangler, M., Stern, M., and McNew, J.A. (2014). Peroxisomes Are Required for Lipid

Metabolism and Muscle Function in Drosophila melanogaster. PLOS ONE *9*, e100213.

https://doi.org/10.1371/journal.pone.0100213.

176. Lodhi, I.J., and Semenkovich, C.F. (2014). Peroxisomes: a Nexus for Lipid

Metabolism and Cellular Signaling. Cell Metab. *19*, 380–392.

https://doi.org/10.1016/j.cmet.2014.01.002.

177. Pridie, C., Ueda, K., and Simmonds, A.J. (2020). Rosy Beginnings: Studying

Peroxisomes in Drosophila. Front. Cell Dev. Biol. *8*.

https://doi.org/10.3389/fcell.2020.00835.

178. Tuthill, B.F., Searcy, L.A., Yost, R.A., and Musselman, L.P. (2020). Tissue-specific

analysis of lipid species in Drosophila during overnutrition by UHPLC-MS/MS and MALDI-

MSI [S]. J. Lipid Res. *61*, 275–290. https://doi.org/10.1194/jlr.RA119000198.

179. Santoro, C., O'Toole, A., Finsel, P., Alvi, A., and Musselman, L.P. (2022). Reducing

ether lipids improves Drosophila overnutrition-associated pathophysiology phenotypes via

a switch from lipid storage to beta-oxidation. Sci. Rep. *12*, 13021.

https://doi.org/10.1038/s41598-022-16870-4.

180. Alfa, R.W., Park, S., Skelly, K.-R., Poffenberger, G., Jain, N., Gu, X., Kockel, L.,

Wang, J., Liu, Y., Powers, A.C., et al. (2015). Suppression of Insulin Production and

Secretion by a Decretin Hormone. Cell Metab. *21*, 323–333.

https://doi.org/10.1016/j.cmet.2015.01.006.

181. Luo, J., Lushchak, O.V., Goergen, P., Williams, M.J., and Nässel, D.R. (2014).

Drosophila Insulin-Producing Cells Are Differentially Modulated by Serotonin and

Octopamine Receptors and Affect Social Behavior. PLOS ONE *9*, e99732.

https://doi.org/10.1371/journal.pone.0099732.

182. Söderberg, J.A.E., Carlsson, M.A., and Nässel, D.R. (2012). Insulin-Producing Cells

in the Drosophila Brain also Express Satiety-Inducing Cholecystokinin-Like Peptide,

Drosulfakinin. Front. Endocrinol. *3*, 109. https://doi.org/10.3389/fendo.2012.00109.

183. Yurgel, M.E., Kakad, P., Zandawala, M., Nässel, D.R., Godenschwege, T.A., and

Keene, A.C. (2019). A single pair of leucokinin neurons are modulated by feeding state

and regulate sleep–metabolism interactions. PLoS Biol. *17*, e2006409.

https://doi.org/10.1371/journal.pbio.2006409.

184. Kwak, S.-J., Hong, S.-H., Bajracharya, R., Yang, S.-Y., Lee, K.-S., and Yu, K. (2013).

Drosophila Adiponectin Receptor in Insulin Producing Cells Regulates Glucose and Lipid

Metabolism by Controlling Insulin Secretion. PLOS ONE *8*, e68641.

https://doi.org/10.1371/journal.pone.0068641.

185. Buch, S., Melcher, C., Bauer, M., Katzenberger, J., and Pankratz, M.J. (2008).

Opposing Effects of Dietary Protein and Sugar Regulate a Transcriptional Target of

Drosophila Insulin-like Peptide Signaling. Cell Metab. *7*, 321–332.

https://doi.org/10.1016/j.cmet.2008.02.012.

186. Lee, G., Bahn, J.H., and Park, J.H. (2006). Sex- and clock-controlled expression of

the neuropeptide F gene in Drosophila. Proc. Natl. Acad. Sci. *103*, 12580–12585.

https://doi.org/10.1073/pnas.0601171103.

187. Asahina, K., Watanabe, K., Duistermars, B.J., Hoopfer, E., González, C.R.,

Eyjólfsdóttir, E.A., Perona, P., and Anderson, D.J. (2014). Tachykinin-expressing neurons

control male-specific aggressive arousal in Drosophila. Cell *156*, 221–235.

https://doi.org/10.1016/j.cell.2013.11.045.

188. Shankar, S., Chua, J.Y., Tan, K.J., Calvert, M.E.K., Weng, R., Ng, W.C., Mori, K., and

Yew, J.Y. (2015). The neuropeptide tachykinin is essential for pheromone detection in a

gustatory neural circuit. eLife *4*, e06914. https://doi.org/10.7554/eLife.06914.

189. Kim, W.J., Jan, L.Y., and Jan, Y.N. (2013). A PDF/NPF neuropeptide signaling

circuitry of male Drosophila melanogaster controls rival-induced prolonged mating. Neuron

80, 1190–1205. https://doi.org/10.1016/j.neuron.2013.09.034.

190. Liu, C., Zhang, B., Zhang, L., Yang, T., Zhang, Z., Gao, Z., and Zhang, W. (2020). A

neural circuit encoding mating states tunes defensive behavior in Drosophila. Nat.

Commun. *11*, 3962. https://doi.org/10.1038/s41467-020-17771-8.

191. Lebreton, S., Mansourian, S., Bigarreau, J., and Dekker, T. (2016). The Adipokinetic

Hormone Receptor Modulates Sexual Behavior, Pheromone Perception and Pheromone

Production in a Sex-Specific and Starvation-Dependent Manner in Drosophila

melanogaster. Front. Ecol. Evol. *3*. https://doi.org/10.3389/fevo.2015.00151.

192. Rezával, C., Nojima, T., Neville, M.C., Lin, A.C., and Goodwin, S.F. (2014). Sexually

dimorphic octopaminergic neurons modulate female postmating behaviors in Drosophila.

Curr. Biol. CB *24*, 725–730. https://doi.org/10.1016/j.cub.2013.12.051.

193. Wang, T., Jing, B., Deng, B., Shi, K., Li, J., Ma, B., Wu, F., and Zhou, C. (2022).

Drosulfakinin signaling modulates female sexual receptivity in Drosophila. eLife *11*,

e76025. https://doi.org/10.7554/eLife.76025.

194. Ameku, T., Yoshinari, Y., Texada, M.J., Kondo, S., Amezawa, K., Yoshizaki, G.,

Shimada-Niwa, Y., and Niwa, R. (2018). Midgut-derived neuropeptide F controls germline

stem cell proliferation in a mating-dependent manner. PLOS Biol. *16*, e2005004.

https://doi.org/10.1371/journal.pbio.2005004.

195. Zhang, C., Daubnerova, I., Jang, Y.-H., Kondo, S., Žitňan, D., and Kim, Y.-J. (2021).

The neuropeptide allatostatin C from clock-associated DN1p neurons generates the

circadian rhythm for oogenesis. Proc. Natl. Acad. Sci. U. S. A. *118*, e2016878118.

https://doi.org/10.1073/pnas.2016878118.

196. Lin, W.-S., Yeh, S.-R., Fan, S.-Z., Chen, L.-Y., Yen, J.-H., Fu, T.-F., Wu, M.-S., and

Wang, P.-Y. (2018). Insulin signaling in female Drosophila links diet and sexual

attractiveness. FASEB J. *32*, 3870–3877. https://doi.org/10.1096/fsb2fj201800067r.

197. Cong, X., Wang, H., Liu, Z., He, C., An, C., and Zhao, Z. (2015). Regulation of Sleep

by Insulin-like Peptide System in Drosophila melanogaster. Sleep *38*, 1075–1083.

https://doi.org/10.5665/sleep.4816.

198. Isabel, G., Martin, J.-R., Chidami, S., Veenstra, J.A., and Rosay, P. (2005). AKH-

producing neuroendocrine cell ablation decreases trehalose and induces behavioral

changes in Drosophila. Am. J. Physiol. Regul. Integr. Comp. Physiol. *288*, R531-538.

https://doi.org/10.1152/ajpregu.00158.2004.

199. Bednářová, A., Tomčala, A., Mochanová, M., Kodrík, D., and Krishnan, N. (2018).

Disruption of Adipokinetic Hormone Mediated Energy Homeostasis Has Subtle Effects on

Physiology, Behavior and Lipid Status During Aging in Drosophila. Front. Physiol. *9*, 949.

https://doi.org/10.3389/fphys.2018.00949.

200. Post, S., Liao, S., Yamamoto, R., Veenstra, J.A., Nässel, D.R., and Tatar, M. (2019).

Drosophila insulin-like peptide dilp1 increases lifespan and glucagon-like Akh expression

epistatic to dilp2. Aging Cell *18*, e12863. https://doi.org/10.1111/acel.12863.

201. Stephenson, E.J., Stayton, A.S., Sethuraman, A., Rao, P.K., Meyer, A., Gomes, C.K.,

Mulcahy, M.C., McAllan, L., Puchowicz, M.A., Pierre, J.F., et al. (2022). Chronic intake of

high dietary sucrose induces sexually dimorphic metabolic adaptations in mouse liver and

adipose tissue. Nat. Commun. *13*, 6062. https://doi.org/10.1038/s41467-022-33840-6.

202. Mauvais-Jarvis, F., Merz, N.B., Barnes, P.J., Brinton, R.D., Carrero, J.-J., DeMeo,

D.L., Vries, G.J.D., Epperson, C.N., Govindan, R., Klein, S.L., et al. (2020). Sex and

gender: modifiers of health, disease, and medicine. The Lancet *396*, 565–582.

https://doi.org/10.1016/S0140-6736(20)31561-0.

203. Mauvais-Jarvis, F. (2015). Sex differences in metabolic homeostasis, diabetes, and

obesity. Biol. Sex Differ. *6*, 14. https://doi.org/10.1186/s13293-015-0033-y.

204. Chella Krishnan, K., Mehrabian, M., and Lusis, A.J. (2018). Sex differences in

metabolism and cardiometabolic disorders. Curr. Opin. Lipidol. *29*, 404–410.

https://doi.org/10.1097/MOL.0000000000000536.

205. Tramunt, B., Smati, S., Grandgeorge, N., Lenfant, F., Arnal, J.-F., Montagner, A., and

Gourdy, P. (2020). Sex differences in metabolic regulation and diabetes susceptibility.

Diabetologia *63*, 453–461. https://doi.org/10.1007/s00125-019-05040-3.

206. Strack, C., Behrens, G., Sag, S., Mohr, M., Zeller, J., Lahmann, C., Hubauer, U.,

Loew, T., Maier, L., Fischer, M., et al. (2022). Gender differences in cardiometabolic health

and disease in a cross-sectional observational obesity study. Biol. Sex Differ. *13*, 8.

https://doi.org/10.1186/s13293-022-00416-4.

207. Veenstra, J.A., Agricola, H.-J., and Sellami, A. (2008). Regulatory peptides in fruit fly

midgut. Cell Tissue Res. *334*, 499–516. https://doi.org/10.1007/s00441-008-0708-3.

208. Kasturacharya, N., Dhall, J.K., and Hasan, G. (2023). A STIM dependent dopamine-

neuropeptide axis maintains the larval drive to feed and grow in Drosophila. PLOS Genet.

19, e1010435. https://doi.org/10.1371/journal.pgen.1010435.

209. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source

platform for biological-image analysis. Nat. Methods *9*, 676–682.

https://doi.org/10.1038/nmeth.2019.

210. Yu, H., and Huan, T. (2022). MAFFIN: metabolomics sample normalization using

maximal density fold change with high-quality metabolic features and corrected signal

intensities. Bioinformatics *38*, 3429–3437. https://doi.org/10.1093/bioinformatics/btac355.

211. May, C.E., Vaziri, A., Lin, Y.Q., Grushko, O., Khabiri, M., Wang, Q.-P., Holme, K.J.,

Pletcher, S.D., Freddolino, P.L., Neely, G.G., et al. (2019). High Dietary Sugar Reshapes

Sweet Taste to Promote Feeding Behavior in Drosophila melanogaster. Cell Rep. *27*,

1675-1685.e7. https://doi.org/10.1016/j.celrep.2019.04.027.

212. Nunes, R.D., and Drummond-Barbosa, D. (2023). A high-sugar diet, but not obesity,

reduces female fertility in Drosophila melanogaster. Dev. Camb. Engl. *150*, dev201769.

https://doi.org/10.1242/dev.201769.

213. Li, Y., Wang, W., and Lim, H.-Y. (2021). Drosophila Solute Carrier 5A5 Regulates

Systemic Glucose Homeostasis by Mediating Glucose Absorption in the Midgut. Int. J.

Mol. Sci. *22*, 12424. https://doi.org/10.3390/ijms222212424.zf

214. LeDue, E.E., Mann, K., Koch, E., Chu, B., Dakin, R., and Gordon, M.D. (2016).

Starvation-Induced Depotentiation of Bitter Taste in Drosophila. Curr. Biol. CB *26*, 2854–

2861. https://doi.org/10.1016/j.cub.2016.08.028.

215. Marella, S., Mann, K., and Scott, K. (2012). Dopaminergic modulation of sucrose

acceptance behavior in Drosophila. Neuron *73*, 941–950.

https://doi.org/10.1016/j.neuron.2011.12.032.

216. Wobbrock, J. O., Findlater, L., Gergle, D., & Higgins, J. J. (2011). The aligned rank

transform for nonparametric factorial analyses using only anova procedures. Proceedings

of the SIGCHI Conference on Human Factors in Computing Systems, 143–146.

https://doi.org/10.1145/1978942.1978963

1819

Figure 1

Figure 2

A — Sex bias in lipid abundance

Legend: TG, DG, PC, ST, TG-O, PI, PC-O, LPC, PG, PE, PE-P, Cer, PS, PE-O, NAE, SE, Acyl-Car

y-axis: p-value (10^{-8}, 10^{-6}, 10^{-4}, 0.01, 1)
x-axis: log2 fold change (-12 -10 -8 -6 -4 -2 0 2 4 6 8 10 12)

46 lipids   117 lipids

Male-biased   Female-biased

B — Sex bias of differentially abundant lipids

Category		Ratio
TG-O		1/13
TG		48/57
ST		4/4
SE		4/4
PI		3/3
PG		3/3
PE		17/20
PC-O		4/6
PC		16/25
LPC		1/3
DG		6/9
Cer		4/7
Acyl-Car		4/5

x-axis: Proportion (0.0, 0.5, 1.0)

■ Female   ■ Male

C — Sex bias in metabolite abundance

y-axis: log2 fold change (-5, 0, 5, 10)

Female (top), Male (bottom)

x-axis metabolites: (-)-Riboflavin, LPC, 2-Aminopyridine, ADP, Asparagine, Aspartic acid, Butyryl carnitine, Cytidine, D-(+)-Pantothenic acid, Glucose-6-phosphate, Glutamic acid, Glutamine, Guanosine, Histamine, Hypoxanthine, Inosine, Isonicotinic acid, Kynurenine, L-5-Oxoproline, L-Pipecolic acid, Lysine, Methioninesulfoxide, N,N-Dimethylglycine, N-Acetylhistamine, Trimethyllysine, Ornithine, Oxidized glutathione, Phosphocholine, Propionylcarnitine, Pyroglutamic acid, Sweroside, Thermopsine, Thiazolidine-4-carboxylic acid, trans-4-Hydroxy-L-proline, Uracil, Undine, UDP-acetylglucosamine

D — Sugar-dependent change in lipid abundance (females)

Legend: TG, DG, PC, ST, TG-O, PI, PC-O, SE, PG, PE, PE-P, Cer, PS, PE-O, NAE, Hex-Cer, Acyl-Car

y-axis: p-value (10^{-8}, 10^{-6}, 10^{-4}, 0.01, 1)
x-axis: log2 fold change (-12 -10 -8 -6 -4 -2 0 2 4 6 8 10 12)

64 lipids   154 lipids

upregulated   downregulated

E — Sugar-dependent change in lipid abundance (males)

Legend: TG, ST, PI, PC, PE, TG-O, Acyl-CAR, Cer, PC-O, Hex-Cer

y-axis: p-value (10^{-8}, 10^{-6}, 10^{-4}, 0.01, 1)
x-axis: log2 fold change (-12 -10 -8 -6 -4 -2 0 2 4 6 8 10 12)

10 lipids   17 lipids

upregulated   downregulated

F — Sugar-dependent change in metabolite abundance (females)

y-axis: p-value (10^{-8}, 10^{-6}, 10^{-4}, 10^{-2}, 0.05, 1)
x-axis: log2 fold change (-12 -10 -8 -6 -4 -2 0 2 4 6 8 10 12)

Labels: Asparagine, Glutamic acid, Kynurenine, Glucose-6-phosphate, L-Ascorbic Acid, N-Acetylhistamine

upregulated   downregulated

G — Sugar-dependent change in metabolite abundance (males)

y-axis: p-value (10^{-8}, 10^{-6}, 10^{-4}, 10^{-2}, 0.05, 1)
x-axis: log2 fold change (-12 -10 -8 -6 -4 -2 0 2 4 6 8 10 12)

Labels: Adenosine-3'-moonophosphate, Kynurenine, Trigonelline, 3-Hydroxy-kynurenine

upregulated   downregulated

H — Glucose levels in Excreta

y-axis: Glucose concentration (per fly mg/ml) (0.000, 0.002, 0.004, 0.006, 0.008, 0.010)

*

Female   Male

1S diet

Figure 3

Figure 4