Pathogen threat proximity shapes host extracellular vesicle production in pre-infection response

Klaudia Kołodziejska[1], Agata Szczepańska[1], Satya Vadlamani[1#], Ramakrishnan Ponath Sukumaran[1#], Mariusz Radkiewicz[2], Henrik Bringmann[3], Nathalie Pujol[4], Wojciech Pokrzywa[5], Michał Turek[1]

Affiliations:

[1]*Laboratory of Animal Molecular Physiology, Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Warsaw, Poland*

[2]*Mass Spectrometry Facility, Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Warsaw, Poland*

[3]*Biotechnology Center, Center for Molecular and Cellular Bioengineering, Technische Universität Dresden, Dresden, Germany*

[4]*Aix Marseille Univ, INSERM, CNRS, CIML, Turing Centre for Living Systems, 163 Avenue de Luminy, case 906, 13009 Marseille, France*

[5]*Laboratory of Protein Metabolism, International Institute of Molecular and Cell Biology in Warsaw, Poland*

[#] These authors contributed equally

Correspondence should be directed to:

MT: m.turek@ibb.waw.pl

Abstract

Extracellular vesicles (EVs) play a crucial role in immune responses, yet it remains unclear whether pathogen metabolites alone can stimulate EV production prior to infection. Using *Caenorhabditis elegans*, we investigate this question through the lens of exophers – large, evolutionarily conserved EVs known to enhance proteostasis, extend lifespan, and improve reproductive fitness. Our study uncovers distinct regulatory mechanisms driving EV production in response to pathogen-derived volatile metabolites and non-volatile secretome, providing insights into host-pathogen signaling before physical interaction. We reveal a sophisticated network that adjusts EV production based on pathogen proximity: non-volatile secretome (including tripeptide Ile-Pro-Pro), signaling an immediate threat, activate immune-dependent EV pathways, while volatile metabolites, forewarning potential danger, initiate immunity-independent exopher production. Both responses rely on sensory neurons, with ASK, ADL, and AWC neurons playing central roles in each, and additional input from ASI, AWB, and ASH neurons specifically enhancing non-volatile secretome-induced EV production. Downstream signal integration is mediated by a circuit involving the RMG hub and AIB/AIA interneurons, with the AIB-specific NPR-9 receptor playing a critical role. Molecularly, we identified multiple GPCRs, including SRI-19, SRI-36/39, and SRR-6, as crucial for the non-volatile response, and revealed that SRR-6 functions in the intestine to regulate muscle exopher production, demonstrating a novel gut-muscle signaling axis. Notably, exposure to volatile pathogen metabolites significantly boosts offspring fitness during infection in SRI-19-dependent manner, albeit at the cost of maternal survival when conditions are being harsher. Thus, our study shows that *C. elegans* distinguishes pathogen proximity through compound types, activating EV-dependent physiological responses tailored to either immediate or anticipated threats to optimize survival across generations.

Keywords

C. elegans, extracellular vesicles, exophers, pathogens, immune response, olfactory neurons,

GPCRs

Introduction

Extracellular vesicles (EVs), including exosomes and microvesicles, are pivotal in the host

immune response by mediating intercellular communication and regulating immune functions.

Released by various cell types, EVs carry a diverse array of bioactive molecules such as

antigens, signalling molecules, and genetic material, thus influencing immune cell behaviour

and activity[1,2]. Pathogen-derived EVs often contain pathogen-associated molecular patterns

(PAMPs) that can activate immune responses[3], while host-derived EVs help modulate

inflammation and immune activation, maintaining homeostasis and enhancing defence

mechanisms against pathogens[4]. These roles for EVs have been well-documented during active

infection, where pathogens are already invading the host. However, pathogens release both

volatile and non-volatile compounds, which can alert and prepare the host to their exposure[5–7].

Whether such compounds can modulate EV production prior to infection is not known.

The nematode *Caenorhabditis elegans* is a widely used model for studying host-pathogen

interactions, particularly valuable for examining its responses to clinically significant human

pathogens such as *Pseudomonas aeruginosa*, *Serratia marcescens*, and *Staphylococcus*

aureus[8]. This organism offers several advantages due to its simplicity, genetic tractability, and

well-characterized immune responses. *C. elegans* relies solely on innate immune pathways,

making it an ideal organism for dissecting fundamental aspects of innate immunity without the

confounding influence of an adaptive immune system[9,10]. Additionally, its short lifespan and

large brood size enable high-throughput genetic and pharmacological screens[11]. The

evolutionary conservation of many immune pathways between *C. elegans* and higher organisms

further enhances its relevance for studying host-pathogen interactions[12,13]. Moreover, *C.*

elegans uses specialised G protein-coupled receptor (GPCR)-mediated sensory circuits to

detect environmental pathogens and orchestrate systemic outputs, including immune activation

and behavioural avoidance, in response to microbial metabolites such as phenazine-1-

carboxamide, pyochelin[14], and volatile 1-undecene[15].

C. elegans has also emerged as a valuable model for studying EV production. Due to its

transparent body and well-characterized genetics, *C. elegans* allows tracking of EV production

in real-time and investigation of the regulatory pathways involved[16]. One of the examples of an

evolutionarily conserved class of particularly large EVs that is investigated in worms are

exophers. Exophers have been shown to be involved in waste removal and material exchange

between tissues, with key functions in cellular stress responses, tissue maintenance, and

reproduction[17–22]. Previous studies have demonstrated that exophers help protect neuronal

activity in *C. elegans* by expelling toxic cellular debris[17], a process also observed in mammals,

where mouse cardiomyocytes release exophers to discard defective mitochondria, maintaining

metabolic balance and preventing inflammation[23]. In addition to waste disposal, exophers have

been shown to facilitate inter-tissue transport to facilitate reproductive fitness[18], a process that

can be regulated by social signalling derived from other animals within the population[22].

In this context, we aimed to investigate whether EV production in a host organism can be

modulated by the mere presence of pathogens in the environment. Importantly, we focused on

the response to pathogen volatile and non-volatile secretomes, allowing us to decouple this

regulation from direct infection-related responses. Using exophers released by *C. elegans* body

wall muscles as a model, we sought to explore the molecular mechanisms underlying pathogen

secretomes-induced EV dynamics.

Our study uncovers a complex regulatory network governing exopher production in *C. elegans*,

shaped by the type of interaction with pathogens. Pathogen infection or exposure to their

secretomes increases exopher production. However, the molecular mechanisms regulating

volatile metabolite-dependent and non-volatile secretome-dependent increases in EV

production are distinctly controlled at immune, sensory, and neuronal levels. Our research

demonstrates that worms can differentiate the proximity of a pathogen threat based on the type

of compound they detect. Non-volatile secreted substances, typically sensed at close range,

signal the pathogen's immediate presence, whereas volatile metabolites, detected from a greater

distance, indicate a nearby pathogen that may pose a lower threat. This ability to distinguish

between signals triggers a context-specific physiological response, which may or may not

involve immune activation, depending on the nature of the detected compounds. These findings

provide new insights into the physiological mechanisms of EV production in *C. elegans* and

suggest that similar mechanisms may be conserved across species.

Results

Infection with pathogens induces exopher production in hermaphrodite muscles

To test if pathogen infection influences production of extracellular vesicles in *C. elegans*, we

exposed worms at adulthood day 1 to live *Pseudomonas aeruginosa* PA14 or *Serratia*

marcescens Db10 bacteria for 3 hours. Next, we moved them back to plates seeded with a non-

pathogenic *Escherichia coli* OP50 bacteria and, 24 hours later, we quantified the number of

exophers released by body-wall muscle cells (BWMs) following a previously described

methodology[24] (**Fig. 1a**). Our results show that worms exposed to both types of pathogenic

bacteria produce significantly more exophers than worms that were kept on *E. coli* OP50

bacteria throughout the whole experiment (**Fig. 1b**).

Given that extracellular vesicles in other organisms are often part of the immune response[1,25],

we next examined whether basal production of exophers as well as increased exopher

production upon exposure to pathogenic bacteria depends on specific immune pathways. We

tested various mutants deficient in key nematode immune response pathways[10,26], including the

insulin pathway (*akt-1*, *daf-16*)[27], TGFβ pathway (*dbl-1*)[28], p38 MAPK pathway (*tir-1*, *pmk-*

1)[29,30], and Intracellular Pathogen Response (IPR) pathway (*cul-6*)[31,32] (**Fig. 1c**). In all tested

immune response pathways, we observed altered levels of exophergenesis, either decreased or

increased, compared to wild-type worms. This was observed under standard growing conditions

on non-pathogenic *E. coli* OP50 bacteria after knocking out at least one component of each

pathway (**Fig. 1d**). Surprisingly, despite their altered basal levels, most of the tested mutants

still exhibited a robust, statistically significant increase in exopher production upon pathogen

exposure (**Fig. 1e-g**). The sole exception was the *cul-6* mutant with a compromised IPR

pathway, which showed an increase in exopher production upon exposure to *P. aeruginosa* and

a decrease in exopher production upon exposure to *S. marcescens* when compared to the control

(**Fig. 1h**). Among the immune-deficient strains we tested, the magnitude of pathogen-induced

exopher upregulation differed. Following *P. aeruginosa* exposure, the *dbl-1*, *pmk-1*, and *cul-6*

mutants exhibited increases in exopher production comparable to wild-type levels, whereas

tir-1 mutants showed an even greater induction. In contrast, *daf-16* mutants displayed a

significantly diminished response relative to wild-type, while *akt-1* mutants, defective in the

negative regulator of DAF-16, exhibited an exaggerated increase (**Supplementary Fig. 1a**).

Because AKT-1 inhibits DAF-16 in the insulin pathway, these data indicate that although

exopher induction by pathogen presence is broadly robust, its amplitude is fine-tuned by

DAF-16-dependent transcriptional programs as well as by p38 MAPK pathway.

A similar pattern emerged when worms were challenged with *S. marcescens*, with the exception

that *tir-1* mutants responded like wild-type and *cul-6* mutants showed a significant reduction in

exopher induction (**Supplementary Fig. 1b**).

Together, these findings indicate that pathogen-triggered exopher upregulation does not rely on

any single canonical immune pathway, although we cannot rule out redundant or compensatory

contributions, and that the insulin/DAF-16 axis plays a primary role in modulating the response

magnitude. Other pathways appear to play less central roles in determining the extent of the

upregulation of exopher formation in response to pathogen.

Previous work has established that exopher biogenesis requires fertility and the presence of

embryos in the uterus[18,33]. To determine whether pathogen exposure alone can bypass this

requirement, we exposed sterile, FUdR-treated worms to *P. aeruginosa* and *S. marcescens*. In

both cases, sterile animals failed to produce exophers (**Supplementary Fig. 1c**), demonstrating

that fertility remains a prerequisite for pathogen-induced exophergenesis. When we examined

embryo accumulation, *S. marcescens* exposure led to an increase in uterine embryos, whereas

P. aeruginosa-exposed worms showed no change in embryo accumulation (**Fig. 1i**). Therefore,

although both pathogens stimulate exopher production in fertile animals, only *S. marcescens*

does so alongside embryo accumulation. This observation is critically important, as it shows

that *P. aeruginosa* activates exophergenesis through a novel signalling pathway that is

independent of the classic mechanism linked to embryo accumulation. This model allows for

the isolation and study of the direct response to a pathogenic signal without the confounding

influence of maternal signals. For this reason, we decided to focus our subsequent mechanistic

analyses specifically on exophergenesis induced by *P. aeruginosa*.

Pathogen non-volatile and volatile secretomes trigger exopher production

Recent reports indicate that *C. elegans* can detect pathogen volatile metabolites (all airborne

compounds emitted by the pathogen), which influence their physiology in a cell non-

autonomous manner[15,34]. Previous studies have also shown that exposing *C. elegans* to the *P.*

aeruginosa secretome (the full complement of water-soluble factors present in clarified culture

supernatant including small metabolites, peptides, proteins, and extracellular vesicles) leads to

a robust immune response[35]. We hence aimed to determine whether the increased production

of EVs in response to pathogens requires direct infection or can be triggered simply by exposure

to volatile and soluble bacterial components.

We first investigated whether exposure to pathogen odours alone would be sufficient to induce

exopher production in muscle cells. To test this, we covered an NGM plate seeded with OP50

and containing age-synchronized L4-stage worms with another NGM plate containing *P.*

aeruginosa and sealed both plates together with parafilm. Worms were incubated in this setup

for 48 hours until they reached adulthood day 2, after which exophers were counted (**Fig. 2a**).

Our results demonstrate that exposure to *P. aeruginosa* volatile metabolites indeed induces

exopher production (**Fig. 2b and c**) and also causes a slight embryo accumulation *in utero* (**Fig.**

2d).

Second, we aimed to determine whether the increased production of EVs in response to

pathogens is triggered by exposure to the bacterial non-volatile secretome (**Fig. 2e**). As shown

in **Fig. 2f and g**, direct infection with *P. aeruginosa* is not necessary; worms grown on plates

with *P. aeruginosa* secretome also produced more exophers. Notably, this increase in exopher

production is not associated with increased embryo accumulation in the worm's uterus (**Fig.**

2h). The fact that *P. aeruginosa* non-volatile secretome fails to increase embryo accumulation

in utero, in contrast to the retention induced by volatile metabolites, suggests that non-volatile

pathogen cues may actively suppress the embryo-accumulation response typically triggered by

volatiles. This opposing outcome points to distinct downstream signalling branches that

modulate reproductive physiology in response to different classes of pathogen-derived

compounds. This also aligns with our observation that simultaneous exposure to volatile

metabolites and the non-volatile secretome is non-additive for exopher production (**Fig. 2i**).

Similar to previous results, worms sterilized by FUdR failed to upregulate exopher production

in response to either *P. aeruginosa* volatile metabolites or non-volatile secretome

(**Supplementary Fig. 2a and d**). This demonstrates that fertility is essential for exopher

upregulation by both volatile and non-volatile pathogen cues. Moreover, because starvation

alone is sufficient to increase exopher production[18,21], we monitored pharyngeal pumping to

rule out reduced feeding as a confounding factor. We found no significant change in pumping

rates following exposure to either pathogen volatile metabolites or non-volatile secretome

(**Supplementary Fig. 2b and e**), confirming that the observed exopher upregulation is not

driven by altered food intake.

We further explored whether neuronal exophers, previously shown to help maintain

proteostasis in neurons, are similarly regulated by the pathogen-derived secretome. Using a

touch-receptor neuron reporter strain[17], we exposed animals to either volatile metabolites or

non-volatile *P. aeruginosa* secretome as described above and quantified neuronal exophers.

Strikingly, we found that volatile metabolites, but not non-volatile secretome, significantly

upregulate exopher formation in neurons (**Supplementary Fig. 2c and f**). This tissue-specific

responsiveness to soluble secretome components suggests that the signalling machinery

governing EV production varies between muscle and neuronal cells, with only volatile cues

triggering the neuronal exopher response.

Volatile and non-volatile cues engage distinct immune-regulatory mechanisms

Next, we investigated whether the increased muscle exopher production in our new

experimental setups is influenced by innate immunity signalling pathways. We first assessed

each immune response mutant against its own unexposed baseline and then compared the

magnitude of PA14-induced changes to wild-type responses. On one hand, our data show that

insulin, TGFβ, p38 MAPK, and IPR pathways are not individually required for the robust

increase in exopher production triggered by volatile metabolites as each mutant mirrored

wild-type induction (**Fig. 3a and c**), although we cannot exclude the possibility of redundant

or compensatory interactions. However, there is a notable discrepancy in the role of these

pathways when considering the non-volatile secretome-dependent increase in EV production.

Specifically, while the insulin pathway does not appear to be involved, the TGFβ, p38 MAPK,

and IPR pathways are essential for the non-volatile secretome-dependent increase in exopher

production (**Fig. 3b and c**). On the other hand, comparison of mutant versus wild-type animals

showed that *pmk-1* and *akt-1* mutants displayed an even greater increase in exophers than wild-

type upon volatile metabolites and non-volatile secretome exposure while *dbl-1* mutants

exhibited a blunted response to both types of compounds (**Supplementary Fig. 3a and b).**

Together, these results indicate that *C. elegans* relies on distinct immune signalling mechanisms

for EV production depending on the chemical nature of the stimulus: volatile *P. aeruginosa*

metabolites trigger exophergenesis that is independent from multiple pathways yet

quantitatively fine-tuned by insulin, TGFβ, and p38 MAPK signalling, whereas non-volatile

secretome-driven upregulation selectively requires TGFβ, p38 MAPK, and IPR pathways.

Tripeptide Ile-Pro-Pro present in *P. aeruginosa* supernatant upregulates exopher

production

During the course of our study, we observed a sudden decline in the reliability of our non-

volatile secretome-based exopher upregulation assay. Troubleshooting traced this inconsistency

to a switch in the membrane filters used for supernatant clarification. We confirmed this by

testing five different filter types (two previously used and three additional ones) and found that

only two permitted robust exopher induction (**Fig. 4a**). We leveraged this unexpected filter

sensitivity as a bioassay to pinpoint the active component(s) in *P. aeruginosa* supernatant (**Fig.**

4b). Using liquid chromatography–mass spectrometry, we compared the metabolite profiles of

exopher-inducing versus non-inducing filtrates from the same bacteria liquid culture and

identified 65 statistically significant, differentially abundant compounds (**Supplementary Data**

1). Interestingly, the active supernatant was enriched in several small peptides, dipeptides,

tripeptides, and tetrapeptides, all of which contained at least one proline residue. Moreover,

inspired by previous work demonstrating that N,N-dimethyltryptophan–containing dipeptide

can modulate *C. elegans* physiology, we selected Ile-Pro-Pro (**Fig. 4c**), a tripeptide known to

be produced by lactic acid bacteria[55], for functional testing.

When synthetic Ile-Pro-Pro was added to standard NGM medium, exopher production

increased in a dose-dependent manner (**Fig. 4d**). At 50 µM, we observed a 17% rise over vehicle

control, rising to 29% increase at 100 µM. These results identify the tripeptide Ile-Pro-Pro as a

specific, pathogen-associated molecule capable of inducing exopher production in the host.

While likely acting in concert with other secreted factors, Ile-Pro-Pro represents a key

molecular signal through which *C. elegans* can sense the metabolic state of surrounding

bacteria.

Multiple types of sensory neurons regulate exopher production in response to pathogen

secretome

Olfactory neurons in *C. elegans* detect numerous environmental signals that influence various

aspects of the worm's life, including behaviour, physiology, and reproduction[36]. Additionally,

our recent findings have shown that olfactory neurons regulate pheromone-dependent increases

in EV production[22]. To identify the sensory neurons responsible for regulating the pathogen

non-volatile secretome-dependent and volatile metabolites-dependent increases in EV

production, we utilized nematodes with genetic ablations of sensory neurons known to mediate

pathogen recognition[37–39] and in exopher regulation[22]. By systematically analysing the effects

of these ablations[40–44] on exopher production, we discovered that the removal of either ADL,

ASK, or AWC neurons completely abolishes both non-volatile secretome-dependent and

volatile metabolites-dependent upregulation of exopher production (**Fig. 5a-c**). Conversely, the

removal of AQR/PQR/URX, previously shown to be responsible for exophergenesis

regulation[22], had no effect in either assay (**Fig. 5a-c**). Interestingly, we observed discrepancies

among ASI, ASH, and AWB neurons. While ASI, ASH, and AWB neurons were crucial for

the non-volatile secretome-induced increase they do not play a role in the volatile metabolites-

dependent increase in EV production (**Fig. 5a-c**). Notably, ablation of ADL or ASK also

reduced baseline exopher levels (**Fig. 5a and c**), consistent with our previous work showing

that these neurons contribute to exophergenesis regulation at the basal level[22]. Finally, when

comparing each ablation to wild type under non-volatile secretome exposure, all

sensory-neuron mutants except AQR/PQR/URX showed a significantly attenuated exopher

response, with ASK and ADL mutants exhibiting the largest reductions, whereas

AQR/PQR/URX-ablated worms produced more exophers than wild type (**Supplementary Fig.**

4b). For volatile metabolites, AQR/PQR/URX and ASI ablations likewise enhanced exopher

induction, AWB ablation had no measurable effect, and the response was reduced in all other

neuron-ablated strains (**Supplementary Fig. 4a**). Altogether, these findings show that multiple

types of sensory neurons regulate exopher production in response to pathogen-produced

compounds in a context-dependent manner, with ADL, ASK, and AWC neurons playing pivotal

roles.

A hierarchical neural circuit distinguishes pathogen threat proximity

To map the downstream interneuron circuit, we took a dual approach. A literature search for a

central integrator of the previously identified sensory neurons pointed to the RMG interneuron,

a known "hub-and-spoke" network center[45]. Separately, a search of the *C. elegans* connectome

for downstream partners highlighted the AIA and AIB interneurons[46]. Notably, AIA and AIB,

along with the AIB-specific neuropeptide receptor NPR-9, have previously been implicated in

regulating *C. elegans* immunity[47,48], making them prime candidates for mediating this

physiological response.

Our functional data confirmed the critical role of the RMG hub. Genetic ablation of RMG not

only reduced basal exophergenesis but also completely abolished the induced response to both

volatile and non-volatile pathogen cues (**Fig. 6a-c**, **Supplementary Fig. 5a and b**), establishing

it as an essential, central integrator.

Next, we examined the roles of the candidate processing interneurons. Ablation of AIA revealed

a striking, context-dependent function. Removal of AIA caused a modest, non-significant

reduction in basal exopher levels (**Fig. 6d**). Upon exposure to volatile metabolites,

AIA-deficient worms still showed a significant increase in exopher production (**Fig. 6e**), albeit

less pronounced than in wild type (**Supplementary Fig. 5a**), whereas non-volatile secretome

failed to induce any upregulation (**Fig. 6f**, **Supplementary Fig. 5b**). These results indicate that

AIA is essential for non-volatile secretome-driven exophergenesis and plays a modulatory role

in the volatile metabolites-mediated response (**Fig. 6k**).

Genetic ablation of AIB did not affect basal exopher production, whereas loss of the inhibitory

receptor NPR-9, a condition that is known to activate AIB[47], reduced baseline exopher levels

(**Fig. 6g**). Consistent with the results from the *npr-9* mutant, optogenetic activation of AIB for

4 hours via ReaChR[49] also significantly decreased exopher production (**Fig. 6h**). Upon exposing

worms to *P. aeruginosa* metabolites, the removal of the AIB neurons or NPR-9 receptor

significantly reduced the increased exophergenesis induced by pathogen volatile metabolites

(**Fig. 6i, Supplementary Fig. 5a**) and non-volatile secretome (**Fig. 6j, Supplementary Fig.**

5b), with a more pronounced effect for volatile metabolites exposure. These results suggest that

the AIB interneurons and NPR-9 receptor are essential for mediating exopher production in

response to pathogen signals (**Fig. 6k**).

In summary, our findings delineate a multi-layered circuit that processes threat-level

information (**Fig. 6k**). Increased exopher production due to volatile metabolites sensing relies

primarily on the ASK, ADL, and AWC sensory neurons, which signal through the AIB and

RMG interneurons, with AIA interneurons acting as a modulator (**Fig. 6l**). In contrast, increased

exopher production due to non-volatile secretome sensing engages an expanded circuit that

includes ASI, AWB, and ASH inputs, alongside ASK, ADL, and AWC, converging on the same

AIB and RMG core, but with stronger AIA involvement (**Fig. 6m**).

Identification and functional characterization of GPCRs mediating pathogen-induced

exophergenesis

G-protein coupled receptors (GPCRs) are essential mediators that enable cells to interact with

their environment by transmitting signals. In both vertebrates and in invertebrates, GPCRs play

a key role in coordinating immune responses[50,51]. In order to pinpoint the receptor(s) responsible

for binding pathogen-released molecule(s) that modulate exophergenesis, we analysed

published transcriptomics data from worms exposed to *P. aeruginosa* PA14 compared to worms

grown on *E. coli* OP50[52]. Among the most significantly upregulated GPCRs were *sri-36* and

sri-39, with fold changes of 480 and 79.5, respectively. According to a study by Vidal et al.[53],

these GPCRs are expressed in ASK and ASI cells, which we have shown to play an important

role in exopher upregulation (**Fig. 5a-b**). Additionally, our prior RNA sequencing, which

identified the GPCR *str-173* as a regulator of socially-mediated exopher production, also ranked

sri-36 and *sri-39* among the top candidates associated with exopher regulation, further

justifying their selection for validation[22]. To investigate the roles of *sri-36* and *sri-39* in exopher

regulation, we employed CRISPR/Cas9 technology to generate a mutant with a deletion

encompassing both receptor-encoding genes, as they are located only 1.9 kb apart, with one

gene situated between them (**Supplementary Fig. 6**).

Moreover, to identify receptors that regulate exophergenesis and may play a role in stimulating

exopher production (potentially through pathogen-released compounds), we conducted RNA

sequencing on age-synchronized wild-type worms that exhibited either many (more than 20) or

few (fewer than 2) exophers on adulthood day 2. A similar strategy, utilizing gene expression

heterogeneity within isogenic populations, was recently employed to study the processes that

generate lifespan variation[54]. The results of this RNA sequencing show that worms producing

many exophers differ significantly at the transcriptomic level from worms producing few

exophers (**Fig. 7a**). Specifically, 609 genes were upregulated and 665 genes were

downregulated (p adj < 0.05; -1 > log2 fold change > 1) (**Fig. 7b and Supplementary Data 2**).

Among the statistically significant GO terms, "cilium assembly" was upregulated in worms

producing many exophers (**Fig. 7d**). This finding aligns with our data showing that ciliated

neurons are crucial for regulating exophergenesis in muscles (as it was previously shown in

Szczepańska, Olek et al.[22]). Notably, three GPCRs; STR-173, SRR-6, and SRI-19, were

significantly altered in animals producing varying numbers of exophers (**Fig. 7c**). Importantly,

we had previously identified STR-173 as a GPCR regulating exophergenesis via the male

pheromone ascr#10, validating our approach[22]. Consequently, we created CRISPR/Cas9-

mediated mutants for the *srr-6* and *sri-19* genes for functional analysis (**Supplementary Fig.**

6).

Among the four GPCRs studied, the basal level of exophergenesis under standard conditions

remained unchanged for *sri-19* and *sri-36/39* mutants (**Fig. 8a and e**), while it was mildly

upregulated in *srr-6* mutant (**Fig. 8i**). Upon exposure to pathogen cues, a clear functional

divergence emerged. All receptor mutants exhibited a statistically significant increase in

exopher production following exposure to volatile metabolites (**Fig. 8b, f, j**). The responses of

sri-36/39 and *srr-6* mutants were comparable to wild type (**Supplementary Fig. 7c, e**), whereas

the *sri-19* mutants showed a significantly attenuated response (**Supplementary Fig. 7a**). In

stark contrast, the response to the non-volatile secretome was completely suppressed in all

tested mutants (*sri-19*, *sri-36/39*, and *srr-6*), indicating they are all essential for this pathway

(**Fig. 8c, g, k**, **Supplementary Fig. 7b, d, f**). Underscoring its central role, the *sri-19* mutant

also failed to upregulate exopher production upon exposure to live *P. aeruginosa* bacteria

(**Supplementary Fig. 7g**).

Expression analysis provided the crucial link between these molecular players and the

previously identified circuits. The wrmScarlet CRISPR/Cas9-mediated transcriptional knock-

in of the *sri-19* gene revealed its expression in ADL and PHA neurons (**Fig. 8d,**

Supplementary Fig. 7h, i). Furthermore, transcriptional reporters using fluorescent proteins

for the *sri-36* and *sri-39* genes indicated expression in anterior ciliated neurons (**Fig. 8h**), which,

based on available literature[53], correspond to ASI and ASK neurons. However, we must note

that we used a double deletion at the *sri-36/39* locus, the observed phenotype in exopher

production could be driven by loss of only one receptor, with the other making little or no

contribution. Nonetheless, the expression pattern of all three receptors is consistent with our

data, underscoring the critical role of ADL, ASK, and ASI neurons in pathogen-induced

exopher production and establishing its molecular mechanism. Additionally, the expression of

the *srr-6* receptor in neurons of the retrovesicular ganglion (RVG), alimentary tract, and

excretory system (**Fig. 8l**), as well as the expression of *sri-36* and *sri-39* in the alimentary tract

(**Fig. 8h**), suggest that one or more of these tissues may also play a role in regulating exopher

production in muscles. To test this, we drove *srr-6* expression specifically in the intestine. This

intestinal rescue not only restored the non-volatile secretome-dependent exopher response but

also reduced basal exopher levels, a result consistent with array-mediated overexpression,

demonstrating that SRR-6 functions in the intestine to regulate muscle exopher production

(**Supplementary Fig. 7j**) and revealing a novel gut-muscle signalling axis.

Offspring of animals exposed to pathogen volatile metabolites show improved survival

post-infection

We aimed to investigate the functional consequences of extracellular vesicles release when

worms detect the pathogen in their environment. Prompted by a recent study showing that

perception of *Pseudomonas vranovensis* volatiles confers intergenerational protection in *C.*

elegans[56], we examined both offspring and maternal outcomes following pre-exposure to *P.*

aeruginosa cues. Specifically, we assessed progeny survival and developmental rate during *P.*

aeruginosa infection following parental pre-exposure to volatile metabolites or non-volatile

secretomes, and evaluated how this preconditioning impacted hermaphrodite mothers' survival

upon subsequent *P. aeruginosa* infection (**Fig. 9a**). For these assays, P0 worms were exposed

to *P. aeruginosa* volatile or non-volatile secretomes for 48 hours beginning at the L4 stage. To

assess progeny outcomes, F1 embryos were collected at the 3-fold stage, transferred to *P.*

aeruginosa–seeded plates at 20°C, and scored after 48 hours for survival and developmental

progression (scored as younger than L4, or L4 and older). For maternal survival, P0 adults

(exposed to secretomes until day 2 of adulthood) were transferred to fresh *P. aeruginosa* plates

and monitored every few hours until death. Pre-exposure of P0 adults to *P. aeruginosa* volatiles

significantly increased F1 survival under infection (**Fig. 9b**) without altering their rate of

development to L4/adulthood (**Fig. 9c**). In contrast, P0 pre-exposure to non-volatile secretomes

had no effect on either progeny survival or developmental timing (**Fig. 9d and e**). For P0

mothers, volatile preconditioning produced a slight, non-significant decrease in survival on *P.*

aeruginosa (**Fig. 9f**), whereas non-volatile preconditioning had no measurable impact (**Fig. 9g**).

We next asked whether the *sri-19* receptor, essential for exopher induction by *P. aeruginosa*

and its secretomes, also influences progeny outcomes. Following P0 pre-exposure to *P.*

aeruginosa volatiles or secretomes, *sri-19* mutants displayed no differences in F1 survival or

developmental rate (**Fig. 9h-k**). This result indicates that the enhanced offspring resilience

conferred by parental exposure to *P. aeruginosa* volatiles depends on the SRI-19-mediated

pathway.

Elevated temperature of 25°C both enhances exopher production in *C. elegans*[22] and increases

P. aeruginosa virulence. Because wild worms encounter pathogens in a range of environmental

temperatures, we assessed how F1 survival and development are affected at 25°C. As shown in

Supplementary Fig. 8a–d, these high-temperature outcomes differ markedly from those at

20°C (**Fig. 9b–e**). Progeny of parents pre-exposed to *P. aeruginosa* volatiles developed faster

(**Supplementary Fig. 8b**) but showed unchanged survival (**Supplementary Fig. 8a**), whereas

progeny of parents exposed to *P. aeruginosa* non-volatile secretome suffered both reduced

survival and delayed development compared to unexposed controls (**Supplementary Fig. 8c**

and d). In P0 adults, pre-exposure to *P. aeruginosa* volatiles significantly decreased survival

upon subsequent *P. aeruginosa* infection (**Supplementary Fig. 8e**), while pre-exposure to non-

volatile secretome had no effect (**Supplementary Fig. 8f**). Notably, *sri-19* mutants,

accumulated fewer embryos in utero (**Supplementary Fig. 8g**), and are unable to upregulate

exophers in response to *P. aeruginosa*, but survived significantly longer during infection

(**Supplementary Fig. 8h**). Likewise, knockdown of the yolk receptor RME-2, which nearly

abolishes exopher production[18], and reduces embryo accumulation *in utero* (**Supplementary**

Fig. 8i) also conferred enhanced resistance to *P. aeruginosa* (**Supplementary Fig. 8j**).

Together, these data show that parental sensing of *P. aeruginosa* volatiles enhances offspring

resilience, albeit at a modest cost to maternal survival. Moreover, environmental temperature

critically shapes this trade-off, underscoring the context-dependent balance between maternal

fitness and progeny protection. Finally, SRI-19's role in this pathway directly links the

machinery of exopher biogenesis to survival outcomes, highlighting how EV production and

organismal fitness are mechanistically intertwined.

Discussion

Our findings support a unified model (**Fig. 10**), in which *C. elegans* employs adaptive

exophergenesis as a sophisticated response mechanism to pathogen-derived environmental

cues. We demonstrate that worms effectively discriminate pathogen cues based on their

chemical nature, volatile versus non-volatile. Volatile metabolites are small,

low-molecular-weight compounds with high vapour pressures, diffusing rapidly through air or

aqueous films to create broad, low-concentration plumes. In contrast, compounds present in

non-volatile secretome are larger, more polar molecules that remain close to their source,

forming steep, localized concentration peaks. We propose that distinguishing between these

two types of compounds enables worms to accurately infer pathogen proximity and

subsequently mount tailored physiological responses. This response is orchestrated by immune

signaling in a stimulus-specific manner: exopher production in response to volatile metabolites

is modulated by insulin, TGFβ, and p38 MAPK but proceeds independently of IPR, whereas

non-volatile secretomes require TGFβ, p38 MAPK, and IPR. The partial overlap of these

pathways[57,58] suggests coordinated or compensatory interactions that fine-tune exophergenesis

according to the nature and intensity of the pathogen cue. The sensory neurons ADL, ASK, and

AWC detect both cues, whereas ASI, ASH, and AWB neurons specialize in detecting non-

volatile compounds. Signal integration occurs downstream via RMG hub, AIA, and AIB

interneurons, with the AIB-specific GPCR NPR-9 functioning as one of the regulatory

components. Both neuronal and non-neuronal GPCRs (SRI-19, SRI-36/39, and SRR-6) are

essential for transducing the non-volatile secretomes signal into exopher release, and SRI-19 is

responsible for modulation of the response to volatile metabolites. Moreover, *P. aeruginosa*

released tripeptide Ile-Pro-Pro functions as an inducer of exopher production. Functionally, we

demonstrate that maternal exposure to volatile pathogen metabolites triggers an SRI-19-

dependent resource shift toward progeny, increasing offspring fitness while compromising

maternal survival under more severe conditions.

Our findings on pathogen-induced exophergenesis gain broader significance when considered

alongside our recent work on its regulation by social cues, revealing a unifying framework in

which *C. elegans* leverages the same physiological process, muscle-derived exopher

production, to navigate radically different environmental contexts. In both cases,

exophergenesis serves as a form of maternal resource allocation that supports offspring fitness,

yet the initiating stimuli represent opposite poles of the ecological spectrum: pathogenic threat

versus reproductive opportunity. The male-produced ascaroside pheromone ascr#10 enhances

maternal provisioning in anticipation of mating, whereas pathogen-derived metabolites provoke

a terminal investment in progeny under threat of infection. This functional reuse of a conserved

output module underscores the evolutionary versatility of the exopher pathway. A striking point

of convergence across these two systems are the ASK and ADL sensory neurons, which emerge

as master integrators capable of detecting important environmental information. For instance,

in the social context, ASK detects ascr#10 via the GPCR STR-173; in the context of pathogen

exposure, ASK, together with ADL, detects non-volatile pathogen secretomes likely via distinct

GPCRs such as SRI-36/39 and SRI-19. This dual role positions ASK and ADL as a key node

in the sensory network that interprets both reproductive cues and pathogen-derived danger

signals, illustrating the remarkable contextual plasticity of the sensory code. However, beyond

this shared component, the upstream sensory receptors and downstream processing circuits

diverge almost completely, demonstrating the evolution of parallel, specialised neural

architectures for environmental decision-making. The pathogen-responsive circuit engages a

distributed network comprising ASK, ADL, and AWC neurons, integrated through the central

RMG interneuron hub, and results in a costly physiological outcome, maternal sacrifice for

progeny survival, mediated through canonical immune pathways such as TGF-β, p38 MAPK,

and the IPR system. In contrast, the social cue circuit recruits the AQR, PQR, and URX

interneurons and is modulated by neuropeptides including FLP-8 and FLP-21. This divergence

reflects a broader principle of sensory neuroscience: *C. elegans* employs combinatorial coding

and distributed receptor repertoires to extract meaning from complex environmental

landscapes. Rather than relying on a single labelled-line logic, the nervous system interprets

unique activation patterns across multiple neurons, allowing for nuanced discrimination

between, for example, distant potential threats and imminent reproductive opportunities. The

evolution of dual sensory systems converging on exopher production, each with distinct inputs

and physiological consequences, illustrates how a single effector process can be co-opted to

serve adaptive roles across diverse ecological scenarios.

Our findings also underscore the dual, context-dependent nature of specific neurons and the

broader physiological trade-offs that shape systemic responses to environmental threats. A

striking example is the ASK neuron, which has been previously implicated in promoting longevity under pathogen-free conditions[59] but also in suppressing survival on pathogens[60]. In our study, ASK participates in a pathway that reduces maternal survival upon pathogen exposure. These highlights ASK as a master environmental sensor whose downstream effects vary dramatically depending on the ecological context. At the organismal level, this manifests as a classic evolutionary trade-off between maternal self-preservation and reproductive investment. The anticipatory production of exophers represents an adaptive "bet" on offspring success that becomes maladaptive when maternal viability is jeopardised by harsh environmental conditions. Importantly, the nature and cost of the exopher response is determined not only by the chemical nature of the stimulus (volatile vs. non-volatile) but also by the associated signalling architecture. Non-volatile cues require canonical immune modules, including TGF-β, p38 MAPK, and the IPR pathway, signalling axes that are deeply conserved across metazoans and underscore an immune-centric, high-cost defensive mobilisation. In stark contrast, volatile-triggered responses proceed largely independently of these immune effectors, instead reflecting a low-cost, anticipatory mode of readiness. This distinction raises the possibility that exophers may constitute distinct subtypes of EVs with unique molecular compositions and functions. The chance that quantitatively similar exophers could be functionally distinct opens new research avenues to isolate and characterize these vesicles in detail. By analysing their content in response to both volatile and non-volatile secretomes, we could uncover their specific roles in organismal health and pathogen defence. This principle echoes findings in mammalian extracellular vesicle (EV) biology, where cargo composition and functional outcome are determined by the upstream signalling context.

Interestingly, while volatile metabolites like 1-undecene have been shown to increase lifespan in *C. elegans* under non-pathogenic conditions[61] or to protect worms against a subsequent infection with *P. aeruginosa*[15], our results demonstrate that volatile-induced exopher

production can lead to decreased survival during infection. The differences in the type, timing, and context of exposure likely contribute to this contradiction. In previous studies[15,61], a specific compound and shorter exposure periods were used. In contrast, our study involved prolonged exposure to a mixture of volatile metabolites from pathogens, which may trigger different physiological responses. The involvement of multiple G protein-coupled receptors in exopher production indicates that exophergenesis is driven by a complex mixture of pathogen-derived signals, rather than a single metabolite, although some receptors may share common ligands. This complexity may allow the organism to fine-tune its response depending on the nature of the threat it perceives, enabling a more nuanced and context-dependent reaction to environmental challenges.

Previous studies have shown that increased exopher release can benefit the organism by removing protein aggregates[17], extending lifespan[20], or promoting faster offspring development[18]. However, when exopher release is elevated in response to pathogen exposure, we observe an opposite effect, leading to decreased survival upon infection. Intriguingly, our data reveal tissue-specific exopher responses: muscle-derived exophers are induced by both volatile and non-volatile cues, whereas neuronal exophers respond exclusively to volatile signals. The apparent paradox between muscle- and neuron-derived exophers may reflect distinct functional roles: neuronal exophers promote longevity via cell-autonomous proteostasis, while muscle exophers constitute a resource-intensive systemic strategy that prioritises offspring at the expense of maternal survival. Moreover, this raises the possibility that exophers, or possibly EV in general, may function under a hormetic response: while a moderate increase in EV production appears beneficial, an excessive increase may overwhelm the organism, negating any positive effects. Thus, while moderate exopher production may be beneficial, too much exopher release might impose a burden on the organism, potentially due to changes in the content of these vesicles under different environmental conditions.

Our findings suggest that pathogen-induced exophergenesis contributes to offspring fitness by

promoting non-genetic maternal investment strategies. One possibility is that exophers serve as

vehicles for protective cargos that are ultimately transferred to developing oocytes.

Alternatively, elevated exopher output may reflect a systemic reconfiguration of maternal

physiology that enhances oocyte provisioning through shared metabolic or stress-response

pathways. Addressing this will require not only profiling exopher content but also functional

testing, such as isolating exophers and injecting them into naïve mothers to assess whether they

confer fitness advantages to offspring in the absence of infection. Resolving this will yield

fundamental insights into the mechanisms of non-genetic, transgenerational immunity and into

the broader role of extracellular vesicles as adaptive mediators of host-pathogen interactions.

In conclusion, our study highlights the complexity of exopher production in response to

pathogen-derived signals. The ability of *C. elegans* to detect and respond to volatile and non-

volatile compounds in a context-dependent manner suggests that exophers and possibly other

EVs play a crucial role in mediating the organism's physiological response to environmental

stressors. Future research should focus on isolating and characterizing exophers produced under

different conditions to determine their specific molecular contents and functions.

Understanding how these vesicles contribute to both the short-term survival of the organism

and the long-term fitness of its offspring will provide deeper insights into the adaptive roles of

EVs in host-pathogen interactions as well as in animals' healthspan and lifespan.

Methods

Worm Maintenance and Strains

C. elegans strains were maintained at 20°C (unless otherwise indicated) on nematode growth

medium (NGM) plates seeded with *Escherichia coli* OP50 as a food source[62]. Only

hermaphrodites were used in the experiments. A comprehensive list of all chemicals and strains

used in the study is provided in Supplementary Table 1 and 2, respectively.

Pathogen procedures

Pathogen bacteria media plates preparation

Worms were exposed to either *Pseudomonas aeruginosa* PA14 or *Serratia marcescens* Db10

strains. Pathogen-seeded plates were prepared following standard protocols[63,64]. All the

experiments with the pathogenic bacteria were conducted in Biosafety level 2 laboratory.

Pseudomonas aeruginosa **PA14 "slow-killing" (SK) plates preparation**

The PA14 strain was revived from frozen stock by streaking onto a 10-cm LB agar plate and

incubating at 37°C for 12–18 hours. Plates were stored at 4°C and used within one week. A

single PA14 colony was inoculated into 5 mL of LB broth and incubated overnight (maximum

16 hours) at 37°C with shaking at 250 rpm. Following incubation, 10 μL of the overnight PA14

culture was spotted onto the center of 35 mm "slow killing" plates (SK plates), which were

incubated at 37°C for 24 hours, followed by 25°C for another 24 hours to maximize virulence.

Serratia marcescens Db10 plates preparation

For *S. marcescens Db10*, the strain was similarly streaked from frozen stock onto a 10-cm LB

agar plate, incubated at 37°C for 12–18 hours, and stored at 4°C for up to one week. A single

Db10 colony was inoculated into 5 mL of LB broth and cultured at 37°C with shaking for 8–

10 hours. To prepare NGM infection plates, 25 μL of the *S. marcescens* culture was seeded onto

35 mm NGM plates, which were incubated at 37°C overnight.

Short-time infection assay

Worms were age-synchronized by isolating 3-fold stage embryos (hand-picked under a

stereomicroscope). Upon reaching the first day of adulthood (AD1), they were transferred either

to control plates seeded with *E. coli* OP50 or to plates seeded with the pathogens PA14 or Db10.

Worms were exposed directly to the pathogens for 3 hours at 25°C, after which they were

transferred back to OP50 seeded plates. Exopher levels were assessed the following day, on the

second day of adulthood (AD2).

Pathogen non-volatile secretome assay

To collect non-volatile secretome, a single colony of PA14 was inoculated into 5 mL of LB

media and cultured overnight for a maximum of 16 hours at 37°C. After incubation, bacterial

cells were removed by passing the culture through a 0.22 μm filter, and the bacterial supernatant

was collected. Age-synchronized L4 stage worms (n = 30), grown on 60 mm NGM plates

seeded with OP50, were treated with 50 μL of the filtered bacterial supernatant, while control

worms received 50 μL of LB broth. Plates were sealed with parafilm, and exophers were scored

48 hours later, when the worms reached AD2.

Volatile metabolites chamber assay

3-fold stage embryos were placed on 35-mm NGM plates seeded with *E. coli* OP50, with 10

embryos per plate. Once the embryos developed into L4 stage worms, a chamber with volatile

metabolites was assembled by placing a 35 mm SK plate containing PA14 or a control 35 mm

NGM plate with OP50 on the bottom. The plate with synchronized worms was positioned on

top, and the two plates were sealed together with parafilm. Worms were incubated with volatile

metabolites of the bacteria for 48 hours. Exophers were evaluated on the second day of

adulthood.

Parental generation survival

The parental generation (P0) of *C. elegans* was synchronized by transferring 10 3-fold stage

embryos onto 35 mm nematode growth medium (NGM) plates seeded with *Escherichia coli*

OP50. When the worms reached the L4 larval stage, they were subjected to the "Volatile

Metabolites Chamber Assay" or "Pathogen non-volatile secretome assay" as previously

described. After 48 hours of exposure to the volatile or non-volatile secretomes, exopher

production was quantified in the P0 generation at AD2. The P0 worms were then moved to SK

plates seeded with PA14 and maintained at 20°C (Fig. 9f, g) or 25°C (Supplementary fig. 8e-

h). The use of 5-fluoro-2′-deoxyuridine (FUdR) was omitted from all plates. Survival of the P0

population was monitored at multiple time points using a touch-provoked movement assay.

Worms that did not respond to touch were recorded as deceased, removed from the killing

plates, and documented. Those that responded to touch were considered alive and were counted

accordingly. Survival data were collected at least twice daily until the death of the entire

population.

Offspring survival and development assay

Once the parental generation (P0) of worms reached the L4 larval stage, they were subjected to

the "Volatile Metabolites Chamber Assay" or "Pathogen non-volatile secretome assay". After

48 hours of secretome exposure, exopher production was quantified in the P0 generation on the

second day of adulthood (AD2).

To assess intergenerational effects, 3-fold stage embryos of the F1 generation, derived from P0

worms were collected and transferred to 35 mm SK plates seeded with PA14 bacteria (10–15

embryos per plate). The plates were incubated at 20°C (Fig. 9b-e, h-k) or 25°C (Supplementary

fig. 8a-d). Offspring survival and development were assessed after 48h by scoring the total

number of worms and determining their developmental stages.

Embryo accumulation assay

First, AD1 age-synchronized worms were placed on either a lawn of pathogenic bacteria (PA14

and Db10) or a control lawn of OP50 for 3 hours. After exposure, worms were transferred to

OP50 plates and cultured until they reached the AD2 stage. In the next step, hermaphrodites

were treated with 1.8% hypochlorite solution. Once the worms dissolved, the embryos retained

in the uterus were counted.

RNA Interference Assay

RNA interference (RNAi) in *C. elegans* was performed using the standard RNAi feeding

method[65]. Briefly, RNAi plates were prepared by supplementing NGM with 12.5 µg/mL

tetracycline, 100 µg/mL ampicillin, and 1 mM IPTG. These plates were seeded with *E. coli*

HT115 bacteria expressing double-stranded RNA (dsRNA) targeting the gene of interest.

Control worms were fed *E. coli* HT115 containing the empty vector L4440.

Age-synchronized 3-fold stage embryos were transferred onto the freshly prepared RNAi plates

and cultured until day 2 of adulthood, at which point exophers were quantified. The

developmental stage of the worms was confirmed at the L4 stage, and any individuals that were

developmentally advanced or delayed were excluded from the experiment.

Metabolomic analysis

Metabolite extraction was achieved by mixing 50 µL of the sample with 250 µL of ice-cold

extraction solution of methanol:acetonitrile:isopropanol (2:9:9, v/v/v). Samples were shaken

for 10 min at 1500 RPM and subsequently centrifuged for 10 min at 14,000 RPM at 4°C.

Supernatants were transferred to new polypropylene sample tubes and evaporated to dryness

under nitrogen stream at 50°C. Dry residue was reconstituted by the addition of 100 µL of 90%

acetonitrile, vortexed and centrifuged.

LC-MS analysis was performed using Acquity UPLC system (Waters) coupled with Orbitrap

High Resolution Mass Spectrometer (Thermo Scientific). Chromatographic separation was

achieved using ACQUITY UPLC BEH HILIC column (130Å, 1.7 µm, 2.1 mm × 100 mm).

Mobile phase A consisted of 0.1% formic acid in MQ-grade water and mobile phase B consisted

of 0.1% formic acid in LC-MS grade acetonitrile. The initial gradient conditions were 100% B

with mobile phase flow rate at 0.4 mL/min and was held for 1 min. The mobile phase

composition changed to 90% B next 6 min, then dropped to 50% B at 9 min and to 10% B at

10 min, kept for next 4 min. Column was re-equilibrated by changing the gradient back to initial

conditions. The total method run time was 20 min. The mass spectrometer operated in separate

positive (ESI+ and negative (ESI-) electrospray ionization. The tune parameters were as given:

sheath gas flow rate 55.0 [a.u.], auxiliary gas flow rate 15.0 [a.u.], sweep gas flow rate 4.0 [a.u.],

spray voltage in ESI+ was 4.0 and in ESI- was 3.5, S-lens RF 30.0 [a.u.]. capillary temperature

was set to 320°C and auxiliary gas temperature to 350°C. Data acquisition was proceeded in

Data Dependent Acquisition (DDA) mode, MS1 resolution set to 70,000, MS1 AGC Target

1.0×106, MS 1 Maximum Injection Time 90 ms, scan range 120 to 1600 m/z, and MS2

resolution was set to 35,000, MS2 AGC Target to 1.0×105, MS2 Maximum Injection Time 100

ms, TopN was 8, isolation window 2.0 m/z and normalized stepped collision energy of 20, 30

and 40 was used to collect fragmentation spectra.

Raw data were uploaded to Compound Discoverer 3.3.3 (Thermo Scientific). An in-house

protocol for metabolomic search and identification was employed. The workflow tree consisted

of: Input Files → Select Spectra → Align Retention Times → Detect Compounds → Merge

Features → Group Compounds → Fill Gaps → Normalize Areas → Mark Bacground.

Metabolomika Pathways, mzVault, mzCloud and ChemSpider databases were selected for

analytes identification and assignment.

Ile-Pro-Pro tripeptide assay

Worms were age synchronized from 3-fold stage embryos and cultured on 60 mm NGM plate

seeded with *E.coli* OP50 bacteria. A total of 50 µL Ile-Pro-Pro solution (containing 50µM or

100 µM) was applied to plates with worms on adulthood day 1. Plates were left open under the

hood to dry for 30 minutes and incubated for further 24 hours at 20°C. BWM exophers were

scored on day 2 of adulthood, after 24h exposure to tripeptide. Synthetic, concentrated stock

solution of Ile-Pro-Pro was stored in H2O at -80 °C. The stock was diluted to working solutions

with M9 buffer.

Transcriptome analysis

C. elegans RNA extractions, library preparations, and sequencing were performed by Azenta

US, Inc (South Plainfield, NJ, USA) as follows:

RNA extraction

Total RNA was extracted using Qiagen RNeasy Plus mini kit (Qiagen, Hilden, Germany)

following the manufacturer's instructions.

Library preparation with polyA selection and Illumina sequencing

RNA samples were quantified using the Qubit 2.0 Fluorometer (Life Technologies, Carlsbad,

CA, USA), and their integrity was assessed with the Agilent TapeStation 4200 (Agilent

Technologies, Palo Alto, CA, USA). RNA sequencing libraries were constructed using the

NEBNext Ultra II RNA Library Prep Kit for Illumina, following the manufacturer's protocol

(NEB, Ipswich, MA, USA). In brief, mRNA was enriched using Oligo(dT) beads and

subsequently fragmented at 94°C for 15 minutes. First and second-strand cDNA synthesis was

performed, followed by end repair and adenylation of the cDNA fragments at their 3' ends.

Universal adapters were then ligated to the cDNA fragments, and libraries were enriched

through a limited-cycle PCR with the addition of indexing sequences. The resulting sequencing

libraries were validated using the Agilent TapeStation and quantified with both the Qubit 2.0

Fluorometer (Invitrogen, Carlsbad, CA, USA) and quantitative PCR (KAPA Biosystems,

Wilmington, MA, USA). The libraries were multiplexed and loaded onto the Illumina NovaSeq

6000 platform according to the manufacturer's instructions. Sequencing was performed with a

2×150 paired-end (PE) configuration using v1.5 reagents. Image analysis and base calling were

carried out using the NovaSeq Control Software v1.7. The raw sequence data (.bcl files)

generated were converted into fastq files and de-multiplexed using the Illumina bcl2fastq

software (version 2.20), allowing for one mismatch in index sequence identification.

Sequencing data analysis

Following quality assessment of the raw sequencing data, reads were trimmed to remove

potential adapter sequences and low-quality nucleotides using Trimmomatic v0.36. The cleaned

reads were then aligned to the Caenorhabditis elegans reference genome from ENSEMBL using

the STAR aligner v2.5.2b. STAR is a splice-aware aligner that identifies splice junctions and

integrates them into the alignment process to enhance the accuracy of read mapping. This step

resulted in the generation of BAM files. Unique gene hit counts were calculated using the

featureCounts function from the Subread package v1.5.2, considering only reads uniquely

aligned to exon regions.

The gene hit counts were subsequently used for downstream differential expression analysis.

DESeq2 was employed to compare gene expression levels between different sample groups.

The Wald test was used to calculate p-values and log2 fold changes. Genes with an adjusted p-

value < 0.05 and an absolute log2 fold change > 1 were considered differentially expressed in

each comparison.

Scoring of Muscle Exophers and Fluorescence Microscopy

Exopher quantification was performed as described in Turek et al.[18] and Banasiak et al.[24] (Bio-

Protocol, detailed, step-by-step guide). In brief, exophers were scored using a Zeiss Axio

Zoom.V16 stereomicroscope equipped with filter sets 63 HE and 38 HE. Worms were age-

synchronized by isolating 3-fold stage embryos (hand-picked under a stereomicroscope) and

maintained until day 2 of adulthood. On the second day of adulthood, worms were observed on

NGM plates, and the number of exophers per freely moving worm was counted.

Fig. 1e-h experiments were performed using different experimental setup. Exophers were

scored using a Leica DMRBE microscope equipped with a 20× objective and RFP filter. Worms

were age-synchronized by isolating 3-fold stage embryos or L4 stage larvae and maintained

until day 2 of adulthood. During the infection period with PA14 or Db10 (or OP50 as a control),

worms were briefly incubated at 25°C for three hours. On day 2 of adulthood, animals were

immobilized on 2% agarose pads with a 50 mM NaCl drop (10–20 µl), and the number of

exophers per worm was scored.

Scoring of Neuronal Exophers

Neuronal exophers were scored using a Leica STELLARIS 8 FALCON confocal system with

a 40× oil immersion objective. For imaging, worms were immobilized on 5% agarose pads

(dissolved in M9), 10 µM tetramisole as an anesthetic.

Optogenetics assays

For optogenetic activation, both control and experimental 35 mm NGM plates seeded with *E.*

coli OP50 bacteria were covered with 0.2 µM all-trans retinal (ATR) and left to dry at 20°C

and darkness for 16 hours. Ten to twelve age-synchronized worms were picked per plate from

optogenetic strains expressing ReaChR in AIB at adulthood day 2. After a 3-hour incubation at

20°C and in darkness, BWM exophers were scored. Next, experimental plates were placed

under the source of white light (light intensity measured at 594nm = 0.02 mW/mm2) for 4

hours. Control plates were shielded from light. The muscle exophers were counted immediately

after illumination. Control and treated groups were randomized before the start of the

experiment.

Fluorescence Microscopy

Representative fluorescence images in Fig. 2b and 2f were captured using a Leica STELLARIS

8 FALCON confocal system, equipped with a 40× oil immersion objective; GFP and RFP

fluorescence were excited using 494 nm and 575 nm lasers, respectively. Representative

fluorescence images in Fig. 7d, 7h, 7l and Supplementary Fig. 7i were captured using a Zeiss

LSM800 laser-scanning confocal microscope equipped with a 40× oil immersion objective;

GFP and RFP fluorescence were excited using 488 nm and 561 nm lasers, respectively. For

imaging, worms were immobilized on 5% agarose pads, using either 10 μl of PolySciences 0.05

μm polystyrene microspheres or 25 μM tetramisole as an anesthetic.

Generation of *sri-19*, *sri-36/39*, and *srr-6* mutant strains

CRISPR/Cas9 genome editing was employed to generate the *sri-19* mutants (*sri-19(wwa15)*

and *sri-19(wwa16)*) following previously established protocols[66]. The crRNA sequence used to

target the *sri-19* locus was ATAATTGGTGGATATACAAATGG. Mutations were mapped to

the first exon, resulting in frame shifts that likely produce molecular null alleles.

To create *srr-6* mutants, a previously described STOP-IN strategy[67] was used, which introduces

an early stop codon within the first exon. The crRNA sequence used to target the *srr-6* locus

was ACTCCAAGTCCTGAAGTCGT, and the STOP-IN cassette sequence was

GGGAAGTTTGTCCAGAGCAGAGGTGACTAAGTGATAA, flanked by

AATCTTTTTCAACAATGATTACTCCAAGTCCTGAAGTCGT (left) and

GGGTGACGAGGTTTCGTGTCCATCTATTATTGAAATACCT (right).

For the deletion of both the *sri-36* and *sri-39* genes, the entire 5968 bp region encompassing

both genes (including the intercalated gene F33H12.7) was removed and replaced with a STOP-

IN cassette (GGGAAGTTTGTCCAGAGCAGAGGTGACTAAGTGATAA). The flanking

sequences were TGGTTACTTTTTATTTTTATTTTTC (left) and

AATCCTAATCTAATATGAATCAATT (right). The crRNA sequences used for this deletion

were ACAATGCTAGTTGTAGGTAC and ATTCATATTAGATTAGGATT.

All mutant strains were backcrossed twice with N2 wild-type worms to remove background

mutations and verified using Sanger sequencing.

Generation of *srr-6* transgenic rescue strains

Plasmids cloning

Rescue construct was generated by simultaneously inserting two fragments, promoter and gene

of interest, into the linearized pMT27 vector using the SLIC method (Sequence and Ligation

Independent Cloning). The pMT27 plasmid contains the *unc-54*-3'UTR and an *unc-119(+)*

rescuing sequence. Restriction sites used for plasmid linearization were HpaI and PciI).

Promoter region was amplified by PCR from plasmid templates. For the gene insert, cDNA

from *C. elegans* was used as a template.

The SLIC reaction (10 µl total volume) consisted of 100 ng of linearized vector and

promoter/gene inserts in a molar ratio approximately equimolar to the vector, ensuring ~30 bp

overlapping sequences between vector and inserts. The reaction mixture also contained

NEBuffer 2.1 (NEB) and 0.5 µl of T4 DNA polymerase (NEB). The mixture was incubated at

room temperature for 2 minutes, then placed on ice to stop the reaction, followed by bacterial

transformation using the MH1 cloning strain. Plasmid sequences were verified for correctness

using Nanopore sequencing.

Transgenic lines preparation

Transgenic strain was generated by germline transformation by injecting rescuing construct for

srr-6 gene (100 ng/µl of eft-3p::srr-6::unc-54-3'UTR and 10 ng/µl of pGLOW77(myo-

2p::mNeonGreen) from Addgene as co-injection marker) to *srr-6(wwa49)* mutant.

Data Analysis and Visualization

807 Data analysis was performed using Microsoft Excel and GraphPad Prism 9. Graphical

808 representations of the results were generated in GraphPad Prism 9.

Statistical Analysis

No prior statistical methods were applied to determine sample sizes. Worms were randomly

allocated to experimental groups, and all experiments were conducted in a blinded fashion for

the datasets presented in Fig. 2d, i; Fig. 3a, b; Fig. 4a, d; Fig. 5a; Fig. 6a, b, c, d, e, f, g, h, i, j;

Fig. 8a, b, c, e, f, g, i, j, k; Fig. 9b, c, d, e, f, g, h, i, j k; Supplementary Fig. 2a, d; Supplementary

Fig. 3a, b; Supplementary Fig. 4a, b; Supplementary Fig. 5a, b; Supplementary Fig. 7a, b, c, d,

e, f, g, j; Supplementary Fig. 8a, b, c, d, e, f, g, h. Because the data were not normally distributed,

we employed nonparametric statistical tests. For pairwise comparisons, two-tailed Mann–

Whitney tests were used, while Kruskal-Wallis tests followed by Dunn's multiple comparisons

test were applied for multi-group comparisons. A p-value of < 0.05 was considered statistically

significant.

Data availability statement

821 The authors declare that the main data supporting the findings of this study are available within

822 the article and its supplementary files. Source data are provided with this paper. All raw data

823 have been deposited at RepOD repository and can be accessed at

824 https://doi.org/10.18150/YBLA2I.

References

1. Buzas, E. I. The roles of extracellular vesicles in the immune system. *Nat. Rev. Immunol.* **23**, 236–250 (2023).

2. Robbins, P. D. & Morelli, A. E. Regulation of immune responses by extracellular vesicles. *Nat. Rev. Immunol.* **14**, 195–208 (2014).

3. Peregrino, E. S. *et al.* The Role of Bacterial Extracellular Vesicles in the Immune Response to Pathogens, and Therapeutic Opportunities. *Int. J. Mol. Sci.* **25**, 6210 (2024).

4. Berumen Sánchez, G., Bunn, K. E., Pua, H. H. & Rafat, M. Extracellular vesicles: mediators of intercellular communication in tissue injury and disease. *Cell Commun. Signal.* **19**, 104 (2021).

5. Naznin, H. A. *et al.* Systemic Resistance Induced by Volatile Organic Compounds Emitted by Plant Growth-Promoting Fungi in Arabidopsis thaliana. *PLoS One* **9**, e86882 (2014).

6. McEwan, D. L., Kirienko, N. V. & Ausubel, F. M. Host Translational Inhibition by Pseudomonas aeruginosa Exotoxin A Triggers an Immune Response in Caenorhabditis elegans. *Cell Host Microbe* **11**, 364–374 (2012).

7. Pradel, E. *et al.* Detection and avoidance of a natural product from the pathogenic bacterium Serratia marcescens by Caenorhabditis elegans. *Proc. Natl. Acad. Sci.* **104**, 2295–2300 (2007).

8. Marsh, E. K. & May, R. C. Caenorhabditis elegans, a Model Organism for Investigating Immunity. *Appl. Environ. Microbiol.* **78**, 2075–2081 (2012).

9. Irazoqui, J. E., Urbach, J. M. & Ausubel, F. M. Evolution of host innate defence: insights from Caenorhabditis elegans and primitive invertebrates. *Nat. Rev. Immunol.* **10**, 47–58 (2010).

10.   Martineau, C. N., Kirienko, N. V & Pujol, N. Innate immunity in C. elegans. in

Current topics in developmental biology vol. 144 309–351 (2021).

11.   Peterson, N. D. & Pukkila-Worley, R. Caenorhabditis elegans in high-throughput

screens for anti-infective compounds. *Curr. Opin. Immunol.* **54**, 59–65 (2018).

12.   Iwama, R. E. & Moran, Y. Origins and diversification of animal innate immune

responses against viral infections. *Nat. Ecol. Evol.* **7**, 182–193 (2023).

13.   Harding, B. W. & Ewbank, J. J. An integrated view of innate immune mechanisms in

C. elegans. *Biochem. Soc. Trans.* **49**, 2307–2317 (2021).

14.   Meisel, J. D., Panda, O., Mahanti, P., Schroeder, F. C. & Kim, D. H. Chemosensation

of bacterial secondary metabolites modulates neuroendocrine signaling and behavior

of C. elegans. *Cell* **159**, 267–280 (2014).

15.   Prakash, D. *et al.* 1-Undecene from Pseudomonas aeruginosa is an olfactory signal for

flight-or-fight response in Caenorhabditis elegans. *EMBO J.* **40**, e106938 (2021).

16.   Wang, J., Barr, M. M. & Wehman, A. M. Extracellular vesicles. *Genetics* **227**, (2024).

17.   Melentijevic, I. *et al.* C. elegans neurons jettison protein aggregates and mitochondria

under neurotoxic stress. *Nature* **542**, 367–371 (2017).

18.   Turek, M. *et al.* Muscle-derived exophers promote reproductive fitness. *EMBO Rep.*

22, e52071 (2021).

19.   Arnold, M. L. *et al.* Intermediate filaments associate with aggresome-like structures in

proteostressed C. elegans neurons and influence large vesicle extrusions as exophers.

Nat. Commun. **14**, 4450 (2023).

20.   Yang, Y. *et al.* Autophagy protein ATG-16.2 and its WD40 domain mediate the

beneficial effects of inhibiting early-acting autophagy genes in C. elegans neurons. *Nat.*

Aging **4**, 198–212 (2024).

21.   Cooper, J. F., Guasp, R. J., Arnold, M. L., Grant, B. D. & Driscoll, M. Stress increases

in exopher-mediated neuronal extrusion require lipid biosynthesis, FGF, and EGF

RAS/MAPK signaling. *Proc. Natl. Acad. Sci.* **118**, e2101410118 (2021).

22.   Szczepańska, A. *et al.* Pheromone-based communication influences the production of

somatic extracellular vesicles in C. elegans. *Nat. Commun.* **15**, 2715 (2024).

23.   Nicolás-Ávila, J. A. *et al.* A Network of Macrophages Supports Mitochondrial

Homeostasis in the Heart. *Cell* **183**, 94-109.e23 (2020).

24.   Banasiak, K., Turek, M. & Pokrzywa, W. Preparation of Caenorhabditis elegans for

Scoring of Muscle-derived Exophers. *Bio-protocol* **13**, (2023).

25.   Schorey, J. S., Cheng, Y., Singh, P. P. & Smith, V. L. Exosomes and other extracellular

vesicles in host–pathogen interactions. *EMBO Rep.* **16**, 24–43 (2015).

26.   Kim, Dennis H, Ewbank, J. J. Signaling in the innate immune response. *WormBook*

2018, 1–35 (2018).

27.   Murphy, C. T. Insulin/insulin-like growth factor signaling in C. elegans. *WormBook* 1–

43 (2013) doi:10.1895/wormbook.1.164.1.

28.   Gumienny, T. L. TGF-β signaling in C. elegans. *WormBook* 1–34 (2013)

doi:10.1895/wormbook.1.22.2.

29.   Kim, D. H. *et al.* A conserved p38 MAP kinase pathway in Caenorhabditis elegans

innate immunity. *Science* **297**, 623–626 (2002).

30.   Liberati, N. T. *et al.* Requirement for a conserved Toll/interleukin-1 resistance domain

protein in the Caenorhabditis elegans immune response. *Proc. Natl. Acad. Sci.* **101**,

6593–6598 (2004).

31.   Lažetić, V., Batachari, L. E., Russell, A. B. & Troemel, E. R. Similarities in the

induction of the intracellular pathogen response in Caenorhabditis elegans and the type

I interferon response in mammals. *BioEssays* **45**, 2300097 (2023).

32.   Balla, K. M. & Troemel, E. R. Caenorhabditis elegans as a model for intracellular

pathogen infection. *Cell. Microbiol.* **15**, 1313–1322 (2013).

33.   Wang, G. *et al.* Mechanical force of uterine occupation enables large vesicle extrusion

from proteostressed maternal neurons. *Elife* **13**, RP95443 (2024).

34.   Dishart, J. G. *et al.* Olfaction regulates peripheral mitophagy and mitochondrial

function. *Sci. Adv.* **10**, eadn0014 (2024).

35.   Peterson, N. D. *et al.* Non-canonical pattern recognition of a pathogen-derived

metabolite by a nuclear hormone receptor identifies virulent bacteria in C. elegans.

Immunity **56**, 768-782.e9 (2023).

36.   Ferkey, D. M., Sengupta, P. & L'Etoile, N. D. Chemosensory signal transduction in

Caenorhabditis elegans. *Genetics* **217**, iyab004 (2021).

37.   Filipowicz, A., Lalsiamthara, J. & Aballay, A. Dissection of a sensorimotor circuit

underlying pathogen aversion in C. elegans. *BMC Biol.* **20**, 229 (2022).

38.   Styer, K. L. *et al.* Innate Immunity in Caenorhabditis elegans Is Regulated by Neurons

Expressing NPR-1/GPCR. *Science (80-.).* **322**, 460–464 (2008).

39.   Foster, K. J. *et al.* Innate Immunity in the C. elegans Intestine Is Programmed by a

Neuronal Regulator of AWC Olfactory Neuron Development. *Cell Rep.* **31**, 107478

(2020).

40.   Ludewig, A. H. *et al.* Pheromone sensing regulates Caenorhabditis

elegans lifespan and stress resistance via the deacetylase SIR-2.1. *Proc.*

Natl. Acad. Sci. **110**, 5522 LP – 5527 (2013).

41.   Masoudi, N. *et al.* Unconventional function of an Achaete-Scute homolog as a terminal

selector of nociceptive neuron identity. *PLOS Biol.* **16**, e2004979 (2018).

42.   Beverly, M., Anbil, S. & Sengupta, P. Degeneracy and Neuromodulation among

Thermosensory Neurons Contribute to Robust Thermosensory Behaviors in

Caenorhabditis elegans. *J. Neurosci.* **31**, 11718–11727 (2011).

43.   Yoshida, K. *et al.* Odour concentration-dependent olfactory preference change in C.

927   elegans. *Nat. Commun.* **3**, 739 (2012).

928   44.   Chang, A. J., Chronis, N., Karow, D. S., Marletta, M. A. & Bargmann, C. I. A

929   Distributed Chemosensory Circuit for Oxygen Preference in C. elegans. *PLOS Biol.* **4**,

930   e274 (2006).

931   45.   Macosko, E. Z. *et al.* A hub-and-spoke circuit drives pheromone attraction and social

932   behaviour in C. elegans. *Nature* **458**, 1171–1175 (2009).

933   46.   White, J. G., Southgate, E., Thomson, J. N. & Brenner, S. The structure of the nervous

934   system of the nematode Caenorhabditis elegans. *Philos. Trans. R. Soc. London. B, Biol.*

935   *Sci.* **314**, 1–340 (1986).

936   47.   Yu, Y., Zhi, L., Wu, Q., Jing, L. & Wang, D. NPR-9 regulates the innate immune

937   response in Caenorhabditis elegans by antagonizing the activity of AIB interneurons.

938   *Cell. Mol. Immunol.* **15**, 27–37 (2018).

939   48.   Cao, X., Kajino-Sakamoto, R., Doss, A. & Aballay, A. Distinct Roles of Sensory

940   Neurons in Mediating Pathogen Avoidance and Neuropeptide-Dependent Immune

941   Regulation. *Cell Rep.* **21**, 1442–1451 (2017).

942   49.   Lin, J. Y., Knutsen, P. M., Muller, A., Kleinfeld, D. & Tsien, R. Y. ReaChR: a red-

943   shifted variant of channelrhodopsin enables deep transcranial optogenetic excitation.

944   *Nat. Neurosci.* **16**, 1499–1508 (2013).

945   50.   Reboul, J. & Ewbank, J. J. GPCRs in invertebrate innate immunity. *Biochem.*

946   *Pharmacol.* **114**, 82–87 (2016).

947   51.   Dubyak, G. R. GPCRs in innate and adaptive immune responses. in *GPCRs* (eds.

948   Jastrzebska, B. & Park, P. S.-H. B. T.-Gpcr.) 429–461 (Elsevier, 2020).

949   doi:10.1016/B978-0-12-816228-6.00021-0.

52.   Sellegounder, D. *et al.* Neuronal GPCR NPR-8 regulates C. elegans defense against

pathogen infection. *Sci. Adv.* **5**, eaaw4717 (2019).

53.   Vidal, B. *et al.* An atlas of Caenorhabditis elegans chemoreceptor expression. *PLOS*

Biol. **16**, e2004218 (2018).

54.   Eder, M. *et al.* Systematic mapping of organism-scale gene-regulatory networks in

aging using population asynchrony. *Cell* **187**, 3919-3935.e19 (2024).

55.   Pessione, E. & Cirrincione, S. Bioactive Molecules Released in Food by Lactic Acid

Bacteria: Encrypted Peptides and Biogenic Amines. *Front. Microbiol.* **7**, 876 (2016).

56.   Pender, C. L. *et al.* Perception of a pathogenic signature initiates intergenerational

protection. *Cell* **188**, 594-605.e10 (2025).

57.   Yamamoto, K. K. & Savage-Dunn, C. TGF-β pathways in aging and immunity: lessons

from Caenorhabditis elegans. *Front. Genet.* **14**, 1220068 (2023).

58.   Madhu, B., Lakdawala, M. F. & Gumienny, T. L. The DBL-1/TGF-β signaling

pathway tailors behavioral and molecular host responses to a variety of bacteria in

Caenorhabditis elegans. *Elife* **12**, e75831 (2023).

59.   Alcedo, J. & Kenyon, C. Regulation of C. elegans longevity by specific gustatory and

olfactory neurons. *Neuron* **41**, 45–55 (2004).

60.   Venkatesh, S. R., Gupta, A. & Singh, V. Amphid sensory neurons of

Caenorhabditis elegans orchestrate its survival from infection

with broad classes of pathogens. *Life Sci. Alliance* **6**, e202301949 (2023).

61.   De-Souza, E. A., Thompson, M. A. & Taylor, R. C. Olfactory chemosensation extends

lifespan through TGF-β signaling and UPR activation. *Nat. Aging* **3**, 938–947 (2023).

62.   Brenner, S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71–94 (1974).

63.   Foster, K. J., McEwan, D. L. & Pukkila-Worley, R. Measurements of Innate Immune

Function in C. elegans. in *Methods in molecular biology (Clifton, N.J.)* vol. 2144 145–

160 (2020).

64.   Powell, J. R. & Ausubel, F. M. Models of Caenorhabditis elegans Infection by

977   Bacterial and Fungal Pathogens. in *Innate Immunity* (eds. Ewbank, J. & Vivier, E.)

978   403–427 (Humana Press, 2008). doi:10.1007/978-1-59745-570-1_24.

979   65.   Kamath, R. Genome-wide RNAi screening in Caenorhabditis elegans. *Methods* **30**,

980   313–321 (2003).

981   66.   Dokshin, G. A., Ghanta, K. S., Piscopo, K. M. & Mello, C. C. Robust Genome Editing

982   with Short Single-Stranded and Long, Partially Single-Stranded DNA Donors in

983   Caenorhabditis elegans. *Genetics* **210**, 781–787 (2018).

984   67.   Wang, H., Park, H., Liu, J. & Sternberg, P. W. An Efficient Genome Editing Strategy

985   To Generate Putative Null Mutants in Caenorhabditis elegans Using CRISPR/Cas9. *G3*

986   *Genes|Genomes|Genetics* **8**, 3607–3616 (2018).

987   68.   Witvliet, D. *et al.* Connectomes across development reveal principles of brain

988   maturation. *Nature* **596**, 257–261 (2021).

989   69.   Taylor, S. R. *et al.* Molecular topography of an entire nervous system. *Cell* **184**, 4329-

990   4347.e23 (2021).

Acknowledgements

Some strains were provided by the CGC, which is funded by NIH Office of Research

Infrastructure Programs (P40 OD010440). We thank Monika Woźniak, Natalia Wasiak,

Małgorzata Śledź, Tomasz Strumiński for assistance with worms maintenance; Adam Kłosin

for discussions and comments on the manuscript. Work in the MT Laboratory was mainly

funded by a National Science Centre SONATA grant (2019/35/D/NZ3/04091 to MT) and

additionally supported by a National Science Centre SONATA BIS grant

(2021/42/E/NZ3/00358 to MT). Work in the WP Laboratory was funded by the Norwegian

Financial Mechanism 2014-2021 operated by the Polish National Science Centre, Poland

(project contract number 2019/34/H/NZ3/00691 to WP). Also supported by the French National

Research Agency ANR-22-CE13-0037-01 to NP.

1003

Author Contributions

Conceptualization: MT; Data curation: MT; Formal analysis: KK, AS, SV, RPS, MR, MT, WP,

NP, HB; Funding acquisition: MT, WP, NP; Investigation: KK, AS, SV, RPS, MR, MT;

Methodology: MT, NP, WP, KK, AS, SV, RPS, MR, HB; Project administration: MT;

Resources: MT, NP, MR, WP, HB; Supervision: MT; Validation: MT, WP, KK, AS, NP, SV,

RPS, MR; Visualization: KK, AS, SV, RPS, MR, MT; Writing – original draft: MT; Writing –

review & editing: MT, WP, KK, AS, NP, SV, RPS, HB, MR.

1011

Competing Interests

1013   The authors declare no competing interests.

Figures and Figure Legends

Figure 1. Production of *C. elegans* muscle exophers is upregulated during pathogen infection.

a   Experimental Setup: Synchronized AD1 worms were exposed to *Pseudomonas aeruginosa* (PA14) and *Serratia marcescens* (Db10), or to *Escherichia coli* (OP50) for 3 hours at 25°C, followed by transfer to OP50-seeded plates. Exopher production was assessed on the second day of adulthood (AD2). Created with BioRender.com.

b   Short time infection with *P. aeruginosa* PA14 or *S. marcescens* Db10 leads to increased exopher production (n = 60 worms, N = 2 independent experiments).

c   Investigated innate immunity pathways: Various innate immunity pathways were assessed for their roles in regulating exopher production.

d   Worms deficient in key immune response pathways exhibited changes in exopher levels (n = 90 worms, N = 3 independent experiments; n = 90-180 worms, N = 3-6 independent experiments).

e - g   Infection-induced exopher production is not dependent on the (e) insulin signalling, (f) p38 MAPK, or (g) TGF-β pathways (n = 37-60 worms, N = 2 independent experiments). The data shown in graphs e–h were collected together and share the same controls but are presented in separate panels to improve visualization and enhance clarity. Baseline differences between panels d and e – g arise from methodological variations which are detailed in the Materials and Methods section "Scoring of Muscle Exophers and Fluorescence Microscopy"

h   Exopher production in response to *S. marcescens* Db10 infection is fully dependent on the IPR pathway, while production in response to *P. aeruginosa* PA14 infection is partially dependent on this pathway (n = 51-58 worms, N = 2 independent experiments). The data shown in graphs e–h were collected together and share the same controls but are presented in separate panels to improve visualization and enhance clarity.

1040   i   Exposure to *S. marcescens* Db10, but not *P. aeruginosa* PA14, results in embryo

1041   accumulation in the uterus (n = 54-75 worms, N = 3-4 independent experiments).

1042   Data are presented as medians with interquartile ranges. Non-significant p-values ($p > 0.05$) are

1043   highlighted in pink, while significant p-values ($p < 0.05$) are shown in blue. Statistical

1044   significance was assessed using the Kruskal-Wallis test with Dunn's multiple comparisons test.

1045

Figure 2. The production of muscle exophers in *C. elegans* is regulated by both volatile and non-volatile metabolites from pathogens.

a   Experimental setup schematic: Synchronized L4-stage worms were exposed to PA14 volatile metabolites or OP50 for 48 hours, and exopher production was evaluated on the second day of adulthood (AD2). Created with BioRender.com.

b   *C. elegans* hermaphrodites increase exopher production upon PA14 volatile metabolites, shown with red (RPN-5::wrmScarlet proteasome subunit) and green (mitochondrial outer membrane::GFP) fluorescence. Arrows: white - exopher, blue - exophers with mitochondria; dashed line circles - coelomocytes. Scale bars are 20 µm.

c   Exposure to PA14 volatile metabolites significantly increases exopher levels in worms (n = 77-90 worms, N = 3 independent experiments).

d   Volatile metabolites exposure causes mild embryo accumulation in the uterus (n = 90 worms, N = 3 independent experiments).

e   Experimental setup schematic: L4-stage worms were exposed to 50 µL of filtered pathogen supernatant, while controls received 50 µL of LB broth. Exophers were scored after 48 hours, at AD2.

f   *C. elegans* hermaphrodites increase exopher production upon PA14 non-volatile secretome, shown with red (RPN-5::wrmScarlet proteasome subunit) and green (mitochondrial outer membrane::GFP) fluorescence. Arrows: white - exopher, blue - exophers with mitochondria. Scale bars are 20 µm.

g   Exposure to PA14 supernatant results in increased exopher production (n = 90 worms, N = 3 independent experiments).

h   PA14 supernatant exposure does not induce embryo accumulation (n = 67-70 worms, N = 3 independent experiments).

i   Exposure to PA14 volatile metabolites induces higher exopher production compared to

PA14 secretome. No additive effect is observed when worms are exposed to both volatile and

non-volatile secretomes (n = 90 worms, N = 3 independent experiments).

Data are presented as medians with interquartile ranges. Statistical analyses were performed

using the two-tailed Mann-Whitney test (c,d,g,h) and the Kruskal-Wallis test with Dunn's

multiple comparisons test (i), with non-significant p-values (p > 0.05) in pink and significant

p-values (p < 0.05) in blue.

1079

c

	genes of the pathway	PA14 volatile metabolites	PA14 supernatant
WT	-	p < 0.0001	p < 0.0001
p38 MAPK	*tir-1*	p < 0.0001	p = 0.4776
	pmk-1	p < 0.0001	p = 0.0503
Insulin Signaling	*akt-1*	p = 0.0059	p = 0.0203
	daf-16	p < 0.0001	p = 0.0174
TGF-β	*dbl-1*	p < 0.0001	p = 0.5061
IPR	*cul-6*	p = 0.0007	p = 0.4151

Figure 3. Exopher production in response to pathogen volatile metabolites and non-

volatile secretome is governed by distinct regulatory mechanisms.

a   Increased exopher production after exposure to PA14 volatile metabolites does not

depend on innate immunity pathways (n = 80-90 worms, N = 3 independent experiments).

b   Increased exopher production after exposure to PA14 non-volatile metabolites depends

on p38, TGF-β, and IPR pathways (n = 85-120 worms, N = 3-4 independent experiments).

c   Summary of experimental results: A table summarizing the experimental findings,

including p-values. Created with BioRender.com.

Data are presented as medians with interquartile ranges. Statistical analyses were performed

using the two-tailed Mann-Whitney test, with non-significant p-values (p > 0.05) in pink and

significant p-values (p < 0.05) in blue.

1093

1094

Figure 4. Untargeted metabolomics revealed Ile-Pro-Pro tripeptide role in exopher

upregulation.

a   Evaluation of the robustness of pathogen supernatant in exopher induction revealed that

filters 2 and 5 allowed efficient clarification while retaining strong exopher-inducing activity

(n = 60-150, N = 3-5 independent experiments).

b   Experimental setup schematic: Metabolite profiles of exopher-inducing (filter 2) and

non-inducing (filter 1) filtrates from *P. aeruginosa* PA14 liquid cultures were compared using

liquid chromatography–mass spectrometry. Created with BioRender.com.

c   The fragmentation mass spectra of Isoleucyl-prolyl-proline acquired from the certified

standard (below) and from the biological sample (above). Each significant signal is described

by assigning the respective structure and mass of the fragment. Some of the fragments were

generated as a result of neutral loss, others as [M+H]+ adducts.

d   The synthetic Ile-Pro-Pro tripeptide induces a dose-dependent increase in exopher

production in exposed animals (n = 80, N = 3 independent experiments).

Data are presented as median with interquartile range. Statistical analyses were performed using

the Kruskal-Wallis test with Dunn's multiple comparisons test; non-significant p values (p >

0.05) are in pink colour, significant p values (p < 0.05) are in blue colour.

a PA14 volatile metabolites 〰〰

b PA14 non-volatile secretome ◊

c

	WT	ASK(-)	ADL(-)	ASI(-)	AQR/PQR/URX(-)	AWB(-)	AWC(-)	ASH(-)	ASJ(-)
PA14 volatile metabolites	p < 0.0001	p = 0.9039	p = 0.5055	p < 0.0001	p = 0.0004	p = 0.0014	p = 0.3608	p < 0.0001	p < 0.0001
PA14 supernatant	p < 0.0001	p = 0.3163	p = 0.5272	p = 0.1678	p = 0.0095	p = 0.4886	p = 0.1583	p = 0.1579	p = 0.0034

1114

1115

Figure 5. Chemosensory neurons regulate exopher production in response to pathogen

metabolite exposure.

a   Exopher production upon exposure to PA14 volatile metabolites is dependent on ASK,

ADL, and AWC neurons (n = 80-110 worms, N = 3-4 independent experiments). The data are

presented in three separate graphs because three different exopher reporter lines, each

exhibiting varying levels of basal exopher production, were used. The usage of different

exopher reporter lines was due to crossing incompatibilities with specific neuronal ablation

mutants.

b   Regulation of exopher production upon PA14 supernatant exposure involves multiple

neurons, including ASK, ADL, ASI, AWB, AWC, and ASH (n = 90-120 worms, N = 3-4

independent experiments). The data are presented in three separate graphs because three

different exopher reporter lines, each exhibiting varying levels of basal exopher production,

were used. The usage of different exopher reporter lines was due to crossing incompatibilities

with specific neuronal ablation mutants.

c   Summary of experimental results: A table summarizing the experimental findings,

including p-values. Created with BioRender.com.

Data are presented as medians with interquartile ranges. Statistical analyses were performed

using the two-tailed Mann-Whitney test, with non-significant p-values ($p > 0.05$) in pink and

significant p-values ($p < 0.05$) in blue.

1135

Figure 6. AIA, RMG, AIB interneurons and AIB-specific NPR-9 receptor integrate

sensory responses to mediate exopher production.

a   Exopher production is reduced in worms with ablated RMG interneuron (n = 88-90

worms, N = 3 independent experiments).

b   RMG interneuron is required for exopher production triggered by pathogen volatile

metabolites (n = 78-88 worms, N = 3 independent experiments).

c   Exopher production induced by pathogen-derived non-volatile secretome requires the

RMG interneuron (n = 90 worms, N = 3 independent experiments).

d   Animals lacking the AIA interneuron produce a similar number of exophers as wild-

type worms (n = 90 worms, N = 3 independent experiments).

e   Increased exopher production after exposure to PA14 volatile metabolites partially

depends on AIA interneuron (n = 90 worms, N = 3 independent experiments).

f   Exopher production induced by PA14 non-volatile secretome is mediated by the AIA

interneuron (n = 90 worms, N = 3 independent experiments).

g   Exopher production in strain lacking the AIB interneuron is comparable to that in wild-

type worms and lower in worms lacking NPR-9 receptor (n = 90 worms, N = 3 independent

experiments).

h   ReaChR-mediated optogenetic activation of AIB interneuron decreases exopher

production (n = 150 worms, N = 5 independent experiments).

i   AIB interneuron and AIB-specific NPR-9 receptor control exopher release upon

pathogen volatile metabolites (n = 88-91 worms, N = 3 independent experiments).

j   AIB interneuron and the NPR-9 receptor are required for exopher production induced

by pathogen-derived non-volatile secretome. (n = 90 worms, N = 3 independent experiments).

k   Summary of experimental results: A table summarizing the experimental findings,

including p-values. Created with BioRender.com.

l　　　Neuronal circuit controlling the exopher release upon volatile PA14 metabolites

exposure (connetome data from White et al. 1986[46] and Witvliet et al. 2021[68]). Created with

wormwideweb.org

m　　　Neuronal circuit regulating exopher release upon exposure to PA14 non-volatile

metabolites (connetome data from White et al. 1986[46] and Witvliet et al. 2021[68]). Created with

wormwideweb.org

Data are presented as median with interquartile range (a-g, i-j) or mean with SEM (h). Statistical

analyses were performed using the Kruskal-Wallis test with Dunn's multiple comparisons test

(g) and the two-tailed Mann-Whitney test (a-f, h-j), with non-significant p-values ($p > 0.05$) in

pink and significant p-values ($p < 0.05$) in blue.

1173

1174

Figure 7. Comparative transcriptomic analysis of worms producing many and few exophers per individual.

a   Principal component analysis (PCA) comparing worms with high exopher production (>20 exophers) and low exopher production (<2 exophers).

b   Volcano plot showing differential gene expression between worms producing many exophers (>20 exophers) and few exophers (<2 exophers).

c   Table listing 7TM GPCR transcripts with significant changes in expression levels.

d   Gene Ontology (GO) term analysis plot based on the transcriptome results.

1183

Figure 8. Exopher production induced by PA14 non-volatile metabolites is regulated by multiple GPCRs.

a   Exopher production in worms lacking the SRI-19 receptor is comparable to that in wild-type worms (n = 90-180, N = 3-6 independent experiments).

b   Increased exopher production after exposure to PA14 volatile metabolites does not depend on SRI-19 receptor (n = 90-180, N = 3-6 independent experiments).

c   The SRI-19 receptor is required for exopher production induced by PA14 non-volatile metabolites (n = 90-180, N = 3-6 independent experiments).

d   The *sri-19* gene encoding 7TM receptor is expressed in ADL neurons and PHA tail neuron, based on position and scRNAseq data[69], consistent with observations in at least 20 animals from three replicates. Square bracket mark gut autofluorescence.

e   Exopher production in worms lacking the SRI-36 and SRI-39 receptors is comparable to that in wild-type worms (n = 120, N = 4 independent experiments).

f   Increased exopher production after exposure to PA14 volatile metabolites does not depend on SRI-36 and SRI-39 receptors (n = 90, N = 3 independent experiments).

g   The SRI-36 and SRI-39 receptors are required for exopher production induced by PA14 non-volatile metabolites (n = 90-120, N = 3-4 independent experiments).

h   The *sri-36* and *sri-39* genes encoding 7TM receptors are expressed in neurons (including ASK# and ASI#) and the alimentary tract (red arrow – pharynx, green arrow – intestine), consistent with observations in at least 20 animals from three replicates. #based on position and Vidal et al.[53].

i   Basal exopher levels in *srr-6* mutants is increased compared to wild-type worms (n = 90-180, N = 3-6 independent experiments).

j   Increased exopher production after exposure to PA14 volatile metabolites does not depend on SRR-6 receptor (n = 90-180, N = 3-6 independent experiments).

k The SRR-6 receptor is required for exopher production induced by PA14 non-volatile

metabolites (n = 90-180, N = 3-6 independent experiments).

l The *srr-6* gene encoding 7TM receptor is expressed in neurons of the retrovesicular

ganglion (encircled), pharynx neurons (black arrow), alimentary tract (green arrow – intestine),

and excretory system (white arrow), consistent with observations in at least 20 animals from

three replicates.

Data information: Scale bars are 20 μm. Data are presented as median with interquartile range.

Statistical analyses were performed using the Kruskal-Wallis test with Dunn's multiple

comparisons test (a, e, i) and the two-tailed Mann–Whitney test (b, c, f, g, j, k); non-significant

p values (p > 0.05) are in pink colour, significant p values (p < 0.05) are in blue colour.

1220

Figure 9. The SRI-19 receptor is essential for the increased progeny resilience observed

after maternal pre-exposure to *P. aeruginosa* volatile cues.

a   Experimental setup schematic: Age-synchronized L4-stage worms (P0 generation) were

exposed for 48 hours to either PA14 and OP50 volatile metabolites (top) or PA14 non-volatile

secretome and LB media (bottom). After this exposure, exopher production was quantified at

the AD2 stage, and embryos from the F1 generation were collected. The P0 worms were

subsequently transferred to Slow Killing (SK) plates seeded with PA14, where their survival

was monitored. F1 embryos were subsequently transferred to SK plates seeded with PA14,

where their survival and development was quantified after 48h post-infection. Created with

BioRender.com.

b   Offspring of hermaphrodites exposed to PA14 volatile metabolites have better survival

rate upon PA14 infection (n = 195, N = 5 independent experiments).

c   No significant differences were observed in the developmental rate of F1 progeny

between control animals and those whose parents were exposed to PA14 volatile metabolites

(n = 195, N = 5 independent experiments).

d   Exposure of parents to PA14 non-volatile secretome had no significant impact on F1

survival during subsequent PA14 infection (n = 150, N = 4 independent experiments).

e   Developmental progression of F1 progeny during PA14 infection was not significantly

affected by parental exposure to PA14 non-volatile secretome (n = 150, N = 4 independent

experiments).

f   Wild-type P0 worms exposed prior to infection to PA14-derived volatile metabolites

had similar survival rate comparing to the control group (n = 163-165, N = 6 independent

experiments).

1246 g  Wild-type P0 worms pre-exposed to PA14-derived secretome showed no significant

1247 difference in survival compared to the control group upon subsequent PA14 infection (n = 52,

1248 N = 2 independent experiments).

1249 h  F1 progeny of *sri-19* mutant worms pre-exposed to PA14-derived volatile metabolites

1250 showed no difference in survival upon PA14 infection compared to the control group (n = 135,

1251 N = 3 independent experiments).

1252 i  Parental exposure to PA14 volatile metabolites did not affect the developmental timing

1253 of F1 progeny in the *sri-19* mutant background (n = 135, N = 3 independent experiments).

1254 j  No significant difference in survival was observed in the F1 generation of *sri-19* mutants

1255 following parental exposure to PA14 non-volatile secretome (n = 150, N = 4 independent

1256 experiments).

1257 k  Developmental progression of F1 progeny was unchanged in *sri-19* mutants whose

1258 parents were exposed to PA14 non-volatile secretome (n = 150, N = 4 independent

1259 experiments).

1260 Data are presented as stacked bar plots (b-e, h-k) and Kaplan–Meier survival curves (f, g);

1261 Statistical analyses were performed using the Fisher's exact test (b-e, h-k) and long-rank

1262 (Mantel-Cox) test (f, g); non-significant p values ($p > 0.05$) are in pink colour, significant p

1263 values ($p < 0.05$) are in blue colour.

1264

Figure 10. Model of exopher regulation from muscles by pathogen non-volatile secretome

and volatile metabolites.

1267

Supplementary Figure 1. Worms deficient in key immune signalling pathways exhibit

altered exopher production in response to pathogenic bacterial exposure.

a   Exopher production in immune signaling mutants following exposure to PA14 (n = 37-

60 worms, N = 2 independent experiments).

b   Exopher production in immune signaling mutants following exposure to Db10 (n = 37-

60 worms, N = 2 independent experiments).

c   Treatment with FUdR significantly reduced exopher production in worms, regardless of

exposure to PA14 or Db10 (n = 85-90 worms, N = 3 independent experiments).

Data are presented as median with interquartile range. Statistical analyses were performed using

the Kruskal-Wallis test with Dunn's multiple comparisons test; non-significant p values (p >

0.05) are in pink colour, significant p values (p < 0.05) are in blue colour.

1281

Supplementary Figure 2. PA14 volatile and non-volatile secretomes exposure

differentially affects exopher production in neurons and muscles without altering feeding

behavior.

a   FUdR treatment significantly reduced muscular exopher production upon exposure to

PA14 volatile metabolites (n = 87-91 worms, N = 3 independent experiments).

b   Exposure to PA14 volatile metabolites does not affect worm feeding behavior (n = 30

worms, N = 3 independent experiments).

c   ALMR neurons show increased exopher production after exposure to PA14 volatile

metabolites (n = 142-147 worms, N = 5 independent experiments).

d   FUdR treatment significantly reduced exopher production upon exposure to PA14 non-

volatile secretome (n = 79-91 worms, N = 3 independent experiments).

e   Exposure to PA14 non-volatile secretome does not affect worm feeding behavior (n =

30 worms, N = 3 independent experiments).

f   PA14 non-volatile secretome has no effect on exopher release from ALMR neurons (n

= 200 worms, N = 5 independent experiments).

Data are presented as medians with interquartile ranges. Statistical analyses were performed

using the Kruskal-Wallis test with Dunn's multiple comparisons test (a, d), the two-tailed

Mann-Whitney test (b,e) and Fisher's exact test (c, f), with non-significant p-values (p > 0.05)

in pink and significant p-values (p < 0.05) in blue.

Supplementary Figure 3. Animals deficient in key immune signaling pathways exhibit

altered exopher production in response to pathogenic bacterial metabolites.

a   Exopher production in immune signaling mutants following exposure to PA14 volatile

metabolites (n = 80-177 worms, N = 3-6 independent experiments).

b   Immune signaling mutants show altered exopher production in response to PA14 non-

volatile secretome (n = 90-175 worms, N = 3-6 independent experiments).

Data are presented as median with interquartile range. Statistical analyses were performed using

the Kruskal-Wallis test with Dunn's multiple comparisons test; non-significant p values (p >

0.05) are in pink colour, significant p values (p < 0.05) are in blue colour.

1314

Supplementary Figure 4. Neuronal ablation of sensory neurons alters exopher production

in response to PA14 metabolites.

a   Exopher production in neuronal ablation strains upon exposure to PA14 volatile

metabolites (n = 90-170 worms, N = 3-6 independent experiments).

b   Exopher production in neuronal ablation strains upon exposure to PA14 non-volatile

secretome (n = 90-210 worms, N = 3-7 independent experiments).

Data are presented as median with interquartile range. Statistical analyses were performed using

the Kruskal-Wallis test with Dunn's multiple comparisons test; non-significant p values (p >

0.05) are in pink colour, significant p values (p < 0.05) are in blue colour.

1327

Supplementary Figure 5. Exopher production in response to PA14 volatile and non-

volatile secretomes is decreased after removal of AIB, AIA and RMG interneurons and

NPR-9 receptor.

a   Animals lacking AIB, AIA, or RMG interneurons, as well as the NPR-9 receptor,

produce fewer exophers following exposure to PA14-derived volatile metabolites (n = 78-90

worms, N = 3 independent experiments).

b   Animals lacking AIA or RMG interneurons, as well as the NPR-9 receptor, produce

fewer exophers following exposure to PA14 non-volatile secretome, whereas AIB-ablated

animals show no significant change (n = 90 worms, N = 3 independent experiments).

Data are presented as median with interquartile range. Statistical analyses were performed using

the Kruskal-Wallis test with Dunn's multiple comparisons test and the two-tailed Mann–

Whitney test; non-significant p values (p > 0.05) are in pink colour, significant p values (p <

0.05) are in blue colour.

1343

Supplementary Figure 6. Diagrams of (a) *sri-19***, (b)** *srr-6***, (c)** *sri-36***, and** *sri-39* **genomic**

structures and genetic lesions (marked in pink) used in the study (related to Materials and

Methods).

1347

1348

1349

Supplementary Figure 7. Deletion of SRI-19, SRI-36, SRI-39, and SRR-6 receptors

modifies exopher release triggered by PA14-produced secretomes.

a-b   *sri-19* deletion alters exopher production in response to PA14 volatile and non-volatile

secretomes (n = 90-180 worms, N = 3-6 independent experiments).

c-d   *sri-36/39* deletion alters exopher production in response to PA14 non-volatile secretome

but not to volatile metabolites (n = 90-180 worms, N = 3-6 independent experiments).

e-f   *srr-6* deletion alters exopher production in response to PA14 non-volatile secretome but

not to volatile metabolites (n = 90-180 worms, N = 3-6 independent experiments).

g   SRI-19 receptor is required for exopher release during PA14 infection (n = 75-90

worms, N = 3 independent experiments).

h   Single-cell RNA-seq data from CeNGENApp[69] – *sri-19* is expressed mostly in ADL

sensory neurons. Expression also occurs in PHA and URY neurons. The circle diameter

represents the proportion of neurons in each cluster that express the sri-19 gene.

i   Expression of SRI-19 receptor co-localizes with *gap-15* GFP-based reporter line in

ADL neurons, consistent in at least 20 animals from three replicates. Scale bar is 10 μm.

j   Expression of *srr-6* under intestine-specific promoter rescues exopher production in *srr-*

6 loss-of-function mutants (n = 81-90 worms, N = 3 independent experiments).

Data are presented as median with interquartile range. Statistical analyses were performed using

the Kruskal-Wallis test with Dunn's multiple comparisons test (a-f) and the two-tailed Mann–

Whitney test (g,j); non-significant p values ($p > 0.05$) are in pink colour, significant p values (p

< 0.05) are in blue colour.

1371

Supplementary Figure 8. F1 development at 25°C is accelerated by P0 pre-exposure to volatile metabolites and impaired by non-volatile secretome.

a   F1 progeny of wild-type worms exposed to PA14 volatile metabolites show no significant difference in survival after 48 hours of PA14 infection compared to the control group (n = 140 worms, N = 3 independent experiments).

b   Parental exposure to PA14-derived volatile metabolites significantly improves developmental progression of the F1 generation after 48 hours on PA14 compared to the control group (n = 140 worms, N = 3 independent experiments).

c   Parental exposure to PA14 non-volatile secretome significantly reduces the survival of F1 progeny upon PA14 infection compared to the control group (n = 150 worms, N = 4 independent experiments).

d   F1 progeny of parents exposed to PA14 non-volatile secretome show reduced developmental progression during PA14 infection compared to the control group (n = 150 worms, N = 4 independent experiments).

e   Pre-exposure to PA14 volatile metabolites reduces survival of P0 worms during subsequent PA14 infection (n = 153 worms, N = 6 independent experiments).

f   Pre-exposure to PA14 non-volatile secretome does not significantly affect survival of P0 generation during PA14 infection (n = 79-80 worms, N = 3 independent experiments).

g   Loss of SRI-19 reduces embryo accumulation (n = 90 worms, N = 3 independent experiments).

h   The SRI-19 receptor that regulate exopher production upon exposure to PA14 non-volatile metabolites also regulates survival in response to PA14 infection (n = 102-108 worms, N = 4 independent experiments).

i   *rme-2* knockdown lowers embryo accumulation (n = 90 worms, N = 3 independent experiments).

j   Worms with *rme-2* gene knockdown, which produce low number of exophers,

demonstrated enhanced survival rates upon PA14 infection (n = 68-75 worms, N = 4

independent experiments).

Experiments conducted at 25°C. Data are presented as stacked bar plots (a-d) and Kaplan–

Meier survival curves (e, f, h, j); Statistical analyses were performed using the Fisher's exact

test (a-d), long-rank (Mantel-Cox) test (e, f, h, j), and the two-tailed Mann–Whitney test (g, i);

non-significant p values (p > 0.05) are in pink colour, significant p values (p < 0.05) are in

blue colour.