A histomorphological atlas of resected mesothelioma from 3446 whole-slide images discovered by self-supervised learning

Farzaneh Seyedshahi[1,2], Kai Rakovic[1,2,7,+], Nicolas Poulain[2,+], Adalberto Claudio Quiros[1,8], Ian R. Powley[2], Cathy Richards[3], Hussein Uraiby[3], Sonja Klebe[4], Apostolos Nakas[3], Claire Wilson[6], Marco Sereno[6], Leah Officer-Jones[2], Catherine Ficken[2], Ana Teodosio[5], Fiona Ballantyne[2], Daniel Murphy[1,2], Ke Yuan[1,2,8,*], and John Le Quesne[1,2,7,*]

[1]School of Cancer Sciences, University of Glasgow, Glasgow, Scotland, UK
[2]Cancer Research UK Scotland Institute, Glasgow, Scotland, UK
[3]University Hospitals of Leicester, Leicester, UK
[4]Flinders Health and Medical Research Institute, Adelaide, Australia
[5]Birmingham Tissue Analytics, University of Birmingham, Birmingham, UK
[6]University of Leicester, Leicester, UK
[7]Pathology Department, Queen Elizabeth University Hospital, NHS Greater Glasgow and Clyde, Glasgow, Scotland, UK
[8]School of Computing Science, University of Glasgow, Glasgow, Scotland, UK
[+]These authors contributed equally to this work
[*]ke.yuan@glasgow.ac.uk, john.lequesne@glasgow.ac.uk

1   Abstract

Mesothelioma is a highly lethal and poorly biologically understood disease which presents diagnostic challenges due to its morphological complexity. This study uses self-supervised AI (Artificial Intelligence) to map the histomorphological landscape of the disease. The resulting atlas consists of recurrent patterns identified from 3446 Hematoxylin and Eosin (H&E) stained images scanned from resected tumour slides. These patterns generate highly interpretable predictions, achieving state-of-the-art performance with 0.65 concordance index (c-index) for outcomes and 85% AUC in subtyping. Their clinical relevance is endorsed by comprehensive human pathological assessment. Furthermore, we characterise the molecular underpinnings of these diverse, meaningful, predictive patterns. Our approach both improves diagnosis and deepens our understanding of mesothelioma biology, highlighting the power of this self-learning method in clinical applications and scientific discovery.

2   Main

Mesothelioma is a highly lethal cancer almost always caused by asbestos exposure [1, 2]. Early detection, critical for effective treatment, remains challenging [3, 4]. Mesothelioma's biological diversity complicates histopathological diagnosis, as early malignancy can be difficult to distinguish from reactive changes. Diagnosing mesothelioma from H&E (Hematoxylin and Eosin) images is a subjective and time-intensive process even for skilled subspecialty histopathologists. Definitive diagnosis often remains elusive, even with immunohistochemistry or FISH (fluorescence in situ hybridisation). These diagnostic challenges are at least in part due to the difficulties in devising robust, manually applicable systems of morphological characterisation, in addition to the well-known issues of inter-pathologist agreement. The emergence of AI methods provides an opportunity to comprehensively describe the morphological complexity of mesothelioma to generate a quantitative visual dictionary of the disease.

Recent AI methods in mesothelioma primarily focus on tile-based or cell-based approaches, using supervised or weakly supervised learning. MesoNet [5] used Whole Slide Images (WSI) tiles to predict patient survival through a risk score based on malignant morphologies but lacked insight into the diversity of the tumour microenvironment. Cell-based methods like MesoGraph [6] and SpindleMesoNET [7] quantified malignancy through tumour cell shapes, especially spindle cells, but required extensive annotations and were computationally intensive for slide-level applications. While these approaches provide valuable insights, their findings are

restricted by the nature and quality of their human annotations. Recent self-supervised models, such as Hierarchical Image Pyramid Transformer (HIPT)[8], CTransPath [9], HistoSSLscaling[10], UNI[11], and Histomorphological Phenotype Learning(HPL)[12], have been developed for H&E WSI histopathological analysis. Uniquely among these approaches, HPL focusses upon identifying recurrent histomorphological patterns through clusters known as histomorphological phenotype clusters (HPCs). HPL leverages the Barlow Twins self-supervised framework [13], using ResNet for feature extraction from 224x224 WSI patches, followed by clustering of tile feature vectors via the Leiden algorithm [14]. Each HPC represents a unique morphological pattern that can be associated with specific molecular landscapes or used to predict patient outcomes and mesothelioma subtypes by quantifying HPC frequencies. (Further details in the Online Methods section.) Previously, HPL has been applied to lung cancer, revealing significant underlying patterns and giving impressive prognostic performance[12], but it has not been implemented in mesothelioma.

Self-supervised methods such as HPL depend upon accessing large volumes of training data, preferably from resected tumour material which offers large tissue areas with full morphological variance. This is especially challenging in mesothelioma, which is so often diagnosed from tiny biopsies and subsequently treated medically rather than surgically. So, in this study, we curated 3446 whole slide images of 485 resected mesothelioma cases to generate a uniquely powerful training resource called Leicester Archival Thoracic Tumour Investigation Cohort-Mesothelioma cohort (LATTICe-M), as shown with further details in Figure 1a. We then applied the HPL pipeline to our dataset to build a comprehensive atlas of mesothelioma H&E morphology. (Figure1b)

3 Results

3.1 Mapping the histomorphological landscape of mesothelioma

We have identified 47 recurrent histomorphological phenotype clusters (HPCs) (2a) based on morphological features encoded by self-learned neural networks. These HPCs are identified from 3,239,939 tiles extracted from 3446 images at 5x equivalent resolution. 42 of 47 HPCs are shared in more than 20% of cases, and none of them are case-specific. The most frequent pattern in our dataset is HPC 11("talc, pleurodesis, epithelioid infiltration") (Figure 2b)

A team of subspecialty expert pathologists from 3 centres, who had no access to the WSI images or labels (blinded assessment), examined every HPC to achieve consensus morphological annotations for each one, derived from their defining features: epithelioid vs spindled morphology, inflammation, necrosis, cellularity, desmoplasia, atypia, and cluster purity and each HPC was given a summary title. They evaluated inflammation levels in each HPC, categorising them as None-Sparse, Mild-Moderate, or Marked. Most HPCs were None-Sparse, but some displayed notable patterns. (Figure 2c). These annotations reveal areas of the UMAP containing multiple HPCs with broad similarities, such as spindled/collagenous HPCs, epithelioid tumour growth patterns, and lymphocytic infiltration, as well as peripheral and projecting clouds of morphologically highly distinct lung tissue and chest wall muscle tiles (Figure 2d). This grouping is supported by the general co-occurrence of HPCs within each slide in the Supplementary figures. These HPCs enable the detailed automated spatial annotation of any mesothelioma whole slide image, as illustrated in Figure 2e, highlighting two cases with highly divergent outcomes and morphologies. The first case, a sarcomatoid malignancy which resulted in death at 66 days, is highly morphologically diverse and contains abundant tiles in spindle cell-associated clusters, while the second is predominantly made up of a single epithelioid cluster.

3.2 Histomorphological Clusters Predict Mesothelioma Subtypes

The crucial histopathological distinction in mesothelioma subtyping lies between epithelioid and non-epithelioid (i.e. sarcomatoid/biphasic) variants. To classify this, we generated a numerical vector representing the percentage or frequency of each HPC for every WSI. This vector was transformed using the centred log-ratio (clr) transformation to enhance stability and interpretability, then feed into a logistic regression model for classification as either Sarcomatoid/Biphasic or Epithelioid. Tumours labelled as Biphasic and Sarcomatoid were combined into a single group, creating a binary classification task. 8 HPCs are significantly associated with the epithelioid subtype (HPCs 14, 39, 24, 25, 27, 40, 8, and 18 with log-rank test $p - value < 0.05$), containing epithelioid malignancy, predominantly characterised by tubular patterns and solid sheets of epithelioid tumour cell growth. Of the 9 HPCs linked to the non-epithelioid subtype, 3 HPCs (15, 16, and 22— mostly containing disorderly spindle cells) are unanimously classified as non-epithelioid malignancy by our pathologists as well. The other 6 HPCs (6, 7, 35, 37, 45, and 28— mostly solid epithelioid growth pattern) contain more diverse appearances, including epithelioid HPCs, pleural plaque and muscle. This might be due to the inclusion of biphasic cases and lethal epithelioid patterns in this group and also suggests a possible link between sarcomatoid growth and invasion into the chest wall. (Figure3a) Our logistic regression classifier achieved 87% 5-fold cross-validated AUC (Area Under the Curve) Score on the LATTICe-M dataset and 76.5% on The Cancer Genome Atlas (TCGA) mesothelioma dataset and robust across varied clustering configurations (Figure 3b). We visualised HPC compositions at the case level using a PCA plot, colour-coding cases by subtype at diagnosis. The transition from epithelioid cases to sarcomatoid cases through biphasic cases is clearly visible.(Figure 3c)

3.3 Histomorphological Clusters as Predictors of Patient Survival

We aggregated HPC frequencies across all samples per patient, summarising each case into a readily interpretable composition of morphologies. The 5-fold cross-validated c-index values for patient prognosis outcomes were 0.67 and 0.65 for the training and test LATTICe-M primary datasets, respectively, and 0.65 for the fully unseen TCGA cohort as an external dataset. The addition of clinical information, including mesothelioma subtype, TNM stage, and age, only modestly improved the ability of the algorithm to predict outcomes, yielding an increment in C-index of 0.01. We further verified that our approach is robust across other clustering configurations. Compared to similar research on mesothelioma outcome prediction using WSIs, such as MesoNet [5], our model achieves at least a comparable c-index score for the same additional dataset (TCGA). While MesoNet reported a score of 0.656 for TCGA, we matched this performance, with c-index scores ranging from 0.64 to 0.7 across different folds, however prioritising model interpretability through our morphology-based HPCs, also using a fully self-supervised pipeline.

We identified HPCs 10 (Log Hazard Ratio=-0.089, p-value=0.001) and 27 (Log Hazard Ratio=-0.062, p-value=0.008) ("epithelioid nests in bland stroma" and "dense lymphocytes") as positive survival factors, while HPCs 15 (Log Hazard Ratio=0.052, p-value=0.026) and 22 (Log Hazard Ratio=0.042, p-value=0.016) ("disorderly spindle cells" and "transitional mesothelioma") emerge as strong predictors of poor outcome. (Figure4a) A comparison of tile-level UMAP plots colour-coded by hazard ratio reveals a high degree of similarity, further underscoring the strong links between sarcomatoid transformation and poor patient outcomes. The map highlights red areas (higher hazard ratio) like HPC 15 and 16 (Sarcomotiod HPCs) and blue areas (lower hazard ratios) like HPC 45, 27 and 5 (either non-tumourous tissue or lymphocyte HPCs) (Figure 4b).

Next, we categorised patients into high- and low-risk groups based on their calculated hazard ratios for 60 months. Kaplan-Meier plots were generated for LATTICe-M train ($p - value = 2.79e - 15$) and test ($p - value = 9.96e - 15$) datasets, as well as the TCGA-Meso additional cohort ($p - value = 1.38e - 3$), as shown in Figure 4c. The model achieves impressive separation, predicting outcomes with surprising power in this extremely poor-prognosis population who face all the complex hazards of radical surgery and impaired respiratory physiology alongside the biology of their tumour burden. Figure 4d shows a comparison between classical histological grading of epithelioid pleural mesothelioma (as suggested in [15]) and our model, both applied to the TCGA mesothelioma just epithelioid sample cases. Our pipeline demonstrates superiority (p-value=0.025) against human grading for patient outcomes in this dataset. Figure 4e shows the SHAP (SHapley Additive exPlanations) [16] decision plots for our Cox model, comparing a high-risk sarcomatoid case (red) and a relatively low-risk epithelioid case (blue). The plot shows how the model assigns high or low-risk labels for these patients based on the abundance/scarcity of influential HPCs, such as the highly lethal HPC 15 ("disorderly spindle cells"), or the protective HPC 27 ("dense lymphocytes"), which both contribute to the calculated risk in these two cases.

We further demonstrate the ability of our model to predict patient outcomes within disease subtypes (epithelioid vs sarcomatoid/biphasic) groups. HPC frequencies were calculated, and survival was predicted separately for each group, identifying HPCs which underscored subtype-specific traits (Figure 5a). For epithelioid cases, HPC 10 ("epithelioid nests in bland stroma") and HPC 22 ("transitional mesothelioma") emerged as significant predictors of good and bad outcomes, respectively. HPC 10 is likely to represent relatively indolent well-differentiated 'classically' epithelioid disease. Interestingly, HPC 22, which is very enriched for the appearances of transitional mesothelioma, is not uncommon in cases diagnostically subtyped as epithelioid and is strongly predictive of poor outcomes in this group. This supports the view that transitional appearances signal early stages of transition to sarcomatoid growth [17], and its presence in this group highlights the difficulty in human identification of this pattern[18]. For biphasic/sarcomatoid cases, HPC 23 ("bland spindle cells and collagen") predicts poor outcome, perhaps identifying areas of cytologically bland desmoplastic differentiation, while the good prognostic association of HPC 27 ("dense lymphocytes") suggests towards a particular role for the immune system in sarcomatoid disease. We also show example tiles for specific HPC groups of interest based on pathologist annotations, including inflamed clusters, classical desmoplastic appearances, and necrosis (Figure5c).

3.4 Molecular underpinnings of HPCs

We next investigated the biological underpinnings of HPC morphology to further explain our model's predictive capabilities in mesothelioma prognosis and subtyping. This was achieved by quantifying associations between gene expression signatures and HPC composition in the TCGA Mesothelioma RNASeq dataset [19].

3.4.1 Ki67 Expression and MCPCounter Immune Features

We used the MCPcounter algorithm to estimate cell types, including fibroblasts, endothelial cells, T cells, B lineage cells, myeloid dendritic cells, NK cells, and CD8 T cells from RNASeq data. (Figure 6a) Expression of the proliferation marker Ki67 is also mapped, revealing especially high proliferation in spindle cell-enriched HPCs (HPCs 22, 16, 15, and 6). Critically, these HPCs are the most predictive of sarcomatoid/biphasic subtype (FIG XX). In contrast, HPCs defined by well-differentiated epithelioid disease and normal tissue (e.g., HPCs 17, 3, 10, and 19) show low proliferation.

Method	AUC-ROC	Avg. Precision	Sensitivity	Specificity
max-MIL	0.70 ± 0.01	0.54 ± 0.12	0.54 ± 0.07	73 ± 0.09
naive-MIL	0.84 ± 0.05	0.72 ± 0.11	0.72 ± 0.08	0.71 ± 0.1
PINS	0.85 ± 0.05	0.80 ± 0.07	0.82 ± 0.1	0.71 ± 0.13
CLAM	0.85 ± 0.07	0.74 ± 0.11	0.75 ± 0.11	0.77 ± 0.02
MesoGraph	0.90 ± 0.007	0.86 ± 0.02	0.88 ± 0.015	0.72 ± 0.01
HPL-MIL (Ours)	**0.93 ± 0.12**	**0.92 ± 0.11**	**0.94 ± 0.09**	**0.86 ± 0.19**

Table 1: AUC-ROC, Average Precision, Sensitivity, and Specificity scores for the HPL model performance using MIL approach to generate unique risk/sarcomatoidness scores. The St. George's Hospital TMA core dataset was used for benchmarking without any prior training, as reported in [6], through internal cross-validation, demonstrating the robustness of tile vector representations from our self-supervised network trained on mesothelioma WSIs. Additionally, we report HPL-MIL performance on our primary dataset (LATTICe-M) for comparison over different sets, using external (unseen) cross-validation for the subtyping task.

Fibroblast signatures are strongly pronounced in multiple clusters which either significantly determine sarcomatoid disease or contain fibroblastic/collagenous/stroma-rich morphology indicative of fibroblast-like mesenchymal dedifferentiation of mesothelioma cells. Fibroblast signatures are minimal in HPCs representing papillary or micropapillary epithelioid mesothelioma, solid pattern disease, lung tissue, and large-vessel-rich HPCs consistent with more specialised epithelioid/tissue-specific phenotypes.

Lymphocyte-rich HPC 27 shows strong correlations with T cells, B lineage cells, and myeloid dendritic cells. Similarly, inflamed HPCs (HPCs 29, 1, 24, 27), identified by Hover-Net, indicate active immune environments linked to better prognosis and likely improved immunotherapy response. HPCs 1 (inflamed fat) and 27 (dense lymphocytes) are high in both B- and T-cell signatures and show strong inter-correlation, suggesting dense inflammation and tertiary lymphoid structure formation in chest wall fatty tissues, supporting previous observations that tertiary lymphoid structures are related to good outcome.[20]

3.4.2 Pathways and Hallmarks

KEGG pathway correlations across HPCs again show clear separation between sarcomatoid/biphasic and epithelioid subtypes (Figure 6b), with generally heightened mitogenic signalling pathway activity in a group of sarcomatoid and fibroblastic clusters associated with aggressive biology. In contrast, HPCs linked to epithelioid growth and normal tissues exhibit relative down-regulation.

An analysis of cancer hallmark pathways further identifies the most sarcomatoid HPCs as a group with strong positive links to multiple proliferation-associated pathways (Figure 6c), in addition to mitogenic signalling and multiple EMT-related pathways. Notably, the same group exhibits downregulation of oxidative phosphorylation components, indicating a metabolic shift towards hypoxia.

3.5 Validation on Tiny Tissue Fragments

To assess the generalisability of our self-supervised model trained on the LATTICe-M dataset, we benchmarked its performance on the St. George's Hospital TMA dataset from the MesoGraph study [6]. This external evaluation is significant for two reasons. First, no additional training was applied to the new dataset, so the results represent the pre-trained model's performance on a fully unseen wholly exterior cohort. Second, although the model was trained on WSIs, it maintained strong performance on tissue microarray cores. These fragments are not only tiny ($\approx$1 millimetre) but are selected to represent 'pure' tumour tissue. In contrast, our model was trained on large diagnostic WSIs including background tissues, and used unsupervised clustering to filter artefacts.

As the image size in TMA cores is insufficient to support the previous frequency-based method, we employed multiple instance learning (MIL) to predict mesothelioma subtypes by summarising information across tiles from a core or a biopsy. We called this method HPL-MIL and benchmarked HPL-MIL against state-of-the-art methods, max-MIL and naive-MIL (patch-based MILmethods), PINS [21], CLAM[22], MesoGraph, on the 235 cores from St. George's Hospital TMA cohort [6] (Table 1). HPL-MIL achieved higher AUC, Average Precision, Sensitivity and specificity scores across all the methods without pre-training on the cohort tissues.

4 Discussion

In this study, we trained our self-supervised HPL pipeline on the LATTICe-M cohort, which we believe to be the largest single mesothelioma digital image dataset in existence. We achieved state-of-the-art accuracy in two key clinically important tasks: a C-index of 0.65 in survival prediction across a 5-fold cross-validation and 85% accuracy in subtype classification (epithelioid vs sarcomatoid/biphasic) and our method outperformed human grading for epithelioid cases. Furthermore, we identified survival-linked histomorphological patterns within each subtype, emphasising the interpretability of self-supervised methods and identifying recurrent morphologies worthy of future study. Quantitative visual maps of HPCs (Figure 2e) and SHAP decision plots (Figure 4e) offer clinical utility for understanding AI diagnoses and future selection of therapies.

Our approach eliminates the need for retraining on new mesothelioma datasets, addressing a key computational challenge in self-supervised models, as proven by its efficacy across three independent cohorts. It effectively extracts relevant morphological patterns from small TMA cores (e.g., St. George's dataset) and WSIs of varied origins and quality (e.g., TCGA and LATTICe-M), enabling real-time clinical decision-making without extensive preprocessing. This suggests the possibility of a robust diagnostic tool for both resection material and mesothelioma biopsies, which remain a major diagnostic challenge.

Our model's linkage of the immune microenvironment to patient survival suggests a potential for predicting responses to immunotherapy. Furthermore, sequence data contextualising histomorphological clusters strengthens connections between immune features, molecular pathways, and survival, offering valuable molecular insights into the biology of the disease.

The model also generates hypotheses for future investigation. We hypothesised clinicopathologically significant associations between low proliferation and hallmark pathways underpinning the improved survival observed in epithelioid cases. Additionally, subtle transitional morphologies in epithelioid cases appear to have significant prognostic value in our survival analysis, suggesting more research on this association. These findings could aid in identifying epithelioid subgroups potentially responsive to immunotherapy (notably effective in sarcomatoid cases at the moment [23]), demonstrating the utility of our approach in guiding targeted treatment strategies.

Sarcomatoid mesothelioma, known to proliferate rapidly under hypoxic conditions [24], corroborates our finding of a negative correlation between sarcomatoid HPCs and the oxidative phosphorylation (OXPHOS) hallmark (Figure 6c). We also observed a link between sarcomatoid growth and proliferation, with the fibroblast-like nature of sarcomatoid disease suggesting co-option of a physiological EMT pathway. Furthermore, inflammation of fat at the invasive front was associated with favourable outcomes.

5   Online Methods

Histomorphological Phenotype Learning (HPL) is a tool developed to detect and categorise visible patterns within large collections of whole-slide images, which are digital representations of tissue samples. HPL employs an automated, self-supervised deep learning approach, eliminating the need for expert pathologists to prelabel or manually define histomorphological phenotype patterns. Once these patterns are identified, new whole-slide images can be introduced to the trained model and classified according to the pre-established patterns. This feature allows pathologists to quantify specific patterns in new patient samples precisely. The clustering of each whole-slide image into meaningful histomorphological patterns follows several sequential steps, which are described below (Figure 1b)

- **Whole-Slide Images Pre-Processing:** In this first step, whole-slide images are segmented into non-overlapping 224×224-pixel tiles at 5X magnification, which corresponds to a pixel size of approximately 1.8 micrometres. Tiles that do not contain at least 60% tissue coverage are filtered out to maintain relevance. Consistent pixel size and magnification are ensured during the tile processing phase to guarantee uniformity in the resulting tiles. The tiling code used for this process is accessible on https://github.com/ncoudray/DeepPATH [25] for further details.

- **Feature extraction:** HPL employs a self-supervised learning technique known as Barlow Twins [13], which matches or even exceeds the performance of other self-supervised methods. What sets Barlow Twins apart is its efficiency, avoiding the need for large batches, as required by SimCLR[26], or complex architectural changes seen in models like BYOL[27] and Momentum Contrast[28]. Despite its simplicity, Barlow Twins consistently delivers state-of-the-art results, making it an ideal choice for HPL.

  One key feature of HPL is its ability to maintain consistent image representations, even with slight colour or zoom level variations. This capability ensures that differences in image scanning or processing across datasets do not affect the results. The aim is to capture diverse visual patterns in tissue samples and represent them as feature vectors, capturing distinct characteristics like texture. Each 224×224-pixel tile is converted into a vector representation, denoted as $\{z \in R^D; D = 128\}$. During training, the model is optimised to produce consistent outputs for twin inputs, ensuring robustness in vector representation.

- **Clustering:** After generating the vector representations, we employed the Leiden community detection algorithm [14] to cluster tiles or vector representations with similar histomorphological features. Since neighbouring vector representations in high-dimensional space exhibit similarity, this method effectively groups the tiles based on shared morphological patterns captured by their feature vector representations.

  We began with a subsample of 750,000 vectors and constructed a nearest-neighbour graph between the tiles. From this initial set of detected clusters, we assigned the remaining vector representations to these clusters (or graph nodes) based on their distance. The number of clusters identified depends on the chosen Leiden algorithm resolution. For our analysis, multiple resolutions were applied to capture varying levels of granularity. The resulting histomorphological phenotype clusters (HPCs) enabled the quantification of patients or whole-slide images (WSIs) based on these clusters, streamlining further analysis and simplifying the understanding of complex tissue patterns.

- **Preparing Compositional Vectors:** At this stage, using the identified HPCs, we can characterise the entire tissue or patient by quantifying the frequency of each Leiden cluster. To achieve this, each whole-slide image (WSI) is transformed into a compositional vector **A**, where the dimensionality is equal to the total number of histomorphological phenotype clusters (c). Each element within the vector represents the percentage of the tissue area attributed to a specific HPC. This approach quantifies the contribution of each HPC to the overall tissue composition, allowing for a detailed analysis of the histomorphological landscape within a patient sample.

$$A = \{a_0, a_1, a_2, \cdots, a_{c-1}\} \text{ s.t. } \sum_{i=0}^{c-1} a_i = 1 \text{ and } a_i \in [0,1]$$

For statistical compositional analysis and to prepare for the use of linear models, we apply the Centred Log-Ratio (clr) transformation [29] to our compositional vector **A**. This transformation maps the vector composition from the c-part Aitchison simplex into a c-dimensional Euclidean vector space, also known as real space. The clr transformation improves the handling of co-dependencies between variables, facilitating the use of linear models such as logistic regression and Cox survival models. Additionally, to address zero elements in the dataset, we use multiplicative replacement [30].

$$A \to U \text{ and } U = \left\{ x : \sum_{i=1}^{c} x = 0, \forall x \in \mathrm{R}^c \right\}$$

which comes to:

$$clr(A) = \left\{ log\frac{a_1}{g(A)}, log\frac{a_2}{g(A)}, ..., log\frac{a_c}{g(A)} \right\} \text{ where } g(A) = \left(\prod_{i=1}^{c} a_i \right)^{\frac{1}{c}}$$

- **Subtype Classification:** For the diagnostic task, we employed the compositional vectors derived from whole-slide images (WSI) and fed them into a logistic regression model. This approach is weakly supervised, utilising patient-level labels assigned by pathologists, where each patient label is applied to both the patient and their corresponding slides. We combined Sarcomatoid and Biphasic mesothelioma into a single class and compared it against the majority class, primarily consisting of Epithelioid samples. Also, to address the class imbalance in the primary dataset (1:3 ratio for Sarcomatoid/Biphasic to Epithelioid subtypes), we applied an undersampling strategy using the Edited Nearest Neighbour (ENN) technique [31]. This method reduced the majority class by removing redundant and noisy samples.

Ultimately, the logistic regression model used the compositional vectors of WSIs to classify mesothelioma subtypes based on the contributions of histomorphological phenotype clusters. In this approach, individual HPCs serve as distinct features for our logistic regression classifier, enabling us to rank the importance of each HPC and its role in predicting specific tumour subtypes within each sample.

$$Y = \frac{e^{b_0 + b_1 A}}{1 + e^{b_0 + b_1 A}}$$

Where b_0 is the bias or intercept term and b_1 is the coefficient for compositional vector(A)

To improve the accuracy of mesothelioma subtype prediction, we applied Multiple Instance Learning (MIL), a modern technique that combines tile-level vector representations from a self-supervised network. Using gated attention in MIL, we predicted the probability of each WSIs for a specific mesothelioma subtype [32], and named our model HPL-MIL.

While MIL is less interpretable, it enabled us to benchmark subtype classification scores against other methods using MIL, demonstrating the generalisability and scalability of the HPL pipeline. Despite its success in benchmarking, the remainder of the paper focuses on the more interpretable histomorphological clusters and their morphologies to better understand this biologically-complex cancer.

- **Survival Analysis:** In the clinical outcome aspect of our study, we created a compositional vector for each patient, reflecting the overall HPCs composition. We then used the Cox proportional hazards regression model [33] to analyse patient survival in relation to the HPC composition vector. Finally, Kaplan-Meier plots [34] were employed to visually distinguish between high-risk and low-risk patient groups within each dataset.

Additionally, we engaged three expert pathologists to independently and blindly annotate HPCs without access to patient clinical data or additional HPC details. Each HPC was classified into one of three categories: epithelioid tumour, spindle cells/extracellular

matrix, or non-tumour. The pathologists assessed each HPC's primary and secondary architectural features, HPC purity, inflammation, necrosis, nuclear atypia and biphasic components (in malignant groups). Also, they evaluated patterns such as desmoplastia and cellularity in spindle cell HPCs, as well as the tumour-stroma ratio and stromal cellularity in epithelioid HPCs. (**??**)

To confirm our findings through the HPL framework and gain further insights into mesothelioma, we undertook the following steps:

1. We used the deep learning model HoVer-Net [35] to segment lymphocytes in each tile within the HPCs and calculate the abundance of inflammatory cells in every HPC. While the tiles used in the HPL framework were at 5x magnification, the HoVer-Net model was trained on 20x tiles. To bridge this difference, we first applied HoVer-Net to 20x tiles, then combined 16 tiles (arranged in a 4x4 grid) to create 5x equivalents. This allowed us to map HoVer-Net's segmentation results, particularly the inflammation annotations, to specific tiles. Each tile was subsequently assigned to a corresponding HPC by calculating the average number of inflamed cells detected across the related tiles. Approximately 900 whole-slide images (WSIs) at 20x magnification were annotated using the HoVer-Net model for this analysis.

2. Three expert pathologists annotated the HPC morphologies and provided differentiation insights. Additionally, we correlated HPCs with immune features, hallmark pathways, and relevant biomarkers (Section 3.4). All WSIs from each patient were used to calculate the clr-transformed HPC compositional vectors. We then applied the single-sample Gene Set Enrichment Analysis (ssGSEA) to quantify pathway expression in both the Kyoto Encyclopedia of Genes and Genomes (KEGG) [36] and Molecular Signatures Database (MSigDB) [37] hallmark datasets. We also estimated immune cell subpopulation abundance using MCPcounter [38]. Ki67 RNA expression levels were also utilised across the entire sample set to assess cellular proliferation in each case, providing further insight into tumour growth activity within different morphological HPCs. Correlations between the clr-transformed HPC compositions and pathway expression levels were calculated, with only correlations having a p-value below 0.01 retained, ensuring statistical significance.

3. We benchmarked and compared the HPL network on WSIs against other state-of-the-art AI methods using a separate small dataset of tissue microarray (TMA) cores to demonstrate the robustness of our HPL approach (3.5).

5.1   Datasets

The primary dataset for this study, the Leicester Archival Thoracic Tumour Investigation Cohort-Mesothelioma (LATTICe-M) [39], includes 512 patients. These images are from patients diagnosed with plural mesothelioma, resected after surgery, and categorised into epithelioid (372 cases), sarcomatoid (107 cases), and biphasic (33 cases). The dataset predominantly features male patients (436 cases) over female patients (76 cases). Figure 1a presents additional clinical details. The WSIs were sectioned and stained with Hematoxylin and Eosin (H&E) at Leicester University Hospital, scanned at 10X, 20X, or 40X magnifications. After tiling and background removal, WSIs with fewer than 100 tiles were excluded, leaving 485 patients and 3446 WSIs for the pipeline and downstream analysis. To identify significant clinical factors in this cohort, we employed a Cox proportional hazards model and found that age, mesothelioma subtype, and TNM stage significantly contributed to survival prediction. (Figure 1a)

To validate our results, we used the publicly available Cancer Genome Atlas (TCGA)-mesothelioma cohort [19], an entirely differently-scanned dataset, still comprising WSIs but obtained from multiple centres. It includes 86 samples from 74 patients with both WSIs and RNAseq data available. This cohort was primarily used to discover links between HPCs and immune features, pathways, and hallmarks. All HPL pipeline steps were performed on the primary dataset (LATTICe-M), and evaluation scores were reported on the fully unseen additional TCGA dataset, without any further training.

Finally, we utilised the St. George's Hospital dataset, consisting of H&E-stained TMAs from tumour biopsies collected at St. George's Hospital, London. This dataset includes four TMA slides scanned at 20× magnification using a Hamamatsu Nanozoomer S360 scanner, comprising 235 cores labelled as epithelioid, biphasic, or sarcomatoid, as the only available clinical information. The dataset, introduced in the Mesograph study [6], was used for training and testing. We employed it to demonstrate the robustness and generalisability of our trained WSI model by benchmarking and comparing its performance on TMA cores against different methods reported in the study. (Section 3.5)

6   Code and Data Availability

The code for our study can be accessed from the HPL paper on this GitHub [12]. For clustering the vector representations, we utilised the Leiden algorithm [14] and the PAGA algorithm [40] from the ScanPy library [41]. For survival analysis, including Cox proportional hazards and Kaplan-Meier models, we relied on the Lifelines [42] and SciPy libraries [43]. To classify mesothelioma subtypes, we used logistic regression from the statsmodels library [44], with dataset balancing supported by the imbalanced-learn library [45]. Additional machine learning tasks were facilitated using scikit-learn [46]. For immune cell population analysis, we applied the single-sample Gene Set Enrichment Analysis (ssGSEA) method, as implemented in the GSVA R package [47], and

estimated immune cell abundance with the MCPcounter R package [38]. The LATTICe cohort was initially constructed under REC 14/EM/1159 (East Midlands REC), with ongoing resource management by the Greater Glasgow and Clyde Biorepository approved via an amendment granted by the Leicester South REC. Current use of the collection is managed under REC 16/WS/0207. The dataset will be made available upon request for individual academic research purposes.

7   Acknowledgements

The authors would like to express their gratitude to the CRUK Scotland Institute, the NHS Greater Glasgow and Clyde Biorepository, the University of Leicester, and the University of Glasgow for their invaluable support and contributions to this study. The manuscript was critically reviewed by Catherine Winchester (CRUK Scotland Institute). CRUK Early Detection Programme, IAMMED-Meso, EDDPGM-Nov21\100001 supported the research.

References

[1] J Christopher Wagner, Christopher A Sleggs, and Paul Marchand. "Diffuse pleural mesothelioma and asbestos exposure in the North Western Cape Province". In: *British journal of industrial medicine* 17.4 (1960), p. 260.

[2] Linda Molinari. *Mesothelioma survival rates & factors that affect patients*. Nov. 2023. URL: https://www.mesothelioma.com/mesothelioma/prognosis/survival-rate/.

[3] Sara Ricciardi et al. "Surgery for malignant pleural mesothelioma: an international guidelines review". In: *Journal of thoracic disease* 10.Suppl 2 (2018), S285.

[4] Arnaud Scherpereel et al. "ERS/ESTS/EACTS/ESTRO guidelines for the management of malignant pleural mesothelioma". In: *European respiratory journal* 55.6 (2020).

[5] Pierre Courtiol et al. "Deep learning-based classification of mesothelioma improves prediction of patient outcome". In: *Nature medicine* 25.10 (2019), pp. 1519–1525.

[6] Mark Eastwood et al. "MesoGraph: Automatic Profiling of Malignant Mesothelioma Subtypes from Histological Images". In: *arXiv preprint arXiv:2302.12653* (2023).

[7] Julia R Naso et al. "Deep-learning based classification distinguishes sarcomatoid malignant mesotheliomas from benign spindle cell mesothelial proliferations". In: *Modern Pathology* 34.11 (2021), pp. 2028–2035.

[8] Richard J Chen et al. "Scaling vision transformers to gigapixel images via hierarchical self-supervised learning". In: *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*. 2022, pp. 16144–16155.

[9] Xiyue Wang et al. "Transformer-based unsupervised contrastive learning for histopathological image classification". In: *Medical image analysis* 81 (2022), p. 102559.

[10] Alexandre Filiot et al. "Scaling self-supervised learning for histopathology with masked image modeling". In: *medRxiv* (2023), pp. 2023–07.

[11] Richard J Chen et al. "Towards a General-Purpose Foundation Model for Computational Pathology". In: *Nature Medicine* (2024).

[12] Adalberto Claudio Quiros et al. "Mapping the landscape of histomorphological cancer phenotypes using self-supervised learning on unannotated pathology slides". In: *Nature Communications* 15.1 (2024), p. 4596.

[13] Jure Zbontar et al. "Barlow twins: Self-supervised learning via redundancy reduction". In: *International Conference on Machine Learning*. PMLR. 2021, pp. 12310–12320.

[14] Vincent A Traag, Ludo Waltman, and Nees Jan Van Eck. "From Louvain to Leiden: guaranteeing well-connected communities". In: *Scientific reports* 9.1 (2019), p. 5233.

[15] Andrew G Nicholson et al. "EURACAN/IASLC proposals for updating the histologic classification of pleural mesothelioma: towards a more multidisciplinary approach". In: *Journal of Thoracic Oncology* 15.1 (2020), pp. 29–49.

[16] Scott Lundberg. "A unified approach to interpreting model predictions". In: *arXiv preprint arXiv:1705.07874* (2017).

[17] Francoise Galateau Salle et al. "Comprehensive molecular and pathologic evaluation of transitional mesothelioma assisted by deep learning approach: a multi-institutional study of the International Mesothelioma Panel from the MESOPATH Reference Center". In: *Journal of thoracic oncology* 15.6 (2020), pp. 1037–1053.

[18] Sanja Dacic et al. "Interobserver variation in the assessment of the sarcomatoid and transitional components in biphasic mesotheliomas". In: *Modern Pathology* 33.2 (2020), pp. 255–262.

[19] Julija Hmeljak et al. "Integrative molecular characterization of malignant pleural mesothelioma". In: *Cancer discovery* 8.12 (2018), pp. 1548–1565.

[20] Laura Mannarino et al. "Epithelioid pleural mesothelioma is characterized by tertiary lymphoid structures in long survivors: results from the MATCH study". In: *International Journal of Molecular Sciences* 23.10 (2022), p. 5786.

[21] Mark Eastwood et al. "Malignant mesothelioma subtyping of tissue images via sampling driven multiple instance prediction". In: *International Conference on Artificial Intelligence in Medicine*. Springer. 2022, pp. 263–272.

[22] Ming Y Lu et al. "Data-efficient and weakly supervised computational pathology on whole-slide images". In: *Nature biomedical engineering* 5.6 (2021), pp. 555–570.

[23] Kentaro Hashimoto et al. "Sarcomatoid malignant pleural mesothelioma treated with nivolumab: A case series". In: *Oncology Letters* 24.5 (2022), pp. 1–7.

[24] Darya Karatkevich et al. "Chemotherapy increases CDA expression and sensitizes malignant pleural mesothelioma cells to capecitabine treatment". In: *Scientific Reports* 14.1 (2024), p. 18206.

[25] Nicolas Coudray et al. "Classification and mutation prediction from non–small cell lung cancer histopathology images using deep learning". In: *Nature medicine* 24.10 (2018), pp. 1559–1567.

[26] Ting Chen et al. "A simple framework for contrastive learning of visual representations". In: *International conference on machine learning*. PMLR. 2020, pp. 1597–1607.

[27] Jean-Bastien Grill et al. "Bootstrap your own latent-a new approach to self-supervised learning". In: *Advances in neural information processing systems* 33 (2020), pp. 21271–21284.

[28] Kaiming He et al. "Momentum contrast for unsupervised visual representation learning". In: *Proceedings of the IEEE/CVF conference on computer vision and pattern recognition*. 2020, pp. 9729–9738.

[29] Juan José Egozcue et al. "Isometric logratio transformations for compositional data analysis". In: *Mathematical geology* 35.3 (2003), pp. 279–300.

[30] Josep A Martín-Fernández, Carles Barceló-Vidal, and Vera Pawlowsky-Glahn. "Dealing with zeros and missing values in compositional data sets using nonparametric imputation". In: *Mathematical Geology* 35 (2003), pp. 253–278.

[31] Dennis L Wilson. "Asymptotic properties of nearest neighbor rules using edited data". In: *IEEE Transactions on Systems, Man, and Cybernetics* 3 (1972), pp. 408–421.

[32] Maximilian Ilse, Jakub Tomczak, and Max Welling. "Attention-based deep multiple instance learning". In: *International conference on machine learning*. PMLR. 2018, pp. 2127–2136.

[33] Danyu Y Lin and Lee-Jen Wei. "The robust inference for the Cox proportional hazards model". In: *Journal of the American statistical Association* 84.408 (1989), pp. 1074–1078.

[34] Bradley Efron. "Logistic regression, survival analysis, and the Kaplan-Meier curve". In: *Journal of the American statistical Association* 83.402 (1988), pp. 414–425.

[35] Simon Graham et al. "Hover-net: Simultaneous segmentation and classification of nuclei in multi-tissue histology images". In: *Medical Image Analysis* (2019), p. 101563.

[36] Minoru Kanehisa and Susumu Goto. "KEGG: kyoto encyclopedia of genes and genomes". In: *Nucleic acids research* 28.1 (2000), pp. 27–30.

[37] Arthur Liberzon et al. "Molecular signatures database (MSigDB) 3.0". In: *Bioinformatics* 27.12 (2011), pp. 1739–1740.

[38] Etienne Becht et al. "Estimating the population abundance of tissue-infiltrating immune and stromal cell populations using gene expression". In: *Genome biology* 17 (2016), pp. 1–20.

[39] David Allan Moore et al. "In situ growth in early lung adenocarcinoma may represent precursor growth or invasive clone outgrowth—a clinically relevant distinction". In: *Modern Pathology* 32.8 (2019), pp. 1095–1105.

[40] F Alexander Wolf et al. "PAGA: graph abstraction reconciles clustering with trajectory inference through a topology preserving map of single cells". In: *Genome biology* 20 (2019), pp. 1–9.

[41] F Alexander Wolf, Philipp Angerer, and Fabian J Theis. "SCANPY: large-scale single-cell gene expression data analysis". In: *Genome biology* 19 (2018), pp. 1–5.

[42] Cameron Davidson-Pilon. "lifelines: survival analysis in Python". In: *Journal of Open Source Software* 4.40 (2019), p. 1317.

[43] Pauli Virtanen et al. "SciPy 1.0: fundamental algorithms for scientific computing in Python". In: *Nature methods* 17.3 (2020), pp. 261–272.

[44] Skipper Seabold and Josef Perktold. "Statsmodels: econometric and statistical modeling with python." In: *SciPy* 7.1 (2010).

[45] Guillaume Lemaître, Fernando Nogueira, and Christos K. Aridas. "Imbalanced-learn: A Python Toolbox to Tackle the Curse of Imbalanced Datasets in Machine Learning". In: *Journal of Machine Learning Research* 18.17 (2017), pp. 1–5. URL: http://jmlr.org/papers/v18/16-365.

[46] F. Pedregosa et al. "Scikit-learn: Machine Learning in Python". In: *Journal of Machine Learning Research* 12 (2011), pp. 2825–2830.

[47] Sonja Hänzelmann, Robert Castelo, and Justin Guinney. "GSVA: gene set variation analysis for microarray and RNA-seq data". In: *BMC bioinformatics* 14 (2013), pp. 1–15.

Figure 1: (a) Overview of the LATTICe-M dataset clinical information, highlighting key fields. Also, a forest plot for clinical data−Age, Subtype, TNM Stage, as well as Kaplan-Meier for the mesothelioma subtype variable, significantly affect outcome prediction for this cohort. **(b)** HPL Pipeline: Each WSI was divided into 224x224 px tiles. After applying various distortions, these tiles were used as input for the Barlow Twins self-supervised model. In the second step, once the model was trained, the ResNet backbone generated 128D vectors per tile, representing prominent features. These vectors were then clustered using the Leiden community clustering algorithm. In the final step, at the patient level, the clusters (HPCs) were analysed to quantify the proportion of each morphologically significant HPC within the WSI. This information was used to predict mesothelioma subtypes and outcomes using logistic regression and Cox hazard ratio models.

Figure 2: **(a)** Colour-coded UMAP plot for 47 HPCs. **(b)**Percentage of patients from the LATTICe-M cohorts associated with each HPC. **(c)** Pathologists' panel annotations for HPCs; three pathologists independently annotated the clusters. The table shows the majority opinion for histopathological features identified in each cluster. **(d)** Main patterns – supercluster – colour-coded UMAP; annotations were given by a pathologist following the clustering step to group HPCs with similar morphologies.**(e)**A WSI from a poor outcome and good outcome cases overlaid and quantified by HPCs, representing the differences in HPC composition and the percentage of the top 10 HPCs for each sample, alongside morphology annotations.

Figure 3: **(a)** Forest plot for the logistic regression model used in subtype classification, showing each HPC's contribution (regression coefficients), along with significant HPCs for each class (epithelioid vs. sarcomatoid/biphasic), including their p-values, confidence intervals, log odds ratios, and pathologist annotations on HPC morphology. **(b)** The ROC (Receiver operating characteristic) curve for LATTICe-M and TCGA-MESO Datasets showing their subtype classification performance, AUC-ROC, Precision, Sesitivity and Specifity Scores **(c)** The PCA plot shows patient vector representations, colour-coded by mesothelioma subtypes.

Figure 4: (a) Forest plot for the Cox proportional hazards model used in subtype classification, showing each HPC's contribution (regression coefficients), along with significant HPCs for each class (epithelioid vs. sarcomatoid/biphasic), including their p-values, confidence intervals, log odds ratios, and pathologist annotations on HPC morphology.**(b)** UMAP plot of tile vector representations, colour-coded by log odds ratio and hazard ratio assigned to each HPC. **(c)**Kaplan-Meier plots showing low-risk and high-risk patient groups for LATTICe-M and TCGA-Meso datasets, with their reported p-value performance scores.**(d)** Kaplan-Meier plots comparing the performance of our risk and survival predictor, pre-trained on the LATTICe-M dataset, tested on TCGA-MESO (only epithelioid cases), versus predictions based on the traditional systemic overall grade of the epithelioid cases.**(e)**An example of a SHAP decision plot for a high-risk (red) and low-risk (blue) patient, displaying the percentage of HPCs within their samples that contribute to their outcomes.

Figure 5: (a) HPC frequency plot for biphasic/sarcomatoid and epithelioid cases, illustrating the co-occurrence of HPCs for each subtype in comparison to the other. Based on a Cox model trained on subtype-filtered cases, significant HPCs in the model ($p-value < 0.05$) are highlighted in bold colours with respect to their log hazard ratio, with sample tiles shown in the legend. **(b)**Grouped HPCs based on pathologist annotations, highlighting key histopathological features such as necrosis, desmoplastic components, and inflammation in the tumour microenvironment. These HPCs provide insights into the different morphologies and their potential roles in disease progression and patient prognosis.

Figure 6: Spatial sequencing data correlations with HPCs; supercluster annotations were provided by expert pathologists. The survival score shows the log odds ratio from the logistic regression model, while survival scores are based on the partial hazards ratio predicted by the Cox model. The inflammation score row was generated using inference from the pre-trained HoVer-Net model. **(a)** Ki67 expression and MCPcounter immune features. **(b)** Kyoto Encyclopedia of Genes and Genomes (KEGG). **(c)** Molecular Signatures Database (MSigDB) hallmarks.