Back to the horns: a reconstruction of the ancestral horn state through distinct types of recombination events.

M. Upadhyay[1], A. Graf[2], N. Pogorevc[1], D. Seichter[3], I. Russ[3], S. Krebs[2], S. Meier[4], I. Medugorac[1*]

[1] Population Genomics Group, Department of Veterinary Sciences, LMU Munich, Martinsried, 82152, Germany.
[2] Laboratory for Functional Genome Analysis, Gene Center, Ludwig-Maximilians-University Munich, 80539, Munich, Germany.
[3] Tierzuchtforschung e.V. München, 85586, Grub, Germany.
[4] SYNETICS Germany GmbH, Research & Genetic Solutions Development, Osterkrug 20, 27283 Verden (Aller), Germany.

*Corresponding author

E-mail addresses:

IM: I.medugorac@lmu.de

Abstract.

Background

Breeding of the genetically polled animals is the desirable approach in modern cattle husbandry. At least four different genetic variants associated with polledness in cattle have been identified, suggesting genetic heterogeneity. These dominant variants have been identified on chromosome 1 between the regions of approx. 2.42 to 2.73 Mb (reference: ARS-UCD1.2), also called the *POLLED* locus. Among these variants, Friesian (P_F, 80 kbp duplication) and Celtic (P_C, 212 bp complex InDel) are the most observed in the majority of breeds in the production systems globally, such as in Holstein-Friesian (HF) and Fleckvieh (FV). While a putative causal association of P_C with polledness is proven, the presence of large duplication in P_F makes it difficult to prove the causality.

Results

In this study, we conduct whole-genome sequencing (WGS) analysis of two trios exhibiting unexpected inheritance patterns related to P_C and P_F variants. In both instances, horned offspring were produced from mating pairs where one parent was homozygous for the polled variant, and the other was homozygous for the ancestral horned variant. By analyzing the WGS data generated using long-read sequence technology, we show that *de-novo* generation of the ancestral horned variant in both the offspring was the result of distinct recombination events. Specifically, in case of the HF trio, it was the result of non-allelic homologous recombination in the gametes of the sire (P_F/P_F), while in case of the FV trio, it was the result of allelic homologous recombination in the gametes of the dam (P_C/P_F). The findings from the HF trio support the hypothesis that the 80-kbp duplication is the genetic variant responsible for the polled phenotype of Friesian origin.

Conclusion

Here we show that different genomic arrangements in *POLLED* locus can lead to the emergence of *de-novo* ancestral horn phenotypes. These kinds of arrangements can make a reliable gene test less reliable and the derivation of the phenotype difficult to predict. Therefore, it is important for the large *POLLED*-locus that any deviation from the expected result is critically analyzed. Possibly some of these cases can further narrow down the sequence motif that is essential for polledness in cattle.

Background

Based on the size, shape, and characteristics of horns, the head phenotype in cattle can be categorized into these three classes: (i) true horns in bovids contain a pneumatized bony inner core that is attached to the frontal bone and have an outer keratin sheath [1]; (ii) scurs are rudimentary horn-like appendages that are loosely attached to the skin rather than the frontal bone with not pneumatized bony inner core. (iii) complete absence of horns-like appendages, a condition called polledness. Like their wild ancestors, the aurochs, and all members of the Bovidae family, cattle are naturally horned. Therefore, the presence of true horns is an ancestral characteristic of Bovidae, and scurs and polledness are derived traits. Horns have been associated with a diverse range of functions such as in exerting territorial dominance, competing for mating and other resources, protecting against predators, and maintaining thermoregulation [2-4]. The process of domestication and subsequent breed formation has largely rendered horns redundant. In fact, in modern industrial settings, horned animals are undesirable as they pose injury risks to themselves, other cattle as well as their handlers. Further, it is commonly assumed that because of its special requirements, horned cattle are associated with high barn investment and labor costs [5]. Therefore, polled conditions in cattle have become highly desirable. To this end, a variety of chemical and surgical methods are used for disbudding or dehorning. Across the cattle sector, these practices are commonplace. According to a survey published in 2015, in Europe, more than 80% of dairy, 46% of beef, and 67% of suckler calves are disbudded or dehorned [6]. All these procedures are labor intensive and cause pain and distress to animals; further, they also leave the wound sites exposed to infection, adding to the distress [5, 7]. Therefore, the animal welfare guidelines recommend adapting husbandry systems to the ancestral characteristics of cattle or breeding genetically hornless animals rather than dehorning them.

Early and accurate detection of animals having genetic predisposition for polled phenotype is key to selective breeding for polled animals [8]. Polled phenotype is a Mendelian trait governed by multiple variants (genetic heterogeneity) that are located on BTA1 (bovine autosome 1) between the region of approx. 2.42 to 2.73 Mb (reference: ARS-UCD1.2), also referred to as *POLLED* locus throughout the manuscript. Phenotypic penetration suggests that all these variants are dominant and cause polledness. Mainly, these four polled variants have been identified so far: (i) Celtic (P_C): This complex insertion-deletion variant, "g.[2429326_2429335del;2429109_2429320dupins]", was first identified in several European breeds; since the geographical distribution of these breeds overlaps mainly with the area in which Celtic culture was practiced, it was named as "Celtic" [9]. (ii). Friesian (P_F): this tandem duplication variant, "g.2629116_2709243dup (80,128 bp)", was first identified in Holstein-Friesian (HF) cattle [9, 10]. The duplicated segment has the same orientation as the reference sequence. It differs from the reference sequence by a SNP (single Nucleotie polymorphism) ($P_{T->A}$) and 2 bp deletion (P_{F2D}, TG) on 1'st bp and 38th bp positions respectively on the duplicated sequence. (iii). Mongolian (P_M): this complex insertion-deletion variant, "[2695261_2695267delinsTCTGAA;2695889_2696047dupins]", was first identified in Mongolian Turano cattle and Mongolian yaks [11]. The P_M variant is located within the 80,128 bp segment duplicated by P_F duplication. (iv). Gaurani (P_G): This duplication variant,

"g.2614828_2724315dup", was first identified in Nellore cattle from Brazil [12] and completely
overlaps with P_F and P_M. Interestingly, none of these four variants are directly involved in the gene
coding. While several hypotheses have been proposed [13, 14], the exact pathways through which
these complex variants disrupt the migration or proliferation or differentiation of cells in the horn
bud region is not known so far [15].

Among these four variants, P_C and P_F are the most observed polled variants in the majority of
breeds in the production systems globally. Of these variants, a putative causal association of P_C
with polledness is proven by introgression of P_C allele from genome-edited cell lines resulting in
the birth of polled calves [16]. For the P_F variant, the presence of large duplication (~80 kb) makes
it difficult to prove the causality by genome editing or to pinpoint the exact genomic region
responsible for the polledness condition. The relatively small P_M structural variant (SV), that is
embedded in P_F, however, suggests that the more distal part of the 80 kb region could be essential
to the polledness. The question still remains whether any SV along the entire *POLLED* locus (i.e.
from the start of the P_C duplication (2,429,326) to the end of the P_G segment (2,724,315) is
sufficient to disrupt the process of horn bud formation.

In this study, we conduct whole-genome sequencing (WGS) analysis of two trios exhibiting
unexpected inheritance patterns related to P_C and P_F variants. In both instances, horned offspring
were produced from mating pairs where one parent was homozygous for the polled variant, and the
other was homozygous for the ancestral horned variant. The polled gene test targeting 2-bp deletion
(P_{F2D}) and 212 bp insertion (entire P_C variant) was used to confirm the presence of P_F and P_C
variants, respectively. Specifically, in the HF trio, the sire was homozygous P_F/P_F, while in the FV
trio, the dam carried P_C/P_F genotype. Therefore, resulting offspring were expected to exhibit
polledness by inheriting either P_C or P_F polled variant. To our surprise, here investigated offspring
from each trio, showed the regular horn condition and *p/p* genotype instead. After confirming
parentage with BovineSNP50 BeadChip, we hypothesized that recombination events in the
POLLED locus of the paternal germ cells in HF trio and maternal germ cells in FV trio may have
resulted in a novel genomic rearrangement, giving rise to the variant similar to the wild type. These
trios provided us an opportunity to shed light on the genomic composition of the P_C and P_F variants.

Methods

Sample information

In this study, we investigated two trios, one from the HF breed and one from the FV breed. In the
HF trio, the sire was confirmed to be homozygous for the P_F variant, while the dam expressed a
horned phenotype and according to genetic test *p/p* genotype. Notably, the sire has over 13,000
registered offspring, of which only one examined here was declared to exhibit the horned
phenotype. In the FV trio, the dam carried one copy of both the P_C and P_F variants, while the sire
was tested as horned with *p/p* genotype. The presence of the P_F variant in the FV population was
expected, as this breed is, since 1960s, introgressed by HF and thus carries a minor but significant
genetic ancestry from the HF breed. In both trios, offspring were genotyped as *p/p* and exhibited

regular horns fused to frontal bone instead of expected polled phenotypes and polled genotypes
(Figure 1).

DNA extraction and whole genome long-read sequencing

The DNA was extracted from the frozen blood or semen (sires) samples immediately before the
sequencing to get an increased fraction of long DNA fragments in the sequencing. The DNA
extraction was carried out from 100 µl blood using the QiaAmp Blood Mini Kit kit protocol
(Qiagen, Hilden, Germany). Briefly, the resuspended blood was lysed and digested using proteinase
K and RNase A, followed by binding to a column, washing, and elution through centrifugation.

To carry out long-read sequencing using the Oxford PromethION sequencer, one µg of unsheared
DNA was end-repaired and A-tailed with NEBnext UltraII End-repair and A-tailing module (New
England Biolabs, Ipswich USA). According to the manufacturer's protocols, this mixture was later
purified with AmpureXP magnetic beads (Beckman Coulter, Brea USA), ligated to 1D sequencing
adapters (LSK114 kit, Oxford Nanopore Technologies (ONT), Oxford, UK), and again purified
with AmpureXP beads. In the last Ampure cleanup step, instead of washing with 70% ethanol, a
washing buffer supplied by Oxford Nanopore Technologies (LSK114 kit) was used. Approximately
200 ng of adapted DNA library was loaded to a primed PromethION R10.4.1M flow cell and
sequenced for 72 h on a PromethION P24 sequencer.

SNP genotyping, haplotyping and parentage and Polled test

Both trios were genotyped with the Illumina BovineSNP50 BeadChip (Illumina San Diego, USA)
according to the manufacturer's protocol. All 54,157 markers were used to test the parentage in
both trios. Subsequently, we confirmed the expected pedigree in both the trios and for FV trio, even
traced the pedigree to much deeper level than the first generation pedigree because our genotypic
database included four FV grandparents. To investigate the recombination events, the inference of
haplotypes and imputation of missing genotypes were carried out using hidden Markov models as
implemented in BEAGLE (v 5.0) [17, 18]. In order to improve the accuracy of the phasing,
genotype and pedigree information of additional trios and pairs of both breeds genotyped with
BovineSNP50K BeadChip, otherwise not included in the analysis of this paper, were also included
in this step of phasing.

Polled test was carried out according to the procedure and primers described in Medugorac *et al.*[9].
Briefly, the primer binding sites flanking the P_C variant InDel events (5'-
TCAAGAAGGCGGCACTATCT-3' and 5'-TGATAAACTGACCCTCTGCCTATA-3') were PCR
amplified (94 °C 30 sec, 58 °C 60 sec, 72 °C 60 sec for 31 cycles). Next, PCR products were size-
separated and visualized using 2% ethidium-bromide stained agarose gel electrophoresis.
Genotyping of the P_F 80 kb duplication event was carried out using two primers flanking the
variable site P_{F2D} (5'-GAAGTCGGTGGTCTGAAAGG-3' and 5'-
TGTTCTGTGTGGGTTTGAGG-3'). Next, PCR amplification (32 cycles 94 °C 30 sec, 59 °C 60
sec, 72 °C 60 sec) generated a RefSeq related product (p_{ref}); this product was obtained in all

animals, be it horned or polled. Additionally, a second product related to P_F variant, containing the
two-base pair (TG) deletion was only observed in the animals that carried one or two copies of the
P_F variant (P_{FDUP}), see (Figure 1). To discriminate between these two products related to P_F
variant, ABI Prism 3130XL DNA sequencer was used. Animals carrying P_F/P_F were distinguished
from P_F/p using quantitative evaluation of the obtained signals. The animals carrying P_F/P_F yielded
signals of similar peak heights as the P_F and p_{ref} products, while heterozygous products (P_F/p)
resulted in signal intensities of approximately double height for the p_{ref} product when compared to
the product specific to P_F.

Bioinformatics pipeline to analyze WGS data

The samples were sequenced on a PromethION sequencer from Oxford Nanopore and were
basecalled and demultiplexed during sequencing using Dorado (Dorado basecall server:
v.7.3.11+0112dde09). After basecalling, basic quality checks of the sequences were carried out
using NanoPlot (v 1.41.6) [19]. Subsequently, the sequencing data were aligned to the cattle
reference genome sequence (ARS-UCD1.2) using minimap2 (v 2.28) with secondary alignments
suppressed [20, 21]. Small genomic variants (SNPs and short InDels) were identified using Clair3
(v 1.0.10) with the appropriate trained model for the basecaller. Structural variations (SVs) were
detected using Sniffles2 (v 2.4), cuteSV (v 2.1.1), Dysgu (v 1.6.7) and SVIM (v 2.0.0)[22-26].
Moreover, SVs were also detected using a de novo genome assembly-based approach. For this
purpose, Flye (v 2.9.5) [27] was used for de novo assembly of the trios, and SVIM-asm (v 1.3.0)
[28] was used for the detection of SVs. Finally, all genomic variants were phased using LongPhase
(v 1.7.3) [29].

To investigate the split-read alignments in the POLLED locus region and to identify recombination
breakpoints in the offspring's genome using the phased information of the parents' genotype,
custom python scripts were developed using the pysam library (https://github.com/pysam-
developers/pysam) for efficient manipulation of the alignment data. The entire bioinformatics
pipeline was implemented in the Nextflow workflow management system [30], which is available
on GitHub at https://github.com/Popgen48/gvdlr/tree/main. To visualize alignment patterns in the
BAM files, we used JBROWSE2 [31] and IGV [32].

Primer design and validation of candidate variants

By examining the POLLED locus in both trios, we searched for short variants that would be
specific for them. We found these putative variants: (a) 1 bp deletion (p_{ref1D}) and a SNP with
transition ($p_{refT->C}$) in the HF trio (b) a SNP with transition ($P_{G->A}$) in the Fleckvieh trio. For
additional sequencing of the regions containing the mentioned three variants, we designed primers
with a tool Primer3Plus [33]. We also sequenced both ends of the duplicated segment P_{FDUP}.
Details about PCR products are explained in Table 1. The PCR amplification was performed with
35 cycles of 30 seconds on 94 °C, 60 seconds on 59 °C and 60 seconds on 72 °C. PCR products of
the trios were converted to barcoded nanopore sequencing libraries (SQK-NBD114.96) and
sequenced on a Promethion R10.4.1 flowcell for approximately 2h. The resulting fastq files were

aligned on the cattle reference genome sequence (ARS-UCD1.2) using minimap2 (v 2.28) with
secondary alignments suppressed. Then, the bam files were visualized using Jbrowse2.

Results

207 The trios were sequenced with ONT and on average produced 36.03 Gbp sequencing data per
208 sample with an average read length N50 of 9.70 Kbp. A detailed overview of the sequencing
209 statistics and alignment statistics is provided in Table 2.

Analysis of HF trio

In the HF trio, the sire was homozygous for the P_F variant. This variant referred to the entire
duplicated segment (Fig. 1): the original segment, p_{ref} (between 2,629,113 and 2,709,240 bp on
BTA1) and its tandem duplicate (P_{FDUP}). The absence of P_{F2D} deletion in the offspring pointed
towards a *de-novo* deletion of at least a part of the P_F variant involving P_{FDUP}. Concerning this
deletion, five hypotheses were synthesized (Fig. 2). Next, each of these five hypotheses were
investigated based on the results of comprehensive variant calling and manual inspections of the
entire region.

Analysis of HF sire

Using the Nextflow pipeline developed specifically for this study to identify comprehensive
genomic variants from the ONT sequences, a total of two structural variants (SVs) and two SNPs
were detected (Fig. 1 and Supplementary file S1 Table S1). Expectedly, the sire carried the P_F
variant (Fig. 1). Interestingly, the visualization of the breakpoints identified 2 split reads that
suggested that a part of the variant is possibly duplicated one more time (Fig. S1). Indeed, a *de-novo*
assembly of the sire also supported this observation (Fig. S2).

Analysis of HF offspring

The Nextflow pipeline identified a total of one structural variant (SV), an insertion, and eight single
nucleotide polymorphisms (SNPs) in the sequence that could maintain parts of segments p_{ref} and
P_{FDUP} in the offspring. The SV (Supplementary file S1 Table S1) was identified solely by the
Dysgu tool; however, it was classified as a low-quality variant, as it was detected on reads with a
mapping quality of 1, indicating multiple read alignments. Notably, no other tools identified any
SVs supported by two or more reads. Furthermore, our custom Python tool, designed to detect split
reads, did not find any regions with two or more split reads sharing common breakpoints.
Additionally, no regions larger than 1 kbp showed significantly higher or lower coverage (with
more than one split read) compared to the average coverage of the region. These findings
collectively suggest that the hypotheses H1, H2 and H3 (Fig. 2) could be rejected.

Next to investigate the hypothesis H4 and H5, it was important to differentiate between reads that
originated from p_{ref} and those originating from P_{FDUP}. However, apart from two variants (one SNP
and a 2 bp deletion, P_{F2D}) detected at the very beginning of the duplicated regions, no other variant

that could differentiate the duplicated segment (P_{FDUP}) from the reference sequence (p_{ref}) has been
identified in previous studies [9, 10]. To assess whether any trio-specific short variants could be
used for this purpose, extensive manual inspection of the region was carried out in the bam files of
sire as well as offspring. A 1 bp deletion (p_{ref1D}, located on 2,682,320 bp) present in the alignment
of both sire and offspring was identified as one such candidate (Fig. S3). The long reads containing
reference and alternative alleles at the candidate's 1-bp deletion and SNP suggested their
localisation in the duplicated segment. However, the sequencing of PCR products to validate p_{ref1D}
and the de-novo SNP suggested that these variants were false positives (Supplementary file S2
Table S2). Thus, to conclude, based on these observations, it is most likely that the genomic region
showing the sequence composition of p/p in the offspring may have been the results of a
recombination event which have happened anywhere in the P_F segments and deleted ~80,128 bp of
the genomic segments covering either a small part of the 3' region of p_{ref} and large part of the 5'
region of P_{FDUP} segment or the entire P_{FDUP} segment (H4 or H5 hypothesis in Fig. 2). Since no
insertions, duplications or multiple split reads with common breakpoints were detected in the
bioinformatic and manual (window-by-window) examination of the entire 80,128 bp in the HF
offspring and this offspring sequence was identical to that of an animal of the p/p-genotype (e.g.
dam), it is obvious that the deletion in the paternal germ cell was exactly 80,128 bp long.

Analysis of the Fleckvieh trio

In the Fleckvieh trio, the dam carried P_C and P_F variants each. Since the P_F allele is inherited from
the sire and the P_C from the dam, these two variants are in *trans* phases in the homologous
chromosome pair of the dam, i.e. located on different members of a homologous pair of
chromosomes. Further, an allelic recombination might have resulted in recombinant haplotype (with
"gene conversion") in which the sequences roughly until the beginning of the P_{FDUP} originated from
the chromosome containing P_F variants and the sequences after that originated from the
chromosome containing P_C variant (Fig. 3).

Analysis of Fleckvieh dam

The Nextflow pipeline identified a total of two SVs and 179 SNPs in the combined region of P_C and
P_F variants. These two SVs were P_C and P_F , confirming the genotypes of the dam. Among these
SVs, only P_C was detected by all the tools implemented in the workflow, while P_F variant was not
detected by Sniffles2, one of the widely used tools to detect SV and it was detected as having low
quality by cuteSV. These variants were also visually confirmed using Jbrowse2 (Fig. S4 and Fig.
S5).

Analysis of Fleckvieh offspring

The Nextflow pipeline identified a total of three SVs and 810 SNPs in the combined region of P_C
and P_F variants. None of the identified SVs corresponded to the polled variants, P_C and P_F,
confirming the results of genotyping (pp). The higher number of SNPs was due to the combination
of higher coverage (than dam); the sire also had high coverage and 910 SNPs.

The analysis of the phased SNPs in the offspring showed that one of the haplotypes, which
originated from dam, had recombinant haplotype that switched from being of one parental origin (a
chromosome containing P_F variant) to the other (a chromosome containing P_C variant) roughly
around the chromosomal position, 2,694,883 bp on BTA1 (Fig. 3). This resulted in skipping of both
the polled variants, P_C and P_F in one of the haplotype of the offspring. It should be noted that
although the sequencing of PCR products did not validate the SNP reported by Clair3 at this
position, the switching in the haplotype was obvious from this position onwards (Supplementary
file S3). This was also supported by the recombination inferred using the 50K SNP genotypes of the
trio (Supplementary file S4).

Discussion

There is growing evidence that different POLLED variants in a heterozygous state suppress horn
growth to varying degrees. For example, [34] found that almost every heterozygous P_C/p Fleckvieh
(FV) animal forms some kind of scurs, ranging from a minimal keratin crust to a horn fused to the
frontal bone. The author concluded that the visibility and strength of the scurs depends on the sex,
age and polygenic background of the FV animals. Later, we compared the frequency of scurs in
heterozygous P_F/p and P_C/p animals and confirmed that P_F suppresses the development of horns
significantly better than P_C [35]. Furthermore, there is evidence that the P_M variant completely
suppresses the development of horns in heterozygous (P_M/p) Mongolian Turano cattle and
Mongolian yaks [11]. In our sample of 58 hornless Mongolian yaks and 40 hornless Mongolian
Turano cattle, only 16 (30 %) yaks and 5 (12.5 %) Turano cattle were homozygous P_M/P_M and all
others were heterozygous P_M/p, but all were smooth-polled. After this observation, we tried to
collect samples of Turano cattle and yaks with scurs. However, we found that breeders in Mongolia
and China had never observed scurs in their animals and informed us that the animals were either
horned or smooth polled. This supports our observation that the P_M variant embedded in a large P_F
duplication could be most effective at suppressing the development of horns in heterozygous yak
and cattle.

In our previous study [10], we used recombination within the *POLLED* locus to prove that
segmental duplication p_{ref}-P_{FDUP} is the only remaining causal candidate for the polledness of the
Friesian origin. The back-mutation of P_F in the offspring of the here investigated HF trio offers the
opportunity for additional confirmation of P_F segmental duplication as causal variant. The HF trio
was analyzed with the aim to decipher the genomic composition of the region containing the P_F
variant. Specifically, after observing the missing P_{F2D} deletion in the offspring of a HF trio
investigated here, we hypothesized that the most parsimonious explanation of this could be that a
portion of P_F in the paternal germ cell that led to the offspring must have been deleted. The deleted
portion include P_{F2D} and SNP $P_{T->A}$ at the transition from p_{ref} to P_{FDUP} (Fig. 2) but we assumed that
some remaining parts of P_{FDUP} and p_{ref} should be still present in paternal haplotype of HF offspring,
i.e. one of the hypotheses H1 to H3 should apply. This would suggest that the part removed de novo
in the P_{FDUP} segment could be responsible for the polledness in cattle. Surprisingly, our analysis
neither identified any deletion nor the duplication in the entire segment in the offspring. In fact, our

observations ruled out hypotheses H1, H2 and H3. The perfect reconstruction of the entire p_{ref}

segment could be achieved by a deletion of exactly 80,128 bp and then fusing of the remaining parts

of p_{ref} and P_{FDUP} segments back to wild-type sequence (H4, Fig. 2). It should also be considered

that wild-type variant created in such an arrangement is indistinguishable to the structural

arrangement in which the complete P_{FDUP} sequence is deleted (H5, Fig. 2): This is because P_{FDUP} is

the almost exact duplicated sequence of p_{ref} except the two variants (P_{F2D} and SNP $P_{T->A}$) which are

located at the beginning of the P_{FDUP} segment and not present in HF offspring. Therefore, any

deletion of 80,128 bp in the P_F gamete, starting at any position between g.2629153 and g.2709243,

will reconstruct the wild-type sequences. The only way to distinguish between H4 and H5 is to find

and confirm some new variants that are present in P_{FDUP} but lacking in the p_{ref} segment of the HF

sire or vice-versa. Our analysis did not identify any such high-quality candidate variant.

Nevertheless, these results confirm that a duplication of at least part of the p_{ref} segment (as in the P_M

variant) or of the entire p_{ref} segment is required for the polled state in cattle.

Segmental duplications (SDs) are duplicated blocks of genomic DNA typically ranging in size from

1–200 kb with a sequence identity of over 90% [36]. Thus, the duplicated blocks p_{ref} and P_{FDUP} can

be classified by definition as long and almost perfect SDs. Typically, SDs and its flanking repeats

are the sites of non-allelic homologous recombination (NAHR) [37]. The direction and orientation

as well as the chromosomal localization (intra-chromosomal or inter-chromosomal) of the

duplicated segments involved in NAHR determine the type of SVs that may arise[37-39]. Meiotic

recombination is initiated by DNA double-strand breaks (DSB). In case of allelic homologous

recombination (AHR), the damaged sequence around the DSB is usually repaired by replacement

with the allelic sequences of the sister chromatids as template. Mostly this recombination process is

highly faithful and ensures accurate distribution of alleles and chromosomes as long as the DSB

occurs in the unique regions of the genome. In the case of a DSB within the SDs regions, it is

possible that the replication machinery finds the segment's paralogs instead of the actual allelic

segment for repair, resulting in NAHR [40]. All types of NAHR (intra-chromatid, inter-sister

chromatid or interhomologue) can generate a deletion or other recurrent genomic rearrangements.

Thus, based on the results, it is likely that, during the DSB repair in the region of P_F variant, the

replication machinery found the paralogous instead of allelic segment, thus resulting in hybrid SD

[39]. During the process, as the repeats are very similar, it skipped the genomic region of the same

size as that of the duplicated segment, resulting in a perfect deletion of one of two segments. It is

noteworthy that the HF sire has more than 13,000 offspring and therefore, it could be that a gamete

containing such a rare recombination event contributed to an offspring investigated here.

The polled variants P_F and P_C are the result of two independent mutations in different

subpopulations of the bovine species. Therefore, they segregate on different haplotype backgrounds.

In some breeds, such as Jersey, Fleckvieh and Holstein, both variants are inherited independently in

different lines and, in rare cases, animals inherit P_C from one parent and P_F from another. These

animals are homozygous for *POLLED*, denoted as *PP*, but heterozygous for the Celtic and for

Frisian mutations, which are 200 kbp apart. The dam of the Fleckvieh trio is such an animal with P_C

and P_F allele in trans phase. According to [41] on average 1.085 Mb corresponds to 1.0 cM in the

female cattle. Therefore, on average, we expect two out of 1000 oocytes to show recombination
between the P_C and P_F allele. It is expected that this proportion is even lower for the *POLLED* locus
because in cattle all autosomal chromosomes are acrocentric [42] and regions near the centromere
show the lowest recombination rate [42-44]. According to the results obtained from the analysis of
the long-read sequences of FV-Trios, the following scenarios seems to be most probable: during
prophase of the first meiotic division in the dam of the FV trio, a DSB was introduced in one of the
sister chromatids carrying the P_F variant. This most likely occurred closer to the distal end of the
p_{ref} segment ~14 kbp upstream from the start of the P_{FDUP} segment. This DSB was repaired by
crossover with the allelic position of a non-sister chromatid containing the P_C variant proximal to
the DSB and p_{ref} segment distal to the DSB. The crossover fused the proximal portion of the P_F
chromatid to the distal portion of the P_C chromatid, thereby reconstructing the wild-type haplotype.

The results of our study point toward some interesting observations and open up intriguing
possibilities that recombination events in the *POLLED* locus might result in. First, the
recombination breakpoint inferred in the gametes of a dam is within 1-kb distance of P_M variant.
Next, in both the trios, the resulting hybrid SD has a similar composition (i.e., the wild-type
arrangement was reconstructed), but are generated by different recombination types. In theory, it is
equally likely that the crossover in FV-Trio fuses the P_C and P_F variants into a *cis*-haplotype. The
offspring with P_C and P_F in *cis*-arrangement would be phenotypically polled and would most likely
not be observed as something unexpected and therefore falsely declared as homozygous polled, *PP*.
Similarly, the recombination within the SD of the P_F variant could lead to different arrangements,
which would make a reliable gene test less reliable. Therefore, it is important for large *POLLED*-
locus that any deviation from the expected results is critically analyzed. Possibly some of these
cases can further narrow down the sequence motive that is essential for polledness in cattle.

Our results also indicated that comprehensive detection of variants using the ONT sequences should
involve application of multiple tools that implement a variety of approaches. For instance, in the FV
dam, while the P_C variant was identified confidently by all the tools implemented in the workflow,
the P_F variant was missed by Sniffles2 and detected as a low quality variant by cuteSV. Further,
comprehensive detection of short variants like InDels and SNPs with high confidence are difficult
using long-read sequencing technology like ONT because of its relatively high error rates.
Therefore, our data suggest that the short variants identified using ONT sequencing technology
should be thoroughly validated even after applying the variety of algorithms to identify such
variants comprehensively.

Conclusions

Here, using the 50k genotyping data and long-read sequencing data generated from HF and
Fleckvieh trios, we comprehensively investigated the genomic composition of Friesian and Celtic
polled variants. Our data support the hypothesis that the 80-kbp duplication is the genetic variant
responsible for the polled phenotype of Friesian origin. Next, we also showed that different
genomic arrangements because of allelic homologous recombination and non-allelic homologous

recombination in *POLLED* locus can lead to the emergence of *de-novo* ancestral horn phenotypes.

These kinds of arrangements can make a reliable gene test less reliable and the derivation of the

phenotype difficult to predict. Therefore, it is important for the large *POLLED*-locus that any

deviation from the expected result is critically analyzed. Possibly some of these cases can further

narrow down the sequence motif that is essential for polledness in cattle.

List of abbreviations

AHR : Allelic homologous recombination

BTA1 : Bovine autosome 1

DSB : Double-strand break

NAHR : Non-allelic homologous recombination

ONT : Oxford nanopore technology

SD : Segmental duplication

SNP : Single nucleotide polymorphism

SV : Structural variation

WGS : Whole genome sequencing

Declarations

Not applicable

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

414 The WGS data of the trios will be deposited in the public repository upon acceptance of the

416 manuscript. All the scripts and workflows used for the analysis can be accessed on the github page:

417 https://github.com/Popgen48/gvdlr.

Competing interests

419 The authors declare that they have no competing interests

Funding

421 The first author MU received a funding from Deutsche Forschungsgemeinschaft (DFG)—

422 Projektnummer 526367023.

Authors' contributions

MU**:** Formal Analysis; investigation; methodology; visualization; writing – original draft

AG**:** Formal analysis; investigation; editing

NP**:** Visualization; editing

DS**:** Resources; editing

IR**:** Resources; editing

SK**:** Formal analysis; resources; editing

SM**:** Resources; editing

IM**:** Conceptualization; funding acquisition; project administration; resources; supervision;

Visualization; writing – review & editing

Acknowledgements

434 Johann Robeis LfL Bayern and breeder of the investigated HF bull (Bokelmann GbR, Sanne,
435 Germany) for sample taking and providing phenotypic information.

References

1. Zhang Y, Huang W, Hayashi C, Gatesy J, McKittrick J. Microstructure and mechanical
properties of different keratinous horns. Journal of The Royal Society Interface.
2018;15(143):20180093.
2. Aldersey JE, Sonstegard TS, Williams JL, Bottema CDK. Understanding the effects of the
bovine POLLED variants. Animal Genetics. 2020;51(2):166-76.
3. Schafberg R, Swalve HH. The history of breeding for polled cattle. Livestock Science.
2015;179:54-70.
4. Stankowich T, Caro T. Evolution of weaponry in female bovids. Proceedings of the Royal
Society B: Biological Sciences. 2009;276(1677):4329-34.
5. Knierim U, Irrgang N, Roth BA. To be or not to be horned—Consequences in cattle. Livestock
Science. 2015;179:29-37.
6. Cozzi G, Gottardo F, Brscic M, Contiero B, Irrgang N, Knierim U, et al. Dehorning of cattle in the
EU Member States: A quantitative survey of the current practices. Livestock Science. 2015;179:4-
11.
7. Stafford KJ, Mellor DJ. Dehorning and disbudding distress and its alleviation in calves. The
Veterinary Journal. 2005;169(3):337-49.
8. Spurlock DM, Stock ML, Coetzee JF. The impact of 3 strategies for incorporating polled genetics
into a dairy cattle breeding program on the overall herd genetic merit. J Dairy Sci.
2014;97(8):5265-74.
9. Medugorac I, Seichter D, Graf A, Russ I, Blum H, Gopel KH, et al. Bovine polledness--an
autosomal dominant trait with allelic heterogeneity. PLoS One. 2012;7(6):e39477.
10. Rothammer S, Capitan A, Mullaart E, Seichter D, Russ I, Medugorac I. The 80-kb DNA
duplication on BTA1 is the only remaining candidate mutation for the polled phenotype of Friesian
origin. Genetics Selection Evolution. 2014;46(1):44.
11. Medugorac I, Graf A, Grohs C, Rothammer S, Zagdsuren Y, Gladyr E, et al. Whole-genome
analysis of introgressive hybridization and characterization of the bovine legacy of Mongolian yaks.
Nature Genetics. 2017;49(3):470-5.
12. Utsunomiya YT, Torrecilha RBP, Milanesi M, Paulan SdC, Utsunomiya ATH, Garcia JF.
Hornless Nellore cattle (Bos indicus) carrying a novel 110 kbp duplication variant of the polled
locus. Animal Genetics. 2019;50(2):187-8.
13. Allais-Bonnet A, Grohs C, Medugorac I, Krebs S, Djari A, Graf A, et al. Novel Insights into the
Bovine Polled Phenotype and Horn Ontogenesis in Bovidae. PLOS ONE. 2013;8(5):e63512.
14. Randhawa IAS, Burns BM, McGowan MR, Porto-Neto LR, Hayes BJ, Ferretti R, et al.
Optimized Genetic Testing for Polledness in Multiple Breeds of Cattle. G3
Genes|Genomes|Genetics. 2020;10(2):539-44.
15. Aldersey J, Chen T, Petrovski K, Williams J, Bottema CK. Histological characterisation of the
horn bud region in 58 day old bovine fetuses. Int J Dev Biol. 2024;In Press(In Press):40-.
16. Carlson DF, Lancto CA, Zang B, Kim E-S, Walton M, Oldeschulte D, et al. Production of
hornless dairy cattle from genome-edited cell lines. Nature Biotechnology. 2016;34(5):479-81.
17. Browning BL, Tian X, Zhou Y, Browning SR. Fast two-stage phasing of large-scale sequence
data. The American Journal of Human Genetics. 2021;108(10):1880-90.
18. Browning BL, Zhou Y, Browning SR. A One-Penny Imputed Genome from Next-Generation
Reference Panels. The American Journal of Human Genetics. 2018;103(3):338-48.
19. De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. NanoPack: visualizing and
processing long-read sequencing data. Bioinformatics. 2018;34(15):2666-9.

20. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.
2018;34(18):3094-100.
21. Rosen BD, Bickhart DM, Schnabel RD, Koren S, Elsik CG, Tseng E, et al. De novo assembly
of the cattle reference genome with single-molecule sequencing. Gigascience. 2020;9(3).
22. Cleal K, Baird DM. Dysgu: efficient structural variant calling using short or long reads. Nucleic
Acids Res. 2022;50(9):e53.
23. Jiang T, Liu S, Cao S, Wang Y. Structural Variant Detection from Long-Read Sequencing Data
with cuteSV. Methods Mol Biol. 2022;2493:137-51.
24. Smolka M, Paulin LF, Grochowski CM, Horner DW, Mahmoud M, Behera S, et al. Publisher
Correction: Detection of mosaic and population-level structural variants with Sniffles2. Nat
Biotechnol. 2024;42(10):1616.
25. Smolka M, Paulin LF, Grochowski CM, Horner DW, Mahmoud M, Behera S, et al. Detection of
mosaic and population-level structural variants with Sniffles2. Nat Biotechnol. 2024;42(10):1571-
80.
26. Heller D, Vingron M. SVIM: structural variant identification using mapped long reads.
Bioinformatics. 2019;35(17):2907-15.
27. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat
graphs. Nature Biotechnology. 2019;37(5):540-6.
28. Heller D, Vingron M. SVIM-asm: structural variant detection from haploid and diploid genome
assemblies. Bioinformatics. 2021;36(22-23):5519-21.
29. Lin JH, Chen LC, Yu SC, Huang YT. LongPhase: an ultra-fast chromosome-scale phasing
algorithm for small and large variants. Bioinformatics. 2022;38(7):1816-22.
30. Di Tommaso P, Chatzou M, Floden EW, Barja PP, Palumbo E, Notredame C. Nextflow enables
reproducible computational workflows. Nat Biotechnol. 2017;35(4):316-9.
31. Diesh C, Stevens GJ, Xie P, De Jesus Martinez T, Hershberg EA, Leung A, et al. JBrowse 2: a
modular genome browser with views of synteny and structural variation. Genome Biology.
2023;24(1):74.
32. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al. Integrative
genomics viewer. Nature Biotechnology. 2011;29(1):24-6.
33. Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al. Primer3--new
capabilities and interfaces. Nucleic Acids Res. 2012;40(15):e115.
34. Heidrich K. Phänotypische und molekulargenetische Untersuchungen zur Vererbung von
Wackelhörnern beim Fleckvieh. Munich: LMU Munich; 2018.
35. Gehrke LJ, Capitan A, Scheper C, König S, Upadhyay M, Heidrich K, et al. Are scurs in
heterozygous polled (Pp) cattle a complex quantitative trait? Genetics Selection Evolution.
2020;52(1):6.
36. Vervoort L, Vermeesch JR. Low copy repeats in the genome: from neglected to respected.
Exploration of Medicine. 2023;4(2):166-75.
37. Gu W, Zhang F, Lupski JR. Mechanisms for human genomic rearrangements. Pathogenetics.
2008;1(1):4.
38. Chen JM, Cooper DN, Férec C, Kehrer-Sawatzki H, Patrinos GP. Genomic rearrangements in
inherited disease and cancer. Semin Cancer Biol. 2010;20(4):222-33.
39. Parks MM, Lawrence CE, Raphael BJ. Detecting non-allelic homologous recombination from
high-throughput sequencing data. Genome Biol. 2015;16(1):72.
40. Sasaki M, Lange J, Keeney S. Genome destabilization by homologous recombination in the
germ line. Nat Rev Mol Cell Biol. 2010;11(3):182-95.
41. Ma L, O'Connell JR, VanRaden PM, Shen B, Padhi A, Sun C, et al. Cattle Sex-Specific
Recombination and Genetic Control from a Large Pedigree Analysis. PLoS Genet.
2015;11(11):e1005387.
42. McFeely R, Popescu P. Chromosomes of the Cow and Bull. In: McFeely R, Popescu P, editors.
Domestic Animal Cytogenetics. 34: Academic Press; 1990. p. 41-71.

43. Stapley J, Feulner PGD, Johnston SE, Santure AW, Smadja CM. Variation in recombination
frequency and distribution across eukaryotes: patterns and processes. Philos Trans R Soc Lond B
Biol Sci. 2017;372(1736).
44. Stapley J, Feulner PGD, Johnston SE, Santure AW, Smadja CM. Correction to 'Variation in
recombination frequency and distribution across eukaryotes: patterns and processes'. Philos Trans
R Soc Lond B Biol Sci. 2018;373(1739).

584 # Figures

Figure 1. Overview of the inheritance patterns in the trios and genomic composition of the P_F variant.

(A) a schematic representation of the trios, including their genotypes, (B) a schematic
representation of the wild type locus and friesian polled variant (P_F); note that P_F variant consists of
a reference segment p_{ref} and its duplication P_{FDUP} segment. Two variants, namely, $P_{T->A}$, P_{F2D} ,
depicted on the 5` side of P_{FDUP} segment refer to a SNP and a 2 bp InDel already reported to be
present on P_{FDUP}, (C) a schematic representation of the patterns in which the reads sequenced from
a sample containing P_F variant would align to the reference, (D) actual screenshot of the split reads
sequenced from the HF sire which is homozygous for P_F variant.

Figure 2. Overview of the hypotheses investigated in the HF trio.

Schematic representation of the hypotheses synthesized to explain the absence of P_{F2D} variant in the offspring of the HF trio.

Figure 3. Overview of the hypothesis investigated in the Fleckvieh trio

Schematic representation of the haplotypes in the Fleckvieh trio. Note the presence of P_F and P_C variants in the trans-phase. The arrow between two maternal haplotypes indicates the most likely position of the recombination in the maternal germ cell.

Tables

Table 1: Information about PCR primers and its products used to validate candidate variants. The location refers to the position on chromosome 1 on ARS-UCD1.2 assembly.

Target	Location (in bp)	Trio	Primers	Product length	Annealing temperature
$P_{G->A}$	2,694,883	FV	5´-GGAACTGGAACTACTGAAGTCCT-3`  5´-TGTTTATCCCACAGCTGTTGGA-3`	2943 bp	59°C
$P_{T->C}$	2,684,110	HF	5´-TGCATTGCTGTAGTCTGTTTGC-3`  5´-GGTAAGAGGTGCAGCCATGT-3`	1424 bp	58°C
p_{ref1D}	2,682,320	HF	5´-TATGTGCAGTGTGAGGCAGG-3`  5´-ATGAATCAGCCATGGGTGCA-3`	1281 bp	65°C
P_{FDUP} left-end	2,629,116	HF	5´-GGGATCAGTCTTGGGCCCTA-3`  5´-TGACTTGGCCACAGACATCC-3`	1268 bp	59°C
P_{FDUP} right-end	2,709,243	HF	5´-ATGCAGCTGGTCTTTCCTCC-3`  5´-ACCTTTCTACCATGCGGCAA-3`	1338 bp	59°C

613   **Table 2.** Basic sequencing and alignment statistics of the HF and FV trio

Sample id	genotypes	Relationship	Total bases sequenced (in Gbp)	Total number of reads sequenced	Read length N50 (in bp)	Mean alignment coverage
HF964	$P_F P_F$	Sire	59.20	15,907,973	7,786	22.98x
HF963	*pp*	Dam	25.00	2,670,919	13,986	9.68x
HF962	*pp*	Offspring	59.28	20,506,948	3,957	22.50x
FV2831	*pp*	Sire	18.92	3,564,540	8,379	7.24x
FV2822	$P_F P_C$	Dam	31.34	4,180,434	10,001	12.01x
FV2823	*pp*	Offspring	22.46	2,528,157	14,093	8.80x

Additional files

Supplementary file S1 Table S1

Format: word file (.docx)

Title : Table S1

Description: Summary of the comprehensive genomic variants identified using *Clair3* and *longshot* tools in the offspring of HF trio and its corresponding genotypes in its sire and dam.

Supplementary file S2 Table S2

Format: word file (.docx)

Title : Table S2

Description: The result of very high coverage sequencing of PCR products using ONT technology.

Supplementary file S3

Format: text file (.txt)

Title: Phasing and inference of recombination point in the FV trio using the variants identified from the long read sequences

Description: The tab-delimited file shows the phasing information of the variants identified in the FV dam using long-read sequences. The last two column refers to the genotypes of the offspring and sire that correspond to the respective position in the dam; the "NA" indicates that the variant was not called at this position because the sample is homozygous for the reference allele. The arrow ">" indicates the allele inherited in the offspring. Notice the switch in the position of arrows from the genomic position, 2,694,883 bp.

Supplementary file S4

Format: word file (.docx)

Title: Phasing and inference of recombination point in the FV trio using the variants genotyped with Bovine SNP 50K array.

Description: The file shows the phased data of the FV trio using the Bovine SNP 50K array genotypes. The recombinant haplotype and its respective positions are highlighted.