Dendritic synaptic integration modes under *in vivo*-like states

Cesar C. Ceballos[1], Rodrigo F. O. Pena[1,2] *

[1] Department of Biological Sciences, Florida Atlantic University, Jupiter, FL 33458, USA

[2] Stiles-Nicholson Brain Institute, Florida Atlantic University, Jupiter, FL 33458, USA

* Corresponding author

E- mail: penar@fau.edu

Abstract

The neural code remains undiscovered and understanding synaptic input integration under *in vivo*-like conditions is just the initial step toward unraveling it. Synaptic signals generate fast dendritic spikes through two main modes of temporal summation: coincidence detection and integration. In coincidence detection, dendrites fire only when multiple incoming signals arrive in rapid succession, whereas integration involves summation of postsynaptic potentials over longer periods with minimal membrane leakage. This process is influenced by ionic properties, especially as the membrane potential approaches the firing threshold, where inactivating currents play a critical role. However, the modulation of temporal summation by these currents under *in vivo*-like conditions has not been thoroughly studied. In our research, we used computer simulations to investigate how three inactivating currents — A-type potassium, T-type calcium, and transient sodium — affect temporal summation. We found that calcium and sodium currents promote integrative behavior in dendrites, while potassium currents enhance their ability to act as coincidence detectors. By adjusting the levels of these currents in dendrites, neurons can flexibly switch between integration and coincidence detection modes, providing them with a versatile mechanism for complex tasks like multiplexing. This flexibility could be key to understanding how neural circuits process information in real time.

Introduction

It is well known that dendrites can fire sodium, calcium and NMDA dependent action potentials (dendritic spikes) under *in vivo* conditions at higher frequency than found at the soma (Moore et al 2017). These suggests the existence of dendritic synaptic integration and transformation into dendritic spikes as a first stage of neuronal processing (Ujfalussy et al 2018). Understanding the neural code involves learning how dendrites process synaptic inputs and transform them into dendritic spikes via temporal summation (Stuart et al 2015, Dembrow et al 2022). Two key summation types, coincidence detection and integration, are critical for neural coding (Brette 2015, Ratté et al 2013, Hong et al 2012, Stoll et al 2023). Coincidence detection occurs when a dendrite fires a spike after receiving multiple signals within a short interval, while integration mode allows voltage accumulation over time until a spike fires (Dembrow et al 2015, Dembrow et al 2022, Roome et al 2020). Dendrites exhibit both modes, with Purkinje neurons dendrites functioning as coincidence detectors and L5 dendrites in medial prefrontal cortex acting as integrators (Dembrow et al 2015, Roome et al 2020). These modes influence the transmission of information via firing rates or synchronized spikes (Hong et al. 2012).

These synaptic summation modes form a continuum from coincidence detection to perfect integration (Rudolph et al. 2003, Prince et al. 2021). The specific mode of summation depends on the composition of ionic currents in the dendritic membrane, which modulate synaptic temporal properties (Azouz et al. 2000, Prince et al. 2021). Strong potassium currents are linked to coincidence detection, while sodium and calcium currents support perfect integration (Mathews et al. 2010, Farries et al. 2010, Branco et al. 2016, Prescott and De Koninck 2005). Adjusting the expression of these currents allows a range of temporal summation modes.

Dendritic inactivating currents are crucial for near-threshold synaptic summation, as shown by

studies on A-type potassium currents influencing theta oscillations and temporal summation

(Bourdeau et al. 2007; Rathour et al. 2016). EPSPs are amplified by T-type calcium currents,

enhancing temporal precision (Fukaya et al. 2018), while transient sodium currents act at

subthreshold potentials (Carter et al. 2012; Stuart et al. 2015). Further research is needed to

explore inactivation's role in dendritic properties.

A complete understanding requires examining currents with (i) established, (ii) less established,

and (iii) inconsistent effects. (i) Non-inactivating currents like M-type potassium reduce EPSP

amplitude and shorten decay (Hönigsperger et al. 2015; Peng et al. 2017; Liu et al. 2008), while

persistent sodium currents amplify and prolong EPSPs (Ceballos et al. 2017b; Liu et al. 2008;

Hsu et al. 2018). (ii) Inactivating currents, such as A-type potassium, consistently reduce EPSP

amplitude (Hoffman et al. 1997; Mathews et al. 2010; Bourdeau et al. 2007; Rathour et al. 2016).

(iii) Calcium currents show mixed effects: some studies report no impact on amplitude or decay

(Stuart et al. 1995; González-Burgos et al. 2001; Andreasen et al. 1999; Hsu et al. 2018; Seong et

al. 2014), others show effects on amplitude (Fukaya et al. 2018; Gillessen et al. 1997), only on

decay (Prescott et al. 2005), or both (Liu et al. 2008; Connelly et al. 2016; Urban et al. 1998;

Evans et al. 2017; Deleuze et al. 2012). Transient sodium currents also show variable effects:

some studies report no effect (Connelly et al. 2016; Grunditz et al. 2008), while others show

amplification of EPSPs (Carter et al. 2012; Hage et al. 2016) or TTX-sensitive prolonged decay

(Ceballos et al. 2018). Sodium-dependent EPSP amplification has been observed *in vivo* (Hsu et

al. 2018).

Most EPSP studies are not easily transferable to *in vivo* situations, where neural coding is better

understood. *In vivo*, large depolarization states occur during neuronal oscillations like theta

oscillations (Adam et al. 2019; Malezieux et al. 2020). These states bring the membrane potential

near the threshold, suggesting the activation of inactivating currents (Hsu et al. 2018), though

little is known about how these conductances affect synaptic summation *in vivo*. This raises

questions about how these currents filter synaptic signals and determine synaptic summation

modes.

Understanding how ionic currents shape membrane temporal properties is essential for dendritic

computation, influencing whether dendrites function as integrators or coincidence detectors. This

paper explores the effects of A-type potassium, T-type calcium, and transient sodium currents on

synaptic temporal summation *in vivo*-like conditions, focusing on their impact on voltage traces

and summation properties (integrator vs. coincidence detector).

Results

Consistent membrane charging effects in the presence of potassium and sodium currents,

but mixed response effects in the presence of calcium current

We begin our paper testing the effect of inactivating currents on τ_m. For that, we built a single

compartment dendritic model. We focused on the most relevant inactivating currents expressed

in pyramidal neurons dendrites and activated near the spiking threshold: A-type potassium (I_{KA}),

T-type calcium (I_{CaT}), and transient sodium (I_{NaT}) (see Methods). A depolarizing pulse of current

was applied, and both exponential rise and decay followed by this pulse were analyzed. Fig. 1A

shows that I_{KA} hyperpolarizes the resting membrane potential, reduces the amplitude of the

response, and accelerates the capacitor's charge. In contrast, I_{NaT} amplifies the response

amplitude and slows down the capacitor's charge but has no effect on the resting membrane

potential due to its activation at higher membrane potentials (Fig 1B).

Opposed to this pattern, the effects of I_{CaT} were mixed. I_{CaT} depolarizes the resting membrane

potential. However, there is only amplification of the voltage change at the onset of the current

pulse, while there is a reduction of the voltage change at the end of the pulse due to slow

inactivation kinetics (Fig 1C). We also observed a clear phase delay during the rising phase of

the EPSP (excitatory postsynaptic potential), but the decay phase showed mixed behavior,

starting with a phase delay in one direction and eventually reversing its direction (Fig 1D). We

believe this mixed behavior in amplification/reduction of the voltage change and

lengthening/shortening of the voltage provides insight into the inconsistent reports found in the

literature about I_{CaT} (Stuart et al. 1995; González-Burgos et al. 2001; Andreasen et al. 1999; Hsu

et al. 2018; Seong et al. 2014, Fukaya et al. 2018; Gillessen et al. 1997, Prescott et al. 2005, Liu

et al. 2008; Connelly et al. 2016; Urban et al. 1998; Evans et al. 2017; Deleuze et al. 2012).

Voltage changes scaling monotonically during theta oscillations

The previous section studied the effect of the inactivating currents on the temporal properties of

a dendritic model. The model was stimulated using simple current pulses and simulated EPSPs.

Here, we wondered whether the previous results also apply under *in vivo*-like conditions. Here,

we investigated the effects of the inactivating currents on scaling voltage traces from *in vivo*-like

conditions. We injected synaptic currents, consisting of evenly distributed EPSCs and IPSCs

over time, to generate sparse synaptic backgrounds as found *in vivo* (Waters et al., 2006). To add

realistic background to the synaptic currents, we simulated theta oscillations at 4 Hz and

analyzed four cycles of voltage traces (Fig. 2; cycles identified by different colors). Our results

show that I_{KA} causes a negative shift of the voltage changes (Fig 2A), whereas I_{CaT} and I_{NaT}

cause a positive shift (Fig 2C, 2E). The scaling from above -60 mV is linearly correlated with the

membrane voltage for I_{KA} and I_{NaT} (Fig 2B, 2F) but no clear tendency is found for I_{CaT}, only high

variability (Fig 2D).

Fig 3 shows the nonlinear scaling for input frequency oscillations from 1 to 30 Hz. The set of

Figs 3A-C demonstrate a clear difference in the voltage scaling when comparing the ionic

currents. More specifically, we observed a consistent decrease for I_{KA} across depolarizing

voltages and a consistent increase for I_{NaT} across depolarizing voltages (see heatmaps in Fig 3D

and Fig 3F). In contrast, I_{CaT} showed nonmonotonic behavior of the scaling for low frequencies,

but monotonic for high frequencies (Fig 3E). Then we evaluated the behavior of the scaling for

different conductance values at 4 Hz. We observed a consistent decrease for I_{KA} across

depolarizing voltages and a consistent increase for I_{NaT} across depolarizing voltages (Fig 3G and

Fig 3I). In contrast, I_{CaT} showed similar values across voltage (horizontal lines, Fig 3H).

These results suggest that most of the knowledge from the Fig 1 can be transferred to the *in vivo-*

like states with except to the I_{CaT}, with ionic currents scaling the voltage differently in a

frequency-dependent manner.

Changes in activation are linearly correlated with changes in inactivation for T-type

calcium and A-type potassium currents

We previously observed the effect of the inactivating currents on the scaling of the voltage. I_{KA}

and I_{NaT} showed linear correlation with voltage, but no consistent tendency was observed for

I_{CaT}. In this section, to investigate the biophysical mechanisms underlying these discrepancies we

present phase plots of activation versus inactivation variables for theta oscillations (Fig. 4). This

analysis demonstrated a clear negative correlation, where increases in activation correspond with

decreases in inactivation (Fig 4G, 4H, 4J and 4K). Interestingly, the window current properties of

I_{NaT} are markedly different than I_{KA} and I_{CaT} and produced a highly nonlinear correlation

between activation and inactivation variables, where large values of one variable correlate with

small values of the other (Fig 4I). We verified that this relationship is well-fitted by a power law

function (Fig 4L).

Analysis of the activation vs. inactivation phase plots showed that fitting trajectories of I_{KA} and

I_{CaT} by a linear regression returns a curve very similar to the trajectories when $\tau_{act} = \tau_{inact} = 0$ (Fig

4J and 4K). This allows us to approximate the functional relationship between activation or

inactivation. Therefore, we use the linear regression to substitute the inactivation variable by a

linear function dependent on the activation variable: $q_{Inact} = - p_{Act} + 0.85$ for I_{KA} and, $q_{Inact} = -$

$0.84 p_{Act} + 0.8$ for I_{CaT}. In contrast, the trajectories for I_{NaT} are highly nonlinear and, as explained

above, were fitted using a power law function with 4/3 exponent ($q_{Inact} = 10^{-3} p_{Act}^{-4/3}$) (Fig 4I

and 4L). This functional behavior is the result of the way activation and inactivation curves

interact within the window current.

Our results showed similar activation/inactivation linear trajectories for I_{KA} and I_{CaT}, although

these two currents differ in their inactivation kinetics (e.g. $\tau_{inact} = 5$ ms for I_{KA} but 10 ms for I_{CaT}).

This suggests that under *in vivo*-like conditions, voltage changes are robust to inactivation

kinetics.

Slow inactivation plays nearly no role on summation during *in vivo*-like states

Based on the results above, we interrogate whether *in vivo*-like states require kinetics. Therefore,

we investigated whether we could get similar voltage traces when using the actual kinetics vs.

when using fast kinetics, i.e. $\tau_{act} = \tau_{inact} = 0.1$ ms. Surprisingly, for all simulated currents we

obtained traces that were similar with average differences less than 0.6 mV for voltages below

spiking threshold (V = -40 mV; Fig 5D). These results suggest that the system of equations can

be reduced when assuming $\tau_{act} = \tau_{inact} = 0$, hinting that the dynamics provided by Eq 3 is no

longer necessary for *in vivo*-like conditions. Suprathreshold spiking activity also displayed

similar results (Suppl Fig 1).

Further, and following the relationships from the previous section, we can also reduce the

inactivation variable using the linear relation obtained from the linear regression above for I_{KA}

and I_{CaT} and power law approximation for I_{NaT} inactivation. As a result of all the reductions, we

can express the behavior of each current as dependent on only a single activation variable.

Therefore, we use the linear regression to substitute the inactivation variable by a linear function

dependent on the activation variable for I_{KA} and I_{CaT} and power law approximation for I_{NaT}. We

also consider $\tau_{act} = \tau_{inact} = 0$. Fig 5H demonstrates that this approximation can obtain similar

voltage traces with average values below 0.7 mV for all currents.

The results from this section allowed us to reduce the dendrite system of equations to a single

variable that resolves the activation variable only. Our approximation is akin to the Morris-Lecar

reduction (Izhikevich 2007). However, in our case, the assumption of instantaneous kinetics

allows us to avoid coupling differential equation.

In the next set of experiments, we wondered about the dependency of our results with the input

frequency. Fig 6 shows the voltage difference between the model with slow kinetics vs.

instantaneous for input frequency oscillations from 1 to 30 Hz. We observed a consistent

increase for all currents across depolarizing voltages, with values below 0.7 mV for I_{KA}, 1.2 mV

for I_{CaT} and 0.35 mV for I_{NaT} (Fig 6D, 6E and 6F). Then we evaluated the voltage difference

between the model with slow kinetics vs. instantaneous kinetics for different conductance values

with input frequency fixed at 4 Hz. We observed a consistent increase for all currents across

depolarizing voltages, with lowest values at conductance close to $g_{KA} = 1$, $g_{CaT} = 0.1$ and $g_{NaT} =$

50 (Fig 6G, 6H and 6I).

These results suggest that the voltage differences between the model with slow kinetics and fast

kinetics is independent on the frequency input, but dependent on the voltage. As to the latter, the

stronger the depolarization the higher was the voltage difference identified, showcasing that the

kinetics becomes more prominent as the voltage gets closer to the threshold of the neuron.

Inactivation opposes activation effects on voltage scaling and membrane time constant

We know from above that activation and inactivation have negative correlation, but it is still

unknown how this correlation determines the voltage scaling and temporal properties of the

voltage traces in the presence of I_{KA}, I_{CaT} and I_{NaT}. On the other hand, it is well known that

scaling is inversely related to the input conductance and that the temporal properties are related

to the membrane time constant. We first used an analytical approach to determine the opposed

effect of the inactivation on voltage scaling and temporal properties compared to activation.

When comparing the input conductance for inactivating current (as shown in Eq 8) vs non-

inactivating current (as shown in Eq 10), we know from Eq 9 that $p_i^\infty q_i^\infty < p_i^\infty$ and $q_i^\infty \frac{\partial p_i^\infty}{\partial v} +$

$p_i^\infty \frac{\partial q_i^\infty}{\partial v} < \frac{\partial p_i^\infty}{\partial v}$. Thus, input conductance of inactivating currents is always smaller compared to

the non-inactivating case, suggesting that the effect of inactivation is always to oppose to the

activation.

To test the opposite effects of activation and inactivation, we investigated the temporal

summation of a train of 3 EPSPs separated by 10 ms. We tested the effect of the inactivating

current and their respective non-inactivating version (Fig 7). The plots show that noninactivating

currents enhance the effect compared to their inactivating counterpart. These results suggest that

inactivation counteracts the effect of the activation, reducing its effect.

While in this section we showed that inactivation opposes activation effects on voltage scaling

and membrane time constant with analytical derivations, together with previous sections we can

conclude that I_{KA} activation shortens τ_m whereas inactivation prolongs it (Fig 7A). Meanwhile,

an opposite effect occurs for I_{CaT} and I_{NaT} (Fig 7B and 7C). Together, these observations have

profound implications for dendritic computation, in particular to their temporal summation

properties. In the next section, we will show how our results are useful to study synchrony- vs.

rate-code.

A-type potassium current enhances synchrony-code but T-type calcium and transient

sodium currents enhance rate-code

Temporal summation of synaptic potentials can be used to identify integrators or coincidence

detector neurons. Integrators, characterized by a long τ_m, allow summation of synaptic potentials

even when they occur far apart in time. In contrast, coincidence detectors, with a short τ_m,

integrate synaptic potentials only when they occur closely together in time. Previous studies have

shown the association between high expression of potassium currents and coincidence detection

mode, and high expression of calcium/sodium currents and integration mode (Mathews et al

2010, Branco et al. 2016). We tested this hypothesis using an analytical approach to demonstrate

that for instantaneous inactivating currents, potassium currents such as I_{KA} always decrease the

input resistance and shorten the membrane time constant, meanwhile calcium/sodium currents

such as I_{CaT}, and I_{NaT} always increase the input resistance and prolong the membrane time

constant (see Methods, section Mathematical analysis of reducing/contracting and

amplifying/prolonging currents).

Moreover, coincidence detection and integration on the level of dendritic compartments can be

seem translated to synchrony-code and rate-code on the level of network computations, with

coincidence detectors being more likely to fire with synchronized inputs and poorly integrate

sparse inputs. In contrast, integrators can summate sparse inputs to generate dendritic spikes but

respond less to coincident inputs compared to coincidence detectors. Thus, a natural question we

investigate in this section is how the modulation of the I_{KA}, I_{CaT} and I_{NaT} affect the synchrony vs.

rate-code.

Accordingly, we randomly generated 100 sparse and 100 synchronous input currents and used

them to generate 100 voltage traces also stimulated by theta input (4 Hz). The sparse and

synchronous input were developed to create distinction between these codes (synchrony- vs.

rate-code). Both sparse and synchronous inputs had the same amount of 200 EPSCs and 200

IPSCs randomly distributed in the course of a simulation. For the synchronous case, we removed

randomly 75% of the EPSCs and the remaining were amplified by a factor of 4 with the rationale

of reproducing a highly synchronous set of inputs although sparse.

While a consistent decrease in firing occurs in the presence of the I_{KA} (compare Fig 8A to B), an

increase in firing occurs for I_{CaT} and I_{NaT} (compare Fig 8A to C and D). This aligns with the

shrinkage and amplification of voltage changes seen above. When all three currents were

present, firing rates remained unchanged (not shown), indicating that the effects of I_{KA} were

counteracted by I_{CaT} and I_{NaT} as expected.

To further quantify our results in Fig. 8, we require a metric that can determine the preferred

synaptic summation mode (coincidence vs. integration). Thus, we calculated the relative change

in spiking frequency of a dendrite when presented with sparse versus synchronized synaptic

inputs. We expect that coincidence detectors will show an increase in relative spiking frequency,

while integrators will show a decrease. We tested this new metric in a single passive

compartment dendrite model by changing the conductance value or capacitance density (Suppl

Fig 2). The heatmap shows the percentage of relative frequency change when dendrites were

stimulated using sparse vs synchrony input. We used this index (% relat freq) as a measure of the

strength of the preferred neural code mode employed by the dendrite. For instance, % relat freq

of 100% suggests the dendrite acts as a perfect coincident detector, meanwhile 0% suggests the

dendrite acts as a perfect integrator. According to the values obtained from the index, we found

that the dendrite would act closer to a perfect integrator when g_{Leak} was low or when capacitance

density was high, both conditions consistent with long τ_m.

As shown in Fig 8, synchronous input increases firing rate. The relative increase in frequency is

highest at the onset of the rise and decay of the theta cycle and lowest at the peak (Fig 8E, 8F,

8G and 8H). The exception is the potassium current where no large relative increase was

observed at the decay (Fig 8F). We also calculated the relative difference of frequency between

sparse vs synchronous stimulation (Fig 8E, 8F, 8G and 8H). We found a significant increase in

the presence of I_{KA} relative to the passive dendrite (17.44 $\pm$ 1.38 % for passive, 38.86 $\pm$ 4.65 %

for I_{KA}, p <0.001). We also found a significant decrease in the presence of I_{CaT} and I_{NaT} relative

to the passive dendrite (9.77 $\pm$ 0.81 % for I_{CaT}, 10.53 $\pm$ 0.8 % for I_{NaT}, p <0.001). Also, I_{KA} was

significantly higher than I_{CaT} and I_{NaT} (p<0.001). Finally, I_{CaT} and I_{NaT} were not significantly

different (p = 0.5). These results suggest that I_{KA} enhances synchrony-code, but I_{CaT} and I_{NaT}

enhance rate-code.

We then expand our analysis to investigate the tuning range of specific neural code modes and

how they depend on the conductance values of I_{KA}, I_{CaT} and I_{NaT}. We use the same dendritic

model used above, but now we added all 3 currents to the model, while changing their

conductance values selectively. According to the values obtained from the index, we found that

the dendrite would act closer to a perfect integrator when g_{KA} was low (Fig 9C) or when

g_{CaT}/g_{NaT} were high (Fig 9G, H). Under these conditions, the variability of the other

conductances did not affect much the index value (the heatmaps are blueish). These results

suggest I_{KA} is responsible for changing the dendrite to a more coincidence detection mode

meanwhile I_{CaT} and I_{NaT} take the dendrite closer to an integration mode. This can also be

observed for high values of g_{KA} (see Fig 9I) or when g_{CaT}/g_{NaT} is low (Fig 9A, B). Under these

conditions the variability of the other conductances play a significant role that affects the index

value (the heatmaps change from dark blue to light yellow).

Discussion

This study examines an important aspect of the neural coding by focusing on different modes of

in vivo dendritic synaptic integration. We show the role of inactivating currents on dendritic

temporal properties, and how the modes vary from integration of synaptic potentials over time

(long τ_m) to coincidence detection of closely timed potentials (short τ_m) (Mathews et al. 2010,

Prescott and De Koninck 2005, Ceballos et al. 2018, Farries et al. 2010, Branco et al. 2016).

Our results point to the role of A-type potassium currents in shortening τ_m, while T-type calcium

and transient sodium currents lengthen it, similar to non-inactivating currents. This suggests

inactivation reduces current strength without altering its profile. Additionally, activation and

inactivation affect τ_m in opposite ways, with currents that shorten τ_m upon activation prolonging

it upon inactivation and vice versa, consistent with previous studies (Richardson et al. 2003).

These findings also apply to fast-activating, non-inactivating currents, e.g. M-type potassium and

persistent sodium currents.

In terms of firing, we found that A-type potassium currents consistently reduce firing rates,

whereas T-type calcium and transient sodium currents increase them. This is due to potassium

currents hyperpolarizing the membrane, reducing voltage amplitude, and shortening temporal

changes, which decreases excitability. In contrast, calcium and sodium currents depolarize the

membrane, enhance voltage amplitude, and extend temporal changes, thereby increasing

excitability.

In relation to the effects of inactivating currents under *in vivo*-like conditions with oscillatory

inputs and synaptic fluctuations, both I_{KA} and I_{NaT}, were observed scaling changes above -60

mV, showing monotonic trends with variability. The rising and decay phases followed distinct

pathways, indicating that these currents differentiate between depolarization and

hyperpolarization. No clear trend was found for I_{CaT}, but its amplification pattern remained.

Activation and inactivation trajectories were linearly correlated for I_{KA} and I_{CaT}, but nonlinear for

I_{NaT}, independent of kinetics. Surprisingly, activation/inactivation kinetics had minimal impact

on voltage changes, with differences of less than 1 mV, suggesting potential misinterpretations

when extrapolating *in vitro* data to *in vivo* contexts. This underscores the importance of *in vivo*

recordings to fully understand dendritic behavior (Moore et al. 2017, Ujfalussy et al. 2018,

Dembrow et al. 2022, Liao et al. 2024).

We can regard our results as a step towards understanding how dendrites process information. In

particular, we have also tested temporally distinct patterns. Synaptic summation in the presence

of A-type potassium currents promoted firing with synchronized input, while T-type calcium and

transient sodium currents supported firing with sparse input. Thus, A-type potassium currents

enhance synchrony-coding, whereas calcium and sodium currents favor rate-coding. These

currents modulate dendritic integration in cortical and hippocampal dendrites (Stuart et al.,

2015). Our findings align with experimental evidence showing that inactivating T-type calcium

currents extend synaptic potentials (Connelly et al. 2016; Gillessen and Alzheimer 1997; Liu and

Shipley 2008; Prescott and De Koninck 2005; Urban et al. 1998), and inactivating A-type

potassium channels shorten them (Hoffman et al. 1997; Mathews et al. 2010). Notably, we are

the first to show that transient sodium currents have a lengthening effect.

Future work should explore other factors influencing the activation/inactivation properties of

ionic currents and systematize findings across species. For example, sodium and potassium

current voltage-dependence is more depolarized in humans than mice (Wilbers, 2023), and

isoflurane hyperpolarizes sodium current inactivation (Qiu et al., 2023). Neuromodulation also

enhances sodium current inactivation (Chen et al., 2006). Though we focus on *in vivo*-like states,

future studies could investigate how alternating states affect synaptic integration.

Methods

Dendrite model

The most general case of a Hodgkin-Huxley (HH) current involves two gating variables: p for

activation and q for inactivation. This type of current is referred to as "inactivating" and acts as a

break to close the channel (Hille, 2001). Each gating process has its own time constant, denoted

as τ_{act} for activation and τ_{inact} for inactivation.

The current equation is

$C\frac{dv}{dt} = \sum_{i=1}^{n} I_i(v) + I_{inj},$ (Eq 1)

where C is the capacitance, v is the voltage, $I_i(v)$ is the voltage-dependent current, n is the

number of ionic currents and I_{inj} is the injected current. In our paper, ionic currents will be

represented using the HH formalism:

$I_i = \bar{g}_i p_i q_i (V - E_{rev}^i),$ (Eq 2)

where $\bar{g}_i$ is the maximum conductance, E_{rev}^i the reversal potential, and for $(x=p,q)$ we have

$\frac{dx}{dt} = \frac{x^\infty - x}{\tau_{act/inact}}.$ (Eq 3)

In Eq. 3, x^∞ is the steady-state activation variable and it is represented by a sigmoid function

following

$$x^\infty(v) = \frac{1}{1+e^{(\frac{V-V_{1/2}}{-k})}} \cdot \qquad\qquad \text{(Eq 4)}$$

Where $V_{1/2}$ and k are constants. Importantly, $p = p(v,t)$ and $q = q(v,t)$, i.e., activation and

inactivation variables are dependent on voltage and time.

In this paper we only address inactivating currents activated by depolarization such as transient

sodium, A-type potassium and T-type calcium. One important mathematical property of p and q

in this situation that can be derived from the fact that they are activation and inactivation

functions is that $\frac{\partial p}{\partial v} > 0$ and $\frac{\partial q}{\partial v} < 0$, although p > 0 and q > 0. This behavior occurs for currents

activated by depolarization, opposed to when the currents are activated by hyperpolarization

($\frac{\partial p}{\partial v} < 0$ and $\frac{\partial q}{\partial v} > 0$).

Importantly, many inactivating potassium, sodium, and calcium currents follow Equation 2,

while some variations of this equation exist where the activation or inactivation variables are

raised to an exponent (such as 2, 3 or 4). Thus, the equation can be represented as $I_i = $

$\bar{g}_i p_i{}^x q_i{}^y (V - E_{rev}^i)$. Although we will use this representation in our simulations, we will use the

simple version as shown in Eq.2 for the analytical approximation.

We performed computer simulations using custom-made codes in Matlab. A single compartment

dendritic model with ionic currents using the HH formalism. Three types of currents were

investigated: A-type potassium, T-type calcium and transient sodium. Capacitance density was

1μF/cm^2. Activation variables were defined by the following parameters: for potassium: $V_{1/2}$ = -

50 mV, k = 15, for calcium: $V_{1/2}$ = -50 mV, k = 15, and for sodium: $V_{1/2}$ = -30 mV, k = 5.

Inactivation variables were defined by the following parameters: for potassium: $V_{1/2}$ = -60, k =

15, for calcium: $V_{1/2}$ = -60 mV, k = 15, and for sodium: $V_{1/2}$ = -60 mV, k = 5. Activation kinetics

was τ_{act} = 0.2, 0.5 and 0.05 ms for potassium, calcium and sodium respectively. Inactivation

kinetics was τ_{inact} = 5, 10 and 0.5 ms for potassium, calcium and sodium respectively.

Conductance densities were expressed as mS/cm^2 with values: g_K = 1, g_{Ca} = 0.1 and g_{Na} = 50.

Leak conductance density was 0.3 mS/cm^2. Reversal potential was E_{rev} = -80, 50 and 50 mV, for

potassium, calcium and sodium respectively. Leak conductance reversal potentials was -80 mV.

Models of A-type potassium, T-type calcium and transient sodium currents are a modified

version from Engbers et al 2012, Jaffe et al 1994, Korngreen et al 2000, Dougalis et al 2017, and

Hoffman et al. 1997. A-type potassium current equation is: $I_{kA} = g_{ak}n_{ka}^4 l_{ka}(E_K - V)$. T-type

calcium current equation is: $I_{CaT} = g_{CaT}m_{CaT}{}^2 h_{CaT}(E_{Ca} - V)$. Transient sodium current

equation is: $I_{NaT} = g_{Na}m_{Na}^2 h_{Na}(E_{Na} - V)$. Leak current equation is: $I_L = g_L(E_L - V)$.

Simulations

Differential equations were solved using the Euler method and time step 0.05 ms. Synaptic

background activity was generated of evenly distributed excitatory postsynaptic currents

(EPSCs) and inhibitory postsynaptic currents (IPSCs) over time, to generate sparse synaptic

backgrounds as found *in vivo* (Waters et al., 2006). EPSCs and IPSCs were generated using

biexponential events: V_{Hold} = -50 mV, E_{AMPA} = 0 mV, E_{GABAA} = -60 mV and g = 0.1 nS. The time

constants are $\tau_{rise/AMPA}$ = 1 ms, $\tau_{decay/AMPA}$ = 10 ms, $\tau_{rise/GABAA}$ = 2 ms, $\tau_{decay/AMPA}$ = 20 ms.

We simulated theta oscillations at 4 Hz and analyzed four cycles of voltage traces. We generated

dendritic spikes using a leaky-integrate and fire model with resting potential -60 mV and spiking

threshold -40 mV.

Mathematical analysis of reducing/contracting and amplifying/prolonging currents

Here we used an analytical approach to determine the effect of I_{KA}, I_{CaT}, and I_{NaT} on the input

resistance and membrane time constant. It is well known that τ_m is a function of I_i:

$$\tau_m = \frac{C}{\sum_{i=1}^{n} G_i},$$ (Eq 5)

where the input conductance G_i can be obtained by differentiating Eq.2:

$$G_i = \frac{\partial I_i}{\partial v} = \bar{g}p_i q_i + \bar{g}(V - E)q_i \frac{\partial p_i}{\partial v} + \bar{g}(V - E)p_i \frac{\partial q_i}{\partial v}.$$ (Eq 6)

As discussed in the methods section, in this study all currents are inactivating. As shown in the

results section (see Slow inactivation plays nearly no role on summation during *in vivo*-like

states allowing for model reduction), all the inactivating currents produce similar voltage

changes when it is assumed fast activation/inactivation kinetics. Under these circumstances, it is

reasonable to assume $\tau_{act} = \tau_{inact} \rightarrow 0$. Consequently, p and q become functions of voltage alone,

denoted as p = p(V) and q = q(V). In this state, they lose their time dependence and are solely

voltage dependent. Specifically, $p = p^\infty(v)$ and $q = q^\infty(v)$. Therefore, when $\tau_{act} = \tau_{inact} \to 0$,

Eq 6 can be expressed as:

$$\frac{\partial I_i}{\partial v} = \bar{g}p_i^\infty q_i^\infty + \bar{g}(V-E)q_i^\infty \frac{\partial p_i^\infty}{\partial v} + \bar{g}(V-E)p_i^\infty \frac{\partial q_i^\infty}{\partial v} \qquad\qquad \text{(Eq 7)}$$

$$\frac{\partial I_i}{\partial v} = \bar{g}p_i^\infty q_i^\infty + \bar{g}(V-E)\left(q_i^\infty \frac{\partial p_i^\infty}{\partial v} + p_i^\infty \frac{\partial q_i^\infty}{\partial v}\right). \qquad\qquad \text{(Eq 8)}$$

The term $\alpha = (V-E)\left(q_i^\infty \frac{\partial p_i^\infty}{\partial v} + p_i^\infty \frac{\partial q_i^\infty}{\partial v}\right)$ can assume positive or negative values. According to

Eq. 5, when $\alpha > 0$, τ_m decreases and voltage changes are reduced, whereas when $\alpha < 0$, τ_m

increases, and voltage changes are amplified. Currents with negative α are termed

"amplifying/prolonging" as they lengthen τ_m, while those with a positive α are termed

"reducing/contracting", as they shorten τ_m (Ceballos et al., 2017b).

The question arises: under what circumstances do the currents "amplify/prolong" or

"reduce/contract"? To answer this, we need to investigate the conditions when α is positive or

negative. We know that:

$$\frac{\partial p^\infty}{\partial v} > 0, \ \frac{\partial q^\infty}{\partial v} < 0, \ p^\infty > 0, \text{and } q^\infty > 0. \qquad\qquad \text{(Eq 9)}$$

Therefore, $q_i^\infty \frac{\partial p_i^\infty}{\partial v}$ is positive and $p_i^\infty \frac{\partial q_i^\infty}{\partial v}$ is negative. We also know that the membrane potential

of dendrites typically ranges between -90 mV and +60 mV. The reversal potential for potassium

(E_k) is usually assumed to be around -90 mV, while the reversal potential for sodium and

calcium ($E_{Na,Ca}$) is assumed to be around +60 mV. Thus, for I_K ocurrs $V - E_k > 0$. Similarly, for

I_{CaT} and I_{NaT}, V- $E_{Na,Ca}$ <0.

Substituting these into Eq. 7, we find that for I_K the first term is positive, the second is positive

and the third is negative. Meanwhile, for I_{CaT} and I_{NaT}, the second is negative and the third is

positive. This demonstrates that activation and inactivation have opposite effects on τ_m: one

prolongs it while the other shortens it. This observation is consistent with the findings of Remme

et al. (2011) and Richardson et al. (2003) that I_{KA} activation reduces, and inactivation amplifies

the voltage scaling, and that I_{KA} activation shortens, and inactivation prolongs τ_m. Meanwhile,

the opposite effect occurs for I_{CaT} and I_{NaT}.

A special case arises when inactivation is removed. This occurs for some voltage-dependent

currents that exhibit either no inactivation or very slow inactivation (also known as nonactivating

currents). This scenario can be simulated by setting q = 1. Therefore, we ask: when can $\alpha =$

$(V - E)\frac{\partial p_i^\infty}{\partial v}$ be positive or negative, i.e., when does it contract or prolong τ_m, after inactivation

is removed? We can easily deduce that, $(V - E_k)\frac{\partial p_i^\infty}{\partial v} > 0$ but $\left(V - E_{Na,Ca}\right)\frac{\partial p_i^\infty}{\partial v} < 0$. This

suggests that when removing the inactivation from I_{KA}, α is positive i.e., it is always a

"contracting" current, meanwhile for I_{CaT} and I_{NaT} , α is negative i.e., it is always a "prolonging"

current.

Reduction of dendritic model

Here, we show how we approximated I_{KA}, I_{CaT}, and I_{NaT} to polynomial functions of the voltage.

Our results suggest that under *in vivo*-like conditions, where the dendritic model receives theta

oscillations and synaptic background noise, activation and inactivation variables can be

simplified to a combination of polynomial functions. This derives from the observation that

phase plots reveal inactivation variable of I_{KA} negatively correlated with its activation variable,

and the same for I_{CaT}, allowing the reduction of these two variables into one. Similarly, sodium

inactivation variable follows a power law with a 4/3 exponent of the activation variable.

Based on these findings, we propose a reduced model in which subthreshold activity is governed

by the approximated dynamics of the HH currents (I_{KA}, I_{CaT}, and I_{NaT}), while suprathreshold

activity is modeled by a fire-and-reset rule, i.e. integrate-and-fire model, which we assume as

having a threshold at -40 mV. To recapitulate, we modeled the currents using HH formalism,

where A-type potassium current equation $I_{kA} = g_{ak} n_{ka}^4 l_{ka} (E_K - V)$, T-type calcium current

equation $I_{CaT} = g_{CaT} m_{CaT}{}^2 h_{CaT} (E_{Ca} - V)$, transient sodium current equation $I_{NaT} = $

$g_{Na} m_{Na}^2 h_{Na} (E_{Na} - V)$, and leak current equation $I_L = g_L (E_L - V)$. When we apply all the

approximations, I_{KA} is expressed as a 6th order polynomial of the voltage

$I_{KA} = g_{kA}(aV^5 + bV^4 + cV^3 + dV^2 + eV + f)(E_K - V),$ (Eq. 11)

I_{CaT} as a 5th order polynomial of the voltage

$I_{CaT} = g_{CaT}(aV^4 + bV^3 + cV^2 + dV + e)(E_{Ca} - V),$ (Eq. 12)

and I_{NaT} as a power law

$I_{NaT} = g_{Na} m_{Na}^2 k m_{Na}^{-4/3}(E_{Na} - V) = g_{Na} k m_{Na}^{2/3}(E_{Na} - V)$

$= g_{Na}(aV^3 + bV^2 + cV + c)^{2/3}(E_{Na} - V)$ (Eq. 13)

The currents I_{KA} and I_{CaT} can be added in a single 6^{th} order polynomial of the voltage. The key

advantage of this approach is that it allows this equation alone to fully determine the dendrite's

voltage dynamics during *in vivo*-like conditions:

$C\frac{dv}{dt} = I_{KA} + I_{CaT} + I_{NaT} + I_L.$ (Eq. 14)

We call this dendritic model as 6^{th} degree polynomial integrate-and-fire (SIF).

References

1. Remme, M. W., & Rinzel, J. (2011). Role of active dendritic conductances in subthreshold input integration. Journal of computational neuroscience, 31, 13-30.

2. Ceballos, C. C., Roque, A. C., & Leão, R. M. (2017a). The role of negative conductances in neuronal subthreshold properties and synaptic integration. Biophysical reviews, 9, 827-834.

3. Ceballos, C. C., Roque, A. C., & Leao, R. M. (2017b). A negative slope conductance of the persistent sodium current prolongs subthreshold depolarizations. Biophysical Journal, 113(10), 2207-2217.

4. Ceballos, C. C., Pena, R. F., Roque, A. C., & Leão, R. M. (2018). Non-Decaying postsynaptic potentials and delayed spikes in hippocampal pyramidal neurons generated by a zero slope conductance created by the persistent Na+ current. Channels, 12(1), 81-88.

5. Richardson, M. J., Brunel, N., & Hakim, V. (2003). From subthreshold to firing-rate resonance. Journal of neurophysiology, 89(5), 2538-2554.

6. Liu S, Shipley MT (2008) Intrinsic conductances actively shape excitatory and inhibitory postsynaptic responses in olfactory bulb external tufted cells. J Neurosci 28:10311–10322

7. Prescott SA, De Koninck Y (2005) Integration time in a subset of spinal lamina I neurons
    is lengthened by sodium and calcium currents acting synergistically to prolong
    subthreshold depolarization. J Neurosci 25:4743–4754

8. Hoffman DA, Magee JC, Colbert CM, Johnston D (1997) K+ channel regulation of signal
    propagation in dendrites of hippocampal pyramidal neurons. Nature 387:869–875

9. Mathews PJ, Jercog PE, Rinzel J, Scott LL, Golding NL (2010) Control of submillisecond
    synaptic timing in binaural coincidence detectors by Kv1 channels. Nat Neurosci 13:601–
    609

10. Farries MA, Kita H, Wilson CJ (2010) Dynamic spike threshold and zero membrane slope
    conductance shape the response of subthalamic neurons to cortical input. J Neurosci
    30:13180–13191

11. Branco T, Tozer A, Magnus CJ, Sugino K, Tanaka S, Lee AK, Wood JN, Sternson SM (2016)
    Near-perfect synaptic integration by Na v 1.7 in hypothalamic neurons regulates body
    weight. Cell 165:1749–1761

12. Carter, B. C., Giessel, A. J., Sabatini, B. L., & Bean, B. P. (2012). Transient sodium current
    at subthreshold voltages: activation by EPSP waveforms. Neuron, 75(6), 1081-1093.

13. González-Burgos, G., & Barrionuevo, G. (2001). Voltage-gated sodium channels shape

subthreshold EPSPs in layer 5 pyramidal neurons from rat prefrontal cortex. Journal of

Neurophysiology, 86(4), 1671-1684.

14. Deleuze, C., David, F., Béhuret, S., Sadoc, G., Shin, H. S., Uebele, V. N., ... & Bal, T. (2012).

T-type calcium channels consolidate tonic action potential output of thalamic neurons to

neocortex. Journal of Neuroscience, 32(35), 12228-12236.

15. Rudolph, M., & Destexhe, A. (2003). Tuning neocortical pyramidal neurons between

integrators and coincidence detectors. Journal of computational neuroscience, 14, 239-

251.

16. Brette, R. (2015). Philosophy of the spike: rate-based vs. spike-based theories of the

brain. Frontiers in systems neuroscience, 9, 140675.

17. Hong, S., Ratté, S., Prescott, S. A., & De Schutter, E. (2012). Single neuron firing

properties impact correlation-based population coding. Journal of Neuroscience, 32(4),

1413-1428.

18. Azouz, R., & Gray, C. M. (2000). Dynamic spike threshold reveals a mechanism for

synaptic coincidence detection in cortical neurons in vivo. Proceedings of the National

Academy of Sciences, 97(14), 8110-8115.

19. Wilbers, R., Metodieva, V. D., Duverdin, S., Heyer, D. B., Galakhova, A. A., Mertens, E.
    J., ... & Goriounova, N. A. (2023). Human voltage-gated Na+ and K+ channel properties
    underlie sustained fast AP signaling. Science advances, 9(41), eade3300.

20. Qiu, J., Yang, Y., Liu, J., Zhao, W., Li, Q., Zhu, T., ... & Zhou, C. (2023). The volatile
    anesthetic isoflurane differentially inhibits voltage-gated sodium channel currents
    between pyramidal and parvalbumin neurons in the prefrontal cortex. Frontiers in Neural
    Circuits, 17, 1185095.

21. Chen, Y., Frank, H. Y., Surmeier, D. J., Scheuer, T., & Catterall, W. A. (2006).
    Neuromodulation of Na+ channel slow inactivation via cAMP-dependent protein kinase
    and protein kinase C. Neuron, 49(3), 409-420.

22. Fukaya, R., Yamada, R., & Kuba, H. (2018). Tonotopic variation of the T-type Ca2+
    current in avian auditory coincidence detector neurons. Journal of Neuroscience, 38(2),
    335-346.

23. Waters, J., & Helmchen, F. (2006). Background synaptic activity is sparse in
    neocortex. Journal of Neuroscience, 26(32), 8267-8277.

24. Bourdeau, M. L., Morin, F., Laurent, C. E., Azzi, M., & Lacaille, J. C. (2007). Kv4. 3-

mediated A-type K+ currents underlie rhythmic activity in hippocampal

interneurons. Journal of Neuroscience, 27(8), 1942-1953.

25. Rathour, R. K., Malik, R., & Narayanan, R. (2016). Transient potassium channels augment

degeneracy in hippocampal active dendritic spectral tuning. Scientific reports, 6(1),

24678.

26. Adam, Y., Kim, J. J., Lou, S., Zhao, Y., Xie, M. E., Brinks, D., ... & Cohen, A. E. (2019).

Voltage imaging and optogenetics reveal behaviour-dependent changes in hippocampal

dynamics. Nature, 569(7756), 413-417.

27. Malezieux, M., Kees, A. L., & Mulle, C. (2020). Theta oscillations coincide with sustained

hyperpolarization in CA3 pyramidal cells, underlying decreased firing. Cell Reports, 32(1).

28. Connelly, W. M., Crunelli, V., & Errington, A. C. (2016). Passive synaptic normalization and

input synchrony-dependent amplification of cortical feedback in thalamocortical neuron

dendrites. Journal of Neuroscience, 36(13), 3735-3754.

29. Gillessen, T., & Alzheimer, C. (1997). Amplification of EPSPs by low Ni2+-and amiloride-

sensitive Ca2+ channels in apical dendrites of rat CA1 pyramidal neurons. Journal of

neurophysiology, 77(3), 1639-1643.

30. Peng, H., Bian, X. L., Ma, F. C., & Wang, K. W. (2017). Pharmacological modulation of the

voltage-gated neuronal Kv7/KCNQ/M-channel alters the intrinsic excitability and

synaptic responses of pyramidal neurons in rat prefrontal cortex slices. Acta

Pharmacologica Sinica, 38(9), 1248-1256.

31. Urban, N. N., Henze, D. A., & Barrionuevo, G. (1998). Amplification of perforant-path

EPSPs in CA3 pyramidal cells by LVA calcium and sodium channels. Journal of

Neurophysiology, 80(3), 1558-1561.

32. Stuart, G., & Sakmann, B. (1995). Amplification of EPSPs by axosomatic sodium channels

in neocortical pyramidal neurons. Neuron, 15(5), 1065-1076.

33. Andreasen, M., & Lambert, J. D. (1999). Somatic amplification of distally generated

subthreshold EPSPs in rat hippocampal pyramidal neurones. The Journal of

Physiology, 519(1), 85-100.

34. Evans, R. C., Zhu, M., & Khaliq, Z. M. (2017). Dopamine inhibition differentially controls

excitability of substantia nigra dopamine neuron subpopulations through T-type calcium

channels. Journal of Neuroscience, 37(13), 3704-3720.

35. Hönigsperger, C., Marosi, M., Murphy, R., & Storm, J. F. (2015). Dorsoventral differences

in Kv7/M-current and its impact on resonance, temporal summation and excitability in

rat hippocampal pyramidal cells. The Journal of physiology, 593(7), 1551-1580.

36. Hsu, C. L., Zhao, X., Milstein, A. D., & Spruston, N. (2018). Persistent sodium current

mediates the steep voltage dependence of spatial coding in hippocampal pyramidal

neurons. Neuron, 99(1), 147-162.

37. Hage, T. A., Sun, Y., & Khaliq, Z. M. (2016). Electrical and Ca^{2+} signaling in dendritic

spines of substantia nigra dopaminergic neurons. Elife, 5, e13905.

38. Seong, H. J., Behnia, R., & Carter, A. G. (2014). Impact of subthreshold membrane

potential on synaptic responses at dendritic spines of layer 5 pyramidal neurons in the

prefrontal cortex. Journal of neurophysiology, 111(10), 1960-1972.

39. Grunditz, Å., Holbro, N., Tian, L., Zuo, Y., & Oertner, T. G. (2008). Spine neck plasticity

controls postsynaptic calcium signals through electrical compartmentalization. Journal of

Neuroscience, 28(50), 13457-13466.

40. Izhikevich, E. M. (2007). Dynamical systems in neuroscience. MIT press.

41. Prince, L. Y., Tran, M. M., Grey, D., Saad, L., Chasiotis, H., Kwag, J., ... & Richards, B. A. (2021). Neocortical inhibitory interneuron subtypes are differentially attuned to synchrony-and rate-coded information. Communications Biology, 4(1), 935.

42. Hille, B. (2001). Ion Channels of Excitable Membranes Third Edition.

43. Stuart, G. J., & Spruston, N. (2015). Dendritic integration: 60 years of progress. Nature neuroscience, 18(12), 1713-1721.

44. Engbers, J. D., Anderson, D., Asmara, H., Rehak, R., Mehaffey, W. H., Hameed, S., ... & Turner, R. W. (2012). Intermediate conductance calcium-activated potassium channels modulate summation of parallel fiber input in cerebellar Purkinje cells. Proceedings of the National Academy of Sciences, 109(7), 2601-2606.

45. Jaffe, D. B., Ross, W. N., Lisman, J. E., Lasser-Ross, N., Miyakawa, H., & Johnston, D. (1994). A model for dendritic Ca2+ accumulation in hippocampal pyramidal neurons based on fluorescence imaging measurements. Journal of Neurophysiology, 71(3), 1065-1077.

721   46. Korngreen, A., & Sakmann, B. (2000). Voltage-gated K+ channels in layer 5 neocortical

722   pyramidal neurones from young rats: subtypes and gradients. The Journal of

723   physiology, 525(3), 621-639.

725   47. Dougalis, A. G., Matthews, G. A., Liss, B., & Ungless, M. A. (2017). Ionic currents

726   influencing spontaneous firing and pacemaker frequency in dopamine neurons of the

727   ventrolateral periaqueductal gray and dorsal raphe nucleus (vlPAG/DRN): A voltage-

728   clamp and computational modelling study. Journal of Computational Neuroscience, 42,

729   275-305.

731   48. Moore, J. J., Ravassard, P. M., Ho, D., Acharya, L., Kees, A. L., Vuong, C., & Mehta, M. R.

732   (2017). Dynamics of cortical dendritic membrane potential and spikes in freely behaving

733   rats. Science, 355(6331), eaaj1497.

735   49. Ujfalussy, B. B., Makara, J. K., Lengyel, M., & Branco, T. (2018). Global and multiplexed

736   dendritic computations under in vivo-like conditions. Neuron, 100(3), 579-592.

738   50. Dembrow, N. C., & Spain, W. J. (2022). Input rate encoding and gain control in dendrites

739   of neocortical pyramidal neurons. Cell reports, 38(7).

741   51. Liao, Z., Gonzalez, K. C., Li, D. M., Yang, C. M., Holder, D., McClain, N. E., ... & Negrean, A.

742   (2024). Functional architecture of intracellular oscillations in hippocampal

743   dendrites. Nature Communications, 15(1), 6295.

52. Roome, C. J., & Kuhn, B. (2020). Dendritic coincidence detection in Purkinje neurons of
    awake mice. elife, 9, e59619.

53. Ratté, S., Hong, S., De Schutter, E., & Prescott, S. A. (2013). Impact of neuronal properties
    on network coding: roles of spike initiation dynamics and robust synchrony
    transfer. Neuron, 78(5), 758-772.

54. Stoll, A., Maier, A., Krauss, P., Gerum, R., & Schilling, A. (2023). Coincidence detection
    and integration behavior in spiking neural networks. *Cognitive Neurodynamics*, 1-13.

55. Dembrow, N. C., Zemelman, B. V., & Johnston, D. (2015). Temporal dynamics of L5
    dendrites in medial prefrontal cortex regulate integration versus coincidence detection
    of afferent inputs. *Journal of Neuroscience*, *35*(11), 4501-4514.

Fig 1. Membrane charging in the presence of the different ionic currents. Each panel has the

actual trace with (coloured) and without the current (black) (upper), after baseline subtraction

(middle), and after normalization (bottom). A. Constant pulse injected in the presence of I_{KA}. B.

Constant pulse injected in the presence of I_{NaT}. C. Constant pulse injected in the presence of I_{CaT}.

D. Actual traces of EPSP application, middle: after baseline subtraction, bottom: after

normalization. Initial resting membrane potential is -50 mV.

Fig 2. Voltage changes for dendritic ionic currents under *in vivo*-like conditions. Voltage traces

with 4 Hz theta oscillations application and spikes removed. Black solid line: passive dendrite.

Red/Green/Blue/Cyan represent first to fourth cycle of the theta sequence with the added ionic

current. A. Voltage traces of dendrite expressing only leak current plus I_{KA}. B. phase plot of

nonlinear scaling vs membrane potential. C. Voltage traces of dendrite expressing only leak

current plus I_{CaT}. D. phase plot of nonlinear scaling vs membrane potential. E. Voltage traces of

782   dendrite expressing only leak current plus I_{NaT}. F. phase plot of nonlinear scaling vs membrane

783   potential.

Fig 3. Frequency and ionic effect on voltage scaling. The nonlinear scaling is computed when

input frequency oscillations change range from 1 to 30 Hz. A. Curves for added I_{KA}. B. Curves

for added I_{CaT}. C. Curves for added I_{NaT}. D. Heatmap of the nonlinear scaling for voltages from -

70 to -40 mV vs. input frequency oscillations change from 1 to 30 Hz for I_{KA}. E. Same as in D

but for I_{CaT}. F. Same as in D but for I_{NaT}. G. Heatmap of nonlinear scaling for voltages from -70

to -40 mV vs. I_{KA} conductance. E. Same as in G but for I_{CaT} conductance. F. Same as in G but

for I_{NaT} conductance.

Fig 4. Activation vs. inactivation functional relationships. A-F. The curves represent activation

and inactivation variables under 4 Hz theta oscillations. Red/Green/Blue/Cyan represent first to

fourth cycle of the theta sequence with the added ionic current: activation for A. I_{KA}. B. I_{CaT}. C.

I_{NaT}; inactivation for D. I_{KA}. E. I_{CaT}. F. I_{NaT}. G-I: Phase plots of activation vs inactivation for G.

I_{KA}. H. I_{CaT}. I. I_{NaT}. Notice the filtering of the inactivation variable due to the slow inactivation

kinetics of I_{CaT} and I_{KA}. J-L: Linear fitting of activation vs. inactivation phase plots for J. I_{KA}. K.

I_{CaT} (magenta) and power-law fitting for L. I_{NaT}. Trajectories when $\tau_{act} = \tau_{inact} = 0$ (black).

Fig 5. Comparison of traces using actual kinetics vs. fast kinetics (i.e. $\tau_{act} = \tau_{inact} = 0.1$ ms). A.

I_{KA}. B. I_{CaT}. C. I_{NaT}. D. Average voltage difference between the two tested conditions (I_{KA} in

red, I_{CaT} in green, I_{NaT} in magenta). Comparison of traces using actual activation/inactivation

818   variables and kinetics vs. fast kinetics (i.e. $\tau_{act} = \tau_{inact} = 0$ ms) and after linearizing inactivation

819   variables using activation variable for I_{KA} and I_{CaT} and power law approximation for I_{NaT}. E. I_{KA}.

820   F. I_{CaT}. G. I_{NaT}. H. Average voltage difference between the two tested conditions (I_{KA} in red, I_{CaT}

821   in green, I_{NaT} in magenta).

Fig 6. Voltage difference between the model with slow kinetics vs. instantaneous, when input

frequency oscillations change from 1 to 30 Hz. A. I_{KA}. B. I_{CaT}. C. I_{NaT}. D. Heatmap of voltage

difference between the model with slow kinetics vs instantaneous for voltages from -70 to -40

mV vs. input frequency oscillations change from 1 to 30 Hz for I_{KA}. E. I_{CaT}. F. I_{NaT}. G. Heatmap

of the voltage difference between the model with slow kinetics vs instantaneous for voltages

from -70 to -40 mV vs. I_{KA} conductance. E. I_{CaT} conductance. F. I_{NaT} conductance.

841   Fig 7. Inactivation and activation effects on summation. The voltage traces show summation for

842   a train of 3 EPSPs. A. Model with I_{KA}. B. Model with I_{CaT}. C. Model with I_{NaT}. Initial resting

843   membrane potential is -50 mV. Top: after baseline subtraction, bottom: after normalization.

844   Three conditions were analyzed: only leak current (black), leak + inactivating current (blue), leak

845   + noninactivating current (red).

Fig 8. Synchrony- vs. rate-code and ionic currents dependency. Firing pattern under 4 Hz theta

oscillations in a dendrite expressing A. only leak current. B. dendrite expressing only leak

current plus I_{KA}. C. neuron expressing only leak current plus I_{CaT}. D. dendrite expressing only

leak current plus I_{NaT}. (E, F, G, H) Raster plots and PSTH when stimulated with sparse input vs

synchronous input.

Fig 9. Heatmaps of % relat freq computed as difference between the sparse versus synchronized synaptic inputs spanned over different conductance values (see axis). A. I_{KA} vs I_{CaT} when gNa = 0. B. I_{KA} vs I_{NaT} when gCa = 0. C. I_{NaT} vs I_{CaT} when gK = 0. D. I_{KA} vs I_{CaT} when gNa = 25. E. I_{KA} vs I_{NaT} when gCa = 0.05. F. I_{NaT} vs I_{CaT} when gK = 0.5. G. I_{KA} vs I_{CaT} when gNa = 50. H. I_{KA} vs I_{NaT} when gCa = 0.1. I. I_{NaT} vs I_{CaT} when gK = 1. The % relat freq is an indicator of whether the dendrite acts as an integrator of coincidence detector (100% for perfect coincident detector and 0% for perfect integrator).

875 Suppl Fig 1. Comparison of spiking activity using actual kinetics vs fast kinetics (i.e. $\tau_{act} = \tau_{inact} =$

876 0.1 ms). A. I_{KA}. B. I_{CaT}. C. I_{NaT}.

885   Suppl Fig 2. Heatmaps of % relat freq computed as difference between the sparse vs

886   synchronized synaptic inputs spanned over different leak conductance values (x axis) vs

887   capacitance density (y axis). The % relat freq is an indicator of whether the dendrite acts as an

888   integrator of coincidence detector (100% for perfect coincident detector and 0% for perfect

889   integrator).