The scaffolding and activation of NLRP3 on acidic vesicles depends on their biophysical properties and is independent of intermediate filaments

Guillermo de Paz Linares[1,2], Jan Mossemann[1], Blayne Sayed[1,3], and Spencer A. Freeman[1,2*]

[1]Program in Cell and Systems Biology, SickKids Research Institute, Toronto, Ontario M5G 0A4, Canada

[2]Department of Biochemistry and the [3]Institute of Medical Science, University of Toronto, Toronto, ON M5S 1A8, Canada.

*Corresponding author: spencer.freeman@sickkids.ca

Key words: vimentin, V-ATPase, LAMP1, PI(4)P, PI(3,5)P2, PI(3)P, membrane tension, membrane potential, ischemia reperfusion injury

Running title: Biophysical determinants of NLRP3 activation

Abstract

Inflammasomes are nucleated by receptors that become activated upon cellular stresses

including ionic dyshomeostasis. Rather than forming in the cytosol, recent evidence suggests

that inflammasomes are nucleated at specific sites in the cell including on cytoskeletal polymers

and the membrane surfaces of organelles. The NLRP3 inflammasome, which is formed upon

the loss of cytosolic K^+, had been proposed to form on intermediate filaments as well as on

vesicles along the endocytic pathway. To determine the necessary requirement of either

mechanism, we used vimentin knockout macrophages which do not have intermediate filaments

and compared the formation and function of NLRP3 inflammasomes. We report that vimentin

was dispensable for the activation of caspase-1, IL-1β cleavage and release, and inflammatory

responses in mice attributed to the inflammasome. Instead, NLRP3 was recruited to PI(3,5)P2,

PI(4)P- and LAMP1-positive compartments undergoing osmotic swelling. Swelling of these

compartments was dependent on the V-ATPase, the inhibition of which curtailed NLRP3

recruitment and inflammasome activation. Similarly, decreasing the hydrostatic pressure on

these vesicles prevented NLRP3 recruitment, IL-1β release and pyroptosis. The results suggest

that NLRP3 is activated by biophysical features of acidic organelles in the endocytic pathway.

Introduction

Inflammasomes are supramolecular organizing centres that assemble in response to diverse threats to cell and tissue homeostasis. Such threats often involve the loss of membrane integrity and/or the breach of the cytosol by microbial-associated molecules (Greaney, Leppla, and Moayeri 2015). The damage that triggers inflammasome activation ultimately leads to cytokine processing and release, which in turn promotes inflammatory cascades and the recruitment of immune cells. Inflammasome assembly is a stepwise process which begins with a transcriptional 'priming' step that upregulates expression of its necessary components. This is followed by the sensing of intracellular perturbances via cytosolic pattern recognition receptors and their recruitment of the effector protease, caspase-1. Once concentrated, caspase-1 initiates autolytic cleavage/activation, and subsequent processing of inflammatory mediators including IL-1β and gasdermin D (Brough and Rothwell 2007; He et al. 2015). The NLRP3 inflammasome, which has been extensively studied, is activated in response to insults that ostensibly cause the collapse of the Na^+:K^+ gradient between the cytosol and the external medium (Muñoz-Planillo et al. 2013; Perregaux and Gabel 1994; Mangan et al. 2024). Its activation is therefore initiated upon increased permeability of the plasma membrane, which can occur following exposure to diverse bacterial toxins and damage-associated molecular patterns, like extracellular ATP (Franchi and Núñez 2008; Perregaux and Gabel 1994; Laliberte, Eggler, and Gabel 1999). The accumulation of damaging particulates in the endocytic pathway, including alum and monosodium urate crystals (MSU), also triggers NLRP3 inflammasome activation (Hanfen Li et al. 2008; Hornung et al. 2008; Martinon et al. 2006), though mechanisms in this case –and whether these converge on ionic homeostasis– are less clear (Mangan et al. 2024).

Pharmacology aimed at inhibiting NLRP3 has shown enormous potential in mitigating conditions with an underlying chronic inflammatory etiology (Hao Li et al. 2022). Despite the

translational relevance of NLRP3 nucleation, recent studies on the location of NLRP3

inflammasome assembly have been seemingly at odds with one another. One report by dos

Santos et al., argued that vimentin-based intermediate filaments are required for the assembly

of the NLRP3 inflammasome in primary murine macrophages (dos Santos et al. 2015). NLRP3

was found to colocalize and co-sediment with vimentin and the loss of vimentin prevented

inflammasome assembly and inflammation. This would seem to be in contrast to other,

subsequent reports (Chen and Chen 2018; Zhang et al. 2023; Schmacke et al. 2022) that have

found that NLRP3 begins to aggregate on intracellular vesicles enriched for

phosphatidylinositol-4-phosphate (PI4P). A stretch of cationic residues found in NLRP3 is

thought to facilitate interactions with the negatively charged PI4P. This motif, known as the

polybasic region, was essential in mediating the initial assembly of the NLRP3 inflammasome in

cell lines (Chen and Chen 2018; Schmacke et al. 2022). Thus, the two models appear to be

diametrically opposed, since vimentin is not known to be required for any PI4P-mediated

processes.

In this report, we revisit the role of intermediate filaments in the inflammatory response

of macrophages, where cytosolic intermediate filaments are solely comprised of vimentin, to

NLRP3-activating stimuli. Despite reports of vimentin knockout ($Vim^{-/-}$) mice as being protected

against a plethora of inflammatory insults (Jiang et al. 2012; dos Santos et al. 2015; Mor-Vaknin

et al. 2013; Håversen et al. 2018), we do not find a role for vimentin in the nucleation of the

NLRP3 inflammasome nor the subsequent release of inflammatory cytokines from

macrophages. Supporting this, we found that $Vim^{-/-}$ mice were not protected against

inflammation in an *in vivo* model of liver ischemia reperfusion injury (IRI), an NLRP3-dependent

process (Wu et al. 2022). Therefore, it appears that the protective effects observed in $Vim^{-/-}$

mice are attributed to functions independent of NLRP3.

Instead, we find that NLRP3 is recruited to PI4P-, PI(3,5)P_2, and LAMP1-positive

vesicles undergoing osmotic swelling that appear shortly after collapsing K^+ gradients. NLRP3

recruitment to vesicles (and the resultant IL-1β release) was reversed upon relieving their

hydrostatic pressure. Rather than swelling alone being a trigger of assembly, NLRP3

inflammasome activity was dependent on the V-ATPase, implicating K^+-H^+ transport on acidic

organelles. This suggests that underappreciated biophysical features and ion transport across

endomembranes play critical roles in triggering the formation of the NLRP3-inflammasome.

Results

ASC specks form independently of vimentin

To test the proposed requirement for vimentin filaments in the nucleation of the NLRP3

inflammasomes, we derived primary macrophages from the bone marrow of *Vim$^{-/-}$* mice. Our

bone marrow-derived macrophages (BMDM) did not express vimentin as judged by western

blotting (**Fig 1A**) and immunofluorescence (**Fig 1B**). NLRP3-mediated pyroptosis is triggered by

the depletion of intracellular [K^+], including for primary macrophages (Muñoz-Planillo et al.

2013). Experimentally, this is most conveniently achieved with nigericin, a K^+ ionophore.

Alternatively, we also used ATP, which engages purinergic receptors to cause their

multimerization and formation of a non-selective pore that conducts K^+ down its concentration

gradient, i.e., out of the cell (Pelegrin and Surprenant 2006). We treated macrophages with

nigericin or ATP at standard concentrations and times known to instigate NLRP3 inflammasome

formation in a manner dependent on net K^+-efflux (Perregaux and Gabel 1994; Muñoz-Planillo

et al. 2013). Using atomic absorption spectroscopy, we found that the loss of intracellular K^+

upon the addition of nigericin or ATP was not different between wt and *Vim$^{-/-}$* BMDM, as

expected **(Fig. 1C)**. Therefore, any differences in NLRP3 inflammasome activation would not be

caused by protection against intracellular [K^+] depletion in *Vim$^{-/-}$* BMDM.

Next, we visualized ASC speck formation in lipopolysaccharide (LPS)-primed BMDM,

which upregulates the expression of NLRP3 (Bauernfeind et al. 2009), upon K^+ efflux. ASC is a

critical adaptor that enables the recruitment of caspase-1 to NLRP3 and its subsequent

activation (Man et al. 2014). We stained endogenous ASC following nigericin or ATP treatment

and, surprisingly, found no difference in the incidence of ASC-speck positive BMDM between wt

and $Vim^{-/-}$ cells **(Fig. 1D,E).** Even more remarkable, when analyzing images using a Pearson's

coefficient, we found there to be no correlation between the localization of vimentin intermediate

filaments and ASC specks following NLRP3 inflammasome activation **(Fig. 1F,G)**. This is

despite the fact that the intermediate filaments persisted in activated cells. Altogether, these

observations would seem to indicate that the assembly of the NLRP3 inflammasome is not

dependent on the presence of vimentin intermediate filaments.

Vimentin is not required for NLRP3-mediated pyroptosis

Once the NLRP3 inflammasome forms, active caspase-1 then cleaves gasdermin D

(GSDMD) and pro-IL-1β. Subsequently, the N-terminus of GSDMD inserts into the plasma

membrane to form a pore that allows for the passage and release of the processed (mature) IL-

1β (Evavold et al. 2018). The involvement of intermediate filaments in GSDMD pore assembly

was not assessed in previous studies; conceivably, if vimentin controlled the release of

proinflammatory mediators it would explain some results in mice in which KOs are protected

from inflammation (Jiang et al. 2012; Håversen et al. 2018; Mor-Vaknin et al. 2013; dos Santos

et al. 2015). We treated LPS-primed wt and $Vim^{-/-}$ BMDM with nigericin along with propidium

iodide (PI) in order to identify potential differences in the rate of plasma membrane

permeabilization **(Fig. 2A,B)**. However, we did not see appreciable differences between the

vimentin KO macrophages when compared to their wt counterparts.

Sustained GSDMD pore activity eventually leads to an irreversible form of lytic cell death

known as pyroptosis that further promotes IL-1β release into the interstitium (Fink and Cookson

2006; He et al. 2015). Thus, we investigated whether the absence of vimentin in BMDM could

prevent or stagnate pyroptosis by measuring the release of lactate dehydrogenase (LDH). In

addition to instigating pyroptosis with nigericin or ATP, we also employed alum and MSU crystals

to trigger the process. As shown by this assay, nigericin, ATP, alum, or MSU induce lytic cell

death in both wt and *Vim*^{-/-} LPS-primed BMDM **(Fig. 2C)**. Moreover, the amount of LDH released

from *Vim*^{-/-} BMDM was not significantly different than wt BMDM, irrespective of the NLRP3

inflammasome activator used. Based on our results, we conclude that *Vim*^{-/-} BMDM are not

protected against pyroptotic cell death.

Vimentin is not required for NLRP3-dependent cleavage and release of caspase-1 nor IL-1β

161   During pyroptosis, caspase-1 activation precedes the cleavage of IL-1β and has been

reported to exit the cell as an active protease following cell rupture (Laliberte, Eggler, and Gabel

1999). Indeed, we found cleaved (i.e., active) caspase-1 in the supernatants of LPS-primed

BMDM that were treated with either nigericin or ATP **(Fig. 3A)**. Remarkably, we saw that *Vim*^{-/-}

BMDM were activating and releasing comparable amounts of caspase-1 compared to their wt

BMDM counterparts upon pyroptotic stimuli **(Fig. 3A,B)**. Ultimately, IL-1β is a major effector

cytokine that results in the propagation of inflammatory responses. We found that BMDM

derived from the *Vim*^{-/-} mice had no impairment in the upregulation of pro-IL-1β upon LPS

stimulation **(Fig. 3C)** and released similar levels of active IL-1β into the supernatant after

stimulation with either nigericin or ATP **(Fig 3C)**. Furthermore, we saw no difference in the

mature IL-1β concentration found in the extracellular medium of LPS-primed BMDM treated with

nigericin or ATP, or upon lysosomal stress (i.e., alum or MSU) **(Fig. 3D)**. Overall, we did not

observe any impairment in capase-1 and IL-1β processing and release from *Vim*^{-/-} BMDM.

Vim^{-/-} *mice are not protected against hepatic ischemia/reperfusion injury*

176   Despite our *in vitro* observations, there is a plethora of studies reporting that *Vim*^{-/-} mice

display reduced tissue damage upon numerous inflammatory stimuli (Jiang et al. 2012;

Håversen et al. 2018; Mor-Vaknin et al. 2013). In fact, others have reported that *Vim$^{-/-}$* mice are

protected against cerebral IRI (Jiang et al. 2012). Thus, we probed whether this protection

translated to a hepatic IRI model, which is reportedly mediated, at least in part, by NLRP3

(Jaeschke 2003; Jiménez-Castro et al. 2019; Zhu et al. 2011). Following 1 h of ischemia and 6 h

blood reperfusion, we collected samples from mice to look at different aspects of inflammatory

injury. First, we did not observe any cohort differences in plasma levels of liver enzymes alanine

aminotransferase (ALT) and aspartate aminotransferase (AST) **(Fig. 4A,B)**, indicating that liver

damage was not ameliorated in the absence of vimentin. Our histological analyses in which we

categorized the percentage of necrotic tissue found in the livers after surgery (*see demarcated*

yellow areas in **Fig. 4C**) revealed that, surprisingly, *Vim$^{-/-}$* livers were more necrotic compared to

wt **(Fig. 4D)**. Based on these observations, we conclude that the absence of vimentin does not

confer a protective effect against the inflammatory damage associated with hepatic IRI.

NLRP3 is recruited to PI4P- and PI(3,5)P$_2$-containing vesicles

An opposing view to the role of intermediate filaments in scaffolding the NLRP3

inflammasome is that the complex is instead assembled on vesicular membranes requiring

electrostatic interactions between NLRP3 and phosphoinositides (Schmacke et al. 2022; Chen

and Chen 2018; Zhang et al. 2023). NLRP3 is reported to recognize PI3P, PI4P, and PI(3,5)P$_2$ *in*

vitro, though only its interaction with PI4P has been demonstrated in cells (Chen and Chen

2018). We therefore expressed an mCherry-NLRP3 fusion protein in HeLa cells, noting that

nigericin treatment resulted in its clear enrichment at vesicular structures over the course of 1

hour **(Fig. 5A and Supplemental movie 1)**. Using biosensors for relevant phosphoinositides,

we reperformed this experiment and used Statistical Object Distance Analysis (SODA) to

determine coincidence of the signals, which provides statistical maps of the assemblies in a

population of objects by using morphology and the distance separating objects (Lagache et al.

2018). By this method, we identified objects that were significantly coupled at a radius of 6

pixels from each object centre. We noted no appreciable association between NLRP3 and PI3P

(PX-GFP) both at 15 min and at 1 hour **(Fig. 1S)**. Interestingly, by using a recently developed

probe for PI(3,5)P_2, SnxA-GFP, we found clear association between NLRP3 and PI(3,5)P_2 15

min post nigericin treatment **(Fig. 1S)**. The association between NLRP3 and PI4P (GFP-P4M)

was higher and followed a similar trend to that of PI(3,5)P_2. As the vesicles increased in size

over time, we noted less enrichment of the phosphoinositides on these compartments, which

would be consistent with defects in membrane traffic likely occurring well after NLRP3

activation. These data corroborate the finding that NLRP3 assembly is, initially, spatially

organized around sites of enriched with PI4P and reveal the possibility that PI(3,5)P_2 plays a

role in NLRP3 activation.

Nigericin causes the swelling of LAMP1-positive vesicles that recruit NLRP3

To further characterize the relevant organelle(s) that recruit NLRP3, we co-expressed

mCherry-NLRP3 together with early endosomal (GFP-EEA1) and late endosomal (LAMP1-

mEmerald) markers. The same SODA analysis was applied which revealed that NLRP3

associates preferentially with LAMP1-positive vesicles following loss of cytosolic potassium

(Fig. 5B,C). This association was present at both 15 min and 1 h post nigericin treatment. As

with PI(4)P and PI(3,5)P_2, we noticed that the association of NLRP3 with LAMP1 vesicles once

they were swollen tended to be 'low' for their expression of LAMP1 while less swollen

compartments that associated with NLRP3 were 'high' for LAMP1 expression. At later timepoints

(i.e., 1 h) NLRP3 associated exclusively with swollen, 'low' LAMP1 membrane-bound

compartments. This could either reflect the extent to which these vesicles had undergone

swelling or the nature of the compartment itself. In contrast, we did not see appreciable NLRP3

aggregation on EEA1-positive compartments at either timepoint. This would suggest that an

initiating event in NLRP3 activation occurs at vesicular compartments containing LAMP1 that

are beginning to undergo swelling.

Increases in hydrostatic pressure and the V-ATPase are required for NLRP3 inflammasome

assembly and IL-1β release

Swelling of organelles in the endocytic pathway alone would seem unlikely to activate

NLRP3; lysosomal storage disorders or osmotic imbalances of endolysosomes do not lead to

IL-1β release or pyroptosis (Cai et al. 2024). Supporting this, incubation of cells with a non-

digestible, non-transportable disaccharide, sucrose, did not lead to the vesicular recruitment of

NLRP3 or pyroptosis, despite leading to the swelling of LAMP1 positive compartments **(Fig.**

6A,C). Vesicular swelling may nevertheless be required for NLRP3 recruitment. To test this, we

prevented the gain in hydrostatic pressure incurred upon nigericin treatment by shifting the cells

to hypertonic solutions at the time of its addition. The increase in extracellular osmolarity causes

water to leave the cell, and organelles, concomitantly (Lang et al. 1998; Hohmann 2002). Under

such conditions, we saw the expected inhibition of vesicular swelling. Importantly, preventing the

gain in hydrostatic pressure eliminated the recruitment of NLRP3 to vesicular compartments.

Instead, NLRP3 remained largely soluble **(Fig. 6B)** and pyroptosis as well as IL-1β release were

averted in BMDM **(Fig. 6C,D)**.

The swelling caused by nigericin, a membrane-permeable ionophore, may be owed to its

direct targeting of endomembranes. Once inserted into the limiting membrane of acidic

organelles, nigericin would serve to exchange H^+ for K^+, the latter of which may drive swelling. If

this is the case, inhibition of the H^+ gradient (by first inhibiting the V-ATPase) would prevent the

accumulation of K^+. We therefore inhibited the V-ATPase with concanamycin A prior to nigericin

addition. This treatment prevented the swelling of LAMP1-positive structures. Critically, this

treatment also prevented the recruitment of NLRP3 **(Fig. 6B).** Moreover, concanamycin A pre-

treatment led to a reduction in pyroptosis and IL-1β release in LPS-primed BMDM after nigericin

addition **(Fig. 6C,D)**. Taken together, we conclude that vesicles endowed with active V-ATPases

are the sites of NLRP3 scaffolding and activation upon disruptions to organellar ion transport.

Discussion

NLRP3 plays a crucial role by acting as a sensor of intracellular dyshomeostasis to then

invoke inflammation. K^+ efflux, indicative of lost plasma membrane integrity or inhibition of the

Na^+/K^+-ATPase, is a major trigger of NLRP3 inflammasome assembly (Muñoz-Planillo et al.

2013). Solid particles/crystals also activate the NLRP3 inflammasome by causing stress and

dysregulation along the endocytic pathway (Hanfen Li et al. 2008; Hari et al. 2014; Groß et al.

2016; Hornung et al. 2008). Whether or not these two, seemingly different stimuli converge

mechanistically has not been clear, but this is discussed later on below. Rather than directly

sensing differences in alkali cation (Na^+, K^+) concentrations in the cytosol, disruption of their

normal gradients leads to changes in the cell which are then sensed by NLRP3.

In this study, we demonstrate that vimentin is dispensable for NLRP3-inflammasome

assembly and instead find that the V-ATPase and vesicular swelling are required. While many

studies implicate vimentin in inflammatory cascades (Mor-Vaknin et al. 2013; Jiang et al. 2012;

Eckes et al. 2000; Håversen et al. 2018), this may be owed to impaired immune cell migration

and infiltration to inflamed tissues (Barberis et al. 2009; Battaglia et al. 2018; Ivaska et al. 2007;

Nieminen et al. 2006). In line with other studies (Zhang et al. 2023; Chen and Chen 2018), our

data show that the formation of NLRP3 inflammasomes occurs at vesicular sites enriched in

PI4P and, interestingly, $PI(3,5)P_2$. Normally these lipids are found in compartments that are

distinct from one another, which could be the case here as well since we were not able to image

both PIPs together with NLRP3. Alternatively, the immediate impact that nigericin has on

membrane traffic could cause the loss of organellar identity in the cells, causing these PIPs to

be on the same compartment.

PI4P has recently been shown to mediate other important immune processes pertaining

to intracellular danger signals. The cGAS-STING pathway, responsible for sensing cytosolic

DNA, was shown to be affected by changes to the intracellular pool of Golgi-PI4P and its

associated kinase, PI4KB (Fang et al. 2023; Luteijn et al. 2024). This suggests that the

availability of PI4P plays broad and diverse roles in facilitating innate immune responses. Under

normal conditions, the majority of PI4P is in the Golgi apparatus and the plasma membrane, as

tightly regulated by lipid kinases and phosphatases (Tan and Brill 2014). Nucleation of the

inflammasome does not occur at these sites, suggesting that while PI4P may be an essential

component of NLRP3 inflammasome activation, there are likely coincidently detected

determinants in vesicles that recruit the receptor. The location of PI4P to compartments in which

it is not normally found may therefore be an important trigger of inflammation. In contrast to

PI4P, little is known about roles for $PI(3,5)P_2$ in innate immune sensing and signaling. Lipid-

protein overlay assays previously found that NLRP3 can bind $PI(3,5)P_2$ (Chen and Chen 2018),

though the importance of this interaction remains to be tested.

Others have reported that NLRP3 can bind to vesicles along the endocytic pathway

marked with EEA1, Rab5, and/or LAMP1 (Lee et al. 2023; Zhang et al. 2023). We found NLRP3

to preferentially associate with swollen, LAMP1-low vesicles and smaller, LAMP1-high vesicles

shortly after nigericin treatment, with this association shifting towards the former over the course

of an hour. The nature of these compartments is not yet known. What is clear is that the initial

acidification of these vesicles is required for the recruitment of NLRP3. Importantly, dissipation

of the H^+ gradient needs to occur <u>before</u> the addition of nigericin to prevent NLRP3 activation,

suggesting that the activity of the V-ATPase is not directly sensed by NLRP3. Nigericin acts as

an ionophore for H^+/K^+; its insertion into the plasma membrane drives K^+ loss for H^+ gain which

is an electroneutral process. Acidosis of the cytosol is then relieved by Na^+/H^+ exchangers,

causing the cell to gain Na^+. The ionophore also readily inserts into the membranes of

organelles. In acidic organelles, the driving forces for ion transport are entirely different from that

of the plasma membrane; here, H^+ will exchange for K^+ and drive K^+ into the lumen. In addition

to the swelling caused, this also has implications for the electrical potential of the membrane. It

is therefore possible that electrical, in addition to biophysical properties of the organelle, are

then sensed by NLRP3. While K^+-efflux is a common trigger of the NLRP3 inflammasome,

agents that damage lysosomes and/or disrupt endosomal traffic such as alum, monosodium

urate crystals, and silica particles also trigger its assembly (Hari et al. 2014; Groß et al. 2016).

Whether or not these damaging agents have similar effects on these electrical/biophysical

parameters warrants systematic inquiry.

Methods and materials

Animals

Vim$^{-/-}$ (129S-*Vim*^{tm1Cba}/MesDmarkJ, stock #025692) and wt 129S1 mice (129S1/SvImJ,

strain #002448) were obtained from Jackson Labs as previously described (Colucci-Guyon et al.

1994). C57Bl/6 wt mice from Jackson Labs (strain #000664) were used where indicated. All

procedures were approved by the Animal Care Committee at the Hospital for Sick Children and

were performed in accordance with regulations from the Animals for Research Act of Ontario

and the Canadian Council on Animal Care. Mice were kept at the Animal Facility in the Peter

Gilgan Centre for Research and Learning.

Liver ischemia/reperfusion surgery

Cohorts of 8- to 12-week-old Vimentin knockout and wt (C57Bl/6) control mice were

used in a 70% segmental warm ischemia-reperfusion model (Selzner et al. 2003). In brief, a

sagittal midline laparotomy was made under isoflurane anesthesia, and an atraumatic vascular

clamp was placed on the portal vein and the hepatic artery to block blood flow to the left and

medial lobes of the liver which account for about 70% of the liver parenchyma. The clamp was

removed after 1 h to allow for reperfusion. Following 6 h of reperfusion, mice were euthanized

before blood and tissue collection for analysis. Male mice were used to control for known sex

differences in IRI responses (W. Li et al. 2018; Ostadal et al. 2019).

Reagents

Mammalian expression vectors were obtained from the following sources: 2xSidM-GFP

as described in (Levin et al. 2017); pCAG mEmerald-LAMP1 was a gift from Franck Polleux

(Addgene plasmid # 168513); Human NLRP3 from pEGFP-C2-NLRP3 (gift from Christian

Stehlik, Addgene plasmid # 73955) was inserted into an mCherry-C2 mammalian vector by

Gibson assembly cloning (NEB, cat #E2611S); pJSK659 (SNX-A-GFP) was a gift from Jason

King (Addgene plasmid # 205128); and GFP-EEA1 was a gift from Silvia Corvera (Addgene

plasmid # 42307). Primary antibodies used were rabbit anti-cleaved caspase-1 (Asp296)

(E2G2I) (Cell Signaling, cat #89332), mouse anti-ASC (BioLegend, cat #676502), rabbit anti-

vimentin (Cell Signaling, cat #5741), rabbit anti-pro caspase-1 + p10 +p12 (Abcam, cat.

#ab179515) and rabbit anti-GAPDH (Cell Signalling, cat #2118S). Primary antibody validation

was done by western blot. Secondary antibodies conjugated with AlexaFluor-405, -488, -555, -

647 and donkey anti-goat HRP secondary antibodies were obtained from Jackson

ImmunoResearch. Secondary mouse and rabbit antibodies conjugated with HRP were obtained

from Cell Signaling.

Cell culture reagents used include Dulbecco's modified eagle medium (DMEM, Wisent,

cat. #319-007-CL), fetal bovine serum (Atlanta Biologicals, cat. #S12450), Gibco™ DMEM, high

glucose, HEPES, no phenol red (FischerScientific, cat. #21063029), HBSS w/out phenol red,

calcium and magnesium, supplemented with FBS (Wisent, cat. #311-512 CL**),** 100x antibiotic-

antimycotic solution (Wisent, cat. #450-115-CL), FuGENE 6 Transfection reagent (Promega,

cat. #E2691), phosphate buffered saline (Wisent, Sigma-Aldrich, cat#311-010-CL), nigericin

sodium salt (Cayman chemicals, cat. #11437), adenosine triphosphate (pH=7.4, cat. #A2383-

5G), propidium iodide (Invitrogen, cat. #P1304MP), alum hydroxide (Invivogen, cat. #21645-51-

2), monosodium urate crystals (Invivogen, cat. #1198-77-2), lipopolysaccharide (Sigma-Aldrich,

cat. # L9764-5MG), concanamycin A (Cayman, cat. # 11050), Prolong Gold Mounting medium

with DAPI (Invitrogen, cat #P36935). .

Cell culture

Bone marrow derived macrophages from the long bones of 6-12-wk-old wt and *Vim$^{-/-}$*

mice was harvested by centrifugation and grown in DMEM + 10% FBS + 10% L929 conditioned

medium containing macrophage colony stimulating factor (MCSF) + 1% penicillin-streptomycin

at 37^{o}C and under 5% CO_2. Cells were used 5-7 days after isolation. HeLa cells were grown

and passaged in DMEM + 5% FBS + 1% penicillin-streptomycin at 37^{o}C and 5% CO_2.

Transfections

HeLa cells were seeded on top of coverslips in 12-well plates and left to attach overnight

at 37^{o}C and 5% CO_2. FuGENE6 Transfection reagent was added to room temperature serum-

free DMEM for 5 min, and then plasmid DNA was added at a 1:3 ratio of DNA (μg) to FuGENE6

(μL). Mixture was incubated at room temperature for 15 min and added to each well. Media was

changed 4 h post-transfection. Cells were transfected at ~60% confluency and incubated for at

least 18 h at 37^{o}C and under 5% CO_2 after transfection.

Intracellular [K$^+$] measurements

BMDM were grown to 90% confluency in 2 separate 6-well plates with DMEM

supplemented with 10% FBS and 10% MCSF. Following nigericin or ATP stimulation, BMDM

from one plate were washed 3x in 150 mM LiCl (pH=7.4) and lysed in 5 mM HNO_3. Cell lysates

were incubated at 4^{o}C overnight and centrifuged at 12,000 g for 15 min to remove any debris.

Samples and standards of known [K$^+$] were analyzed using a PinAAcle 900T Atomic Absorption

Spectrometer (PerkinElmer, Waltham, MA, USA) to assess the total amount of K$^+$ ions. In

parallel, BMDM from the second plate were washed as previously described and lifted in

isotonic solution. Cell concentration and volumes were determined electronically using a Z2

Coulter particle count and size analyzer (Beckman Coulter) to calculate total cell number and

volume for each sample. Intracellular concentrations of K+ were calculated by dividing the total

ion content by the total cell volume of each well.

Immunostaining

BMDM were grown to 70-80% confluency in 12-well plates with DMEM supplemented

with 10% FBS and 10% MCSF. Then, BMDM were primed with LPS (1 μg/mL, 4 h) and

subsequently treated with nigericin (10 μM, 30 min) or ATP (3 mM, 1 h). Cells were then fixed

with 4% PFA for 15 min, permeabilized in 0.1% triton for 5 min and blocked in 5% BSA

overnight. Primary and secondary antibodies were diluted to 1:100 and 1:500 in 2.5% BSA in

PBS, respectively, and added to coverslips at room temperature for 1 h. Coverslips were

washed 3 times with PBS following incubation. Slides were mounted using mounting media

containing DAPI.

PI assay

BMDM were grown to 70-80% confluency in 12 well plates with DMEM supplemented

with 10% FBS and 10% MCSF. Cells were primed with LPS (1 μg/mL) for 4 h before

simultaneously adding nigericin (10 μM) and propidium iodide (0.3 μg/mL). Pictures of each well

were taken in both brightfield and RFP setting of an EVOS microscope (ThermoFischer). At

least 3 images of different fields of view were obtained per well per timepoint. Quantification of

PI positivity was done in ImageJ using the "Cell counter" feature.

LDH assay

Cells were grown to 80-90% confluency in 12 well plates with DMEM supplemented with

10% FBS and 10% MCSF. Priming with LPS (1 μg/mL) was done for 4 h, then cells were treated

with nigericin (10 μM) for 30 min, ATP (3 mM) for 1 h, alum (100 μg/ml) for 6 h, or MSU (100

μg/ml) for 16 h in 0.1% FBS DMEM. The supernatants of each well were collected and

centrifuged at 13,000 g for 5 min, and cells in each well were washed with 1x PBS and lysed in

RIPA buffer with protease inhibitor. LDH release was assayed according to manual instructions

(CyQUANT LDH Cytotoxicity Assay, Invitrogen, cat #C20301).

Western blotting

BMDM were grown to 80-90% confluency in 6 well plates with 10% FBS and 10%

MCSF. Cells were primed with LPS (1 μg/mL) for 4 h, then treated in reduced serum medium

(0.1% FBS) with either nigericin (10 μM) for 30 min or ATP (3 mM) for 1 h. Supernatants were

prepared following the TCA precipitation protocol from (Volchuk et al. 2020). Briefly, samples

were centrifuged at 13,000 g for 5 min at 4°C to remove any debris and precipitated using 100%

TCA. Samples were then washed with ice cold acetone, then heated at 95° C, and finally

resuspended in 2x SDS-PAGE sample buffer. For the cell lysates, cells were washed with 1x D-

PBS and lysed with RIPA buffer with protease inhibitors. Lysates were centrifuged at 12,000 g

for 15 min at 4°C and prepared after determining protein concentration by BCA assay. Lysate

proteins were resolved in a 12% SDS-PAGE gel and supernatant proteins using a 15% SDS-

PAGE gel. Proteins were transferred to a PVDF membrane and immunoblotted with the

appropriate antibodies. Images were analyzed using ImageJ to determine the adjusted mean

gray intensity value of each band over the same area.

IL-1β Enzyme linked immunosorbent assay (ELISA)

BMDM were grown on 12-well plates to 80-90% confluency using DMEM supplemented

with 10% FBS and 10% MCSF. BMDM were stimulated and supernatants were collected and

centrifuged for 15 min (13,000 rpm, 4°C). Samples were diluted at a ratio of 1:5 of supernatant

to the provided dilution buffer. ELISAs were performed according to the manufacturer's

instructions (Mouse IL-1 beta ELISA kit, Abcam, cat. #ab100705). Sample absorbance was read

using a Varioskan LUX Plate Reader 1 w/Dual Injectors (Thermofischer Scientific)

Microscopy

Cells immunostained for ASC and vimentin, as well as live cell imaging was performed

by confocal microscopy using a spinning-disk Zeiss Axiovert 200 M microscope (Quorum

Technologies Inc.) with an ORCA-Fusion BT cMOS camera, and a five-line laser module with

405-, 443-, 491-, 561-, and 655-nm lines. The microscope operating software was Volocity v6.3

(Perkin Elmer). Images were acquired as Z-stacks with a 63×/1.4 NA for Vimentin/ASC images

and 25x/0.8 NA water objective for ASC-speck quantification images. For video acquisition of

NLRP3-expressing HeLa cells, a Zeiss Axio Observer microscope with a PCO camera and four-

line laser module with 405-, 443-, 491-, and 561-nm lines was used.

For live cell microscopy, coverslips were placed in a chamber with HBSS and imaged on

a pre-warmed stage and 63×/1.4 NA oil objective to 37° C. Coverslips used to acquire movies

were placed in a chamber along with phenol red-free DMEM and an environmental control

chamber set to 37° C and 5% CO_2. Movies were acquired as single-slice images over the

course of one hour.

Image analysis

Liver tissue was fixed in PFA and stained with hematoxylin and eosin. Slides were

digitalized and ischemic score was determined using the DenseNetV2 identifier available in the

HALO software. ASC-speck positivity and Pearson's correlation analysis were done using the

Volocity software over the ROI of individual cells. Pearson's correlation image thresholding was

done according to the (Costes et al. 2004) thresholding method to avoid any bias.

SODA analysis of HeLa cells expressing mCherry-NLRP3 and different probes was done

using the free image analysis software called Icy (http://icy.bioimageanalysis.org/) and using a

modified protocol from (Lagache et al. 2018). Briefly, detection of bright spots on single slice

two-channel images was calibrated using the Wavelet Spot Detector. Then, ROIs were made for

each individual cell and the statistical probability that spots from both channels were associated

with each other by random chance was calculated (as defined by $p > 0.05$). Probability was

found for different radii (r) distances from the centre of each spot. Represented in our figure is

the coupled puncta of NLRP3 within r = 6 pixels (~516 nm). For each biological replicate, 5-10

cells were analyzed using SODA per probe used.

Data analysis

All experiments and data were conducted with at least 3 biological replicates. Data is

represented as mean ± SEM or individual replicate values ± SD. Statistical significance was

determined using two-tailed unpaired Student's *t* test (significant difference threshold $p < 0.05$).

For multiple comparisons, two-way ANOVA was used with Tukey's correction for all wt vs *Vim$^{-/-}$*

analysis. All other multiple comparisons were done using a one-way ANOVA with Tukey's

correction. All experiments shown were conducted at least 3 independent times (N≥3). All data

analysis was done using GraphPad Prism 6 and Excel.

Figures

Figure 1. K$^+$-dependent ASC aggregation remains unaffected by the absence of vimentin
intermediate filaments. Validation of *Vim$^{-/-}$* BMDM by A) western blot and B) vimentin
immunostaining as compared to wt control BMDM. (C) Intracellular K$^+$ concentration
measurements of BMDM following treatment with nigericin for 15 min or ATP for 30 min. Replicate
values $\pm$ SD are shown (N=3). (D-G) wt and *Vim$^{-/-}$* BMDM primed with LPS (1 µg/mL, 4 h) and
treated with nigericin (10 µM, 30 min) or ATP (3 mM, 1 h). In (D,E), ASC was immunostained in
wt and Vim$^{-/-}$ BMDM, along with F-actin (phalloidin) and DAPI. Representative images shown in
D). Data are shown as mean of each replicate $\pm$ SEM in E (N=5). In (F,G) cells were
immunostained for vimentin and ASC in wt and *Vim$^{-/-}$* BMDM. Representative images shown in F).
(G) Pearson's correlation coefficient per cell, grey dots and mean, pink dots (N=3, minimum 7
cells per replicate).

Figure 2. Pyroptotic cell death downstream of NLRP3 inflammasome activation is not dependent
on vimentin intermediate filaments. (A,B) LPS-primed (1 μg/mL, 4 h) wt and *Vim*⁻/⁻ BMDM stained
with propidium iodide (1 μg/mL) after nigericin treatment (10 μM) over 1 h. Percent PI positivity is
displayed in B) as the mean ± SEM of each timepoint. (N=3). (C) LDH release in wt and *Vim*⁻/⁻
BMDM primed with LPS (1 μg/mL, 4 h) and treated with nigericin (10 μM) for 30 min, ATP (3 mM)
for 1 h, alum (100 μg/ml) for 6 h, or MSU (100 μg/ml) for 16 h. Data represented as replicate mean
± SD (N=3)

Figure 3. *Vim$^{-/-}$* BMDM cleave and release caspase-1 and IL-1β following NLRP3 inflammasome activation. (A-C) LPS-primed (1 μg/mL, 4 h) BMDM were treated with nigericin (10 μM, 30 min) or ATP (3 mM, 1 h). Samples were visualized by SDS-PAGE against (A,B) uncleaved and cleaved capase-1, and C) immature and mature IL-1β. In B), quantification of the ratio of cleaved to uncleaved caspase-1 normalized to wt treatments is shown. Individual values ± SD (N=4). (D) Extracellular IL-1β measured in LPS-primed wt and *Vim$^{-/-}$* BMDM treated with nigericin (10 μM) for 30 min, ATP (3 mM) for 1 h, alum (100 μg/ml) for 6 h, or MSU (100 μg/ml) for 16 h. Mature IL-1β concentration was measured by ELISA. Data represented as replicate means ± SD (minimum N=3).

Figure 4. *Vim*[-/-] mice are not ostensibly protected against hepatic ischemia/reperfusion injury. Warm ischemia/reperfusion injury was modeled using male wt (C57Bl/6) control and *Vim*[-/-] mice between 6-12 weeks of age. (A,B) Levels of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) in the serum of mice are graphed. (wt N=5, Vim[-/-] N=6 mice per group.) (C) Hematoxylin and eosin (H&E) staining of liver tissue samples along with Halo DenseNetV2 mask to detect healthy and necrotic areas. (D) Quantification of percent necrosis of liver samples (wt N=6, Vim[-/-] N=7 mice per group). All data shown as biological replicates ± SD.

Figure 5. NLRP3 aggregates on LAMP1-positive vesicular compartments. (A) HeLa cells expressing mCherry-NLRP3 before and after treatment with nigericin (10 µM, 15 min and 1h). (B) Co-expression of mCherry-NLRP3 with GFP-EEA1 or LAMP1-mEmerald in nigericin-treated cells after 15 min or 1 h. (C) SODA analysis for object association. Data represented as distribution of percent NLRP3 objects coupled with EEA1- or LAMP1-positive objects. (N=3, with 5-10 cells per technical replicate). Data shown as replicate means and individual technical replicate values.

Figure 6. NLRP3 recruitment and activation are dependent on the activity of the V-ATPase. (A) HeLa cells co-expressing mCherry-NLRP3 and LAMP1-mEmerald following treatment with either nigericin (10 µM, 1 h) or sucrose (30 mM, 16 h). (B) HeLa cells co-expressing LAMP1-mEmerald and mCherry-NLRP3 treated with nigericin in either isotonic (300 mOsm) or hypertonic (680 mOsm) medium, or pre-treated with concanamycin A (1 µM, 30 min) prior to nigericin treatment in isotonic medium. (C, D) LPS-primed BMDM treated with nigericin (10 µM, 30 min), sucrose, or concanamycin A (1 µM, 30 min) followed by nigericin. In C), the percent PI positive cells normalized to LPS + nigericin treatment is shown (N=3). In D) the release of IL-1β measured by ELISA and normalized to LPS + nigericin treatment is shown (N=4). All data shown as individual replicate values ± SD.

595
Supplemental Figure 1. NLRP3 associates with PI4P- and PI(3,5)P$_2$-positive vesicles following
nigericin treatment. (A) SODA analysis for object association between NLRP3-mCherry and PX-
GFP, SnxA-GFP, or GFP-P4M phosphoinositide probes following nigericin treatment (10 µM, 15
min and 1h). Data represented as distribution of percent NLRP3 objects coupled with
phosphoinositide probe objects. (N=3, with 5-10 cells per technical replicate). All data shown as
replicate means and individual technical replicate values.

602
603

References

Barberis, Laura, Christian Pasquali, Dominique Bertschy-Meier, Alessandra Cuccurullo, Carlotta Costa, Chiara Ambrogio, Francis Vilbois, et al. 2009. "Leukocyte Transmigration Is Modulated by Chemokine-Mediated PI3Kγ-Dependent Phosphorylation of Vimentin." *European Journal of Immunology* 39 (4): 1136–46. https://doi.org/10.1002/eji.200838884.

Battaglia, Rachel A., Samed Delic, Harald Herrmann, and Natasha T. Snider. 2018. "Vimentin on the Move: New Developments in Cell Migration." *F1000Research* 7 (November):1796. https://doi.org/10.12688/f1000research.15967.1.

Bauernfeind, Franz G., Gabor Horvath, Andrea Stutz, Emad S. Alnemri, Kelly MacDonald, David Speert, Teresa Fernandes-Alnemri, et al. 2009. "Cutting Edge: NF-κB Activating Pattern Recognition and Cytokine Receptors License NLRP3 Inflammasome Activation by Regulating NLRP3 Expression." *The Journal of Immunology* 183 (2): 787–91. https://doi.org/10.4049/jimmunol.0901363.

Brough, David, and Nancy J. Rothwell. 2007. "Caspase-1-Dependent Processing of pro-Interleukin-1β Is Cytosolic and Precedes Cell Death." *Journal of Cell Science* 120 (5): 772–81. https://doi.org/10.1242/jcs.03377.

Cai, Ruiqi, Ori Scott, Gang Ye, Trieu Le, Ekambir Saran, Whijin Kwon, Subothan Inpanathan, et al. 2024. "Pressure Sensing of Lysosomes Enables Control of TFEB Responses in Macrophages." *Nature Cell Biology* 26 (8): 1247–60. https://doi.org/10.1038/s41556-024-01459-y.

Chen, Jueqi, and Zhijian J. Chen. 2018. "PtdIns4P on Dispersed Trans-Golgi Network Mediates NLRP3 Inflammasome Activation." *Nature* 564 (7734): 71–76. https://doi.org/10.1038/s41586-018-0761-3.

Colucci-Guyon, Emma, Marie-Madeleine Portier, Irene Dunia, Denise Paulin, Sandrine Pournin, and Charles Babinet. 1994. "Mice Lacking Vimentin Develop and Reproduce without an Obvious Phenotype." *Cell* 79 (4): 679–94. https://doi.org/10.1016/0092-8674(94)90553-3.

Costes, Sylvain V., Dirk Daelemans, Edward H. Cho, Zachary Dobbin, George Pavlakis, and Stephen Lockett. 2004. "Automatic and Quantitative Measurement of Protein-Protein Colocalization in Live Cells." *Biophysical Journal* 86 (6): 3993–4003. https://doi.org/10.1529/biophysj.103.038422.

Eckes, B., E. Colucci-Guyon, H. Smola, S. Nodder, C. Babinet, T. Krieg, and P. Martin. 2000. "Impaired Wound Healing in Embryonic and Adult Mice Lacking Vimentin." *Journal of Cell Science* 113 (13): 2455–62. https://doi.org/10.1242/jcs.113.13.2455.

Evavold, Charles L., Jianbin Ruan, Yunhao Tan, Shiyu Xia, Hao Wu, and Jonathan C. Kagan. 2018. "The Pore-Forming Protein Gasdermin D Regulates Interleukin-1 Secretion from Living Macrophages." *Immunity* 48 (1): 35-44.e6. https://doi.org/10.1016/j.immuni.2017.11.013.

Fang, Run, Qifei Jiang, Xinying Jia, and Zhengfan Jiang. 2023. "ARMH3-Mediated Recruitment of PI4KB Directs Golgi-to-Endosome Trafficking and Activation of the Antiviral Effector STING." *Immunity* 56 (3): 500-515.e6. https://doi.org/10.1016/j.immuni.2023.02.004.

Fink, Susan L., and Brad T. Cookson. 2006. "Caspase-1-Dependent Pore Formation during Pyroptosis Leads to Osmotic Lysis of Infected Host Macrophages." *Cellular Microbiology* 8 (11): 1812–25. https://doi.org/10.1111/j.1462-5822.2006.00751.x.

Franchi, Luigi, and Gabriel Núñez. 2008. "The Nlrp3 Inflammasome Is Critical for Aluminium Hydroxide-Mediated IL-1β Secretion but Dispensable for Adjuvant Activity." *European Journal of Immunology* 38 (8): 2085–89. https://doi.org/10.1002/eji.200838549.

Greaney, Allison J., Stephen H. Leppla, and Mahtab Moayeri. 2015. "Bacterial Exotoxins and the Inflammasome." *Frontiers in Immunology* 6 (November). https://doi.org/10.3389/fimmu.2015.00570.

Groß, Christina J., Ritu Mishra, Katharina S. Schneider, Guillaume Médard, Jennifer Wettmarshausen, Daniela C. Dittlein, Hexin Shi, et al. 2016. "K + Efflux-Independent NLRP3 Inflammasome Activation by Small Molecules Targeting Mitochondria." *Immunity* 45 (4): 761–73. https://doi.org/10.1016/j.immuni.2016.08.010.

Hari, Aswin, Yifei Zhang, Zhongyuan Tu, Pascal Detampel, Melanie Stenner, Anutosh Ganguly, and Yan Shi. 2014. "Activation of NLRP3 Inflammasome by Crystalline Structures via Cell Surface Contact." *Scientific Reports* 4 (1): 7281. https://doi.org/10.1038/srep07281.

Håversen, Liliana, Jeanna Perman Sundelin, Adil Mardinoglu, Mikael Rutberg, Marcus Ståhlman, Ulrika Wilhelmsson, Lillemor Mattsson Hultén, et al. 2018. "Vimentin Deficiency in Macrophages Induces Increased Oxidative Stress and Vascular Inflammation but Attenuates Atherosclerosis in Mice." *Scientific Reports* 8 (1): 16973. https://doi.org/10.1038/s41598-018-34659-2.

He, Wan-ting, Haoqiang Wan, Lichen Hu, Pengda Chen, Xin Wang, Zhe Huang, Zhang-Hua Yang, Chuan-Qi Zhong, and Jiahuai Han. 2015. "Gasdermin D Is an Executor of Pyroptosis and Required for Interleukin-1β Secretion." *Cell Research* 25 (12): 1285–98. https://doi.org/10.1038/cr.2015.139.

Hohmann, Stefan. 2002. "Osmotic Stress Signaling and Osmoadaptation in Yeasts." *Microbiology and Molecular Biology Reviews* 66 (2): 300–372. https://doi.org/10.1128/MMBR.66.2.300-372.2002.

Hornung, Veit, Franz Bauernfeind, Annett Halle, Eivind O Samstad, Hajime Kono, Kenneth L Rock, Katherine A Fitzgerald, and Eicke Latz. 2008. "Silica Crystals and Aluminum Salts Activate the NALP3 Inflammasome through Phagosomal Destabilization." *Nature Immunology* 9 (8): 847–56. https://doi.org/10.1038/ni.1631.

Ivaska, Johanna, Hanna-Mari Pallari, Jonna Nevo, and John E. Eriksson. 2007. "Novel Functions of Vimentin in Cell Adhesion, Migration, and Signaling." *Experimental Cell Research* 313 (10): 2050–62. https://doi.org/10.1016/j.yexcr.2007.03.040.

Jaeschke, Hartmut. 2003. "Molecular Mechanisms of Hepatic Ischemia-Reperfusion Injury and Preconditioning." *American Journal of Physiology-Gastrointestinal and Liver Physiology* 284 (1): G15–26. https://doi.org/10.1152/ajpgi.00342.2002.

Jiang, Susan X., Jacqueline Slinn, Amy Aylsworth, and Sheng T. Hou. 2012. "Vimentin Participates in Microglia Activation and Neurotoxicity in Cerebral Ischemia: Vimentin Controls Microglia Activation." *Journal of Neurochemistry* 122 (4): 764–74. https://doi.org/10.1111/j.1471-4159.2012.07823.x.

Jiménez-Castro, Mónica B., María Eugenia Cornide-Petronio, Jordi Gracia-Sancho, and Carmen Peralta. 2019. "Inflammasome-Mediated Inflammation in Liver Ischemia-Reperfusion Injury." *Cells* 8 (10): 1131. https://doi.org/10.3390/cells8101131.

Lagache, Thibault, Alexandre Grassart, Stéphane Dallongeville, Orestis Faklaris, Nathalie Sauvonnet, Alexandre Dufour, Lydia Danglot, and Jean-Christophe Olivo-Marin. 2018. "Mapping Molecular Assemblies with Fluorescence Microscopy and Object-Based Spatial Statistics." *Nature Communications* 9 (1): 698. https://doi.org/10.1038/s41467-018-03053-x.

Laliberte, Ronald E., James Eggler, and Christopher A. Gabel. 1999. "ATP Treatment of Human Monocytes Promotes Caspase-1 Maturation and Externalization." *Journal of Biological Chemistry* 274 (52): 36944–51. https://doi.org/10.1074/jbc.274.52.36944.

Lang, Florian, Gillian L. Busch, Markus Ritter, Harald Völkl, Siegfried Waldegger, Erich Gulbins, and Dieter Häussinger. 1998. "Functional Significance of Cell Volume Regulatory Mechanisms." *Physiological Reviews* 78 (1): 247–306. https://doi.org/10.1152/physrev.1998.78.1.247.

Lee, Bali, Christopher Hoyle, Rose Wellens, Jack P. Green, Fatima Martin-Sanchez, Daniel M. Williams, Billie J. Matchett, et al. 2023. "Disruptions in Endocytic Traffic Contribute to the Activation of the NLRP3 Inflammasome." *Science Signaling* 16 (773): eabm7134. https://doi.org/10.1126/scisignal.abm7134.

Li, Hanfen, Stephen B. Willingham, Jenny P.-Y. Ting, and Fabio Re. 2008. "Cutting Edge: Inflammasome Activation by Alum and Alum's Adjuvant Effect Are Mediated by NLRP3." *Journal of Immunology (Baltimore, Md.: 1950)* 181 (1): 17–21. https://doi.org/10.4049/jimmunol.181.1.17.

Li, Hao, Yanling Guan, Bo Liang, Peng Ding, Xin Hou, Wei Wei, and Yang Ma. 2022. "Therapeutic Potential of MCC950, a Specific Inhibitor of NLRP3 Inflammasome." *European Journal of Pharmacology* 928 (August):175091. https://doi.org/10.1016/j.ejphar.2022.175091.

Li, Wujun, Dong Li, Liankang Sun, Zeyu Li, Liang Yu, and Shengli Wu. 2018. "The Protective Effects of Estrogen on Hepatic Ischemia-Reperfusion Injury in Rats by Downregulating the Ang II/AT1R Pathway." *Biochemical and Biophysical Research Communications* 503 (4): 2543–48. https://doi.org/10.1016/j.bbrc.2018.07.013.

Luteijn, Rutger D., Sypke R. Van Terwisga, Jill E. Ver Eecke, Liberty Onia, Shivam A. Zaver, Joshua J. Woodward, Richard W. Wubbolts, David H. Raulet, and Frank J. M. Van Kuppeveld. 2024. "The Activation of the Adaptor Protein STING Depends on Its Interactions with the Phospholipid PI4P." *Science Signaling* 17 (827): eade3643. https://doi.org/10.1126/scisignal.ade3643.

Man, Si Ming, Lee J. Hopkins, Eileen Nugent, Susan Cox, Ivo M. Glück, Panagiotis Tourlomousis, John A. Wright, Pietro Cicuta, Tom P. Monie, and Clare E. Bryant. 2014. "Inflammasome Activation Causes Dual Recruitment of NLRC4 and NLRP3 to the Same Macromolecular Complex." *Proceedings of the National Academy of Sciences* 111 (20): 7403–8. https://doi.org/10.1073/pnas.1402911111.

Mangan, Matthew S.J., Anil Akbal, Sophie Rivara, Agnes Schröder, Gabor Horvath, Philipp Walch, Jan Hegermann, et al. 2024. "Cytosolic Sodium Accumulation Is a Common Danger Signal Triggering Endocytic Dysfunction and NLRP3 Inflammasome Activation." https://doi.org/10.1101/2024.03.26.586774.

Martinon, Fabio, Virginie Pétrilli, Annick Mayor, Aubry Tardivel, and Jürg Tschopp. 2006. "Gout-Associated Uric Acid Crystals Activate the NALP3 Inflammasome." *Nature* 440 (7081): 237–41. https://doi.org/10.1038/nature04516.

Mor-Vaknin, Nirit, Maureen Legendre, Yue Yu, Carlos H. C. Serezani, Sanjay K. Garg, Anna Jatzek, Michael D. Swanson, et al. 2013. "Murine Colitis Is Mediated by Vimentin." *Scientific Reports* 3 (1): 1045. https://doi.org/10.1038/srep01045.

Muñoz-Planillo, Raúl, Peter Kuffa, Giovanny Martínez-Colón, Brenna L. Smith, Thekkelnaycke M. Rajendiran, and Gabriel Núñez. 2013. "K+ Efflux Is the Common Trigger of NLRP3 Inflammasome Activation by Bacterial Toxins and Particulate Matter." *Immunity* 38 (6): 1142–53. https://doi.org/10.1016/j.immuni.2013.05.016.

Nieminen, Mikko, Tiina Henttinen, Marika Merinen, Fumiko Marttila–Ichihara, John E. Eriksson, and Sirpa Jalkanen. 2006. "Vimentin Function in Lymphocyte Adhesion and Transcellular Migration." *Nature Cell Biology* 8 (2): 156–62. https://doi.org/10.1038/ncb1355.

Ostadal, B., Z. Drahota, J. Houstek, M. Milerova, I. Ostadalova, M. Hlavackova, and F. Kolar. 2019. "Developmental and Sex Differences in Cardiac Tolerance to Ischemia–Reperfusion Injury: The Role of Mitochondria." *Canadian Journal of Physiology and Pharmacology* 97 (9): 808–14. https://doi.org/10.1139/cjpp-2019-0060.

Pelegrin, Pablo, and Annmarie Surprenant. 2006. "Pannexin-1 Mediates Large Pore Formation and Interleukin-1β Release by the ATP-Gated P2X7 Receptor." *The EMBO Journal* 25 (21): 5071–82. https://doi.org/10.1038/sj.emboj.7601378.

Perregaux, D., and C. A. Gabel. 1994. "Interleukin-1 Beta Maturation and Release in Response to ATP and Nigericin. Evidence That Potassium Depletion Mediated by These Agents Is a Necessary and Common Feature of Their Activity." *The Journal of Biological Chemistry* 269 (21): 15195–203.

Santos, Gimena dos, Micah R. Rogel, Margaret A. Baker, James R. Troken, Daniela Urich, Luisa Morales-Nebreda, Joseph A. Sennello, et al. 2015. "Vimentin Regulates Activation of the NLRP3 Inflammasome." *Nature Communications* 6 (1): 6574. https://doi.org/10.1038/ncomms7574.

Schmacke, Niklas A., Fionan O'Duill, Moritz M. Gaidt, Inga Szymanska, Julia M. Kamper, Jonathan L. Schmid-Burgk, Sophia C. Mädler, et al. 2022. "IKKβ Primes Inflammasome Formation by Recruiting NLRP3 to the Trans-Golgi Network." *Immunity* 55 (12): 2271-2284.e7. https://doi.org/10.1016/j.immuni.2022.10.021.

Selzner, Nazia, Markus Selzner, Wolfram Jochum, and Pierre-Alain Clavien. 2003. "Ischemic Preconditioning Protects the Steatotic Mouse Liver against Reperfusion Injury: An ATP Dependent Mechanism." *Journal of Hepatology* 39 (1): 55–61. https://doi.org/10.1016/S0168-8278(03)00147-8.

Tan, Julie, and Julie A. Brill. 2014. "Cinderella Story: PI4P Goes from Precursor to Key Signaling Molecule." *Critical Reviews in Biochemistry and Molecular Biology* 49 (1): 33–58. https://doi.org/10.3109/10409238.2013.853024.

Volchuk, Allen, Anna Ye, Leon Chi, Benjamin E. Steinberg, and Neil M. Goldenberg. 2020. "Indirect Regulation of HMGB1 Release by Gasdermin D." *Nature Communications* 11 (1): 4561. https://doi.org/10.1038/s41467-020-18443-3.

Wu, Tong, Cheng Zhang, Tianfeng Shao, Jianzhong Chen, and Diyu Chen. 2022. "The Role of NLRP3 Inflammasome Activation Pathway of Hepatic Macrophages in Liver Ischemia–

785   Reperfusion Injury." *Frontiers in Immunology* 13 (June):905423.
786   https://doi.org/10.3389/fimmu.2022.905423.
Zhang, Zhirong, Rossella Venditti, Li Ran, Zengzhen Liu, Karl Vivot, Annette Schürmann, Juan S.
Bonifacino, Maria Antonietta De Matteis, and Romeo Ricci. 2023. "Distinct Changes in
Endosomal Composition Promote NLRP3 Inflammasome Activation." *Nature*
Immunology 24 (1): 30–41. https://doi.org/10.1038/s41590-022-01355-3.
Zhu, Ping, Lihua Duan, Jie Chen, Ali Xiong, Qin Xu, Hongwei Zhang, Fang Zheng, Zheng Tan, Feili
Gong, and Min Fang. 2011. "Gene Silencing of NALP3 Protects Against Liver Ischemia–
Reperfusion Injury in Mice." *Human Gene Therapy* 22 (7): 853–64.
https://doi.org/10.1089/hum.2010.145.
795