Mutualism mediates legume response to microbial climate legacies

<u>Abstract</u>

Climate change is altering both soil microbial communities and the ecological context of plant-
microbe interactions. Heat, drought, and their legacies can alter soil microbiomes and potential
plant symbionts, but the direct consequences of these microbial changes on plant performance
and plant investment in symbiosis remain underexplored. Predicting how soil microbes modulate
plant resilience to heat and drought is critical to mitigating the negative effects of climate change
on ecosystems and agriculture. In this study, we conducted growth chamber experiments to
isolate the microbially-mediated indirect effects of heat and drought on plant performance and
symbiosis. In the first experiment, focused on drought, we found that drought and drought-
treated microbes, along with their interaction, significantly decreased the biomass of *Medicago*
lupulina plants compared to well-watered microbiomes and conditions. In a second experiment,
we then tested how the addition of a well-known microbial mutualist, *Sinorhizobium meliloti*,
affected heat- and drought-treated microbiomes' impact on *M. lupulina*. We found that drought-
adapted microbiomes negatively impacted legume performance by increasing mortality and
reducing branch number, but that adding rhizobia erased differences in plant response to climate
treated soils. In contrast, heat-adapted microbiomes did not differ significantly from control
microbiomes in their effects on a legume. Our results suggest microbial legacy effects, mutualist
partners, and their interactions are important in mediating plant responses to drought, with some
mutualists equalizing plant responses across microbial legacies.

<u>Introduction</u>

How plant-microbe symbioses respond to climate change is hard to predict (Classen et al. 2015,
Rudgers et al. 2020) because plants and microbes are each affected by climate directly and
indirectly through their interactions (Figure 1) (Allison and Treseder 2008, Castro et al. 2010,
Sheik et al. 2011, Franklin et al. 2016, Melillo et al. 2017, Zhou et al. 2020, Pec et al. 2021,
Seaton et al. 2021). Plants and microbes cycle nutrients such as nitrogen and phosphorus, often
while engaged in symbiosis, so if these relationships are altered by components of climate
change like excessive heat and drought, ecosystem function may be altered (Kardol et al. 2010).
In addition, because microbes can strongly affect plant fitness, outcomes of symbiosis are critical

for predicting plant performance and evolution under climate change (Classen et al. 2015,

Hawkes et al. 2020). Here, we experimentally tested the microbially-mediated indirect effects of

heat and drought on plant performance and symbiosis.

Soil microbiomes respond to heat and drought through changes in microbial diversity and

community composition (Allison and Treseder 2008, Castro et al. 2010, Melillo et al. 2017, Pec

et al. 2021), and the response of symbiotic microbes may not be in the same direction or of the

same magnitude as in the broader microbial community (Boyle et al. 2024). Changes in a soil's

microbial community composition can alter rhizosphere microbiomes and the benefit that

rhizosphere microbes provide to plants (Rubin et al. 2018, Lu et al. 2018, Fitzpatrick et al. 2019)

with reproducible effects (Panke-Buisse et al. 2015), although elucidating the links between

climate-driven changes in the soil and changes in the rhizosphere's benefit is still needed. In

addition to shifts in community composition, soil microbes may also respond to climate stressors

via within-strain evolution or plasticity (Sachs et al. 2011, Simonsen 2021), and potentially push

plant-microbe interactions along the mutualism-parasitism continuum (Davis et al. 2015, Shantz

et al. 2016, Rudgers et al. 2020, González et al. 2021). Whether multiple climate stressors have

compounding effects on soil microbiomes and plant-microbe symbioses is infrequently tested

(Kivlin et al. 2013), but remains important given that multiple climatic axes are changing

simultaneously (Thuiller 2007).

Determining the importance and scale of direct versus indirect effects of climate on plant-

microbe symbioses is challenging if both direct and indirect effects are occurring at the same

time. To address indirect effects only, studies can leverage the microbial legacy effects left

behind by climate stressors (Averill et al. 2016, Martiny et al. 2017, de Nijs et al. 2019). Most

studies examining the effects of climate-treated microbes suggest that they tend to improve plant

performance in conditions that match those they evolved in (Lau and Lennon 2012, Allsup et al.

2023, Ricks and Yannarell 2023); these studies demonstrate that indirect effects of climate

change on plants, mediated through microbial symbionts, may contribute to plant performance

under changing climates. Yet shifts in the microbiome due to climate change have the potential

to be either beneficial or detrimental to plants. On the one hand, microbiomes may facilitate a

faster adaptive response than relying solely on host evolution, since microbiomes can rapidly

change in composition (Copeland et al. 2015, Sepulveda and Moeller 2020) and this may offer

new or more robust beneficial functions to the host (Kolodny and Schulenburg 2020). On the

other hand, changes in the microbiome could also lead to an increase of mismatched or harmful

symbiotic partners (Delgado-Baquerizo et al. 2020).

In a previous study, Boyle et al. (2024) factorially implemented heat and drought treatments on

controlled field plots, and characterized changes in the communities of potentially symbiotic

microbes found in the bulk soil. Known bacterial symbionts showed a strong response to climate

treatments, including rhizobia, which form a specialized structure on legume roots called a

nodule, where they fix nitrogen in exchange for fixed carbon (Beringer et al. 1979). Drought

increased the diversity of both rhizobia and plant beneficial bacteria (PBB) and shifted

phytopathogen community composition. The increased diversity of rhizobia and PBBs could

mean that there are more potential strains for a plant to associate with, however given the high

specificity of legume-rhizobium interactions (Chen et al. 2021), increased diversity could also

make it harder for legumes to find compatible, high-quality partners (Burghardt and diCenzo

2023). Fungi also had strong community-wide responses to climate treatments, with mycorrhizal

abundance decreasing in heat-treated plots and parasitic fungi increasing in drought-treated plots

(Boyle et al. 2024). Mycorrhizae can increase plant stress tolerance (Quiroga et al. 2017;

Kakouridis et al. 2022), provide additional nutrients (Koide 1991), and act as 'fungal highways'

for bacteria to travel along to the plant root (Zhang et al. 2020). Thus, the previous results

suggest heat and drought-induced variation in the soil microbiome could have a variety of

consequences for plant performance.

The role of mutualism in buffering plants against the effects of climate change remains an open

empirical question (Keeler et al. 2021, Duarte and Maherali 2022). If mutualistic symbionts

stabilize plant fitness during times of fluctuating environment or microbiome quality (Mendes et

al. 2011, Porter and Sachs 2020), this could affect how plants evolve and respond to climate

change. Additionally, the presence of beneficial microbes in climate-treated soils may alter the

magnitude of microbially-mediated indirect effects of climate change on plants, although to our

knowledge, this has not been tested. How mutualisms buffer partners against climate change also

has implications for the maintenance of mutualism (Kiers et al. 2010). For example, given

stressful conditions, mutualists may lose one or more partners, switch to antagonism or new

partners, or abandon mutualism altogether (Sachs and Simms 2006, Kiers et al. 2010). Whether

heat or drought facilitates any of these outcomes may be detectable in clearly defined

partnerships, such as the nutritional mutualism between legumes and rhizobia.

We experimentally tested the indirect effects of heat and drought on legumes mediated through

changes in the soil microbiome, and whether the addition of rhizobia buffered legumes against

these microbiome effects. Given the large impact of drought on the soil microbial community

(Boyle et al. 2024), we first tested the effect of control and drought-treated microbes under both

drought and well-watered conditions. We predicted that drought-treated microbiomes would

improve legume tolerance of drought stress compared to control microbiomes, since microbes in

the soil should be robust to drought, but that drought-treated microbes would be worse for plants

under well-watered conditions. Then, in a second experiment, we added soil from the plots that

experienced heat, drought, both, or neither to legumes in a growth chamber with and without

added rhizobia, and quantified plant performance and plant investment in symbiosis. We

predicted that adding climate-treated microbes to plants would decrease plant performance and

weaken plant investment into symbiosis, but that adding rhizobia would buffer against these

effects. Within this second experiment, we used 16S and ITS amplicon sequencing of

rhizosphere and nodule microbes, in combination with previously published 16S and ITS data on

the climate-treated bulk soil microbial communities used as inocula, to understand the impact of

heat and drought legacy on plant microbiomes.

<u>Methods</u>

Study system

The temperature-free-air-controlled enhancement (T-FACE) experiment is in an old field at

Koffler Scientific Reserve (KSR, www.ksr.utoronto.ca) in Ontario, Canada (44°01'48"N,

79°32'01"W). The experimental design, treatment effectiveness, and soil microbiome analysis

are fully described in Boyle et al. (2024), but we give a brief overview here. Plots in the T-FACE

experiment grew only seedlings of white spruce, *Picea glauca*, and were either heated,

droughted, heated and droughted, or ambient (3 plots/treatment). Boyle et al. (2024) applied

treatments during the growing seasons of 2020 and 2021, with rainout structures present for 8

months and heaters activated for 9 months. During active treatment, the mean soil temperature of

heated plots was 3.7 ▢ or 3.6 ▢ hotter than un-heated plots in 2020 and 2021, respectively. In

2020, the mean soil volumetric water content (VWC) during droughts was 0.28 (m^3/m^3) in non-

drought plots and 0.25 (m^3/m^3) in drought plots, and in 2021, mean VWC was 0.26 (m^3/m^3) in

non-drought plots and 0.21 (m^3/m^3) in drought plots. We collected sifted bulk soil (passed

through a 4.75 mm metal sieve) from each plot on June 5th, 2022, sterilizing tools between each

collection. We kept soil bags stored slightly open in the dark at 4 ☐ until application to the

experimental legumes.

We used seeds of a single genotype of *Medicago lupulina*, an annual legume that forms

indeterminate root nodules with rhizobia in the genus *Sinorhizobium* (Batstone et al. 2020b).

Seeds were collected from KSR and selfed for two generations in the University of Toronto

greenhouses (Simonsen and Stinchcombe 2014). *Medicago lupulina* is naturalized at KSR near

the experimental warming array, and its growing season peaks in June. The growth chamber

experiments were not designed to mimic ecological interactions in nature, but rather to test the

general hypothesis that soil microbial changes, including known changes in rhizobia in response

to heat and drought (Boyle et al. 2024), can by themselves affect plant performance and

symbiosis. We chose the legume *M. lupulina* for experimental tractability and because it partakes

in multiple microbial symbioses, including symbiosis with rhizobia and arbuscular mycorrhizal

fungi.

Experiment 1: Effect of drought-treated microbes on legume performance in drought

Experimental design and data collection

We tested whether drought-treated microbes affected legumes in both dry and well-watered soil

conditions, because past work showed that drought was particularly important for shifting

microbial community composition (Boyle et al. 2024). We implemented a 2×2 factorial design

with plants receiving biotic (e.g. live) drought soil or biotic control soil, and experiencing a

terminal drought or well-watered conditions, with 30 replicates per treatment, for a total of 120

plants. We pooled soil within treatment types to generate inocula. We surface-sterilized seeds

with bleach and ethanol then stratified them for two weeks at 4 °C. We planted them into

Magenta boxes containing autoclaved Profile® fine-grained turface and a nylon rope wick,

spatially blocked by light canopy in the growth chamber. The experiment took place in a growth

chamber room set to cycle between 22 °C day/20 °C night with 16 hours of light.

Upon planting, we applied 10 ml of soil to each plant. We added soil directly to maintain the

integrity of fungal structures and the diversity of microbes. Additionally, we applied 1 ml of low-

nitrogen fertilizer (Supplemental material) to each plant, and we reapplied 1 ml of fertilizer every

4 weeks. All pots were covered with plastic wrap to maintain moisture, and when covered plants

grew tall enough, we made a hole in the cover to accommodate their growth (Supplemental

Figure 1). Four weeks post-planting, we uncovered the terminal drought plants to allow more

drying and kept the well-watered plants covered. We filled the bottom compartments of well-

watered plants with 300 ml of autoclaved dH_2O. We began the gradual terminal drought

treatment with 2 weeks of well-watered conditions, followed by 2 weeks of ⅔ volume water,

then 3 weeks of ⅓ volume water, and finally no further water. We refilled all plant water

compartments with the appropriate volumes whenever the well-watered plants used all their

water. Ten weeks post-planting, just over 80% of terminal drought plants were dead and we

ended the experiment. We counted branch number approximately every 10 days, and measured

the nodule number and dry weight of above- and below- ground biomass of each plant, including

dead plants, at the end of the experiment.

Statistical analysis

We fit linear models with log_{10}-transformed above- and below-ground biomasses as response

variables and the main and interactive effects of soil origin and experimental water condition as

fixed effects and spatial block as a random effect. Final branch number was tested in a

generalized linear model with a Poisson distribution. We included data from both live and

measurable dead plants in the models. Only a single plant made nodules, and thus we did not fit a

statistical model to nodule number.

Experiment 2: Soil climate legacy effects with and without an additional mutualist

Experimental design

In a separate experiment, we next tested whether heat- and drought-treated microbes affected

legumes and rhizobia, under well-watered conditions only. We factorially tested the four climate

treatments, each with 3 replicate plots from the warming array, and with sterilized or fresh soil.

The same collection of soil was used in Experiment 1 and 2. We used autoclaved sterilized soil

(Supplemental Material) to test for abiotic differences in soil from different field plots and

climate treatments. The field soil resulted in poor nodulation in Experiment 1, so, in Experiment

2, we also inoculated half of the plants with *Sinorhizobium meliloti* 1021-71 tagged with green

fluorescent protein (GFP) (courtesy of Daniel Gage, Gage et al. 1996) to determine whether

additional rhizobia buffer legumes against the negative effects of drought-treated microbiomes.

In total, we had 48 treatments replicated 15 times for a total of 720 plants (Figure 1). We also

included an additional 15 sterile control plants with no additional soil.

We prepared the same genotype of seeds as before and blocked the plants by light canopy under

the same growth chamber conditions as previously described. We placed cone-tainers with a

wicking nylon rope and 130 ml of autoclaved Profile® fine-grained turface into 50 ml falcon

tubes with autoclaved distilled water, and refilled as needed. We added 1 ml of low-nitrogen

fertilizer upon planting, two weeks after planting, and every 3 weeks subsequently. We applied

biotic and sterilized soil treatments five days after planting, and applied *S. meliloti* 1021-71 or

sham inoculations six days after planting, as described in the Supplemental Material. Only two

sterile plants in the experiment made nodules, indicating minimal rhizobial contamination.

Autoclaved soil, however, was very susceptible to ambient fungal colonization, and 154

sterilized soil replicates showed mold growth. We removed all plants that showed signs of

contamination from further analysis.

Data collection

Over almost 12 weeks, we surveyed branch number and mortality. At the end of the experiment,

we randomly sampled five live plants per treatment receiving biotic soil to sequence the

microbial communities in their rhizosphere and nodules (n = 120). We did not use fluorescence

to determine rhizobia strain identity in nodules because of the possibility of bacterial

autofluorescence (Yang et al. 2012) and because we sequenced nodule bacteria. As described in

the Supplemental Material, we extracted DNA from the rhizosphere and nodule microbes, then

sequenced the fungal ITS region and bacterial 16S V4 region using Illumina MiSeq paired end

sequencing. For all plants, we counted the number of nodules then dried and weighed above and

below ground biomass.

 Statistical analysis: Plant phenotypic responses

We initially tried to fit a full four-way ANOVA (heat * drought * sterility * rhizobia) with

random effects of array plot and block to our data, but found that these models did not converge

(or had error warnings) for some of our response variables, precluding the use of a single model

structure for all the phenotypes we measured. As such, we tested the effects of added rhizobia

and soil sterility separately from the effects of climate (heat and drought treatment). In the

analyses described below, our inferences about rhizobia mediating plant responses to climate

legacies come from testing for significant differences due to climate in biotic soils, with or

without additional rhizobia (Figure 1C); because the only difference between these treatments is

the presence or absence of rhizobia addition, we attribute the differences to rhizobia.

First, to test how added rhizobia and soil sterility affected plant outcomes, we used a generalized

linear mixed-effects model with sterility treatment and rhizobia treatment as interacting fixed

effects, and block and array plot as random effects. As an exception, to test nodule data we used

a zero-inflated Poisson model in the GLMMadaptive package (Rizopoulos 2023) with heat,

drought, and block as fixed effects, and array plot as a random effect; we removed the "sterile no

added rhizobia" treatment group from this analysis because its values were only zero. The zero-

inflated model included both a model of the zeros (nodule presence or absence), and a model of

the non-zeros (nodule number).

Then, we tested for effects of soil climate origin within each of the four broad treatment

categories using generalized linear models with heat and drought as interacting fixed effects, and

with block and array plots as random effects, with exceptions and modifications described

below. To test how climate-treated microbes affected plant performance, we used repeated

branch counts as the response variable, with an additional fixed effect of days since germination

and random effect of unique plant ID. We tested branch count in the biotic no added rhizobia and

biotic added rhizobia treatments separately. In the biotic soil with no added rhizobia treatments,

to test nodule data, we used a zero-inflated Poisson model with heat, drought, and block as fixed

effects and array plot as a random effect.

We normalized response variables as needed or used generalized linear models. We $\log_{10}$-

transformed above- and below-ground biomasses and nodule number, and $\log_{10}(y+1)$-

transformed leaf counts to use a Gaussian family. We tested mortality and nodule presence with

a binomial family. We performed analysis in R v4.2.0 (R Core Team 2022), with tidyverse

(Wickham et al. 2019), cowplot (Wilke 2017), lme4 (Bates et al. 2015), and lmerTest

(Kuznetsova et al. 2017). We used type 3 ANOVAs from the car (Fox and Weisberg 2011)

package to calculate *Wald X^2* values and significance for all linear and generalized linear models.

We excluded plants that did not successfully germinate as well as plants with suspected

contamination from the analysis.

Statistical analysis: Plant microbiomes

We tested the effects of climate legacy and added rhizobia on the rhizosphere and nodule

microbial communities, and linked microbes to plant performance. We analyzed nodule,

bacterial, and fungal rhizosphere datasets separately, using the phyloseq (McMurdie and Holmes

2013), and microbiome (Lahti and Shetty 2019) packages. We excluded any potential root

endophytes from our nodule samples by subsetting reads to include only *Sinorhizobium* ASVs.

We used generalized linear models with heat and drought as interacting fixed effects, the fixed

effect of added rhizobia, and random effects of block and array plot; when we tested nodule data

we added total nodule number as a fixed effect. In the rhizosphere data, we tested observed

richness, Shannon diversity, and evenness using Gaussian distributions. To test alpha diversity in

nodules, we used a Poisson distribution for observed richness, a Gaussian distribution for

evenness, and a binomial distribution for Shannon diversity (due to the frequency of a single

ASV being observed, we categorized Shannon diversity as 0 or > 0). Next, we tested how

climate legacy affected microbial composition using adonis2 (vegan package, (Oksanen et al.

2020) permutational analysis of variance models (PERMANOVA; Bray-Curtis method, 9999

permutations) with heat and drought as interacting fixed effects, added rhizobia as another fixed

effect, and permutations done within blocks. To test nodule composition, we added total nodule

number as a fixed effect.

Finally, to identify microbes correlated to plant performance, we subset the top 20% best

performing (n = 23) and bottom 20% worst performing (n = 23) plants from the sequenced plants

after rarefaction in both the bacterial and fungal datasets. Then, using the microviz package

(Barnett et al. 2021), we used the tax_fix function, center log ratio transformed abundances of

the rarefied rhizosphere reads, and used a principal components analysis (PCA) to plot points

and identify the top 10 genera with the largest effect sizes. We tested whether the best and worst

performing plants had significantly different rhizospheres using a PERMANOVA (Aitchison

method, 9999 permutations accounting for block). To match our PCA, we aggregated taxa

abundance to genus, then used performance category and added rhizobia as fixed predictors, and

heat and drought as interacting predictors.

Results

Drought-treated microbes reduced legume performance

In both experiments, drought-treated microbes harmed plants. In Experiment 1, plants grew

largest under well-watered conditions, but only when inoculated with soil microbes from the

control, not the droughted, field plots. There were significant effects of both current water

condition and the historical soil water condition on aboveground biomass (*Wald X^2* > 18.5, *p* <

0.001 for all effects; Supplemental Table 1, Figure 2A), with a significant interaction (*Wald X^2* =

5.98, *p* < 0.05; Supplemental Table 1, Figure 2A). When plants received ample water, drought-

treated microbes reduced plant aboveground biomass by 51.3% compared to control microbes,

but in terminal drought conditions, drought-treated microbes reduced plant aboveground biomass

by only 9.18% compared to control microbes. Belowground biomass also significantly decreased

with drought-treated microbiomes (*Wald X^2* = 18.0, *p* < 0.001; Supplemental Table 1, Figure

2B), and there was no main effect of the terminal drought treatment (Supplemental Table 1,

Figure 2B), although there was a current conditions x historical conditions interaction (*Wald X^2* =

7.44, *p* < 0.01; Supplemental Table 1, Figure 2B). The interaction term appears to be driven by

the different responses to well-watered conditions between plants given control or drought-

treated soil, when they had the same response in terminal drought (Figure 2B). The belowground

biomass of plants with control soil decreased by 40.7% in terminal drought, compared to well-

watered conditions. In contrast, plants with drought-treated microbes had 134% higher

belowground biomass in terminal drought conditions, compared to well-water conditions.

Similarly, drought-treated microbes and terminal drought significantly decreased branch number

(*Wald X^2* > 4.64, *p* < 0.05 for all effects; Supplemental Table 1, Supplemental Figure 2), but did

not significantly interact (Supplemental Table 1). Terminal drought also significantly increased

mortality (*Wald X^2* = 9.58, *p* < 0.01; Supplemental Table 1), but there were no mortality

differences between microbe treatments (Supplemental Table 1). Only one plant made nodules,

and it received control soil under well-watered conditions.

In our second experiment, conducted under well-watered conditions only, we found that *M.*

lupulina performance was unaffected by heat-treated vs. control microbes, but again drought-

treated microbes reduced plant performance. Plants with biotic, drought-treated microbes with no

added rhizobia had a significantly reduced branch count compared to plants given biotic control

microbes with no added rhizobia (*Wald X^2* = 15.4, *p* < 0.001; Supplemental Table 2, Figure 3A).

There was a significant interaction between drought and heat legacies on branch count (*Wald X^2*

= 5.37, *p* < 0.05; Supplemental Table 2, Figure 3A), where heat counteracted the negative effect

of drought. In biotic soil with no added rhizobia, drought-treated microbes significantly

increased mortality from 2.2% when inoculated with control microbes to 18% when inoculated

with drought-treated microbes (*Wald X^2* > 4.33, *p* < 0.05; Supplemental Table 2, Figure 3B).

However, by the end of Experiment 2, the above- and below-ground biomasses of surviving

plants did not differ significantly among microbial treatments (Supplemental Table 2, Figure

3CD). While mean nodule number (including plants that made no nodules) decreased by 44% in

plants inoculated with heat-treated field soil compared to control soil, the reductions in nodule

presence and number was not statistically significant (Supplemental Table 2, Figure 3E).

Differences in plant performance were not due to abiotic soil differences, because there were no

significant differences in plant performance due to climate origin in the autoclaved, no added

rhizobia soil treatments (Supplemental Table 3).

Rhizobia reduced differences due to climatic legacy effects

Adding rhizobia to the biotic soil erased the significant plant performance differences that arose

from inoculation with biotic drought-treated soils. In this treatment, mortality was 12-16% across

climate treatments, with no significant differences due to climate legacy (Supplemental Table 4).

Furthermore, there were no significant differences in branch counts, nodule number, nodule

presence, or plant biomass among climate treatments in biotic soil with added rhizobia

(Supplemental Table 4). There also were no differences due to microbial climate in the added

rhizobia, sterilized soil treatment (Supplemental Table 5).

Biotic soil improved plant performance more than adding rhizobia alone

When we tested the effects of soil sterility and added rhizobia, we found that plants performed best in unaltered field soil. Above- and below-ground biomasses significantly increased with added rhizobia inoculation (*Wald X^2* > 5.23, $p < 0.05$ for all effects; Supplemental Table 6, Figure 4AB), but with a significant interaction between rhizobia and sterility condition (*Wald X^2* > 6.64, $p < 0.01$ for all effects; Supplemental Table 6, Figure 4AB). Unsurprisingly, adding rhizobia increased plant biomass in sterilized soil, but adding rhizobia decreased plant biomass when plants were inoculated with biotic field soils. The mean aboveground biomass of plants in the biotic soil with the natural soil rhizobia community was 23% higher compared to plants in biotic soil with added rhizobia. Mortality was not affected by soil sterility or added rhizobia (Supplemental Table 6, Figure 4C). Nodule presence and number were significantly affected by soil sterility, added rhizobia, and spatial block (|*z-value*| > 2.04, $p < 0.05$ for all effects; Supplemental Table 6, Figure 4DE). Nodule presence decreased with no added rhizobia and no access to the natural rhizobia community. More nodules were made per plant with the unmodified field rhizobia, and plants made more nodules in the added rhizobia treatment only in the sterilized soil, when no other rhizobia were available.

Root microbiomes correlated to performance and changed by adding rhizobia

After rarefaction, bacterial rhizospheres contained 7,907 ASVs and had a high abundance of rhizobial taxa such as *Rhizobium* and *Bradyrhizobium*, but also high amounts of *Arthrobacter*, *Streptomyces*, and unknown taxa (Supplemental Figure 3). After rarefaction, fungal rhizospheres contained 992 ASVs, and were mainly composed of *Trichoderma* and *Fusarium*, with high occurrences of *Aspergillus*, *Neocosmospora*, and *Penicillium* (Supplemental Figure 4). Bacterial rhizosphere composition was not significantly altered by climate legacy (Supplemental Table 7) but was significantly altered by the addition of *S. meliloti* 1021-71 ($F_{1,108} = 1.35$, $p < 0.05$; Supplemental Table 7). Only heat-treated soils led to significantly higher observed bacterial richness in the rhizosphere compared to control soil (*Wald X^2* = 4.08, $p < 0.05$; Supplemental Table 7), but Shannon diversity and evenness were not significantly different among treatments (Supplemental Table 7). Fungal rhizosphere community composition and diversity were not significantly affected by adding rhizobia or by the legacies of climate (Supplemental Table 8).

The best and worst performing plants had significantly different bacterial rhizospheres ($F_{1,39}$ = 4.26, $p < 0.001$; Figure 5, Supplemental Table 7), but not fungal rhizospheres (Supplemental Table 8; Supplemental Figure 5). Of the 10 microbes with the largest effect, all but one were known plant beneficial bacteria (Li et al. 2023). The diazotrophs *Cellulomonas*, *Cryocola*, and *Sinorhizobium* (Hrynkiewicz et al. 2019) were positively correlated with plant performance (Figure 5). *Bacillus* was also positively correlated with plant performance (Figure 5) and can induce systemic resistance in plants, reducing the incidence and severity of phytopathogens (Kloepper et al. 2004). *Emticicia*, *Niastella*, *Variovorax*, and *Bdellovibrio,* as well as the families *Verrumicrobiaceae* and *Chitinophagaceae*, correlated with decreased plant performance (Figure 5).

Plants were very good at isolating *Sinorhizobium* from the soil (Supplemental Figure 3), and sequenced nodules were predominantly composed of 13 ASVs in the genus *Sinorhizobium*. *Sinorhizobium* ASVs were very similar to one another and to the ASV of *S. meliloti* 1021-71; we observed three different base pair substitutions, with other ASV diversity coming from 1-3 fewer or additional base pairs at the beginning and end of the sequence. The composition of those ASVs was not significantly affected by climate legacy, added rhizobia, or total nodule number (Supplemental Table 9). Increasing nodule number significantly increased the Shannon diversity of *Sinorhizobium* strains (*Wald X^2* = 4.89, $p < 0.05$; Supplemental Table 9), but no other metrics of alpha diversity (Supplemental Table 9).

<u>Discussion</u>

Climate events leave lasting effects on soil microbiomes, generating indirect microbially-mediated effects on plant performance and symbiosis. We found that drought-treated microbes negatively impacted legume performance in two experiments. In the first, drought-treated microbes significantly reduced plant biomass, and microbe history significantly interacted with experimental water availability to affect plant performance. In the second, drought-treated microbes increased mortality rates. When we added rhizobia, the differences in plant performance due to climate legacy became nonsignificant. Symbiosis outcomes were altered in other ways, with added rhizobia altering the bacterial rhizosphere composition and heat-legacy increasing the diversity of bacteria in the rhizosphere of live plants. Here we discuss the

378   consequences of plant performance being shaped by these microbial indirect effects during

379   climate change, how mutualism quality may remain stable in the face of climate change, and

380   how mutualisms could potentially provide a stabilizing effect on plant performance across

381   climate-driven differences in the microbiome.

Negative effects of drought-treated microbes and their ecological implications

Plants respond to climate change through both direct and indirect pathways (Porter et al. 2020,

Trivedi et al. 2022), and the relative importance and impact of these effects is necessary to

predict plant performance and adaptation. Strikingly, the strength of microbially-mediated

indirect effects of drought on plant performance was equal in magnitude to the direct effects of

imposed drought (Figure 2). It is concerning that drought-treated microbiomes led to a worse

plant performance because droughts are becoming more frequent and intense (Thuiller 2007,

Allan and Soden 2008), and our results suggest that we cannot rely on soil microbiomes to

contribute positively to plant resistance and resilience to climate change. Further, strong

microbially-mediated indirect effects of climate change may impose additional ecological

pressures on plants that must already respond to the direct effects of climate change. Previous

studies found that drought microbiomes can buffer plants against drought stress (Lau and Lennon

2012, Allsup et al. 2023, Ricks and Yannarell 2023), but our results and recent results from Swift

et al. (2024) indicate that this effect is not universal and that repeated drought events do not

always select for microbes that improve plant performance relative to control microbes.

There are several possible mechanisms by which the drought-treated soil microbiomes may have

decreased plant performance. The higher risk to plants could have been due to a lack of

compatible mutualists, as the higher rhizobia diversity in drought (Boyle et al. 2024) may have

increased competitive interference amongst rhizobia (Burghardt and diCenzo 2023, Rahman et

al. 2023), lengthening the time needed for plants to locate compatible partners, and ultimately

increasing risk of death. Given that plant performance was strongly positively correlated with

Sinorhizobium abundance in the rhizosphere (Figure 5), and adding rhizobia erased climate-

driven differences in plant performance, access to compatible rhizobia is clearly important for

the plant. While the rhizobia diversity mechanism is formally plausible, we failed to detect other

clear signals of it, such as corresponding differences in nodule number and diversity between

surviving plants that received different climate-treated soils; as such, we think it is less likely.

Alternatively, our observed patterns of performance would occur if the prevalence of a

detrimental microbe was higher in drought-treated soil. Indeed, the higher mortality may have

been because of the 36% increase in relative abundance of fungal parasites and the altered

bacterial phytopathogen composition in the drought inoculation soil (Boyle et al. 2024). Included

amongst the plant fungal parasites in the inoculation soils are *Verticillium* and

Magnaporthiopsis, which can cause wilt in legumes and non-legumes (Klosterman et al. 2009,

Dor and Degani 2019), and *Paraphoma*, which can cause root rot in alfalfa and other plants

using phytotoxins (Dang and Li 2022). Global warming and drought are projected to increase the

proportion of soil-borne pathogens (Preece et al. 2019, Delgado-Baquerizo et al. 2020), and how

potential pathogens interact with the rest of the plant microbiome under drought and heat stress

is not well understood (Trivedi et al. 2022). Our results novelly suggest drought increases plants'

risk of encountering combinations of deadly microbes, but as we discuss more below, that the

harm could be diluted if microbial mutualists are present at high enough frequency. Additionally,

our sequencing results suggest that the microbes in the drought-treated soil and rhizospheres

were less metabolically active and provided fewer nutrients to plants. The bacterial genus

Verrucomicrobium was significantly enriched in the drought-treated inoculation soil (Boyle et al.

2024), and the *Verrucomicrobiaceae* family abundance in the rhizosphere was negatively

correlated with plant performance in Experiment 2 (Figure 5). Possibly due to its slow growth,

the phylum *Verrucomicrobia* is linked to reduced soil enzyme activity and loss of soil nutrients

like nitrogen (Xie et al. 2023), and as such, it is considered as an indicator of poor nutrient

conditions (Navarrete et al. 2015). Given the significant differences in the microbial

inoculations, it is possible that a combination of reduced access to rhizobia, fewer aggregate

microbial benefits, as well as increased pathogen prevalence may have contributed to the worse

plant performance in drought-treated soil.

Our root microbiome analysis did not find strong compositional and diversity differences due to

drought-treated soil, likely because we only sequenced the microbiomes of live plants, which

reduced our ability to detect microbes that may have contributed to differential mortality

between climate-treated soils (e.g., survivorship bias). Furthermore, it is possible that there were

more differences in the rhizosphere of plants early in the experiment, reflected in the strong

decrease in performance of younger plants, but that these compositional differences were

reduced over time by the mortality of affected plants and the selective environment of the growth

chamber and the plants.

440 Rhizobia inoculation reduced differences in plant performance due to climate-legacy

We added rhizobia to the soil to determine if mutualists could buffer plants against microbially-

mediated negative effects of climate and found that the inoculation erased plant performance

differences between climate-treated soils. If abundant mutualistic microbes reduce variation in

plant performance and growth due to climate-driven differences in the microbiome, other

organisms relying on those plants for nutrition or habitat will also experience less disruption, and

this stabilizing effect could cascade across trophic levels (Kagata and Ohgushi 2006, Scherber et

al. 2010). Thus, when partners are prevalent, mutualism can stabilize ecological dynamics

against the indirect effects of climate change.

The relative abundance of compatible rhizobia in the un-inoculated field soil was very low

(Boyle et al. 2024, Supplemental Figure 3), and in these conditions, climate-treated microbes

altered plant performance. So, performance differences may have disappeared because of the

increased abundance and accessibility of rhizobial mutualists to the host early in life, or

potentially due to microbe-microbe interactions, both of which can affect host performance

(Bakker et al. 2014, May and Nelson 2014, Burghardt et al. 2022). For example, a relatively high

abundance of added rhizobia could have outcompeted detrimental microbes for resources or

access to the host, or even altered the host as a niche via priority effects (Boyle et al. 2021,

Debray et al. 2022, Burghardt and diCenzo 2023).

While performance was homogenized across climate treatments with the added rhizobia, plants

with added rhizobia tended to do worse than plants with no added rhizobia in biotic soil,

suggesting that the inoculated rhizobium strain was not a high-quality partner. This was not

unexpected, as *S. meliloti* 1021 is well-characterized as a mediocre partner with *Medicago*

truncatula (Terpolilli et al. 2008, Batstone et al. 2020a). Furthermore, the inserted green

fluorescence protein (GFP) in our added strain of *Sinorhizobium meliloti* 1021-71 may make the

microbe grow more slowly than non-modified strains (Rang et al. 2003) and may have required

relatively more energy and resources from the plant. Another possibility is that our collected *M.*

lupulina genotype is locally adapted to the rhizobia with which it co-occurs, and local rhizobia

provided a greater benefit than the added strain (Batstone et al. 2020a). Our results indicate

mutualists can reduce the effect of background variation in the microbiome if they are prevalent

enough, however, for a truly protective effect they may need to match or supersede the quality of

local mutualists.

Mutualism quality and symbiotic patterns were robust to climate legacy

The maintenance of mutualism under changing environmental contexts will be important for

interacting species' fitnesses under future climate change, as well as continued ecosystem

services (Kiers et al. 2010). Climate change is predicted to make mutualisms vulnerable to shifts

towards antagonism, partner switches and losses, and mutualism abandonment (Brown 1997,

Sachs and Simms 2006, Kiers et al. 2010). Our results demonstrated that heat- and drought-

treated field rhizobia maintained their partner quality and ability to nodulate a legume host

(Figure 3). The most immediate threat to legume-rhizobia symbioses may be the lower

abundance or loss of a preferred partner, where the loss of a specialized mutualist means the

other partner has few replacements and replacements are lower-quality (Froehlich et al. 2023).

In general, differences in the soil were not very predictive of differences in the root microbiomes

(Supplemental Figures 3-4). Previous results found the most significant differences from climate

treatments were due to drought and in the fungal community (Boyle et al. 2024), yet surprisingly,

this did not manifest in significantly different fungal rhizospheres. One reason for this difference

may be that many strongly affected fungi did not form a close symbiosis with *Medicago*

lupulina, as most fungi in the soil were saprotrophs and most mycorrhizal fungi were

ectomycorrhizal (Boyle et al. 2024). Moreover, it appears that there is increasingly strong host

selection from the soil to the rhizosphere to the nodule; a pattern that has been observed in other

legumes (Miranda☐Sánchez et al. 2016), suggestive of a deterministic microbiome (Lundberg et

al. 2012). This pattern may arise due to factors such as host morphology, host rewards and

byproducts, and host immune response, all of which can select for a subset of microbes from the

broader microbial community (Zipfel and Oldroyd 2017, VanWallendael et al. 2022). Filtering

mechanisms may increase plant performance if it also ensures recruitment of beneficial microbes

and the exclusion of harmful microbes (Werner and Kiers 2015), but, in and across changing

environments, the microbes to target and filter may be different (Caballol et al. 2024). An

interesting next step would be to investigate the differences in how generalist or specialist

legumes respond to microbially-mediated effects of climate, and the benefits and drawbacks of each strategy. While we observed a strongly deterministic root microbiome of living *M. lupulina* plants, it is also clear that when the 'right' microbes are affected by climate, legumes can still suffer lower plant performance and higher mortality.

Conclusions

Plants experienced detrimental indirect effects of climate change through their microbiome, with the strongest effects early in life and due to the legacy of drought. Repeated drought events eroded the soil microbiome's benefits under both drought stress and wet conditions, potentially reducing future plant resistance and resilience to drought. Indirect negative effects of drought on plant growth and fitness may challenge ecosystem productivity and function as well as agricultural yield. Additional access to rhizobia mutualists reduced microbially-mediated differences in plant performance due to climate. Our results motivate further inquiry into whether the addition of good-quality mutualists could buffer plants against this variation in the microbiome and potentially provide plants with increased stability under new climate regimes, with relevant applications in managed systems like agriculture and restoration sites. Our study highlights the complexity of plant response to components of climate change like extreme drought, and the threat of indirect microbially-mediated climate effects.

Acknowledgements

We thank past and present KSR staff, Anna Simonsen for generating the seed line, Daniel Gage for providing the rhizobia strain, Chris Carlson for his help collecting soil and thoughtful discussions, Jessica Underwood for her help planting and applying treatments, and Bill Cole and Tom Gludovacz for growth chamber help. This work was funded by the Government of Canada through Genome Canada and Ontario Genomics (OGI-211), NSERC Discovery grants to IE, JRS, and MEF, the Canada Foundation for Innovation, and the University of Toronto's Center for Global Change Science. We also acknowledge Genome Quebec for their sequencing facilities.

524 **Figures**

525
526
Figure 1. Conceptual motivation and treatment design for the two experiments. A: Plants
and microbes experience the direct effects of climate (fully red arrows), as well as the indirect
effects of climate mediated through plant-microbe interactions (partially red arrows). Here, we
investigate the indirect effect of climate on plants, mediated through the microbiome (yellow to
red arrow). B: We pooled soil from control and drought-treated field plots and added them to
plants in well-watered or terminal drought conditions. C: For soil from each field plot, we
factorially tested the effect of biotic vs. autoclaved soil and added rhizobia vs. no added rhizobia.

Figure 2. Climate legacy effects in soil microbiomes have context dependent effects on plants. Current and historical conditions, along with their interaction, significantly affected plant aboveground biomass (A) and belowground biomass (B) in the first experiment. Points are means with ±1 standard error. The y-axes are on a $\log_{10}$ scale.

Figure 3. Drought-treated microbes decreased *Medicago lupulina* performance in the biotic, no added rhizobia treatment. In the second experiment, drought-treated microbes significantly decreased branch number when plants were young (A) and increased mortality (B), but by the end of the experiment, surviving plants had similar biomasses and nodule number across climate-treated soils (C, D, E). Points represent means ± 1 standard error. The y-axes in C and D are on a $\log_{10}$ scale.

546

Figure 4. Plants performed best in the biotic soil with no added rhizobia treatments. In the second experiment, plants given unaltered field soil had the highest biomasses, while plants that received sterile soil with no additional rhizobia had the lowest biomass (A, B). The effect of adding rhizobia on biomass was dependent on the presence of other microbes, causing significant interaction terms (A, B). Mortality was not significantly different due to sterilized soil or added rhizobia (C). Nodule presence was highest when added rhizobia were the only microbes available, and second highest with added rhizobia and field microbes (D), but when plants did make nodules, they made the most with biotic unaltered field soil (E). Points represent means ± 1 standard error. The y-axes in A and B are on a $\log_{10}$ scale.

Figure 5. Microbes were correlated to performance differences in the bacterial rhizosphere. Here a PCA shows the center log ratio transformed bacterial rhizosphere composition of the top 20% best performing and bottom 20% worst performing plants in the second experiment. Performance was quantified by aboveground biomass (g).

557
558
559
560

Literature cited

Allan, R. P., and B. J. Soden. 2008. Atmospheric Warming and the Amplification of Precipitation Extremes. Science 321:1481–1484.

Allison, S. D., and K. K. Treseder. 2008. Warming and drying suppress microbial activity and carbon cycling in boreal forest soils. Global Change Biology 14:2898–2909.

Allsup, C. M., I. George, and R. A. Lankau. 2023. Shifting microbial communities can enhance tree tolerance to changing climates. Science 380:835–840.

Averill, C., B. G. Waring, and C. V. Hawkes. 2016. Historical precipitation predictably alters the shape and magnitude of microbial functional response to soil moisture. Global Change Biology 22:1957–1964.

Bakker, M. G., D. C. Schlatter, L. Otto☐Hanson, and L. L. Kinkel. 2014. Diffuse symbioses: roles of plant–plant, plant–microbe and microbe–microbe interactions in structuring the soil microbiome. Molecular Ecology 23:1571–1583.

Barnett, D., I. Arts, and J. Penders. 2021. microViz: an R package for microbiome data visualization and statistics. Journal of Open Source Software 6:3201.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software 67:1–48.

Batstone, R. T., A. M. O'Brien, T. L. Harrison, and M. E. Frederickson. 2020a. Experimental evolution makes microbes more cooperative with their local host genotype. Science 370:476–478.

Batstone, R. T., M. A. E. Peters, A. K. Simonsen, J. R. Stinchcombe, and M. E. Frederickson. 2020b. Environmental variation impacts trait expression and selection in the legume–rhizobium symbiosis. American Journal of Botany 107:195–208.

Beringer, J. E., N. Brewin, A. W. B. Johnston, H. M. Schulman, and D. A. Hopwood. 1979. The *Rhizobium*-legume symbiosis. Proceedings of the Royal Society of London. Series B. Biological Sciences 204:219–233.

Boyle, J. A., B. K. Murphy, I. Ensminger, J. R. Stinchcombe, and Megan. E. Frederickson. 2024. Resistance and resilience of soil microbiomes under climate change. Ecosphere 15: e70077.

Boyle, J. A., A. K. Simonsen, M. E. Frederickson, and J. R. Stinchcombe. 2021. Priority effects alter interaction outcomes in a legume–rhizobium mutualism. Proceedings of the Royal Society B 288.

Brown, B. E. 1997. Coral bleaching: causes and consequences. Coral Reefs 16:129–138.

Burghardt, L. T., and G. C. diCenzo. 2023. The evolutionary ecology of rhizobia: multiple facets of competition before, during, and after symbiosis with legumes. Current Opinion in Microbiology 72:102281.

Burghardt, L. T., B. Epstein, M. Hoge, D. I. Trujillo, and P. Tiffin. 2022. Host-Associated Rhizobial Fitness: Dependence on Nitrogen, Density, Community Complexity, and Legume Genotype. Applied and Environmental Microbiology 88.

Caballol, M., M. Á. Redondo, N. Catalán, T. Corcobado, T. Jung, B. Marçais, I. Milenković, M. Nemesio-Gorriz, J. Stenlid, and J. Oliva. 2024. Climate acts as an environmental filter to plant pathogens. The ISME Journal 18.

Castro, H. F., A. T. Classen, E. E. Austin, R. J. Norby, and C. W. Schadt. 2010. Soil microbial community responses to multiple experimental climate change drivers. Applied and Environmental Microbiology 76:999–1007.

Chen, W. F., E. T. Wang, Z. J. Ji, and J. J. Zhang. 2021. Recent development and new insight of diversification and symbiosis specificity of legume rhizobia: mechanism and application. Journal of Applied Microbiology 131:553–563.

Classen, A. E., M. K. Sundqvist, J. A. Henning, G. S. Newman, J. A. Moore, M. A. Cregger, L. C. Moorhead, C. M. Patterson, M. K. Sundqvist, J. A. Henning, and G. S. Newman. 2015. Direct and indirect effects of climate change on soil microbial and soil microbial-plant interactions: What lies ahead? Ecosphere 6:1–21.

Copeland, J. K., L. Yuan, M. Layeghifard, P. W. Wang, and D. S. Guttman. 2015. Seasonal Community Succession of the Phyllosphere Microbiome. Molecular Plant-Microbe Interactions 28:274–285.

Dang, S.-Z., and Y.-Z. Li. 2022. The Characterization and the Biological Activity of Phytotoxin Produced by *Paraphoma radicina*. Journal of Fungi 8:867.

Davis, T. S., N. A. Bosque-Pérez, N. E. Foote, T. Magney, and S. D. Eigenbrode. 2015. Environmentally dependent host–pathogen and vector–pathogen interactions in the Barley yellow dwarf virus pathosystem. Journal of Applied Ecology 52:1392–1401.

Debray, R., R. A. Herbert, A. L. Jaffe, A. Crits-Christoph, M. E. Power, and B. Koskella. 2022. Priority effects in microbiome assembly. Nature Reviews Microbiology 20:109–121.

Delgado-Baquerizo, M., C. A. Guerra, C. Cano-Díaz, E. Egidi, J.-T. Wang, N. Eisenhauer, B. K. Singh, and F. T. Maestre. 2020. The proportion of soil-borne pathogens increases with warming at the global scale. Nature Climate Change 10:550–554.

Dor, S., and O. Degani. 2019. Uncovering the Host Range for Maize Pathogen *Magnaporthiopsis maydis*. Plants 8:259.

Duarte, A. G., and H. Maherali. 2022. A meta☐analysis of the effects of climate change on the mutualism between plants and arbuscular mycorrhizal fungi. Ecology and Evolution 12.

Fitzpatrick, C. R., Z. Mustafa, and J. Viliunas. 2019. Soil microbes alter plant fitness under competition and drought. Journal of Evolutionary Biology 32:438–450.

Fox, J., and S. Weisberg. 2011. An R companion to applied regression. Second edition. Sage Publications.

Franklin, J., J. M. Serra-Diaz, A. D. Syphard, and H. M. Regan. 2016. Global change and terrestrial plant community dynamics. Proceedings of the National Academy of Sciences 113:3725–3734.

Froehlich, C. Y. M., O. S. Klanten, M. L. Hing, M. Dowton, and M. Y. L. Wong. 2023. Delayed recovery and host specialization may spell disaster for coral☐fish mutualism. Ecology and Evolution 13.

Gage, D. J., T. Bobo, and S. R. Long. 1996. Use of Green Fluorescent Protein To Visualize the Early Events of Symbiosis between *Rhizobium meliloti* and Alfalfa (*Medicago sativa*). JOURNAL OF BACTERIOLOGY 178:7159–7166.

González, R., A. Butković, F. J. Escaray, J. Martínez-Latorre, Í. Melero, E. Pérez-Parets, A. Gómez-Cadenas, P. Carrasco, and S. F. Elena. 2021. Plant virus evolution under strong drought conditions results in a transition from parasitism to mutualism. Proceedings of the National Academy of Sciences 118.

Hawkes, C., J. Bull, and J. Lau. 2020. Symbiosis and stress: how plant microbiomes affect host evolution. Philosophical Transactions Royal Society B 375.

Hrynkiewicz, K., S. Patz, and S. Ruppel. 2019. *Salicornia europaea* L. as an underutilized saline-tolerant plant inhabited by endophytic diazotrophs. Journal of Advanced Research 19:49–56.

Kagata, H., and T. Ohgushi. 2006. Bottom☐up trophic cascades and material transfer in terrestrial food webs. Ecological Research 21:26–34.

Kakouridis, A., J. A. Hagen, M. P. Kan, S. Mambelli, L. J. Feldman, D. J. Herman, P. K. Weber, J. Pett-Ridge, and M. K. Firestone. 2022. Routes to roots: direct evidence of water transport by arbuscular mycorrhizal fungi to host plants. New Phytologist 236:210–221.

Kardol, P., M. A. Cregger, C. E. Campany, and A. T. Classen. 2010. Soil ecosystem functioning under climate change: plant species and community effects. Ecology 91:767–781.

Keeler, A. M., A. Rose-Person, and N. E. Rafferty. 2021. From the ground up: Building predictions for how climate change will affect belowground mutualisms, floral traits, and bee behavior. Climate Change Ecology 1:100013.

Kiers, T. E., T. M. Palmer, A. R. Ives, J. F. Bruno, and J. L. Bronstein. 2010. Mutualisms in a changing world: an evolutionary perspective. Ecology Letters 13:1459–1474.

Kivlin, S. N., S. M. Emery, and J. A. Rudgers. 2013. Fungal symbionts alter plant responses to global change. American Journal of Botany 100:1445–1457.

Kloepper, J. W., C.-M. Ryu, and S. Zhang. 2004. Induced Systemic Resistance and Promotion of Plant Growth by *Bacillus* spp. Phytopathology 94:1259–1266.

Klosterman, S. J., Z. K. Atallah, G. E. Vallad, and K. V. Subbarao. 2009. Diversity,
Pathogenicity, and Management of *Verticillium* Species. Annual Review of Phytopathology
47:39–62.

Koide, R. T. 1991. Nutrient supply, nutrient demand and plant response to mycorrhizal infection.
New Phytologist 117:365–386.

Kolodny, O., and H. Schulenburg. 2020. Microbiome-mediated plasticity directs host evolution
along several distinct time scales. Philosophical Transactions of the Royal Society B:
Biological Sciences 375:20190589.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest Package: Tests in
Linear Mixed Effects Models . Journal of Statistical Software 82:1–26.

Lahti, L., and S. Shetty. 2019. microbiome R package.
https://bioconductor.org/packages/release/bioc/html/microbiome.html.

Lau, J. A., and J. T. Lennon. 2012. Rapid responses of soil microorganisms improve plant fitness
in novel environments. Proceedings of the National Academy of Sciences of the United
States of America 109:14058–14062.

Li, P., L. Tedersoo, T. W. Crowther, A. J. Dumbrell, F. Dini-Andreote, M. Bahram, L. Kuang, T.
Li, M. Wu, Y. Jiang, L. Luan, M. Saleem, F. T. de Vries, Z. Li, B. Wang, and J. Jiang. 2023.
Fossil-fuel-dependent scenarios could lead to a significant decline of global plant-beneficial
bacteria abundance in soils by 2100. Nature Food 4:996–1006.

Lu, T., M. Ke, M. Lavoie, Y. Jin, X. Fan, Z. Zhang, Z. Fu, L. Sun, M. Gillings, J. Peñuelas, H.
Qian, and Y. G. Zhu. 2018. Rhizosphere microorganisms can influence the timing of plant
flowering. Microbiome 6:1–12.

Lundberg, D. S., S. L. Lebeis, S. H. Paredes, S. Yourstone, J. Gehring, S. Malfatti, J. Tremblay,
A. Engelbrektson, V. Kunin, T. G. del Rio, R. C. Edgar, T. Eickhorst, R. E. Ley, P.
Hugenholtz, S. G. Tringe, and J. L. Dangl. 2012. Defining the core Arabidopsis thaliana root
microbiome. Nature 488:86–90.

Martiny, J. B., A. C. Martiny, C. Weihe, Y. Lu, R. Berlemont, E. L. Brodie, M. L. Goulden, K. K.
Treseder, and S. D. Allison. 2017. Microbial legacies alter decomposition in response to
simulated global change. The ISME Journal 11:490–499.

May, G., and P. Nelson. 2014. Defensive mutualisms: do microbial interactions within hosts
drive the evolution of defensive traits? Functional Ecology 28:356–363.

McMurdie, P. J., and S. Holmes. 2013. phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. PLOS ONE 8:e61217.

Melillo, J. M., S. D. Frey, K. M. DeAngelis, W. J. Werner, M. J. Bernard, F. P. Bowles, G. Pold,
M. A. Knorr, and A. S. Grandy. 2017. Long-term pattern and magnitude of soil carbon
feedback to the climate system in a warming world. Science 358:101–105.

Mendes, R., M. Kruijt, I. de Bruijn, E. Dekkers, M. van der Voort, J. H. M. Schneider, Y. M.
Piceno, T. Z. DeSantis, G. L. Andersen, P. A. H. M. Bakker, and J. M. Raaijmakers. 2011.
Deciphering the Rhizosphere Microbiome for Disease-Suppressive Bacteria. Science
332:1097–1100.

Miranda☐Sánchez, F., J. Rivera, and P. Vinuesa. 2016. Diversity patterns of Rhizobiaceae
communities inhabiting soils, root surfaces and nodules reveal a strong selection of
rhizobial partners by legumes. Environmental Microbiology 18:2375–2391.

Navarrete, A. A., T. Soares, R. Rossetto, J. A. van Veen, S. M. Tsai, and E. E. Kuramae. 2015.
Verrucomicrobial community structure and abundance as indicators for changes in chemical
factors linked to soil fertility. Antonie van Leeuwenhoek 108:741–752.

de Nijs, E. A., L. C. Hicks, A. Leizeaga, A. Tietema, and J. Rousk. 2019. Soil microbial moisture
dependences and responses to drying–rewetting: The legacy of 18 years drought. Global
Change Biology 25:1005–1015.

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. Mcglinn, P. R. Minchin, R. B.
O'hara, G. L. Simpson, P. Solymos, M. Henry, H. Stevens, E. Szoecs, and H. W. Maintainer.
2020. Package "vegan" Version 2.5-7.

Panke-Buisse, K., A. C. Poole, J. K. Goodrich, R. E. Ley, and J. Kao-Kniffin. 2015. Selection on
soil microbiomes reveals reproducible impacts on plant function. ISME Journal 9:980–989.

Pec, G. J., L. T. A. Van Diepen, M. Knorr, A. S. Grandy, J. M. Melillo, K. M. Deangelis, J. L.
Blanchard, S. D. Frey. 2021. Fungal community response to long-term soil warming with
potential implications for soil carbon dynamics. Ecosphere 12:e03460.

Porter, S. S., R. Bantay, C. A. Friel, A. Garoutte, K. Gdanetz, K. Ibarreta, B. M. Moore, P. Shetty,
E. Siler, and M. L. Friesen. 2020. Beneficial microbes ameliorate abiotic and biotic sources
of stress on plants. Functional Ecology 34:2075–2086.

Porter, S. S., and J. L. Sachs. 2020. Agriculture and the Disruption of Plant–Microbial
Symbiosis. Trends in Ecology & Evolution 35:426–439.

Preece, C., E. Verbruggen, L. Liu, J. T. Weedon, and J. Peñuelas. 2019. Effects of past and
current drought on the composition and diversity of soil microbial communities. Soil
Biology and Biochemistry 131:28–39.

Quiroga, G., Erice, G., Aroca, R., Chaumont, F., and J.M. Ruiz-Lozano. 2017. Enhanced Drought
Stress Tolerance by the Arbuscular Mycorrhizal Symbiosis in a Drought-Sensitive Maize
Cultivar Is Related to a Broader and Differential Regulation of Host Plant Aquaporins than
in a Drought-Tolerant Cultivar. Frontiers in Plant Science 8.

R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Rahman, A., M. Manci, C. Nadon, I. A. Perez, W. F. Farsamin, M. T. Lampe, T. H. Le, L. Torres
Martínez, A. J. Weisberg, J. H. Chang, and J. L. Sachs. 2023. Competitive interference
among rhizobia reduces benefits to hosts. Current Biology 33:2988-3001.e4.

Rang, C., J. E. Galen, J. B. Kaper, and L. Chao. 2003. Fitness cost of the green fluorescent
protein in gastrointestinal bacteria. Canadian Journal of Microbiology 49:531–537.

Ricks, K. D., and A. C. Yannarell. 2023. Soil moisture incidentally selects for microbes that
facilitate locally adaptive plant response. Proceedings of the Royal Society B: Biological
Sciences 290.

Rizopoulos, D. 2023. GLMMadaptive: Generalized Linear Mixed Models using Adaptive
Gaussian Quadrature.

Rubin, R. L., G. W. Koch, A. Martinez, R. L. Mau, M. A. Bowker, and B. A. Hungate. 2018.
Developing climate-smart restoration: Can plant microbiomes be hardened against heat
waves? Ecological Applications 28:1594–1605.

Rudgers, J. A., M. E. Afkhami, L. Bell-Dereske, Y. A. Chung, K. M. Crawford, S. N. Kivlin, M.
A. Mann, and M. A. Nuntildez. 2020. Climate Disruption of Plant-Microbe Interactions.
Annual Review of Ecology, Evolution, and Systematics 51:561–586.

Sachs, J. L., J. E. Russell, and A. C. Hollowell. 2011. Evolutionary Instability of Symbiotic
Function in *Bradyrhizobium japonicum*. PLOS ONE 6:e26370.

Sachs, J., and E. Simms. 2006. Pathways to mutualism breakdown. Trends in Ecology &
Evolution 21:585–592.

Scherber, C., N. Eisenhauer, W. W. Weisser, B. Schmid, W. Voigt, M. Fischer, E.-D. Schulze, C.
Roscher, A. Weigelt, E. Allan, H. Beßler, M. Bonkowski, N. Buchmann, F. Buscot, L. W.
Clement, A. Ebeling, C. Engels, S. Halle, I. Kertscher, A.-M. Klein, R. Koller, S. König, E.
Kowalski, V. Kummer, A. Kuu, M. Lange, D. Lauterbach, C. Middelhoff, V. D. Migunova,
A. Milcu, R. Müller, S. Partsch, J. S. Petermann, C. Renker, T. Rottstock, A. Sabais, S.
Scheu, J. Schumacher, V. M. Temperton, and T. Tscharntke. 2010. Bottom-up effects of
plant diversity on multitrophic interactions in a biodiversity experiment. Nature 468:553–
556.

Seaton, F. M., S. Reinsch, T. Goodall, N. White, D. L. Jones, R. I. Griffiths, S. Creer, A. Smith,
B. A. Emmett, and D. A. Robinson. 2021. Long-Term Drought and Warming Alter Soil
Bacterial and Fungal Communities in an Upland Heathland. Ecosystems:1–16.

Sepulveda, J., and A. H. Moeller. 2020, March 10. The Effects of Temperature on Animal Gut
Microbiomes. Frontiers Media S.A.

Shantz, A. A., N. P. Lemoine, and D. E. Burkepile. 2016. Nutrient loading alters the performance
of key nutrient exchange mutualisms. Ecology Letters 19:20–28.

Sheik, C. S., W. H. Beasley, M. S. Elshahed, X. Zhou, Y. Luo, and L. R. Krumholz. 2011. Effect
of warming and drought on grassland microbial communities. ISME Journal 5:1692–1700.

Simonsen, A. K. 2021. Environmental stress leads to genome streamlining in a widely distributed
species of soil bacteria. The ISME Journal 16:423–434.

Simonsen, A. K., and J. R. Stinchcombe. 2014. Herbivory eliminates fitness costs of mutualism
exploiters. New Phytologist 202:651–661.

Swift, J. F., M. R. Kolp, A. Carmichael, N. E. Ford, P. M. Hansen, B. A. Sikes, M. Kleiner, and
M. R. Wagner. 2024. Drought stress homogenizes maize growth responses to diverse natural
soil microbiomes. Plant and Soil.

Terpolilli, J. J., G. W. O'Hara, R. P. Tiwari, M. J. Dilworth, and J. G. Howieson. 2008. The
model legume *Medicago truncatula* A17 is poorly matched for N2 fixation with the
sequenced microsymbiont *Sinorhizobium meliloti* 1021. New Phytologist 179:62–66.

Thuiller, W. 2007. Climate change and the ecologist. Nature 448:550–552.

Trivedi, P., B. D. Batista, K. E. Bazany, and B. K. Singh. 2022. Plant–microbiome interactions
under a changing world: responses, consequences and perspectives. New Phytologist
234:1951–1959.

VanWallendael, A., G. M. N. Benucci, P. B. da Costa, L. Fraser, A. Sreedasyam, F. Fritschi, T. E.
Juenger, J. T. Lovell, G. Bonito, and D. B. Lowry. 2022. Host genotype controls ecological
change in the leaf fungal microbiome. PLOS Biology 20:e3001681.

Werner, G. D. A., and E. T. Kiers. 2015. Partner selection in the mycorrhizal mutualism. New
Phytologist 205:1437–1442.

Wickham, H., M. Averick, J. Bryan, W. Chang, L. McGowan, R. François, G. Grolemund, A.
Hayes, L. Henry, J. Hester, M. Kuhn, T. Pedersen, E. Miller, S. Bache, K. Müller, J. Ooms,
D. Robinson, D. Seidel, V. Spinu, K. Takahashi, D. Vaughan, C. Wilke, K. Woo, and H.
Yutani. 2019. Welcome to the Tidyverse. Journal of Open Source Software 4:1686.

Wilke, C. 2017. cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2."

Xie, Y., W. Liu, P. Li, S. Bai, D. Li, L. Zhang, H. Sun, Y. Zheng, Q. Cheng, C. Wang, E. Du, and
W. Gou. 2023. Soil bacterial community structure at different plant maturity stages in an
annual grass–legume production system. Frontiers in Sustainable Food Systems 7.

Yang, L., Y. Zhou, S. Zhu, T. Huang, L. Wu, and X. Yan. 2012. Detection and Quantification of
Bacterial Autofluorescence at the Single-Cell Level by a Laboratory-Built High-Sensitivity
Flow Cytometer. Analytical Chemistry 84:1526–1532.

Zhang, W., X. G. Li, K. Sun, M. J. Tang, F. J. Xu, M. Zhang, and C. C. Dai. 2020. Mycelial
network-mediated rhizobial dispersal enhances legume nodulation. ISME Journal 14:1015–
1029.

Zhou, Z., C. Wang, and Y. Luo. 2020. Meta-analysis of the impacts of global change factors on
soil microbial diversity and functionality. Nature Communications 11:3072.

Zipfel, C., and G. E. D. Oldroyd. 2017. Plant signalling in symbiosis and immunity. Nature
543:328–336.

Bacterial rhizosphere

46 samples & 425 taxa (Genus). PCA tax_transform=clr