Remote-control meiotic drive of sex chromosomes

Naomi L. Greenberg[1,#] (nlg34@georgetown.edu; ORCID: 0009-0008-6880-4562)

Manus M. Patten[1] (mmp64@georgetown.edu; ORCID: 0000-0003-2890-7977)

Martijn A. Schenkel[1,2,3,*] (maschenkel@gmail.com; ORCID: 0000-0002-7721-319X)

<u>Affiliations</u>

[1] Department of Biology, Georgetown University, 37th and O St. NW, Washington DC, 20057, USA

[2] Groningen Institute of Evolutionary Life Sciences, University of Groningen, Nijenborgh 7, 9747 AG,
Groningen, The Netherlands

[3] Institute of Organismic and Molecular Evolution, Johannes Gutenberg-Universität Mainz, Hanns-Dieter-Hüsch-
Weg 15, 55128, Mainz, Germany.

[#] Current address: Department of Life Sciences, Imperial College, London SW7 2AZ, UK

[*] Correspondence: Martijn A. Schenkel (maschenkel@gmail.com)

Abstract

Some selfish genetic elements drive at meiosis to achieve transmission distortion, breaking the rules of Mendelian
segregation to enhance their own evolutionary success. It has been shown that enhancers of drive must act in *cis*
in order to gain the selfish benefit of drive and that suppressors of drive will be selected at unlinked loci. Here,
we model the evolution of an autosomal *trans*-acting gene (*Distorter*) that causes the Y-chromosome (or even 0-
chromosome) to drive without driving itself, a phenomenon we call "remote-control meiotic drive". We show that
such a gene may spread in the population when linked to a second locus, *Assister*, whose alleles are transmitted
at different frequencies through sperm as compared to eggs, for which we consider various scenarios, e.g. sexually
antagonistic selection or sex-limited drive. Depending on the mechanistic details of sex-chromosome drive,
Distorter's spread can additionally facilitate transitions between XY and X0 sex determination. Our results
provide a proof of principle that stretches the current understanding of segregation modifier and sex allocation
theory. Moreover, we identify alternative evolutionary trajectories that could also lead to remote control drive,
and discuss its potential applications in developing synthetic sex-ratio-distorting elements for use in e.g. pest
management.

Introduction

Mendel's Law of Segregation states that in a diploid individual, the two alleles at a locus segregate to an equal (i.e., 50/50) proportion of the gametes (Mendel, 1865). The fair transmission of genetic material can be distorted in myriad ways (Burt & Trivers, 2006; Patten *et al.*, 2023), such as by true meiotic drivers (Clark & Akera, 2021) and, to similar effect, post-segregational killers (Bravo Núñez *et al.*, 2018), which often fall under the moniker of "meiotic drive" themselves. The former achieve their advantage in asymmetric meiosis by segregating preferentially to the egg pole during meiosis and avoiding the polar body (Sandler & Novitski, 1957). The latter target their allelic counterpart just after meiosis, e.g. by preventing the formation of functional gametes that lack the transmission-distorting gene (Lindholm *et al.*, 2016; Price *et al.*, 2020). Consequently, the gametes produced by heterozygotes do not feature equal proportions of both alleles; instead the transmission-distorting allele becomes overrepresented, and the non-distorting allele underrepresented. Meiotic drivers typically harm the fitness of their bearers, often by causing reduced fertility (Lyttle, 1991; Larracuente & Presgraves, 2012). In the case of sex-chromosome drive, additional harm comes from overproducing the common sex (Lyttle, 1991). In either case, their ability to cheat the Law of Segregation offsets this cost to themselves, enabling them to spread and persist in the population.

In males, meiotic drivers may achieve their higher-than-Mendelian transmission using two mechanisms, both of which entail a supergene that consists of two distinct loci. First, the toxin-antidote system (Burga *et al.*, 2020) features a toxin that is expressed during or prior to meiosis, so that all resulting gametes receive a dose. In gametes that bear the toxin-antidote supergene, the antidote is expressed post-meiosis to prevent the harmful effects of the toxin and rescue the supergene-bearing gametes. The second system is the killer-responder system, which produces a killer element that targets a responder locus (Meiklejohn & Tao, 2010). In this case, the killer allele is linked to an insensitive variant of the locus that is targeted by the killer, whereas the non-killer allele is assumed to be linked to a sensitive variant and is therefore eliminated. Gametes that lack the supergene (or more specifically, the antidote allele or the insensitive responder allele) are eliminated, so that the gamete pool produced consists predominantly of gametes bearing the segregation-distorting complex at the supergene.

A key feature of both mechanisms is that the targeted genetic element is the chromosomal complement of the driver, meaning segregation distortion acts to remove these from the gamete pool to enhance its own odds of being transmitted. Theoretical models have explored how this dynamic affects the potential for genes that modify the severity of segregation distortion. According to modifier theory, genetic modifiers that arise linked to the segregation-distorting complex would typically act to enhance the degree of segregation distortion, whereas modifiers linked to the targeted gene complex would be expected to diminish its effect (Prout *et al.*, 1973; Feldman & Otto, 1991). Effectively, this constitutes the formation of antagonistic, multigenic collaborations to cheat Mendelian transmission or enforce it, respectively. Unlinked modifiers, i.e. those occurring on other linkage groups, are also typically thought to favor diminished segregation distortion owing to the deleterious effects segregation distorters have on their bearer's fitness (Eshel, 1985; Veller, 2022). These dynamics underlie the so-called "parliament of genes" effect which seeks to explain why so few genes succeed in being selfish (Leigh, 1971). Only the select few genes linked to a selfish genetic element like a segregation distorter are selected to enhance it, but the remainder of the genome is selected to suppress it. Through sheer numbers, the latter typically ends up as the winner.

As a variation on the above, we could consider a segregation distorter that targets not its own
chromosomal complement, but rather one of another linkage pair. One hypothetical example of this is an
autosomal killer gene that targets a responder gene on one of the sex chromosomes; this might also be considered
a modifier gene that affects the segregation of the sex chromosome pair (Maffi & Jayakar, 1981). Such a gene
would normally be counter-selected for two reasons. The first is that by biasing the sex ratio among its bearer's
offspring, it ultimately ends up in the overrepresented sex (Lyttle, 1991; Jaenike, 2001). The second is that the
segregation distorter would target gametes containing copies of itself half of the time (i.e., "suicide combinations";
Hartl, 1974). Such *trans*-acting or "remote-control" meiotic drive has been largely ignored, as the lack of benefit
obtained from eliminating the competition and potential additional costs to organismal fitness of such a
segregation distorter are thought to render it unlikely to evolve and spread. In this manuscript, we identify a route
by which such a mechanism may overcome these costs and lead to the evolution of autosomal genes for sex
chromosome meiotic drive.

Drive may be found more commonly on sex chromosomes than autosomes for two reasons, one owing
to genetics and the other to ascertainment bias. First, the X- and Y-chromosomes do not typically recombine for
much of their length, which prevents the separation of the alleles that bring about segregation distortion when
found in tandem (Hartl, 1974; Haig & Grafen, 1991). Repeated cycles of drive and suppression, which may be
common on the X- and Y-chromosomes for this reason, may lead to the evolution of many cryptic drivers on the
sex chromosomes (Ellison & Bachtrog, 2019; Bachtrog, 2020). Second, before suppressors of sex-chromosome
drive evolve, they are more readily ascertained than autosomal segregation distorters, as skewed transmission of
the sex chromosomes brings about a conspicuous shift in the offspring sex ratio (Hamilton, 1967; Jaenike, 2001,
2008). Further, this sex ratio effect can maintain drivers in a polymorphic state owing to the frequency dependence
of fitness. The sex chromosomes may therefore be enriched for both fixed cryptic drivers as well as polymorphic
drivers.

Sex ratio biases brought about by drive can promote evolutionary transitions between sex determination
mechanisms (Kozielska *et al.*, 2010). For example, a driving X-chromosome may lead to female-biased sex ratios,
promoting the evolution of a male-determining gene on an autosome, converting it into a proto-Y-chromosome.
Autosomal driving elements may also promote the origin of sex chromosomes (Úbeda *et al.*, 2014). Owing to
differences between oogenesis and spermatogenesis, most meiotic drivers exhibit sex-specific (or sex-biased)
segregation distortion, so that one sex may transmit it at a Mendelian rate, whereas in the other it distorts
segregation to achieve higher transmission. If a locus linked to this sex-specific driver evolves a novel, sex-
determining mutation, this confines the driving allele to the sex in which it achieves segregation distortion more
frequently. As this supergene spreads in the population, the segregation-distorting effect causes sex-ratio biases
(similar to Kozielska *et al.* (2010)), leading to a series of selectively favored mutations that eventually yield a set
of sex-determining chromosomes with normal, Mendelian segregation.

The origin of a sex-determining gene next to a segregation-distorting gene effectively restricts the
distorter to one sex, forming a synergistically-acting supergene. We posit here that achieving this effect does not
explicitly require a sex-determination gene, but rather that biasing transmission to one sex more generally enables
this to occur. That is, any gene that skews the sex ratio of its bearers' offspring would produce a similar effect.
Ordinarily, a sex-ratio-biasing autosomal gene would be harmful to its bearers, and hence be purged (per Fisher

(1930), but see West (2009)). But a sex-ratio-biasing gene paired with a sex-specific meiotic driver establishes a
synergism that offsets such costs and might therefore be able to spread and persist.

Here, we model the evolution of an autosomal, *trans*-acting segregation distorter (*Distorter*) that biases
the transmission of the sex chromosomes. We consider two potential proximate mechanisms by which it achieves
this effect, which we dub X-shredding and X-conversion, further explained below. Linked to this *trans*-acting
segregation distorter is a secondary locus (*Assister*), which may represent: (1) a locus under sexually antagonistic
selection, where different alleles are favored in females as compared to males (van Doorn, 2009; Schenkel *et al.*,
2018) or (2) a sex-specific meiotic driver, an analog to the situation modeled by Úbeda *et al.* (2014). In the
Supplement we consider two other scenarios, one in which the *Assister* locus is parentally antagonistic, wherein
an allele's fitness depends on its parental origin (Patten *et al.*, 2013; Haig *et al.*, 2014), and a second in which the
Assister locus simply causes co-segregation of the autosome and the Y-chromosome. The underlying rationale for
these scenarios is that in all cases, the *trans*-acting segregation-distorting gene and the linked gene have the same
optimal destination—namely, a male offspring (Gardner & Úbeda, 2017). In addition, these phenomena have
previously been shown capable of spurring transitions between sex-determining mechanisms and of affecting sex
ratio adjustment (van Doorn & Kirkpatrick, 2007, 2010; Blackburn *et al.*, 2010; Kozielska *et al.*, 2010; Úbeda *et*
al., 2014; Muralidhar & Veller, 2018; Schenkel, 2024). We use our model to determine the conditions under which
such a gene can spread in the population, what frequency it reaches at equilibrium, and what level of sex-ratio
distortion is achieved once equilibrium is attained by causing increased transmission of the Y-chromosome over
the X-chromosome. We also consider whether it can spur a transition between genetic sex-determination
mechanisms, where this interference results in the formation of 0-chromosomes, whose spread in the population
leads to a loss of the Y-chromosome. Through this, we show that autosomal genes may interfere with the
segregation of the sex chromosomes to promote their own spread in the population. This results in the sex
chromosomes appearing to drive through no effort of their own—i.e., through remote control. We discuss our
results in light of previous modifier theory of segregation distortion, sex allocation theory, definitional issues of
gene drives, and in the context of potential applications.

Methods
Our model assumes an infinite population with XY sex determination (females XX, males XY), with an initially
equal sex ratio. The X- and Y-chromosomes determine sex and do not carry any other genes relevant to our model.
Terms such as 'female' and 'male', and similarly 'XY' versus 'ZW' can be exchanged to consider cases with
female heterogamety, rather than male heterogamety. In addition to the XY sex chromosomes, individuals carry
an autosome carrying two loci, dubbed *Distorter* (with alleles D_1 and D_2) and *Assister* (with alleles A_1 and A_2);
recombination between *Distorter* and *Assister* occurs at a rate r in both sexes (Figure 1A). The D_1 allele makes
for equal segregation of the sex chromosomes, whereas the D_2 allele interferes with the normal (i.e., Mendelian)
segregation of the XY sex chromosomes in males. Further details on D_2's mechanism are provided under "*Trans*-
segregation distortion by *Distorter*" below.

For the *Assister* locus, we focus in the main text on two selective scenarios, sexually antagonistic
selection and male-limited *cis*-segregation distortion, and examine the cases of parentally antagonistic selection
and of co-segregation in the Supplementary Material. More details of the two schemes we examine in the main
text are provided in "Sexually antagonistic selection on *Assister*" and "*Cis*-segregation distortion by *Assister*",

Figure 1: Model overview. (A) Our model incorporates two linked autosomal loci, *Assister* (with alleles A_1 and A_2) and *Distorter* (with alleles D_1 and D_2); recombination between *Assister* and *Distorter* occurs at a rate r in both sexes. *Assister* may represent either a sexually antagonistic locus (A_1 beneficial in females/detrimental in males, A_2 detrimental in females/beneficial in males) or a locus experiencing *cis*-segregation distortion (D_2 overtransmitted in D_1/D_2 heterozygotes). The D_2 allele interferes with the Mendelian transmission of the X-, Y-, and/or 0-chromosomes during spermatogenesis to achieve *trans*-segregation distortion.

respectively. The crux of each of these selective scenarios is that the A_2 allele experiences higher long-term fitness
by segregating to males, either because: (1) this is the sex in which it exhibits higher-than-Mendelian transmission
(segregation distortion); or (2) it enhances the sex of its bearer (sexual antagonism); or (3) it comes to be
transmitted in a manner that enhances its fitness in the subsequent generation (parental antagonism). This favors
the buildup of linkage disequilibrium between *Assister* and *Distorter*, as well as statistical linkage between this
pair of loci and the Y-chromosome. Consequently, the *trans*-segregation distorting allele D_2 can spread via its
association with the A_2 allele and its overrepresentation in males. We measure linkage disequilibrium in sperm to
track these non-random associations, using the following equations (Rydzewski *et al.*, 2016):

$$Ds_{AD} = (A_1D_1X + A_1D_1Y)(A_2D_2X + A_2D_2Y) - (A_2D_1X + A_2D_1Y)(A_1D_2X + A_1D_2Y)$$
$$D_{sAY} = (A_2D_1Y + A_2D_2Y)(A_1D_2X + A_1D_1X) - (A_1D_1Y + A_1D_2Y)(A_2D_1X + A_2D_2X)$$
$$D_{sDY} = (A_1D_2Y + A_2D_2Y)(A_1D_1X + A_2D_1X) - (A_1D_1Y + A_2D_1Y)(A_1D_2X + A_2D_2X)$$

And for the X-conversion model, which additionally includes 0-chromosomes as a third allele at the sex
chromosomes:

$$Ds_{AD} = (A_1D_1X + A_1D_1Y + A_1D_10)(A_2D_2X + A_2D_2Y + A_2D_20)$$
$$- (A_2D_1X + A_2D_1Y + A_2D_10)(A_1D_2X + A_1D_2Y + A_1D_20)$$
$$D_{sAY0} = (A_2D_1Y + A_2D_2Y + A_2D_10 + A_2D_20)(A_1D_2X + A_1D_1X) - (A_1D_1Y + A_1D_2Y + A_1D_10$$
$$+ A_1D_20)(A_2D_1X + A_2D_2X)$$
$$D_{sDY0} = (A_1D_2Y + A_2D_2Y + A_1D_20 + A_2D_20)(A_1D_1X + A_2D_1X)$$
$$- (A_1D_1Y + A_2D_1Y + A_1D_10 + A_2D_10)(A_1D_2X + A_2D_2X)$$

The overall life cycle is as follows: mature males and females undergo meiosis to produce their respective
gamete pools; during this time, segregation distortion takes place; gametes are then united at random; the resulting
juvenile offspring undergo viability selection during maturation; and upon reaching maturity they replace their
parents as the reproductive population so that generations are non-overlapping. At this point, the cycle resets. Our

population is initially fixed for the D_1 allele, and with A_1 and A_2 at or near their equilibrium as achieved by 500

"burn-in" generations of simulation. We then mutate a small ($p=0.001$) proportion of D_1 alleles to D_2 alleles on

each genetic background in Y-bearing sperm simultaneously (proportional to their frequency in the population at

the time) to prevent any advantage D_2 might experience from arising under linkage disequilibrium. The eventual

results did not differ when we introduced D_2 on alternative genetic backgrounds instead. We then continue the

simulation procedure for 1,000 generations before we evaluate the terminal frequency of D_2.

In our simulations, we vary the parameter values for the selective effects of the *Assister* and *Distorter*

loci, as well as their capacity to distort allelic segregation during meiosis (where applicable); details on these are

provided under "Results". We focus here in the main text primarily on the impact of the strength of selection on

Assister (for sexually antagonistic selection) or its capacity to achieve higher-than-Mendelian transmission (for

cis-segregation distortion), as well as the strength of *trans*-segregation distortion achieved by *Distorter*. We

consider the role of recombination between the *Assister* and the *Distorter* loci and direct selection against *Distorter*

more fully in the Supplement.

An overview of all model variables, their descriptions, and standardized values is included in

Supplementary Table S1.

<u>Sexually antagonistic selection on *Assister*</u>

Under sexually antagonistic selection, the A_1 and A_2 alleles differently affect the fitness of female versus male

bearers. We assume that A_1 is beneficial to female fitness but costly to male fitness, and vice versa for A_2. The

partial fitness score $w_{F_{ij}}$ of a female with a genotype $A_i A_j$ is given by:

$$w_{F_{11}} = 1$$

$$w_{F_{12}} = w_{F_{21}} = 1 - h_A t$$

$$w_{F_{22}} = 1 - t$$

The partial fitness score $w_{M_{ij}}$ of a male with a $A_i A_j$ genotype is given by:

$$w_{M_{11}} = 1 - s$$

$$w_{M_{12}} = w_{M_{21}} = 1 - (1 - h_A)s$$

$$w_{M_{22}} = 1$$

Here, t and s indicate the homozygous fitness cost of A_2 in females and A_1 in males, respectively, and $h_A \in [0,1]$

is the dominance of A_2 in heterozygotes.

<u>*Cis*-segregation distortion by *Assister*</u>

As a segregation-distorting gene, the A_2 allele of the *Assister* locus is transmitted at a higher-than-Mendelian rate

in heterozygotes. Specifically, we assume that an $A_2 A_1$ (or $A_1 A_2$) heterozygote transmits the A_2 allele at a rate of

$\frac{1}{2} + k_A$ and the A_1 allele at a rate $\frac{1}{2} - k_A$ where $k_A \in (0, \frac{1}{2}]$ indicates the strength of *cis*-segregation distortion by A_2

and $k_A = 0$ represents Mendelian segregation. We assume that the A_2 allele negatively affects its bearer's fitness

in females and males alike. The partial fitness score w_{ij} of an individual with genotype $A_i A_j$ is given by:

$$w_{11} = 1$$

$$w_{12} = w_{21} = 1 - h_A s$$

$$w_{22} = 1 - s$$

Here, $s \in [0,1]$ represents the fitness effect of the A_2 allele in homozygotes, $h_A \in [0,1]$ its dominance in

heterozygotes. We chose this fitness scheme for *Assister* to reflect some well-known autosomal drivers (e.g., the

t-haplotype in mouse and *SD* in *Drosophila*, which have linked recessive lethality/sterility mutations) and also

because it maintains polymorphism at the *Assister* locus, which is essential for our model.

Trans-segregation distortion by *Distorter*

The D_2 allele of the *Distorter* locus acts to distort the segregation rate of the sex chromosomes. We assume that

D_2 acts only in males, i.e. the heterogametic sex, as a dominant *trans*-acting segregation distorter, such that D_2D_1

heterozygotes and D_2D_2 homozygotes transmit their sex chromosomes at equally distorted frequencies. We

consider two mechanisms of *trans*-segregation distortion, denoted X-shredding and X-conversion, respectively.

X-shredding represents a skew in the transmission of X- versus Y-chromosomes to future offspring, where the Y-

chromosome is transmitted at a rate $\frac{1}{2} + k_D$ and the X chromosome at a rate $\frac{1}{2} - k_D$, where $k_D \in \left(0, \frac{1}{2}\right]$; this is

achieved by the killing off of some fraction of X-bearing sperm. Under X-conversion, D_2 acts to eliminate the X-

chromosome from a certain number of X-bearing sperm to convert them into 0-bearing gametes that, when fused

with an X-bearing egg yield a viable, fertile X0 male (note that this assumes that maleness is determined by lack

of a second X-chromosome, rather than some dominant male-determining gene on the Y-chromosome, in which

case X0 individuals would develop as females). Consequently, the Y-chromosome is transmitted at a Mendelian

rate of $\frac{1}{2}$, the X-chromosome is transmitted at a rate $\frac{1}{2} - k_D$, and the remaining k_D represents the proportion of

sperm that have been converted from being X-bearing to 0-bearing. In X0 males, the 0-chromosome would be

transmitted at a rate $\frac{1}{2} + k_D$ and the X-chromosome at a rate $\frac{1}{2} - k_D$. We assume that lacking the Y-chromosome

comes at some fitness cost $v \in (0,1)$ to males, so XY males have a relative fitness score of 1 and X0 males have

a fitness of $1 - v$. Potentially, Y-chromosome-bearing males may also experience reduced fitness owing to the

accumulation of mutations on the Y-chromosome, resulting in an effective fitness benefit for X0 males ($v < 0$);

we consider this scenario in the Discussion.

We assume that the D_2 allele is harmful to male fertility through its sperm-killing or X-conversion efforts.

The resulting partial fitness score w_{ij} of a male with a D_iD_j genotype is given by:

$$w_{11} = 1$$

$$w_{12} = w_{21} = 1 - z$$

$$w_{22} = 1 - z$$

Here, $z \in [0,1]$ measures the loss of total fertility, and we assume that the *Distorter* locus is neutral in females.

Finally, we assume that the partial fitness effects of the *Assister* and *Distorter* loci are multiplicative;

Supplementary Table S2 provides an overview of the fitnesses of all potential genotypes in our population.

Data analysis and visualization

Simulations, data analysis, and visualization were carried out using R (R Development Core Team, 2023; v.4.4.1)

and RStudio (RStudio Team, 2023; v.2024.09.0+375) using the 'tidyverse' and 'viridis' packages (Garnier, 2018;

Wickham *et al.*, 2019).

Results

<u>Sexually antagonistic selection on *Assister* enables the spread of *Distorter*</u>

We first explored whether an autosomal *trans*-acting segregation distorter (i.e., the D_2 allele of the *Distorter* locus)

of the Y-chromosome could spread in the population when linked to a sexually antagonistic locus (the *Assister*

Figure 2. Dynamics and scope of Y-chromosome drive via the *trans*-acting *Distorter*. Here, *Distorter* is linked to a sexually antagonistic *Assister* locus and causes drive via the X-shredding mechanism. (A) Changes in the frequency of the A_2 and D_2 alleles and the X- and Y-chromosomes in males across 500 generations. (B) Changes in the linkage disequilibrium of *Assister-Distorter*, *Assister*-XY, and *Distorter*-XY across 500 generations. (C) Haplotype frequencies over 500 generations. A_1D_2X and A_1D_2Y have near-zero frequencies. The shaded area represents the frequency of D_2 and primarily comprises the haplotypes A_2D_2Y and A_2D_2X. (D) D_2 equilibrium frequency after 1,000 generations. (E) Population sex ratio after 1,000 generations. Parameter values (if not otherwise specified): $k_D = 0.25, s, t = 0.5, h_A = 0.5, z = 0.001, r = 0.001$.

locus) under rather favorable conditions. With strong sexually antagonistic selection ($s, t = 0.5$), a high level of *trans*-acting segregation distortion ($k_D = 0.25$) but low fitness costs thereof ($z = 0.001$), and tight linkage between *Assister* and *Distorter* ($r = 0.001$), we found that D_2 can invade and reach a frequency of ~0.13, enhancing Y-chromosome transmission so that it reaches a frequency – and hence a population sex ratio – of 0.57 (Figure 2A). Meanwhile, the *Assister* locus remains polymorphic, with A_2's frequency changing only slightly from 0.58 at its initial equilibrium to a frequency of 0.55 after 500 generations (Figure 2A).

We simultaneously tracked the linkage between *Assister*, *Distorter*, and the XY chromosomes (Figure 2B). These three loci come into close statistical linkage and are inherited non-independently—effectively, as a group—which creates the following reinforcing benefits: D_2 to benefit from the male-beneficial fitness effects of A_2; the Y-chromosome to benefit from its increased transmission thanks to D_2; and A_2 to benefit from being assigned to its fitness-enhancing state (i.e., males) thanks to a stronger association with the Y chromosome.

To better characterize the conditions that favor the spread of D_2, we fully explored the parameter space of sexual antagonism (s and t) at the *Assister* locus and the strength of drive (k) of the D_2 allele. We found that D_2 can spread under most parameter value combinations considered here (Figure 2D), though it achieves the highest equilibrium frequency when *trans* drive is relatively weak ($k_D \approx 0.1$) and sexually antagonistic selection is strong ($s, t \cong 1$). The presence of D_2 in the population leads to biased sex ratios, which are highest for high levels of *trans* drive and strong sexually antagonistic selection (Figure 2E). When sexually antagonistic selection is relatively weak, we see that higher *trans*-segregation distortion does not lead to concomitantly higher frequencies of D_2. Under these conditions, the cost to D_2 of assigning itself to the common sex outstrip the benefits linkage to A_2.

We also investigated the constraints that higher recombination rates and higher D_2 fertility costs place on the spread of D_2. We found that higher recombination ($r = 0.01$ or $r = 0.1$) reduced the opportunity for male-biased sex ratios, likely by reducing the necessary *Assister*-*Distorter*-XY linkage disequilibrium; similarly, a higher D_2 fertility cost reduced its success (Supplementary Figure S1).

Cis- and *trans*-segregation distortion can work in tandem to achieve sex ratio biases

We then explored whether a male-limited, *cis*-acting segregation distorter at the *Assister* locus could similarly favor the spread of D_2. We found that for a large range of values of *cis* drive strength (k_A) and *trans* drive strength (k_D), D_2 spreads and the population skews male. Similar to the sexual antagonism model above, the Y-chromosome as well as D_2 increase in frequency, whereas A_2's frequency remains relatively stable (Figure 3A). The linkage disequilibrium between *Assister* and *Distorter* is strong, as associations between the Y-chromosome and both D_2 and A_2 develop (Figure 3B). D_2 occurs primarily in combination with A_2 (Figure 3C). We investigated the full range of drive strengths ($k_A \in [0, 0.5], k_D \in [0, 0.5]$) at both the *Distorter* locus and the *Assister* locus (Figures 3D, 3E). We found that most of the parameter space permits invasion of D_2, and some parameter combinations enable high frequencies of D_2 and a high sex-ratio skew. With freer recombination between *Assister* and *Distorter* and with elevated fitness costs of D_2, the opportunity for D_2's spread diminishes, and the resulting sex-ratio skew is less extreme (Supplementary Figure 2).

Figure 3. Dynamics and scope of Y-chromosome drive via the *trans*-acting *Distorter* causing X-shredding, linked to a male-limited meiotic driver *Assister*. (A) Changes in the allele frequency of A_2, D_2, and the X- and Y-chromosome in males across 500 generations. (B) Changes in the linkage disequilibrium of *Assister-Distorter*, *Assister*-XY, and *Distorter*-XY across 500 generations. (C) Haplotype frequencies over 500 generations. A_1D_2X and A_1D_2Y have near-zero frequencies. The shaded area represents the frequency of D_2 and primarily comprises the haplotypes A_2D_2Y and A_2D_2X. (D) D_2 equilibrium frequency after 1,000 generations. (E) Population sex ratio after 1,000 generations. Parameter values (if not otherwise specified): $k_D = 0.25, k_A = 0.25, h_A = 0.0, z = 0.01, r = 0.001, s = 0.5$.

288 <u>X-conversion leads to the emergence of X0 sex determination through loss of the Y-chromosome</u>

289 In the previous sections, we assumed that D_2 distorts the segregation of the X- and Y-chromosomes by the

290 selective killing of X-bearing sperm. An alternative mechanism by which D_2 might generate a male-biased sex

291 ratio among its bearer's offspring is to selectively eliminate the X-chromosome from sperm to convert them to 0-

Figure 4. Dynamics and scope of 0-chromosome drive via the *trans*-acting *Distorter* causing X-conversion, linked to a sexually-antagonistic *Assister*. (A) Changes in the allele frequency of A_2, D_2, and the X- and Y-chromosome in males across 500 generations. (B) Changes in the linkage disequilibrium of *Assister-Distorter*, *Assister*-XY0, and *Distorter*-XY0 across 500 generations. (C) Haplotype frequencies over 500 generations. A_1D_2X, A_1D_2Y, A_2D_2Y, and A_1D_20 have near-zero frequencies. The shaded area represents the frequency of D_2 and primarily comprises the haplotypes A_2D_2Y and A_2D_2X. (D) D_2 equilibrium frequency after 1,000 generations. (E) Population sex ratio after 1,000 generations. Parameter values (if not otherwise defined): $k_D = 0.25, s, t = 0.5, h_A = 0.5$, $z = 0.01, r = 0.001, v = 0.001$.

292   bearing sperm, yielding X0 sons instead of XY sons (note however that this assumes that sex determination occurs

293   through some dosage effect of the X-chromosomes as in *Drosophila* fruit flies (Pomiankowski *et al.*, 2004); if

294   maleness instead is conferred by a Y-chromosomal dominant male-determiner, an X0 genotype would be expected

295   to lead to femaleness (Graves, 2006), constraining the spread of D_2). This mechanism might not harm male fertility

296   as severely as the X-shredding mechanism, and so is poised to spread more easily. This form of *trans*-segregation

Figure 5: Dynamics and scope of 0-chromosome drive via the *trans*-acting *Distorter* causing X-conversion, linked to a male-limited meiotic driver *Assister*. (A) Changes in the allele frequency of A_2, D_2, and the X-, Y-, and 0-chromosome in males across 500 generations. (B) Changes in the linkage disequilibrium of *Assister-Distorter*, *Assister*-XY0, and *Distorter*-XY0 across 500 generations. (C) Haplotype frequencies over 500 generations. A_1D_2X, A_1D_2Y, A_2D_2Y, and A_1D_20 have near-zero frequencies. The shaded area represents the frequency of D_2 and primarily comprises the haplotypes A_2D_20 and A_2D_2X. (D) D_2 equilibrium frequency after 1,000 generations. (E) Population sex ratio after 1,000 generations. Parameter values (if not otherwise defined): $k_D = 0.25$, $k_A = 0.25$, $h_A = 0$, $z = 0.01$, $r = 0.001$, $v = 0.001$.

distortion also provides a mechanism for an XY sex determination system to transition to an X0 sex determination system. We find that when the D_2 allele causes X-conversion, the resulting 0-chromosomes accumulate and become the predominant male-determining factor. This holds true when *Assister* experiences sexual antagonism (Figure 4), as well as when it exhibits male-limited *cis* drive (Figure 5). In both cases, we find that the 0-chromosome can entirely replace the Y-chromosome. There is only a small range of parameter space in which the

Y-chromosome escapes loss and is maintained alongside the 0 chromosome in the population (Supplementary
Figure S5).

Discussion

Here, we have shown that some *trans*-acting segregation distorters, which target an unlinked locus on the X-
chromosome, can spread when they arise on a suitable genetic background; we refer to this novel form of
segregation distortion as "remote-control meiotic drive". Thus, sex-chromosomal segregation distortion need not
be caused solely by sex-linked loci. Additionally, our results run counter to the conclusion one may draw from
previous theory, which finds that unlinked loci should ordinarily suppress drive (Eshel, 1985; Veller, 2022). We
also found that such *trans*-acting distorters could spur a transition in sex chromosome systems from XY to X0
male heterogamety.

We focused on two particular scenarios, where a *trans*-acting segregation distorter (*Distorter*) arises near
a locus (*Assister*) that is either under sexually antagonistic selection or exhibits sex-specific *cis*-segregation
distortion (in the supplement we consider a third scenario, where the nearby locus experiences parentally
antagonistic selection, and a fourth scenario, where the autosomal locus co-segregates with the Y-chromosome).
Both genetic backgrounds lead to linkage disequilibrium between *Distorter* and *Assister*, where the *trans*-acting
segregation-distorting allele D_2 becomes associated with A_2, the male-beneficial allele or the *cis*-segregation
distorting allele in the two respective scenarios. Under either scenario, A_2 has the same optimal destination: a male
individual (Figure 6A). If *Assister* is sexually antagonistic, A_2 will act to enhance its bearer's fitness—and hence
its own fitness—when it ends up in a male as opposed to a female. If *Assister* is parentally antagonistic (see
Supplementary Material), A_2 will come to be paternally transmitted, and hence will enhance the fitness of its future
bearers; this is effectively the same effect as for sexual antagonism, but with the fitness effects delayed by one
generation. Lastly, when *Assister* is a sex-specific segregation distorter, A_2 will exhibit a higher-than-Mendelian
transmission rate when its bearer is a male, as opposed to a fair transmission rate of 50% in females. In all
scenarios, D_2 acts to increase its own effective transmission to males by skewing the sex ratio of its bearer's
offspring. The cost to D_2 of assigning itself to the more common sex is offset by the linked benefit of delivering
an A_2 allele to its favored destination (Figure 6B).

The two main scenarios for *Assister* (sexually antagonistic selection *versus* *cis*-acting drive) have
different implications. Empirical studies of these phenomena find that many genes may be under sexually
antagonistic selection, but the selective effects of singular loci might be small (Rice, 1992; Innocenti & Morrow,
2010; Schenkel *et al.*, 2018). In contrast, segregation distorters are thought to be less common (though many may
be cryptic; Werren, 2011), but often involve strong levels of segregation distortion and/or more severe fitness
effects (Burt & Trivers, 2006; Lindholm *et al.*, 2016). For these reasons, sexually antagonistic selection may
enable the evolution of many *Distorter*-like loci throughout the genome, each with either low frequencies of the
trans-driving alleles or with relatively weak effects thereof. In contrast, segregation distorters might provide fewer
opportunities for such loci to evolve, but those that do might harbor stronger or more frequently-occurring *trans*-
acting drivers.

Mechanistically, a *trans*-acting sex-chromosome driver may seem rather unlikely. But, on the contrary,
it is actually quite easy to obtain. Consider a traditional Y-linked driver that results in biased sex ratios but reaches
fixation despite assigning itself to the more common sex. Such sex-ratio biases cause selection to favor the

Figure 6: (A) A_2 experiences fitness benefits by preferentially segregating to males under each evolutionary scenario for *Assister*. Sexual antagonism (SA): A_2 confers a fitness benefit so that male bearers have higher-than-average fitness (w), whereas female bearers have lower-than-average fitness. Male-limited *cis*-segregation distortion (SD): A_2 has a higher-than-Mendelian transmission (p) in males, but regular Mendelian transmission in females. Parental antagonism (PA): A_2 confers a fitness benefit to its bearers when paternally inherited so that they have higher-than-average fitness, whereas it confers a fitness cost when maternally-inherited. (B) A_2 and D_2 each benefit from the other. A_2 provides D_2 a fitness or transmission benefit, and D_2 provides A_2 a "destination benefit," i.e. indirectly delivers it to a destination where it fares better (i.e., in a male). This synergy is responsible for the gene frequency changes we observe.

342 evolution of suppressors on other chromosomes, including the autosomes (Hamilton, 1967; Leigh, 1971; Hurst &
343 Pomiankowski, 1991), to restore the 50:50 sex ratio. This renders the Y-chromosomal driver cryptic, as is
344 commonly seen in e.g. *Drosophila* (Ellison & Bachtrog, 2019; Bachtrog, 2020). Loss-of-function mutations to
345 fixed autosomal suppressors of Y-chromosomal drive should re-enable the initial Y-chromosomal segregation
346 distortion in such mutants. While normally such mutations would be selected against, our model suggests that
347 they may spread if they arise when linked to a locus experiencing balancing selection from either sexually
348 antagonistic selection or sex-specific drive (see also Supplementary Figure S6). In this scenario, the loss-of-
349 function allele at the autosomal suppressor is functionally equivalent to a *trans*-acting distorter of the Y-
350 chromosome. Elsewhere in this paper we have described the *trans*-acting distorter as though it alone caused the
351 drive of what would have otherwise been well behaved, Mendelian sex chromosomes. But in the scenario just
352 outlined the *trans*-acting distorter doesn't so much cause as *permit* the sex-chromosome drive that had been lying
353 latent. This may also help explain why some suppressors of segregation distortion remain polymorphic rather than
354 become fixed, as is seen in e.g. *Drosophila* (Vaz & Carvalho, 2004; Lyth *et al.*, 2023; Gupta & Unckless, 2024).
355
356 <u>Remote-control meiotic drive in relation to sex determination systems</u>
357 Although we presented a male heterogametic system above, these results might be even more important in female
358 heterogametic (ZW) systems. A key parameter in our models, z, measures the fertility effect of sex-chromosome
359 drive. We assume that a male who experiences the loss of some X-bearing sperm is able to compensate those lost
360 sperm with functional Y-bearing sperm, at least to some extent. However, if the fertility effect of drive is
361 sufficiently steep, a *trans*-acting driver may never get off the ground. In ZW species, this sort of compensation is
362 likely automatic. Sex-chromosome drive in ZW systems takes place in an asymmetric meiosis, where some loss
363 of meiotic products (the polar bodies) is a given. Drive here is a matter of navigating to the egg pole during meiosis

I rather than sabotaging or incapacitating the half of meiotic products that inherit the alternative allele, as is the

case in male meiosis. Because drive in ZW systems may leave fertility unaffected, we might expect *trans*-acting

drivers to be more common here.

When remote-control drive occurs through X-conversion, we see that this leads to the evolution of X0

sex determination (females XX, males X0). X0 sex determination is commonly considered to evolve via the loss

of the Y-chromosome (Furman *et al.*, 2020), owing to its predisposition for accumulating mutations and reduced

rates of adaptive evolution (Bachtrog, 2008, 2013). Our model identifies a novel trajectory for this transition that

does not rely on the protracted decay of the Y-chromosome, but rather can occur at any point after the original Y-

chromosome has evolved. The genetic decay of the Y-chromosome may even make this transition selectively

favorable. The "hot potato" model of sex determination transitions posits that the accumulation of deleterious

mutations on the Y-chromosome results in a genetic load in males (Blaser *et al.*, 2014). This favors males that

lack the Y-chromosome, which are normally thought to occur through the evolution of a novel male-determining

locus on a (former) autosome. In our model, the conversion of X-bearing sperm into 0-bearing sperm effectively

reflects the conversion of a former X-chromosome into a new Y-chromosome (with regard to its male-determining

function) (Meisel, 2020). In our analysis, X0 males were assumed to suffer a fitness cost $v > 0$ relative to XY

males, but if X0 males actually experience a fitness benefit (i.e. $v < 0$) this should expand the parameter space

enabling remote-control drive.

<u>Remote-control drive has potential practical applications in e.g. pest control</u>

Fundamentally, these results challenge the current view of sex-chromosome drive as a system in which one of a

dimorphic pair of sex chromosome skews its own transmission at the expense of another. In our system, the genetic

"actors" that encode drive are autosomal, with sex chromosomes involved only for their sex-determining function.

This reveals the possibility that the field has previously overlooked cryptic sex chromosome drive systems

encoded by autosomal sex ratio distorters and challenges the view that the evolution of Y-chromosome drive is

limited by the Y-chromosome's diminutive size. Moreover, autosomally-encoded sex-chromosome drivers bypass

one major obstacle to sex-chromosome meiotic drive: meiotic sex-chromosome inactivation (Meiklejohn & Tao,

2010). This inactivation would prevent X- or Y-chromosomal meiotic drive genes from being expressed during

meiosis and is therefore often used to explain the relative dearth of killer sex chromosomes in taxa with meiotic

sex chromosome inactivation (MSCI) as opposed to taxa without it (e.g. dipterans) (Burt & Trivers, 2006).

However, if the relevant sex-chromosome–meiotic-drive genes are encoded by an autosome rather than a sex

chromosome, meiotic sex-chromosome inactivation would not prevent drive.

These observations prompt us to consider the potential application of a *trans*-drive system in the realm

of synthetic biology, as autosomally-encoded Y-drives may help overcome current technical barriers. In vector

control research, synthetic sex-ratio distortion is an attractive strategy for population suppression of a disease

vector such as *Aedes aegytpi*, and researchers have proposed X-shredders to distort population sex ratios, both via

autosomally encoded mechanisms (see Alcalay *et al.*, 2021) and Y-encoded mechanisms (see Burt & Deredec,

2018). Standard autosomal X-shredders are highly self-limiting (i.e. they only persist for a short period of time),

as the autosome encoding drive follows Mendelian inheritance. Autosomal X-shredders therefore require repeated

releases of modified mosquitoes to be an effective form of population suppression (Alcalay *et al.*, 2021). Y-linked

X-shredders are theoretically much more effective than their autosomal counterparts, given their supermendelian

inheritance, but are functionally hampered by MSCI (Lifschytz & Lindsley, 1972; Tolosana *et al.*, 2024). These
shortcomings generate desire for a Y-drive system that both avoids MSCI and is more effective than autosomal
drive.

407   Alongside existing research to circumvent MSCI on the Y-chromosome by inducing expression of the
X-shredder before it occurs (Haber *et al.*, 2024; Tolosana *et al.*, 2024), we suggest additionally considering the
potential for remote-control drive as discussed here. Specifically, inserting the synthetic X-shredding mechanism
(e.g. CRISPR[SRD], see Galizi et al. 2016) near either 1) a strongly male-beneficial autosomal allele or 2) a male-
limited autosomal driver, could provide a self-limiting synthetic drive more effective than standard autosome-
linked X-shredders and yet simpler than Y-linked X-shredders that need to subvert meiotic sex-chromosome
inactivation. While ecological modelling is outside of the scope of this paper, we encourage further exploration
of the potential application of *trans*-acting drive in synthetic systems.

<u>Conceptual implications of remote-control drive</u>
Leigh (1971) analogized the genome to a political body, where cabals of a few genes, such as meiotic drivers,
would be overwhelmed into fair segregation by a so-called parliament of genes. Specifically, he considered the
case of a sex-chromosome driver, whose distortions of the sex ratio would unite any unlinked genes into an
alliance to suppress sex-chromosome drive, thus pitting the driving sex chromosome against every other faction
of the genome. Ordinarily, this is how selfish genetic elements are viewed: as a small cabal, capable of being put
down by the parliament of genes (Scott & West, 2019). But our results call to mind a different political metaphor:
politics makes strange bedfellows. We find that unlinked loci may form a cabal, finding a shared interest in
skewing the sex ratio. The alignment between autosomal loci and sex chromosomes that we demonstrate here is
unexpected on the view that selfish genetic elements must co-locate on what Cosmides and Tooby (1981) called
a "coreplicon". Our findings show that it is not necessarily the transmission pattern that promotes collaboration.
Instead, as Gardner and Úbeda (2017) have described, a shared destination (in this case, the male germline), can
also enable collaboration among genetic parties.

429   Finally, the observation that sex chromosomes can theoretically experience *trans*-acting meiotic drive
raises definitional and conceptual questions about what meiotic "drive" means. The term "drive," when first
introduced, referred specifically to a gene's preferential segregation into an egg during female meiosis (Sandler
& Novitski, 1957). In the nearly 70 years since, the term "meiotic drive" has taken on a broader meaning, now
referring to segregation distorters as well as gamete killers or gamete converters (Burt & Trivers, 2006). Across a
broad range of meiotic drive systems, the "driver" is generally taken to mean the element that increases its own
transmission and is both the actor and direct beneficiary of drive capabilities. In *trans*-acting drive, however, when
the actor (autosome) and the direct beneficiary (Y-chromosome) of drive are physically distinct, it becomes
unclear which element is the "driver". Considering a range of options, if we take the autosome to be the driver,
then what is the Y-chromosome? The Y-chromosome could arguably be considered a hitchhiker—though
hitchhikers are generally inert rather than an integral part of the system, as the Y-chromosome is here. By contrast,
if the Y-chromosome is the driver, then how do we explain its reliance on the autosome? This leads us to our
colloquial description of *trans*-acting drive as "remote-control drive," as this recognizes that the Y-chromosome
is the object being driven, while the autosome is the enabler of this system, much like a toy car and its remote.

Conclusion

Here we have presented a model describing the evolution of a *trans*-acting segregation distorter. Previous theory has assumed that the benefits of segregation distortion must be experienced directly by the genes causing such interference, owing to the costs that come with meiotic meddling. We show that this assumption can be violated; autosomal loci that interfere with the transmission of the sex chromosomes can evolve, provided that they originate in the vicinity of a suitable locus, such as one under sexually antagonistic selection or one that exhibits sex-specific segregation distortion itself. These results show that intragenomic conflict may engender even more exotic forms of selfish genetic elements than previously considered.

Author contributions

Conceptualization: N.L.G., M.M.P., and M.A.S. Software: N.L.G., M.M.P., and M.A.S. Formal analysis: N.L.G. and M.A.S. Writing – original draft: N.L.G. and M.A.S.. Writing – review & editing: N.L.G., M.M.P., and M.A.S. Visualization: N.L.G. and M.A.S. Supervision: M.M.P. and M.A.S.

Data availability

The model source code, as well as the data processing, analysis, and visualization scripts are available through https://github.com/MartijnSchenkel/RemoteControlMeioticDrive.

Acknowledgments

We thank Arvid Ågren for fruitful discussion and helpful comments on an earlier version of the manuscript. The authors declare no conflict of interest.

References

Alcalay, Y., Fuchs, S., Galizi, R., Bernardini, F., Haghighat-Khah, R.E., Rusch, D.B., *et al.* (2021). The potential for a released autosomal X-shredder becoming a driving-Y chromosome and invasively suppressing wild populations of malaria mosquitoes. *Front. Bioeng. Biotechnol.* 9:1–11.

Bachtrog, D. (2008). The temporal dynamics of processes underlying Y chromosome degeneration. *Genetics* 179:1513–1525.

Bachtrog, D. (2020). The Y chromosome as a battleground for intragenomic conflict. *Trends Genet.* 36:510–522.

Bachtrog, D. (2013). Y-chromosome evolution: emerging insights into processes of Y-chromosome degeneration. *Nat. Rev. Genet.* 14:113–124.

Blackburn, G.S., Albert, A.Y.K. & Otto, S.P. (2010). The evolution of sex ratio adjustment in the presence of sexually antagonistic selection. *Am. Nat.* 176:264–275.

Blaser, O., Neuenschwander, S. & Perrin, N. (2014). Sex-chromosome turnovers: The hot-potato model. *Am. Nat.* 183:140–146.

Bravo Núñez, M.A., Nuckolls, N.L. & Zanders, S.E. (2018). Genetic villains: killer meiotic drivers. *Trends Genet.* 34:424–433.

Burga, A., Ben-David, E. & Kruglyak, L. (2020). Toxin-antidote elements across the tree of life. *Annu. Rev. Genet.* 54:387–415.

Burt, A. & Deredec, A. (2018). Self-limiting population genetic control with sex-linked genome editors. *Proc. R.*

Soc. B Biol. Sci. 285.

Burt, A. & Trivers, R. (2006). *Genes in Conflict: The Biology of Selfish Genetic Elements.* Harvard University
Press, Cambridge, MA, USA.

Clark, F.E. & Akera, T. (2021). Unravelling the mystery of female meiotic drive: Where we are. *Open Biol.*
11:210074.

Cosmides, L.M. & Tooby, J. (1981). Cytoplasmic inheritance and intragenomic conflict. *J. Theor. Biol.* 89:83–
129.

Ellison, C. & Bachtrog, D. (2019). Recurrent gene co-amplification on Drosophila X and Y chromosomes. *PLoS*
Genet. 15:1–23.

Eshel, I. (1985). Evolutionary genetic stability of mendelian segregation and the role of free recombination in the
chromosomal system. *Am. Nat.* 125:412–420.

Feldman, M.W. & Otto, S.P. (1991). A comparative approach to the population-genetics theory of segregation
distortion. *Am. Nat.* 173:443–456.

Fisher, R.A. (1930). *The Genetical Theory of Natural Selection*. Oxford University Press, Oxford, United
Kingdom, United Kingdom.

Furman, B.L.S., Metzger, D.C.H., Darolti, I., Wright, A.E., Sandkam, B.A., Almeida, P., *et al.* (2020). Sex
chromosome evolution: so many exceptions to the rules. *Genome Biol. Evol.* 12:750–763.

Gardner, A. & Úbeda, F. (2017). The meaning of intragenomic conflict. *Nat. Ecol. Evol.* 30:1807–1815.

Garnier, S. (2018). Default color maps from "matplotlib."

Graves, J.A.M. (2006). Sex chromosome specialization and degeneration in mammals. *Cell* 124:901–914.

Gupta, A. & Unckless, R.L. (2024). Autosomal suppression of sex-ratio meiotic drive influences the dynamics of
X and Y chromosome coevolution. *J. Hered.* 660–671.

Haber, D.A., Arien, Y., Lamdan, L.B., Alcalay, Y., Zecharia, C., Krsticevic, F., *et al.* (2024). Targeting mosquito
X-chromosomes reveals complex transmission dynamics of sex ratio distorting gene drives. *Nat. Commun.*
15:1–12.

Haig, D. & Grafen, A. (1991). Genetic scrambling as a defence against meiotic drive. *J. Theor. Biol.* 153:531–
558.

Haig, D., Úbeda, F. & Patten, M.M. (2014). Specialists and generalists: the sexual ecology of the genome. *Cold*
Spring Harb. Perspect. Biol. 6:a017525.

Hamilton, W.D. (1967). Extraordinary sex ratios. *Science (80-.).* 156:477–488.

Hartl, D.L. (1974). Genetic dissection of segregation distortion. I. Suicide combinations of *SD* genes. *Genetics*
76:477–486.

Hurst, L.D. & Pomiankowski, A. (1991). Causes of sex ratio bias may account for unisexual sterility in hybrids:
A new explanation of Haldane's rule and related phenomena. *Genetics* 128:841–858.

Innocenti, P. & Morrow, E.H. (2010). The sexually antagonistic genes of *Drosophila melanogaster. PLoS Biol.*
8:e1000335.

Jaenike, J. (2001). Sex chromosome meiotic drive. *Annu. Rev. Ecol. Syst.* 32:25–49.

Jaenike, J. (2008). X chromosome drive. *Curr. Biol.* 18:508–511.

Kozielska, M., Weissing, F.J., Beukeboom, L.W. & Pen, I. (2010). Segregation distortion and the evolution of
sex-determining mechanisms. *Heredity (Edinb).* 104:100–112.

Larracuente, A.M. & Presgraves, D.C. (2012). The selfish *Segregation Distorter* gene complex of *Drosophila*
melanogaster. *Genetics* 192:33–53.

Leigh, E.G. (1971). The Parliament of Genes. In: *Adaptation and Diversity. Natural History and the Mathematics*
of Evolution, pp. 245–254. Freeman, Cooper & Company, San Francisco, CA, USA.

Lifschytz, E. & Lindsley, D.L. (1972). The role of X-chromosome inactivation during spermatogenesis. *Proc.*
Natl. Acad. Sci. U. S. A. 69:182–186.

Lindholm, A.K., Dyer, K.A., Firman, R.C., Fishman, L., Forstmeier, W., Holman, L., *et al.* (2016). The Ecology
and Evolutionary Dynamics of Meiotic Drive. *Trends Ecol. Evol.* 31:315–326.

Lyth, S., Betancourt, A.J., Price, T.A.R. & Verspoor, R.L. (2023). The suppression of a selfish genetic element
increases a male's mating success in a fly. *Ecol. Evol.* 13:1–13.

Lyttle, T.W. (1991). Segregation distorters. *Annu. Rev. Genet.* 25:511–557.

Maffi, G. & Jayakar, S.D. (1981). A two-locus model for polymorphism for sex-linked meiotic drive modifiers
with possible applications to Aedes aegypti. *Theor. Popul. Biol.* 19:19–36.

Meiklejohn, C.D. & Tao, Y. (2010). Genetic conflict and sex chromosome evolution. *Trends Ecol. Evol.* 25:215–
223.

Meisel, R.P. (2020). Evolution of Sex Determination and Sex Chromosomes: A Novel Alternative Paradigm.
BioEssays 42:1–11.

Mendel, G. (1865). Versuche über Pflanzenhybriden. In: *Verhandlungen des naturforschenden Vereines in Brünn,*
Bd. IV für das Jahr 1865, pp. 3–47.

Muralidhar, P. & Veller, C. (2018). Sexual antagonism and the instability of environmental sex determination.
Nat. Ecol. Evol. 2:343–351.

Patten, M.M., Schenkel, M.A. & Ågren, J.A. (2023). Adaptation in the face of internal conflict: the paradox of
the organism revisited. *Biol. Rev.* 98:1796–1811.

Patten, M.M., Úbeda, F. & Haig, D. (2013). Sexual and parental antagonism shape genomic architecture. *Proc.*
R. Soc. London Ser. B-Biological Sci. 280:20131795.

Pomiankowski, A., Nöthiger, R. & Wilkins, A. (2004). The evolution of the *Drosophila* sex-determination
pathway. *Genetics* 166:1761–1773.

Price, T.A.R., Windbichler, N., Unckless, R.L., Sutter, A., Runge, J.N., Ross, P.A., *et al.* (2020). Resistance to
natural and synthetic gene drive systems. *J. Evol. Biol.* 33:1345–1360.

Prout, T., Bundgaard, J. & Bryant, S. (1973). Population genetics of modifiers of meiotic drive I. The solution of
a special case and some general implications. *Theor. Popul. Biol.* 4:446–465.

R Development Core Team. (2023). A Language and Environment for Statistical Computing. R Foundation for
Statistical Computing, Vienna, Austria.

Rice, W.R. (1992). Sexually antagonistic genes: experimental evidence. *Science (80-.).* 256:1436–1439.

RStudio Team. (2023). RStudio: Integrated Development Environment for R. RStudio, Inc., Boston MA, USA.

Rydzewski, W.T., Carioscia, S.A., Liévano, G., Lynch, V.D. & Patten, M.M. (2016). Sexual antagonism and
meiotic drive cause stable linkage disequilibrium and favour reduced recombination on the X chromosome.
J. Evol. Biol. 29:1247–1256.

Sandler, L. & Novitski, E. (1957). Meiotic drive as an evolutionary force. *Am. Nat.* 857:105–110.

Schenkel, M.A. (2024). Transitions in sex determination mechanisms through parental and sexual antagonism.

Heredity (Edinb)., doi: 10.1038/s41437-024-00717-x.

Schenkel, M.A., Pen, I., Beukeboom, L.W. & Billeter, J.-C. (2018). Making sense of intralocus and interlocus
sexual conflict. *Ecol. Evol.* 8:13035–13050.

Scott, T.W. & West, S.A. (2019). Adaptation is maintained by the parliament of genes. *Nat. Commun.* 10:5163.

Tolosana, I., Willis, K., Burt, A., Gribble, M., Nolan, T., Crisanti, A., *et al.* (2024). A Y chromosome-linked
genome editor for efficient population suppression in the malaria vector *Anopheles gambiae*. *bioRxiv*
2024.05.14.594116.

Úbeda, F., Patten, M.M. & Wild, G. (2014). On the origin of sex chromosomes from meiotic drive. *Proc. R. Soc.*
B Biol. Sci. 282:20141932.

van Doorn, G.S. (2009). Intralocus sexual conflict. In: *Year in Evolutionary Biology 2009*, pp. 52–71.

van Doorn, G.S. & Kirkpatrick, M. (2010). Transitions between male and female heterogamety caused by sex-
antagonistic selection. *Genetics* 186:629–645.

van Doorn, G.S. & Kirkpatrick, M. (2007). Turnover of sex chromosomes induced by sexual conflict. *Nature*
449:909–912.

Vaz, S.C. & Carvalho, A.B. (2004). Evolution of autosomal suppression of the *Sex-Ratio* trait in *Drosophila*.
Genetics 166:265–277.

Veller, C. (2022). Mendel's First Law: partisan interests and the parliament of genes. *Heredity (Edinb).* 129:48–
55.

Werren, J.H. (2011). Selfish genetic elements, genetic conflict, and evolutionary innovation. *Proc. Natl. Acad.*
Sci. U. S. A. 108:10863–10870.

West, S.A. (2009). *Sex Allocation*. Princeton University Press, Princeton, NJ, USA.

Wickham, H., Averick, M., Bryan, J., Chang, W., D'Agostino McGowan, L., François, R., *et al.* (2019). Welcome

Supplementary Material

The results presented in the main text sough to consider the effect of varying the strength of the *Assister* and *Distorter* loci, while keeping other parameters constant and set to optimal conditions (e.g. low fitness cost, low recombination rate). Here, we investigate the effect of varying the fitness cost, z, of the D_2 allele, and the recombination rate, r, between the *Assister* locus and the *Distorter* locus (Supplementary Figure S1 and S2). In both cases, we find that the sex ratio is most heavily skewed toward males when z and r are both minimized. The effect of recombination (r) seems to be slightly larger than the cost of drive (z).

We also present two further selective scenarios for the *Assister* locus, parental antagonism and co-segregation. We consider these scenarios to be less biologically likely than those presented in the main text, though an interesting extension of the same logic that an autosome that favors being in males will enable remote-control Y-chromosome drive.

Supplementary Figure S1. Low recombination rate (r) and low *trans*-drive cost (z) favor sex ratio distortion when *Assister* is under sexually antagonistic selection. Parameter values: $G = 1,000, h_A = 0.0$.

Supplementary Figure S2. Low recombination rate (r) and low *trans*-drive cost (z) favor sex ratio distortion when *Assister* exhibits male-limited *cis* segregation distortion. Parameter values: $G = 1,000, h_A = 0.0$.

598 <u>Parentally antagonistic selection on *Assister*</u>

599 Parental antagonism refers to the scenario in which the fitness effect of a gene depends on the parent from which

600 it was inherited. Under parentally antagonistic selection, A_1 and A_2 experience different fitness depending on the

601 parent of origin. We assume that A_1 is beneficial to fitness when inherited maternally and costly when inherited

602 paternally, and vice versa for A_2. The partial fitness score of an individual with genotype A_iA_j is given by:

603
$$w_{11} = 1 - bt$$

604
$$w_{12} = 1$$

605
$$w_{21} = 1 - t$$

606
$$w_{22} = 1 - at$$

607 Here, $t \in [0,1]$ represents the fitness effect of the wrong parent of origin, $a, b \in [0,1]$ indicates the dominance of

608 a single costly allele when inherited maternally or paternally, respectively.

Supplementary Figure S3. Dynamics and scope of Y-chromosome drive via the *trans*-acting *Distorter* causing X-shredding, linked to a parentally antagonistic *Assister* locus. (A) Changes in the allele frequency of A_2, D_2, and the X- and Y-chromosome across 2,000 generations. (B) Changes in the linkage disequilibrium of *Assister-Distorter*, *Assister*-XY, and *Distorter*-XY across 2,000 generations. (C) Haplotype frequencies over time. A_1D_2X and A_1D_2Y have near-zero frequencies. The shaded area represents the frequency of D_2 and primarily comprises the haplotypes A_2D_2Y and A_2D_2X. (D) D_2 terminal frequency after 2,000 generations. (E) Population sex ratio after 2,000 generations. Parameter values (if not otherwise defined): $a, b = 0.5, t = 0.3, k_D = 0.25, z = 0.01, r = 0.001$.

Cosegregation of *Assister*

Our final mechanism for *trans*-acting sex chromosome drive is the most direct: cosegregation of the autosome with the Y chromosome. Here, A_2 encodes a structural variant that creates a physical attachment between the autosome and the Y chromosome. We assume that an A_1A_2 (or A_2A_1) heterozygote, A_2 is co-inherited with Y at a

Supplementary Figure S4. Dynamics and scope of Y-chromosome drive via the *trans*-acting *Distorter* causing X-shredding and exhibiting co-segregation with the Y-chromosome. (A) Changes in the allele frequency of A_2D_2, and the X- and Y-chromosome across 500 generations. (B) Changes in the linkage disequilibrium of *Assister-Distorter*, *Assister*-XY, and *Distorter*-XY across 500 generations. The shaded area represents the frequency of D_2 and primarily comprises the haplotypes A_2D_2Y and A_2D_2X. (C) D_2 equilibrium frequency after 500 generations. (D) Population sex ratio after 1,000 generations. (E) Population sex ratio after 1,000 generations. Parameter values (if not otherwise defined): $c = 0.75, s = 0.01, k_D = 0.25, z = 0.01, r = 0.001$.

rate of c and A_2 with X at a rate of $1 - c$, where $c \in (0,1]$ is the cosegregation rate. We assume that A_2 negatively
affects its bearer's fitness, considering that this co-segregation mechanism may induce aneuploidy.
The partial fitness score of a male with genotype A_iA_j is given by:

$$w_{M_{11}} = 1$$
$$w_{M_{12}} = w_{M_{21}} = 1 - s$$
$$w_{M_{22}} = 1 - s$$

Here, $s \in [0,1]$ represents the fitness effect of the A_2 allele in males. The partial fitness score of a male with

genotype $D_i D_j$ is given by:

$$w_{M_{11}} = 1$$

$$w_{M_{12}} = w_{M_{21}} = 1 - z$$

$$w_{M_{22}} = 1 - z$$

Here, $z \in [0,1]$ measures the loss of total fertility, and we assume that the *Distorter* locus is neutral in females.

<u>Polymorphism of XY and X0 sex determination</u>

Considering the XY to X0 transition models presented in the main text, we sought to understand whether

polymorphism for the 0- and Y-chromosomes as the male-determining genetic factor was possible. We found a

narrow range of possibility, represented in green in Supplementary Figure S5.

Supplementary Figure S5. Range of parameter space for which Y-/0-chromosome polymorphism is possible; polymorphism is defined as both the Y- and 0-chromosome having frequencies above 1%. (A) Sexually antagonistic *Assister,* where parameter values: $s, t = 0.5; h_A = 0.5; z = 0.01; ; r = 0.001$. (B) Male-limited *cis-*segregation distorting *Assister*, where parameter values: $k_A = 0.25; h_A = 0.0; z = 0.01; r = 0.001$.

631
632

Supplementary Figures

Supplementary Figure S6: Loss of autosomal suppression of segregation distortion mimics remote control sex
chromosome meiotic drive. (A) In the ancestral state, there is no meiotic drive or suppression thereof by any
element in the genome. (B) a Y-chromosomal segregation distorter Y_D evolves and causes male-biased sex ratios.
(C) An autosomal locus S evolves to reduce Y-chromosomal segregation distortion and restore the sex ratio to
equality via the spread of the S_1 allele. (D) A second autosomal locus (A), nearby the S locus, mutates to yield a
novel allele A_1 which exhibits sex-specific segregation distortion. (E) Presence of A_1 renders loss-of-function at
the S locus advantageous to the S-A complex; evolution of the non-suppressing S_0 variant re-enables segregation
distortion by Y_D, and by extension male-biased sex ratios.

Supplementary Tables

Supplementary Table S1: Overview of model parameters.

Variable	Description	Standard value or range
k_D	strength of *trans* segregation distortion	$(0, 0.5]$ standard value: 0.25
z	fitness effect of the D_2 allele in homozygotes	$[0, 1]$ standard value = 0.01
r	Recombination between the *Assister* locus and *Distorter* locus	$[0, 0.5]$ standard value = 0.001
h_A	Dominance of A_2	$[0, 1]$ standard value = 0.5 for sexual antagonism, 0.0 for *cis* drive
t (SA only)	Fitness cost of A_2 in females	$[0, 1]$ standard value = 0.5
s	SA: Fitness cost of A_1 in males *Cis* drive: fitness cost of A_2 in both males and females	$[0, 1]$ standard value = 0.5
k_A (*cis* drive only)	Strength of *cis* drive	$(0, 0.5]$ standard value = 0.25
b	Scalar of less fit paternally inherited allele	$(0, 1]$ standard value = 0.5
a	Scalar of less fit maternally inherited allele	$(0, 1]$ standard value = 0.5
t	Cost of less fit allele from either parent	$(0, 1]$ standard value = 0.3
v	Fitness deficit of X0 males relative to XY males (X-conversion only)	$(0, 1]$ standard value = 0.01

Supplementary Table S2: Overview of fitness scores for all combinations of *Assister* and *Distorter* genotypes.

Assister type	Assister genotype	Distorter genotype	Female fitness	Male fitness
Sexual antagonism	A_1A_1	D_1D_1	1	$1 - s$
		D_1D_2, D_2D_1	1	$(1 - s)(1 - z)$
		D_2D_2	1	$(1 - s)(1 - z)$
	A_1A_2, A_2A_1	D_1D_1	$1 - h_A t$	$1 - h_A s$
		D_1D_2, D_2D_1	$(1 - h_A t)$	$(1 - h_A s)(1 - z)$
		D_2D_2	$(1 - h_A t)$	$(1 - h_A s)(1 - z)$
	A_2A_2	D_1D_1	$1 - t$	1
		D_1D_2, D_2D_1	$(1 - t)$	$1 - z$

		D_2D_2	$(1-t)$	$1-z$
Cis-segregation distortion	A_1A_1	D_1D_1	1	
		D_1D_2, D_2D_1	1	$1-z$
		D_2D_2	1	$1-z$
	A_1A_2, A_2A_1	D_1D_1	$(1-h_As)$	
		D_1D_2, D_2D_1	$(1-h_As)$	$(1-h_As)(1-z)$
		D_2D_2	$(1-h_As)$	$(1-h_As)(1-z)$
	A_2A_2	D_1D_1	$(1-s)$	
		D_1D_2, D_2D_1	$(1-s)$	$(1-s)(1-z)$
		D_2D_2	$(1-s)$	$(1-s)(1-z)$
Parental antagonism	A_1A_1	D_1D_1	$(1-bt)$	
		D_1D_2, D_2D_1	$1-bt$	$(1-bt)(1-z)$
		D_2D_2	$(1-bt)$	$(1-bt)(1-z)$
	A_1A_2	D_1D_1	1	1
		$D_1D_2,$	1	$1-z$
		D_2D_1	1	$1-z$
		D_2D_2	1	$1-z$
	A_2A_1	D_1D_1	$(1-t)$	
		$D_1D_2,$	$(1-t)$	$(1-t)(1-z)$
		D_2D_2	$(1-t)$	$(1-t)(1-z)$
	A_2A_2	D_1D_1	$(1-at)$	
		D_1D_2, D_2D_1	$(1-at)$	$(1-at)(1-z)$
		D_2D_2	$(1-at)$	$(1-at)(1-z)$