Remote-control meiotic drive of sex chromosomes

Naomi L. Greenberg[1,#] (nlg34@georgetown.edu; ORCID: 0009-0008-6880-4562)

Manus M. Patten[1] (mmp64@georgetown.edu; ORCID: 0000-0003-2890-7977)

Martijn A. Schenkel[1,2,3,*] (m.a.schenkel@rug.nl; ORCID: 0000-0002-7721-319X)

<u>Affiliations</u>

[1] Department of Biology, Georgetown University, 37th and O St. NW, Washington DC, 20057,

USA

[2] Groningen Institute of Evolutionary Life Sciences, University of Groningen, Nijenborgh 7,

9747 AG, Groningen, The Netherlands

[3] Institute of Organismic and Molecular Evolution, Johannes Gutenberg-Universität Mainz,

Hanns-Dieter-Hüsch-Weg 15, 55128, Mainz, Germany.

[#] Current address: Department of Life Sciences, Imperial College, London SW7 2AZ, UK

[*] Correspondence: Martijn A. Schenkel (maschenkel@gmail.com; m.a.schenkel@rug.nl)

Keywords: genetic conflict; meiotic drive; segregation distortion; sex chromosomes; sex

chromosome turnover; sex determination.

Abstract (174 words)

Some selfish genetic elements drive at meiosis to achieve transmission distortion, breaking the

rules of Mendelian segregation to enhance their own evolutionary success. It has been

previously shown that positive modifiers of drive must act in *cis* in order to gain the selfish

benefit of drive and that suppressors of drive will be selected at any unlinked loci. Here, we

model the evolution of an autosomal *trans*-acting gene (*Distorter*) that causes the Y-

chromosome (or even 0-chromosome) to drive. We show that such a gene may spread in the

population when linked to a second locus, *Assister*, whose alleles are transmitted at different

frequencies through sperm as compared to eggs, for which we consider various scenarios, e.g.

genes subject to sexually antagonistic selection or sex-limited drive. Depending on the

mechanistic details of sex-chromosome drive, *Distorter*'s spread can additionally facilitate

transitions between XY and X0 sex determination. We discuss these results in relation to sex

allocation theory, and as proof of principle for a novel type of naturally occurring or

synthetically useful sex-ratio-distorting selfish genetic element.

Introduction

Mendel's Law of Segregation states that in a diploid individual, the two alleles at a locus

segregate to an equal (i.e., 50/50) proportion of the gametes (Mendel, 1865). The fair

transmission of genetic material can however be distorted in myriad ways (Burt & Trivers,

2006; Patten *et al.*, 2023), such as by true meiotic drivers (Clark & Akera, 2021) and, to similar

effect, post-segregational killers (Bravo Núñez *et al.*, 2018), which often fall under the moniker

of "meiotic drive" themselves. The former achieve their advantage in asymmetric meiosis by

segregating preferentially to the egg pole during meiosis and avoiding the polar body (Sandler

& Novitski, 1957). The latter target their allelic counterpart just after meiosis, e.g. by

preventing the formation of functional gametes that lack the transmission-distorting gene

(Lindholm *et al.*, 2016; Price *et al.*, 2020). Consequently, the gametes produced by heterozygotes do not feature equal proportions of both alleles; instead the transmission-distorting allele becomes overrepresented, and the non-distorting allele underrepresented. Meiotic drivers typically harm the fitness of their bearers (Lyttle, 1991; Larracuente & Presgraves, 2012), often by causing reduced fertility. In the case of sex-chromosome drive, additional harm comes from overproducing the common sex (Lyttle, 1991). In either case, their ability to cheat the Law of Segregation offsets this cost to themselves, enabling them to spread and persist in the population.

In males, meiotic drivers may achieve their higher-than-Mendelian transmission using a variety of mechanisms, of which two are most relevant here. Both of these entail a supergene that consists of two distinct loci. First, the toxin-antidote system (Burga *et al.*, 2020) features a toxin that is expressed during or prior to meiosis, so that all resulting gametes receive a dose. In gametes that bear the toxin-antidote supergene, the antidote is expressed post-meiosis to prevent the harmful effects of the toxin and rescue the supergene-bearing gametes. The second system is the killer-responder system, which produces a killer element that targets a responder locus (Meiklejohn & Tao, 2010). In this case, the killer allele is linked to an insensitive variant of the locus that is targeted by the killer, whereas the non-killer allele is assumed to be linked to a sensitive variant and is therefore eliminated. Gametes that lack the supergene (or more specifically, the antidote allele or the insensitive responder allele) are eliminated, so that the gamete pool produced consists predominantly of gametes bearing the segregation-distorting allele at the supergene.

Drive may be found more commonly on sex chromosomes than autosomes for two reasons, one owing to genetics and the other to ascertainment bias. First, the X- and Y-chromosomes do not typically recombine for much of their length, which prevents the separation of the alleles that bring about segregation distortion when found in tandem (Hartl,

1974; Haig & Grafen, 1991). Repeated cycles of drive and suppression, which may be common

on the X- and Y-chromosomes for this reason, may lead to the evolution of many cryptic drivers

on the sex chromosomes (Ellison & Bachtrog, 2019; Bachtrog, 2020). Second, before

suppressors of sex-chromosome drive evolve, they are more readily ascertained than autosomal

segregation distorters, as skewed transmission of the sex chromosomes brings about a

conspicuous shift in the offspring sex ratio (Hamilton, 1967; Jaenike, 2001, 2008). Further, this

sex ratio effect can maintain drivers in a polymorphic state owing to the frequency dependence

of fitness. The sex chromosomes may therefore be enriched for both fixed cryptic drivers as

well as polymorphic drivers.

Sex ratio biases brough about by drive can promote evolutionary transitions between

sex determination mechanisms (Kozielska *et al.*, 2010). For example, a driving X-chromosome

may lead to female-biased sex ratios, promoting the evolution of a male-determining gene on

an autosome, converting it into a proto-Y-chromosome. Autosomal driving elements may also

promote the origin of sex chromosomes (Úbeda *et al.*, 2014). Owing to differences between

oogenesis and spermatogenesis, most meiotic drivers exhibit sex-specific (or sex-biased)

segregation distortion, so that one sex may transmit it at a Mendelian rate, whereas in the other

it distorts segregation to achieve higher transmission. If a locus linked to this sex-specific driver

evolves a novel, sex-determining mutation, this confines the driving allele to the sex in which

it achieves segregation distortion more frequently. As this supergene spreads in the population,

the segregation-distorting effect causes sex-ratio biases (similar to Kozielska *et al.* (2010)),

leading to a series of selectively favored mutations that eventually yield a set of sex-

determining chromosomes with normal, Mendelian segregation.

In the model by Úbeda *et al.*(2014), the segregation-distorting/sex-determining

supergene benefits from assigning the driving allele to the sex in which it achieves said drive.

However, we posit that achieving this effect does not require a sex-determination gene per se.

In principle, any gene that skews the sex ratio of its bearers' offspring would capture a similar benefit. Ordinarily, a sex-ratio-biasing autosomal gene would be harmful to its bearers, and hence be purged (per Fisher(1930), but see West (2009)). But a sex-ratio-biasing gene paired with a sex-specific meiotic driver—a slight modification of Úbeda *et al.*'s supergene—establishes a synergism that offsets such costs and might therefore be able to spread and persist.

Here, we model the evolution of an autosomal, *trans*-acting segregation distorter that biases the transmission of the sex chromosomes. We consider two potential proximate mechanisms by which it achieves this effect, which we dub X-shredding and X-conversion, further explained below. Linked to this *trans*-acting segregation distorter is a secondary locus, which may represent: (1) a locus under sexually antagonistic selection, where different alleles are favored in females as compared to males (van Doorn, 2009; Schenkel *et al.*, 2018) or (2) a sex-specific meiotic driver, an analog to the situation modeled by Úbeda *et al.* (2014). In the Supplement we consider two other scenarios, one in which the *Assister* locus is parentally antagonistic, wherein an allele's fitness depends on its parental origin (Patten *et al.*, 2013; Haig *et al.*, 2014), and a second in which the *Assister* locus simply causes co-segregation of the autosome and the Y-chromosome. The underlying rationale for these scenarios is that in all cases, the *trans*-acting segregation-distorting gene and the linked gene have the same optimal destination—namely, a male offspring (Gardner & Úbeda, 2017).

In addition, these phenomena have previously been shown capable of spurring transitions between sex-determining mechanisms (van Doorn & Kirkpatrick, 2007, 2010; Kozielska *et al.*, 2010; Úbeda *et al.*, 2014; Schenkel, 2024). We use our model to determine the conditions under which such a gene can spread in the population, what frequency it reaches at equilibrium, and what level of sex-ratio distortion is achieved once equilibrium is attained. We also consider whether it can spur a transition between genetic sex-determination mechanisms. Through this, we show that autosomal genes may interfere with the segregation

of the sex chromosomes to promote their own spread in the population. This results in the sex

chromosomes appearing to drive through no effort of their own—i.e., through remote control.

We discuss our results in light of sex allocation theory, definitional issues of gene drives, and

in the context of potential applications.

Methods

Our model assumes an infinite population with XY sex determination (females XX, males

XY), with an initially equal sex ratio. The X and Y chromosomes determine sex and do not

carry any other genes relevant to our model. Terms such as 'female' and 'male', and similarly

'XY' versus 'ZW' can be exchanged to consider cases with female heterogamety, rather than

male heterogamety. In addition to the XY sex chromosomes, individuals carry an autosome

carrying two loci, dubbed *Distorter* (with alleles D_1 and D_2) and *Assister* (with alleles A_1 and

A_2); recombination between *Distorter* and *Assister* occurs at a rate r in both sexes (Figure 1A).

The D_1 allele makes for equal segregation of the sex chromosomes, whereas the D_2 allele

interferes with the normal (i.e., Mendelian) segregation of the XY sex chromosomes in males.

Further details on D_2's mechanism are provided under "*Trans*-segregation distortion by

Distorter" below.

For the *Assister* locus, we focus in the main text on two selective scenarios, sexually

antagonistic selection and male-limited *cis*-segregation distortion, and examine the cases of

parentally antagonistic selection and of co-segregation in the Supplementary Material. More

details of the two schemes we examine in the main text are provided in "Sexually antagonistic

selection on *Assister*" and "*Cis*-segregation distortion by *Assister*", respectively.

The crux of each of these selective scenarios is that the A_2 allele experiences higher

long-term fitness by segregating to males, either because: (1) this is the sex in which it exhibits

higher-than-Mendelian transmission (segregation distortion); or (2) it enhances the sex of its

bearer (sexual antagonism); or (3) it comes to be transmitted in a manner that enhances its

fitness in the subsequent generation (parental antagonism). This favors the buildup of linkage

disequilibrium between *Assister* and *Distorter*, as well as statistical linkage between this pair

of loci and the Y-chromosome. Consequently, the *trans*-segregation distorting allele D_2 can

spread via its association with the A_2 allele and its overrepresentation in males.

The overall life cycle is as follows: mature males and females undergo meiosis to

produce their respective gamete pools; during this time, segregation distortion takes place;

gametes are then united at random; the resulting juvenile offspring undergo viability selection

during maturation; and upon reaching maturity they replace their parents as the reproductive

population so that generations are non-overlapping. At this point, the cycle resets. Our

population is initially fixed for the D_1 allele, and with A_1 and A_2 at or near their equilibrium as

achieved by 500 "burn-in" generations of simulation. We then mutate a small ($p=0.001$)

proportion of D_1 alleles to D_2 alleles on an A_2Y background, turning them into A_2D_2Y males.

We then continue the simulation procedure for 1,000 generations before we evaluate the

terminal frequency of D_2.

In our simulations, we vary the parameter values for the selective effects of the *Assister*

and *Distorter* loci, as well as their capacity to distort allelic segregation during meiosis (where

applicable); details on these are provided under "Results". We focus here in the main text

primarily on the impact of the strength of selection on *Assister* (for sexually antagonistic

selection) or its capacity to achieve higher-than-Mendelian transmission (for *cis*-segregation

distortion), as well as the strength of *trans*-segregation distortion achieved by *Distorter*. But

we also consider the role of recombination between the *Assister* and the *Distorter* loci and

direct selection against *Distorter* more fully in the Supplement.

An overview of all model variables, their descriptions, and standardized values is

included in Supplementary Table S1.

Figure 1: Model overview. (A) Our model incorporates two linked autosomal loci, *Assister* (with alleles A_1 and A_2) and *Distorter* (with alleles D_1 and D_2); recombination between *Assister* and *Distorter* occurs at a rate r in both sexes. *Assister* may represent either a sexually antagonistic locus (A_1 beneficial in females/detrimental in males, A_2 detrimental in females/beneficial in males) or a locus experiencing *cis*-segregation distortion (D_2 overtransmitted in D_1/D_2 heterozygotes). The D_2 allele interferes with the Mendelian transmission of the X-, Y-, and/or O-chromosomes during spermatogenesis to achieve *trans*-segregation distortion.

Sexually antagonistic selection on *Assister*

Under sexually antagonistic selection, the A_1 and A_2 alleles differently affect the fitness of female versus male bearers. We assume that A_1 is beneficial to female fitness but costly to male fitness, and vice versa for A_2. The partial fitness score $w_{F_{ij}}$ of a female with a genotype $A_i A_j$ is given by:

$$w_{F_{11}} = 1$$

$$w_{F_{12}} = w_{F_{21}} = 1 - h_A t$$

$$w_{F_{22}} = 1 - t$$

The partial fitness score $w_{M_{ij}}$ of a male with a $A_i A_j$ genotype is given by:

$$w_{M_{11}} = 1 - s$$

$$w_{M_{12}} = w_{M_{21}} = 1 - (1 - h_A)s$$

$$w_{M_{22}} = 1$$

Here, t and s indicate the homozygous fitness cost of A_2 in females and A_1 in males, respectively, and $h_A \in [0,1]$ is the dominance of A_2 in heterozygotes.

Cis-segregation distortion by *Assister*

As a segregation-distorting gene, the A_2 allele of the *Assister* locus is transmitted at a higher-than-Mendelian rate in heterozygotes. Specifically, we assume that an A_2A_1 (or A_1A_2) heterozygote transmits the A_2 allele at a rate of k_A and the A_1 allele at a rate $(1- k_A)$ where $k_A \in$ (0,1] indicates the strength of *cis*-segregation distortion by A_2 and $k_A = \frac{1}{2}$ represents Mendelian segregation. We assume that the A_2 allele negatively affects its bearer's fitness. The partial fitness score w_{ij} of an individual with genotype A_iA_j is given by:

$$w_{11} = 1$$

$$w_{12} = w_{21} = 1 - h_As$$

$$w_{22} = 1 - s$$

Here, $s \in [0,1]$ represents the fitness effect of the A_2 allele in homozygotes, $h_A \in [0,1]$ its dominance in heterozygotes. We chose this fitness scheme for *Assister* to reflect some well-known autosomal drivers (e.g., the *t* haplotype in mouse and *SD* in *Drosophila*, which have linked recessive lethality/sterility mutations) and also because it maintains polymorphism at the *Assister* locus, which is essential for our model.

Trans-segregation distortion by *Distorter*

The D_2 allele of the *Distorter* locus acts to distort the segregation rate of the sex chromosomes; we assume that D_2 acts only in males, i.e. the heterogametic sex, as a dominant *trans*-acting

segregation distorter, such that D_2D_1 heterozygotes and D_2D_2 homozygotes transmit their sex chromosomes at equally distorted frequencies. We consider two mechanisms of *trans*-segregation distortion, denoted X-shredding and X-conversion, respectively. X-shredding represents a skew in the transmission of X- versus Y-chromosomes to future offspring, where the Y-chromosome is transmitted at a rate $\frac{1}{2} + k_D$ and the X chromosome at a rate $\frac{1}{2} - k_D$, where $k_D \in (0, 1/2]$; this is achieved by the killing off of some fraction of X-bearing sperm. Under X-conversion, D_2 acts to eliminate the X-chromosome from a certain number of X-bearing sperm to convert them into 0-bearing gametes that, when fused with an X-bearing egg yield a viable, fertile X0 male. Consequently, the Y-chromosome is transmitted at a Mendelian rate of $\frac{1}{2}$, the X-chromosome is transmitted at a rate $\frac{1}{2} - k_D$, and the remaining k_D represents the proportion of sperm that have been converted from being X-bearing to 0-bearing. In X0 males, the 0-chromosome would be transmitted at a rate $\frac{1}{2} + k_D$ and the X-chromosome at a rate $\frac{1}{2} - k_D$.

We assume that the D_2 allele is harmful to male fertility through its sperm-killing or X conversion efforts. The resulting partial fitness score w_{ij} of a male with a D_iD_j genotype is given by:

$$w_{11} = 1$$

$$w_{12} = w_{21} = 1 - z$$

$$w_{22} = 1 - z$$

Here, $z \in [0,1]$ measures the loss of total fertility, and we assume that the *Distorter* locus is neutral in females. Finally, we assume that the partial fitness effects of the *Assister* and *Distorter* loci are multiplicative; Supplementary Table S2 provides an overview of the fitnesses of all potential genotypes in our population.

 <u>Data analysis and visualization</u>

Simulations, data analysis, and visualization were carried out using R (R Development Core

Team (2023), v. 4.4.1) and RStudio (RStudio Team (2023), v. 2024.09.0+375) using the

'tidyverse' and 'viridis' packages (Garnier, 2018; Wickham *et al.*, 2019).

Results

<u>Remote control drive with sexually antagonistic selection</u>

We first explored whether an autosomal *trans*-acting segregation distorter (i.e., the D_2 allele of

the *Distorter* locus) of the Y-chromosome could spread in the population when linked to a

sexually antagonistic locus (the *Assister* locus). With strong sexually antagonistic selection

($s, t = 0.5$) as well as a high level of *trans*-acting segregation distortion ($k_D = 0.75$), we

found that D_2 can invade and enhance Y-chromosome transmission (Figure 2). Under these

conditions, and with minimal fertility costs of D_2 ($z = 0.001$) and tight linkage between

Assister and *Distorter* ($r = 0.001$), we found that D_2 could reach an equilibrium frequency of

0.138, resulting in a population sex ratio (proportion of males) of 56.5%. Meanwhile, the

Assister locus remains polymorphic, with A_2's frequency changing only slightly from 0.58 at

its initial equilibrium to a frequency of 0.55 after 500 generations (Figure 2A).

We also tracked the linkage between A_2, D_2, and the Y chromosome (Figure 2B). These three

loci (A_2, D_2, and Y) come into close statistical linkage and are inherited non-independently—

effectively, as a group—allowing D_2 to benefit from the male-beneficial fitness effects of A_2,

Y to benefit from its increased transmission thanks to D_2, and A_2 to benefit from being assigned

to its fitness-enhancing state (i.e. in males) thanks to a stronger association with the Y

chromosome. We measure linkage disequilibrium to track these non-random associations,

using the following equations:

$$D_{AD} = (A_1 D_1 X + A_2 D_2 X + A_1 D_1 Y + A_2 D_2 Y) - (A_2 D_1 X + A_1 D_2 X + A_2 D_1 Y + A_1 D_2 Y)$$

$$D_{AY} = (A_2D_1Y + A_2D_2Y + A_1D_1X + A_2D_1X) - (A_1D_1Y + A_2D_1X + A_2D_1Y + A_2D_2X)$$

$$D_{DY} = (A_2D_1Y + A_2D_2Y + A_1D_1X + A_2D_1X) - (A_1D_2X + A_2D_2X + A_1D_1Y + A_2D_1Y)$$

To better characterize the conditions that favor the spread of D_2, we fully explored the

parameter space of sexual antagonism (s and t) at the *Assister* locus and the strength of drive

(k) of the D_2 allele. We found that D_2 can spread under most parameter value combinations

considered here (Figure 2D), though it achieves the highest equilibrium frequency when *trans*

drive is relatively weak ($k \approx 0.6$) and sexually antagonistic selection is strong ($s, t \cong 1$). The

presence of D_2 in the population leads to biased sex ratios, which are highest for high levels of

trans drive and strong sexually antagonistic selection (Figure 2E). When sexually antagonistic

selection is relatively weak, we see that higher *trans*-segregation distortion does not lead to

concomitantly higher frequencies of D_2. Under these conditions, the cost to D_2 of assigning

itself to the common sex outstrip the benefits linkage to A_2.

We also investigated the constraints that higher recombination rates and higher D_2

fertility costs place on the spread of D_2. We found that higher recombination ($r = 0.01$ or $r =$

0.001) reduced the opportunity for male-biased sex ratios, likely by reducing the necessary

A_2-D_2-Y linkage disequilibrium; similarly, a higher D_2 fertility cost reduced the success of the

distorter (Supplementary Figure S1).

Figure 2. Dynamics and scope of Y-chromosome drive via the *trans*-acting *Distorter*. Here,

Distorter is linked to a sexually antagonistic *Assister* locus and causes drive via the X-

shredding mechanism. (A) Changes in the frequency of the A_2 and D_2 alleles and the X and Y

chromosomes across 500 generations. (B) Changes in the linkage disequilibrium of *Assister-*

Distorter, *Assister-XY*, and *Distorter-XY* across 500 generations. (C) Haplotype frequencies

over 500 generations. A_1D_2X and A_1D_2Y are near-zero, as D_2 is introduced on an A_2 background

and recombination is low. The shaded area represents the frequency of D_2 and primarily

comprises the haplotypes A_2D_2Y and A_2D_2X. (D) D_2 equilibrium frequency after 1,000

generations. (E) Population sex ratio after 1,000 generations. Parameter values (if not otherwise

specified): $k_D = 0.25, s, t = 0.5, h_A = 0.5, z = 0.01, r = 0.001$.

<u>Remote control drive by male-limited *cis*-segregation distortion</u>

We then explored whether a male-limited, *cis*-acting segregation distorter at the *Assister* locus

could similarly favor the spread of D_2. We found that for a large range of values of *cis* drive

strength (k_A) and *trans* drive strength (k_D), D_2 spreads and the population skews male. Similar

to the sexual antagonism model above, the Y chromosome as well as D_2 increase in frequency,

whereas A_2's frequency remains relatively stable (Figure 3A). The linkage disequilibrium

between *Assister* and *Distorter* is strong, as associations between the Y chromosome and both

D_2 and A_2 develop (Figure 3B). D_2 occurs primarily in combination with A_2 (Figure 3C). We

investigated the full range of drive strengths ($k_A \in [0.5, 1], k_D \in [0.5, 1]$) at both the *Distorter*

locus and the *Assister* locus (Figures 3D, 3E). We found that most of the parameter space

permits invasion of D_2, and some parameter combinations enable high levels of D_2 and high

sex-ratio skew. With freer recombination between *Assister* and *Distorter* and with elevated

fitness costs of D_2, the opportunity for D_2's spread diminishes, and the resulting sex-ratio skew

is less extreme (Supplementary Figure 2).

Figure 3. Dynamics and scope of Y-chromosome drive via the *trans*-acting *Distorter* causing X-shredding, linked to a male-limited meiotic driver *Assister*.. (A) Changes in the allele frequency of A_2, D_2, and the X- and Y-chromosome across 500 generations. (B) Changes in the linkage disequilibrium of *Assister-Distorter*, *Assister-XY*, and *Distorter-XY* across 500 generations. (C) Haplotype frequencies over 500 generations. A_1D_2X and A_1D_2Y are near-zero, as D_2 is introduced on an A_2 background and recombination is low. The shaded area represents the frequency of D_2 and primarily comprises the haplotypes A_2D_2Y and A_2D_2X. (D) D_2 equilibrium frequency after 1,000 generations. (E) Population sex ratio after 1,000 generations.

314 | Parameter values (if not otherwise specified): $k_D = 0.25, k_A = 0.25, h_A = 0.0, z =$
315 | $0.01, r = 0.001$.

X-conversion and the emergence of X0 sex determination

In the previous sections, we assumed that D_2 distorts the segregation of the X- and Y-

chromosomes by the selective killing of X-bearing sperm. An alternative mechanism by which

D_2 might generate a male-biased sex ratio among its bearer's offspring is to selectively

eliminate the X-chromosome from sperm to convert them to 0-bearing sperm, yielding X0 sons

instead of XY sons. This mechanism might not harm male fertility as severely as the X-

shredding mechanism, and so is poised to spread more easily. This form of *trans*-segregation

distortion also provides a mechanism for an XY sex determination system to transition to an

X0 sex determination system. We find that when the D_2 allele causes X-conversion, the

resulting 0-chromosomes accumulate and become the predominant male-determining factor.

This holds true when *Assister* experiences sexual antagonism (Figure 4), as well as when it

exhibits male-limited *cis* drive (Figure 5). In both cases, we find that the 0-chromosome can

entirely replace the Y-chromosome. There is only a small range of parameter space in which

the Y-chromosome escapes loss and is maintained alongside the 0 chromosome in the

population (Supplementary Figure S5).

Figure 4. Dynamics and scope of 0-chromosome drive via the *trans*-acting *Distorter* causing X-conversion, linked to a sexually-antagonistic *Assister*. (A) Changes in the allele frequency of A_2, D_2, and the X- and Y-chromosome across 500 generations. (B) Changes in the linkage disequilibrium of *Assister-Distorter*, *Assister-XY*, and *Distorter-XY* across 500 generations. (C) Haplotype frequencies over 500 generations. A_1D_2X and A_1D_20 are near-zero, as D_2 is introduced on an A_2 background and recombination is low. The shaded area represents the frequency of D_2 and primarily comprises the haplotypes A_2D_2Y and A_2D_2X. (D) D_2 equilibrium frequency after 1,000 generations. (E) Population sex ratio after 1,000 generations. Parameter

values (if not otherwise defined): $k_D = 0.25, s, t = 0.5, h_A = 0.0, z = 0.01, r = 0.001,$ $w = 0.01$.

Figure 5: Dynamics and scope of 0-chromosome drive via the *trans*-acting *Distorter* causing X-conversion, linked to a male-limited meiotic driver *Assister*. (A) Changes in the allele frequency of A_2, D_2, and the X-, Y-, and 0-chromosome across 500 generations. (B) Changes in the linkage disequilibrium of *Assister-Distorter*, *Assister-XY*, and *Distorter-XY* across 500

generations. (C) Haplotype frequencies over 500 generations. A_1D_2X and A_1D_2Y are near-zero,

as D_2 is introduced on an A_2 background and recombination is low. The shaded area represents

the frequency of D_2 and primarily comprises the haplotypes A_2D_20 and A_2D_2X. (D) D_2

equilibrium frequency after 1,000 generations. (E) Population sex ratio after 1,000 generations.

Parameter values (if not otherwise defined): $k_D = 0.25, k_A = 0.25, h_A = 0.5, z =$

$0.01, r = 0.001, w = 0.01.$

Discussion

Here, we have shown that some *trans*-acting segregation distorters, which target an unlinked

locus on the X-chromosome, can spread when they arise on a suitable genetic background.

Thus, sex-chromosomal segregation distortion need not be caused solely by sex-linked loci.

Additionally, our results contradict the conclusion from previous theory, which finds that

unlinked loci should ordinarily suppress drive (Eshel, 1985; Veller, 2022). We also found that

such *trans*-acting distorters could spur a transition in sex chromosome systems from XY to X0

male heterogamety.

We focused on two particular scenarios, where a *trans*-acting segregation distorter

(*Distorter*) arises near a locus (*Assister*) that is either under sexually antagonistic selection or

exhibits sex-specific *cis*-segregation distortion (in the supplement we consider a third scenario,

where the nearby locus experiences parentally-antagonistic selection, and a fourth scenario,

where the autosomal locus co-segregates with the Y-chromosome). Both genetic backgrounds

lead to linkage disequilibrium between *Distorter* and *Assister*, where the *trans*-acting

segregation-distorting allele D_2 becomes associated with A_2, the male-beneficial allele or the

cis-segregation distorting allele in the two respective scenarios. Under either scenario, A_2 has

the same optimal destination: a male individual (Figure 6A). If *Assister* is sexually antagonistic,

A_2 will act to enhance its bearer's fitness—and hence its own fitness—when it ends up in a

male as opposed to a female. If *Assister* is parentally antagonistic (see Supplementary Material), A_2 will come to be paternally transmitted, and hence will enhance the fitness of its future bearers; this is effectively the same effect as for sexual antagonism, but with the fitness effects delayed by one generation. Lastly, when *Assister* is a sex-specific segregation distorter, A_2 will exhibit a higher-than-Mendelian transmission rate when its bearer is a male, as opposed to a fair transmission rate of 50% in females. In all scenarios, D_2 acts to increase its own effective transmission to males by skewing the sex ratio of its bearer's offspring. The cost to D_2 of assigning itself to the more common sex is offset by the linked benefit of delivering an A_2 allele to its favored destination (Figure 6B).

Figure 6: (A) A_2 experiences fitness benefits by preferentially segregating to males under each evolutionary scenario for *Assister*. Sexual antagonism (SA): A_2 confers a fitness benefit so that male bearers have higher-than-average fitness (w), whereas female bearers have lower-than-average fitness. Male-limited *cis*-segregation distortion (SD): A_2 has a higher-than-Mendelian transmission (p) in males, but regular Mendelian transmission in females. Parental antagonism (PA): A_2 confers a fitness benefit to its bearers when paternally inherited so that they have higher-than-average fitness, whereas it confers a fitness cost when maternally-inherited. (B) A_2 and D_2 each benefit from the other. A_2 provides D_2 a fitness or transmission benefit, and D_2 provides A_2 a "destination benefit," i.e. indirectly delivers it to a destination where it fares better (i.e., in a male). This synergy is responsible for the gene frequency changes we observe.

Mechanistically, a *trans*-acting sex-chromosome driver may seem rather unlikely. But,

on the contrary, it is actually quite easy to obtain. Consider a traditional Y-linked driver that

results in biased sex ratios but reaches fixation despite assigning itself to the more common

sex. Such sex-ratio biases cause selection to favor the evolution of suppressors on other

chromosomes, including the autosomes (Hamilton, 1967; Leigh, 1971; Hurst & Pomiankowski,

1991), to restore the 50:50 sex ratio. This renders the Y driver cryptic, as is commonly seen in

e.g. *Drosophila* (Ellison & Bachtrog, 2019; Bachtrog, 2020). Loss-of-function mutations to

fixed autosomal suppressors of Y drive should re-enable the initial Y-chromosomal segregation

distortion in such mutants. While normally such mutations would be selected against, our

model suggests that they may spread if they arise when linked to a locus experiencing balancing

selection from either sexually antagonistic selection or sex-specific drive (see also

Supplementary Figure S6). In this scenario, the loss-of-function allele at the autosomal

suppressor functions as a *trans*-acting distorter of the Y chromosome. This may also help

explain why some suppressors of segregation distortion remain polymorphic rather than

become fixed, as is seen in e.g. *Drosophila* (Vaz & Carvalho, 2004; Lyth *et al.*, 2023; Gupta

& Unckless, 2024).

<u>Diverse Sex Determination Systems</u>

Although we presented a male heterogametic system above, we think these results might be

even more important in female heterogametic (ZW) systems. A key parameter in our models,

z, measures the fertility effect of sex-chromosome drive. We assume that a male who

experiences the loss of some X-bearing sperm is able to compensate those lost sperm with

functional Y-bearing sperm, at least to some extent. However, if the fertility effect of drive is

sufficiently steep, a *trans*-acting driver may never get off the ground. In ZW species, this sort

of compensation is likely automatic. Sex-chromosome drive in ZW systems takes place in an

asymmetric meiosis, where some loss of meiotic products (the polar bodies) is a given. Drive

here is a matter of navigating to the egg pole during meiosis I rather than sabotaging or

incapacitating the half of meiotic products that inherit the alternative allele, as is the case in

male meiosis. Because drive in ZW systems may leave fertility unaffected, we might expect

trans-acting drivers to be more common here.

<u>Practical applications of remote-control drive</u>

Fundamentally, these results challenge the current view of sex-chromosome drive as a system

in which one of a dimorphic pair of sex chromosome skews its own transmission at the expense

of another. In our system, the genetic "actors" that encode drive are autosomal, with sex

chromosomes involved only for their sex-determining function. This reveals the possibility that

the field has previously overlooked cryptic sex chromosome drive systems encoded by

autosomal sex ratio distorters and challenges the view that the evolution of Y-chromosome

drive is limited by the Y-chromosome's diminutive size. Moreover, autosomally encoded sex-

chromosome drivers bypass one major obstacle to sex-chromosome meiotic drive: meiotic sex-

chromosome inactivation (Meiklejohn & Tao, 2010). This inactivation would prevent X- or Y-

chromosomal meiotic drive genes from being expressed during meiosis and is therefore often

used to explain the relative dearth of killer sex chromosomes in taxa with meiotic sex

chromosome inactivation (MSCI) as opposed to taxa without it (e.g. dipterans) (Burt & Trivers,

2006). However, if the relevant sex-chromosome–meiotic-drive genes are encoded by an

autosome rather than a sex chromosome, meiotic sex-chromosome inactivation would not

prevent drive.

These observations prompt us to consider the potential application of a *trans*-drive

system in the realm of synthetic biology, as autosomally encoded Y-drives may help overcome

current technical barriers. In vector control research, synthetic sex-ratio distortion is an

attractive strategy for population suppression of a disease vector such as *Aedes aegytpi*, and

researchers have proposed X-shredders to distort population sex ratios, both via autosomally

encoded mechanisms (see (Alcalay *et al.*, 2021)) and Y-encoded mechanisms (see (Burt &

Deredec, 2018)). Standard autosomal X-shredders are highly self-limiting (i.e. they only persist

for a short period of time), as the autosome encoding drive follows Mendelian inheritance.

Autosomal X-shredders therefore require repeated releases of modified mosquitoes to be an

effective form of population suppression (Alcalay *et al.*, 2021). Y-linked X-shredders are

theoretically much more effective than their autosomal counterparts, given their

supermendelian inheritance, but are functionally hampered by MSCI (Tolosana *et al.*, 2024).

These shortcomings generate desire for a Y-drive system that both avoids MSCI and is more

effective than autosomal drive.

Alongside existing research to circumvent MSCI on the Y-chromosome by inducing

expression of the X-shredder before it occurs (see (Haber *et al.*, 2024; Tolosana *et al.*, 2024),

we suggest additionally considering the potential for remote-control drive as discussed here.

We propose that inserting the synthetic X-shredding mechanism (e.g. CRISPR[SRD], see Galizi

et al. 2016) near either 1) a strongly male-beneficial autosomal allele or 2) a male-limited

autosomal driver, could provide a self-limiting synthetic drive more effective than standard

autosome-linked X-shredders and yet simpler than Y-linked X-shredders that need to subvert

meiotic sex-chromosome inactivation. While ecological modelling is outside of the scope of

this paper, we encourage further exploration of the potential application of *trans*-acting drive

in synthetic systems.

<u>Conceptual implications of remote-control drive</u>

Leigh (1971) analogized the genome to a political body, where cabals of a few genes, such as

meiotic drivers, would be overwhelmed into fair segregation by a so-called parliament of genes.

Specifically, he considered the case of a sex-chromosome driver, whose distortions of the sex

ratio would unite any unlinked genes into an alliance to suppress sex-chromosome drive, thus

pitting the driving sex chromosome against every other faction of the genome. Ordinarily, this

is how selfish genetic elements are viewed: as a small cabal, capable of being put down by the

parliament of genes (Scott & West, 2019). But our results call to mind a different political

metaphor: politics makes strange bedfellows. We find that unlinked loci may form a cabal,

finding a shared interest in skewing the sex ratio. The alignment between autosomal loci and

sex chromosomes that we demonstrate here is unexpected on the view that selfish genetic

elements must co-locate on what Cosmides and Tooby (1981) called a "coreplicon". Our

findings show that it is not necessarily the transmission pattern that promotes collaboration.

Instead, as Gardner and Úbeda (2017) have described, a shared destination (in this case, the

male germline), can also enable collaboration among genetic parties.

Finally, the observation that sex chromosomes can theoretically experience *trans*-acting

meiotic drive raises definitional and conceptual questions about what meiotic "drive" means.

The term "drive," when first introduced, referred specifically to a gene's preferential

segregation into an egg during female meiosis (Sandler & Novitski, 1957). In the nearly 70

years since, the term "meiotic drive" has taken on a broader meaning, now referring to

segregation distorters as well as gamete killers or gamete converters (Burt & Trivers, 2006).

Across a broad range of meiotic drive systems, the "driver" is generally taken to mean the

element that increases its own transmission and is both the actor and direct beneficiary of drive

capabilities. In *trans*-acting drive, however, when the actor (autosome) and the direct

beneficiary (Y-chromosome) of drive are physically distinct, it becomes unclear which element

is the "driver". Considering a range of options, if we take the autosome to be the driver, then

what is the Y-chromosome? The Y-chromosome could arguably be considered a hitchhiker—

though hitchhikers are generally inert rather than an integral part of the system, as the Y

chromosome is here. By contrast, if the Y chromosome is the driver, then how do we explain

its reliance on the autosome? This leads us to our colloquial description of *trans*-acting drive

as "remote-control drive," as this recognizes that the Y is the object being driven, while the

autosome is the enabler of this system, much like a toy car and its remote.

Conclusion

Here we have presented a model describing the evolution of a *trans*-acting segregation

distorter. Previous theory has assumed that the benefits of segregation distortion must be

experienced directly by the genes causing such interference, owing to the costs that come with

meiotic meddling. We show that this assumption can be violated; autosomal loci that interfere

with the transmission of the sex chromosomes can evolve, provided that they originate in the

vicinity of a suitable locus, such as one under sexually antagonistic selection or one that

exhibits sex-specific segregation distortion itself. These results show that intragenomic conflict

may engender even more exotic forms of selfish genetic elements than previously considered.

Author contributions

Conceptualization: N.L.G., M.M.P., and M.A.S. Software: N.L.G., M.M.P., and M.A.S.

Formal analysis: N.L.G. and M.A.S. Writing – original draft: N.L.G. and M.A.S.. Writing –

review & editing: N.L.G., M.M.P., and M.A.S. Visualization: N.L.G. and M.A.S. Supervision:

M.M.P. and M.A.S.

Data availability

The model source code, as well as the data processing, analysis, and visualization scripts will

be made freely available upon acceptance

Acknowledgments

We thank Arvid Ågren for fruitful discussion and helpful comments on an earlier version of

the manuscript. The authors declare no conflict of interest.

References

Alcalay, Y., Fuchs, S., Galizi, R., Bernardini, F., Haghighat-Khah, R.E., Rusch, D.B., *et al.*

(2021). The potential for a released autosomal X-shredder becoming a driving-Y

chromosome and invasively suppressing wild populations of malaria mosquitoes. *Front.*

Bioeng. Biotechnol. 9:1–11.

Bachtrog, D. (2020). The Y chromosome as a battleground for intragenomic conflict. *Trends*

Genet. 36:510–522.

Bravo Núñez, M.A., Nuckolls, N.L. & Zanders, S.E. (2018). Genetic villains: killer meiotic

drivers. *Trends Genet.* 34:424–433.

Burga, A., Ben-David, E. & Kruglyak, L. (2020). Toxin-antidote elements across the tree of

life. *Annu. Rev. Genet.* 54:387–415.

Burt, A. & Deredec, A. (2018). Self-limiting population genetic control with sex-linked

genome editors. *Proc. R. Soc. B Biol. Sci.* 285.

Burt, A. & Trivers, R. (2006). *Genes in Conflict: The Biology of Selfish Genetic Elements.*

Harvard University Press, Cambridge, MA, USA.

Clark, F.E. & Akera, T. (2021). Unravelling the mystery of female meiotic drive: Where we

are. *Open Biol.* 11:210074.

Cosmides, L.M. & Tooby, J. (1981). Cytoplasmic inheritance and intragenomic conflict. *J.*

Theor. Biol. 89:83–129.

Ellison, C. & Bachtrog, D. (2019). Recurrent gene co-amplification on Drosophila X and Y

chromosomes. *PLoS Genet.* 15:1–23.

Eshel, I. (1985). Evolutionary genetic stability of mendelian segregation and the role of free

recombination in the chromosomal system. *Am. Nat.* 125:412–420.

Fisher, R.A. (1930). *The Genetical Theory of Natural Selection*. Oxford University Press,

Oxford, United Kingdom, United Kingdom.

Gardner, A. & Úbeda, F. (2017). The meaning of intragenomic conflict. *Nat. Ecol. Evol.*

30:1807–1815.

Garnier, S. (2018). Default color maps from "matplotlib."

Gupta, A. & Unckless, R.L. (2024). Autosomal suppression of sex-ratio meiotic drive

influences the dynamics of X and Y chromosome coevolution. *J. Hered.* 660–671.

Haber, D.A., Arien, Y., Lamdan, L.B., Alcalay, Y., Zecharia, C., Krsticevic, F., *et al.* (2024).

Targeting mosquito X-chromosomes reveals complex transmission dynamics of sex

ratio distorting gene drives. *Nat. Commun.* 15:1–12.

Haig, D. & Grafen, A. (1991). Genetic scrambling as a defence against meiotic drive. *J.*

Theor. Biol. 153:531–558.

Haig, D., Úbeda, F. & Patten, M.M. (2014). Specialists and generalists: the sexual ecology of

the genome. *Cold Spring Harb. Perspect. Biol.* 6:a017525.

Hamilton, W.D. (1967). Extraordinary sex ratios. *Science (80-.).* 156:477–488.

Hartl, D.L. (1974). Genetic dissection of segregation distortion. I. Suicide combinations of

SD genes. *Genetics* 76:477–486.

Hurst, L.D. & Pomiankowski, A. (1991). Causes of sex ratio bias may account for unisexual

sterility in hybrids: A new explanation of Haldane's rule and related phenomena.

Genetics 128:841–858.

Jaenike, J. (2001). Sex chromosome meiotic drive. *Annu. Rev. Ecol. Syst.* 32:25–49.

Jaenike, J. (2008). X chromosome drive. *Curr. Biol.* 18:508–511.

Kozielska, M., Weissing, F.J., Beukeboom, L.W. & Pen, I. (2010). Segregation distortion and

the evolution of sex-determining mechanisms. *Heredity (Edinb).* 104:100–112.

Larracuente, A.M. & Presgraves, D.C. (2012). The selfish *Segregation Distorter* gene

complex of *Drosophila melanogaster. Genetics* 192:33–53.

Leigh, E.G. (1971). The Parliament of Genes. In: *Adaptation and Diversity. Natural History*

and the Mathematics of Evolution, pp. 245–254. Freeman, Cooper & Company, San

Francisco, CA, USA.

Lindholm, A.K., Dyer, K.A., Firman, R.C., Fishman, L., Forstmeier, W., Holman, L., *et al.*

(2016). The Ecology and Evolutionary Dynamics of Meiotic Drive. *Trends Ecol. Evol.*

31:315–326.

Lyth, S., Betancourt, A.J., Price, T.A.R. & Verspoor, R.L. (2023). The suppression of a

selfish genetic element increases a male's mating success in a fly. *Ecol. Evol.* 13:1–13.

Lyttle, T.W. (1991). Segregation distorters. *Annu. Rev. Genet.* 25:511–557.

Meiklejohn, C.D. & Tao, Y. (2010). Genetic conflict and sex chromosome evolution. *Trends*

Ecol. Evol. 25:215–223.

Mendel, G. (1865). Versuche über Pflanzenhybriden. In: *Verhandlungen des*

naturforschenden Vereines in Brünn, Bd. IV für das Jahr 1865, pp. 3–47.

Patten, M.M., Schenkel, M.A. & Ågren, J.A. (2023). Adaptation in the face of internal

conflict: the paradox of the organism revisited. *Biol. Rev.* 98:1796–1811.

Patten, M.M., Úbeda, F. & Haig, D. (2013). Sexual and parental antagonism shape genomic

architecture. *Proc. R. Soc. London Ser. B-Biological Sci.* 280:20131795.

Price, T.A.R., Windbichler, N., Unckless, R.L., Sutter, A., Runge, J.N., Ross, P.A., *et al.*

(2020). Resistance to natural and synthetic gene drive systems. *J. Evol. Biol.* 33:1345–

1360.

R Development Core Team. (2023). A Language and Environment for Statistical Computing.

R Foundation for Statistical Computing, Vienna, Austria.

RStudio Team. (2023). RStudio: Integrated Development Environment for R. RStudio, Inc.,

Boston MA, USA.

Sandler, L. & Novitski, E. (1957). Meiotic drive as an evolutionary force. *Am. Nat.* 857:105–

110.

Schenkel, M.A. (2024). Transitions in sex determination mechanisms through parental and

sexual antagonism. *Heredity (Edinb).*, doi: 10.1038/s41437-024-00717-x.

Schenkel, M.A., Pen, I., Beukeboom, L.W. & Billeter, J.-C. (2018). Making sense of

intralocus and interlocus sexual conflict. *Ecol. Evol.* 8:13035–13050.

Scott, T.W. & West, S.A. (2019). Adaptation is maintained by the parliament of genes. *Nat.*

Commun. 10:5163.

Tolosana, I., Willis, K., Burt, A., Gribble, M., Nolan, T., Crisanti, A., *et al.* (2024). A Y

chromosome-linked genome editor for efficient population suppression in the malaria

vector *Anopheles gambiae*. *bioRxiv* 2024.05.14.594116.

Úbeda, F., Patten, M.M. & Wild, G. (2014). On the origin of sex chromosomes from meiotic

drive. *Proc. R. Soc. B Biol. Sci.* 282:20141932.

van Doorn, G.S. (2009). Intralocus sexual conflict. In: *Year in Evolutionary Biology 2009*,

pp. 52–71.

van Doorn, G.S. & Kirkpatrick, M. (2010). Transitions between male and female

heterogamety caused by sex-antagonistic selection. *Genetics* 186:629–645.

van Doorn, G.S. & Kirkpatrick, M. (2007). Turnover of sex chromosomes induced by sexual

conflict. *Nature* 449:909–912.

Vaz, S.C. & Carvalho, A.B. (2004). Evolution of autosomal suppression of the *Sex-Ratio* trait

in *Drosophila*. *Genetics* 166:265–277.

Veller, C. (2022). Mendel's First Law: partisan interests and the parliament of genes.

Heredity (Edinb). 129:48–55.

West, S.A. (2009). *Sex Allocation*. Princeton University Press, Princeton, NJ, USA.

Wickham, H., Averick, M., Bryan, J., Chang, W., D'Agostino McGowan, L., François, R., *et*

al. (2019). Welcome to the Tidyverse. *J. Open Source Softw.* 4:1686.

Supplementary Material

The results presented in the main text sough to consider the effect of varying the strength of

the Assister and the Driver, while keeping other parameters constant and set to optimal

conditions (e.g. low fitness cost, low recombination rate). Here, we investigate the effect of

varying the fitness cost, z, of the D_2 allele, and the recombination rate, r, between the *A* locus

and the *D* locus (**Supplementary Figure S1** and **S2**).

Supplementary Figure S1. Low recombination rate (r) and low *trans*-drive cost (z) favor sex

ratio distortion in the sexual antagonism model**.** Sex ratio skews the most heavily toward males

when z and r are both minimized (G=1,000, $h_A = 0.5$). The effect of the cost of drive (z)

seems to be slightly larger than the effect of recombination (r).

Supplementary Figure S2. Low recombination rate (r) and low *trans*-drive cost (z) favor sex

ratio distortion in the male-limited *cis* segregation distorter model. Sex ratio skews the most

heavily toward males when z and r are both minimized ($G = 1,000, h_A = 0.0$). The effect of

the cost of drive (z) seems to be slightly larger than the effect of recombination (r).

We also present two further selective scenarios for the Assister locus, parental antagonism and

co-segregation. We consider these scenarios to be less biologically likely than those presented

in the main text, though an interesting extension of the same logic that an autosome that favors

being in males will enable remote-control Y drive.

<u>Parentally antagonistic selection on *Assister*</u>

Parental antagonism refers to the scenario in which the fitness effect of a gene depends on the

parent from which it was inherited. Under parentally antagonistic selection, A_1 and A_2

experience different fitness depending on the parent of origin. We assume that A_1 is beneficial

to fitness when inherited maternally and costly when inherited paternally, and vice versa for

A_2. The partial fitness score of an individual with genotype $A_i A_j$ is given by:

658
$$w_{11} = 1 - bt$$

659
$$w_{12} = 1$$

660
$$w_{21} = 1 - t$$

661
$$w_{22} = 1 - at$$

662 Here, $t \in [0,1]$ represents the fitness effect of the wrong parent of origin, $a, b \in [0,1]$ indicates

663 the dominance of a single costly allele inherited maternally or paternally, respectively.

Supplementary Figure S3. Dynamics and scope of Y-chromosome drive via the *trans*-acting

Distorter causing X-shredding, linked to a parentally antagonistic *Assister* locus. (A) Changes

in the allele frequency of A_2D_2, and the X- and Y-chromosome across 2,000 generations. (B)

Changes in the linkage disequilibrium of *Assister-Distorter*, *Assister-XY*, and *Distorter-XY*

across 500 generations. (C) Haplotype frequencies over time. A_1D_2X and A_1D_2Y are near-zero,

as D_2 is introduced on an A_2 background and recombination is low. The shaded area represents

the frequency of D_2 and primarily comprises the haplotypes A_2D_2Y and A_2D_2X.. (D) D_2

equilibrium frequency after 2,000 generations. (E) Population sex ratio after 2,000 generations.

Parameter values (if not otherwise defined): $a, b = 0.5, t = 0.3, k_D = 0.25, z =$

$0.01, r = 0.001$.

Cosegregation of *Assister*

Our final mechanism for *trans*-acting sex chromosome drive is the most direct: cosegregation

of the autosome with the Y chromosome. Here, A_2 encodes a structural variant that creates a

physical attachment between the autosome and the Y chromosome. We assume that an A_1A_2

(or A_2A_1) heterozygote, A_2 is co-inherited with Y at a rate of c and A_2 with X at a rate of (1-c),

where $c \in [0,1]$ is the cosegregation rate. We assume that A_2 negatively affects its bearer's

fitness, considering that this co-segregation mechanism may induce aneuploidy.

The partial fitness score of a male with genotype A_iA_j is given by:

$$w_{M_{11}} = 1$$

$$w_{M_{12}} = w_{M_{21}} = 1 - s$$

$$w_{M_{22}} = 1 - s$$

Here, $s \in [0,1]$ represents the fitness effect of the A_2 allele in males.

When paired with A_2 in the co-segregation model, D_2 is rarely inherited with the X chromosome

and does not have the opportunity to shred it. Therefore, the partial fitness score of a male with

genotype D_iD_j, where D_2 is not linked with A_2, is:

$$w_{M_{11}} = 1$$

$$w_{M_{12}} = w_{M_{21}} = 1 - z$$

$$w_{M_{22}} = 1 - z$$

Here, $z \in [0,1]$ measures the loss of total fertility, and we assume that the *Distorter* locus is

neutral in females. However, if D_2 is linked with A_2, then the partial fitness score of a male

with genotype D_iD_j is 1.

$$697 \qquad w_{M_{11}} = 1$$

$$698 \qquad w_{M_{12}} = w_{M_{21}} = 1$$

$$699 \qquad w_{M_{22}} = 1$$

Here, c is the frequency of cosegregation, as defined above.

Supplementary Figure S4. Dynamics and scope of Y-chromosome drive via the *trans*-acting

Distorter causing X-shredding and exhibiting co-segregation with the Y-chromosome. (A)

Changes in the allele frequency of A_2D_2, and the X- and Y-chromosome across 500 generations.

(B) Changes in the linkage disequilibrium of *Assister-Distorter*, *Assister-XY*, and *Distorter-*

XY across 500 generations. The shaded area represents the frequency of D_2 and primarily

comprises the haplotypes A_2D_2Y and A_2D_2X. (C) D_2 equilibrium frequency after 500

generations. (D) Population sex ratio after 1,000 generations. (E) Population sex ratio after

1,000 generations. Parameter values (if not otherwise defined): $c = 0.75, s = 0.01, k_D = $

$0.25, z = 0.01, r = 0.001$.

Considering the XY to XO transition models presented in the main text, we sought to

understand whether polymorphism for O and Y as the male-determining genetic factor was

possible. We found a narrow range of possibility, represented in green in Figure S5.

Supplementary Figure S5. Range of parameter space for which Y-/O-chromosome

polymorphism is possible; polymorphism is defined as both the Y- and O-chromosome having

frequencies above 1%. (A) Sexually antagonistic *Assister*, where parameter values: $k_D = $

$0.25; s, t = 0.5; h_A = 0.5; z = 0.01; w = 0.001; r = 0.001$. (B) Male-limited *cis*-segregation

distorting *Assister*, where parameter values: $k_D = 0.25; k_A = 0.25; h_A = 0.0; z = 0.01; w = $

$0.001; r = 0.001$.

Supplementary Figure S6: Loss of autosomal suppression of segregation distortion mimics remote control sex chromosome meiotic drive. (B) a Y-chromosomal segregation distorter Y_D evolves and causes male-biased sex ratios. (C) An autosomal locus S evolves to reduce Y-chromosomal segregation distortion and restore the sex ratio to equality. (D) A second autosomal locus (A) evolves nearby the S locus, and experiences sex-specific segregation distortion. (E) Presence of A renders loss-of-function at the S locus advantageous to the S-A complex; evolution of the non-suppressing S_0 variant re-enables segregation distortion by Y_D, and by extension male-biased sex ratios.

733 **Supplementary Tables**

734 **Supplementary Table S1:** Overview of model parameters.

Variable	Description	Standard value or range
k_D	strength of trans segregation distortion	(0,0.5] standard value: 0.25
z	fitness effect of the D_2 allele in homozygotes	[0,1] standard value = 0.01
r	Recombination between the A locus and D locus	[0,1] standard value = 0.001
h_A	Dominance of A_2	[0.5,1] standard value = 0.5 for sexual antagonism, 0.0 for cis drive
t (SA only)	Fitness cost of A_2 in females	[0,1] standard value = 0.5
s	SA: Fitness cost of A_1 in males Cis drive: fitness cost of A_2 in both males and females	[0,1] standard value = 0.5
k_A (*cis* drive only)	Strength of *cis* segregation distortion	(0,0.5] standard value = 0.25
b	Scalar of less fit paternally inherited allele	(0,1] standard value = 0.5

a	Scalar of less fit maternally inherited allele	(0,1] standard value = 0.5
t	Cost of less fit allele from either parent	(0,1] standard value = 0.3

736 **Supplementary Table S2:** Overview of fitness scores for all combinations of *Assister* and

737 *Distorter* genotypes.

Assister type	Assister genotype	Distorter genotype	Female fitness	Male fitness
Sexual antagonism	A_1A_1	D_1D_1	1	$1-s$
		$D_1D_2,$ D_2D_1	1	$(1-s)(1-z)$
		D_2D_2	1	$(1-s)(1-z)$
	$A_1A_2,$ A_2A_1	D_1D_1	$1-h_At$	$1-h_As$
		$D_1D_2,$ D_2D_1	$(1-h_At)$	$(1-h_As)(1-z)$
		D_2D_2	$(1-h_At)$	$(1-h_As)(1-z)$
	A_2A_2	D_1D_1	$1-t$	1
		$D_1D_2,$ D_2D_1	$(1-t)$	$1-z$
		D_2D_2	$(1-t)$	$1-z$
Cis-segregation distortion	A_1A_1	D_1D_1	1	
		$D_1D_2,$ D_2D_1	1	$1-z$

		D_2D_2	1	$1-z$
	$A_1A_2,$ A_2A_1	D_1D_1	$(1-h_As)$	
		$D_1D_2,$ D_2D_1	$(1-h_As)$	$(1-h_As)(1-z)$
		D_2D_2	$(1-h_As)$	$(1-h_As)(1-z)$
	A_2A_2	D_1D_1	$(1-s)$	
		$D_1D_2,$ D_2D_1	$(1-s)$	$(1-s)(1-z)$
		D_2D_2	$(1-s)$	$(1-s)(1-z)$
Parental antagonism	A_1A_1	D_1D_1	$(1-bt)$	
		$D_1D_2,$ D_2D_1	$1-bt$	$(1-bt)(1-z)$
		D_2D_2	$(1-bt)$	$(1-bt)(1-z)$
	A_1A_2	D_1D_1	1	1
		$D_1D_2,$	1	$1-z$
		D_2D_1	1	$1-z$
		D_2D_2	1	$1-z$
	A_2A_1	D_1D_1	$(1-t)$	
		$D_1D_2,$	$(1-t)$	$(1-t)(1-z)$
		D_2D_2	$(1-t)$	$(1-t)(1-z)$
	A_2A_2	D_1D_1	$(1-at)$	
		$D_1D_2,$ D_2D_1	$(1-at)$	$(1-at)(1-z)$

		D_2D_2	$(1-at)$	$(1-at)(1-z)$