Iso-orientation bias of layer 2/3 connections: the unifying mechanism of spontaneous, visually and optogenetically driven V1 dynamics

Tibor Rózsa[1†], Rémy Cagnol[1†], Ján Antolík[1]

[1]*Faculty of Mathematics and Physics, Charles University, Czech Republic*
† These authors contributed equally to this work.

Abstract

Functionally specific long-range lateral connectivity in layer 2/3 of the adult primary visual cortex (V1) supports the integration of visual information across visual space and shapes spontaneous, visual and optogenetically driven V1 activity. However, a comprehensive understanding of how these diverse cortical regimes emerge from this underlying cortical circuitry remains elusive. Here we address this gap by showing how the same model assuming moderately iso-orientation biassed long-range cortical connectivity architecture explains diverse phenomena, including (i) range of visually driven phenomena, (ii) modular spontaneous activity, (iii) the propagation of spontaneous cortical waves, and (iv) neural responses to patterned optogenetic stimulation. The model offers testable predictions, including presence of slower and iso-tropic spontaneous wave propagation in layer 4 and non-monotonicity of optogenetically driven cortical response to increasingly larger disk of illumination. We thus offer a holistic framework for studying how cortical circuitry governs information integration across multiple operating regimes.

Introduction

A fundamental function of sensory systems is integrating information across sensory space. In vision, this is often studied by sensory stimulation and analysis of lateral interactions between spatially offset cortical sites (*1–3*). However, such sensory-driven approaches often conflate the complexities of feed-forward processing with recurrent lateral interactions. The primary visual cortex (V1) exhibits complex dynamics even without visual input, either during spontaneous activity, or in response to artificial external stimulation, offering an alternative means to study cortical integration without feed-forward confounds. Several phenomena highlight the interconnectedness of visually and non-visually driven dynamics:

Spontaneous travelling waves (STW): Waves of activity propagating across V1 (*4–6*) at speeds consistent with unmyelinated lateral axons (*7*), with propagation biases aligned with cortical functional organisation (*8*), and influencing behaviour (*7, 9*).

Modular spontaneous activity: Spontaneously emerging activity patterns in V1 are correlated with functional maps (*10*), linking resting-state dynamics with visual coding.

Optogenetically evoked patterns of activity: Optogenetic perturbations of ongoing V1 activity bridge the resting and visually evoked dynamics (*11–14*), offering a way to study cortical interactions without engaging retino-cortical pathways.

Ultimately, cortical dynamics across these different driving regimes (spontaneous, visually
driven, and optogenetically driven) rely on a shared neural substrate. Yet, a systematic
understanding of how this common substrate supports the diverse phenomena involved in
visual integration, and under different cortical regimes, remains elusive.

Bridging this gap requires a comprehensive approach capable of simultaneously addressing
anatomy, network dynamics, and function, which computational models are uniquely suited
for. While offering valuable insights, previous computational studies fall short of unifying the
full range of dynamical regimes and levels of integration. Recent phenomenological models
have shown how modular activity arises spontaneously or through optogenetic stimulation
in young animals (*10*, *14*), while others investigated travelling waves (*15*, *16*), optogenetic
stimulation (*17*) or statistics of spontaneous activity (*18*) in adult V1. But all these models
did not consider visually specific processing or anatomy. The model of Davis et al. (*9*)
included structured connectivity and demonstrated functionally biassed spontaneous
activity, but did not link to visually or optogenetically mdriven processing. Finally, Cai et al.
(*19*) demonstrated modular spontaneous activity in a single-layer V1 model, but did not
address spontaneous wave propagation or external stimulation.

To explain how V1's shared neural substrate supports spatial visual integration across
diverse dynamical regimes, we developed a unified model of generic adult columnar V1
(Figure 1A) that accurately simulates visual processing, spontaneous travelling waves,
modular spontaneous activity, and functionally specific optogenetic modulation (Figure 1C).
By comparing to data from multiple species, we demonstrate that the model replicates (i)
distributions of wavelength and propagation speed in spontaneous waves observed in
marmoset, along with biases toward iso-oriented propagation seen in macaque and cat V1,
(ii) modular spontaneous activity correlated with orientation maps in both excitatory and
inhibitory neurons in ferret V1, (iii) impulse-response properties to both full-field
homogeneous and patterned optogenetic stimulation in ferret V1, and (iv) range of visually
driven phenomena. The model yields testable predictions including slower, functionally
independent spontaneous wave propagation in layer 4 and non-monotonic response of
neurons in the surround of a disk of optogenetic stimulation that is increasing in diameter.
Parametric analysis of the model reveals that functional specific lateral interactions in layer
2/3 are critical for emergence of these spontaneous, visually driven and optogenetically
driven phenomena (Figure 1C), unifying an unprecedented range of previously fragmented
experimental findings (*4*, *8*, *10*, *12–15*, *20*). This study advances understanding of how
these diverse processes are interlinked, moving towards a holistic understanding of sensory
processing in the adult brain.

Results

Here we present a generic model of adult columnar V1 derived from our previous work (*21*) (see Methods for full list of changes), which we will compare to data from multiple species (macaque, cat, ferret and marmoset). It considers the first two stages of V1 processing - the cortical layers 4 and 2/3, each populated by excitatory and inhibitory cells modelled as adaptive exponential integrate-and-fire neurons (Figure 1A). Layer 4 receives input from a phenomenological model of the retino-thalamic pathway, with feed-forward receptive field templates globally organised according to pre-computed retinotopic and orientation preference maps (see Methods). The cortical model circuit follows multiple connectivity rules: the probability of connectivity falls off with distance, intra-layer 4 connectivity follows push-pull organisation, and the long-range layer 2/3 connections exhibit a moderate iso-orientation bias (*22–24*); Figure 1B). The model was calibrated using anatomical and physiological data from cat and macaque, but we demonstrate here that it also explains activity dynamics in marmoset and ferret V1. As with its predecessor (*21, 25*), the new model offers quantitatively accurate account of a broad range of visually evoked phenomena, including orientation tuning of spikes, membrane potential and conductances, distribution of modulation ratios, differences between flashed vs. continuous stimulation, and cross-trial spiking precision modulation by stimulus (see Supplementary figures S1-5).

Spontaneous travelling waves exhibit iso-oriented bias of propagation dependent on functionally biassed connectivity in layer 2/3

Voltage-sensitive dye and local field potential (LFP) studies have shown that cortical activity can propagate as travelling waves on both macroscopic and mesoscopic scales under various conditions (*4–6*). To determine if the model's spontaneous activity exhibits wave-like patterns we generated a synthetic LFP signal from the neurons' input current (*26*), which we then convolved with a 2D Gaussian (SD=250 µm) to account for the spatial pooling of LFPs in electrode recordings (*27*) (see Methods). The signal was then wide-band filtered from 5 to 100 Hz. As illustrated in Figure 2A, the synthetic LFP signal propagates across the model cortical surface as a STW.

We next wished to study the relationship between spontaneous wave propagation and the intrinsic functional maps identified by Nauhaus et al. in cats and macaques (8). They used spike-triggered LFP (stLFP), which enables investigating how spiking activity at one reference recording location affects the LFP activity across all other recording sites. They found that the delay of the peak of the stLFP increased and that the amplitude of the peak decreased with distance from the reference electrode. We repeated their analysis using the spike trains of 207 model neurons as reference to compute the stLFPs (see Methods).

As in the experimental data, we observe that the delay of the peak of stLFP increases with distance between recording sites (Figure 2B), and that its amplitude decreases exponentially based on distance (Figure 2C), confirming the validity of the stLFP method implementation in our model. The mean propagation speed of the stLFP signal is 0.52 m/s, within one standard deviation from the value of 0.31 m/s reported in cat V1 by Nauhaus et

al. (Figure 2D). For each reference electrode, Nauhaus et al. used an exponential function to fit the relationship between the stLFP amplitude and the distance to the reference electrode, and then computed the stLFP residuals by subtracting the value predicted by the exponential fit from the stLFP amplitude at each electrode. We applied the same fitting procedure (Figure 2C) and computed stLFP residuals for all but two reference neurons for which the fitting performances were poor (see Methods). Crucially, Nauhaus et al. observed significantly higher amplitudes of stLFP residuals at recording sites with orientation preference congruent to the reference electrode, than for sites with orthogonal preference, indicating functionally specific wave propagation. We find that, for a given reference neuron, iso-oriented recording sites has significantly higher stLFP residuals than recording sites in orthogonal domains (Iso-oriented: 0.0124 ± 0.0008, ortho-oriented: -0.0129 ± 0.0008, p = 4.74×10^{-31}, n = 205, two-sided Wilcoxon signed-rank test) (Figure 2E). This is inline with the observations of Nauhaus et al., although the magnitude of the iso-orientation bias they report is higher. Overall these findings indicate that cortical STW propagate preferentially towards iso-oriented domains in the model as observed experimentally, confirming that we successfully account for functionally specific spontaneous V1 dynamics. Furthermore, we performed the same analysis after removing the functional bias in the long-range excitatory connectivity in the model layer 2/3. After this manipulation, the stLFP residuals are no longer higher for iso-oriented recording sites than for orthogonal recording sites (Iso-oriented: -0.0020 ± 0.0008, ortho-oriented: -0.0001 ± 0.0008, p = 0.07, n = 201, two-sided Wilcoxon signed-rank test) (Figure 2E), demonstrating that functionally specific connectivity is critical for the iso-oriented bias of STW propagation.

Next, Davis et al. offered a detailed characterization of the spontaneous wave properties in marmoset MT using the Generalized Phase analysis (7, 15). We applied the Generalized Phase analysis to compute the instantaneous phases of the wideband filtered signal (Figure 2F), yielding the distribution of wavelengths and propagation speeds of the synthetic LFP (see Methods) which we compared to Davis et al. marmoset data. Prior to convolution, the wavelengths of our synthetic LFP signal are significantly higher than in the spatially shuffled signal, demonstrating that the signal is indeed spatially structured (Figure 2J). After applying the Gaussian convolution to account for spatial pooling, the distribution of the wavelengths in the signal closely matches experimental data in marmoset MT at high and intermediate wavelengths. However, we observe a greater proportion of low wavelengths, likely due to the limited simulated cortical area and the absence of extra-areal input which increases global network synchronisation. The distribution of propagation speeds also lies within the inter-subject variability identified by Davis *et al.* (7, 15) (Figure 2K). The peaks of both the distribution of firing rates and coefficients of variation (CV) match well the experimental data from marmoset MT by Davis et al. (Figure 2GH), ensuring that our network is in a realistic regime of activity similar to that of animal visual cortex.

Due to inaccessibility to imaging techniques, STWs have not been yet analysed in layer 4 of V1. We hypothesised that due to the absence of functionally specific long-range lateral connectivity in layer 4 (28), the propagation of STWs in layer 4 would lack orientation bias. Indeed we find that the stLFP residuals for recording sites of orientation preference congruent to the reference neurons' were not significantly different to stLFP at orthogonally

oriented sites (Iso-oriented: 0.0001 ± 0.0003, ortho-oriented: -0.0005 ± 0.0003, p = 0.23, n
= 207, two-sided Wilcoxon signed-rank test) (Figure 2E). Interestingly, we also find that
propagation speeds and wavelengths in model layer 4 are lower than in layer 2/3 (Figure
2IL). We attribute this to shorter-range connectivity in layer 4 which requires activity to make
multiple synaptic jumps to propagate across the same distance. Polysynaptic transmission
of activity has previously been hypothesized to slow down traveling waves propagation (*6*).
177

Functional specificity of cortical connections dictate correlation structure of
spontaneous activity

Activity of both inhibitory and excitatory neurons is correlated over multiple millimetre
distances, with excitatory activity resembling orientation preference maps in young and adult
carnivore/primate V1 (*10, 20, 29, 30*). To test the presence of this phenomenon in the model
we have first replicated the analysis of Smith et al. (*10*), who recorded from the V1 of young
ferrets: first, we accounted for the spatiotemporal blurring of the underlying neural signal
due to calcium imaging (Figure 3A; Methods; Figure S6) and subsequently calculated
correlation maps (CMs) for each cortical position (Figure 3C), by correlating the activity time
course at the given "seed" cortical position to the activity time courses of all other cortical
positions.

We observed long-range correlations across the model cortex, with excitatory CMs with
seeds at iso-oriented positions in the orientation map being near-identical (Figure 3C), in
line with ferret data (*10*). The model also replicates the finding that excitatory CMs match
inhibitory CMs (Figure 3D). When we quantify their similarity (see Methods) it is consistently
high regardless of seed position (Figure 3E, mean similarity=0.457 ± 0.002, range (-1,1))
(*20*). Additionally, excitatory CMs exhibit high similarity (see Methods) to the orientation
preference map, regardless of seed position (Figure 3F (left), mean similarity=0.744 ±
0.001, range (0,1)). Note that this similarity metric is different from the aforementioned CM
to CM similarity due to the periodicity of orientation preference (*10*). Crucially, the orientation
map similarity diminishes when we remove the orientation bias of Layer 2/3 long-range
horizontal connections (Figure 3E (right), 4F), further underlining the importance of the
functional biassed lateral connectivity in shaping spontaneous cortical dynamics.

Next, replicating the findings of Kenet et al. (*29*) in cat V1, we reconstructed the orientation
preference map from excitatory spontaneous activity by fitting a circular Self-Organizing
Map (SOM) to excitatory CMs (Figure 3G), demonstrating that the orientation preference
encoding can be fully recovered from the correlations of the model spontaneous activity.

Finally, we quantitatively assess how well the model captures the spatial properties of in-
vivo spontaneous activity using several key metrics of spontaneous V1 activity (*10, 20*)
(Figure 3H), overall finding a good match with young ferret data (see Methods for a detailed
description of all metrics). The model reproduces the degree of similarity of excitatory and
inhibitory CMs, long-range correlations and correlation wavelength for both excitatory and
inhibitory activity, while the inhibitory dimensionality and local correlation eccentricity lie only
slightly out of experimental data ranges. Our model does exhibit somewhat higher similarity

between spontaneous activity and orientation preference maps than observed in young
ferrets, which could be due to the absence of measurement noise in the model. Alternatively,
this discrepancy could be explained by the young age of the imaged ferrets, whereby the
orientation representation becomes more refined as the animal matures (see figure 5C in
(10)), reflecting the stronger alignment between orientation maps and spontaneous activity
we observe in the model of adult V1.
222
To validate our model also against data from adult animals, we assess how well the model
reproduces the level of correlation between the orientation preference map and
spontaneous activity measured in adult cat (Kenet et al. (29), Tsodyks et al. (31)) using
alternative metrics. The distribution of correlation coefficients between spontaneous activity
frames and single-condition orientation maps (activity maps in response to gratings of a
single orientation; see Methods) matches that of Kenet et al. (29)), including the control
distribution computed using horizontally flipped single-condition maps (Figure 3I). Next, the
model matches the correlation distributions of spontaneous activity frames to the Preferred
Cortical State (PCS) of neurons (Figure 3J), a proxy for single-condition orientation maps
(Tsodyks et al. (31), see Methods). Additionally, the subset of the above distribution
consisting of only spontaneous frames during which the neuron fired an action potential
(Figure 3K), exhibits a significant positive bias (p<10^{-16}, n=367169, 1-sided one-sample T-
test; mean = 0.1007 ± 0.0003 for model, 0.1591 ± 0.0045 for experiment). Thus, the
predicted firing rate (Figure 3L, ratio of histograms K,J) increases with stronger correlations
to the orientation map, in line with cat data.
238
239 **Iso-orientation biassed layer 2/3 connectivity boosts response to optogenetic**
240 **stimulation congruent with orientation maps**
241
We next investigated the V1 model's response to optogenetic stimulation from the cortical
surface (Figure 4A). To do so, we used a brain-machine-interface (BMI) model we recently
developed (32), that incorporates an LED array, light propagation through neural tissue, and
channelrhodopsin dynamics (see Methods; Figure S8). The BMI model determines the
current injected into each V1 neuron through the channelrhodopsin channels in response to
arbitrary light stimulation patterns. Subsequently the network's response to the optogenetic
input can be simulated.
249
First, we replicated findings from a study that utilised an analogous optogenetic stimulation
setup to modulate young ferret V1 with homogeneous full-field stimulation (14). We
calibrated the intensity of optogenetic stimulation such that the time course of a synthetic
analog of the calcium imaging signal derived from the model's activity (see Methods) aligned
with its biological counterpart ((14); Figure 4B). Experimentally, responses to full-field
optogenetic stimulation vary from trial to trial but CMs derived from such optogenetically
evoked activity closely resemble those derived from spontaneous activity (14). Consistent
with these results, the model full-field optogenetic responses were variable (Supplementary
Figure S9) and the derived CMs closely matched spontaneous CMs (Figure 4C),
irrespective of seed position (Figure 4D). The mean similarity of spontaneous and opto-
response correlation maps was within the range of experimentally measured values (Figure

4E). As with spontaneous activity, the full-field opto-response CMs showed high similarity to the orientation map, but removing the orientation bias of long-range connections from the model decreased the similarity to chance level, demonstrating that the functional biases in later-connectivity mediate the link between orientation maps and optogenetically evoked activity (Figure 4F; see Methods).

Next we replicated Mulholland et al.'s (*12*) experiments that used a thresholded CM seeded at an arbitrary position (endogenous stimulation pattern - see Methods) as optogenetic stimulation pattern (Figure 4G bottom left). Since CMs are highly correlated with the orientation map, they predominantly stimulate cortical positions within a narrow band of orientation preferences (Figure 4G, top left). Following Mulholland et al. (*13*), we also created surrogate stimulation patterns that match the spatial statistics of endogenous patterns, but are unaligned with the spontaneous CMs (see Methods) to serve as controls. Finally, we define the region-of-interest (ROI) for the given optogenetic stimulus as the area of cortex beneath the stimulation pattern (Figure 4G; Methods).

Model responses to stimulation with endogenous patterns matched the maximum-normalised population response time course of young ferret (*12*), both within and outside the stimulation ROIs (Figure 4H, left). This result suggests that our model can account for the spatio-temporal dynamics of V1 responses to patterned optogenetic stimulation. Consistent with the results of Mulholland et al. (*13*), we found that endogenous patterns evoked responses in the model that are stronger and more correlated to the stimulation patterns than those evoked by surrogate stimuli (Figure 4HI). We hypothesised that this is because aligned stimulation amplifies network responses via functionally specific recurrent connectivity, while unaligned stimulation fails to effectively engage this circuitry. This recurrent amplification thus acts as a 'soft winner-take-all' mechanism across the orientation domain, amplifying the dominant orientations while inhibiting the rest, leading to stronger, more correlated responses. In support of our hypothesis, removing the orientation bias of layer 2/3 lateral connections reduced the stimulus-response correlation for endogenous stimuli to a level close to that evoked by surrogate stimuli (Figure 4I, endogenous no bias condition).

To confirm that our model captures the optogenetically driven regime also in adult animals , we replicated the combined visual and optogenetic stimulation experiment of Chernov et al. (*33*), who recorded intrinsic imaging responses in adult macaques. Following Chernov et al., we optogenetically stimulated the horizontal orientation preference column in a circle of 100 µm radius, while simultaneously visually presenting full-field drifting square gratings either iso- or ortho-oriented relative to the stimulated column (Figure 4J, see Methods). We also presented the same gratings without optogenetic stimulation, as the baseline condition. We then identified cortical positions significantly (two-tailed paired t-test for dependent samples, n = 10, $P<0.05$) preferring either the orientation of the optogenetically targeted column (iso-regions) or the orthogonal orientation (ortho-regions) (Figure 4K). We observed that the response in iso-regions increased significantly (two-sided Wilcoxon signed-rank test, $p<0.006$, n = 10) during optogenetic stimulation compared to the visual-only condition (Figure 4L, left), while the response of ortho-regions decreased ($p<0.004$, n = 10). Chernov

et al. (*33*) observed an analogous iso-activity increase and ortho-decrease of the intrinsic imaging signal when stimulating the vertical orientation column (Figure 4L, right).

Orientation specific connectivity is necessary for sharpening of V1 response to optogenetic stimulation

Having access to the underlying spiking responses of all neurons within the 5×5 mm^2 section of modelled cortex, we wished to gain deeper insight into how optogenetic stimulation engages the functionally specific cortical network.

First we noticed that the spatiotemporal blurring effect of the calcium imaging readout underestimates the precision of neural responses in relation to stimulation, both spatially and temporally (Figure 5A,B). Notably, this temporal smoothing effect of calcium imaging has been recently demonstrated by Kettlewell et al. (*34*), who deconvolved the ferret calcium imaging spontaneous activity recordings using Prior Frame Subtraction (see Methods), revealing faster temporal dynamics. Our model spike data matches the timescale of the example deconvolved signal provided in their article (*34*), with the first peak of the temporal autocorrelation function occurring at ~226ms in ferret and ~200ms in our model (Figure 5C). Given this alignment, we opted to perform all subsequent analyses directly on model spike responses.

We have shown that model responses to endogenous (OP map correlated) optogenetic stimuli are stronger than to equivalent surrogate stimuli. We hypothesise this is due to orientation-biassed recurrent connectivity selectively boosting the activity of the dominantly active orientation, resulting in a sharper orientation tuning profile than would be expected from the underlying orientation preference map. To quantify how the population activity is distributed across the orientation domain during patterned stimulation, we introduced the measure of Distribution of Activity across the Orientation Domain (DAOD). For each frame of activity, we binned cortical positions by their orientation preference, summed the activity within each orientation bin, and normalised the resulting distribution to a unit-integral density function. For each of the 8 endogenous and 8 surrogate stimulation patterns, we calculated DAOD within the optogenetically stimulated regions and averaged it across all frames of the response and 5 trials (see Figure 5D for example patterns, Supplementary Figure S10 for all patterns).

To quantify the sharpening of DAODs over the distribution of orientation preferences in stimulated regions, we compared the Shannon entropies of these two distributions (Figure 5E), with lower entropies indicating sharper tuning in the orientation domain. For all of the 16 tested stimulation patterns (8 endogenous, 8 surrogate), DAOD entropy was lower than that of the distribution of stimulated orientations. The entropy difference between the DAOD and stimulated orientation distribution was significant compared to spatially shuffled controls ($p<10^{-17}$, two-tailed Z-test, n=1000; for all stimulation patterns, see Methods). This result supports our hypothesis that the network amplifies the response of the dominant orientation preference subpopulation, resulting in a more sharply oriented response than the orientation

distribution of the stimulation pattern, suggesting soft-winner-takes-all dynamics often hypothesised in cortical networks (*35*).

Finally, we found an inverse relationship between DAOD entropy and firing rate across both spontaneous activity and responses to endogenous/surrogate stimulation - the higher the firing rate, the lower the DAOD entropy (Figure 5F). This effect was strongest for endogenous stimulation (Pearson correlation r=-0.55), but still present with surrogate stimulation (r=-0.20) and spontaneous activity (r=-0.27). This relationship is also consistent with soft-winner-takes-all dynamics, which would predict recurrent boosting of any spontaneous or externally induced patterns of activity aligned with the network's functional organisation due to the iso-orientation biassed connectivity.

Complex nonlinear interactions across the cortical space underlie population responses to optogenetic stimulation in V1

Experimental and modelling results discussed in the previous section suggest that stimulation patterns aligned with V1 functional organisation are more effective at driving cortical activity than equivalent unaligned patterns, which we hypothesise is underpinned by the functionally specific connectivity in layer 2/3. To understand these network phenomena more mechanistically, we performed simulations focusing on how lateral centre-surround interactions emanating from a single column shape the response to optogenetic stimulation. To investigate this, we stimulated the model in a circular area of variable diameter centred on a central orientation domain (Figure 6A), and analysed the resulting activations in the surrounding cortical region. We defined the not-directly-stimulated surround as the cortex beyond the largest tested circular stimulation area (300μm) plus an extra 100μm margin to exclude neurons potentially directly activated by scattered light (Figure 6B, left). Crucially, we kept the total luminance input to the cortex constant by rescaling the stimulation intensity for each radius accordingly.

In line with iso-orientation biassed long-range lateral connectivity, we found that activated neurons in the surround of the directly stimulated cortex are more likely to have similar orientation preference to that of the stimulated column (Figure 6B, left). As we increase the radius of illumination, neurons from an increasingly broader range of orientation domains are directly stimulated (Figure 6C). Yet, the activity and the overall orientation selectivity of the excitatory response in the surround is non-monotonic (Figure 6D), in contrast with the monotonic increase of inhibitory activity (Figure 6E). Initially, increasing stimulation radius raises the mean activation of neurons in the surround (Figure 6D, red line), while decreasing their DAOD entropy (Figure 6D, black line). This mirrors the relationship between firing rates and selectivity observed in patterned stimulation (Figure 5F). This initial rise in selectivity is driven by the recurrent boosting of the dominantly stimulated orientation mediated by the iso-oriented long-range connectivity, and consequent inhibition of the non-dominant orientation domains, in analogous manner to the previous results on patterned optogenetic stimulation (Figure 4I). Crucially, as the stimulation radius, and hence the range of stimulated orientations increases, both the mean activity level and DAOD in the surround concomitantly flip their trends. This flip cannot be explained by the total input coming from

the directly stimulated central area, where the total spiking output of excitatory cells increases monotonically (Figure S11DE). Instead, this non-monotonic effect is due to the expanding range of directly stimulated orientations, which gradually diminishes the aforementioned recurrent iso-orientation boosting, leading to reduced activity and less sharp orientation tuning in the surround (Figure 6D). The precise location of this inflection point is the result of a complex interplay between the spatial scale of the orientation map, the spatial extents of long-range excitatory and short-range inhibitory connections, and balance of excitation and inhibition in the network. Nevertheless, the existence of the inflection point at around 100µm radius is an important prediction of the model.

We next wished to understand the difference between how the stimulated centre engages populations of iso-oriented vs. ortho-oriented neurons in the surround, which receive substantially different input from the centre. We found that in the iso-oriented regions, inhibitory activity rose continuously, while mean excitatory activity mirrored the non-monotonic profile of the pooled analysis (Figure 6F, full lines). In contrast, in the ortho-oriented subpopulations, both excitatory and inhibitory responses followed the non-monotonic U-shaped curve (Figure 6F, interrupted lines). This demonstrates the complexity of intra-cortical interactions that can demonstrate qualitatively different behavior, depending on the relative functional properties between the stimulated and recorded neurons.

Afferent vs. lateral interactions ratio, conduction delays, and functional connectivity bias impact cortical waves and modular spontaneous activity

To gain understanding on how key components of the model impact the expression of investigated model phenomena we systematically manipulated model parameters (see Methods), while monitoring the various measures introduced earlier. Here we pinpoint the most interesting findings. More exhaustive results of this parameter search can be found in Figures S12 and S13. Increasing the strength of the afferent thalamo-cortical connections (TC strength) increases the spontaneous activity of the network (Figure S12A) and the mean propagation speeds of spontaneous waves (Figure 7A). We hypothesise this is due to the increase of the resting voltage level, which raises the likelihood of fast monosynaptic propagation of the signals over long distances. Because we found that the overall model activity has a strong impact on its dynamics (Figures 5, S12AEIM), in the following, we dissociate the impact of activity level from the impact of the remaining parameters. We do so by adjusting TC strength for each investigated parameter value combination to match the overall activity level of the base model (see Methods) (Figure S12B-D).

As expected, the propagation speed of the cortical activity decreases with increasing intra-laminar distance-dependent conduction delays, highlighting its association with horizontal connections (Figure 7B). Next, in Figure 7C-F we show that the orientation bias of travelling waves concomitantly changes with both the spontaneous and opto-evoked CMs similarity to the orientation map as we change model parameters, indicating a close association between these phenomena and the lateral connectivity substrate. Specifically, we find that decreasing the relative impact of the lateral vs. feed-forward connectivity (through increasing TC strength) is negatively associated with these three metrics (Figure 7C). They

are also dependent on the E:I balance as increasing the strength of the layer 2/3 excitatory-
to-inhibitory connections (L2/3 E→I strength) also induces a decrease in propagation bias
and both similarity measures (Figure 7E). Finally, all three measures sharply increase as
the orientation bias of long-range horizontal connections increases (L2/3 E→E bias, Figure
7F), underlining the importance of the functional selectivity of the long-range connections
for the coexistence of STWs, modular spontaneous activity and optogenetically evoked
responses.

Discussion

In this study we presented a unified computational model of adult V1 grounded in biologically
established mechanisms and anatomical data, that explains an unprecedented range of
phenomena across multiple dynamical regimes including visually driven processing,
external optogenetic stimulation and spontaneous emerging activity patterns. The model for
the first time reproduces: (i) iso-orientation biassed propagation of spontaneous waves, (ii)
similarity between excitatory and inhibitory correlation maps, (iii) modular activity response
to uniform full-field optogenetic stimulation that is correlated with modular spontaneous
activity, (iv) response ratio inside/outside of optogenetically stimulated area for endogenous
stimuli, and (v) the iso-facilitation and ortho-suppression during simultaneous visual and
single-column optogenetic stimulation.

Our findings further the hypothesis that functionally specific lateral interactions in cortical
layer 2/3 are key to the emergence of modular spontaneous activity, functionally specific
propagation of spontaneous cortical waves and functionally specific responses to patterned
optogenetic stimulation, bridging previously fragmented experimental observations into a
coherent understanding of adult V1 dynamics.

This hypothesis is first corroborated by the finding that preferential propagation of STW into
iso-oriented regions is abolished by removing the orientation bias of the long-range lateral
connections in the model layer 2/3 (Figure 2E). Next, spontaneous activity in V1 (*10*) and
other cortical areas (*36*) reflects the underlying functional organisation of the cortical
circuitry. This manifests in modular events of higher activity which are correlated with the
feature maps encoded across the cortical surface, such as orientation preference or ocular
dominance maps in V1 (*30*). In the macaque, these events are more frequently aligned with
ocular dominance maps (*30*), while in the ferret spontaneous activity events aligned with
orientation preference are more frequent (*10*). In our model, lacking binocular processing,
we demonstrate the encoding of orientation maps in spontaneous activity, and show that
the long-range iso-orientation biassed lateral connectivity in layer 2/3 is necessary for the
emergence of this phenomenon (Figure 3F). Finally, the similarity of the orientation map to
both correlation maps derived from spontaneous activity and from activity evoked by full-
field optogenetic stimulation in ferrets (*14*) suggests that spontaneous and optogenetically
evoked activities are governed by the same underlying substrate. Indeed, we find that the
iso-orientation bias of the layer 2/3 long-range connections is necessary for the emergence
of modular activity in both stimulation conditions (Figure 4F).

Diving deeper into mechanisms of cortical dynamics, our analysis of the spatial scale of
optogenetic stimulation revealed a non-linear relationship between the radius of optogenetic
stimulation and cortical response (Figure 6DEF). We propose that this non-linear
relationship is a consequence of the network operating within an inhibitory stabilised regime
(37, 38). We have shown that our model operates in this regime under both visual (10) and
optogenetic stimulation (Figure S14), where increased input can paradoxically reduce the
activity of both excitatory and inhibitory populations (37, 38). Note that excitatory neurons
in regions iso-oriented with the column at the centre of the opto-stimulated area receive
stronger long-range input from this column due to the iso-orientation bias of the long-range
E-to-E connectivity. In contrast, in ortho-oriented regions it is the inhibitory neurons that
receive relatively stronger input, explaining the simultaneous reduction in the activity of both
excitatory and inhibitory neurons in the ortho-oriented populations at the smaller stimulation
radii which effectively drive a narrow range of orientations in the centre. These findings
demonstrate that in cortical networks with heterogeneous, functionally specific connectivity,
multiple functionally differentiated parts of the network can simultaneously operate in
different regimes, leading to highly complex dynamical effects. This highlights the necessity
to supplement simplified phenomenological models with simulations taking into account the
full complexity of cortical interactions.

The critical assumption of this study is the existence of moderately iso-orientation biassed
long-range connectivity in layer 2/3 of V1. The strongest evidence for functionally biassed
intra-areal connectivity in V1 comes from mouse studies using a combination of
optogenetics and patch-clamp recordings, which demonstrate a higher likelihood of
connections between neurons with similar receptive fields (39, 40). In mammals with
columnar organisation of orientation preference, early studies combining intrinsic imaging
and anatomical labelling provided evidence for long-range iso-orientation biassed
connectivity in both cat (22, 23) and tree-shrew (41). More recently, an optogenetics-based
study confirmed these findings in ferrets, showing that the average orientation preference
of synaptic inputs to layer 2/3 neurons aligns with their orientation preference (24).
Moreover, while highlighting the heterogeneity of long-range connection targets, Martin et
al. (42) show a clear bias toward synapses forming on iso-oriented targets in cat V1 neurons
(see Figure 3, top-right histogram with SI peak above 0.5, indicating an iso-orientation bias).
Overall, there is strong evidence across multiple species for a moderate iso-orientation bias
in long-range layer 2/3 connectivity.

Several other models aimed at studying some of the individual phenomena we reproduce
in our model. Spontaneous travelling waves have been generated in small-scale spiking
neural networks with local distant-dependent recurrent connectivity (43, 44), but these
waves are too dense in comparison to the sparse traveling waves observed in vivo (4). In a
simple model with ring architecture, Muller et al. have shown that the oscillatory activity of
the model could propagate laterally only when the length of the model was within a specific
range (5). Davis et al. designed a topologically organised spiking neural network which also
exhibited spontaneous traveling waves with realistic wavelengths and propagation speeds
properties (15). Importantly, they showed that the presence of the topologically-organized
recurrent connectivity is essential for the emergence of these properties. In a follow up
study, they present an iteration of the model with functionally-biassed long-range

connectivity (*9*). They demonstrate that this connectivity scheme is needed to reproduce
activity phase patterns associated with functionally biassed spiking activity which they
observed in vivo. Their work is complementary to ours as they focus on the functional bias
of modulation of spiking activity induced by phase patterns of waves whereas we focus on
the functional bias of the propagation of the traveling waves, with both studies emphasizing
the importance of functionally-biassed connectivity. Unlike our study, however, Davis et al.
do not investigate iso-orientation bias of activity propagation, do not consider retino-cortical
pathway and hence visual driven regime, and do not investigate optogenetic stimulation.

Next, spontaneous structured activity was modeled by Cai et al. (*19*) in a single-layer spiking
V1 model. Just as our study, Cai et al. assumed iso-orientation biassed excitatory long-
range lateral connectivity and demonstrated spontaneous events that were dominated by a
narrow range of orientations, but did not provide direct quantitative comparison of these
spontaneous correlations to experimental data. Interestingly, spontaneous structured
activity is present already in young animals before eye-opening (*10*), before long-range
correlations are fully formed (*45*). Smith et al. offered a possible resolution to this conundrum
by demonstrating that a model with only short range mexican-hat-like interactions can
explain long-range correlations in spontaneous activity (*10*), but their model did not account
for afferent processing, and hence could not demonstrate the direct correlation of the
spontaneous activity with orientation maps. It is important to emphasize that even if during
development long-range connections might not be necessary for inducing long-range
correlations, this does not imply this is also the case in the adult cortex. For example, the
cortex is much more driven by the afferent pathway in adult animals than in the young in
which visual input is ineffective in evoking reliable neuronal responses (*10, 46*). This is
further corroborated by modeling in Smith et al. (*10*) that shows that the long-range
correlations emerge only if the model dynamics are strongly dominated by lateral
interactions, i.e. when the afferent input is about 1/100th of the lateral interaction strength.
Under the weaker lateral interactions in the adult condition, the long-range connections
might be necessary for the emergence of the long-range correlations in spontaneous
activity, as our model suggests.

As with any model, we identify several key limitations, some of which are primary candidates
for future work. With the increasing range of phenomena we wish to explain within one
model we face the challenge that data from no animal preparation offer full coverage of
phenomena of interest. Thus, our model comparisons involve data from multiple species.
While this approach encourages a focus on principles common across mammals with
columnar V1, rather than species-specific details, it inevitably limits precise quantitative fits
across all measures due to inter-species differences. Nonetheless, our quantitative
comparisons reveal consistently good matches across diverse measures, with only
occasional moderate deviations, confirming the model operates within a biologically
plausible regime. Similarly, due to the lack of suitable data from adult animals, some of the
model validations were against studies that pool data from young ferrets recorded on days
around eye-opening (*10, 12–14, 20*). While the spontaneous structured activity largely
stabilises at eye-opening (see e.g. Figure 4D,E, 5C, SF2F, SF9BDFH of Smith et al. (*10*)),
justifying the comparison to our adult model, this could explain some of the moderate
quantitative discrepancies between our adult model and the ferret data. Next, due to the

absence of feedback from higher visual areas, which has recently been shown to influence
spontaneous activity (*47*), we cannot rule out that some of the studied phenomena could
not be, at least partially, mediated by extra-areal interactions. The present model also does
not consider layers 5 and 6, which, with their extensive lateral arborisation, likely also
contribute to the spatial integration of information in V1. Apical dendrites of layer 5 cells
extend up to the supragranular layers and participate in the generation of the LFP signal in
these layers. Their absence in the model therefore impacts the magnitude of the LFP signal
and might contribute to the discrepancy between the magnitude of the stLFP residuals of
our model and the ones observed in Nauhaus et al. data (*8*). Next, although the model also
provides opportunity to study the interactions between spontaneous and visually evoked
activity and their implications for perception (*7, 9, 21*), we have not explored these yet in the
present manuscript. We have also presented parameter exploration analysis of a number
of key empirically identified model parameters, however given the complexity of the model
there is opportunity to broaden the scope to many more parameters, albeit at exceptional
computational cost given the size of the model and the sheer range of experimental
protocols against which we test it. Similarly, deeper investigation of which model
components contribute to which of the studied phenomena would be of great interest.
Finally, while we made first steps in that direction in Figures 5 and 6, the present model
offers an unparalleled opportunity for deeper analysis of the dynamics of V1 activity, their
relationship to the underlying neural circuits, and the relationship between different cortical
driving regimes, that are yet to be exploited.
594

Materials & Methods

1. V1 Model

The V1 model is a direct iteration of our previous modelling effort (*21*). It is composed of two layers corresponding to 5 × 5 mm patches of layers 4 and layers 2/3 of parafoveal V1. The new model has higher density, with each layer containing 125 000 excitatory neurons and 31 250 inhibitory neurons. Layer 4 gets its input from a simplified model of LGN covering 6° × 6° of visual field and containing 14 400 neurons (see Methods: Retina & Lateral Geniculate Nucleus (LGN)). Layer 2/3 receives its feedforward input from Layer 4. In each layer, the connectivity is dominated by recurrent connections. The connections in Layer 4 are only local, while the excitatory connections in Layer 2/3 are both local and long-range.

We offer detailed justification of modelling choices in our previous study (*21*). Here we will provide justification only for new or modified parameters (see Methods: Modifications from the previous model version).

1.1 Neuron modelling

We simulate cortical neurons with adaptive exponential integrate-and-fire models (*48*):

$$\tau_m \frac{dV}{dt} = -(V - E_L) + \Delta_T \exp\left(\frac{V - V_T}{\Delta_T}\right) + R_m g_{exc}(E_{exc} - V) + R_m g_{inh}(E_{inh} - V) - w \qquad (1)$$

Every time the membrane potential of a neuron crosses a threshold of -40 mV, a spike is generated and the membrane potential of the neuron is reset to a value V_r = -70 mV for a refractory period of 2 ms and 0.5 ms for excitatory and inhibitory neurons respectively. τ_m is the time constant of the membrane and is equal to 8ms for excitatory neurons and to 9 ms for inhibitory neurons. The leak potential E_L and the spike initiation threshold V_T are set respectively to -80 mV and -57 mV for excitatory neurons, and to -78 mV and -58 mV for inhibitory neurons. The threshold slope factor Δ_T of the exponential function is equal to 0.8 mV. The membrane resistance R_m is equal to 250 MΩ for excitatory neurons and to 300 MΩ for inhibitory neurons. The excitatory and inhibitory reversal potentials E_{exc} and E_{inh} are respectively set to 0 mV and -80 mV. The adaptation current w is further detailed in Table S2.

1.2 Retina & Lateral Geniculate Nucleus (LGN)

ON and OFF-center LGN neurons are modelled as leaky integrate-and-fire cells. They are driven by injected current computed by convolving stimuli in the model's visual field with a 3D spatiotemporal receptive field kernel based on LGN measurements by Cai et al. (*49*). This convolution is subsequently scaled by contrast and luminance gain to match experimental observations (*50, 51*) of contrast and luminance saturation, as described by Antolik et al. (*21*). The baseline firing rate of LGN neurons is achieved by injecting Gaussian noise current into the neurons, such that the baseline firing rates of both ON and OFF LGN neurons match experimentally observed values (*51*). Due to the relatively small region of

640 visual space our model covers, we do not model the systematic changes in RF parameters
641 with foveal eccentricity and thus assume that all ON and OFF LGN neurons have identical
642 parameters.
643
644 **1.3 Thalamo-cortical connections**
645
646 Every neuron in Layer 4 receives an input from the LGN. The spatial pattern of the thalamo-
647 cortical connectivity is based on a template that is the sum of a Gabor distribution and of a
648 2 dimensional Gaussian.
649
650
651
$$g(x, y, \lambda, \theta, \psi, \sigma, \gamma) = \exp\left(\frac{x'^2 + y'^2\gamma^2}{2\sigma^2}\right)(G + \cos(2\pi x'\lambda + \psi)) \quad (2)$$
652
653
$$x' = x\cos\theta + y\sin\theta$$
654
$$y' = -x\sin\theta + y\cos\theta$$
655
656
Where G represents the relative weight of the Gaussian, and is set at 0.085. In all sheets
neurons are populated at random physical positions that correspond to retinotopic positions
assuming a magnification factor of 1 between LGN and cortex, corresponding to retinotopic
eccentricity parafoveally of roughly 3°. The coordinates x and y represent the retinotopic
position of the LGN neurons relative to the retinotopic center of the layer 4 cell. The phase
preference ψ, is generated randomly. The orientation of the Gabor distribution □ is
determined by the position of the neuron on a pre-computed orientation map (Figure 1A).
We set the aspect ratio □ of the Gabor distribution to 2.5, the spatial frequency □ to 0.8
cycles per degree, and the size σ to 0.17°. To generate connections from LGN cells to a
given layer 4 neuron, we take the absolute value of the positive (ON cells) or negative (OFF
cells) part of the template, renormalize it to form distribution, and sample from it the
respective connections to ON or OFF cells.
669
670 Individual layer 4 excitatory and inhibitory cells receive connections from the LGN whose
671 number is drawn from uniform distributions of respective boundaries of [90, 190] and [112,
672 168]. A limitation of the model is that we do not model the dominance of the OFF pathway
673 and systematic variations of the ON-OFF pattern of RF following the orientation maps
674 revealed in recent electrophysiological experiments (*52, 53*).
675
676 **1.4 Cortico-cortical connectivity**
677
Excitatory neurons in layer 4 receive 1000 synapses, whereas layer 2/3 neurons receive
2300 synapses. Inhibitory neurons receive 40% less connections than their excitatory
counterparts. Cortical neurons receive excitatory and inhibitory connections following a ratio
of 4:1. Layer 2/3 neurons receive 22% of their excitatory input through the feedforward
connection originating in Layer 4. Furthermore, layer 4 neurons receive a direct feedback
connection from Layer 2/3 corresponding to 20% of their total excitatory inputs, while the
remaining excitatory synapses are formed by intra-laminar connections. Although there is
little evidence of such direct connections from supragranular layers to layer 4, an important

feedback projection from supragranular layers reaches layer 4 via infragranular layers.
Because we believe that it is important to close this cortico-cortical loop and as infragranular
layers are absent in the model, we have therefore decided to create this direct projection
from layer 2/3 to layer 4 in the model. Removing this feedback projection doesn't
significantly impact the results presented in this study (Figure S15).
691
692 Recurrent connectivity in Layer 4, inhibitory connectivity in Layer 2/3, and excitatory
693 feedforward connectivity from layer 4 to layer 2/3 are only local and distance-dependent.
694 For each of these types of projections, the probability distribution of connections between
695 pairs of neurons is given by the following equation:
696

$$\text{pdf} = p_{\text{dist}} + p_{\text{func}} \tag{3}$$

698 Where p_{func} and p_{dist} are respectively the functional and spatial components of the probability
699 of connection. For these connections, p_{dist} follows a hyperbolic distribution:
700

$$p_{\text{dist}}(d) = \exp(-\alpha\sqrt{\theta^2 + d^2}) \tag{4}$$

where d is the lateral distance between the two neurons, and α and θ are the parameters
shown in table 1 which were obtained by fitting connection probability data (see (*21*) for
details) from Stepanyants et al. (*54*).
For recurrent connectivity in layer 4, the functionally specific component is modelled as a
push-pull scheme, with excitatory connections being more likely to occur between neurons
with correlated afferent receptive field templates (see Methods: Thalamo-cortical
connections), and with inhibitory connections biassed towards neurons with anti-correlated
receptive fields. A correlation c is computed between their receptive fields, here defined
solely based on their afferent inputs from the LGN. This functional component is then the
following:

$$p_{\text{func\_l4}}(c) = \frac{1}{\sigma\sqrt{2\pi}} e^{\frac{-(c-\mu)^2}{2\sigma^2}} \tag{5}$$

713
714 With σ = 1.3, and μ = 1 or -1 respectively when the pre-synaptic neuron is excitatory or
715 inhibitory.
716
717 For inhibitory connections in layer 2/3 there is no functional component and $p_{\text{func\_L23\_inh}}$ is
718 therefore equal to 0 in equation (3). Feedforward projections from layer 4 to layer 2/3 are
719 biassed for connecting neurons with similar orientation preferences. Their functional
720 component is:

$$p_{\text{func\_l4}\rightarrow\text{l23}}(\Delta o) = \frac{1}{\sigma\sqrt{2\pi}} e^{\frac{\Delta o^2}{2\sigma^2}} \tag{6}$$

722
Where Δo is the difference of orientation preference between the pre-synaptic and post-
synaptic cells based on the pre-computed orientation map and with a standard deviation σ
of 1.3 radians. The feedback connections from layer 2/3 to layer 4 are modelled in the same
way as in equation (3). Their functional component has the same formula as equation (6)
with σ is equal to 1.3 and 3 radians when targeting respectively excitatory and inhibitory
neurons. Their spatial component is Gaussian:

$$p_{\mathrm{dist\_l23 \to l4}}(d) = \exp(-\frac{d^2}{2\sigma^2}) \tag{7}$$

with σ = 100 μm. Finally, recurrent excitatory connections in Layer 2/3 consist of both local connections and long-range horizontal projection, the latter being orientation biassed (see Supplementary Figure S16). We model these connections with two spatial components as in Buzás et al. (*23*):

$$\mathrm{pdf}_{\mathrm{l23}} = p_{\mathrm{loc\_l23}} + \alpha p_{\mathrm{lr\_l23}} p_{\mathrm{funct\_l23}} \tag{8}$$

where $p_{\mathrm{loc\_l23}}$ and $p_{\mathrm{lr\_l23}}$ represent the local and long-range spatial component, and follow the same formula as in equation (7) with σ respectively equal to 270 μm and 1000 μm. The functional factor $p_{\mathrm{funct\_l23}}$ has the same formula as in equation (6) with σ equal to 0.45 and to 1.3 radians for connections targeting respectively excitatory and inhibitory cells. □ defines the relative weight of the long range component and is equal to 4.

1.5 Synapses

All synapses of the model are modelled with an exponential decay according to the following formula:

$$g_{\mathrm{syn}}(t) = \overline{g}_{\mathrm{syn}} e^{\frac{-(t-t^*)}{\tau_{\mathrm{syn}}}} \tag{9}$$

Where t* is the spike time and τ_{syn} is the time constant of the exponential decay and is equal to 1.5 ms for excitatory synapses and to 4.2 ms for inhibitory synapses. $\underline{g}_{\mathrm{syn}}$ represents the weight of the synapses and differs according to the type of connection as shown in table S1C

Every synapse of the model also exhibits short-term plasticity through the Tsodyks-Markram model (*55*), with parameters shown in table S3. The delays of the intra-cortical connections of the model are implemented as the sum of two components, one constant and one distance-dependent. The propagation constant of the distance-dependent component is equal to 0.2 s/m for all intra-cortical connections. The constant delays are 1.4 ms and 0.5 ms for excitatory connections targeting respectively excitatory and inhibitory neurons, whereas inhibitory connections have constant delays equal to 1 ms. The delays of thalamo-cortical connections are drawn randomly from an interval of [1.4, 2.4] ms.

1.6 Modifications from the previous model version

The following changes from the previous model version (*21*) were made: The density was increased from 1500 neurons to 5000 neurons per square millimeters to reduce the downsampling of the number of neurons in the model relative to biological densities. This density roughly represents a one fifth downscaling relative to the neuronal density in cat V1 (*56*), which was necessary for our computational simulations to run in a tractable amount of time and memory. Previous studies have shown that both properties of waves (15) and correlation of activity (*57*) are negatively affected by downscaling the neuronal density. We find that downscaling indeed affects the properties we investigated, but that this effect

774 saturates for a downscaling by a factor 5 or less, showing that the density used in this study
775 represents a good compromise between fidelity and computational efficiency (Figure S17).
776
To match the activity regime of the experimental data of Davis et al. (*15*) (see figure 2) and
to compensate the reduction in spontaneous activity caused by the higher density, the
strength of the thalamo-cortical connections was increased from 1.2 nS to 2.2 nS and the
spike initiation threshold was decreased from -58 mV to -56 mV for inhibitory neurons.
Furthermore, to compensate the excess of evoked activity that these changes created, we
increased the strength of the inhibitory synapses from 1 nS to 2 nS as well as reduced their
short-term plasticity recovery time constant in L4 from 70 ms to 30 ms, such that it now
matches the other recurrent cortical connections, hence reducing the number of model
parameters. Additionally, we have increased the strength of the recurrent and feedback
excitatory-to-inhibitory connections targeting layer 4 from 0.22 nS to 0.35 nS so that it
matches the strengths of their equivalent in layer 2/3 further simplifying the model.
Unfortunately, the effective synaptic strength has not been accurately determined
experimentally at such a layer-to-layer and neuron type to neuron type level of granularity.
We therefore tuned these parameters manually to achieve desired dynamical properties of
the cortical network, but such that they remain within physiologically realistic ranges.
792
We have also decreased the standard deviation of the orientation bias of the layer 2/3
excitatory-to-excitatory connections long-range component from 1.3 radians to 0.45 radians,
therefore making them more tuned to orientation in comparison to the previous model
iteration. We found this increased bias was necessary to quantitatively accurately capture
the functional propagation bias of STW and the alignment of the modular spontaneous
activity on the network underlying functional organisation (see Figure 7F). Unfortunately,
the exact strength of the orientation bias was never experimentally quantified due to the
complexity of designing such experiments, but has been qualitatively demonstrated in
several studies (*23, 58*). We have also decreased the conduction delays from 3.3 s/m to 2
s/m, taking it closer to the mean value of 1.67 s/m but further away from the median value
of 3.3 s/m observed in macaques that we used previously (*59*).
804
805

2. Data analysis

Quantitative values in this article are expressed as mean ± SEM.

2.1 Coefficient of variation of the inter-spike-interval

For each neuron, we computed the coefficient of variation (CV) of its inter-spike-interval (ISI) as:

$$CV_{ISI}^2 = \frac{Var[ISI]}{\langle ISI \rangle^2}$$

2.2 LFP signal

To synthesise LFP signal from our point neuron population of L2/3 excitatory neurons we used a proxy method that approximates an LFP signal as the sum of the absolute values of the excitatory and inhibitory currents to estimate the participation of single neurons in the LFP signal of the population (*26*).

$$LFP = \sum_i^n |I_{exc}^{(i)}(t)| + \sum_i^n |I_{inh}^{(i)}(t)|$$

I_{exc} and I_{inh} represent respectively the input excitatory and inhibitory currents. We computed the LFP signal in a 100 x 100 grid with 50 μm spacing representing the 5 mm x 5 mm area of the model, with each portion of the grid corresponding to the signal generated independently by a pool of n=12.5 neurons in average. We then normalised the LFP signal to transform it into a z-score.

To simulate the spatial reach of the LFP, we convolved our signal with a 2D Gaussian with a standard deviation of 250 μm based on the previous modeling study of Davis et al. (*15*). A modeling study from Lindén et al. has estimated that without any input correlations the spatial reach of the LFPs is around 200 μm, corresponding to roughly half the gaussian kernel size used in this study, but found that this value increases with the level of input correlations (*60*). For the low level of pairwise correlations in our model of 0.013 (Figure S1C), Lindén et al. find that the spatial reach of LFP starts increasing by a few hundreds of micrometers. Therefore the choice of 250 μm as the standard deviation of the gaussian kernel is a plausible estimate in our context.

Ultimately, the current understanding allows only for rough choice of the kernel width. We therefore verified the impact of the kernel width on the key metrics of the spontaneous waves in V1 for which the synthetic LFP is used, that we present in Figure S18. We find that halving the kernel width moderately but significantly reduces the estimated wavelength and speed of the spontaneous waves, while the magnitude of the iso-orientation propagation bias increases. Doubling the kernel width has the opposite effect on all metrics.

2.3 Wave properties analysis

We applied the Generalised Phase (GP) analysis (*7, 15*) on the computed three dimensional LFP signal of the 4mm x 4mm central portion of the model. We first applied a band-pass Butterworth filter (5-100 Hz, 4th order, forward-backward) on the computed LFP signal. Our results are however not significantly affected if using a narrower band of frequency in the filtering (Figure S19). We then computed the analytic signal after applying an Hilbert transform, based on fast Fourier transform, on the wideband filtered LFP signal. The phase ϕ of the signal was computed as the argument of the analytic signal. The instantaneous frequency of the signal was calculated as the time derivative of its phase:

$$f_{\mathrm{inst}} = \frac{\partial \phi}{\partial t}$$

As high frequencies in the signal induce sequences of n_t points with negative frequencies, we found these data points with negative instantaneous frequencies and performed a cubic interpolation along the temporal dimension on the phase of these n_t as well as on the phase of the next $2n_t$ time points.

The wavelengths of the signal were computed as the inverse of the magnitude of the spatial gradient of the phase of the signal at each time and space points:

$$\lambda_{\mathrm{LFP}} = \frac{1}{|\nabla_{x,y}\phi|}$$

We then computed the propagation speed of the signal as the product of the wavelengths and of the instantaneous frequency of the signal:

$$v_{\mathrm{prop}} = \lambda_{\mathrm{LFP}} f_{\mathrm{inst}}$$

2.4 Spike-triggered LFP

We computed the spike-triggered LFP (stLFP) based on the spiking activity of 207 reference excitatory neurons in the Layer 2/3 of the model following the procedure of Nauhaus et al. (*8*). Every reference neuron was located in the central 4 mm x 4 mm portion of the model. For each reference neuron, we then computed the stLFP as the spike-triggered average of the LFP signal over a 13 x 13 sub-grid of the original LFP grid, keeping a cell size of 50 μm x 50 μm but with now 400 μm of spacing. We then fitted the relationship between stLFP peak amplitude, and the distance between the neuron and the corresponding electrode with an exponential (see Figure 2C). We discarded from further analysis 2 neurons for which the explained variance of the fitting was lower than 0.5. We then computed the stLFP residuals by subtracting the expected stLFP amplitude (determined by the exponential fit) at each electrode from the actually measured amplitude. We assigned an orientation preference to each recording site based on the pre-computed orientation map of the model. Next, for each reference neuron, we binned the residual stLFP of the recording site based on the difference between their assigned orientation preference and the orientation preference of the neuron. We then averaged the stLFP residuals in each bin across recording sites and seed neurons to compute the orientation bias of the stLFP residuals.

2.5 Simulated calcium imaging signal

To facilitate comparisons with calcium imaging experimental data, we constructed a synthetic calcium imaging signal from our model recordings (see Supplementary Figure S6).

891   We recorded model spikes on a 4 × 4 mm grid with 40 µm spacing and binned them into 50
892   ms time bins, forming the basis of our neural signal. We next convolved the neural signal at
893   each spatial position with a temporal kernel (*61*) that characterises the exponential decay
894   of calcium fluorescence following spikes ($\square$=1s): $K = e^{-\frac{t}{\tau}}$
895
Next, to model light absorption and scattering of the fluorescence signals in the cortex, we
convolved the signal with a spatial kernel generated using a beam-spread function model
(*62*). Fluorescence was modelled as isotropic emission from a disk of 20µm diameter
(representing the neuron) at a depth corresponding to the superficial region of cortical Layer
2/3 (150µm). Due to the lack of available scattering data for the ferret cortex, we based our
coefficients on experimental measurements in mouse cortex (*62*). Finally, we applied a
spatial Gaussian bandpass filter to our signal (σ_{low} = 26 µm, σ_{high} = 200 µm), to match the
experimental calcium image processing pipeline of (*10*).
904
905   For simplicity, we omitted the screening for large-scale events used by Smith et al. (*10*), as
906   it had minimal impact on results (see Supplementary Figure S7).
907
908   **2.6 Correlation maps (CMs) and CM similarity**
909
910   We constructed correlation maps (CMs) from the synthetic calcium imaging signal by
911   calculating pairwise correlations across time between all pairs of spatial positions following
912   Smith et al. (*10*). This process resulted in a correlation map for each spatial position,
913   effectively capturing the spatial organisation of spontaneous cortical activity.
914
915   Next, to quantify the similarity of each correlation map (CM) to the orientation map we
916   computed the magnitude of the pairwise correlation coefficient between the CM and the real
917   and imaginary components of the orientation map (Smith et al. (*10*)). Specifically, for a
918   selected "seed" cortical position s, the similarity Sim(s) was calculated as follows:
919
920   $$\text{Sim}(s) = \sqrt{\text{corr}_x(\Re(\text{O}(x)), \text{C}(s,x))^2 + \text{corr}_x(\Im(\text{O}(x)), \text{C}(s,x))^2}$$
921
922   Here, Sim(*s*) represents the similarity at position *s*, C(*s*,*x*) is the correlation value at spatial
923   position *x* of the CM with seed *s*, and O(*x*) is vector-summed orientation map at spatial
924   position *x*.
925
926   We generated the chance level orientation map similarity distribution for our neural signal
927   by calculating the synthetic calcium imaging signal for uniform random noise (with
928   equivalent space and time dimensions as our spontaneous activity recording) and
929   calculating the correlation map similarity for each spatial position in this recording.
930
931   Finally, we calculated the similarity between two distinct correlation maps (CMs) with the
932   same seed position as the Pearson's correlation coefficient between the two maps. To
933   prevent local correlations from inflating the similarity between the networks, we excluded all
934   pixels within a radius of 400 µm around the seed position in both maps. We repeated this
935   process for each seed position, replicating the analysis of Mulholland et al. (*20*).

2.7 Determining orientation preference with Self-Organizing Maps

To reconstruct the orientation preference map of the model cortex, we used a circular Self-
Organizing Map (SOM) with 40 nodes, analogously to Kenet et al. (*29*). We supplied the set
of all correlation maps to the SOM as input features, and ran the algorithm for 1000 iterations
using the publicly available pip *som-pbc* package. We subsequently assigned an orientation
to each node in the SOM with a step of 4.5° (180° / 40), and estimated the orientation
preference of each spatial position by standard vectorial summation.

Since this algorithm only assigns orientations in a circular topology, but cannot recover the
correct phase and direction of the assignment relative to the orientation map, we evaluated
the mean circular distance between the estimated orientations and the true orientation map
for both directions and 1,000 phase shifts from 0 to π. We subsequently selected the phase
and direction with the smallest circular distance.

2.8 Dimensionality of spontaneous activity, correlation wavelength, local correlation eccentricity

955 We calculated all 3 metrics following Mulholland et al. (*20*). We estimated the
956 **dimensionality** of spontaneous activity patterns as follows:

$$d_{\text{eff}} = \frac{(\sum_{i=1}^{N} \lambda_i)^2}{\sum_{i=1}^{N} \lambda^2}$$

Here, λ_i are the eigenvalues of the covariance matrix for the pixels in each frame. Similarly
to Mulholland et al. (*20*), we randomly sub-sampled spontaneous activity frames (n=30
frames, 100 simulations), and took the median of the distribution. We omitted the event
detection step used by Mulholland et al., as it had minimal impact on our results (see
Supplementary Figure S7).

To estimate the **wavelength of correlation maps**, we centered and averaged the
correlation maps in a 1500µm radius across all seed points. To obtain the average
correlation as a function of distance, we then calculated the radial mean with respect to the
center point. The wavelength of the correlation maps was then defined as the distance to
the first local maximum, after the maximum at 0.

To estimate **local correlation eccentricity**, describing the shape of the local correlation
pattern around a seed point, we fit an ellipse (with axes □1, □2) to the contour line at
correlation 0.7 around the seed point. The eccentricity of the ellipse was subsequently
calculated as:

$$\epsilon = \frac{\sqrt{\zeta_1^2 - \zeta_2^2}}{\zeta_1}$$

Here, $\epsilon = 0$ denotes a circle, and increasing values (up to 1) , reflect the increasing
elongation of the ellipse.

2.9 Prediction interval

To determine the range of values in Figure 3H into which new experimental observations
would likely fall, we calculated the 95% prediction interval from the data of each metric. The
prediction interval for data generated from a Gaussian distribution with unknown mean and
variance is given as follows (*63*):

$$\text{PI} = \mu \pm T_a s_n \frac{\sqrt{n+1}}{\sqrt{n}}$$

where μ denotes sample mean, s_n sample standard deviation, n the number of points, T_a
the 100((1 − *p*)/2)th percentile of Student's t-distribution with $n - 1$ degrees of freedom and
$p = 0.95$ the confidence level.

2.10 Single-condition orientation preference map

To construct single-condition orientation preference maps following Kenet et al. (*29*) we
visually stimulated the model cortex with high-contrast full-field sinusoidal gratings of 8
orientations over 10 trials. We then generated frames of model cortical activity by binning
model spikes on a 4×4 mm grid with 40μm spacing, into 50ms time bins, as substitutes for
the Voltage-Sensitive Dye Imaging (VSDI) frames recorded by Kenet et al. The single-
condition orientation preference maps for each orientation were then computed as the mean
over trials of the 50 ms frame from 100 to 150 ms after stimulus onset, matching the analysis
window of Kenet et al. (mean over 5 frames, 100-150 ms after stimulus onset).

1005

1006 Following Kenet et al., we generated control maps by horizontally flipping the 8 single-
1007 condition maps.

1008

1009 The full-field sinusoidal gratings were of 2s duration, 100% contrast with 50 cd/m²
1010 background luminance, 0.8 cyc/° spatial frequency, 2 Hz temporal frequency. We used 10
1011 trials per orientation (compared to 33 in Kenet et al.), due to the lack of measurement noise
1012 in our model.

1013

1014 **2.11 Preferred cortical state**

1015

In the study of Tsodyks et al. (*31*), the Preferred Cortical State (PCS) of each neuron was
calculated by averaging Voltage-Sensitive Dye Imaging (VSDI) frames at time points
corresponding to the neuron's spike times while presenting drifting grating stimuli. In our
analysis, we substituted the VSDI signal frames with model spikes on a 4x4 mm grid with
40μm spacing, binned into 50ms time bins. Spontaneous activity frames were computed
using the same binning procedure, and correlated with the PCS of all Layer 2/3 excitatory
neurons using Pearson's correlation coefficient.

2.12 Timescale of spontaneous activity

We estimated the temporal scale of spontaneous activity in both model and ferret by calculating temporal autocorrelation functions of population activity. For model data, we averaged 120s of raw spiking and simulated calcium imaging signal across space to arrive at population activity.

For ferret data, we extracted the time courses of calcium imaging and deconvolved calcium imaging population activity from Kettlewell et al. (*34*) Figure 1 a,c, and calculated the temporal autocorrelation functions from the extracted traces. As the available deconvolved time course spanned only 4s, the autocorrelation features on longer timescales (>0.5s) might be less reliable.

To verify that our non-convolved model data was indeed equivalent to deconvolved calcium imaging traces, we used the Prior Frame Subtraction method described by Kettlewell et al. (*34*), where each pixel value in each time point was calculated as follows:

$$y_{p,t} = r_{p,t} - \gamma r_{p,t-1}$$

Here, $r_{p,t}$ denotes the calcium imaging signal value at pixel *p* and time *t*, $y_{p,t}$ denotes the deconvolved value, and $\gamma=0.95$ denotes the calcium imaging temporal kernel decay rate over 50ms (frame duration).

2.13 Distribution of Activity across the Orientation Domain (DAOD)

To calculate Distribution of Activity across the Orientation Domain (DAOD) histograms for each frame of activity during the 1-second of simulated optogenetic stimulation we assumed orientation preference binned into 30 intervals (ranging from 0 to π). We added the activity at each cortical position into one of these bins based on the orientation preference at that cortical position, and normalised the binned activity to an integral of 1 to form a distribution of activities across the orientation domain. In some cases (Figure 5D-F), we only included in the analysis positions within a given stimulation pattern's ROI.

To determine if the Shannon entropy difference between the distribution of sampled orientations and the corresponding DAOD is statistically significant we tested the null hypothesis that the activations from which the DAOD is constructed do not depend on spatial position and thus orientation preference. To do so we created a population (n = 1000) of control DAOD distributions by spatially shuffling the activations from which the DAOD is calculated. Next we calculated the absolute difference between the entropy of each control DAOD and the entropy of the distribution of sampled orientations, forming a control distribution. Finally we used a two-tailed Z-test to determine if the absolute entropy difference between the distribution of sampled orientations and the corresponding DAOD is significantly different from the control distribution.

2.14 Parameterizations used for studying the impact of model parameters

We modified the following five parameters of the model to explore how they influence the phenomena we investigate in this study: (i) the strength of the afferent thalamo-cortical connections (TC strength), (ii) the cortico-cortical connections distance-dependent conduction delays, (iii) the orientation bias of the layer 2/3 intra-laminar excitatory-to-excitatory connections (L2/3 E→E bias) computed as the inverse of the standard deviation used to derive the functional component in the layer 2/3 excitatory-to-inhibitory connectivity (see Methods: Cortico-cortical connectivity), (iv) the strength of the layer 2/3 intra-laminar excitatory-to-inhibitory connections (L2/3 E→I strength), and (v) the cortical density of neurons. As parameters (ii-v) impacted the overall activity of the model, we paired these modifications with a change in TC strength to match the mean activity in the model for each model parameterization:

3. Optogenetic stimulation

To simulate the effects of optogenetic stimulation on the V1 model, we utilised a model of BMI setup we developed recently (*32*). Figure S8 summarises the five-tier structure of this optogenetic stimulator construct:

1. Stimulation patterns
2. 2D stimulator array of identical circular light sources at the cortical surface
3. Light propagation in the cortical tissue
4. Illumination-dependent ChrimsonR channelrhodopsin dynamics in transfected cells
5. Large-scale model of the primary visual cortex.

Since the model of the primary visual cortex is detailed in Methods V1 Model, we will only expand on steps 1-4 in this section.

3.1 Stimulation patterns

We define the region-of-interest (ROI) as the cortical area located beneath each stimulation pattern, with uniform stimulation intensity across active pixels in each pattern. We present 3 types of stimuli to the model cortex:

1. **Full-field stimulation** - ROI covers the entire array.
2. **Endogenous patterns** - ROI corresponds to regions in the correlation map (CM) surrounding its local maxima, resulting in orientation-specific stimulation (see Figure 4, pattern construction details below).
3. **Surrogate patterns** - equivalent stimulation patterns to endogenous patterns but uncorrelated with the orientation map.

With the goal of matching the stimulation protocols of Mulholland et al. (*12, 13*) as closely as possible, we create the patterned stimuli as follows:

1113

1. **CM selection:** We select CMs with seeds at positions with 8 equidistant orientation preferences (0, 22.5, 45, 67.5, 90, 112.5, 135, 157.5°) - this ensures a variety of different stimulation patterns all following intracortical structure.
2. **Endogenous ROIs**: We define the ROIs as the 25% highest correlation value pixels in each map. This is in line with the stimulation protocols of Mulholland et al. (*12*), who manually selected stimulation ROIs around CM maxima, resulting in approximately 25% of the cortical surface being stimulated (24.27% based on scraped data from Mulholland et al. (*12*) Figure 5i).
3. **Surrogate ROIs**: we rotate endogenous patterns by 90°, resulting in stimuli uncorrelated with the orientation map (mean orientation map similarity = 0.1 as opposed to 0.54 for endogenous stimuli).

To replicate the relative intensity difference used by Mulholland et al. (*12, 13*) between full-field (10 mW) and patterned (7.6 mW) stimulation, we set the intensity of patterned stimulation to 76% of that used for full-field stimulation.

3.2 Stimulator array

We stimulate our in-silico cortical surface with a 4x4mm array of 10μm diameter circular light sources, centred on the cortex. We only stimulate layer 2/3 as the light intensity falloff from the surface is exponential (see Figure S20), resulting in the amount of light reaching layer 4 to be negligible given the assumed illumination intensities. The light intensity at the cortical surface across the array is 3.0×10^{14} photons/s/cm^2.

3.3 Light propagation in cortical tissue

We simulate the propagation of light in the cortical tissue using the simulations of the 'Human Brain Grey Matter' model implemented in the LightTools software, assuming emitted light at 590 nm. The scattering and absorption properties of brain tissue are calculated using the Henyey-Greenstein model (*64*). Based on the work of Jacques et al. (*65*) we set the two key parameters of the model, g (anisotropy factor) and *MFP* (mean free path) to 0.87 and 0.07 mm, respectively. The simulation results were tabulated into a 2D lookup table of light flux values at depth d and radial distance r relative to the centre of the light source to speed up simulations.

3.4 Channelrhodopsin model

To simulate the dynamics of the ChrimsonR channelrhodopsin in response to 590 nm light stimulation, as used in the experiments of Mulholland et al. (*12–14*), we utilize the model of Sabatier et al. (*66*). The electro-chemical behaviour of the ChrimsonR protein is modelled using a 5-state Markov kinetic model (*67*), representing the conformations the protein can adopt. Transitions between these states, which the real system can chemically switch between, each have their own time constant and can be either thermally or photo-induced. Thermal transition time constants are fixed, while photo-transition time constants vary with

1158 light intensity. These transition time constants describe the linear differential equations
1159 governing the evolution of the proportion of channels in each state. The total conductance
1160 of all channels in a single neuron is then derived from the sum of conductances of open
1161 channels. We use model parameters fitted to 590 nm light stimulation experiments by
1162 Sabatier et al. (*66*) on ChrimsonR-expressing HEK293 cells.

1163

1164 **3.5 Combined optogenetic & visual stimulation**

1165

For the comparison of visual-only and visual+optogenetic stimulation protocols, we
presented full-field square gratings, matching the visual stimulus parameters reported by
Chernov et al. (*33*): 4s duration, high contrast (100%, 50 cd/m^2 mean luminance), 4 cyc/s
temporal frequency and 1 cyc/° spatial frequency. As noise levels when recording from our
model cortex were much smaller than with in vivo intrinsic imaging, we used n=10 trials,
compared to n=40 of Chernov et al.. For the visual+optogenetic condition, we used the same
visual stimuli and additionally optogenetically stimulated the orientation column at the center
of our model cortex for the first 600 ms of stimulus presentation within a circular region
(r=100µm), matching Chernov et al.. Trials were separated by 4s of blank stimuli, and were
presented in succession.

1176

Intrinsic imaging signals reflect slow vascular responses, peaking several seconds after
neural activation depending on stimulus duration, and returning to baseline 4-6s after
stimulus cessation (*68*). As such, the 4s visual+opto intrinsic signal response analysed by
Chernov et al. likely contained the response to optogenetic stimulation in the first 600 ms of
the stimulus across its entire time course. In contrast, spiking neural responses in our model
(Figure 5B) and in vivo (Jun & Cardin (*69*) Figure 2C) return to baseline activity within tens
of milliseconds after optogenetic stimulus cessation. As such, we only used the first 600 ms
of each recording (the durations of optogenetic stimulation) for all further visual and
visual+opto analyses.

Bibliography:

1. J. R. Cavanaugh, W. Bair, J. A. Movshon, Nature and Interaction of Signals From the Receptive Field Center and Surround in Macaque V1 Neurons. *J. Neurophysiol.* **88**, 2530–2546 (2002).

2.   C. D. Gilbert, A. Das, M. Ito, M. Kapadia, G. Westheimer, Spatial integration and cortical
dynamics. *Proc. Natl. Acad. Sci.* **93**, 615–622 (1996).

3.   F. Chavane, D. Sharon, D. Jancke, O. Marre, Y. Frégnac, A. Grinvald, Lateral spread of
orientation selectivity in V1 is controlled by intracortical cooperativity. *Front. Syst. Neurosci.* **5**
(2011).

4.   L. Muller, F. Chavane, J. Reynolds, T. J. Sejnowski, Cortical travelling waves: mechanisms
and computational principles. *Nat. Rev. Neurosci.* **19**, 255–268 (2018).

5.   L. Muller, Propagating waves in thalamus, cortex and the thalamocortical system: Experiments
and models. *J. Physiol.* (2012).

6.   T. Sato, I. Nauhaus, M. Carandini, Traveling Waves in Visual Cortex. *Neuron* **75**, 218–229
(2012).

7.   Z. W. Davis, L. Muller, J. Martinez-Trujillo, T. Sejnowski, J. H. Reynolds, Spontaneous
travelling cortical waves gate perception in behaving primates. *Nature* **587**, 432–436 (2020).

8.   I. Nauhaus, L. Busse, M. Carandini, D. L. Ringach, Stimulus contrast modulates functional
connectivity in visual cortex. *Nat. Neurosci.* **12**, 70–76 (2009).

9.   Z. W. Davis, A. Busch, C. Steward, L. Muller, J. Reynolds, Horizontal cortical connections
shape intrinsic traveling waves into feature-selective motifs that regulate perceptual sensitivity.
Cell Rep. **43**, 114707 (2024).

10.   G. B. Smith, B. Hein, D. E. Whitney, D. Fitzpatrick, M. Kaschube, Distributed network
interactions and their emergence in developing neocortex. *Nat. Neurosci.* **21**, 1600–1608
(2018).

11.   X. Huang, Y. M. Elyada, W. H. Bosking, T. Walker, D. Fitzpatrick, Optogenetic Assessment of
Horizontal Interactions in Primary Visual Cortex. *J. Neurosci.* **34**, 4976–4990 (2014).

12.   H. N. Mulholland, H. Jayakumar, D. M. Farinella, G. B. Smith, All-optical interrogation of
millimeter-scale networks and application to developing ferret cortex. *J. Neurosci. Methods*
403, 110051 (2024).

13.   H. Mulholland, S. Tragenap, M. Kaschube, G. Smith, "Selective amplification of recurrent
subnetworks in the developing visual cortex" in *Poster Presented at: Computational and*
Systems Neuroscience (COSYNE) Conference (Lisbon, Portugal, 2024).

14.   H. N. Mulholland, M. Kaschube, G. B. Smith, Self-organization of modular activity in immature
cortical networks. *Nat. Commun.* **15**, 4145 (2024).

15.   Z. W. Davis, G. B. Benigno, C. Fletterman, T. Desbordes, C. Steward, T. J. Sejnowski, J. H.
Reynolds, L. Muller, Spontaneous traveling waves naturally emerge from horizontal fiber time
delays and travel through locally asynchronous-irregular states. *Nat. Commun.* **12** (2021).

16.   J. Senk, E. Hagen, S. J. van Albada, M. Diesmann, Reconciliation of weak pairwise spike–train
correlations and highly coherent local field potentials across space. *Cereb. Cortex* **34**,
bhae405 (2024).

17.   J. Luboeinski, T. Tchumatchenko, Nonlinear response characteristics of neural networks and
single neurons undergoing optogenetic excitation. *Netw. Neurosci.* **4**, 852–870 (2020).

18.   T. C. Potjans, M. Diesmann, The Cell-Type Specific Cortical Microcircuit: Relating Structure
and Activity in a Full-Scale Spiking Network Model. *Cereb. Cortex* **24**, 785–806 (2014).

19. D. Cai, A. V. Rangan, D. W. McLaughlin, Architectural and synaptic mechanisms underlying
coherent spontaneous activity in V1. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 5868–5873 (2005).

20. H. N. Mulholland, B. Hein, M. Kaschube, G. B. Smith, Tightly coupled inhibitory and excitatory
functional networks in the developing primary visual cortex. *eLife* **10**, e72456 (2021).

21. J. Antolík, R. Cagnol, T. Rózsa, C. Monier, Y. Frégnac, A. P. Davison, A comprehensive data-
driven model of cat primary visual cortex. *PLOS Comput. Biol.* **20**, e1012342 (2024).

22. C. Gilbert, T. Wiesel, Columnar specificity of intrinsic horizontal and corticocortical connections
in cat visual cortex. *J. Neurosci.* **9**, 2432–2442 (1989).

23. P. Buzás, K. Kovács, A. S. Ferecskó, J. M. L. Budd, U. T. Eysel, Z. F. Kisvárday, Model-
based analysis of excitatory lateral connections in the visual cortex. *J. Comp. Neurol.* **499**,
861–881 (2006).

24. D. E. Wilson, D. E. Whitney, B. Scholl, D. Fitzpatrick, Orientation selectivity and the functional
clustering of synaptic inputs in primary visual cortex. *Nat. Neurosci.* **19**, 1003–1009 (2016).

25. M. M. Taylor, D. Contreras, A. Destexhe, Y. Frégnac, J. Antolik, An Anatomically Constrained
Model of V1 Simple Cells Predicts the Coexistence of Push–Pull and Broad Inhibition. *J.*
Neurosci. **41**, 7797–7812 (2021).

26. A. Mazzoni, H. Lindén, H. Cuntz, A. Lansner, S. Panzeri, G. T. Einevoll, Computing the Local
Field Potential (LFP) from Integrate-and-Fire Network Models. *PLOS Comput. Biol.* **11**,
e1004584 (2015).

27. S. Katzner, I. Nauhaus, A. Benucci, V. Bonin, D. L. Ringach, M. Carandini, Local Origin of
Field Potentials in Visual Cortex. *Neuron* **61**, 35–41 (2009).

28. K. S. Rockland, Anatomical organization of primary visual cortex (area 17) in the ferret. *J.*
Comp. Neurol. **241**, 225–236 (1985).

29. T. Kenet, D. Bibitchkov, M. Tsodyks, A. Grinvald, A. Arieli, Spontaneously emerging cortical
representations of visual attributes. *Nature* **425**, 954–6 (2003).

30. D. B. Omer, T. Fekete, Y. Ulchin, R. Hildesheim, A. Grinvald, Dynamic Patterns of
Spontaneous Ongoing Activity in the Visual Cortex of Anesthetized and Awake Monkeys are
Different. *Cereb. Cortex N. Y. N 1991* **29**, 1291–1304 (2019).

31. M. Tsodyks, T. Kenet, A. Grinvald, A. Arieli, Linking Spontaneous Activity of Single Cortical
Neurons and the Underlying Functional Architecture. *Science* **286**, 1943–1946 (1999).

32. J. Antolik, Q. Sabatier, C. Galle, Y. Frégnac, R. Benosman, Assessment of optogenetically-
driven strategies for prosthetic restoration of cortical vision in large-scale neural simulation of
V1. *Sci. Rep.* **11**, 10783 (2021).

33. M. M. Chernov, R. M. Friedman, G. Chen, G. R. Stoner, A. W. Roe, Functionally specific
optogenetic modulation in primate visual cortex. *Proc. Natl. Acad. Sci.* **115**, 10505–10510
(2018).

34. L. Kettlewell, A. Sederberg, G. B. Smith, Stereotyped spatiotemporal dynamics of spontaneous
activity in visual cortex prior to eye-opening. *J. Neurosci.*, e1420242025 (2025).

35. U. Rutishauser, J.-J. Slotine, R. J. Douglas, Competition through selective inhibitory
synchrony. *Neural Comput.* **24**, 2033–52 (2012).

36. N. J. Powell, B. Hein, D. Kong, J. Elpelt, H. N. Mulholland, M. Kaschube, G. B. Smith,

Common modular architecture across diverse cortical areas in early development. *Proc. Natl. Acad. Sci.* **121**, e2313743121 (2024).

37. M. V. Tsodyks, W. E. Skaggs, T. J. Sejnowski, B. L. McNaughton, Paradoxical Effects of External Modulation of Inhibitory Interneurons. *J. Neurosci.* **17**, 4382–4388 (1997).

38. H. Ozeki, I. M. Finn, E. S. Schaffer, K. D. Miller, D. Ferster, Inhibitory Stabilization of the Cortical Network Underlies Visual Surround Suppression. *Neuron* **62**, 578–592 (2009).

39. H. Ko, L. Cossell, C. Baragli, J. Antolik, C. Clopath, S. B. Hofer, T. D. Mrsic-Flogel, The emergence of functional microcircuits in visual cortex. *Nature* **496**, 96–100 (2013).

40. H. Ko, S. B. Hofer, B. Pichler, K. A. Buchanan, P. J. Sjöström, T. D. Mrsic-Flogel, Functional specificity of local synaptic connections in neocortical networks. *Nature* **473**, 87–91 (2011).

41. W. H. Bosking, Y. Zhang, B. Schofield, D. Fitzpatrick, Orientation selectivity and the arrangement of horizontal connections in tree shrew striate cortex. *J. Neurosci. Off. J. Soc. Neurosci.* **17**, 2112–27 (1997).

42. K. a C. Martin, S. Roth, E. S. Rusch, Superficial layer pyramidal cells communicate heterogeneously between multiple functional domains of cat primary visual cortex. *Nat. Commun.* **5**, 5252 (2014).

43. P. Gong, C. Van Leeuwen, Distributed Dynamical Computation in Neural Circuits with Propagating Coherent Activity Patterns. *PLoS Comput. Biol.* **5**, e1000611 (2009).

44. A. Keane, P. Gong, Propagating Waves Can Explain Irregular Neural Dynamics. *J. Neurosci.* **35**, 1591–1605 (2015).

45. V. Borrell, E. M. Callaway, Reorganization of Exuberant Axonal Arbors Contributes to the Development of Laminar Specificity in Ferret Visual Cortex. *J. Neurosci.* **22**, 6682–6695 (2002).

46. B. Chapman, M. P. Stryker, T. Bonhoeffer, Development of Orientation Preference Maps in Ferret Primary Visual Cortex. *J. Neurosci.* **16**, 6443–6453 (1996).

47. A. Morales-Gregorio, A. C. Kurth, J. Ito, A. Kleinjohann, F. V. Barthélemy, T. Brochier, S. Grün, S. J. Van Albada, Neural manifolds in V1 change with top-down signals from V4 targeting the foveal region. *Cell Rep.* **43**, 114371 (2024).

48. W. Gerstner, R. Brette, Adaptive exponential integrate-and-fire model. *Scholarpedia* **4**, 8427 (2009).

49. D. Cai, G. C. DeAngelis, R. D. Freeman, Spatiotemporal receptive field organization in the lateral geniculate nucleus of cats and kittens. *J. Neurophysiol.* **78**, 1045–1061 (1997).

50. V. Bonin, The Suppressive Field of Neurons in Lateral Geniculate Nucleus. *J. Neurosci.* **25**, 10844–10856 (2005).

51. J. Papaioannou, A. White, Maintained activity of lateral geniculate nucleus neurons as a function of background luminance. *Exp. Neurol.* **34**, 558–566 (1972).

52. J. Z. Jin, C. Weng, C.-I. Yeh, J. A. Gordon, E. S. Ruthazer, M. P. Stryker, H. A. Swadlow, J. M. Alonso, On and off domains of geniculate afferents in cat primary visual cortex. *Nat. Neurosci.* **11**, 88–94 (2008).

53. Y. Wang, J. Jin, J. Kremkow, R. Lashgari, S. J. Komban, J. M. Alonso, Columnar organization of spatial phase in visual cortex. *Nat. Neurosci.* **18**, 97–103 (2014).

54. A. Stepanyants, J. A. Hirsch, L. M. Martinez, Z. F. Kisvárday, A. S. Ferecskó, D. B. Chklovskii,
Local Potential Connectivity in Cat Primary Visual Cortex. *Cereb. Cortex* **18**, 13–28 (2008).

55. H. Markram, Y. Wang, M. Tsodyks, Differential signaling via the same axon of neocortical
pyramidal neurons. *Proc. Natl. Acad. Sci.* **95**, 5323–5328 (1998).

56. C. Beaulieu, M. Colonnier, Number of neurons in individual laminae of areas 3B, 4γ, and 6aα
of the cat cerebral cortex: A comparison with major visual areas. *J. Comp. Neurol.* **279**, 228–
234 (1989).

57. S. J. Van Albada, M. Helias, M. Diesmann, Scalability of Asynchronous Networks Is Limited by
One-to-One Mapping between Effective Connectivity and Correlations. *PLOS Comput. Biol.*
11, e1004490 (2015).

58. Z. Kisvarday, Orientation-specific relationship between populations of excitatory and inhibitory
lateral connections in the visual cortex of the cat. *Cereb. Cortex* **7**, 605–618 (1997).

59. P. Girard, J. M. Hupé, J. Bullier, Feedforward and Feedback Connections Between Areas V1
and V2 of the Monkey Have Similar Rapid Conduction Velocities. *J. Neurophysiol.* **85**, 1328–
1331 (2001).

60. H. Lindén, T. Tetzlaff, T. C. Potjans, K. H. Pettersen, S. Grün, M. Diesmann, G. T. Einevoll,
Modeling the Spatial Reach of the LFP. *Neuron* **72**, 859–872 (2011).

61. H. Lütcke, F. Gerhard, F. Zenke, W. Gerstner, F. Helmchen, Inference of neuronal network
spike dynamics and topology from calcium imaging data. *Front. Neural Circuits* **7** (2013).

62. G. Yona, N. Meitav, I. Kahn, S. Shoham, Realistic Numerical and Analytical Modeling of Light
Scattering in Brain Tissue for Optogenetic Applications. *eNeuro* **3** (2016).

63. *Predictive Inference* (Routledge, 2017; https://www.taylorfrancis.com/books/9780203742310).

64. L. C. Henyey, J. L. Greenstein, Diffuse radiation in the Galaxy. *Astrophys. J.* **93**, 70 (1941).

65. S. L. Jacques, Optical properties of biological tissues: a review. *Phys. Med. Biol.* **58**, R37-61
(2013).

66. Q. Sabatier, C. Joffrois, G. Gauvain, J. Chavas, D. Pruneau, S. Picaud, R. Benosman,
"Modeling the Electro-chemical Properties of Microbial Opsin ChrimsonR for Application to
Optogenetics-based Vision Restoration" (preprint, Neuroscience, 2018);
https://doi.org/10.1101/417899.

67. A. Destexhe, Z. F. Mainen, T. J. Sejnowski, Synthesis of models for excitable membranes,
synaptic transmission and neuromodulation using a common kinetic formalism. *J. Comput.*
Neurosci. **1**, 195–230 (1994).

68. D. Malonek, A. Grinvald, Interactions Between Electrical Activity and Cortical Microcirculation
Revealed by Imaging Spectroscopy: Implications for Functional Brain Mapping. *Science* **272**,
551–554 (1996).

69. N. Y. Jun, J. A. Cardin, Activation of Distinct Channelrhodopsin Variants Engages Different
Patterns of Network Activity. *eneuro* **7**, ENEURO.0222-18.2019 (2020).

70. J. Antolík, A. P. Davison, Integrated workflows for spiking neuronal network simulations. *Front.*
Neuroinformatics **7** (2013).

71. A. Sinha, R. de Schepper, J. Pronold, J. Mitchell, H. Mørk, P. Nagendra Babu, J. M. Eppler, M.
Lober, C. Linssen, D. Terhorst, M. A. Benelhedi, A. Morrison, W. Wybo, G. Trensch, R. Deepu,

N. Haug, A. Kurth, S. B. Vennemo, S. Graber, S. Spreizer, J. Gille, J. Vogelsang, M. Krüger,
H. E. Plesser, NEST 3.4, version 3.4, Zenodo (2023).

72. J. Antolík, A. P. Davison, Arkheia: Data Management and Communication for Open
Computational Neuroscience. *Front. Neuroinformatics* **12**, 6 (2018).

73. H. J. Alitto, W. M. Usrey, Influence of Contrast on Orientation and Temporal Frequency Tuning
in Ferret Primary Visual Cortex. *J. Neurophysiol.* **91**, 2797–2808 (2004).

74. Z. W. D. Lyle Muller, Generalized Phase, *GitHub repository* (2020).
https://github.com/mullerlab/generalized-phase.

1363

Data Availability

All simulation data used in this study can be freely generated by running the virtual experiments in the Github repository https://github.com/CSNG-MFF/mozaik-models/tree/main/Rozsa_Cagnol2024

Code Availability

The model, experiments and analyses were implemented through the Mozaik framework (*70*), freely accessible at https://github.com/CSNG-MFF/mozaik (version 0.4). To run the model one first needs to install Mozaik and its dependencies as indicated in the Mozaik Github repository, and then to download the implementation of model at https://github.com/CSNG-MFF/mozaik-models/tree/main/Rozsa_Cagnol2024 (version 1.0) and to follow the instructions to run the experiments presented in this study. We used NEST simulator (version 3.4) as the underlying simulation engine (*71*). We also made our model results and virtual experiment specifications available via a dedicated data store (http://v1model.arkheia.org) implemented in the Arkheia framework (*72*).

Contributions

J.A. conceived of and supervised the project. T.R. and R.C. ran computational simulations, along with analyses and visualisations. The three authors jointly wrote, reviewed and edited the paper.

Figure 1: Common neural substrate underlies a range of visually evoked, optogenetically evoked, and spontaneous phenomena (A) Schematic of the V1 model: thalamocortical connections to cortical Layer 4 from ON and OFF LGN cells are sampled based on a precomputed orientation preference map. All cortical neurons make local connections, while only Layer 2/3 neurons make long-range intra-layer horizontal connections. See Methods for the functionally specific rules of connectivity. **(B)** Hypothesis: the recurrent distance dependent, patchy, orientation-biassed horizontal connectivity in Layer 2/3 underpins spontaneous, visually and optogenetically evoked cortical responses. **(C)** Computational approach: the simulated responses of neurons of a single parameterisation of the V1 model under multiple stimulation paradigms (no stimulus, visual or optogenetic) are recorded and analysed according to a range of experimentally measured criteria and compared to experimental results (illustrative experimental data scraped from (*12*, *73*) or adapted from (*10*, *26*) with permissions obtained via Rightslink and under the GNU licence, respectively).

Figure 2: STW in the V1 model (A) Example frame sequence of the generated LFP signal of the model Layer 2/3 spontaneous activity showing propagation of a STW. A video of this example is available in Supplementary Video 1. **(B)** The peak times of the spike-triggered LFP (stLFP) for each spatial location plotted against their distance from one example reference neuron from layer 2/3. **(C)** The amplitudes of the stLFP for each spatial location in relation to their distance from one example reference neuron from layer 2/3, and the corresponding exponential fit (solid blue line). **(D)** Propagation speeds of the stLFP in cat V1 (black) from Nauhaus et al. (*8*), and for the model. Error bars represent the standard deviations. **(E)** The average of the stLFP residuals for layers 4 (pink) and 2/3 (red, solid: default configuration, dashed: configuration without functional bias in horizontal connections, blue: experimental data from Nauhaus et al. (*8*)) across reference neurons and recording sites, binned according to their orientation preference differences. Error bars: Standard errors of the mean. **(F)** Synthetic LFP signal (black) and the corresponding output of the generalised phase analysis (color-coded with the instantaneous phase), for a single recording site. **(G)** Distribution of the firing rates of the neurons in Layer 2/3 of the V1 model (Red) and analogous experimental data in marmoset MT (Davis et al. 2021 (15); Blue). **(H)** Same as **(G)**, for the coefficient of variation of the inter-spike-interval. **(I)** The average wavelengths for the model layers 4 (pink) and 2/3 (red) and for the experimental data of Davis et al. (15) (blue), as computed with the Generalized phase analysis. **(J)** Cumulative distributions of the wavelengths of the simulated LFP signal in the model Layer 2/3 before (orange) and after (red) its spatial convolution with a Gaussian. The cumulative distributions of the spatially shuffled signal is shown in yellow, and the cumulative distribution of the LFP signal recorded in marmoset MT by Davis et al. is shown in blue (15). **(K)** Propagation speeds distribution of the LFP signal in the model (red) and in 2 marmosets (turquoise: scraped from Extended Data 2d 'Monkey T' in Davis et al. 2020 (7); blue: data from Davis et al. 2021 (15)). **(L)** Same as **(I)**, for the average propagation speeds.

Figure 3: The structure of correlations in spontaneous activity is related to orientation maps (A) Example spontaneous activity events in the simulated visual cortex calcium imaging recordings. The colour border indicates the dominant orientation preference of activated neurons. **(B)** Orientation map with two iso-oriented points (1,2) and **(C)** correlation maps (CMs) of excitatory activity with seeds at those points. The CMs with iso-oriented seeds are similar (example CMs similarity=0.73). **(D)** CMs of inhibitory activity are similar to CMs from excitatory activity (example CMs similarity=0.57). **(E)** Similarity of excitatory and inhibitory CMs is high, regardless of seed position. **(F)** (left) Similarity between CMs at each seed point and the orientation map. Similarity is consistently high regardless of seed position. (right) Removing the orientation bias of lateral connections decreases similarity. **(G)** A circular SOM fitted to spontaneous excitatory CMs predicts the structure of the orientation map, as seen in Kenet et al. (*29*). **(H)** For a wide range of metrics recorded in the ferret cortex (*10*, *20*), excitatory and inhibitory activity in the model lies within experimentally recorded values. 'E', 'I', 'Or.' denote excitatory activity, inhibitory activity and orientation preference map, respectively. **(I)** Validation against Kenet et al. (*29*): The distribution of correlation coefficients between model spontaneous activity frames and single-condition orientation preference map is consistent with experimental data, including correlations to control (horizontally flipped) maps. **(J-L)** Validation against Tsodyks et al. (*31*): **(J)** The distribution of correlations between model spontaneous activity frames and Preferred Cortical States (PCS) (see Methods) closely matches experimental data **(K)** The distribution of correlations to PCS for spontaneous frames at which the monitored neuron fired an action potential exhibits a significant positive bias ($p<10^{-16}$, n=367169, 1-sided one-sample T-test), consistent with experimental data (mean = 0.1007 ± 0.0003 for model, 0.1591 ± 0.0045 for experiment). **(L)** The predicted instantaneous firing rate increases with PCS correlation for both model and experiment. It is calculated as the ratio of histograms (L) and (K) divided by the duration of the time frame. Experimental data scraped from Smith et al. (*10*) Supplementary Figure 5g, Mulholland et al. (*20*) Figure 4c,e,g,j, 5f, Kenet et al. (*29*) Figure 2, Tsodyks et al. (*31*) Figure 3d,e,f .

Figure 4: Full-field and patterned stimulation evokes structured model responses (A) Schematic of optogenetic stimulation of model layer 2/3 neurons using a 400×400 LED array (see Figure S8 for a detailed description). **(B)** Mean simulated population response to full-field optogenetic cortical stimulation matches experimental data (*14*). **(C)** CMs calculated from spontaneous activity (top) and response to optogenetic stimulation (bottom) at the same seed position are similar (example CMs similarity=0.64). **(D)** Spontaneous and optogenetic response CMs are highly correlated to each other, regardless of seed position. **(E)** The mean correlation between spontaneous activity and optogenetic responses matches experimental results. **(F)** The similarity of both spontaneous and optogenetic correlation maps to the orientation map decreases to chance level when we remove the orientation bias of long-range connections. **(G)** (left) Endogenous stimulation patterns based on CMs and (right) statistically matched surrogate patterns uncorrelated with CMs. **(H)** The model matches the experimentally measured (*12*) maximum-normalised mean population response to endogenous stimulation both within and outside the stimulation ROI (left). Stimulating with surrogate patterns evokes a weaker activation compared to the endogenous stimulation response (right). **(I)** Correlation between the stimulation pattern and

evoked network activity pattern is higher for endogenous stimuli than for surrogate stimuli, in line with ferret experiments (*13*). Removing the orientation bias from long-range connections decreases the endogenous stimulus-response correlation towards the level of surrogate stimulus responses. All three populations of correlations have significantly different means (endogenous: 0.7996 ± 0.004, surrogate: 0.7097 ± 0.006, endogenous no bias: 0.7465 ± 0.003; $p_{endo-surr}$ = 2 × 10^{-18}, $p_{endo-endo\_no\_bias}$= 5 × 10^{-16}, $p_{surr-endo\_no\_bias}$ = 3 × 10^{-6}, Welch's two sample T-test: two-sided, assuming unequal variances, n = 40 for all three populations). **(J)** Schematic of Chernov et al. (*33*) experiment: simultaneous visual stimulation with full-field square gratings and optogenetic stimulation of an orientation preference column (circle, r=100μm), **(K)** Spatial map of regions with a significant (two-tailed paired t-test, for dependent samples; n = 10, P < 0.05) preference for visual stimuli iso-oriented (iso-regions, blue) or ortho-oriented (ortho-regions, red) relative to the optogenetically stimulated cortical column, within 2mm from the stimulation site. **(L)** (left) Optogenetic stimulation significantly (two-sided Wilcoxon signed-rank test, p<0.006, n = 10) increases response in iso-regions (blue) and decreases response in ortho-regions (red, p<0.004, n = 10) (right). Intrinsic imaging data from Chernov et al. (*33*) show a qualitatively similar effect. Experimental data scraped from: Mulholland et al. (*14*) Figure 2c for (B), Supplementary Figure S10c for (E); Mulholland et al. (*12*) Figure 5k for (H), Chernov et al. Figure 4l for (L).

Figure 5: Sharper orientation tuning of optogenetic stimulation responses compared to stimulation patterns (A) Example stimulation patterns (endogenous and surrogate) and corresponding responses at the end of the 1s stimulation period shown for raw spiking data (bottom) and simulated calcium imaging signal (top). **(B)** Trial mean of stimulation responses within and outside the stimulated areas, showing the spatiotemporal smoothing effect of the calcium imaging readout. **(C)** The temporal autocorrelation functions of ferret calcium imaging spontaneous activity signal (*34*) (black dashed, see Methods) and model simulated calcium imaging spontaneous activity signal (blue dashed) are of similar timescale. In addition, temporally deconvolved ferret calcium imaging activity (gray) using the Prior Frame Subtraction (PFS) technique is of a similar timescale to model simulated calcium imaging signal where we omitted convolving with the temporal kernel (dark blue). The autocorrelation functions of raw spiking activity and PFS-deconvolved calcium imaging signal (light blue) are close to identical. **(D)** The mean DAOD over the stimulation period and trials is more sharply oriented for both endogenous (red) and surrogate (pink) stimuli, than the distribution of orientation preferences pooled from the orientation map in the stimulated area. The plot shows the DAOD for a single example endogenous/surrogate pattern, see all patterns on Figure S10. **(E)** Entropy of mean DAOD is consistently lower than the distribution of orientation preferences entropy across all stimulation patterns, showing that the activity due to stimulation is more sharply oriented than the stimulated area itself. **(F)** Entropy of DAOD is closely linked with firing rate - activity snapshots with higher activity exhibit sharper activations for both endogenous/surrogate stimulus responses and spontaneous activity (Pearson correlation $r_{endogenous}$=-0.55, $r_{surrogate}$=-0.20, $r_{spontaneous}$=-0.27). The points represent DAOD histograms pooled across all stimuli, trials and stimulation time points. Experimental data scraped from: Kettlewell et al. (*34*) Figure 1a,b for (C).

Figure 6: Central stimulation produces functionally specific nonlinear effects in the cortical surround (A) Localised optogenetic stimulation - we stimulate in a small circle with radii increasing from 50μm to 300μm in the middle of the simulated cortical region, targeting a cortical orientation column. **(B)** (left) Stimulation responses beyond the stimulated area. The central stimulated orientation column has horizontal orientation preference (red colour code). The activity in the surrounding cortex is biassed towards iso-oriented areas (red). (right) DAOD histograms for excitatory and inhibitory neurons in the non-stimulated cortex for the depicted frame (with entropies 2.9, 3.3). Activity of the excitatory population is much more sharply tuned than that of the inhibitory population. **(C)** The range of orientation preferences under stimulation ROIs increases with the increasing stimulation radius. **(D)** Excitatory firing rate rises to a maximum and subsequently begins falling, while **(E)** inhibitory firing rate in the surround scales monotonically with increasing radius. Both excitatory and inhibitory DAOD entropy inversely follows excitatory activation. **(F)** At the stimulation radius maximally exciting the surround, in iso-oriented regions both excitatory (red, solid) and inhibitory (blue, solid) neurons show increased firing rate compared to spontaneous, while in ortho-oriented regions both excitatory (red, dashed) and inhibitory (blue, dashed) neurons exhibit decreased firing rates.

Figure 7: Impact of the parameters on model properties. (A) The distribution of propagation speeds for six parametrizations varying in TC strength. Red font in legends indicates the parameter values used in the base model. **(B)** Same, for six parametrizations varying in intra-laminar distance-dependent conduction delays. **(C)** For six parametrizations varying in TC strength: (blue) stLFP residuals difference between recording sites with orientation preference congruent to the reference electrode and sites with orthogonal preference; (orange) similarity between the excitatory CMs and the orientation map (solid), and between the uniform full-field optogenetically evoked CMs and the orientation map (dashed). The red circles indicate the parameter values used in the base model. **(D)** Same, for six parametrizations varying in intra-laminar distance-dependent conduction delays. **(E)** Same, for five parametrizations varying in L2/3 E→I strength. **(F)** Same, for five parametrizations varying in L2/3 E→E bias.

	α				θ			
	L4 Exc	L4 Inh	L23 Exc	L23 Inh	L4 Exc	L4 Inh	L23 Exc	L23 Inh
L4 Exc	0.0139	0.0148	0.0174	0.0197	207.7	191.8	154.4	131.5
L4 Inh	0.0126	0.0119			237.5	256.4		
L23 Inh			0.0149	0.0150			189.5	188.61

Table 1: Parameters of the hyperbolic distributions used to generate the local connectivity of the model. Rows refer to pre-synaptic populations, and columns to post-synaptic populations.

Delays		L2/3 E→E bias		L2/3 E→I strength		Cortical density	
Delay (s/m)	TC Strength (nS)	Bias (rad^{-1})	TC Strength (nS)	Strength (nS)	TC Strength (nS)	Density (cells mm^{-2})	TC Strength (nS)
0.2	2.2	0.77	2.4	0.35	2.2	500	0.7
0.225	2.1	1	2.8	0.375	2.5	1500	1.6
0.25	2	1.25	2.36	0.4	2.7	3000	2
0.275	1.9	1.67	2.3	0.425	2.9	5000	2.2
0.3	1.75	2.22	2.2	0.45	3.05	7000	2.3
0.333	1.6					10000	2.3

Table 2: Pairing of values of TC strength with other parameters modifications. Cells with red background correspond to the parameters values used for the baseline version of the model.

A Model V1

Layer 2/3
Layer 4
Precomputed orientation map
Sampled thalamocortical connections
orientation color key

LGN ON
LGN OFF

Sample
Sample
Sample

Afferent receptive field template

OFF RF
11° Visual field
ON RF

B Processing in V1

Lateral connectivity properties

Distance dependence
Patchiness
Functional bias

Connection probability
Distance (mm)

Connection probability
0

Connection probability
Orientation pref. difference (°)

No input

LED

Recurrent network

Network activity

Visually evoked

Spontaneous

Optogenetically evoked

C

Simulation   Analysis   Comparison

Stimulus   Response   Experiment   Model

Visual

Orientation tuning

Firing rate (sp/s) 70
Orientation
Firing rate (sp/s) 35
Orientation

Model

No input

Modular activity

Z-score 3 / -3
1 mm
Z-score 3 / -3
1 mm

Generalized phase

50 ms
50 ms

Optogenetic

Endogenous stim. patterns

1 s   Within ROI   Outside ROI
1 s   Within ROI   Outside ROI

Time

A

Normalized spont. activity

Time (s)

B Orientation

π

C Exc

Correlation

-1

D Inh

E Exc - Inh corr. maps

Similarity

-1

F Exc corr. maps - or.map

Or. bias   No bias

Similarity

G Estimated orientation

π

H

Similarity

E → Or.   I → E

Dimensionality

E   I

Correlation at maxima (>2 mm)

E   I

Local correlation eccentricity

E   I

Correlation wavelength (mm)

E   I

○ Experiment   ● Exp. mean   — 95% Prediction Interval (PI)   ● Model

I Correlation of spont. activity to oriented grating responses

Density

Correlation coefficient

Model
Grating
Control

Kenet et al.
Grating
Control

J Correlation of PCS to spont. frames

Density

Correlation coefficient

K Correlation of PCS (spike time spont. frames)

Density

Correlation coefficient

L

Predicted firing rate (sp/s)

Correlation coefficient

■ Tsodyks et al. 1999   ■ Model

A

LED LED LED

Layer 2/3

Layer 4

B

Light ON
Experiment
Model

Population mean ΔF/F

1.5

0.0

0   1   2   3   4
Time (s)

C

Spontaneous

Correlation
1
-1

②

Optogenetic

②

D

Spont-opto

Similarity
1
-1

E

○ Exp. trial
● Exp. mean
— 95% PI
● Model

Spontaneous - opto corr. map similarity

Opto

F

◆ Opto
✕ Spont
▢ Chance 95% PI

Or.map similarity

Model   Model no bias

G

Endogenous   Surrogate

Orientation
π
0

Normalized response
1
0

Stim. ROI

H

Endogenous   Surrogate

Normalized response

0 1 2 3 4   0 1 2 3 4
Time (s)   Time (s)

Experiment   Model
Within ROI   Within ROI
Outside ROI   Outside ROI

I

Stimulus-response correlation

0.8

0.5

Endog   Surr   Endog No bias

▢ Trial   ● Mean
— 95% Confidence Interval

J

visual stimuli

optogenetic stimulator

r

OP map

0.25 mm

K

Visual only

FR diff (sp/s)
30
-30

L

Model

Firing rate (sp/s)

**   **

Visual   Visual+ opto

Experiment

■ Iso ■ Ortho

dR / R (%)

0.15

Visual   Visual+ opto

A

stimulator

r

0.5 mm

B Model response in the surround

π

Orientation preference

0   1
Activity

Stimulus
Margin

Excitatory
Inhibitory

Firing rate (sp/s)

−π/2   0   π/2
Orientation preference

C

π/3

Stimulated orientation std (°)

0   100   200   300
Radius (um)

D

3.39

Exc DAOD entropy

3.25

3.30

Exc Firing rate (sp/s)

3.10

0   100   200   300
Radius (um)

E

3.40

Inh DAOD entropy

3.37

7.00

Inh Firing rate (sp/s)

5.00

0   100   200   300
Radius (um)

F

Mean Firing rate (sp/s)

Iso surround
Ortho surround
Spontaneous

0   100   200   300
Radius (um)