Mechanistic pathways of tick exposure risk in native and invaded plant communities

Drew Hiatt[1*], Whalen W. Dillon[1], Allison Gardner[2], Steven Cabrera[3], Michael Dietze[4], Brian F. Allan[5], and S. Luke Flory[1]

[*]Corresponding author dhiatt@ufl.edu ph:(775) 790-4303
[1]Agronomy Department, University of Florida, Gainesville, FL
[2]School of Biology and Ecology, University of Maine, Orono ME
[3]College of Forestry, Wildlife and Environment, Auburn University, Auburn, AL
[4]Department of Earth and Environment, Boston University, Boston, MA
[5]Department of Entomology, University of Illinois Urbana-Champaign, Urbana, IL

Running title: < Plant invasion effects on ticks >

Keywords: *Amblyomma americanum, Imperata cylindrica*, tick survival, disease ecology, invasion, indirect effects

Statement of authorship:
DH, WD, BA, MD, and SLF conceived the idea and designed the experiment. DH, WD, and SC carried out the experiment. DH, WD, SC, and AG performed the analyses. DH wrote the first draft of the manuscript and all authors contributed substantially to revisions.

Data accessibility (Open Research Statement):
All data and code supporting these results has been archived via data dryad.
https://doi.org/10.5061/dryad.mpg4f4r2k

Section length:
Abstract: 238/ 350 words
Manuscript length: 31 pages
Number of references: 62
Number of figures: 4
Number of tables/ textboxes: 0

Drew Hiatt   ORCiD: 0000-0001-9514-9585
Whalen Dillon   ORCiD: 0000-0001-8171-0407
Allison Gardner   ORCiD: 0000-0001-8438-0634
Steven Cabrera   ORCiD: 0000-0002-4637-7458
Michael Dietze   ORCiD: 0000-0002-2324-2518
Brian F. Allan   ORCiD: 0000-0002-1155-3271
S. Luke Flory   ORCiD: 0000-0003-3336-8613

Abstract

Plant invasions may alter vector abundance by several mechanistic pathways, including modifying microclimates that influence vector survival or changing habitats to influence host use. Here, we used a field experiment and observational data to evaluate multiple mechanistic pathways (tick survival and host abundance) by which plant invasions may alter vector-borne disease risk using the common disease vector lone star tick (*Amblyomma americanum*) and the widespread invasive cogongrass (*Imperata cylindrica*) in the southeastern USA. In the field experiment, ticks survived over 50% longer in areas dominated by the invasive plant compared to those with only native plant species. Invaded areas had lower temperatures and higher humidity that likely reduced tick desiccation. The observational study showed greater average tick abundance in native plant communities and no difference in wildlife host (white-tailed deer) activity between plant communities. There was also a positive relationship between tick abundance and white-tailed deer activity, but only in native areas. Together, these results suggest that more favorable microclimate conditions resulting in greater tick longevity is the dominant driver of tick abundance in invaded areas, while tick abundance in native-dominated areas is promoted, at least in part, by white-tailed deer activity. Our results demonstrate that plant invaders affect multiple, potentially off-setting mechanistic pathways contributing to tick exposure risk. The complexity of these relationships highlights the need for better understanding of how invasive species and other global change drivers influence abundance of disease vectors and, ultimately, disease transmission.

Introduction

Introduction

Human-mediated introductions of species to non-native ranges have resulted in widespread

biological invasions, which are a key driver of global environmental change (Vitousek et al.

1996). Accumulating evidence demonstrates that invasive species can have direct effects on

ecological communities by altering, for example, biodiversity, nutrient cycling, and disturbance

regimes (Vilà et al. 2011, Pyšek et al. 2012). However, they can also elicit indirect effects that

may be larger than their direct effects, such as through changes to habitat structure and

microclimate (Linders et al. 2019). These indirect effects can have cascading impacts on humans

by affecting food production, recreation, and climate mitigation (Pejchar and Mooney 2009).

Invasive species may also indirectly influence human health if they alter interactions among

species that transmit diseases to humans (Rai and Singh 2020), however, to our knowledge such

effects have been rarely explored.

Infectious pathogens transmitted from animals to humans via arthropod vectors such as

mosquitoes and ticks place a major burden on global health (Morens et al. 2004, Jones et al.

2008). Tick-borne disease transmission to humans results from complex species interactions that

are mediated by impacts of plant communities on tick growth, survival, dispersal, questing, and

reproduction (Ostfeld et al. 2018, Morand and Lajaunie 2021) and host habitat conditions that

affect the frequency and duration of host use (Elias et al. 2006, Allan et al. 2010a, Guiden and

Orrock 2019). Plant invasions can alter multiple mechanistic pathways that affect tick abundance

and infectious disease exposure for humans by either modifying host habitats that alter host

abundance or by changing microclimate conditions that promote or inhibit tick survival (Allan et

al. 2010a, Williams and Ward 2010, Mackay et al. 2016, Stewart et al. 2021) (Figure 1).

Entomological metrics of exposure risk include metrics of tick abundance (e.g. density of ticks),

pathogen infection prevalence, or their product (e.g. density of infected ticks) (Fischhoff et al.

2019). Vector abundance often is used as a proxy for tick exposure risk, or a human's likelihood

of encountering a tick upon entering a particular habitat. While there has been research on how

these pathways individually affect tick abundance, studies that simultaneously address multiple

pathways are needed to improve understanding of complex processes by which invasive plants

may alter tick exposure risk, and thus infectious disease.

Collectively, evidence for mechanisms that influence tick-borne disease risk in invaded

areas is sparse and inconsistent. The limited evidence for invasive plant effects on tick-borne

disease varies by tick species, invasive plant species, and habitat conditions, among other factors

(Civitello et al. 2008, Allan et al. 2010a, Williams and Ward 2010, Mack and Smith 2011,

Racelis et al. 2012, Adalsteinsson et al. 2016). Plant invaders have been shown to increase host

habitat use (Guiden and Orrock 2019), leading to greater abundance of ticks (Elias et al. 2006)

and pathogen-infected ticks (Allan et al. 2010a) compared to native areas. Invasive plants may

also increase tick abundances relative to native plant dominated areas by providing favorable

microclimate conditions (i.e., higher humidity and/or lower temperature) that reduce tick

desiccation (Williams and Ward 2010, Racelis et al. 2012). However, the few studies comparing

the abiotic mechanisms affecting tick survival in invaded and native areas show either no

difference (Allan et al. 2010a) or lower tick survival (Civitello et al. 2008) in invaded compared

to native plant dominated communities. Given the ongoing introduction and spread of both

invasive plants (Seebens et al. 2017), and ticks (Ostfeld and Brunner 2015), there is a critical

need to better understand how invasive plants affect host habitat use (Stewart et al. 2021) as well

as the abiotic factors that influence tick survival.

Here, we measured effects of plant invasion on tick survival and host activity to

determine if these pathways were predictive of tick abundance. Specifically, we combined field

experiments and observational field studies to evaluate how an invasive plant (cogongrass;

Imperata cylindrica) affected microclimate conditions and tick survival (lone star tick;

Amblyomma americanum) compared to native species dominated areas, and how differences in

relative host activity (i.e., white-tailed deer, *Odocoileus virginianus*) between invaded and native

areas were related to tick abundance (Figure 1). We measured temperature, relative humidity

(RH), and survival of lab-reared ticks in invaded and nearby native communities in a field

experiment over 11 months and we evaluated relative white-tailed deer activity (via camera

traps) and abundances of host-seeking ticks (via CO_2 trapping) in paired invaded and native sites

in a field survey. Because the invader creates much denser stands compared to native plant

communities, we hypothesized that invaded areas would have lower temperatures and higher

relative humidity, which would be favorable for tick survival relative to native areas. We also

hypothesized that relative white-tailed deer use would be greater in invaded communities

because dense cover created by the invader would provide perceived protection from predators,

as documented with small mammals in cogongrass (Borden et al. 2022), which would in turn

result in greater tick abundance.

Methods

128   *Study system*

129   One of the most widely distributed invasive plant species in the southeastern USA is

130 *Imperata cylindrica,* (L.) Beauv. (cogongrass; hereafter *Imperata*), (Estrada and Flory 2015) and

131 it is one of the top five costliest terrestrial invasive plants to manage in Florida (Hiatt et al.

2019). *Imperata* can invade a wide variety of habitats due in part to its greater trait plasticity

(Hiatt and Flory 2020), and drought tolerance (Fahey et al. 2018) compared to co-occurring

native plant species. *Imperata* invasions can alter the structure of southern pine forests by

producing greater fire intensity compared to native plant communities (Dillon et al. 2021)

leading to fire induced tree mortality (Flory et al. 2022). *Imperata* has also been shown to alter

microclimate conditions, including higher soil moisture and relative humidity, and lower

temperatures at ground level compared to native plant communities due to its higher stem density

and greater leaf area (Alba et al. 2017, Cabrera et al. 2021).

Lone star ticks (*Amblyomma americanum*) are hard ticks in the family Ixodidae (Keirans

and Litwak 1989) that can transmit pathogens, including the causative agents of human

ehrlichiosis, Rocky Mountain spotted fever, and tularemia (Childs and Paddock 2003). Time

between blood meals (off-host), either seeking hosts to parasitize, digesting blood meals, or

dormant, comprises over 90% of an Ixodid tick's life cycle, making the primary cause of off-host

tick mortality desiccation (i.e., loss of water through the cuticle) (Needham and Teel 1991).

Maintaining a constant water balance is one of the most important processes that influence off-

host survival and, ultimately, tick population size (Knülle and Rudolph 1982, Needham and Teel

1991, Perret et al. 2000, Maldonado-Ruiz et al. 2020). Ticks will begin to desiccate when they

are below their critical equilibrium humidity (CEH), which is 80-82% relative humidity for adult

A. americanum (Hair et al. 1975, Knülle and Rudolph 1982). While desiccation is primarily

driven by low relative humidity (at or below CEH), increased temperature leads to an increase in

water molecules escaping the cuticle due to kinetic energy (Sauer and Hair 1986). For *A.*

americanum temperatures in excess of 35 ºC can contribute to desiccation and mortality (Sauer

and Hair 1971), while all terrestrial arthropods, including ticks lose their capacity to retain

155   moisture at temperatures in excess of 40 ºC (Londt and Whitehead 1972). While tick abundance

156   is influenced by microclimate, local abundance of *A. americanum* can also be regulated by host

157   activity. White-tailed deer (*Odocoileus virginianus*) serve as key hosts for all three life stages of

158   *A. americanum* (Bloemer et al. 1986), and are important reservoir hosts for multiple pathogens

159   transmitted by *A. americanum* (Allan et al. 2010b).

160   *Field experiment: Tick survival and microclimate conditions*

161   To examine the effect of *Imperata* invasion on tick survival, we established an

experiment in a planted longleaf pine (*Pinus palustris*) stand with fine sandy soils near La

Crosse, FL (29.897° N, -82.420° W) that had distinct *Imperata*-invaded and -uninvaded areas

(Appendix S1: Figure S1, S2). The invaded area was highly dominated by *Imperata* and spanned

approximately 6300 m^2, while the surrounding area was uninvaded, native understory. The plant

community of the native area was typical of a longleaf pine stand, with *Aristida stricta*

(wiregrass), *Rubus flagellaris* (common dewberry), *Carya sp.* (hickory), and *Andropogon* spp.

There was no recent chemical or mechanical management in either the invaded or native areas,

and the native understory included no other non-native plant species. Because we conducted our

experiment within a forest stand with distinct invaded (*Imperata*) and native areas in close

proximity that shared the same overstory structure, we expected any differences in microclimate

conditions to be due to invasion, not other underlying factors (see Results).

173   We established 24 plots (1m x 1m), 12 in invaded habitat and 12 in native habitat, with at

least 10m between plots. Plots were established every 10m along two transects, one in each plant

community; plots were then randomly placed along each transect by in areas that were

representative of each plant community. At the center of each plot, a 300-micron 10 cm x 20 cm

nylon mesh bag (Skimz, Singapore), containing 10 nymph and 10 adult (5 female and 5 male) *A.*

americanum obtained from the National Tick Research and Educational Resource (Department

of Entomology and Plant Pathology, Oklahoma State University, Stillwater, OK) were placed in

50 cm tall cylindrical wire mesh exclosures to deter wildlife interference. To control for age of

ticks (within life stage) and standardize the prior range of exposure to environmental conditions,

we used lab-reared instead of wild-caught ticks. Each mesh bag containing ticks was oriented

vertically to facilitate natural diurnal tick questing (i.e., host seeking behavior) and, although

there was no substrate within the bags, the bags remained in contact with the litter layer to allow

for ticks to seek refuge below the leaf litter to avoid desiccation stress (Schulze and Jordan 2003)

throughout the experiment (Appendix S1: Figure S1).

The start date of the survival experiment (June 21, 2018) coincided with typical off host

periods (between blood meals) for *A. americanum* in the region (Davidson et al. 1994, Gleim et

al. 2014). During these long non-parasitic phases, ticks seek refuge in areas with microclimates

that promote constant water balance (Knülle and Rudolph 1982, Davidson et al. 1994). Mesh

bags were examined frequently (every 7-10 days) for the first month and then every 10-14 days

afterward to determine days survived for both nymph and adult ticks. During each visit, bags

were removed from the cages but not opened, and survival status of each individual tick was

determined by assessing the response of the tick to CO_2 stimulation from human breath (Steullet

and Guerin 1992). Visits continued to the site until May 23, 2019 when all ticks had died.

To assess plant community structure and abiotic conditions in invaded and native areas,

at the start of the survival experiment we collected understory herbaceous plant height and

standing biomass, light availability at the soil surface, and overstory canopy cover at each plot.

Plant height was measured as the tallest point of living leaf tissue from soil surface. Standing

biomass was collected in 25 cm x 25 cm quadrats and dried at 60 ºC to a constant mass and

weighed (± 0.01g). Photosynthetically active radiation (PAR, i.e., light) was measured at ground

level in each plot using an ACCUPAR LP-80 PAR/LAI ceptometer (Decagon Device Inc.,

Pullman, WA, USA). Overstory canopy cover was estimated using a concave spherical

densiometer (Forestry Suppliers Inc., Jackson, MS, USA) by the average of four readings facing

each cardinal direction.

Throughout the experiment we recorded ground-level microclimate conditions

(temperature and relative humidity) at five-minute intervals immediately adjacent to each

enclosure containing the mesh bags using HOBO U23 Pro v2 loggers (Onset Computer

Corporation, Bourne, MA). Each logger was housed in a capped 18-inch length of 3.8 cm

diameter PVC pipe to protect it from rainfall and direct sunlight. Holes were drilled on four sides

six inches apart along the length of the pipe to allow for airflow and better reflect ambient

temperature and relative humidity (see Appendix S1: Figure S1a-b).

Statistical analysis: Site conditions

We conducted our statistical analysis of the tick survival field experiment in several

phases. First, we conducted two analyses to determine how *Imperata* affected plant community

characteristics that might influence microclimate conditions and if *Imperata* invasion altered tick

survival over time. We used multivariate analysis of variance (MANOVA) to test the hypothesis

that *Imperata* changed plant community characteristics (i.e., vegetation height, standing biomass,

and light availability at soil surface). Analysis of variance (ANOVA) was used to test if

overstory canopy cover was similar in invaded and uninvaded areas (i.e., differences in

understory plant community characteristics were due to invasion, not the tree canopy or other

environmental factors). We also fit polynomial non-linear ordinary least squares regressions to

test the hypothesis that *Imperata* invasions changed microclimate conditions (i.e., higher average

daily minimum relative humidity and lower average daily maximum temperature) during the

experiment. These analyses were done using the "car" package (Fox and Weisberg 2018, Team

2019) in R version 3.6.1, R Core (Fox and Weisberg 2018, Team 2019)).

Statistical analysis: Tick survival and microclimate conditions

We used linear regression models to assess how the number of days survived for

individual *A. americanum* of each life stage was related to the plant community and each

microclimate variable (hours per day under 80% relative humidity and daily maximum

temperature). The first set of mixed effects regression models on number of days survived for

adult or nymph life stage ticks included the fixed effects of treatment (native), hours per day

under 80% relative humidity (total hrs. below 80% relative humidity/ days survived), and their

interaction. The second set of mixed effects regression models for days survived for nymph or

adult life stage ticks included fixed effects of treatment (native), maximum temperature

experienced at mortality, and their interaction. Maximum temperature at mortality was recorded

as the maximum temperature measured at each plot between survival checks (i.e., maximum

temperature between the time when the tick was last observed alive and when it was observed as

dead). We included plot as a random effect in both models to account for repeated measures.

Models were fit for nymph and adult life stage ticks separately using the "lme4" package (Bates

et al. 2014), and degrees of freedom were calculated using "lmertest" (Kuznetsova et al. 2017) in

R 3.6.1 (R Core Team 2019). To further elucidate the effect of both relative humidity and

temperature on tick survival, we also used ordinary least squares linear regression to further

investigate the mechanistic effect of each respective microclimate variable on nymph and adult

survival time (days) (Appendix S1: Table S4). These models provide estimates of effect sizes of

maximum temperature or relative humidity on survival, irrespective of plant community. For

these models, we scaled the microclimate variables using a z-score standardization so that the

effect of each variable could be compared on the same axis.

Field survey: White-tailed deer activity, tick abundance, and microclimate

To examine the impact of *Imperata* invasions on white-tailed deer activity and tick

abundance, we conducted field surveys in invaded and native dominated plant communities

across eight sites in north central Florida (Appendix S1: Figure S2). Invaded plots were selected

if the *Imperata* invasion covered a minimum of 500 m^2, native plots were selected if they had

similar overstory tree canopy structure to that of the invaded plots. Since we encountered many

large invasions with few comparable native plant communities, at some sites we sampled

multiple invaded plots per site. We sampled a total of 34 plots, 22 plots in *Imperata* invaded

areas and 12 in native plant dominated areas. Due to limited supply of camera traps, a sub-set of

the plots were sampled for white-tailed deer activity (28 plots total, 17 in *Imperata* invaded areas

and 11 in native plant dominated areas, Appendix S1: Table S5). Plots were also sampled for tick

abundance and some plots were sampled multiple times, yielding 51 tick sampling events, 32 in

Imperata invaded areas and 19 in native plant communities (Appendix S1: Table S6). Within a

site, invaded and native plots were located at least 20 m away from each other. *Imperata*

invasions varied in size across sites but covered a minimum of 500 m^2 to more than 1 ha. We

established plots (10 m radius) under similar overstory tree species and canopy cover between

invaded and native plots, so the primary difference in habitats was presence or absence of the

invader.

To estimate relative white-tailed deer use of invaded and native areas, we placed two

motion-activated cameras (Primos 12MP Proof Cam 02 HD, Primos Hunting, Flora, MS) in each

plot to monitor wildlife activity. All cameras were set to take a burst of three photos on a 30

second delay, meaning that after the cameras were motion activated there would be a 30 second

delay before the cameras were able to be activated again. We made no effort to correct for

particular individuals as we were interested in measuring white-tailed deer activity per day rather

than number of unique hosts. Cameras were placed simultaneously at paired native and invaded

plots so that deer activity could be calculated across the same time frame. At the end of the

experiment photos were examined by three individuals for presence of white-tailed deer

(Appendix S1: Table S5). Cameras were positioned on opposite sides of the plot at a distance

between 10 m and 15 m from the center of the plot and approximately 1.5 m aboveground.

Cameras were positioned so that their fields of view did not overlap, maximizing the total area of

the plot being sampled. Cameras were active from 11 April to 1 June 2019, which coincided with

questing activity for nymph and adult *A. americanum* in the region (Davidson et al. 1994, Gleim

et al. 2014). In each field site we also recorded ground-level microclimate conditions

(temperature and relative humidity) from 13 June to 1 August 2019 at five-minute intervals using

HOBO U23 Pro v2 logger and PVC housing as described in the field experiment above.

We concurrently estimated tick abundance for each plot using carbon dioxide-baited traps

(hereafter CO_2 trap). The CO_2 trap is one of the most effective means of sampling *A. americanum*

nymph and adult life stage ticks (Schulze et al. 1997) and is less biased by different vegetation

structures (e.g., a plant invasion) than drag sampling (Kensinger and Allan 2011). The CO_2 traps

consisted of a mini cooler mounted on a sheet of plywood (45 cm x 60 cm) with vent holes

drilled along the sides of the cooler near the base to disperse gaseous CO_2 from sublimating dry

ice. For each sampling event, double-sided carpet tape was placed around the perimeter of the

plywood to capture ticks attracted by the CO_2. Each trap was filled with approximately 2 kg of

dry ice and left in the field for 24 hours before ticks were collected. We placed four CO_2 traps

per plot, and traps were placed at least 10m apart from each other within a plot. CO_2 traps can

attract ticks that are within a 5m radius of the trap (Kensinger and Allan 2011). When plots were

resampled, traps were placed in new locations that were greater than 5m from the previous

sampling location. This sampling approach maximized the area of the 10m radius plot that could

be sampled with CO_2 traps (Appendix S1: Figure S3). In total there were 51 tick sampling

events, and with four CO_2 traps per sample, there were over 200 trap nights in the study. Traps

were purposefully not placed in plot centers to ensure that new areas of the plot were examined

during each sampling event. Ticks ensnared on the tape were counted and preserved in 70%

ethanol for identification and to determine life stage.

302   *Statistical analysis: Tick abundance and white-tailed deer activity*

303   We focused our analysis on white-tailed deer activity only due to the tall vegetation in

invaded areas limiting detection of smaller wildlife hosts with camera traps (Appendix S1:

Figure S4). White-tailed deer serve as an important host for all three life stages of *A.*

americanum (Bloemer et al. 1986). Relative white-tailed deer activity was estimated and

standardized by dividing the number of camera trap photos containing deer by the number of

days the camera was deployed and operating. This approach provided data in units of deer

photos/day and allowed for comparison of data from cameras with different lengths of operable

deployment. Our approach was intended to detect relative differences in activity by white-tailed

deer in invaded and native areas, not to estimate white-tailed deer population sizes.

312   We fit negative binomial general linear mixed effects model to examine the effect of

white-tailed deer activity on tick abundance from 11 April to 1 June 2019 in native and invaded

plant communities. We analyzed the fixed effects of plant invasion status (native vs. invaded),

white-tailed deer activity, and their interaction, on tick abundance (nymph + adult ticks). We

included site as a random effect in the model to account for repeated measures. White-tailed deer

activity was log-transformed to meet assumptions of normality. Models were fit using the "lme4"

package in R 3.6.1 (Bates et al. 2014, Team 2019). Negative binomial regression was also used

to test for differences in relative host activity and in tick abundance between native and invaded

plant communities.

Results

Field experiment: Site conditions

Average plant height, plant biomass, and light availability at the soil surface were

influenced by *Imperata* (MANOVA; $F_{3,20} = 74.48$; $p < 0.0001$). Average plant height was 268%

higher and plant biomass was 408% greater in invaded than native areas (Appendix S1: Figure

S5a, b), which corresponded to 91% less light availability at the soil surface in invaded areas

(Appendix S1: Figure S5c). There was no difference in overstory canopy cover in invaded and

native areas (ANOVA; $F_{1,22} = 0.146$; $p = 0.706$), indicating differences in plant communities and

understory light availability was driven by presence of *Imperata* (Appendix S1: Figure S5d).

Field experiment: Tick survival and microclimate conditions

Regardless of sex or life stage, ticks survived longer in invaded compared to native plant

communities (Figure 2 a, b). Time to 100% mortality per plot was, on average, 57% longer for

nymphs and 45% longer for adults in the invaded compared to uninvaded native plots (Figure 2

a, b). Throughout the experiment, there was greater relative humidity as well as lower maximum

temperatures in invaded areas (Figure 2 c, d). Average daily minimum relative humidity in

invaded areas was 79% at the start of the experiment (Appendix S1: Table S1, Intercept) and was

16% greater than native areas throughout the experiment (Appendix S1: Table S1, Invasion

effect). Average daily maximum temperature in the invaded area was 39 ºC at the start of the

experiment (Appendix S1: Table S1, Intercept) and remained roughly 6 ºC cooler compared to

native areas throughout the experiment (Figure 2 c, d; Appendix S1: TableS1, Treatment effect).

Across both life stages, individual tick survival was longer in invaded areas. Mean ± SE

nymph survival was 130 ± 5.4 days in invaded areas versus 80.5 ± 5.0 days in native vegetation,

while mean adult survival was 256 ± 4.8 and 140 ± 7.3 days in invaded and native areas,

respectively. For both nymph and adult life stages, exposure time (hours per day) to relative

humidity below 80% was significantly related to fewer days survived in native plant

communities (Figure 3a, b) (Appendix S1: Table S2). In invaded areas there was less exposure

time to relative humidity below 80% (1.9 h/day invaded vs 3.1 h/day native), and relative

humidity was not related to number of days survived for nymph or adult ticks. Maximum

temperature at mortality (i.e., maximum temperature between the time when the tick was last

observed alive and when it was observed as dead) was negatively correlated with survival time

for nymphal ticks, regardless of invasion (Figure 3c) (Appendix S1: Table S3). For adult ticks,

maximum temperature prior to mortality was negatively correlated with survival time for ticks in

native but not invaded plots (Figure 3d) (Appendix S1: Table S3).

Both microclimate variables had a negative effect on days survived, but maximum

temperature prior to observed mortality had a stronger effect than hours below the CEH

threshold for both life stages (Appendix S1: Table S4, Figure S6). For nymphs, maximum

temperature had a strong negative effect size on days survived (estimate = -46.9; 95% CI = -55.0

$-$ -38.6; $p < 0.0001$), while hours below 80% relative humidity had a smaller, nonsignificant

effect size (estimate = -5.0; 95% CI = -14.0 $-$ 4.1; $p < 0.254$). For adult ticks, maximum

temperature also had a strong negative effect size (estimate = -28.2; 95% CI = -38.3 $-$ -19.6; $p <$

0.0001), and hours below 80% relative humidity had a smaller but still negative effect (estimate

= -14.6; 95% CI = -29.2 – 0.0; p = 0.052) on survival.

364   *Field survey: White-tailed deer activity, tick abundance, and microclimate*

365   There were no differences in relative white-tailed deer activity between native and

invaded plots ($t_{1,44}$ = 1.01; p = 0.318) (Appendix S1: Figure S7). There were also no differences

in average tick abundance (nymph [$t_{1,49}$ = 0.905; p = 0.370]; adult [$t_{1,49}$ = 1.298; p = 0.200] life

stage ticks) between native and invaded areas (Appendix S1: Figure S8). In native communities,

white-tailed deer activity (deer photos/day) was positively correlated with tick abundance

(estimate = 1.4; 95% CI = 0.1 – 2.8; p = 0.033) (Figure 4, Appendix S1: Table S7), while in

invaded habitats, tick abundance was not related to white-tailed deer activity (estimate = -0.6;

95% CI = -1.8 – 0.5; p = 0.273). For plots where no white-tailed deer were detected (i.e.,

intercept term), invaded plant communities had positive tick abundances (estimate = 1.3; 95% CI

= 0.6 – 2.8; p = 0.027) (Figure 4, Appendix S1: Table S7). There was one plot with both high

tick abundance and high deer activity that could have strongly influenced the results, so we ran

the analysis using the same linear mixed effects model but with that data point removed.

Omitting that one data point resulted in no correlation between white-tailed deer activity and tick

abundance (estimate = 0.8; 95% CI = -0.8 – 2.6; p = 0.349) (Appendix S1: Table S8), but there

was a significant intercept term (estimate = 1.3; 95% CI = -0.1 – 2.7; p = 0.036) (Appendix

S1:Table S8), indicating that tick abundance was greater in invaded plant communities even

when no white-tailed deer were detected.

382   Across the eight field sites, microclimate conditions (relative humidity and maximum

temperature) in invaded and native plant communities were similar to what was observed at the

field experiment site (Appendix S1: Figure S9). Mean ± SE of daily minimum relative humidity

(%) was 92.0 ± 0.6 in invaded areas and 68.4 ± 0.9 in native areas, while average daily

maximum temperature (ºC) was 30.0 ± 0.1 and 34.00 ± 0.2 in invaded and native areas,

respectfully (Appendix S1: Figure S9).

Discussion

These results contribute to a growing understanding that introductions of non-native species may

have a myriad of indirect effects on ecological communities (Pejchar and Mooney 2009, Linders

et al. 2019). We demonstrated that an invasive plant species increased survival time of human

disease vectors that may maintain tick abundances in less favorable wildlife host habitat. The

invasive grass *Imperata*, which is widespread in the southeastern USA, likely influences tick

exposure risk through multiple mechanistic pathways. Consistent with our hypothesis about

invader effects on microclimate, lower temperatures and higher humidity corresponded with

greatly prolonged tick survival in invaded areas. We also hypothesized that invaded areas would

have greater tick abundance than native-dominated areas because of longer tick survival and

greater white-tailed deer activity. Contrary to this hypothesis, we observed no overall difference

in white-tailed deer activity between native and invaded areas and detected no difference in

average tick abundance between the two areas during the time period of peak tick abundance

(Davidson et al. 1994, Gleim et al. 2014). However, there was a positive relationship between

tick abundance and white-tailed deer activity observed only in native-dominated areas. Together,

these results suggest the invasive plant is affecting multiple, potentially off-setting, mechanistic

pathways contributing to tick exposure risk.

Our results show that tick abundances in *Imperata*-invaded plant communities may be

maintained even with relatively low white-tailed deer activity due to more favorable

microclimate conditions that prolong off-host tick survival. In contrast, native plant communities

were less favorable for ticks because they had consistently lower humidity at or below the

critical equilibrium humidity (CEH). The cumulative stress of relative humidity at or below the

CEH predisposed ticks to mortality in native-dominated areas during persistently hotter periods

or rapid increases in temperature, which were buffered in invaded areas. Our field experiment

results also validate the thresholds of relative humidity and temperature for tick desiccation and

mortality described in foundational laboratory experiments (Sauer and Hair 1971, Hair et al.

1975).

Deer are widely considered a key contributing factor to abundances of several North

American tick species but the shape of this relationship is often difficult to detect (Kugeler et al.

2016, Hofmeester et al. 2017). Other studies have emphasized the complexity of this

relationship, detecting either a weak positive relationship (Ostfeld et al. 2018), or no correlation

between deer abundance and tick abundance (Hofmeester et al. 2017). In our analysis of data

from eight field sites, we detected a positive relationship between deer and tick abundance in

native plant communities (Figure 4). However, one site appears to be driving this relationship,

and when that site is removed from the analysis the effect is no longer significant. In contrast to

previous studies, we did not find a relationship between tick abundance and deer activity in

invaded plant communities (Williams et al. 2009, Allan et al. 2010a), however, both studies

evaluated non-native invasive shrubs, which may promote deer activity by providing forage or

cover. Although other species are known to use *Imperata* invaded areas for refuge, e.g., wild

turkey (*Meleagris gallopavo,* Cabrera et al. 2021) and cotton rats (*Sigmodon hispidus*, Borden et

al. 2022), our results suggest that *Imperata* invasions aren't preferred over native plant

communities by white-tailed deer.

Climate change is predicted to facilitate the spread of *Imperata,* expanding its range from

southeastern Gulf States to as far north as Virginia along the eastern USA seaboard (Bradley et

al. 2010). As *Imperata* spreads northward, it is likely to interact with other tick species, including

the blacklegged tick *Ixodes scapularis,* the primary vector for Lyme disease in the United States,

which could increase tick-borne disease risk (Steere et al. 2004). The geographic distribution of

the black-legged tick is also expanding, with an increase in the number of USA counties

reporting its presence from 1996-2005 (Eisen et al. 2016), and ongoing landscape change

facilitates this invasion (Gardner et al. 2020). The blacklegged tick has a similar CEH to that of

A. americanum, suggesting other tick species may benefit from the relatively high relative

humidity and low maximum temperature generated by *Imperata* invasions (Berger et al. 2014).

This study uniquely demonstrated how alterations of abiotic and biotic factors by

invasive plants can impact tick exposure risk. Despite the duration and precision of the survival

assay and the scale of the field study, there were limitations. For example, although we

documented much longer tick survival due to a plant invasion, and strong evidence for

microclimate driving that pattern, additional studies replicated across a larger environmental

gradient of invasion would be useful for validating this result. In addition, including sites that

vary in invader density or biomass could inform management of highly invaded areas to reduce

tick-borne disease risk. Finally, while we focused on white-tailed deer activity, in part because of

invader height and difficulty camera trapping rodents or other hosts, quantifying invader effects

on other hosts (small- and medium-sized mammals, birds, etc.) that are known to influence tick-

borne disease risk could help in resolving the relative influence of host abundance and

microclimate-driven survival on tick abundance (Greiner et al. 1984, Clark 2004, Ostfeld et al.

2018). Additionally, since CO_2 traps only capture host-seeking ticks, actual tick abundance could

be greater in invaded areas if cogongrass improves microclimate significantly where ticks might

be more selective about when they expend energy to pursue a host.

Other studies that have evaluated the effects of invasive plants on tick abundance have

focused on woody invaders (*Lonicera macckii*, *Berberis thunbergia, Rosa multiflora*) and were

conducted in the Midwest or Northeast US (Williams et al. 2009, Allan et al. 2010a,

Adalsteinsson et al. 2016). These studies have found that woody invasive plants alter tick

abundance predominately by providing host habitat rather than creating more favorable

microclimates for tick survival (but see Williams and Ward 2010). Tick survival assays

conducted using the invasive grass *Microstegium vimineum* found that the invader created less

favorable microclimate conditions and reduced survival time for off-host ticks (Civitello et al.

2008). The dense herbaceous structure of *Imperata* invasions differ drastically from that of *M.*

vimineum and the woody invaders that have been studied previously, making comparisons across

systems difficult. Impacts of plant invasions on tick abundance will likely differ by location and

species studied, thus there is a need to address how invasive plants impact tick abundance

through changes to host activity as well as how tick survival is altered by invader-driven changes

in microclimate.

Our findings demonstrate potential indirect effects of plant invasions on human health.

Specifically, we provide evidence for how tick exposure risk may be driven by different

mechanistic pathways in invaded and native plant communities. Consistent with Wei et al.

(2020), invaded plant communities appear to play an important, but relatively underappreciated,

role in altering human vector-borne disease risk. Invasive plant species have a wide variety of

traits that could differentially affect off-host tick survival and host activity. While measurement

of microclimate factors and tick survival are straightforward for field ecology research studies,

measurement of activity of large-bodied mammals with large home range sizes, such as white-tailed deer, is notoriously difficult. Thus, we encourage future work to explore these relationships among invasive plants, environmental conditions, ticks, and their hosts more generally across other invasive plant and tick species, and under different climates, landscape conditions, and disturbance regimes. Understanding the relative contribution of host use and microclimate across habitats invaded by different species will aid in our ability to determine current, and forecast future, impacts of plant invasions on tick-borne disease risk. More broadly, these findings highlight the complex and indirect mechanisms by which invasive species may impact species interactions.

Acknowledgements

We thank Taylor Clark for assistance with data collection, Tempest McCabe for feedback and assistance with statistical analysis, and members of the Flory Lab at the University of Florida for comments and feedback on figures and text.

Funding

This research was funded by the Strategic Environmental Research and Development Program (RC-2636).

The authors declare no conflict of interest

497   **References**
498
Adalsteinsson, S. A., V. D'Amico, W. G. Shriver, D. Brisson, and J. J. Buler. 2016. Scale-dependent

effects of nonnative plant invasion on host-seeking tick abundance. Ecosphere 7.

Alba, C., J. E. NeSmith, C. Fahey, C. Angelini, and S. L. Flory. 2017. Methods to test the

interactive effects of drought and plant invasion on ecosystem structure and function

using complementary common garden and field experiments. Ecology and Evolution

7:1442–1452.

Allan, B. F., H. P. Dutra, L. S. Goessling, K. Barnett, J. M. Chase, R. J. Marquis, G. Pang, G. A.

Storch, R. E. Thach, and J. L. Orrock. 2010a. Invasive honeysuckle eradication reduces

tick-borne disease risk by altering host dynamics. Proceedings of the National Academy

of Sciences 107:18523–18527.

Allan, B. F., L. S. Goessling, G. A. Storch, and R. E. Thach. 2010b. Blood meal analysis to identify

reservoir hosts for Amblyomma americanum ticks. Emerging Infectious Diseases

16:433–440.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2014. Fitting linear mixed-effects models using

lme4. arXiv:1406.5823 [stat].

Berger, K. A., H. S. Ginsberg, K. D. Dugas, L. H. Hamel, and T. N. Mather. 2014. Adverse moisture

events predict seasonal abundance of Lyme disease vector ticks (Ixodes scapularis).

Parasites & Vectors 7:181.

Bloemer, S. R., E. L. Snoddy, J. C. Cooney, and K. Fairbanks. 1986. Influence of Deer Exclusion on

Populations of Lone Star Ticks and American Dog Ticks (Acari: Ixodidae). Journal of

Economic Entomology 79:679–683.

Borden, J. B., K. M. San Antonio, G. Tomat-Kelly, T. Clark, and S. L. Flory. 2022. Invasive grass

indirectly alters seasonal patterns in seed predation. Biology Letters 18:20220095.

Bradley, B. A., D. S. Wilcove, and M. Oppenheimer. 2010. Climate change increases risk of plant

invasion in the Eastern United States. Biological Invasions 12:1855–1872.

Cabrera, S., D. Hiatt, W. W. Dillon, T. Clark, B. F. Allan, and S. L. Flory. 2021. Observations of wild

turkey nesting in invasive cogongrass. Southeastern Naturalist 20:N42.

Childs, J. E., and C. D. Paddock. 2003. The ascendancy of Amblyomma americanum as a vector

of pathogens affecting humans in the United States. Annual Review of Entomology

48:307–337.

Civitello, D. J., S. L. Flory, and K. Clay. 2008. Exotic grass invasion reduces survival of

Amblyomma americanum and Dermacentor variabilis ticks (Acari: Ixodidae). Journal of

Medical Entomology 45:6.

Clark, K. 2004. Borrelia species in host-seeking ticks and small mammals in northern Florida.

Journal of Clinical Microbiology 42:5076–5086.

Davidson, W. R., D. A. Siefken, and L. H. Creekmore. 1994. Seasonal and annual abundance of

Amblyomma americanum (Acari: Ixodidae) in central Georgia. Journal of medical

entomology 31:67–71.

Dillon, W. W., D. Hiatt, and S. L. Flory. 2021. Experimental manipulation of fuel structure to

evaluate the potential ecological effects of fire. Forest Ecology and Management

482:118884.

Eisen, R. J., L. Eisen, and C. B. Beard. 2016. County-Scale Distribution of Ixodes scapularis and

Ixodes pacificus (Acari: Ixodidae) in the Continental United States. Journal of Medical

Entomology 53:349–386.

Elias, S. P., C. B. Lubelczyk, P. W. Rand, E. H. Lacombe, M. S. Holman, and R. P. Smith. 2006.

Deer browse resistant exotic-invasive understory: An indicator of elevated human risk of

exposure to Ixodes scapularis (Acari: Ixodidae) in southern coastal Maine woodlands.

Journal of Medical Entomology 43:11.

Estrada, J. A., and S. L. Flory. 2015. Cogongrass (Imperata cylindrica) invasions in the US:

Mechanisms, impacts, and threats to biodiversity. Global Ecology and Conservation 3:1–

10.

Fahey, C., C. Angelini, and S. L. Flory. 2018. Grass invasion and drought interact to alter the

diversity and structure of native plant communities. Ecology 99:2692–2702.

Fischhoff, I. R., F. Keesing, and R. S. Ostfeld. 2019. Risk Factors for Bites and Diseases Associated

With Black-Legged Ticks: A Meta-Analysis. American Journal of Epidemiology 188:1742–

1750.

Flory, S. L., W. Dillon, and D. Hiatt. 2022. Interacting global change drivers suppress a

foundation tree species. Ecology Letters 25:971–980.

Fox, J., and S. Weisberg. 2018. An R Companion to Applied Regression. SAGE Publications.

Gardner, A. M., N. C. Pawlikowski, S. A. Hamer, G. J. Hickling, J. R. Miller, A. M. Schotthoefer, J. I.

Tsao, and B. F. Allan. 2020. Landscape features predict the current and forecast the

future geographic spread of Lyme disease. Proceedings of the Royal Society B: Biological

Sciences 287:20202278.

Gleim, E. R., L. M. Conner, R. D. Berghaus, M. L. Levin, G. E. Zemtsova, and M. J. Yabsley. 2014.

The phenology of ticks and the effects of long-term prescribed burning on tick

population dynamics in southwestern Georgia and northwestern Florida. PLoS ONE

9:e112174.

Greiner, E. C., P. P. Humphrey, R. C. Belden, W. B. Frankenberger, D. H. Austin, and E. P. J.

Gibbs. 1984. Ixodid ticks on feral swine in Florida. Journal of Wildlife Diseases 20:114–

119.

Guiden, P. W., and J. L. Orrock. 2019. Invasive shrubs modify rodent activity timing, revealing a

consistent behavioral rule governing diel activity. Behavioral Ecology 30:1069–1075.

Hair, J. A., J. R. Sauer, and K. A. Durham. 1975. Water balance and humidity preference in three

species of ticks. Journal of Medical Entomology 12:37–47.

Hiatt, D., and S. L. Flory. 2020. Populations of a widespread invader and co-occurring native

species vary in phenotypic plasticity. New Phytologist 225:584–594.

Hiatt, D., K. Serbesoff-King, D. Lieurance, D. R. Gordon, and S. L. Flory. 2019. Allocation of

invasive plant management expenditures for conservation: Lessons from Florida, USA.

Conservation Science and Practice 1:e51.

Hofmeester, T. R., H. Sprong, P. A. Jansen, H. H. T. Prins, and S. E. van Wieren. 2017. Deer

presence rather than abundance determines the population density of the sheep tick,

Ixodes ricinus, in Dutch forests. Parasites & Vectors 10:433.

Jones, K. E., N. G. Patel, M. A. Levy, A. Storeygard, D. Balk, J. L. Gittleman, and P. Daszak. 2008.

Global trends in emerging infectious diseases. Nature 451:990–993.

Keirans, J. E., and T. R. Litwak. 1989. Pictorial key to the adults of hard ticks, family Ixodidae

(Ixodida: Ixodoidea), east of the Mississippi River. Journal of Medical Entomology

26:435–448.

Kensinger, B. J., and B. F. Allan. 2011. Efficacy of dry ice-baited traps for sampling Amblyomma

americanum (Acari: Ixodidae) varies with life stage but not habitat. Journal of Medical

Entomology 48:708–711.

Knülle, W., and D. Rudolph. 1982. Humidity relationships and water balance of ticks. Pages 43–

70 *in* F. D. Obenchain and R. Galun, editors. Physiology of Ticks. Pergamon, Oxford,

United Kingdom.

Kugeler, K. J., R. A. Jordan, T. L. Schulze, K. S. Griffith, and P. S. Mead. 2016. Will culling white-

tailed deer prevent Lyme disease? Zoonoses and Public Health 63:337–345.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest Package: Tests in

Linear Mixed Effects Models. Journal of Statistical Software 82:1–26.

Linders, T. E. W., U. Schaffner, R. Eschen, A. Abebe, S. K. Choge, L. Nigatu, P. R. Mbaabu, H.

Shiferaw, and E. Allan. 2019. Direct and indirect effects of invasive species: Biodiversity

loss is a major mechanism by which an invasive tree affects ecosystem functioning.

Journal of Ecology 107:2660–2672.

Londt, J. G. H., and G. B. Whitehead. 1972. Ecological studies of larval ticks in South Africa

(Acarina: Ixodidae). Parasitology 65:469–490.

Mack, R. N., and M. C. Smith. 2011. Invasive plants as catalysts for the spread of human

parasites. NeoBiota:13–29.

Mackay, A. J., E. J. Muturi, M. P. Ward, and B. F. Allan. 2016. Cascade of ecological

consequences for West Nile virus transmission when aquatic macrophytes invade

stormwater habitats. Ecological Applications 26:219–232.

Maldonado-Ruiz, L. P., Y. Park, and L. Zurek. 2020. Liquid water intake of the lone star tick,

Amblyomma americanum: Implications for tick survival and management. Scientific

Reports 10:6000.

Morand, S., and C. Lajaunie. 2021. Outbreaks of vector-borne and zoonotic diseases are

associated with changes in forest cover and oil palm expansion at global scale. Frontiers

in Veterinary Science 8.

Morens, D. M., G. K. Folkers, and A. S. Fauci. 2004. The challenge of emerging and re-emerging

infectious diseases. Nature 430:242–249.

Needham, G. R., and P. D. Teel. 1991. Off-host physiological ecology of Ixodid ticks. Annual

Review of Entomology 36:659–81.

Ostfeld, R. S., and J. L. Brunner. 2015. Climate change and Ixodes tick-borne diseases of

humans. Philosophical Transactions of the Royal Society B: Biological Sciences

370:20140051.

Ostfeld, R. S., T. Levi, F. Keesing, K. Oggenfuss, and C. D. Canham. 2018. Tick-borne disease risk

in a forest food web. Ecology 99:1562–1573.

Pejchar, L., and H. A. Mooney. 2009. Invasive species, ecosystem services and human well-

being. Trends in Ecology & Evolution 24:497–504.

Perret, J.-L., E. Guigoz, O. Rais, and L. Gern. 2000. Influence of saturation deficit and

temperature on Ixodes ricinus tick questing activity in a Lyme borreliosis-endemic area

(Switzerland). Parasitology Research 86:554–557.

Pyšek, P., V. Jarošík, P. E. Hulme, J. Pergl, M. Hejda, U. Schaffner, and M. Vilà. 2012. A global

assessment of invasive plant impacts on resident species, communities and ecosystems:

the interaction of impact measures, invading species' traits and environment. Global

Change Biology 18:1725–1737.

Racelis, A. E., R. B. Davey, J. A. Goolsby, A. A. P. de León, K. Varner, and R. Duhaime. 2012.

Facilitative ecological interactions between invasive species: Arundo donax stands as

favorable habitat for cattle ticks (Acari: Ixodidae) along the U.S.-Mexico border. Journal

of Medical Entomology 49:410–417.

Rai, P. K., and J. S. Singh. 2020. Invasive alien plant species: Their impact on environment,

ecosystem services and human health. Ecological Indicators 111:106020.

Sauer, J. R., and J. A. Hair. 1971. Water balance in the lone star tick (Acarina: Ixodidae): the

effects of relative humidity and temperature on weight changes and total water

content. Journal of Medical Entomology 8:479–485.

Sauer, J. R., and J. A. Hair, editors. 1986. Morphology, physiology, and behavioral biology of

ticks. E. Horwood ; Halsted Press, Chichester, West Sussex, England : New York.

Schulze, T. L., and R. A. Jordan. 2003. Meteorologically mediated diurnal questing of *Ixodes*

scapularis and *Amblyomma americanum* (Acari: Ixodidae) nymphs. Journal of Medical

Entomology 40:395–402.

Schulze, T. L., R. A. Jordan, and R. W. Hung. 1997. Biases associated with several sampling

methods used to estimate abundance of Ixodes scapularis and Amblyomma

americanum (Acari: Ixodidae). Journal of Medical Entomology 34:615–623.

Seebens, H., T. M. Blackburn, E. E. Dyer, P. Genovesi, P. E. Hulme, J. M. Jeschke, S. Pagad, P.

Pyšek, M. Winter, M. Arianoutsou, S. Bacher, B. Blasius, G. Brundu, C. Capinha, L.

Celesti-Grapow, W. Dawson, S. Dullinger, N. Fuentes, H. Jäger, J. Kartesz, M. Kenis, H.

Kreft, I. Kühn, B. Lenzner, A. Liebhold, A. Mosena, D. Moser, M. Nishino, D. Pearman, J.

Pergl, W. Rabitsch, J. Rojas-Sandoval, A. Roques, S. Rorke, S. Rossinelli, H. E. Roy, R.

Scalera, S. Schindler, K. Štajerová, B. Tokarska-Guzik, M. van Kleunen, K. Walker, P.

Weigelt, T. Yamanaka, and F. Essl. 2017. No saturation in the accumulation of alien

species worldwide. Nature Communications 8:14435.

Steere, A. C., J. Coburn, and L. Glickstein. 2004. The emergence of Lyme disease. The Journal of

Clinical Investigation 113:1093–1101.

Steullet, P., and P. M. Guerin. 1992. Perception of breath components by the tropical bont tick,

Amblyomma variegatum Fabricius (Ixodidae) 170:665–676.

Stewart, P. S., R. A. Hill, P. A. Stephens, M. J. Whittingham, and W. Dawson. 2021. Impacts of

invasive plants on animal behaviour. Ecology Letters 24:891–907.

Team, R. C. 2019. R: A language and environment for statistical computing [Computer software

manual]. Vienna, Austria.

Vilà, M., J. L. Espinar, M. Hejda, P. E. Hulme, V. Jarošík, J. L. Maron, J. Pergl, U. Schaffner, Y. Sun,

and P. Pyšek. 2011. Ecological impacts of invasive alien plants: a meta-analysis of their

effects on species, communities and ecosystems. Ecology letters 14:702–708.

Vitousek, P. M., C. M. D'Antonio, L. L. Loope, and R. Westbrooks. 1996. Biological invasions as

global environmental change. American scientist 84:468–478.

Wei, C.-Y., J.-K. Wang, H.-C. Shih, H.-C. Wang, and C.-C. Kuo. 2020. Invasive plants facilitated by

socioeconomic change harbor vectors of scrub typhus and spotted fever. PLOS

Neglected Tropical Diseases 14:e0007519.

Williams, S. C., and J. S. Ward. 2010. Effects of Japanese barberry (Ranunculales: Berberidaceae)

removal and resulting microclimatic changes on Ixodes scapularis (Acari: Ixodidae)

abundances in Connecticut, USA. Environmental Entomology 39:1911–1921.

Williams, S. C., J. S. Ward, T. E. Worthley, and K. C. Stafford. 2009. Managing Japanese barberry

(Ranunculales: Berberidaceae) infestations reduces blacklegged tick (Acari: Ixodidae)

abundance and infection prevalence with Borrelia burgdorferi (Spirochaetales:

Spirochaetaceae). Environmental Entomology 38:977–984.

684
685 **Figure 1.** Conceptual diagram of two pathways that contribute to tick-borne disease risk. Native
686 and invaded plant communities may have different effects on both host (e.g., deer, rodent)
687 abundances and tick survival, which together determine tick abundance and, in turn, influence
688 human tick exposure and tick-borne disease risk.
689

Figure 2. Tick survival per plot and microclimate conditions in native and invaded areas. Percent survival (mean ± SE) of (a) nymph and (b) adult Amblyomma americanum in native (blue) and invaded (red) plots over time (start date of experiment June 21, 2018). Box plots below panels (a) and (b) are distributions of time to 100% mortality across plots. Microclimate conditions: (c) average daily minimum relative humidity (RH) (%), dotted line indicates critical equilibrium humidity for Amblyomma americanum (80% RH), and (d) average daily maximum temperature (ºC). Points in (c) and (d) are daily averages across all plots (n = 12) in each treatment. Curved lines in panels (c and d) are polynomial non-linear least squares regression (solid) with 95% CI (dashed) for native (blue) and invaded (red) areas.

701
Figure 3. Number of days survived for individual nymph (a, b) and adult (c, d) life stage ticks
and microclimate conditions that were experienced in native (blue) and invaded (red) plots.
Relationship between days survived and hours per day below 80% RH, and the relationship
between days survived and the maximum temperature that was experienced prior to mortality for
nymphs (a, b) and adults (c, d). Best fit lines (solid) with 95% confidence interval (dashed) are
from linear mixed effects models. Box plots indicate median and 25th and 75th percentile, with
10th and 90th percentile whiskers for days survived (y-axes) and microclimate conditions
experienced by individual ticks in each treatment (x-axes).
710

711
712 **Figure 4.** Relationship between host use (deer activity) and tick abundance (nymph and adult
713 ticks per plot) in native (blue) and invaded (red) plant communities. Best fit lines (solid) from
714 linear mixed effect models with 95% confidence interval (dashed).
715
716