1 **Mechanistic pathways of tick exposure risk in native and invaded plant communities**
2

3 Drew Hiatt[1*], Whalen W. Dillon[1], Allison Gardner[2], Steven Cabrera[3], Michael Dietze[4], Brian F.
4 Allan[5], and S. Luke Flory[1]
5
[*]Corresponding author dhiatt@ufl.edu ph:(775) 790-4303
[1]Agronomy Department, University of Florida, Gainesville, FL, USA
[2]School of Biology and Ecology, University of Maine, Orono ME, USA
[3]College of Forestry, Wildlife and Environment, Auburn University, Auburn, AL, USA
[4]Department of Earth and Environment, Boston University, Boston, MA, USA
[5]Department of Entomology, University of Illinois Urbana-Champaign, Urbana, IL, USA

12
13 **Running title: < Plant invasion effects on ticks >**
14
15 **Keywords:** *Amblyomma americanum, Imperata cylindrica*, tick survival, disease ecology,
16 invasion, indirect effects
17
18 **Statement of authorship:**
19 DH, WD, BA, MD, and SLF conceived the idea and designed the experiment. DH, WD, and SC
20 carried out the experiment. DH, WD, SC, and AG performed the analyses. DH wrote the first
21 draft of the manuscript and all authors contributed substantially to revisions.
22
23 **Data accessibility (Open Research Statement):**
24 All data and code supporting these results has been archived via data dryad and zenodo:
25 https://doi.org/10.5061/dryad.mpg4f4r2k
26 https://doi.org/10.5281/zenodo.6476505
27
Section length:
Abstract: 257/ 350 words
Manuscript length: 34 pages
Number of references: 73
Number of figures: 4
Number of tables/ textboxes: 0
34
35

Drew Hiatt   ORCiD: 0000-0001-9514-9585
Whalen Dillon   ORCiD: 0000-0001-8171-0407
Allison Gardner   ORCiD: 0000-0001-8438-0634
Steven Cabrera   ORCiD: 0000-0002-4637-7458
Michael Dietze   ORCiD: 0000-0002-2324-2518
Brian F. Allan   ORCiD: 0000-0002-1155-3271
S. Luke Flory   ORCiD: 0000-0003-3336-8613

Abstract

Plant invasions may alter disease vector abundance by several mechanistic pathways, including

modifying microclimates that influence vector survival or changing habitats to influence host

use. Here, we used a field experiment and observational data to evaluate multiple mechanistic

pathways (tick survival and host abundance) by which plant invasions may alter vector-borne

disease risk using the common disease vector lone star tick (*Amblyomma americanum*), its

preeminent host white-tailed deer (*Odocoileus virginianus*), and the widespread invasive

cogongrass (*Imperata cylindrica*) in the southeastern USA. In the field experiment, ticks

survived over 50% longer in areas dominated by the invasive plant compared to those with only

native plant species. Invaded areas had lower temperatures and higher relative humidity, yielding

a lower vapor pressure deficit (VPD) that likely reduced tick desiccation. The observational

study showed similar average tick abundance in native and invaded plant communities and no

difference in wildlife host (white-tailed deer) activity between plant communities. However,

there was a positive relationship between tick abundance and white-tailed deer activity, but only

in native areas. Together, these results suggest that more favorable microclimate conditions

resulting in greater tick longevity are the dominant driver of tick abundance in invaded areas,

while tick abundance in native-dominated areas may be promoted, at least in part, by white-tailed

deer activity. Our results demonstrate that plant invasions can affect multiple, potentially

counteracting mechanistic pathways that contribute to tick exposure risk. The complexity of

these relationships highlights the need for better understanding of how invasive species and other

global change drivers influence disease vectors and, ultimately, disease transmission.

Introduction

Human-mediated introductions of species to non-native ranges have resulted in widespread

biological invasions, which are a key driver of global environmental change (Vitousek et al.

1996). Accumulating evidence demonstrates that invasive species can have direct effects on

ecological communities by altering, for example, biodiversity, nutrient cycling, and disturbance

regimes (Vilà et al. 2011, Pyšek et al. 2012). However, they can elicit indirect effects that may be

larger than their direct effects, such as through changes to habitat structure and microclimate

(Linders et al. 2019). These indirect effects can have cascading impacts on humans by affecting

food production, recreation, and climate mitigation (Pejchar and Mooney 2009). Invasive species

may also indirectly influence human health if they alter interactions among species that transmit

diseases to humans (Rai and Singh 2020), however, to our knowledge such effects have been

rarely explored.

Infectious pathogens transmitted from animals to humans via arthropod vectors such as

mosquitoes and ticks place a major burden on global health (Morens et al. 2004, Jones et al.

2008). Tick-borne disease transmission to humans results from complex species interactions that

are mediated by impacts of plant communities on tick growth, survival, dispersal, questing, and

reproduction (Ostfeld et al. 2018, Morand and Lajaunie 2021) and host habitat conditions that

affect the frequency and duration of host use (Elias et al. 2006, Allan et al. 2010a, Guiden and

Orrock 2019). Plant invasions can alter multiple mechanistic pathways that affect tick abundance

and infectious disease exposure for humans by modifying either habitats that alter host

abundance or microclimate conditions that promote or inhibit tick survival (Allan et al. 2010a,

Williams and Ward 2010, Stewart et al. 2021) (Figure 1). Entomological metrics of exposure risk

include measures of tick abundance (e.g., density of ticks), pathogen infection prevalence, or

their product (e.g., density of infected ticks) (Fischhoff et al. 2019). Vector abundance often is

used as a proxy for tick exposure risk, indicating a human's likelihood of encountering a tick

upon entering a particular habitat (Barbour and Fish 1993, Fischhoff et al. 2019). While there has

been research on how these pathways individually affect tick abundance, studies that

simultaneously address multiple pathways are needed to improve understanding of complex

processes by which invasive plants may alter tick exposure risk, and thus infectious disease.

Collectively, evidence for mechanisms that influence tick-borne disease in invaded areas

is sparse and inconsistent, and varies by tick species, invasive plant species, and habitat

conditions, among other factors (Civitello et al. 2008, Allan et al. 2010a, Williams and Ward

2010, Mack and Smith 2011, Racelis et al. 2012, Adalsteinsson et al. 2016). Plant invaders have

been shown to increase habitat use by small mammals (Guiden and Orrock 2019), and white-

tailed deer (Elias et al. 2006, Allan et al. 2010a), leading to greater abundance of ticks (Elias et

al. 2006) and pathogen-infected ticks (Allan et al. 2010a) compared to native areas. Plant

invasions may also increase tick abundances relative to native plant dominated areas by

providing favorable microclimate conditions (i.e., higher humidity, lower temperature, or lower

vapor pressure deficit) that reduce tick desiccation (Williams and Ward 2010, Racelis et al. 2012,

Brunner et al. 2023). However, the few studies comparing the abiotic mechanisms affecting tick

survival in invaded and native areas show either no difference (Allan et al. 2010a) or lower tick

survival (Civitello et al. 2008) in invaded compared to native plant dominated communities.

Given the ongoing introduction and spread of both invasive plants (Seebens et al. 2017), and

ticks (Ostfeld and Brunner 2015), there is a critical need to better understand how invasive plants

affect host habitat use (Stewart et al. 2021) as well as the abiotic factors that influence tick

survival.

Here, we measured differences in tick abundance in native and invaded plant

communities, and how these may be attributable to tick survival and host activity, which are the

two main pathways that determine tick abundance (Figure 1), in the two plant community types.

Specifically, we used field experiments and observational field studies to compare microclimate

conditions and tick survival (lone star tick, *Amblyomma americanum*) in native plant dominated

areas compared to areas invaded by cogongrass (*Imperata cylindrica*). We also measured relative

activity of a key host, white-tailed deer (*Odocoileus virginianus*), in invaded and native areas,

and evaluated how tick abundance was related to host activity. White-tailed deer serve as an

important host for all three life stages of *A. americanum* (Bloemer et al. 1986).

First, in a field experiment, we measured survival of lab-reared ticks in native plant

communities and adjacent invaded areas while concurrently measuring temperature and relative

humidity (RH) to calculate vapor pressure deficit (VPD), a measure of the strength of

atmospheric desiccation (Grossiord et al. 2020). Second, we evaluated relative white-tailed deer

activity (via camera traps) and abundances of host-seeking ticks (via CO_2 trapping) in native and

invaded sites in a field survey. Because the invader creates much denser stands compared to

native plant communities (Appendix S1: Figure S1), we hypothesized that invaded areas would

have lower temperatures and higher relative humidity, yielding lower vapor pressure deficit,

which would be more favorable for tick survival relative to native areas (Alba et al. 2017). We

also hypothesized that relative white-tailed deer use would be greater in invaded communities

because dense cover created by the invader would provide perceived cover from predators, as

documented with small mammals in experimental *Imperata*-invaded plots (Borden et al. 2022),

which would in turn be linked to greater tick abundance.

Methods

Study system

One of the most widely distributed invasive plant species in the southeastern USA is *Imperata cylindrica,* (L.) Beauv. (cogongrass; hereafter *Imperata*) (Estrada and Flory 2015), and it is one of the top five most costly terrestrial invasive plants to manage in Florida (Hiatt et al. 2019). *Imperata* can invade a wide variety of habitats due in part to its greater trait plasticity (Hiatt and Flory 2020), and drought tolerance (Fahey et al. 2018) compared to co-occurring native plant species. *Imperata* invasions can alter the structure of southern pine forests by producing greater fire intensity compared to native plant communities (Dillon et al. 2021), leading to enhanced fire induced tree mortality (Flory et al. 2022). *Imperata* has also been shown to change microclimate conditions, including greater soil moisture and relative humidity, and lower temperatures at ground level compared to native plant communities due to its higher stem density and greater leaf area (Alba et al. 2017, Cabrera et al. 2021).

Lone star ticks (*Amblyomma americanum*) are hard ticks in the family Ixodidae (Keirans and Litwak 1989) that can transmit pathogens, including the causative agents of human ehrlichiosis, Rocky Mountain spotted fever, and tularemia (Childs and Paddock 2003). Time between blood meals (off-host), either seeking hosts to parasitize, digesting blood meals, or dormant, comprises over 90% of an Ixodid tick's life cycle, making desiccation (i.e., loss of water through the cuticle) (Needham and Teel 1991) the primary cause of off-host tick mortality (Ostfeld and Brunner 2015). Maintaining a constant water balance is one of the most important processes that influence off-host survival and, ultimately, tick population size (Knülle and Rudolph 1982, Needham and Teel 1991, Perret et al. 2000, Maldonado-Ruiz et al. 2020). Ticks will begin to desiccate when they are below their critical equilibrium humidity (CEH), which is

80-82% relative humidity for adult *A. americanum* (Hair et al. 1975, Knülle and Rudolph 1982).

While desiccation is primarily driven by low relative humidity (at or below CEH), higher

temperatures lead to more water molecules escaping the cuticle due to kinetic energy (Sauer and

Hair 1986). For *A. americanum,* temperatures in excess of 35 ºC can contribute to desiccation

and mortality (Sauer and Hair 1971), while all terrestrial arthropods, including ticks, lose their

capacity to retain moisture at temperatures in excess of 40 ºC (Londt and Whitehead 1972).

Relative humidity and temperature can be converted to vapor pressure deficit (VPD) using the

formula from Murray (1967). The daily hazard of tick mortality has been shown to increase

drastically with increasing levels of daily maximum VPD (e.g., > 2 kPa) of tick species such as

Ixodes scapularis and *Ixodes ricinus* (Herrmann and Gern 2010, Brunner et al. 2023). Increases

in daily maximum VPD has also been associated with a reduction in tick counts and limiting

questing activity for *Ixodes* spp. (Sambado et al. 2024) and has shown to be an important

predictor of unsuitable habitat for *A. americanum* (Rochlin et al. 2023). Tick abundance,

including local abundance of *A. americanum,* can also be regulated by host activity (Paddock and

Yabsley 2007, Rochlin et al. 2022). White-tailed deer (*Odocoileus virginianus*) serve as key

hosts for all three life stages of *A. americanum* (Bloemer et al. 1986), and are important reservoir

hosts for multiple pathogens transmitted by *A. americanum* (Allan et al. 2010b).

Field experiment: Tick survival and microclimate conditions

To test if ticks survived longer in *Imperata*-invaded areas, we established an experiment

in a planted longleaf pine (*Pinus palustris*) stand with fine sandy soils near La Crosse, FL

(29.897° N, -82.420° W) that had distinct *Imperata*-invaded and native plant dominated areas

(Appendix S1: Figure S2, S3). The invaded area was highly invaded by *Imperata* and spanned

approximately 6300 m^2, while the surrounding area was not invaded and dominated by native

plant species. The native plant community was typical of a longleaf pine stand, with *Aristida*

stricta (wiregrass), *Rubus flagellaris* (common dewberry), *Carya sp.* (hickory), and *Andropogon*

spp. There was no recent chemical or mechanical management in either the invaded or native

areas, and the native plant community included no other non-native plant species. Because we

conducted our experiment within a forest stand with distinct invaded (*Imperata*) and native areas

in close proximity that shared the same overstory structure, we expected any differences in

microclimate conditions to be due to invasion, not other underlying factors (see Results).

We established 24 plots (1m x 1m), 12 in invaded habitat and 12 in native plant

dominated habitat. Plots were randomly established along two transects, one in each plant

community type, with at least 10m spacing. At the center of each plot, a 300-micron 10 cm x 20

cm nylon mesh bag (Skimz, Singapore) containing 10 nymph and 10 adult (5 female and 5 male)

A. americanum obtained from the National Tick Research and Educational Resource

(Department of Entomology and Plant Pathology, Oklahoma State University, Stillwater, OK)

was placed in a 50 cm tall cylindrical wire mesh exclosure to deter wildlife interference

(Appendix S1: Figure S2). We used lab-reared instead of wild-caught ticks to control for tick age

(within life stage) and to standardize the range of tick exposure to environmental conditions prior

to the experiment. Each mesh bag containing ticks was oriented vertically to facilitate natural

diurnal tick questing (i.e., host seeking behavior) and, although there was no substrate within the

bags, the bags remained in contact with the soil surface to allow for ticks to seek refuge below

the leaf litter to avoid desiccation stress (Schulze and Jordan 2003, Ostfeld and Brunner 2015)

throughout the experiment (Appendix S1: Figure S2).

The start date of the survival experiment (June 21, 2018) coincided with typical off host

periods (between blood meals) for *A. americanum* in the region (Davidson et al. 1994, Gleim et

al. 2014). During these long non-parasitic phases, ticks seek refuge in areas with microclimates

that promote water balance (Knülle and Rudolph 1982, Davidson et al. 1994). Mesh bags were

examined every 7-10 days for the first month and then every 10-14 days afterward to determine

days survived for both nymph and adult ticks. During each visit, bags were removed from the

cages but not opened, and survival status of each individual tick was determined by assessing the

response of the tick to CO_2 stimulation from human breath (Steullet and Guerin 1992). Number

of days survived for individual ticks was determined by the time between the start of the survival

experiment and the date when the tick was observed not responding to CO_2 stimulation. Visits

continued to the site until May 23, 2019, when all ticks had died.

To assess plant community structure and abiotic conditions in invaded and native plant

dominated areas, at the start of the survival experiment we collected understory herbaceous plant

height and standing biomass, light availability at the soil surface, and overstory canopy cover at

each plot. Plant height was measured as the tallest point of living leaf tissue from soil surface.

We collected aboveground biomass by clipping at the soil surface in 25 cm x 25 cm quadrats and

dried at 60 ºC to a constant mass and weighed (± 0.01g). The 25 cm quadrats were placed

adjacent to each 1 m experimental plot, in an area representative of the plant communities in the

experimental plots and far enough away (~1-2m) to not alter microclimate conditions in the

experimental plot. Photosynthetically active radiation (PAR, i.e., light) was measured at ground

level in each plot using an ACCUPAR LP-80 PAR/LAI ceptometer (Decagon Device Inc.,

Pullman, WA, USA). To determine if the pine tree overstory was similar in native and invaded

areas and was not a confounding factor with invasion, overstory canopy cover was estimated

using a concave spherical densiometer (Forestry Suppliers Inc., Jackson, MS, USA) by taking

the average of four readings facing each cardinal direction above each plot.

227    Throughout the experiment we recorded ground-level microclimate conditions

228 (temperature and relative humidity) at five-minute intervals immediately adjacent to each

229 enclosure containing the mesh bags using HOBO U23 Pro v2 loggers (Onset Computer

230 Corporation, Bourne, MA). Each logger was housed in a capped 18-inch length of 3.8 cm

231 diameter PVC pipe to protect it from rainfall and direct sunlight. Holes were drilled on four sides

232 six inches apart along the length of the pipe to allow for airflow and better reflect ambient

233 temperature and relative humidity (see Appendix S1: Figure S2a-b). We used the temperature

234 and relative humidity data at each five-minute interval to calculate VPD (Murray 1967).

235         *Statistical analysis: Site conditions*

236    We conducted our statistical analysis of the tick survival field experiment in several

237 phases. First, we conducted analyses to determine a) if *Imperata* invaded plant communities had

238 different characteristics that might influence microclimate conditions and b) if *Imperata* invasion

239 altered tick survival over time. We used multivariate analysis of variance (MANOVA) to test the

240 hypothesis that *Imperata* was associated with different plant community characteristics (i.e.,

241 vegetation height, standing biomass, light availability at soil surface, and overstory canopy

242 cover). We also plotted microclimate conditions (i.e., lower average daily maximum VPD,

243 higher average daily minimum relative humidity, and lower average daily maximum

244 temperature) over time to test the hypothesis that microclimates differed between native plant

245 dominated and invaded plots. These analyses were done using the "car" package (Fox and

246 Weisberg 2018, R Core Team 2020) in R version 3.6.1.

247     *Statistical analysis: Tick survival and microclimate conditions*

248    We used linear models using general least squares regression to assess how the number of

249 days survived for individual *A. americanum* of each life stage was related to the plant community

and VPD (daily maximum VPD). The general linear mixed effects model on number of days

survived for adult or nymph life stage ticks included the fixed effect terms of treatment (native),

maximum daily VPD experienced prior to mortality, and their interaction. Maximum daily VPD

prior to mortality was recorded as the maximum daily VPD measured at each plot between

survival checks (i.e., maximum VPD between the time when the tick was last observed alive and

when it was observed as dead). If data loggers failed or malfunctioned, then data from that time

interval was omitted from analyses. We included plot as a random effect as well as a compound

symmetry covariance structure based on BIC to account for repeated measures over time. Since

nymphs are generally more susceptible to desiccation due to their higher surface:volume ratio

and lower biomass energy reserves compared to adult ixodid ticks (Knülle and Rudolph 1982,

Needham and Teel 1991), we fit models for nymph and adult life stage ticks separately using the

"nlme" package (Pinheiro et al. 2025). Degrees of freedom were calculated using "lmertest"

(Kuznetsova et al. 2017) in R 3.6.1 (R Core Team 2019).

Field survey: White-tailed deer activity, tick abundance, and microclimate

To examine the relationship between *Imperata* invasions and white-tailed deer activity

and tick abundance, we conducted field surveys in invaded and native dominated plant

communities across eight sites in north central Florida (Appendix S1: Figure S3). Invaded plots

were selected if the *Imperata* invasion covered a minimum of 500 m^2, and native plant

dominated plots were selected if they had visually similar overstory tree canopy structure to that

of the invaded plots. Since we encountered many large invasions with few comparable native

plant communities, at some sites we sampled multiple invaded plots per site. We sampled a total

of 34 plots, 22 plots in *Imperata* invaded areas and 12 in native plant dominated areas. Due to

limited supply of camera traps, a sub-set of the plots were sampled for white-tailed deer activity

(28 plots total, 17 in *Imperata* invaded areas and 11 in native plant dominated areas, Appendix

S1: Table S1). Plots were also sampled for tick abundance and some plots were sampled multiple

times, yielding 51 tick sampling events, 32 in *Imperata* invaded areas and 19 in native plant

communities (Appendix S1: Table S2). Corresponding pairs of native and invaded plots that

were resampled were chosen haphazardly and revisited after a minimum of three weeks from the

previous sampling event (Appendix S1: Table S2). Within a site, invaded and native plant

dominated plots were located at least 20 m away from each other. *Imperata* invasions varied in

size across sites but covered a minimum of 500 m^2 to more than 1 ha. We established plots (10 m

radius) under similar overstory tree species and canopy cover between invaded and native plots,

so the primary difference in habitats was presence or absence of the invader.

We placed two motion-activated cameras (Primos 12MP Proof Cam 02 HD, Primos

Hunting, Flora, MS) in each plot to monitor wildlife activity. We focused our analysis on white-

tailed deer activity only due to the tall vegetation in invaded areas limiting detection of smaller

wildlife hosts with camera traps (Appendix S1: Figure S1). All cameras were set to take a burst

of three photos on a 30 second delay, meaning that after the cameras were motion activated there

would be a 30 second delay before the cameras were able to be activated again. From the burst of

three photos, we analyzed only one photo to count occurrences. In the rare event that an

individual or multiple deer were activating the camera by staying in frame for longer than 30

seconds, the deer in these photos were recorded as one occurrence. We made no effort to correct

for particular individuals as we were interested in measuring white-tailed deer activity per day

rather than number of unique hosts. Cameras were placed simultaneously at native and invaded

plots so that deer activity could be calculated across the same time frame. At the end of the

experiment photos were examined by three individuals for presence of white-tailed deer

(Appendix S1: Table S1). Cameras were positioned on opposite sides of the plot at a distance

between 10 m and 15 m from the center of the plot and approximately 1.5 m aboveground

(Appendix S1: Figure S4). Cameras were positioned so that their fields of view did not overlap,

maximizing the total area of the plot being sampled. Cameras were active from 11 April to 1

June 2019, which coincided with questing activity for nymph and adult *A. americanum* in the

region (Davidson et al. 1994, Gleim et al. 2014). In each field site we also recorded ground-level

microclimate conditions (temperature and relative humidity) from 13 June to 1 August 2019 at

five-minute intervals using HOBO U23 Pro v2 logger and PVC housing as described in the field

experiment above.

While the cameras were deployed, we concurrently estimated tick abundance for each

plot using carbon dioxide-baited traps (hereafter CO_2 trap). The CO_2 trap is one of the most

effective means of sampling *A. americanum* nymph and adult life stage ticks (Schulze et al.

1997) and is less biased by different vegetation structures (e.g., a plant invasion) than drag

sampling (Kensinger and Allan 2011). The CO_2 traps consisted of a mini cooler mounted on a

sheet of plywood (45 cm x 60 cm) with vent holes drilled along the sides of the cooler near the

base to disperse gaseous CO_2 from sublimating dry ice (Appendix S1: Figure S5). For each

sampling event, double-sided carpet tape was placed around the perimeter of the plywood to

capture ticks attracted by the CO_2. Each trap was filled with approximately 2 kg of dry ice and

left in the field for 24 hours before ticks were collected. We placed four CO_2 traps per plot, and

traps were placed at least 10m apart from each other within a plot. CO_2 traps can attract ticks that

are within a 5m radius of the trap (Kensinger and Allan 2011). When plots were resampled, traps

were placed in new locations that were greater than 5m from the previous sampling location.

This sampling approach maximized the area of the 10m radius plot that could be sampled with

CO_2 traps (Appendix S1: Figure S4). In total there were 51 tick sampling events, and with four

CO_2 traps per sample, there were 204 trap nights in the study. Traps were purposefully not

placed in plot centers to ensure that new areas of the plot were examined during each sampling

event. Ticks ensnared on the tape were counted and preserved in 70% ethanol for identification

and to determine life stage.

324   *Statistical analysis: Tick abundance and white-tailed deer activity*

325   Relative white-tailed deer activity was estimated and standardized by dividing the

326   number of camera trap photos containing deer by the number of days (total hours / 24) the

327   camera was deployed and operating. This approach provided data in units of deer photos/day and

328   allowed for comparison of data from cameras with different lengths of operable deployment. Our

329   approach was intended to detect relative differences in activity by white-tailed deer in invaded

330   and native areas, not to estimate white-tailed deer population sizes.

331   We fit a negative binomial general linear mixed model to examine the relationship

between white-tailed deer activity and concurrent estimates of tick abundance from 11 April to 1

June 2019 in native and invaded plant communities. We fit a model of tick abundance (nymph +

adult ticks) using plant invasion status (native vs. invaded), white-tailed deer activity, and their

interaction, as fixed-effect terms and site as a random effect term to account for repeated

measures. Tick abundance includes both nymph and adult life stage ticks, because white-tailed

deer serve as an important host for both life stages of *A. americanum* (Bloemer et al. 1986).

Models were fit using the "lme4" package in R 3.6.1 (Bates et al. 2014, Team 2019). Two sided

t-test was also used to test for differences in relative host activity and in tick abundance between

native and invaded plant communities.

Results

Field experiment: Site conditions

There was a significant difference between the native and invaded plant community types

on the combined dependent variables average plant height, plant biomass, light availability at the

soil surface, and overstory canopy cover (MANOVA; Pillai's Trace $_{1,22}$ = 0.922; p < 0.0001). In

separate one-way ANOVAs, average plant height was 268% higher ($F_{1,22}$ = 179.83; p <0.001)

and plant biomass was 408% greater in invaded than native areas ($F_{1,22}$ = 29.29; p <0.001)

(Appendix S1: Figure S6a, b), which corresponded to 91% less light availability at the soil

surface in invaded areas ($F_{1,22}$ = 21.22; p = <0.001) (Appendix S1: Figure S6c). However, there

was no difference in overstory canopy cover in invaded and native areas ($F_{1,22}$ = 0.15; p =

0.706), indicating differences in plant communities and understory light availability were likely

driven by presence of *Imperata* and not differences in overstory canopy cover (Appendix S1:

Figure S6d).

Field experiment: Microclimate conditions and tick survival

Throughout the survival experiment, average daily maximum VPD was approximately

70% lower in invaded (mean ± SE: 0.99 ± 0.04 kPa) compared to native (1.7 ± 0.07, Figure 2a)

areas. Average daily minimum relative humidity was approximately 16% greater in invaded

(mean ± SE: 79.21% ± 0.79) compared to native (67.98% ± 0.87) (Appendix S1: Figure S7a).

Average daily maximum temperature was approximately 13% lower in invaded (mean ± SE:

27.20°C ± 0.34) compared to native (31.29 °C ± 0.40) (Appendix S1: Figure S7b).

Regardless of sex or life stage, ticks survived longer in invaded compared to native plant

communities (Figure 2 c, d). Time to 100% mortality per plot was, on average, 57% longer for

nymphs and 45% longer for adults in the invaded compared to native dominated plots (Figure 2

c, d). Mean ± SE nymph survival was 130 ± 5.4 days in invaded areas versus 80.5 ± 5.0 days in

native vegetation, while mean adult survival was 256 ± 4.8 and 140 ± 7.3 days in invaded and

native areas, respectively. For nymph life stages, daily maximum VPD (kPa) was significantly

related to fewer days survived in both native and invaded plant communities (Figure 3a)

(Appendix S1: Table S3). Daily maximum VPD (kPa) was significantly related to fewer days

survived for adult life stage ticks in the native plant dominated community (Figure 3b)

(Appendix S1: Table S3). However, in invaded areas there was a significant, positive

relationship between daily maximum VPD and days survived (Figure 3b) (Appendix S1: Table

S3).

Field survey: Microclimate, white-tailed deer activity, and tick abundance

Across the eight field sites, microclimate conditions (vapor pressure deficit, relative

humidity and maximum temperature) in invaded and native plant communities were similar to

what was observed at the field experiment site, though over a much shorter time period

(Appendix S1: Figure S8). Mean ± SE of daily maximum VPD (kPa) was 0.4 ± 0.03 in invaded

areas and 1.8 ± 0.07 in native areas, minimum relative humidity (%) was 91.4 ± 0.7 in invaded

areas and 67.5 ± 1.0 in native areas, while average daily maximum temperature (ºC) was 30.0 ±

0.1 and 34.00 ± 0.2 in invaded and native areas, respectfully (Appendix S1: Figure S8).

There were no differences in relative white-tailed deer activity between native and

invaded plots (t_{44} = 1.01, p = 0.318) (Appendix S1: Figure S9). There were also no differences in

average tick abundance (nymph [t_{43} = -0.713, p = 0.480]; adult [t_{43} = -1.031, p = 0.308] life

stage ticks) between native and invaded areas (Appendix S1: Figure S10). In native

communities, tick abundance increased with white-tailed deer activity (deer photos/day)

(estimate = 0.6 ticks per deer photo/day; 95% CI = 0.1 – 1.1) (Figure 4, Appendix S1: Table S4),

while in invaded habitats, white-tailed deer activity was not related to tick abundance (estimate =

-0.4 ticks per deer photo/day; 95% CI = -0.9 – 0.1). For plots where no white-tailed deer were

detected (i.e., intercept term), invaded plant communities had positive tick abundances (estimate

= 1.4 ticks; 95% CI = 0.2 – 2.5) (Figure 4, Appendix S1: Table S4). There was one plot with

both the highest tick abundance and highest deer activity that could have strongly influenced the

results, so we ran the analysis using the same linear mixed effects model but with that data point

removed. Omitting that one data point resulted in no association between white-tailed deer

activity and tick abundance (estimate = -0.4 ticks per deer photo/day; 95% CI = -0.9 – 0.1)

(Appendix S1: Table S5), but there was a significant intercept term (estimate = 1.3 ticks; 95% CI

= 0.1 – 2.6) (Appendix S1:Table S5), indicating that tick abundance was greater in invaded plant

communities even when no white-tailed deer were detected.

Discussion

These results contribute to a growing understanding that introductions of non-native species may

have a myriad of indirect effects on ecological communities (Pejchar and Mooney 2009, Linders

et al. 2019). We documented that the invasive plant species cogongrass was associated with

greater tick survival time, which could maintain tick abundances even though invaded areas may

have less activity by the common host white-tailed deer. Consistent with our hypothesis about

differences in microclimate due to invasion, there was greatly prolonged tick survival in invaded

areas, which had lower vapor pressure deficit corresponding to both lower temperatures and

higher humidity. We had also hypothesized that invaded areas would have greater tick

abundance than native-dominated areas because of longer tick survival times and greater white-

tailed deer activity. Contrary to this hypothesis, we observed no overall difference in white-tailed

deer activity between native and invaded areas and detected no difference in average tick

abundance between the two areas during the time period of peak tick abundance (Davidson et al.

1994, Gleim et al. 2014). There was a weak positive relationship between tick abundance and

white-tailed deer activity, though only in native-dominated areas, and the pattern was driven by

one of the eight sites (Figure 4). Together, these results suggest the invasive plant is influencing

multiple, potentially counteracting, mechanistic pathways contributing to tick exposure risk.

Our results show that tick abundances in *Imperata*-invaded plant communities may be

maintained even with relatively low white-tailed deer activity due to more favorable

microclimate conditions that prolong off-host tick survival. Exploration of the relationship

between tick mortality and VPD revealed that survival of nymphal ticks in both vegetation types

and adult ticks in native vegetation declined with increasing VPD. Conversely, adult tick

survival in invaded habitats was prolonged by increasing VPD. We posit that this

counterintuitive result stems from the physiological buffering provided by *Imperata*, where

invaded sites rarely exhibited conditions that have been shown to produce rapid mortality for

other tick species in laboratory conditions (i.e., VPD > 2 kPa) (Herrmann and Gern 2010).

Instead, adult ticks in invaded habitats lived long enough that they eventually died as a result of

loss of metabolic reserves (Alasmari and Wall 2021) or other stressors that were not measured.

More generally, invasive plant buffering of microclimate conditions that would otherwise be

stressful for ticks suggests that invasions may facilitate tick survival under the hotter and drier

conditions expected with climate change (McDowell et al. 2022).

Deer are widely considered a key contributing factor to abundances of several North

American tick species but the shape of this relationship is often difficult to detect (Kugeler et al.

2016, Hofmeester et al. 2017). Other studies have emphasized the complexity of this

relationship, detecting either a weak positive relationship (Ostfeld et al. 2018) or no association

between deer abundance and tick abundance (Hofmeester et al. 2017). We essentially found the

same pattern: in our analysis of data from all eight field sites, we detected a weak positive

relationship between deer and tick abundance in native plant communities but no relationship

when an influential site was removed (Figure 4). In contrast to previous studies (Williams et al.

2009, Allan et al. 2010a), we did not find a relationship between tick abundance and deer activity

in invaded plant communities; however, both of those studies evaluated non-native invasive

shrubs, which may promote deer activity by providing forage or cover. Although other species

are known to use *Imperata* invaded areas for refuge, e.g., wild turkey (*Meleagris gallopavo,*

Cabrera et al. 2021) and cotton rats (*Sigmodon hispidus*, Borden et al. 2022), we found no

difference in abundance of white-tailed deer based on camera trap detections in *Imperata*

invasions compared to native plant communities.

Climate change is predicted to facilitate the spread of *Imperata,* expanding its range from

southeastern Gulf States to as far north as Virginia along the eastern USA seaboard (Bradley et

al. 2010). As *Imperata* spreads northward, it is likely to interact with other tick species, including

the blacklegged tick *Ixodes scapularis,* the primary vector for Lyme disease in the United States,

which could increase tick-borne disease risk (Steere et al. 2004). The geographic distribution of

the black-legged tick is also expanding, with an increase in the number of USA counties

reporting its presence from 1996-2005 (Eisen et al. 2016) facilitated by ongoing landscape

change (Gardner et al. 2020). The blacklegged tick has similar physiological thresholds to *A.*

americanum (Berger et al. 2014)*,* suggesting it too may benefit from the low vapor pressure

deficit generated by *Imperata* invasions*.*

This study uniquely demonstrated how differences in abiotic conditions in invaded areas

can be associated with differences in tick abundance and thus disease exposure risk. Despite the

duration and precision of the survival assay and the scale of the field study, there were

limitations. For example, although we documented much longer tick survival in invaded areas,

and evidence for microclimate conditions driving that pattern, additional studies replicated across

a larger environmental gradient of invasion would be useful for validating this result. In addition,

including sites that vary in invader density or biomass could inform management of highly

invaded areas to reduce tick-borne exposure risk. While we found evidence for microclimate

driven survival, we found less support for increases in host habitat use within invaded plant

communities. We focused on white-tailed deer activity, in part because of invader height and

difficulty camera trapping rodents or other hosts with our methods; however, quantifying other

hosts (small- and medium-sized mammals, birds, etc.) in invaded areas that are known to

influence tick-borne disease risk could resolve the relative influence of host abundance and

microclimate-driven survival on tick abundance (Greiner et al. 1984, Clark 2004, Ostfeld et al.

2018). Additionally, since CO_2 traps only capture host-seeking ticks, actual tick abundance could

be greater in invaded areas if *Imperata* improves microclimate to the point where ticks might be

more selective about when they expend energy to pursue a host.

The dense herbaceous structure of *Imperata* invasions differ drastically from that of the

woody and grassy invaders that have been studied previously, making comparisons across

species and systems difficult. For example, studies evaluating the effects of woody invasive

plants on tick abundance (*Lonicera maackii*, *Berberis thunbergia, Rosa multiflora*) in the

Midwest or Northeast US (Williams et al. 2009, Allan et al. 2010a, Adalsteinsson et al. 2016)

have found that woody invasive plants alter tick abundance predominately by providing host

habitat rather than creating more favorable microclimates for tick survival (but see Williams and

Ward 2010). By contrast, tick survival assays conducted using the invasive grass *Microstegium*

vimineum found that the invader created less favorable microclimate conditions and reduced

survival time for off-host ticks (Civitello et al. 2008). Impacts of plant invasions on tick

abundance will likely differ by location and species studied, thus there is a need to address how

invasive plants impact tick abundance through changes to host activity as well as how tick

survival is altered by invader-driven changes in microclimate.

Consistent with Wei et al. (2020), our findings suggest that invaded plant communities

play an important, but relatively underappreciated, indirect role in altering human vector-borne

disease risk. Specifically, we provide evidence for how tick exposure risk may be driven by

different mechanistic pathways in invaded and native plant communities. Invasive plant species

have a wide variety of traits that could differentially affect off-host tick survival and host

activity. While measurement of microclimate factors and tick survival are straightforward for

field ecology research studies, measurement of activity of large-bodied mammals with large

home range sizes, such as white-tailed deer, is notoriously difficult. Thus, we encourage future

work to explore these relationships among invasive plants, environmental conditions, ticks, and

their hosts more generally across other invasive plant and tick species, and under different

climates, landscape conditions, and disturbance regimes. Understanding the relative contribution

of host use and microclimate across habitats invaded by different species will aid in our ability to

determine current, and forecast future, impacts of plant invasions on tick-borne disease risk.

More broadly, these findings highlight the complex and indirect mechanisms by which invasive

species may impact species interactions.

Acknowledgements

We thank Taylor Clark for assistance with data collection, Tempest McCabe for feedback and

assistance with statistical analysis, and members of the Flory Lab at the University of Florida for

comments and feedback on figures and text.

Funding

This research was funded by the U.S. Strategic Environmental Research and Development

Program (RC-2636).

The authors declare no conflict of interest

References

Adalsteinsson, S. A., V. D'Amico, W. G. Shriver, D. Brisson, and J. J. Buler. 2016. Scale-dependent

effects of nonnative plant invasion on host-seeking tick abundance. Ecosphere 7.

Alasmari, S., and R. Wall. 2021. Metabolic rate and resource depletion in the tick Ixodes ricinus

in response to temperature. Experimental and Applied Acarology 83:81–93.

Alba, C., J. E. NeSmith, C. Fahey, C. Angelini, and S. L. Flory. 2017. Methods to test the

interactive effects of drought and plant invasion on ecosystem structure and function

using complementary common garden and field experiments. Ecology and Evolution

7:1442–1452.

Allan, B. F., H. P. Dutra, L. S. Goessling, K. Barnett, J. M. Chase, R. J. Marquis, G. Pang, G. A.

Storch, R. E. Thach, and J. L. Orrock. 2010a. Invasive honeysuckle eradication reduces

tick-borne disease risk by altering host dynamics. Proceedings of the National Academy

of Sciences 107:18523–18527.

Allan, B. F., L. S. Goessling, G. A. Storch, and R. E. Thach. 2010b. Blood meal analysis to identify

reservoir hosts for Amblyomma americanum ticks. Emerging Infectious Diseases

16:433–440.

Barbour, A. G., and D. Fish. 1993. The Biological and Social Phenomenon of Lyme Disease.

Science 260:1610–1616.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2014. Fitting linear mixed-effects models using

lme4. arXiv:1406.5823 [stat].

Berger, K. A., H. S. Ginsberg, K. D. Dugas, L. H. Hamel, and T. N. Mather. 2014. Adverse moisture

events predict seasonal abundance of Lyme disease vector ticks (Ixodes scapularis).

Parasites & Vectors 7:181.

Bloemer, S. R., E. L. Snoddy, J. C. Cooney, and K. Fairbanks. 1986. Influence of Deer Exclusion on

Populations of Lone Star Ticks and American Dog Ticks (Acari: Ixodidae). Journal of

Economic Entomology 79:679–683.

Borden, J. B., K. M. San Antonio, G. Tomat-Kelly, T. Clark, and S. L. Flory. 2022. Invasive grass

indirectly alters seasonal patterns in seed predation. Biology Letters 18:20220095.

Bradley, B. A., D. S. Wilcove, and M. Oppenheimer. 2010. Climate change increases risk of plant

invasion in the Eastern United States. Biological Invasions 12:1855–1872.

Brunner, J. L., S. L. LaDeau, M. Killilea, E. Valentine, M. Schierer, and R. S. Ostfeld. 2023. Off-host

survival of blacklegged ticks in eastern North America: A multistage, multiyear, multisite

study. Ecological Monographs 93:e1572.

Cabrera, S., D. Hiatt, W. W. Dillon, T. Clark, B. F. Allan, and S. L. Flory. 2021. Observations of wild

turkey nesting in invasive cogongrass. Southeastern Naturalist 20:N42.

Childs, J. E., and C. D. Paddock. 2003. The ascendancy of Amblyomma americanum as a vector

of pathogens affecting humans in the United States. Annual Review of Entomology

48:307–337.

Civitello, D. J., S. L. Flory, and K. Clay. 2008. Exotic grass invasion reduces survival of

Amblyomma americanum and Dermacentor variabilis ticks (Acari: Ixodidae). Journal of

Medical Entomology 45:6.

Clark, K. 2004. Borrelia species in host-seeking ticks and small mammals in northern Florida.

Journal of Clinical Microbiology 42:5076–5086.

Davidson, W. R., D. A. Siefken, and L. H. Creekmore. 1994. Seasonal and annual abundance of

Amblyomma americanum (Acari: Ixodidae) in central Georgia. Journal of medical

entomology 31:67–71.

Dillon, W. W., D. Hiatt, and S. L. Flory. 2021. Experimental manipulation of fuel structure to

evaluate the potential ecological effects of fire. Forest Ecology and Management

482:118884.

Eisen, R. J., L. Eisen, and C. B. Beard. 2016. County-Scale Distribution of Ixodes scapularis and

Ixodes pacificus (Acari: Ixodidae) in the Continental United States. Journal of Medical

Entomology 53:349–386.

Elias, S. P., C. B. Lubelczyk, P. W. Rand, E. H. Lacombe, M. S. Holman, and R. P. Smith. 2006.

Deer browse resistant exotic-invasive understory: An indicator of elevated human risk of

exposure to Ixodes scapularis (Acari: Ixodidae) in southern coastal Maine woodlands.

Journal of Medical Entomology 43:11.

Estrada, J. A., and S. L. Flory. 2015. Cogongrass (Imperata cylindrica) invasions in the US:

Mechanisms, impacts, and threats to biodiversity. Global Ecology and Conservation 3:1–

10.

Fahey, C., C. Angelini, and S. L. Flory. 2018. Grass invasion and drought interact to alter the

diversity and structure of native plant communities. Ecology 99:2692–2702.

Fischhoff, I. R., F. Keesing, and R. S. Ostfeld. 2019. Risk Factors for Bites and Diseases Associated

With Black-Legged Ticks: A Meta-Analysis. American Journal of Epidemiology 188:1742–

1750.

Flory, S. L., W. Dillon, and D. Hiatt. 2022. Interacting global change drivers suppress a

foundation tree species. Ecology Letters 25:971–980.

Fox, J., and S. Weisberg. 2018. An R Companion to Applied Regression. SAGE Publications.

Gardner, A. M., N. C. Pawlikowski, S. A. Hamer, G. J. Hickling, J. R. Miller, A. M. Schotthoefer, J. I.

Tsao, and B. F. Allan. 2020. Landscape features predict the current and forecast the

future geographic spread of Lyme disease. Proceedings of the Royal Society B: Biological

Sciences 287:20202278.

Gleim, E. R., L. M. Conner, R. D. Berghaus, M. L. Levin, G. E. Zemtsova, and M. J. Yabsley. 2014.

The phenology of ticks and the effects of long-term prescribed burning on tick

population dynamics in southwestern Georgia and northwestern Florida. PLoS ONE

9:e112174.

Greiner, E. C., P. P. Humphrey, R. C. Belden, W. B. Frankenberger, D. H. Austin, and E. P. J.

Gibbs. 1984. Ixodid ticks on feral swine in Florida. Journal of Wildlife Diseases 20:114–

119.

Grossiord, C., T. N. Buckley, L. A. Cernusak, K. A. Novick, B. Poulter, R. T. W. Siegwolf, J. S.

Sperry, and N. G. McDowell. 2020. Plant responses to rising vapor pressure deficit. New

Phytologist 226:1550–1566.

Guiden, P. W., and J. L. Orrock. 2019. Invasive shrubs modify rodent activity timing, revealing a

consistent behavioral rule governing diel activity. Behavioral Ecology 30:1069–1075.

Hair, J. A., J. R. Sauer, and K. A. Durham. 1975. Water balance and humidity preference in three

species of ticks. Journal of Medical Entomology 12:37–47.

Herrmann, C., and L. Gern. 2010. Survival of Ixodes ricinus (Acari: Ixodidae) Under Challenging

Conditions of Temperature and Humidity Is Influenced by Borrelia burgdorferi sensu lato

Infection. Journal of Medical Entomology 47:1196–1204.

Hiatt, D., and S. L. Flory. 2020. Populations of a widespread invader and co-occurring native

species vary in phenotypic plasticity. New Phytologist 225:584–594.

Hiatt, D., K. Serbesoff-King, D. Lieurance, D. R. Gordon, and S. L. Flory. 2019. Allocation of

invasive plant management expenditures for conservation: Lessons from Florida, USA.

Conservation Science and Practice 1:e51.

Hofmeester, T. R., H. Sprong, P. A. Jansen, H. H. T. Prins, and S. E. van Wieren. 2017. Deer

presence rather than abundance determines the population density of the sheep tick,

Ixodes ricinus, in Dutch forests. Parasites & Vectors 10:433.

Jones, K. E., N. G. Patel, M. A. Levy, A. Storeygard, D. Balk, J. L. Gittleman, and P. Daszak. 2008.

Global trends in emerging infectious diseases. Nature 451:990–993.

Keirans, J. E., and T. R. Litwak. 1989. Pictorial key to the adults of hard ticks, family Ixodidae

(Ixodida: Ixodoidea), east of the Mississippi River. Journal of Medical Entomology

26:435–448.

Kensinger, B. J., and B. F. Allan. 2011. Efficacy of dry ice-baited traps for sampling Amblyomma

americanum (Acari: Ixodidae) varies with life stage but not habitat. Journal of Medical

Entomology 48:708–711.

Knülle, W., and D. Rudolph. 1982. Humidity relationships and water balance of ticks. Pages 43–

70 *in* F. D. Obenchain and R. Galun, editors. Physiology of Ticks. Pergamon, Oxford,

United Kingdom.

Kugeler, K. J., R. A. Jordan, T. L. Schulze, K. S. Griffith, and P. S. Mead. 2016. Will culling white-

tailed deer prevent Lyme disease? Zoonoses and Public Health 63:337–345.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest Package: Tests in

Linear Mixed Effects Models. Journal of Statistical Software 82:1–26.

Linders, T. E. W., U. Schaffner, R. Eschen, A. Abebe, S. K. Choge, L. Nigatu, P. R. Mbaabu, H.

Shiferaw, and E. Allan. 2019. Direct and indirect effects of invasive species: Biodiversity

loss is a major mechanism by which an invasive tree affects ecosystem functioning.

Journal of Ecology 107:2660–2672.

Londt, J. G. H., and G. B. Whitehead. 1972. Ecological studies of larval ticks in South Africa

(Acarina: Ixodidae). Parasitology 65:469–490.

Mack, R. N., and M. C. Smith. 2011. Invasive plants as catalysts for the spread of human

parasites. NeoBiota:13–29.

Maldonado-Ruiz, L. P., Y. Park, and L. Zurek. 2020. Liquid water intake of the lone star tick,

Amblyomma americanum: Implications for tick survival and management. Scientific

Reports 10:6000.

McDowell, N. G., G. Sapes, A. Pivovaroff, H. D. Adams, C. D. Allen, W. R. L. Anderegg, M. Arend,

D. D. Breshears, T. Brodribb, B. Choat, H. Cochard, M. De Cáceres, M. G. De Kauwe, C.

Grossiord, W. M. Hammond, H. Hartmann, G. Hoch, A. Kahmen, T. Klein, D. S. Mackay,

M. Mantova, J. Martínez-Vilalta, B. E. Medlyn, M. Mencuccini, A. Nardini, R. S. Oliveira,

A. Sala, D. T. Tissue, J. M. Torres-Ruiz, A. M. Trowbridge, A. T. Trugman, E. Wiley, and C.

Xu. 2022. Mechanisms of woody-plant mortality under rising drought, CO_2 and vapour

pressure deficit. Nature Reviews Earth & Environment 3:294–308.

Morand, S., and C. Lajaunie. 2021. Outbreaks of vector-borne and zoonotic diseases are

associated with changes in forest cover and oil palm expansion at global scale. Frontiers

in Veterinary Science 8.

Morens, D. M., G. K. Folkers, and A. S. Fauci. 2004. The challenge of emerging and re-emerging

infectious diseases. Nature 430:242–249.

Murray, F. W. 1967. On the Computation of Saturation Vapor Pressure. Journal of Applied

Meteorology and Climatology 6:203–204.

Needham, G. R., and P. D. Teel. 1991. Off-host physiological ecology of Ixodid ticks. Annual

Review of Entomology 36:659–81.

Ostfeld, R. S., and J. L. Brunner. 2015. Climate change and Ixodes tick-borne diseases of

humans. Philosophical Transactions of the Royal Society B: Biological Sciences

370:20140051.

Ostfeld, R. S., T. Levi, F. Keesing, K. Oggenfuss, and C. D. Canham. 2018. Tick-borne disease risk

in a forest food web. Ecology 99:1562–1573.

Paddock, C. D., and M. J. Yabsley. 2007. Ecological Havoc, the Rise of White-Tailed Deer, and the

Emergence of Amblyomma americanum-Associated Zoonoses in the United States.

Pages 289–324 in J. E. Childs, J. S. Mackenzie, and J. A. Richt, editors. Wildlife and

Emerging Zoonotic Diseases: The Biology, Circumstances and Consequences of Cross-

Species Transmission. Springer Berlin Heidelberg, Berlin, Heidelberg.

Pejchar, L., and H. A. Mooney. 2009. Invasive species, ecosystem services and human well-

being. Trends in Ecology & Evolution 24:497–504.

Perret, J.-L., E. Guigoz, O. Rais, and L. Gern. 2000. Influence of saturation deficit and

temperature on Ixodes ricinus tick questing activity in a Lyme borreliosis-endemic area

(Switzerland). Parasitology Research 86:554–557.

Pinheiro, J. S., D. Bates, S. Debroy, D. Sarkar, and R Core Team. 2025, March 31. nlme: Linear

and Nonlinear Mixed Effects Models.

Pyšek, P., V. Jarošík, P. E. Hulme, J. Pergl, M. Hejda, U. Schaffner, and M. Vilà. 2012. A global

assessment of invasive plant impacts on resident species, communities and ecosystems:

the interaction of impact measures, invading species' traits and environment. Global

Change Biology 18:1725–1737.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria.

Racelis, A. E., R. B. Davey, J. A. Goolsby, A. A. P. de León, K. Varner, and R. Duhaime. 2012.

Facilitative ecological interactions between invasive species: Arundo donax stands as

favorable habitat for cattle ticks (Acari: Ixodidae) along the U.S.-Mexico border. Journal

of Medical Entomology 49:410–417.

Rai, P. K., and J. S. Singh. 2020. Invasive alien plant species: Their impact on environment,

ecosystem services and human health. Ecological Indicators 111:106020.

Rochlin, I., A. Egizi, and H. S. Ginsberg. 2023. Modeling of historical and current distributions of

lone star tick, Amblyomma americanum (Acari: Ixodidae), is consistent with ancestral

range recovery. Experimental and Applied Acarology 89:85–103.

Rochlin, I., A. Egizi, and A. Lindström. 2022. The Original Scientific Description of the Lone Star

Tick (*Amblyomma americanum* , Acari: Ixodidae) and Implications for the Species' Past

and Future Geographic Distributions. Journal of Medical Entomology 59:412–420.

Sambado, S., A. J. MacDonald, A. Swei, and C. J. Briggs. 2024. Climate-associated variation in the

within-season dynamics of juvenile ticks in California. Ecosphere 15:e70064.

Sauer, J. R., and J. A. Hair. 1971. Water balance in the lone star tick (Acarina: Ixodidae): the

effects of relative humidity and temperature on weight changes and total water

content. Journal of Medical Entomology 8:479–485.

Sauer, J. R., and J. A. Hair, editors. 1986. Morphology, physiology, and behavioral biology of

ticks. E. Horwood ; Halsted Press, Chichester, West Sussex, England : New York.

Schulze, T. L., and R. A. Jordan. 2003. Meteorologically mediated diurnal questing of *Ixodes*

scapularis and *Amblyomma americanum* (Acari: Ixodidae) nymphs. Journal of Medical

Entomology 40:395–402.

Schulze, T. L., R. A. Jordan, and R. W. Hung. 1997. Biases associated with several sampling

methods used to estimate abundance of Ixodes scapularis and Amblyomma

americanum (Acari: Ixodidae). Journal of Medical Entomology 34:615–623.

Seebens, H., T. M. Blackburn, E. E. Dyer, P. Genovesi, P. E. Hulme, J. M. Jeschke, S. Pagad, P.

Pyšek, M. Winter, M. Arianoutsou, S. Bacher, B. Blasius, G. Brundu, C. Capinha, L.

Celesti-Grapow, W. Dawson, S. Dullinger, N. Fuentes, H. Jäger, J. Kartesz, M. Kenis, H.

Kreft, I. Kühn, B. Lenzner, A. Liebhold, A. Mosena, D. Moser, M. Nishino, D. Pearman, J.

Pergl, W. Rabitsch, J. Rojas-Sandoval, A. Roques, S. Rorke, S. Rossinelli, H. E. Roy, R.

Scalera, S. Schindler, K. Štajerová, B. Tokarska-Guzik, M. van Kleunen, K. Walker, P.

Weigelt, T. Yamanaka, and F. Essl. 2017. No saturation in the accumulation of alien

species worldwide. Nature Communications 8:14435.

Steere, A. C., J. Coburn, and L. Glickstein. 2004. The emergence of Lyme disease. The Journal of

Clinical Investigation 113:1093–1101.

Steullet, P., and P. M. Guerin. 1992. Perception of breath components by the tropical bont tick,

Amblyomma variegatum Fabricius (Ixodidae) 170:665–676.

Stewart, P. S., R. A. Hill, P. A. Stephens, M. J. Whittingham, and W. Dawson. 2021. Impacts of

invasive plants on animal behaviour. Ecology Letters 24:891–907.

Team, R. C. 2019. R: A language and environment for statistical computing [Computer software

manual]. Vienna, Austria.

Vilà, M., J. L. Espinar, M. Hejda, P. E. Hulme, V. Jarošík, J. L. Maron, J. Pergl, U. Schaffner, Y. Sun,

and P. Pyšek. 2011. Ecological impacts of invasive alien plants: a meta-analysis of their

effects on species, communities and ecosystems. Ecology letters 14:702–708.

Vitousek, P. M., C. M. D'Antonio, L. L. Loope, and R. Westbrooks. 1996. Biological invasions as

global environmental change. American scientist 84:468–478.

Wei, C.-Y., J.-K. Wang, H.-C. Shih, H.-C. Wang, and C.-C. Kuo. 2020. Invasive plants facilitated by

socioeconomic change harbor vectors of scrub typhus and spotted fever. PLOS

Neglected Tropical Diseases 14:e0007519.

Williams, S. C., and J. S. Ward. 2010. Effects of Japanese barberry (Ranunculales: Berberidaceae)

removal and resulting microclimatic changes on Ixodes scapularis (Acari: Ixodidae)

abundances in Connecticut, USA. Environmental Entomology 39:1911–1921.

Williams, S. C., J. S. Ward, T. E. Worthley, and K. C. Stafford. 2009. Managing Japanese barberry

(Ranunculales: Berberidaceae) infestations reduces blacklegged tick (Acari: Ixodidae)

abundance and infection prevalence with Borrelia burgdorferi (Spirochaetales:

Spirochaetaceae). Environmental Entomology 38:977–984.

732
733

Figure 1. Conceptual diagram of two pathways that contribute to tick-borne disease risk. Native

and invaded plant communities may have different effects on both host (e.g., deer, rodent)

abundances and tick survival, which together determine tick abundance and, in turn, influence

human tick exposure and tick-borne disease risk. Differences between invaded and native areas

in microhabitat variables such as relative humidity and temperature, which determine vapor

pressure deficit (VPD), can affect off-host tick survival. Vector graphics are copyright-free

images obtained from stock.adobe.com via the University of Florida's enterprise license.

Figure 2. Microclimate conditions and tick survival per plot in native and invaded areas.

Average daily maximum vapor pressure deficit (VPD, kPa) (a), points are the average of

maximum daily VPD across all plots in each plant community type (n = 12). Solid lines in panel

(a) indicate best fit quadratic polynomials for native (blue) and invaded (red) areas. Box plots to

the right of panel (a) indicate the median and 25th and 75th percentile, with 10th and 90th

percentile whiskers for daily maximum VPD (kPa). Percent survival (mean ± SE) of (b) nymph

and (c) adult *Amblyomma americanum* in native (blue) and invaded (red) plots over time; the

second x-axis shows the month of the year. Box plots below panels (b) and (c) are distributions

of time to 100% mortality across the 12 plots showing the median and 25th and 75th percentile,

with 10th and 90th percentile whiskers.

Figure 3. Number of days survived for individual nymph (a) and adult (b) life stage ticks and the daily maximum VPD experienced in native (blue) and invaded (red) plots between survival checks (the time interval when the tick was last seen alive and when it was recorded dead). Best fit lines (dashed) with 95% confidence interval (solid) are from linear mixed effects models. Box plots indicate median and 25th and 75th percentile, with 10th and 90th percentile whiskers for days survived (y-axes) and daily maximum VPD experienced by individual ticks prior to mortality in each treatment (x-axes).

766 **Figure 4.** Relationship between deer activity (deer photos/day) and tick abundance (nymph and

767 adult ticks per plot) in native (blue) and invaded (red) plant communities. Best fit lines (solid)

768 from linear mixed effect models with 95% confidence interval (dashed).