Knockdown of TTLL1 reduces Aβ-induced TAU pathology in human iPSC-derived cortical neurons

Mohamed Aghyad Al Kabbani[1,2], Laura Köhler[1,2], Tamara Wied[3], Daniel Adam[1,2], Jennifer Klimek[1,2], Hans Zempel[1,2,*]

1 Institute of Human Genetics, Faculty of Medicine and University Hospital Cologne, University of Cologne, Cologne, Germany

2 Center for Molecular Medicine Cologne (CMMC), Faculty of Medicine and University Hospital Cologne, University of Cologne, Cologne, Germany

3 Current affiliation: Max Planck Institute for the Biology of Ageing, Cologne, Germany

* Correspondence to: Dr. Dr. Hans Zempel, hans.zempel@uk-koeln.de

Abstract

Microtubules play a crucial role in neuronal structure and function, with their stability and dynamics regulated by posttranslational modifications (PTMs) such as polyglutamylation. In Alzheimer disease (AD), the microtubule-associated protein TAU becomes mislocalized into the somatodendritic compartment ('TAU missorting'), dissociates from microtubules, aggregates into neurofibrillary tangles, and contributes to microtubule destabilization and neuronal death. Here, we investigated the role of Tubulin-Tyrosine-Ligase-Like proteins (TTLLs) in TAU missorting and microtubule dysregulation using human induced pluripotent stem cell (hiPSC)-derived cortical neurons treated with oligomeric amyloid-beta (oAβ) to replicate AD-like conditions. TTLL1, TTLL4, TTLL6 were selectively knocked down (KD) to assess their impact on TAU missorting and microtubule stability. Fluorescence resonance energy transfer (FRET) microscopy was used to examine interactions between TAU and TTLL proteins. We observed TAU missorting, increased tubulin polyglutamylation, decreased microtubule stability, and synaptic declustering in oAβ-treated neurons. TTLL1 KD significantly reduced TAU missorting, tubulin polyglutamylation, and

synaptic disintegration, while TTLL4 KD showed moderate effects, and TTLL6 KD restored

microtubule acetylation. Importantly, TTLL KD did not impair neuritic networks, dendritic

complexity, or neuronal activity. FRET microscopy revealed a potential interaction between TAU

and TTLL1, but not other TTLLs, suggesting a direct role of TTLL1 in TAU-mediated toxicity. Our

findings indicate that targeting TTLL1, either alone or in combination with other TTLLs, may be a

promising therapeutic strategy to counteract microtubule and synaptic dysfunction in AD and

related neurodegenerative disorders.

Keywords: TAU, TTLL, Alzheimer disease, hiPSCs, microtubules, synapses

Introduction

Microtubules are cylindrical filamentous heterodimers of α- and β-tubulin that form an essential

part of the cytoskeleton. Microtubules play an important role in maintaining neuronal shape and

facilitating the transportation of organelles and vesicles across neuronal networks (Sakakibara et

al., 2013). Microtubule dynamics and functions are tightly regulated by a complex set of

posttranslational modifications (PTMs) (Janke and Kneussel, 2010), as well as binding to and

interacting with proteins like microtubule-associated proteins (MAPs) and microtubule-severing

enzymes such as spastin (Goodson and Jonasson, 2018). In Alzheimer disease (AD),

microtubules are significantly depleted in affected brains, though the underlying causes remain

unclear (Cash et al., 2003; Jean and Bass, 2013).

TAU, a microtubule-associated protein encoded by the *MAPT* gene, is an axonal-enriched protein

that binds microtubules and promotes their assembly and stability. However, in tauopathies like

AD, TAU dissociates from microtubules, missorts to the somatodendritic compartments, and

aggregates into hyperphosphorylated neurofibrillary tangles, leading to microtubule fragmentation

and neuronal death (Zempel and Mandelkow, 2015). Spastin, an ATP-dependent enzyme

encoded by the *SPAST* gene, mediates microtubule severing, and its activity is catalyzed by a

specific PTM of microtubules called polyglutamylation, which comprises the addition of a

glutamate side chain to a glutamate residue, usually on the C-terminal tail of tubulin (Lacroix et

al., 2010). This modification is carried out by several members of a class of enzymes known as

Tubulin-Tyrosine-Ligase-Like proteins (TTLLs), mainly TTLL 1, 4, 5, 6, 7 and 11 (Magiera et al.,

2018).

Polyglutamylation is reversible, with cytosolic carboxypeptidases (CCPs) removing glutamate

side chains (Rogowski et al., 2010). The balance between adding and removing glutamate chains

is essential for health, with hyperglutamylation linked to neurodegeneration. For instance, Purkinje

cell degeneration (pcd) mice, in which CCP1 is deficient, exhibit severe neurodegeneration, which

is attenuated by knocking out TTLL1 and TTLL4, but not TTLL5 or TTLL7 (Li et al., 2020; Wu et

al., 2022). Another hint towards potential involvement of TTLLs in neurodegeneration came from

primary rat neurons treated with oligomeric amyloid-beta (oAβ), where missorted TAU appeared

to recruit TTLL6 to the somatodendritic compartments, inducing polyglutamylation and spastin-

mediated microtubule severing, leading to extensive microtubule loss (Zempel et al., 2010, 2013).

However, the underlying and potentially druggable disease mechanisms by which TAU missorting

triggers microtubule dysfunction, and the specific TTLLs involved, remain unclear in human

disease-relevant models.

In this study, we first aimed to establish a human tauopathy-relevant model using human induced

pluripotent stem cell (hiPSC)-derived cortical neurons (iNeurons). We treated these neurons with

oAβ and observed TAU missorting, increased tubulin polyglutamylation, and synaptic

declustering. We then aimed to identify the TTLL(s) responsible for mediating the pathological

effects of missorted TAU via individually knocking down TTLL1, TTLL4, or TTLL6. We show that

decreased expression of TTLL1, and to some extent TTLL4, alleviate the pathological effects of

oAβ without disturbing neuritic network, dendritic morphology, or neuronal function. We provide

first evidence that targeting TTLL1 alone or in combination with another TTLL is a potential

therapeutic target in AD and other tauopathies, and should be further validated and investigated.

Methods

hiPSC maintenance

WTC11 cells with a doxycycline-inducible Ngn2 transgene (Miyaoka et al., 2014; Wang et al.,

2017) were cultured on Geltrex-coated plates (Thermofisher Scientific #A1413302) at 37°C, 5%

CO_2 and regularly passaged when almost fully confluent using Versene (Thermofisher Scientific

#15040066) and thiazovivin-supplemented StemMACS iPS-Brew X.F. (Axon Medchem #1535,

Miltenyi Biotec #130-104-368) for the first 24 hours (Buchholz et al., 2024).

Differentiation of hiPSCs into cortical neurons (iNeurons)

Differentiation of hiPSCs into cortical neurons was carried out as described before (Buchholz et

al., 2024; Bachmann et al., 2021; Wang et al., 2017). At the start of differentiation, iPSCs were

harvested using Accutase (Sigma-Aldrich #A6964-100ML) and seeded onto Geltrex-coated

plates pre-differentiation medium (Thermofisher Scientific #12660012) supplemented with

thiazovivin (day before differentiation: d−3). The medium was changed every day for 2 days to

fresh pre-differentiation medium without supplementation. On day 0, 50,000 or 300,000 cells were

seeded onto Poly-D-Lysine- (Sigma-Aldrich #P7886-50MG) and Laminin- (Trevigen #3446-005-

01) coated 24-well-plates or 6-well plates respectively using maturation medium supplemented

with 1:100 GelTrex. Half of the media was exchanged once per week until analysis.

oAβ preparation and treatment

Aβ was prepared and reconstituted into oligomers as described before (Zempel et al., 2013).

Briefly, Aβ40 and Aβ42 powder (rPeptide #A-1153-1 and #A-1163-2) were completely dissolved

in Hexafluoro-2-propanol (HFIP) to a final concentration of 1 mM. Following aliquotation, HFIP

was evaporated completely using a vacuum concentrator, and the lyophilized powder was stored

at -80°C. On the day of treatment of iNeurons, lyophilized Aβ40 and Aβ42 were redissolved in 50

mM NaOH and mixed to produce a Aβ40/Aβ42 ratio of 7:3, and then diluted with PBS and 50 mM

HCl to a final concentration of 100 μM. To induce oligomerization, Aβ mixture was incubated at

37°C for one hour. Subsequently, iNeurons at day 21 were treated with 1 μM oAβ for 3 hours and

analyzed. Fig. S1 shows oAβ-treated iNeurons highlighting the distribution and targeting of the

oligomers.

Short hairpin RNA sequences

For the knockdown of human *TTLL*s in iNeurons, short hairpin RNA (shRNA) oligonucleotides

were inserted into pLKO.3G vector (Addgene #14748) resulting in a multi-cistronic lentiviral

construct expressing Green fluorescent protein (GFP) and the corresponding shRNA. The

following shRNA sequences were used to target human *TTLL*s or as a control:

Scrambled shRNA (control): 5´-TTGTCTTGCATTCGACTAA-3´

sh*TTLL1*: 5´-GTTTGTGTCTCAATCTAATAA-3´

sh*TTLL4*: 5´-GAGCCTTGGCAATAAGTTC-3´

sh*TTLL6*: 5´-CGGACUCATGAUUUCCAGGATT-3´, 5´-AACAACUCCCUCUUCCAGAAU-3´

Lentiviral-based knockdown of TTLLs

Lentivirus particle production and subsequent lentiviral transduction of iNeurons are described in

detail in Buchholz et al., 2024. Briefly, HEK293T cells were co-transfected with the corresponding

pLKO.3G plasmid, the packaging plasmid psPAX, and the envelope plasmid pMD2.G (Addgene

#12259 and #12260). Four and five days after transfection, the culture supernatant containing

lentivirus was collected, filtered and stored at -80°C. iNeurons were transduced with the lentiviral

particles on day 10 and analyzed 11 days after transduction (day 21).

Western blot analysis

For Western blot analysis, iNeurons were lysed in RIPA buffer (Sigma Aldrich #R0278),

centrifuged at 16,000 ×g for 20 minutes at 4°C, diluted in 5× Laemmli buffer, boiled for 10 minutes

at 95°C, and then separated on 10% Sodium dodecyl sulfate (SDS)-polyacrylamide gels.

Afterwards, proteins were transferred to polyvinylidene fluoride (PVDF) membranes overnight at

4°C, and blocked in 5% bovine serum albumin (BSA) in Tris-buffered saline with 0.1% Tween

(TBS-T). Membranes were incubated with the primary antibody overnight at 4°C, washed three

times with TBS-T, and incubated with the corresponding secondary horseradish peroxidase

(HRP)-coupled antibody for 1 hour at room temperature. After three washing rounds with TBS-T,

the immunoreactions were detected by applying the SuperSignal West Pico Chemiluminescent

Substrate (Thermofisher Scientific #34580) using a ChemiDoc XRS + system (Bio-Rad).

Immunofluorescence labeling of iNeurons

For immunocytochemistry, iNeurons were fixed with 3.7% Formaldehyde in PBS containing 4%

sucrose at room temperature for 30 minutes. Afterwards, cells were permeabilized and blocked

in 5% BSA (Carl Roth #8076.4) and 0.2% Triton X-100 (Carl Roth #3051.2) in PBS for 10 minutes.

After fixation, iNeurons were stained with primary antibodies at 4°C overnight. The following day,

coverslips were washed three times with PBS and stained with the corresponding secondary

antibodies coupled to an AlexaFluor dye for two hours at room temperature. Coverslips were then

washed with PBS and stained with NucBlue (Thermofisher Scientific #R37605) for 20 min at room

temperature, followed by mounting onto glass slides using Aqua-Poly/Mount (Polysciences

#18606-20). The slides were dried 24 hours at room temperature and then imaged.

Imaging

Immunostained iNeurons were imaged using a wide-field fluorescence microscope (Axioscope 5,

Zeiss) with ZenBlue Pro imaging software (V2.5, Zeiss). Images were analyzed using ImageJ

software (Version 2.14.0/1.54f, National Institutes of Health and the Laboratory for Optical and

Computational Instrumentation (LOCI), University of Wisconsin, USA). To measure the levels of

TAU or tubulin PTMs in neurons, regions of interest (ROIs) were manually delineated as profiles

of the soma or the dendrite where no other somas or processes overlapped, and the mean

fluorescence intensity (MFI) of each ROI was measured.

Neuronal network analysis

149   Fields of iNeurons cultures were imaged at 10x magnification after immunostaining for the axon-

150   enriched neurofilament light chain (NF-L) and the somatodendritic marker microtubule-associated

151   protein 2 (MAP2). The area of NF-L- or MAP2- positive neurites was calculated using ImageJ

152   software and normalized to the number of transduced nuclei in each field, identified via GFP

153   fluorescence.

Sholl analysis

Sholl analysis (Sholl, 1953) was used to investigate the complexity of dendritic arbouring in

iNeurons immunostained for MAP2. The center of the soma was designated as the center of

concentric circles, and the number of intersections was analyzed via the Neuroanatomy plugin in

ImageJ software. Sholl profiles were created by plotting the number of intersections against the

distance from the soma (µm). For statistical comparison, either the area under the curve (AUC)

or the full-width at half maximum of the Sholl profiles were compared. Data was tested for

normality by Shapiro-Wilk test and compared using a pairwise t-test. Analysis was performed

using R (R Core Team, 2024), AUC was calculated using the DescTools package (Signorell,

2024).

Microelectrode array measurements

For microelectrode array (MEA) measurements, iPSCs were seeded on MEA 24-well plates and

differentiated into cortical neurons as described above. At day 10, iNeurons were transduced with

lentiviral particles carrying the corresponding shRNA. Spontaneous activity was recorded for 2

minutes at day 21 at 37°C.

Fluorescence Resonance Energy Transfer (FRET) assay

Live-cell imaging-based FRET assay was carried out to assess interactions between TAU and

different TTLLs. HEK293T cells were co-transfected for 24 hours with teal fluorescent protein

(TFP)-TAU construct (FRET donor) and yellow fluorescent protein (YFP)-TTLL1, TTLL4, or TTLL6

constructs (FRET acceptor). HEK293T cells were also transfected for 24 hours with empty vectors

expressing TFP or YFP as controls. Live-cell imaging was performed with an inverted Leica DMi8

microscope with the help of Leica LAS X software (v3.7.3). FRET efficiency was analyzed via

FRET and Colocalization Analyzer plugin (Hachet-Hass et al., 2006) on ImageJ software.

Statistical analysis

The GraphPad Prism (v9.5.1, GraphPad Software, Boston, Massachusetts, USA) was used for

statistical analysis. Shapiro-Wilk test was performed to test for normal distribution of the data. In

case of normal distribution, statistical analysis was performed by unpaired t-test to compare the

means of two groups, or one-way ANOVA with correction for multiple comparisons (Tukey's test)

to compare three or more groups. When the data were not normally distributed, Mann–Whitney

U test or Kruskal-Wallis test with correction for multiple comparisons (Dunn's test) were carried

out, respectively. Statistical significance was denoted by a significance level of $P < 0.05$.

Antibodies

The antibodies used in this study are listed in Table 1.

Table 1. List of the primary and secondary antibodies used in the study.

Antibody	Animal species	Clonality	Cat#	Supplier	RRID	Use and dilution
Total TAU (K9JA)	rabbit	polyclonal	A0024	Agilent, USA	AB_1001372 4	ICC (1:1000)
Acetyl-α-Tubulin (Lys40) (D20G3)	rabbit	monoclon al	5335	Cell Signaling, USA	AB_1054446 94	ICC (1:500)
Anti-Tubulin Polyglutamylated antibody	mouse	monoclon al	T9822	Sigma-Aldrich, USA	AB_477598	ICC (1:500)
Anti-tyrosinated-α-Tubulin Antibody	rat	monoclon al	MAB1864-I	Sigma-Aldrich, USA	AB_289065 7	ICC (1:500)
GAPDH Antibody	mouse	monoclon al	sc-365062	Santa Cruz Biotechnology, USA	AB_1084780 62	WB (1:1000)
Anti-MAP2 Antibody	chicken	polyclonal	ab5392	Abcam, UK	AB_213815 3	ICC (1:2000)
Anti-NF-L antibody	rabbit	polyclonal	12998-1-AP	Proteintech, USA	AB_1059730 88	ICC (1:500)
Anti-Homer1 antibody	rabbit	polyclonal	12433-1-AP	Proteintech, USA	AB_229557 3	ICC (1:200)

Anti-synaptophysin antibody	mouse	monoclonal	67864-1-Ig	Proteintech, USA	AB_2918622	ICC (1:200)
Anti-TTLL1 antiibody	rabbit	polyclonal	PA5-27285	Thermofisher Scientific, USA	AB_2544761	WB (1:500)
Anti-TTLL4 antibody	rabbit	polyclonal	HPA027091	Sigma Aldrich, USA	AB_1060128	WB (1:500)
Anti-TTLL6 antibody	rabbit	polyclonal	PA5-100050	Thermofisher Scientific, USA	AB_2815580	WB (1:500)
Anti-Chicken Secondary Antibody, DyLight™ 350	goat	polyclonal	SA5-10069	Thermofisher Scientific, USA	AB_2556649	ICC (1:1000)
Anti-Rabbit Secondary Antibody, Alexa Fluor™ 488	donkey	polyclonal	A-21206	Thermofisher Scientific, USA	AB_2535792	ICC (1:1000)
Anti-Mouse Secondary Antibody, Alexa Fluor™ 568	goat	polyclonal	A-11031	Thermofisher Scientific, USA	AB_144696	ICC (1:1000)
Anti-Rabbit Secondary	donkey	polyclonal	A10042	Thermofisher Scientific, USA	AB_2534017	ICC (1:1000)

Antibody, Alexa Fluor™ 568						IHC (1:400)
Anti-Rat Secondary Antibody, Alexa Fluor™ 568	goat	polyclonal	A-11077	Thermofisher Scientific, USA	AB_2534121	ICC (1:1000)
Anti-Chicken Secondary Antibody, Alexa Fluor™ 647	goat	polyclonal	A21449	Thermofisher Scientific, USA	AB_2535866	ICC (1:1000)
Anti-Mouse Secondary Antibody, Alexa Fluor™ 647	donkey	polyclonal	A-31571	Thermofisher Scientific, USA	AB_162542	ICC (1:1000)
anti-Mouse Secondary Antibody, HRP	goat	polyclonal	115-035-003	Jackson ImmunoResearch Labs, USA	AB_10015289	WB (1:1000)
Anti-Rabbit Secondary Antibody, HRP	goat	polyclonal	7074	Cell Signaling, USA	AB_2099233	WB (1:1000)

Results

Establishment of Aβ-induced TAU pathology in iNeurons

Previous insights into the potential connection between TAU missorting, microtubule instability,

and TTLLs were obtained from rodent-derived models, while a representative human neuronal

model was lacking. In order to establish a human tauopathy-relevant model to study TAU-based

effects on microtubules, human iPSCs were differentiated into cortical neurons. Briefly, genetically

modified WTC11 iPSC line harboring a doxycycline-inducible neurogenin-2 (Ngn2) transgene

were induced to differentiate into pure glutamatergic neuronal cultures (See: Methods in this

paper; Wang et al., 2017; Buchholz et al., 2024).

Day 21 iNeurons treated with 1 μM oAβ showed increased levels of somatic TAU indicative of

TAU missorting (Fig. 1A, E). This was accompanied by a marked increase in the levels of tubulin

polyglutamylation (Fig. 1B, F). Interestingly, levels of polyglutamylated tubulin remained

unchanged in *MAPT* knockout iNeurons (characterized in Buchholz et al., 2025, Fig. S2). To

further evaluate the stability of microtubules in oAβ-treated iNeurons, we investigated the levels

of acetylated and tyrosinated tubulin, and observed reduced acetylation in the dendrites and

decreased tyrosination in the somatic compartments (Fig. 1C, G, H), indicating that microtubule

stability and dynamics were further compromised.

We also wanted to test the effects of oAβ on the synaptic integrity in our iNeurons. To this end,

we triple stained Homer1, synaptophysin, and MAP2 to identify dendritic synapses and quantified

the size and mean fluorescence intensity of synaptophysin-colocalized Homer1. While the

fluorescence intensity of synaptic Homer1 clusters did not change, the size of these clusters was

reduced upon oAβ treatment, indicating synaptic destabilization (Fig. 1D, I). Hence, oAβ treatment

in iNeurons causes TAU missorting, microtubule instability, and synaptic instability, making these

neurons a suitable human model for our study.

Reduction of TTLL1 and TTLL4 expression attenuates oAβ toxicity and TAU missorting

To identify the specific TTLL(s) driving pathological microtubule polyglutamylation upon TAU

missorting, we aimed to knock down three polyglutamylating TTLLs and then to observe the

effects of subsequent oAβ insult on the levels of polyglutamylated tubulin, and on the other toxicity

read-outs established above.

TTLL1, TTLL4, and TTLL6 were individually knocked down using shRNA-lentiviral transduction

of Day 10 iNeurons. Effective knockdown was confirmed via western blotting at Day 21, with

residual expression of targeted TTLLs reduced to 20-50% of the control (iNeurons transduced

with viruses carrying scrambled shRNA) (Fig. 2B).

Following the establishment of efficient lentiviral-based knockdown of TTLL1, TTLL4, and TTLL6,

the effects of each of these knockdowns on oAβ-induced toxicity was investigated. Day10

iNeurons were transduced with the knockdown viruses or scrambled control. At Day 21,

transduced iNeurons were treated with either oAβ or a vehicle control for 3 hours, followed by

fixation and staining as above. (Fig. 2A). Interestingly, oAβ-induced TAU missorting and tubulin

polyglutamylated tubulin were significantly reduced by TTLL1 knockdown to near normal levels,

with TTLL4 knockdown showing a similar effect on polyglutamylation but only a partial effect on

TAU missorting. On the other hand, TTLL6 knockdown significantly restored acetylated tubulin

levels but did not impact TAU missorting or tubulin polyglutamylation. However, the knockdown

of none of these TTLLs was sufficient to counteract the decrease of tyrosinated tubulin induced

by oAβ treatment (Fig. 2C-D).

While neither TTLL1 nor TTLL4 knockdowns fully restored synaptic Homer1 cluster size, they

both prevented significant cluster disassembly by 15-30 % upon oAβ insult, a protective effect not

observed with TTLL6 knockdown (Fig. 3A-B). This indicates that knocking down TTLL1 and

TTLL4 in iNeurons reduced oAβ-induced TAU missorting, tubulin polyglutamylation, and partially

protected synapses, while TTLL6 restored acetylated tubulin but did not affect TAU missorting or

synaptic protection, with none preventing tyrosinated tubulin loss.

Knockdown of TTLLs does not impair neuritic networks or neuronal function

To assess the broader impact of TTLL knockdown on neuronal networks, morphology, and function, TTLL1, TTLL4, or TTLL6 were knocked down at Day 10 and the effects on neuritic networks and dendritic branching were investigated at Day 21. Axonal and dendritic networks were studied via staining of NF-L and MAP2, respectively.

No significant changes were observed in the overall axonal (NF-L staining) or dendritic (MAP2 staining) networks across TTLL knockdown conditions compared to control (Fig. 4A-D). Sholl analysis revealed no significant differences in dendritic complexity although we noted a trend towards longer dendrites in iNeurons with TTLL1 and TTLL6 knockdowns (Fig. 4E-H). Additionally, neuronal activity, measured by microelectrode array (MEA) recordings, showed no significant alterations in spike rate or burst count across the different knockdown conditions (Fig. 4I). This indicates that iNeurons tolerate individual knockdown of the TTLLs studied here without obvious impairments in neuronal morphology and function.

TTLL1 and TAU interact in HEK293T cells

To investigate the mechanistic basis behind the ameliorating effects of TTLL knockdowns against oAβ-induced toxicity and test molecular interaction and proximity, we applied Fluorescence Resonance Energy Transfer (FRET) live-cell imaging to explore potential interactions between TAU and TTLLs. TFP-TAU and YFP-TTLL1, YFP-TTLL4, or YFP-TTLL6 were co-expressed in HEK293T cells. Co-expression of TFP-TAU and YFP alone, or YFP-TTLL1, YFP-TTLL4, or YFP-TTLL6 and TFP alone, served as negative controls.

Cells co-transfected with TFP-TAU and YFP-TTLL1 showed significantly higher FRET efficiency compared to the corresponding negative controls, suggesting direct interaction between TAU and

261   TTLL1. In contrast, no significant FRET signal was observed for TTLL4 or TTLL6 compared to

262   their controls (Fig. 5).

Discussion

This study presents a novel in vitro model of Aβ-induced tauopathy and microtubule impairments

using human iPSC-derived cortical neurons. The model effectively recapitulates key features of

TAU pathology seen in AD, including TAU missorting, microtubule destabilization, and synaptic

defects. By exploring the role of TTLL proteins in this context, we provide new insights into the

molecular mechanisms underlying Aβ-induced tauopathy and identify potential therapeutic

targets.

oAβ are thought to be the upstream disease-causing agent in AD (Cline et al., 2018). Previously,

acute oAβ treatment of rat-derived primary neurons led to TAU missorting, decreased acetylation

and increased polyglutamylation of tubulin (Zempel et al., 2013). In this study, we showed oAβ-

induced TAU missorting to the soma in our human iNeurons, a hallmark of early tauopathy (Thies

and Mandelkow, 2007). This pathological shift in TAU localization was accompanied by

decreased tubulin acetylation and tyrosination, and increased polyglutamylation. Microtubules are

regulated by a complex set of PTMs that govern their dynamics and stability. Decreased

acetylation for instance is a well-established marker of unstable microtubules (Maruta et al.,

1986), while decreased tyrosination negatively affects neuronal polarity and neurite outgrowth

(Erck et al., 2005). Polyglutamylation on the other hand recruits the microtubule-severing enzyme

spastin (Lacroix et al., 2010), with hyperglutamylation linked to neurodegeneration and neuronal

death (Berezniuk et al., 2012). In addition, we observed a significant decrease in the size of

synaptophysin-juxtaposing Homer1 clusters following oAβ treatment. Homer1 is a post-synaptic

density scaffold protein that is classified as an immediate early gene and which is induced by

neuronal activity (Clifton et al., 2019). Synaptic Homer1 cluster disassembly was shown before to

be driven by Aβ in primary rat neurons, which leads to the loss of synaptic structure and function (Roselli et al., 2009). Taken together, oAβ-insulted iNeurons exhibit detrimental changes in the PTMs of their microtubules, alongside TAU missorting and synaptic declustering, all of which is reminiscent of AD and related tauopathies.

We decided to knock down several glutamylating TTLLs in order to pinpoint the one responsible for pathological polyglutamylation, and to observe whether abolishing it would mitigate the harmful effects of oAβ. We opted to focus on TTLL1, TTLL4, and TTLL6 because of their expression levels and previously described pathological relevance: TTLL1 is the major polyglutamylating TTLL in the brain (Janke et al., 2005); TTLL1 and TTLL4 depletion mitigated neurodegeneration in pcd mice, a model that mainly exhibits adult-onset degeneration of cerebellar Purkinje neurons and selected thalamic neurons (Wu et al., 2022); and TTLL6 translocated to dendrites of primary neurons exposed to oAβ where it mediated polyglutamylation, spastin recruitment, and microtubule loss (Zempel et al., 2013). In our study, TTLL1 knockdown significantly protected iNeurons against oAβ-induced TAU missorting, reduced elevated levels of tubulin polyglutamylation, and partially alleviated the dissociation of synaptic clusters, suggesting that TTLL1 plays a critical role in the early stages of Aβ-induced tauopathy. On the other hand, overexpression of TTLL1 in iNeurons resulted in elevated levels of somatic TAU reminiscent of TAU missorting (Fig. S3), highlighting a potential toxic feedback loop between the two proteins. TTLL4 knockdown managed to decrease polyglutamylation levels but its effect on TAU missorting were less pronounced, indicating a rather secondary role for TTLL4. Interestingly, TTLL6 knockdown did not significantly affect TAU missorting but was the only knockdown that restored microtubule acetylation, hinting at a potential compensatory mechanism that stabilizes microtubules independently of TAU.

Polyglutamylating TTLLs are important enzymes that regulate neuronal microtubule organization, dynamics, and interactions with other proteins (Lacroix et al., 2010; Bodakuntla et al., 2020;

Genova et al., 2023). Therefore, we speculated that knockdown of different TTLLs could impact

neuronal morphology, networks, or activity, limiting the possibility of targeting TTLLs

therapeutically. To investigate this further, we decided to study dendritic network and dendritic

branching by staining for MAP2, a long-established dendritic marker (Caceres et al., 1984), and

axonal network by staining for NF-L, an intermediate filament highly concentrated in axons and a

biomarker of neuro-axonal damage (Khalil et al., 2024). However, neither neuritic networks nor

dendritic branching were affected by knockdown of any of the three TTLLs investigated in this

study. Additionally, burst count and spike rate of iNeurons measured by MEA recordings were

also not impacted by TTLL depletion, indicating maintained neuronal activity and making TTLLs

an attractive therapeutic target for reducing TAU pathology while preserving neuronal health.

In order to understand the mechanism through which TTLL1 could mediate the detrimental effects

of oAβ insult and TAU missorting, we decided to investigate potential interactions between TFP-

TAU and YFP-tagged constructs of the three TTLLs investigated in this study via FRET

microscopy in HEK293T cells. It was reported before that TAU and TTLL6 overexpressed in

HEK293T cells may interact directly (Zempel et al., 2013). In this study however, only cells

expressing TFP-TAU and YFP-TTLL1 showed FRET efficiency values higher than negative

controls. FRET phenomena can only occur when the distance between donor (TFP) and acceptor

(YFP) is less than 10 nm, which indicates that TAU and TTLL1 are in close proximity and may

interact directly with each other. This may explain why the most effective protection against oAβ

was observed in iNeurons with TTLL1 knockdown. We think that missorted TAU following oAβ

insult transports TTLL1 to the somatodendritic compartments where it polyglutamylates

microtubules, culminating in decreased microtubule stability and synaptic loss.

Conclusion

We showed that human iPSC-derived neurons subjected to oAβ suffer from TAU missorting,

stability-decreasing changes of microtubule PTMs, and dissociation of synaptic clusters. These

noxious effects are significantly reduced via TTLL1 knockdown, without affecting neuronal

networks and activity. Our findings suggest that TTLL1, alone or with combination with another

TTLL, could be a promising therapeutic target for preventing or slowing the progression of

tauopathy in AD. Future studies should explore the therapeutic potential of genetic or

pharmacological targeting of TTLL1 in vivo and investigate the broader implications of TTLL-

mediated microtubule modifications in neurodegenerative diseases.

List of Abbreviations

AD: Alzheimer disease

AUC: Area under the curve

BSA: bovine serum albumin

CCP: cytosolic carboxypeptidases

FRET: Fluorescence Resonance Energy Transfer

GFP: Green fluorescent protein

HFIP: Hexafluoro-2-propanol

HRP: horseradish peroxidase

hiPSCs: Human induced pluripotent stem cells

iNeurons: hiPSC-derived neurons

KD: Knockdown

MAP2: Microtubule-associated protein 2

MEA: Microelectrode array

MFI: Mean fluorescence intensity

NF-L: Neurofilament light chain

Ngn2: Neurogenin-2

oAβ: Oligomeric amyloid beta

Pcd: Purkinje cell degeneration

PTM: Posttranslational modification

PVDF: polyvinylidene fluoride

ROI: Region of interest

SDS: Sodium dodecyl sulfate

shRNA: short hairpin RNA

TBS-T: Tris-buffered saline with 0.1% Tween

TFP: Teal fluorescent protein

TTLL: Tubulin-Tyrosine-Ligase-Like

YFP: Yellow fluorescent protein

Declarations

Ethics approval and consent to participate: not applicable.

Consent for publication: not applicable

Availability of data and materials: The datasets used and/or analyzed during the current study are

available from the corresponding author on reasonable request.

<u>Competing interests:</u> The authors have no relevant financial or non-financial interests to disclose.

<u>Funding:</u> This study was supported by the Alzheimer Forschung Initiative e.V. (grant #22039, to
HZ).

<u>Authors' contributions:</u> Study design: MAAK, HZ. Experimental work: MAAK. Methodological
support: TW, JK. Data analysis and interpretation: MAAK, DA, HZ. Manuscript writing: MAAK.
Manuscript commenting and proofreading: HZ, DA, TW, JK. All authors read and approved the
final manuscript.

<u>Acknowledgments:</u> We thank Prof. Dr. Li Gan (Weill Cornell Medicine, NY, USA) for providing
Ngn2-WTC11 iPSCs. We thank Prof. Dr. Florian Klein (Institute of Virology, University Hospital
Cologne) for providing lentiviral vectors. We thank Dr. Sarah Buchholz (current address: Max
Planck Institute for Biology of Ageing, Cologne, Germany) for the stimulating discussions and
helpful insights. Stem cell work was performed at the iPSC-Lab core facility of the CMMC
(Cologne, Germany).

References

Bachmann S, Linde J, Bell M, Spehr M, Zempel H, Zimmer-Bensch G. DNA Methyltransferase 1
(DNMT1) shapes neuronal activity of human iPSC-derived glutamatergic cortical neurons. Int J
Mol Sci. 2021;22(4):2034. doi: 10.3390/ijms22042034.

Berezniuk I, Vu HT, Lyons PJ, Sironi JJ, Xiao H, Burd B, et al. Cytosolic carboxypeptidase 1 is
involved in processing α- and β-tubulin. J Biol Chem. 2012;287(9):6503-17. doi:
10.1074/jbc.M111.309138.

Bodakuntla S, Schnitzler A, Villablanca C, Gonzalez-Billault C, Bieche I, Janke C, et al. Tubulin

polyglutamylation is a general traffic-control mechanism in hippocampal neurons. J Cell Sci.

2020;133(3). doi: 10.1242/jcs.241802.

Buchholz S, Bell-Simons M, Cagmak C, Klimek J, Gan L, Zempel H. Cultivation, differentiation,

and lentiviral transduction of human induced pluripotent stem cell (hiPSC)-derived glutamatergic

neurons for studying human TAU. In: TAU protein: Methods and Protocols. Springer; 2024.

Buchholz S, Al Kabbani MA, Bell-Simons M, Kluge L, Cagmak C, Klimek J, Haag N, Iohan LC,

Coulon A, Costa MR, Kilinc D, Zempel H. The tau isoform 1N4R confers vulnerability of MAPT

knockout human iPSC-derived neurons to amyloid beta and phosphorylated tau-induced neuronal

dysfunction. Alzheimers Dement. 2025;21(5):e14403. doi: 10.1002/alz.14403.

Caceres A, Banker G, Steward O, Binder L, Payne M. MAP2 is localized to the dendrites of

hippocampal neurons which develop in culture. Brain Res. 1984;315(2):314-8. doi: 10.1016/0165-

3806(84)90167-6.

Cash AD, Aliev G, Siedlak SL, Nunomura A, Fujioka H, Zhu X, et al. Microtubule reduction in

Alzheimer's disease and aging is independent of TAU filament formation. Am J Pathol.

2003;162(5):1623-7. doi: 10.1016/s0002-9440(10)64296-4.

Clifton NE, Trent S, Thomas KL, Hall J. Regulation and function of activity-dependent Homer in

synaptic plasticity. Mol Neuropsychiatry. 2019;5(3):147-61. doi: 10.1159/000500267.

Cline EN, Bicca MA, Viola KL, Klein WL. The Amyloid-β oligomer hypothesis: Beginning of the

third decade. J Alzheimers Dis. 2018;64(s1). doi: 10.3233/JAD-179941.

Erck C, Peris L, Andrieux A, Meissirel C, Gruber AD, Vernet M, et al. A vital role of tubulin-tyrosine-

ligase for neuronal organization. Proc Natl Acad Sci U S A. 2005;102(22):7853-8. doi:

10.1073/pnas.0409626102.

Genova M, Grycova L, Puttrich V, Magiera MM, Lansky Z, Janke C, et al. Tubulin

polyglutamylation differentially regulates microtubule-interacting proteins. EMBO J. 2023;42(5).

doi: 10.15252/embj.2022112101.

Goodson HV, Jonasson EM. Microtubules and microtubule-associated proteins. Cold Spring Harb

Perspect Biol. 2018;10(6). doi: 10.1101/cshperspect.a022608.

Hachet-Haas M, Converset N, Marchal O, Matthes H, Gioria S, Galzi JL, et al. FRET and

colocalization analyzer--a method to validate measurements of sensitized emission FRET

acquired by confocal microscopy and available as an ImageJ Plug-in. Microsc Res Tech.

2006;69(12):941-56. doi: 10.1002/jemt.20376.

Janke C, Rogowski K, Wloga D, Regnard C, Kajava AV, Strub JM, et al. Tubulin polyglutamylase

enzymes are members of the TTL domain protein family. Science. 2005;308(5729):1758-62. doi:

10.1126/science.1113010.

Janke C, Kneussel M. Tubulin post-translational modifications: encoding functions on the

neuronal microtubule cytoskeleton. Trends Neurosci. 2010;33(8):362-72. doi:

10.1016/j.tins.2010.05.001.

Jean DC, Baas PW. It cuts two ways: microtubule loss during Alzheimer disease. EMBO J.

2013;32(22):2900-2. doi: 10.1038/emboj.2013.219.

Khalil M, Teunissen CE, Lehmann S, Otto M, Piehl F, Ziemssen T, et al. Neurofilaments as
biomarkers in neurological disorders - towards clinical application. Nat Rev Neurol.
2024;20(5):269-87. doi: 10.1038/s41582-024-00955-x.

Lacroix B, van Dijk J, Gold ND, Guizetti J, Aldrian-Herrada G, Rogowski K, et al. Tubulin
polyglutamylation stimulates spastin-mediated microtubule severing. J Cell Biol.
2010;189(6):945-54. doi: 10.1083/jcb.201001024.

Li J, Snyder EY, Tang FH, Pasqualini R, Arap W, Sidman RL. Nna1 gene deficiency triggers
Purkinje neuron death by tubulin hyperglutamylation and ER dysfunction. JCI Insight. 2020;5(19).
doi: 10.1172/jci.insight.136078.

Magiera MM, Singh P, Gadadhar S, Janke C. Tubulin posttranslational modifications and
emerging links to human disease. Cell. 2018;173(6):1323-7. doi: 10.1016/j.cell.2018.05.018.

Maruta H, Greer K, Rosenbaum JL. The acetylation of alpha-tubulin and its relationship to the
assembly and disassembly of microtubules. J Cell Biol. 1986;103(2):571-9. doi:
10.1083/jcb.103.2.571.

Miyaoka Y, Chan AH, Judge LM, Yoo J, Huang M, Nguyen TD, et al. Isolation of single-base
genome-edited human iPS cells without antibiotic selection. Nat Methods. 2014;11:291-3. doi:
10.1038/nmeth.2840.

R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing; 2024. Available from: https://www.R-project.org/.

Rogowski K, van Dijk J, Magiera MM, Bosc C, Deloulme JC, Bosson A, et al. A family of protein-
deglutamylating enzymes associated with neurodegeneration. Cell. 2010;143(4):564-78. doi:
10.1016/j.cell.2010.10.014.

Roselli F, Hutzler P, Wegerich Y, Livrea P, Almeida OF. Disassembly of shank and homer

synaptic clusters is driven by soluble beta-amyloid(1-40) through divergent NMDAR-dependent

signalling pathways. PLoS One. 2009;4(6). doi: 10.1371/journal.pone.0006011.

Sakakibara A, Ando R, Sapir T, Tanaka T. Microtubule dynamics in neuronal morphogenesis.

Open Biol. 2013;3(7):130061. doi: 10.1098/rsob.130061.

Signorell A. DescTools: tools for descriptive statistics. R package version 0.99.54; 2024. Available

from: https://CRAN.R-project.org/package=DescTools.

Sholl DA. Dendritic organization in the neurons of the visual and motor cortices of the cat. J Anat.

1953;87(4):387-406.

Thies E, Mandelkow EM. Missorting of TAU in neurons causes degeneration of synapses that

can be rescued by the kinase MARK2/Par-1. J Neurosci. 2007;27(11):2896-907. doi:

10.1523/JNEUROSCI.4674-06.2007.

Wang C, Ward ME, Chen R, Liu K, Tracy TE, Chen X, et al. Scalable production of iPSC-derived

human neurons to identify TAU-lowering compounds by high-content screening. Stem Cell

Reports. 2017;9(4):1221-33. doi: 10.1016/j.stemcr.2017.08.019.

Wu HY, Rong Y, Bansal PK, Wei P, Guo H, Morgan JI. TTLL1 and TTLL4 polyglutamylases are

required for the neurodegenerative phenotypes in pcd mice. PLoS Genet. 2022;18(4). doi:

10.1371/journal.pgen.1010144.

Zempel H, Thies E, Mandelkow E, Mandelkow EM. Abeta oligomers cause localized Ca(2+)

elevation, missorting of endogenous TAU into dendrites, TAU phosphorylation, and destruction

of microtubules and spines. J Neurosci. 2010;30(36):11938-50. doi: 10.1523/JNEUROSCI.2357-

10.2010. Erratum in: J Neurosci. 32(17):6052.

Zempel H, Luedtke J, Kumar Y, Biernat J, Dawson H, Mandelkow E, et al. Amyloid-β oligomers

induce synaptic damage via TAU-dependent microtubule severing by TTLL6 and spastin. EMBO

J. 2013;32(22):2920-37. doi: 10.1038/emboj.2013.207.

Zempel H, Mandelkow EM. TAU missorting and spastin-induced microtubule disruption in

neurodegeneration: Alzheimer disease and hereditary spastic paraplegia. Mol Neurodegener.

2015;10:68. doi: 10.1186/s13024-015-0064-1.

504   **Figure 1:**

509 **Figure 2:**

511   **Figure 3:**

Figure 4:

Figure 5:

Figure 1. oAβ treatment induces TAU missorting, alters microtubule modifications, and disrupts synaptic clusters in iNeurons. A) Co-immunostaining of TAU and MAP2 in iNeurons treated with oAβ for 3 hours or untreated control (Ctrl). Insets display 4-fold magnifications of areas framed with dashed lines, highlighting TAU missorting into the somatic compartment. Scale bar: 10 μm. **B)** Co-staining of TAU and polyglutamylated tubulin (polyE-tub) in iNeurons treated with oAβ for 3 hours or untreated control. Insets show 4-fold magnifications of areas framed with dashed lines, emphasizing increased polyglutamylation in oAβ-treated neurons. Scale bar: 10 μm. **C)** Immunostaining of acetylated tubulin (ace-tub) and tyrosinated tubulin (tyr-tub) in iNeurons treated with oAβ for 3 hours or untreated control. Insets show 3-fold magnifications of areas framed with dashed lines, demonstrating reduced acetylation and tyrosination post-treatment. Scale bar: 10 μm. **D)** Co-immunostaining of Homer1 and synaptophysin (SYP) in iNeurons treated with oAβ for 3 hours or untreated control. Insets show 2-fold magnifications of the areas framed with dashed lines, illustrating the disassembly of synaptic clusters. Scale bar: 2 μm. **E)** Quantification of somatic TAU levels in (A). (N=3, n=45 neurons). **F)** Quantification of somatic polyE-tub levels in (B). (N=3, n=45 neurons). **G)** Quantification of somatic and dendritic ace-tub levels in (C). (N=3, n=45 neurons). **H)** Quantification of somatic and dendritic tyr-tub levels in (A). (N=3, n=45 neurons). **I)** Quantification of SYP-colocalizing Homer1 cluster size and mean fluorescence intensity (MFI) in (D). (N=3, n=150 puncta). ^{ns} non-significance, * P ≤ 0.05, ** P ≤ 0.01.

Figure 2. Knockdown of TTLL1 and TTLL4 partially ameliorates oAβ-induced pathological changes in iNeurons. A) Schematic representation of the experimental workflow, outlining the knockdown and treatment procedures. Neurons were transduced at Day 10 with shRNA, and then treated and fixed as indicated. **B)** Immunoblotting confirmation of successful knockdown of TTLL1, TTLL4, and TTLL6 in iNeurons. **C)** Immunostaining for TAU, acetylated tubulin (Ace-tub), polyglutamylated tubulin (PolyE-tub), tyrosinated tubulin (Tyr-tub) in Day 21 iNeurons transduced with scrambled RNA (Scr.) or shRNAs targeting TTLL1, TTLL4, or TTLL6 for 11 days, and treated with oAβ or vehicle control for 3 hours. Scale bar: 10 μm. **D)** Quantification of somatic TAU, dendritic ace-tub, somatic polyE-tub, somatic tyr-tub levels of (C). (N=3, n=45 neurons). ^{ns} non-significance, * P ≤ 0.05, ** P ≤ 0.01.

Figure 3. Knockdown of TTLL1 and TTLL4 sightly attenuates oAβ-induced synaptic declustering in iNeurons. A) Co-immunostaining of Homer1 and synaptophysin (SYP) in Day 21 iNeurons transduced with scrambled RNA (Scr.) or shRNAs targeting TTLL1, TTLL4, or TTLL6 for 11 days, and treated with oAβ or vehicle control for 3 hours. Insets show 2-fold magnifications of areas framed with dashed lines, highlighting changes in synaptic cluster size. Scale bar: 2 μm. **B)** Quantification of SYP-colocalizing Homer1 cluster size and mean fluorescence intensity (MFI) in (A). (N=3, n≈300 puncta). ^{ns} non-significance, * P ≤ 0.05.

Figure 4. TTLL Knockdown does not affect neuronal networks, morphology, or function. A-B) Immunostaining of neurofilament-L (NF-L) (A) or MAP2 (B) in Day 21 iNeurons transduced with scrambled RNA (Scr.) or shRNAs targeting TTLL1, TTLL4, or TTLL6 for 11 days. Scale bars: 200 μm. **C)** Quantification of the area covered by the axonal network (NF-L) in (A). (N=3, n=15 fields). ^{ns} non-significance. **D)** Quantification of the area covered by the dendritic network (MAP2) in (B). (N=3, n=15 fields). ^{ns} non-significance. **E)** Quantification of the number of dendritic intersections obtained from Sholl analysis. (N=3, n=15 neurons). ^{ns} non-significance. **F-H)** Sholl profiles showing dendritic branching complexity in Day 21 iNeurons transduced with scrambled RNA (Scr.) or shRNAs targeting TTLL1 (F), TTLL4 (G), or TTLL6 (H) for 11 days. (N=3, n=15 neurons). **I)** Quantification of spike rate and burst count from microelectrode array (MEA) recordings in Day 21 iNeurons transduced with scrambled RNA (Scr.) or shRNAs targeting TTLL1, TTLL4, or TTLL6 for 11 days. (N=3, n=12-15 wells). ^{ns} non-significance.

Figure 5. Fluorescence Resonance Energy Transfer (FRET) analysis reveals potential interaction between TAU and TTLL1. A) FRET microscopy images in HEK293T cells showing various controls (TFP-TAU+YFP, TFP+YFP-TTLL1, TFP+YFP-TTLL4, TFP+YFP-TTLL6) and experimental conditions (TFP-TAU+YFP-TTLL1, TFP-TAU+YFP-TTLL4, TFP-TAU+YFP-TTLL6). Images of the three detection channels: donor (TFP), acceptor (YFP), and FRET, as well as the spectral bleed-through (BT) corrected FRET are depicted. Scale bar: 10 μm. **B)** 7-fold magnification of the area framed by dashed lines in (A) showing enhanced FRET signal in low-expressing cells co-transfected with TFP-TAU and YFP-TTLL1. **C)** Quantification of FRET efficiency from the conditions shown in (A). (N=3, n=15 cells). [ns] non-significance, $* P \leq 0.05$, $*** P \leq 0.001$, $**** P \leq 0.0001$.