**Brain-Cognitive Gaps in relation to Dopamine and Health-related Factors: Insights**

**from AI-Driven Functional Connectome Predictions**

Morteza Esmaeili [1,2]

Erin Beate Bjørkeli[2,3]

Robin Pedersen[4,5,6]

Farshad Falahati[4,5,6]

Jarkko Johansson[4,8]

Kristin Nordin[4,5,6]

Nina Karalija[7,8]

Lars Bäckman[6]

Lars Nyberg[5,7,8]

Alireza Salami[4,5,6,7#]

*Author Affiliations*:

[1] Department of Electrical Engineering and Computer Science, University of Stavanger,

Stavanger, Norway

[2] Department of Diagnostic Imaging, Akershus University Hospital, Lørenskog, Norway

[3] Institute of Clinical Medicine, University of Oslo, Oslo, Norway

[4] Wallenberg Center for Molecular Medicine (WCMM), Umeå University, Umeå, Sweden

[5] Department of Medical and Translational Biology, Umeå University, Umeå, Sweden

[6] Aging Research Center, Karolinska Institute and Stockholm University, Solna, Sweden

[7] Umeå Center for Functional Brain Imaging (UFBI), Umeå University, Umeå, Sweden

[8] Department of Diagnostics and Intervention, Diagnostic Radiology, Umeå University,

Umeå, Sweden

Emails

A.S.: Alireza.Salami@ki.se

M.E.: Morteza.Esmaeili@uis.no

**#Correspondence to:**

*Email:* Alireza.Salami@ki.se

*Address: Aging Research Center; Tomtebodavägen 18a, 17165, Stockholm, Sweden*

*Phone:* +46790682161

**Word count:** 9678 including text of figure legends – reference is not included

**Running title:** AI-based Prediction of Cognitive Function

**Acknowledgments:** This work was funded by the Swedish Research Council (grant number

2016-01936 to A.S.), Knut and Alice Wallenberg Foundation (Wallenberg Fellow grant to

A.S.), bank of Sweden (RJ, grant number P20-0515 to A.S.), StratNeuro grant at Karolinska

Institute (A.S.). Morteza Esmaeili and Erin B. Bjørkeli were supported by the Southern

Eastern Norway Regional Health Authority (Helse Sør-Øst RHF, HSØ, ME: grant number

HSØ 2018047; E.B.B. HSØ 2021023).

**Author Contributions.** M.E., E.B.B., and A.S. (and COBRA PIs) study design, method

development, data measurement, and analysis. All authors participated in manuscript writing

or editing.

**Additional Information and Competing Interests Statement.** The authors have no

competing interests to declare.

**Data availability**

The scripts used for developing the model are available at https://github.com/MorEsm/AI-

based-Prediction-of-Cognitive-Function. (We are working on this channel to provide open-

source scripts).

**ABSTRACT**

A key question in human neuroscience is to understand how individual differences in brain

function are related to cognitive differences. However, the optimal condition of brain function

to study between-person differences in cognition remains unclear. Additionally, there is a lack

of objective biomarkers to accurately predict cognitive function, with brain age emerging as a

potential candidate. Recent research suggests that brain age offers minimal additional

information on cognitive decline beyond what chronological age provides, prompting a shift

toward approaches focused directly on cognitive prediction. Using a novel deep learning

approach, we evaluated the predictive power of the functional connectome during various states

(resting state, movie-watching, and n-back) on episodic memory and working memory

performance. Our findings show that while task-based connectomes, especially during movie

watching, better predict working memory, resting state connectomes are equally effective in

predicting episodic memory. Furthermore, individuals with a negative brain-cognition gap

(where brain predictions underestimate actual performance) exhibited lower physical activity,

lower education, and higher cardiovascular risk compared to those with a positive gap. This

shows that knowledge of the brain-cognition gap provides insights into factors contributing to

cognitive resilience. Further lower PET-derived measures of dopamine binding were linked to

a greater brain-cognition gap, mediated by regional functional variability. Together, our study

introduces the brain-cognitive gap, as a new marker, modulated by the dopamine system, to

identify individuals at risk of compromised brain function.

**Keywords:** Functional connectome, Cognition, Dopamine, Predictive modeling

**1. INTRODUCTION**

Resting state fMRI serves as a tool to map brain function (*1, 2*). Functional connectivity (FC)

at rest, estimated with methods to gauge correlated spontaneous activity between two or more

regions, serves as a measure of functional brain integrity (*1, 3*). Individual differences in FC at

rest are associated with differences in various cognitive domains (*4-6*). Associations of this

nature have been reported for several large-scale networks, including the default mode network

(DMN) (*7-9*), the frontoparietal network (*10-12*), the dorsal attention network (*13*), and the

salience network (*14*). Moreover, such associations have also been identified in more

comprehensive investigations spanning the entire functional brain repertoire of the brain (*15,*

*16*). Most previous reports focused on associations, not predictions, utilizing the correlation

between FC and behavioral phenotypes, which tend to overfit the data and therefore fail to

generalize (*17*). Proper cross-validation, preferably in an independent sample, is therefore

important to assert reliable population-level inferences (*18*). Recent machine learning-based

predictive frameworks offer powerful tools for assessing the predictability of individual

behavioral phenotypes based on brain connectivity (*19-23*). In particular, deep neural networks

(DNN) methods have been successfully applied to behavioral and disease prediction (*24-26*),

and have been found to outperform other machine learning approaches (*27-29*).

The functional connectome has demonstrated predictive utility regarding trait-like cognitive

phenotypes (*30-33*). The predictive-modeling framework of the functional connectome has

been applied to various cognitive domains, including fluid intelligence (*34*), working memory

(WM) (*35*), visuospatial ability (*36*), attention (*37*), creativity (*38*), as well as personality traits

(*39*). Understanding the patterns contributing to predictions could offer insights into the

functional organization underlying cognitive phenotypes, serving as biomarkers indicating

current or prospective health conditions (*40-42*). Moreover, the whole-brain functional

connectome acts as a fingerprint with accurate identification of subjects from a large population

(*43*) within the same cognitive state (e.g., rest-to-rest) but also across different states (e.g., rest-

to-task).  Overall, past research suggests that the functional connectome is relatively robust

within individuals, is unique across individuals, and can predict cognitive and personality

phenotypes.

Despite consensus on the value of resting state functional connectivity for mapping brain

function, there is an ongoing debate about whether rest is the optimal brain state for

investigating individual differences in neurocognitive function (*44-47*). A study using data

from the human connectome project has shown that resting state fMRI predicts differences in

brain activity during various tasks, including social, language, relational, and motor tasks (*48*).

This finding supports the notion that individual differences in neural activation can be predicted

from resting state (*46*). However, results from the same dataset revealed that task-based FC

outperforms resting state FC in predicting individual differences in fluid intelligence, with task-

based data explaining 20 % of the variance compared to just 6 % explained by rest (*49*).

Consistent with this result, previous studies have investigated trait- as well as state-dependent

FC, supporting the utility of an integrative approach (*11, 47*). More recent studies suggested

that naturalistic viewing, such as movie-watching, may serve as a happy medium between

unconstraint rest and overlay-constrained tasks in predicting behavior differences (*50, 51*).

Despite the presence of similar spatiotemporal activity patterns across individuals during

movie-watching (*52*), notable individual differences in activity and functional connectivity

(*53*) persist alongside these idiosyncratic features. This suggests that tasks which align

individuals' functional connectome more closely to an optimal level, neither completely

unconstrained like rest nor overly synchronized like a task, also render them easiest to identify

(*44*). Hence, it is plausible that FC during naturalistic paradigms improves sensitivity to predict

behavioral differences (*50, 54*).

In this study, we first assess the predictive efficacy of the functional connectome across various

states for two commonly used cognitive functions. Specifically, we focused on working

memory i.e., the ability to store and manipulate information over shorter durations of time, as

well as its integral involvement in problem-solving, reasoning, and other key components of

fluid intelligence (e.g., (*55-57*)). In contrast, episodic memory (EM) entails the recollection of

specific experiences and events (*58*), which is regarded as an important element of mind-

wandering (*59*) during resting state.

Past studies have used neuroimaging data, including resting state to predict brain age (*60-62*).

These studies show that brain age, based on biological phenotypes, and their deviation from

the chronological age (known as brain age gap prediction), could serve as a biomarker in

characterizing disease risk (*62-64*). Importantly, an older-appearing brain has been shown to

be associated with exacerbated physiological and cognitive aging, and risk of mortality (*61*).

However, a recent study suggests that brain age predictions contribute minimally compared to

chronological age for explaining cognitive decline (*65*), implying that cognitive predictions are

more reliable. Building upon this study and extending the concept of brain-age gap to brain-

cognition gap, it is plausible that a potential disconnection between the brain's biological state

and observed cognitive state might emerge. If the brain predicts lower performance than is

observed (i.e., a negative brain-cognition gap), it may be compensating for underlying issues

not yet apparent through cognitive assessments. By this view, individuals with negative brain-

cognition gaps should be less healthy than those whose brains predict higher cognitive function

than their actual performance (i.e., a positive brain-cognition gap). Considering the significance

of life-style and cardiovascular risk for maintaining healthy brain function (*66, 67*), the

secondary aim of this study was to assess whether individuals with positive versus negative

brain-cognitive gaps differ in terms of physical activity habits and cardiovascular risk factor

burden.

The third aim of the current study is to uncover the contribution of dopamine (DA) integrity to

brain-cognition gaps. Specifically, we investigate whether individuals with lower DA receptor

levels exhibit higher functional variability, and in turn greater brain-cognition gaps. DA is a

vital neuromodulator with critical implications for motor function, reward-seeking behavior,

and various higher-order cognitive functions (*68-75*). Insufficient DA modulation can affect

neurocognitive functions detrimentally (*69, 74, 76-78*). DA is important for enhancing the

signal-to-noise ratio (SNR) of neural transmission (*79, 80*), and insufficient modulation may

contribute to increased variability in blood-oxygen-level-dependent (BOLD) signal (*81, 82*)

and reduced coherence in the functional connectome (*83*). Indeed, pharmacological studies

have shown that DA depletion increases the variability of the BOLD signal, subsequently

leading to less synchronized connectivity within resting state networks (*84*). Consequently, we

expect individuals with inadequate DA levels to exhibit increased regional signal variability, a

less unique functional connectome, and greater brain-cognition gaps.

Using data from the DopamiNe, Age, connectoMe, and Cognition (DyNAMiC) study (n =180,

20-79 years, 50% female) (*54*) (**Fig. 1**), we evaluated the predictive power of the functional

connectome during resting state, movie-watching, and n-back tasks for two different cognitive

domains: episodic memory and working memory. Based on recent research indicating that

movie-watching enhances predictability by highlighting key features of FC (*50*), we

hypothesized that FC during movie-watching would outperform FC during rest, and possibly

during task, in predicting both cognitive measures. To achieve this objective, we employed a

deep neural network approach, a specific subtype of artificial intelligence (AI), to predict

cognitive scores from the functional connectome. Deep learning approaches overcome the

limitation of predictive techniques that solely rely on linear associations between connectivity

and behavioral phenotypes (*17*). Considering the importance of individual characteristics, such

as age, in predicting behavior from FC (*33*), we conducted cross-validation of our model,

initially derived from an age-heterogeneous sample, in an age-homogeneous sample (from the

Cognition, Brain, and Aging (COBRA) study (*85*)). We subsequently investigated whether

individuals with positive prediction gaps differ from those with negative gaps in terms of

lifestyle and cardiovascular disease risk factors. Moreover, we tested the hypothesis of whether

individuals with lower striatal dopamine D1-like receptor availability (D1DR), the brain's most

abundant DA receptor subtype, have a less distinctive FC pattern (i.e., more regional

variability) and, in turn, a larger brain-cognition gap. Finally, we conducted a cross-validation

of the link between DA and the prediction gap in an independent cohort with estimates of DA

D2-like receptor (D2DR) availability (*85*).

**2. Results**

***2.1. AI-Driven Predictive Modeling of Cognition Scores from the Functional Connectome***

We used fMRI data from the DyNAMiC project, in which each subject underwent rest, movie-

watching, and working memory n-back tasks. These data were parcellated into 273 nodes (264

with 9 additional subcortical nodes) using a previously published whole-brain functional atlas

(*86*). The averaged time series of 273 regions were subsequently correlated, to create the FC

matrix for each participant and cognitive state (rest, movie-watching, and n-back). We found

FC to be significantly correlated between cognitive states ($r^2$ for rest versus movie-watching

=.81; $r^2$ for rest versus n-back =.76; $r^2$ for movie-watching versus n-back =.71), suggesting that

a common functional architecture is present across various tasks.

We trained a convolutional neural network, DenselyAttention, derived from DenseNet (*87*) on

FC matrices from each condition (resting state, movie-watching, and n-back) to generate

cognition-specific prediction models of two memory domains (EM and WM). Model

performance was quantified as the correlation between observed and predicted cognitive

performance. Each model was then tested on all three conditions to examine the

generalizability of each model across cognitive states. For example, the model trained on the

resting state data (orange circles in **Fig. 1**) was used to predict EM scores using the test dataset

derived from resting state (**Fig. 1a**), movie-watching, and n-back. Significance of our main

predictions was assessed via linear correlations, and uncorrected *p* values are presented in

**Tables 1-2**.

**220   *2.2. Resting state and movie-watching models outperform n-back in episodic-memory***

**221   *prediction, with resting state offering the best generalizability***

We first started with all cases in which congruent conditions were used for model building and

prediction. Only models derived from the rest and movie-watching datasets yielded significant

predictions of EM (**Figs. 1a-b** and **Table 1**), with resting state yielding the best-performing

model ($r^2 =25.0\%$, $p <.0001$) followed by movie-watching ($r^2 =24.0\%$, $p <.0001$) (**Table 1**).

While there was no significant difference between resting state and movie watching in

predicting episodic memory (Mann–Whitney $U$ test, $p =.17$), both models yielded better EM

prediction than n-back (Mann–Whitney $U$ test, $p <.0001$; **Fig. 1e**).

To test the generalizability of these models, two types of validation analyses were performed:

cross-condition and cross-data set. In the cross-condition analysis, models trained on one

condition (e.g., rest) were tested on an incongruent condition (e.g., movie-watching, n-back;

**Table 1**). Interestingly, the model trained on resting state significantly predicted EM when

tested on movie-watching ($r^2 =19.36\%$, $p <.001$) and n-back ($r^2 =14.44\%$, $p =.003$) conditions.

In contrast, models trained on movie-watching or n-back could not be generalized to other

conditions, unable to significantly predict EM ($p$'s $>.1$), except for significant generalizability

from the movie to the rest condition (**Table 1**).

In a cross-dataset validation analysis, the best-performing model from the age-heterogenous

DyNAMiC dataset was tested on the corresponding condition in an age-homogeneous cohort

from the COBRA dataset. By doing this, we found that the resting state model derived from

DyNAMiC significantly predicted EM performance in the COBRA dataset ($r^2 =5.76\%$, $p$

$<.0001$).

Overall, our results suggest that both resting state and movie-watching FC effectively predict

EM performance, although resting state models provide better generalizability across different

cognitive states and datasets. This finding could further be replicated using the Schaefer

Parcellation (**Supplementary Figs. 1-2 and Tables S1-S2**).

*Figure 1. Model trained on functional connectivity maps acquired at rest predicts (**a**) episodic memory (EM) of

the test dataset. The models trained on movie-watching (**b**) but not n-back (**c**) datasets predicted the EM scores of

the test dataset. **Table 1** summarizes the p values and correlations for each model. Test datasets were obtained

from the same cohort for rest, movie-watching, and n-back. The winning model trained on EM at rest was

evaluated on the external COBRA cohort and yielded a significant prediction of EM scores (**d**). Bootstrap

distributions of correlations between predicted and actual EM scores indicated no significant difference in the

predictive power of EM between models trained on resting state and movie-watching data (**e**). Additionally, the

bootstrap distribution revealed that models trained on resting state and movie-watching data yielded higher

correlations than those trained on n-back data (**e**). Visualization of features contributing to successful prediction

of EM at rest (**f**). A grad-CAM-derived saliency map displays the features that contributed to the model's

predictions. The hot spots overlaid on the FC map demonstrate noticeable cross-correlation contributions in

"default mode" (DMN) regions. Another important feature visualized by Grad-CAM includes off-diagonal hot

spots reflecting inter-connections of the DMN – "subcortical" node.*

**Table 1.** *Correlation results for episodic memory (EM) score predictions*

| Model trained on | rest | | | movie | | | n-back | | |
|---|---|---|---|---|---|---|---|---|---|
| **Model tested on** | rest | movie | n-back | rest | movie | n-back | rest | movie | n-back |
| Correlation | **.50** | **.44** | **.38** | **.28** | **.49** | **.09** | .05 | .08 | .17 |
| Significance | **<.0001** | **.0004** | **.003** | **.02** | **<.0001** | **.49** | .76 | .84 | .42 |

Next, we aimed to delineate the relative contributions of different brain regions for the best-

performing model—the model trained on the "resting state data" in predicting episodic

memory. Utilizing the Grad-CAM algorithm, saliency maps were generated for the 120 FC

matrices used during the training of the winning model. An averaged and interpolated

description of all saliency maps is depicted in **Figure 1f**. The saliency map highlights specific

edges, especially within the default-mode network, edges between the default-mode network

and subcortical areas, and edges between the default-mode and the cerebellar network. These

edges, indicated by a salience intensity of $\geq.5$, exert a significant influence on the model (**Fig.**

**1f**).

***2.3. Movie-watching and n-back models outperform resting state in working-memory***

***predictions, with movie-watching offering the best generalizability***

We next investigated whether the superiority of resting state in predicting EM was unique to

this domain, considering that previous research demonstrated the advantages of task-based

fMRI and naturalistic viewing in predicting fluid intelligence (*88*). To do so, we compared the

predictive power of different states for WM, which is shown to be more directly associated

with fluid intelligence compared to EM.

The model derived from the resting state failed to predict and generalize regarding WM (*p*'s

$>.10$; **Fig. 2a** and **Table 2**). By contrast, models trained on movie-watching and n-back yielded

significant predictions of WM (**Figs. 2b-c** and **Table 2**), with the movie-watching model

emerging as the best-performing model ($r^2 = 32.49\%$, $p < .0001$) followed by n-back

($r^2 = 22.09\%$, $p < .0001$). While there was no significant difference between movie-watching

and n-back in predicting WM (Mann–Whitney *U* test, $p = .81$), these models yielded better WM

prediction than resting state ($p < 0.0001$, Mann–Whitney *U* test; **Fig. 2e**).

In cross-condition validation (**Table 2**), the movie-watching model yielded a significant WM

prediction during both resting state ($r^2 = 17.64\%$, $p < .0001$) and n-back ($r^2 = 21.16\%$, $p = .004$).

The model derived from n-back yielded a significant WM prediction during movie-watching

($r^2 = 21.16\%$, $p = .0002$), but not during the resting state ($r^2 = 4\%$, $p = .12$).

For cross-dataset validation, the best-performing model in the DyNAMiC dataset (i.e., movie-

watching) was applied to predict WM to the COBRA dataset. However, since COBRA does

not include a movie-watching paradigm, we applied the model to the n-back task in COBRA.

This approach revealed that the model from the DyNAMiC movie-watching condition yielded

a significant WM prediction in the COBRA n-back task ($r^2 = 22.09\%$, $p < 0.0001$).

Overall, our results suggest that movie-watching-based WM predictions generalize across

different cognitive states and datasets. This finding could be further replicated using a different

functional parcellation (**Supplementary Figs. 1-2 and Tables S1-S2**).

***Figure 2***. *Model trained on FC maps acquired at rest did not significantly predict (**a**) the working memory (WM)*

*in the test dataset. The model trained on the movie-watching dataset yielded the best-performing model in*

*predicting WM (**b**) while the model trained on the n-back dataset (**c**) was the second-best model. **Table 2***

*summarizes the p values and correlation power for each model. (**d**) Results of cross-dataset validation, where the*

*best-performing model in the DyNAMiC dataset (i.e., movie-watching) was applied to predict WM to the COBRA*

*dataset. However, since COBRA does not include a movie-watching paradigm, we applied the model to the n-*

*back task in COBRA(**Table 2**). Bootstrap distributions of correlations between predicted and actual WM scores*

*showed no significant difference in predictive power between models trained on movie-watching and n-back data*

*(**e**). The bootstrap distribution revealed that models trained on movie-watching and n-back data exhibited higher*

*correlations than those trained on resting state data (**e**). The Grad-CAM-derived saliency map highlights*

*dominant features in the FC maps that contributed to the model's predictions (**f**). The hot spots overlaid on an FC*

*map demonstrate noticeable cross-correlation contributions in the "VAN", "visual", and to a lesser degree (<.5)*

*"DMN". Other important features visualized by Grad-CAM include off-diagonal hot spots reflecting inter-*

*connections of the "DMN" – with "FPN, Fronto-parietal Task Control", "Subcortical", and "Cerebellar";*

*"Cerebellar" – "FPN" node.*

**Table 2.** *Correlation results for working memory (WM) score predictions.*

| Model trained on | rest | | | movie | | | n-back | | |
|---|---|---|---|---|---|---|---|---|---|
| **Model tested on** | rest | movie | n-back | rest | movie | n-back | rest | movie | n-back |
| Correlation | .02 | .08 | .12 | .42 | .57 | .46 | .20 | .46 | .47 |
| Significance | .88 | .72 | .15 | <.0001 | <.0001 | .004 | .12 | .0002 | <.0001 |
| COBRA | | | | | | | | | |
| Correlation | | | | | | 0.47 | | | |
| Significance | | | | | | <.0001 | | | |

The Grad-CAM algorithm generated saliency maps of 120 FC maps from the DyNAMiC

dataset employed for training. **Figure 2f** depicts an average of all Grad-CAM generated maps.

The saliency map unveils that certain edges, specifically within network connectivity of task-

positive regions such as the frontoparietal task control network, dorsal/ventral attention, visual,

and subcortical networks as well as between-network connectivity. FC between task-positive

dorsal and ventral attention networks, and between the DMN and the fronto-parietal network,

contributed to the best-performing model derived from the movie-watching dataset.

***2.4. The brain-cognition gap is related to physical activity levels and cardiovascular risk***

***factors***

Given the importance of lifestyle and cardiovascular risk for maintaining healthy brain function

(*66, 67, 89*), we examined whether individuals with positive versus negative prediction gaps

differ in physical activity habits, education, and Framingham cardiovascular disease (CVD)

risk score. Our primary focus is on EM predictions derived from resting state data, as this was

the common condition in the DyNAMiC and COBRA datasets. We computed the difference

between predicted and observed EM scores to generate a gap score. A positive brain-cognition

gap indicates that an individual's brain predicted better-than-observed EM performance,

whereas a negative gap indicates more compromised brains relative to actual performance.

In the test sample of the DyNAMiC data (n =60) and the entire COBRA sample (n =177), we

found that individuals with a negative brain-cognition gap exhibited lower levels of physical

activity and higher CVD risk scores compared to those with positive gaps (**Fig. 3**). Moreover,

individuals with negative brain-cognitive gaps were less educated compared to those with
positive brain-cognitive gaps in the DyNAMiC dataset.

347
348   ***Figure 3.*** *The boxplots compare physical activity scores (total hours per week), CVD risk, and educational level*
349   *(years) between two groups with positive and negative brain-cognition gaps in the DyNAMiC (**top row**) and*
350   *COBRA (**bottom row**) datasets. The striped cyan-colored boxes represent scores for subjects with a positive brain-*
351   *cognition gap, while the magenta-colored boxes represent subjects with a negative estimated gap.*

### 2.5. Dopamine D1 and D2 receptor availability are associated with brain-cognition gaps

We hypothesized that inadequate DA levels might adversely affect neurocognitive function by
reducing the neural signal-to-noise ratio, thereby resulting in a less unique functional
connectome, consequently leading to a greater prediction gap. We therefore initially
investigated the relationship between DA receptor levels and predictive gaps across different
types of DA receptors in the DyNAMiC and COBRA samples.
In the DyNAMiC sample, we found a significant correlation between striatal D1DR and
prediction gap (positive gap: $r = -.46$, $p = .03$; negative gap: $r = .38$, $p = .01$) (**Fig. 4a**), suggesting
that lower D1DR is associated with greater brain-cognition gaps.

**Figure 4.** *Relationship between the gap measured from predicted and actual EM scores and dopamine D1 receptor (**a**). Negative gaps indicate that the predicted EM score was lower than actual EM scores, while the positive gaps indicate higher predicted scores than actual EM scores. Partial correlation analysis showed a significant correlation between D1 receptor values and the measured negative and positive gaps. Relationship between the gap measured from predicted and actual EM scores and dopamine D2 receptor (**b**). Negative gaps indicate that the predicted EM score was lower than actual EM scores, while positive gaps indicate higher predicted scores than actual EM scores. Partial correlation analysis showed a significant link between D2 receptor values to negative gaps and positive gaps.*

We replicated our finding in the COBRA sample using D2-like receptor availability (D2DR), revealing a significant relationship between striatal D2DR and prediction gap (positive gap: $r$ =-.38, $p$ =.01; negative gap: $r$ =.39, $p$ =.004) (**Fig. 4b**). Our findings provide support the view that lower D1DR/D2DR is associated with larger brain-cognition prediction gaps.

### 2.6. Regional variability mediates the direct impact of dopamine on brain-cognition gaps

To evaluate whether functional variability mediated the relationship between D1DR and prediction gap connectivity, we conducted additional mediation analyses. We computed BOLD signal entropy, which estimates within-region signal variability (*84*), that was averaged across all regions of the parcellation. In the DyNAMiC sample, for the group exhibiting a negative brain-cognitive gap, we observed a significant indirect effect of D1DR on the gap mediated by entropy, $\beta$ =2.41, 95% CI [0.89, 4.51], $p$ <.0001, explaining 56.81% of the total effect of D1DR on the gap. The direct effect of D1DR, however, was not significant, $\beta$ =1.83, 95% CI [-.86, 3.79], $p$ =.19. Similarly, for those with a positive brain-cognition gap, an indirect effect of D1DR on the gap through entropy was observed, $\beta$ =-6.78, 95% CI [-11.26, -3.05], $p$ <.0001, accounting for 89.41% of the D1DR effect on the gap. The direct effect of D1DR was again non-significant, $\beta$ =-0.803, 95% CI [-.48, 2.27], $p$ =.26. These findings suggest that lower D1DR levels contribute to increased signal variability, which in turn leads to reduced specificity of FC and, consequently, a larger brain-cognition gap.

In the COBRA sample, within the negative gap group, we detected a significant indirect effect

of D2DR on both the negative and positive gap groups through entropy. For the negative gap,

$\beta$ =2.18, 95% CI [0.01, 4.25], $p$ =.04, accounting for 63.43% of the D2DR effect on the gap;

and for the positive gap, $\beta$ =-.2.18, 95% CI [-3.87, -.04], $p$ =.04, accounting for 61.49% of the

D2DR effect on the gap.

**3. DISCUSSION**

Using deep learning models, we examined the predictive power of the functional connectome

during various states (resting state, movie-watching, and n-back) on two different cognitive

domains (EM and WM). Both rest and movie-watching states yielded significant predictions

of EM, with the model derived from resting state generalized across states and datasets.

Differences between the DyNAMiC and COBRA datasets make cross-dataset prediction a

harder problem, as the age ranges of samples significantly vary, and prior studies highlight the

importance of individual characteristics like age in predicting behavior from FC (*33*). Possibly

as a consequence of these age differences, model performance decreased when using deep

learning models to predict EM from COBRA data. Validation outcomes thus affirm that the

models, particularly those constructed from rest data, are robust to the particulars of the dataset.

The saliency map generated from the final layer of the deep learning model indicates that

certain edges within DMN, between DMN and the subcortical network contributed

significantly to the prediction. Building on a recent finding by Kurkela and Ritchy (*90*), our

finding reveals that a fraction of the known-memory subnetwork within the DMN, as well as

whole-brain multivariate pattern, which notably encompasses interactions of the DMN with

other networks, such as the subcortical network, made a more substantial contribution to

prediction. Importantly, this prediction generalizes across conditions and datasets, suggesting

that features derived from resting state FC serve as a stable marker of individual differences in

EM.

Our findings are in contrast to recent work suggesting that task paradigms, in general, and

movie-watching, in particular, outperform resting state data in predicting cognitive

performance (*49, 50, 91*). There are two possible explanations for this discrepancy. First, while

previous studies have often demonstrated a superiority of task and naturalistic viewing over

resting state in predicting fluid intelligence or WM (*49, 50*), there are fewer reports of FC

predicting EM (e.g., (*92, 93*)) and, to our knowledge, no study, has compared rest and movie-

watching. While we acknowledge that the resting state represents a complex amalgamation of

cognitive, emotional, and perceptual processes (*94*), the good prediction power of the resting

state may arise from the presence of mind wandering during rest, which is strongly related to

EM (*95, 96*). EM plays a crucial role in generating mental content during mind wandering,

especially episodes characterized by distinct times and locations (*59, 97*). Second, most

previous studies utilized connectome predictive modeling (CPM), which relies on a linear

relationship between FC and behavioral data. In contrast, deep learning makes no assumption

about the nature of the relationship, and thus, a deep learning model may outperform CPM.

Indeed, the predictive power in the current study is stronger than for CPM-based predictions

reported before. For example, across various studies, the correlation between predicted and

actual fluid intelligence typically hovers around 0.25 (*98-100*), while an average correlation of

approximately 0.11 was reported across 58 different measures covering various domains (*101*).

However, in the current study, we found that deep learning yields a stronger association (resting

state prediction of EM: $.31 < r < .49$; movie-watching prediction of WM: $.42 < r < .57$), even

across data-sets (cross-validation of EM: $r = .24$; cross-validation of WM: $r = .47$, between FC

and cognitive data).

In contrast to the EM prediction, both n-back and movie-watching datasets yielded significant

predictions of offline WM performance. Importantly, the models derived from movie-watching

and n-back outperformed the resting state in WM prediction.  These differences in model

accuracy when predicting the same target behaviors (i.e., WM) suggest the presence of trait-

state interactions. Specifically, movies and n-back enhance individual differences in WM-

relevant connection. Indeed, we found that several WM-related networks (*102-105*), including

the fronto-parietal, the salience, and the dorsal/ventral attention networks contribute to

prediction. Additionally, previous research showed that movie-watching alters the propagation

of activity across cortical pathways (*106*), particularly within and between regions involved in

audiovisual processing and attention. These alterations lead to a less segregated and more

integrated network organization (*107*). Similarly, the n-back task has been associated with

increased integration of task-positive cortico-cortical connectivity (*105, 108*) and striato-

cortical connectivity (*103*). Taken together, our results confirm previous findings that movie

watching paradigm is a great condition for studying individual differences in various cognitive

domains. Nonetheless, if movie watching paradigm is not feasible/ available, the resting state

still provides a robust means of studying individual differences, particularly in self-referential

domains, such as EM.

Our study introduces a novel deep neural network architecture that features dense connections
and incorporates an attentional mechanism. We found that the current method outperformed
widely used approaches, such as kernel regression (*110*) and connectome predictive modeling
(*17*), in predicting cognitive data. Our findings contrast with previous work showing similar
performance between kernel regression and various DMN architectures for predicting
behavioral and demographic variables (*29*). Unlike the BrainNet convolutional neural network,
which focuses on staged transformations, our densely connected model promotes extensive
feature reuse, possibly leading to more robust feature extraction. That is, the model explicitly
uses the Enhanced Residual Block (ERB) and High-Frequency Attention Block (HFAB),
enhancing the focus on relevant features and potentially boosting performance. The
architecture includes multiple layers and blocks, balancing depth and computational efficiency.
472
Applying two different parcellation methods (*86, 111*) to the DyNAMiC data indicated that
parcellation resolution does not significantly impact model performance (see **Figs. S1-S2 and**
**Tables S1-S2**). However, improvements in registration and parcellation, such as individualized
parcellations and areal features for alignment, may enhance the delineation of functionally
homogeneous regions (*112*), improving the predictive performance of deep learning models.
Therefore, the impact of alignment and parcellation on predictive model performance remains
a key area for future research. Additionally, the performance of our approach still needs to be
examined on a large-scale publicly available neuroimaging dataset for predicting other types
of cognitive data.
482
We found a significant link between the brain-cognition prediction gap, life-style, and risk
factors for vascular disease, such that individuals with a negative gap exhibited lower levels of
physical activity and higher cardiovascular risk scores compared to those with a positive gap.
This finding was consistent across both age- heterogeneous and age- homogenous older
samples. Brain-cognition prediction gap could serve as a biomarker for identifying individuals
at risk (e.g., individuals with a negative gap). Previous studies suggest that the brain age
prediction gap is associated with cognitive aging (*62*), some aspects of physiological aging
(*61*), as well as aging in other organs (*113*), and even mortality in older age (*61*). However, a
recent study revealed that brain age accounts for only a small portion of cognitive decline
compared to chronological age (*65*), suggesting that cognitive prediction might be more
informative. Our findings build upon this concept by extending the brain-age gap to behavioral
variables, demonstrating that the brain-cognition gap could provide insights regarding physical

activity status, education, and cardio vascular risk – key factors contributing to cognitive

reserve (*114-117*).

Critically, we found that D1DRs and D2DRs were strongly associated with the brain-cognition

gap, such that lower dopamine receptor levels were associated with greater gaps. More

specifically, in two independent samples, we discovered greater correspondence (i.e., near zero

in the gap) between brain function and cognition in individuals with higher D1DR/D2DR

whereas lower correspondence (i.e., significantly different gap from zero) was found in

individuals with lower D1DR/D2DR availability. Previous computational models proposed

that DA modulates neuronal gain, which improves SNR in neural processing, contributing to

more coherent activity across large-scale networks (e.g., balanced integration and segregation

(*83*)). Past studies also showed that lower D1DR contributed to more BOLD variability in the

subcortical area (*81*) and less functional segregation of the striatum (*118*) and the large-scale

networks in aging (*83*), possibly due to increased noise (lower SNR). In support of this notion,

we found for the first time that regional variability, estimated using entropy, mediated the

impact of DA on the brain-cognition gap. Although the cross-sectional nature of our data

warrants caution, this novel finding suggests that lower DA integrity relates to BOLD

variability, which in turn is associated with a larger brain-cognition gap.

In summary, our findings reveal that while tasks like movie-watching excel in predicting WM

performance, there are features during rest that predict internally-oriented mind-wandering-

type tasks, such as EM. Additionally, individuals whose brains predict poorer cognitive

performance (i.e., negative gap) exhibit lower physical activity and higher cardiovascular risk

compared to those whose brains predict higher cognitive function than their actual performance

(i.e., positive gap). This finding suggests that our prediction model offers a novel marker to

identify individuals at risk of compromised brain maintenance. Furthermore, individuals with

lower DA showed less accurate cognitive prediction (larger brain-cognition gap) due to

increased BOLD variability and less unique and cohesive FC.

**4. MATERIALS AND METHODS**

*4.1. Participants*

All participants provided written informed consent and studies were conducted in accordance

with the Declaration of Helsinki and approved by the Regional Ethical Board and the local

Radiation Safety Committee (reference numbers: 2012-57-31M; 2017-404-32M).

This study used data from DyNAMiC (*54*), which is a longitudinal study with a focus on

changes in the brain connectome and the D1DR system. At baseline,180 participants (20-79

years, 50% female) across the adult lifespan underwent all tests between 2017 and 2020 (*54*)

(**Fig. 5**). Rigorous exclusion criteria were used to recruit a sample without neurological

conditions and medical treatments affecting brain functioning and cognition. Exclusion criteria

included brain injury or neurological disorder, dementia, neurodevelopmental disorder,

psychiatric diagnosis, psychopharmacological treatment, history of severe head trauma,

substance abuse or dependence, and illicit drug use. Individuals with other chronic or severe

medical conditions (e.g., cancer, diabetes, and Parkinson's disease) were also excluded. Here,

we only use data from the baseline measurement.

We used a separate sample as a testing dataset from the COBRA study (*85*) (**Fig. 5**). COBRA

is a longitudinal aging study in which 181 healthy individuals (64-68 years, 45% female)

underwent baseline assessments of brain, cognition, health, and lifestyle during 2012–2014

(*85*). Exclusion criteria at baseline included traumatic brain injury, stroke, dementia,

intellectual disability, epilepsy, psychiatric and neurological disorders, diabetes, and cancer

medications, severe visual or auditory impairment, claustrophobia, and poor Swedish language

skills. In the current study, we used data from 177 subjects from COBRA who underwent both

MRI and PET examinations at baseline (*78, 85, 102, 103, 119*).

**Figure 5.** *Overview of the experimental procedure and the use of datasets. We used a 3-fold within-sample*

*(DyNAMiC) cross-validation where we trained our model on 120 subjects (8:2; 80% training:20% validation*

*during training) and tested it in a separate sample of 60 subjects. The winning within-sample model was used*

*for between-sample (COBRA) cross-validation.*

**4.2. Cognitive Measures**

The same cognitive test battery was used in DyNAMiC and COBRA (Nevalainen et al 2015;

Nordin et al, 2022), and assessed two cognitive domains: episodic memory (EM) and working

memory (WM). Each domain was assessed using three separate tests, including letter-, number-

, and figure-based material, respectively. Participants completed all tasks on a computer and

responded by either typing in words or numbers; using the computer mouse; or pressing keys

marked by different colors. Each test included initial practice runs, after which testing followed

across several trials. For each of the three tests within a domain, scores were summarized across

trials for a measure of overall performance. Finally, these sum scores were z-standardized and

averaged to form one composite score per cognitive domain (*T* score: *mean* =50; *sd* =10)

representing within-domain performance (e.g., EM or WM performance). Missing values were

replaced by the average of the available observed scores.

***4.2.1. Episodic Memory (EM)***

Tests of EM included word recall, number-word recall, and object-location recall. In word

recall, participants viewed 16 Swedish concrete nouns, successively appearing on the computer

screen. Each word was presented for 6 s, with an inter-stimulus interval (ISI) of 1 s. Directly

following encoding, participants reported as many words as they could using the keyboard.

Two trials were completed, with a combined maximum score of 32. In number-word recall,

participants encoded pairs of 2-digit numbers and concrete plural nouns (e.g., 46 dogs). During

encoding, eight number-word pairs were presented, each for 6 s, with an ISI of 1 s. Directly

after encoding, nouns were presented again, in re-arranged order, and participants had to report

the 2-digit number associated with each presented noun (e.g., How many dogs?). This test

included two trials with a total combined maximum score of 16. The third test assessed object-

location memory. Participants viewed a 6 × 6 square grid in which 12 objects were

consecutively shown at distinct locations. Each object-position pairing was displayed for 8 s,

with an ISI of 1 s. Directly following encoding, all objects were simultaneously displayed next

to the grid for the participant to place in their correct position within the grid. If unable to recall

the correct position of an object, participants had to guess to the best of their ability. Two trials

of this test were completed, making the total combined maximum score 24.

***4.2.2. Working Memory (WM)***

Working memory was examined with three tests letter updating, number updating, and spatial

updating. These tests differed from the working memory *n*-back task performed during fMRI

scanning. In letter-updating, capital letters (A–D) were consecutively presented on the

computer screen, with participants instructed to always keep the three final letters in memory.

Each letter was presented for 1 s, with an ISI of 0.5 s. When prompted, at any given moment,

participants provided their responses by typing in three letters using the keyboard and provided

a guessing-based answer if they failed to remember the correct letters. Four practice trials were

followed by 16 test trials consisting of either 7-, 9-, 11- or 13-letter sequences. The combined

maximum number of correct answers across trials was 48 (16 trials × 3 reported letters = 48).

The number-updating test followed a columnized 3-back design. Three boxes were presented

next to each other on the screen throughout the task, in which a single digit (1–9) was

sequentially presented from left to right for 1.5 s with an ISI of 0.5 s. During an ongoing

sequence, participants had to judge whether the number currently presented in a specific box

matched the last number presented in the same box (i.e., appearing three numbers before). For

each presented number they responded yes/no by pressing assigned keys ("yes" = right index

finger; "no" = left index finger). Two practice trials were followed by four test trials each

consisting of 30 numbers. The combined maximum number of correct answers across trials

(after discarding responses to the first three numbers in every trial – as these were not preceded

by any numbers to be matched with) was 108 (27 numbers × 4 trials). In spatial-updating,

participants viewed three 3 × 3 square grids presented next to each other on the computer

screen. At the start of each trial, a blue circular object was displayed at a random location within

each grid. After 4 s, the circular objects disappeared, leaving the grids empty. An arrow then

appeared below each grid, indicating that the circular object in the corresponding grid was to

be mentally moved one step in the direction of the arrow. The arrows appeared stepwise from

left to right grids, each presented for 2.5 s (ISI = 0.5 s). Prompted by three new arrows,

participants mentally moved the circular objects one more time, resulting in each circular object

having moved two steps from its original location at the end of each trial. Participants indicated

which square the circular object in each grid had ended up in using the computer mouse. If

unsure, they were instructed to guess. The test was performed across 10 test trials, preceded by

five practice trials. The combined maximum number of correct location indications was 30.

*4.3. Measure of Physical Activity and Cardiovascular Disease Risk*

*4.3.1. Physical Activity:* An extensive battery of self-rating questionnaires was administered in

DyNAMiC and COBRA (*54, 85*). Participants were asked to indicate the frequency (number

of hours during a typical summer week; options: 1-14 h with 1-h increments, or 15+ hrs) and

the intensity (how physically demanding on a scale from 1 ="not at all" to 5 ="extremely") by

which they typically engage in a selection of activities relevant to life in northern Sweden.

These included 15 specific activities. For the purpose of the present study, we focused

on physical activities, and in particular on those activities that are purely physical and that

individuals are sufficiently engaged in (i.e., physically demanding $>= 2$). Each of these

activities was performed by at least 20% of the participants at least once a week; e.g., walking,

bicycling, jogging, strength training, household tasks, and daily work-related activities. We

computed physical activity frequency (sum hrs/week) to generate physical activity scores

accordingly.

*4.3.2. Cardiovascular Disease risk:* The risk of cardiovascular disease was determined via a

multivariable score, according to the algorithm developed in the Framingham Heart Study

(*120*). Variables include age, sex, hypertension diagnosis, systolic blood pressure, body-mass

index, smoking, and diabetes mellitus. The risk estimates were derived using an algorithm

proposed by D'Agostino et al. (*120*), which employs Cox proportional-hazard regression

models to predict the probability of developing any form of cardiovascular disease within 10

years:

$$\hat{p} = 1 - S_0(t) exp(\sum_{i=1}^{m} \beta_i (X_i - \overline{X_i}))$$          [1]

With $S_0(t)$ being the baseline survival at follow-up $t$ ($t$ =10 years), $\beta i$ the estimated regression

coefficient, $X_i$ the log-transformed value of the $i$th risk factor, $X_i$ the corresponding mean, and

*m* the number of risk factors included. Baseline survival, means, and regression coefficients

were taken from the original algorithm, with DyNAMiC and COBRA participants' risk

variables inserted to compute the final scores. Risk score calculators can be found at

*framinghamheartstudy.org*.

*4.4. Image Acquisitions*

Structural, functional, and neurochemical brain measures were acquired using MRI and PET

at Umeå University Hospital in northern Sweden. For both DyNAMiC and COBRA, all MRI

data were collected using a 3T Discovery MR750 MRI system (General Electric, Healthcare,

Illinois, USA) equipped with a 32-channel phased-array head coil. PET was conducted in 3D

mode with a Discovery PET/CT 690 (General Electric, WI, United States) to assess whole-

brain D1DR with [11C]SCH23390) and D2DR with [11C]raclopride at rest in DyNAMiC and

COBRA, respectively. Comprehensive descriptions of MRI, PET, and cognitive testing

protocols are given elsewhere (*54, 85*). In this study, we primarily focus on those data directly

pertinent to the current investigation.

*4.4.1. Functional MRI*

For the DyNAMiC dataset, high-resolution anatomical T1-weighted images were collected

using a 3-dimensional (3D) fast spoiled gradient-echo sequence with acquisition parameters of

176 sagittal slices, thickness =1 mm, TR =8.2 ms, TE =3.2 ms, flip angle =12°, and a field of

view (FOV) =250×250 mm. Whole-brain functional images were acquired during resting state,

naturalistic viewing, and an n-back WM task. Functional images were acquired using a T2*-

weighted single-shot echo-planar imaging (EPI) sequence, with 330 volumes collected over

12 min. The sequence provided 37 axial slices, slice thickness =3.4 mm, 0.5 mm spacing, TR

=2,000 ms, TE =30 ms, flip angle =80°, and FOV =250×250 mm. Ten dummy scans were

collected at the start of the sequence. During the resting state, participants were instructed to

keep their eyes open and focus on a white fixation cross on a black background displayed on a

computer screen through a tilted mirror attached to the head coil. WM was assessed in the

scanner (12-min) with a numerical *n*-back task which consisted of blocks of 1-back, 2-back,

and 3-back (*102, 103*). During movie-watching, participants viewed and listened to a 12-minute

video consisting of selected and chronologically ordered sections from the Swedish movie

*Cockpit* (*121*). Participants were instructed to view the movie attentively and answered a short

multiple-choice questionnaire about the movie after the scanning session.

In COBRA, data were collected using identical scans for resting state and n-back WM.

However, the resting state scan was shorter, lasting only 6 minutes (*73, 103*).

*4.5. Functional Connectivity Analysis*

Functional data from all conditions (i.e., rest, movie, n-back) were pre-processed using the

Statistical Parametric Mapping software package (SPM12). The functional images were first

corrected for slice-timing differences and in-scanner motion followed by registration to

anatomical T1 images. Distortion correction was performed using subject-specific and T1 co-

registered field maps. The functional time series were subsequently demeaned and detrended

followed by simultaneous nuisance regression and temporal high-pass filtering (threshold at

0.009 Hz) to not re-introduce nuisance signals (*122*). The nuisance regression model included

mean cerebrospinal and white-matter signal and their derivatives, 24-motion parameters (*123*),

a binary vector flagging motion-contaminated volumes exceeding framewise displacement

(FD) of 0.2 mm (*124*), in addition to an 18-parameter RETRICOR model (*125, 126*) of cardiac

pulsation (up to third-order harmonics), respiration (up to fourth-order harmonics), and first-

order cardio-respiratory interactions estimated using the PhysIO Toolbox v.5.0 (*127*).

Regression models for n-back included an additional set of finite impulse response (FIR) task

regressors (*128*) to avoid false positive connectivity due to task-evoked activations (*129*). The

FIR regression approach involved fitting mean cross-block responses for each time point within

a time-locked window of equal duration to each task block (27 blocks of 20 s), extended by an

additional 18 s following each block to account for the duration of the hemodynamic response

function (HRF). Given that this approach linearly fits a set of binary task regressors for each

time point, it is nearly identical to subtracting the mean task response, with the difference being

that FIR regression is better able to handle overlapping task responses and differences in the

shape of the HRF (*129*). The nuisance-regressed images were subsequently normalized to

sample-specific group templates (DARTEL) (*130*) for each dataset respectively, followed by

spatial smoothing using a 6-mm FWHM Gaussian kernel to mitigate DARTEL-induced

aliasing and affine-transformed to stereotactic MNI space (ICBM152NLin2009) (*131, 132*).

Functional images from both DyNAMiC and COBRA were preprocessed according to the steps

described above with the only exception that the RETRICOR parameters were excluded from

the regression models in COBRA due to technical issues related to the respiration and cardiac

traces.

### 710    *4.5.1. Graph Construction*

For all fMRI conditions, functional time series were averaged from 273 cortical and subcortical

regions, represented by 5-mm radius spheres, based on a widely employed FC parcellation

(*86*). In addition to 264 regions reported in the Power parcellation, we added 9 additional

regions including some subcortical regions, such as putamen, caudate, and anterior and

posterior hippocampus, identified using independent component analysis (*8, 133*). These

regions were categorized into 14 resting state networks according to a consensus partition (*86*).

To mitigate sampling from non-gray matter voxels, each parcel underwent erosion by a

permissive gray matter mask (eroding voxels <.1% threshold). The averaged time series were

then subjected to Pearson's correlations, followed by Fisher's r-to-z transformation, resulting

in the creation of a 273×273 adjacency matrix for each participant, with coefficients along the

main diagonals set to zero.

To further investigate the impact of network parcellation, we replicated our prediction analysis

(**Supplementary Material, Figs S1-S2, and Table S1-S2**) using Schaefer parcellation (*111*),

which entails 300 cortical and subcortical regions.

### 727    *4.6. Deep Neural Network Model*

Based on convolutional neural networks, deep learning is an advanced form of artificial
intelligence that uses multiple layers of "hidden" neural networks. Deep learning
methodologies are capable of automatically identifying complex patterns and representations
directly from raw data using these multi-layered networks, thus eliminating the need for
explicit feature engineering or manual intervention (*134*). The success of the training and
learning phases depends on the model's ability to process high-dimensional input data,
extracting meaningful features from complex data. This is done while managing the number of
trainable parameters, which are crucial for automated feature learning during the construction
of the model.

In this study, the inputs to our deep learning models were subject-specific FC maps with a
matrix size of 273×273 (e.g., **Fig. 6a**). We generated different versions of each FC map by
replacing portions of networks. For example, we relocated the DMN network toward the last
nodes, which were assigned as DAN, cerebellar, subcortical, and uncertain in **Figure 6a**. This
approach allowed us to create a diverse set of FC matrices for each individual, each reflecting
a different composition of edges and neighboring while maintaining the linear relationships
exonerated in the original data. Consequently, we augmented the dataset by producing a total
of 3,600 FC maps from the initial set of FC maps.

**Figure 6. (a)** *Example of functional connectivity map across three different cognitive states. SEN hand: SENsory*
*hand; SEN Mouth: SENsory Mouth; CON: Cingulo-Operculum control Network; Aud: Auditory; DMN: Default*
*Mode Network; Memory Ret: Memory Retrieval network; Visual: Visual; FPN: Fronto-Parietal Network;*
*Salience: Salience control network; Subcortical (upper row): subcortical network included in original Power*
*parcellation; VAN: Ventral Attention Network; DAN: Dorsal Attention Network; Cerebellar: Cerebellar network;*
*Subcortical (lower row): additional Subcortical regions, including hippocampus and caudate, added to the*
*original Power Parcellation; Uncertain: Regions with less known network assignment.* **(b)** *DenselyAttention*
*architecture. Enhanced Residual Block (ERB) and High-Frequency Attention Block (HFAB) into the Transition*
*Block. Note that each "D.L." layer in the table corresponds to the sequence BatchNormalization-ReLU-Conv3×3.*

Following the DenseNet framework (*87*), we incorporated the Enhanced Residual Block (ERB)
and High-Frequency Attention Block (HFAB) into the dense layers (**Fig. 6b**), termed
DenselyAttention, to facilitate feature reuse in each layer. DenseNet (*87*) diverges from
traditional methods like deepening layers or widening network structure by focusing on feature
reuse and bypass settings. This results in fewer parameters than similar dense networks such

as ResNet (*135*), enhances feature reuse, improves feature propagation, makes training easier,

and reduces issues of gradient vanishing and model degradation.

The architecture of DenseNet is characterized by its dense connectivity pattern, which entails

direct connections from each layer to all subsequent layers within its dense block (as illustrated

in **Fig. 6b**). This design ensures that every layer has access to the feature maps generated by

preceding layers, thereby facilitating a seamless and efficient gradient flow throughout the

network. In essence, the knowledge acquired at each layer is propagated forward, enabling the

model to effectively capture intricate patterns and dependencies within the data, ultimately

enhancing its ability to learn and generalize (*87*). Additionally, dense blocks provide a

regularizing effect, reducing overfitting, particularly on tasks with smaller training datasets

(*87*). This is suitable for this study which includes relatively small samples. Each sequence

combines operations of batch normalization (BN), rectified linear unit (ReLU), and a 3×3

convolution (Conv). Batch normalization can effectively prevent overfitting, as described by

equation 1:

778   
$$BN(x_n) = \alpha\left(\frac{x_n - \mu_B}{\sqrt{\sigma_\beta^2 + \epsilon}}\right) + \beta,$$
                [1]

780   Where $\mu_\beta$, mean; $\sigma_\beta$, standard deviation; $\epsilon$, random noise; $\alpha$ and $\beta$ are adaptable variables in

781   training. We utilized the Rectified Linear Unit (ReLU) activation function (*136*), which

782   activates neurons by directly outputting the input if it is positive, or zero otherwise, as outlined

783   in equation 2.

784   
$$ReLu(x) = \max(0, x),$$
                [2]

Where $x$ is the input to a neuron. ReLU benefits the performance of networks with dense layers

by decreasing the computation and selectively optimizing parameters.

Four dense blocks facilitate a stepwise down-sampling in the network. These blocks are

connected with transition layers, which consist of a 1×1 convolutional layer followed by a 2×2

average pooling layer.

Additionally, ERB and HFAB pairs are introduced for targeted high-frequency features and

residual block enhancement. The ERB-HFAB pairs are stacked sequentially at the beginning

of each dense block loop (**Fig. 6b**), with ERB and HFAB having 16 feature maps.

**High-Frequency Attention Block (HFAB):** Our approach to attentional mechanisms,
specifically the HFAB, introduces a sequential attention branch inspired by edge detection.
This branch rescales each position based on its neighboring pixels, efficiently focusing on high-
frequency areas. The HFAB employs a 3×3 convolution to enhance both the receptive field and
computational efficiency. Batch normalization is seamlessly integrated into the attention
branch, introducing global statistics during inference without additional computational cost.

**Enhanced Residual Block (ERB):** We present ERB as an alternative to the traditional residual
block. As illustrated in **Figure 6b**, ERB comprises a re-parameterization block (RepBlock) and
a ReLU. In the training stage, the RepBlock utilizes a 1×1 convolution to either expand or
contract the number of feature maps, employing a 3×3 convolution to extract features in a
higher-dimensional space. Furthermore, two skip connections are integrated to mitigate
training complexities. During inference, all linear transformations can be unified, facilitating
the conversion of each RepBlock into a singular 3×3 convolution. Essentially, ERB capitalizes
on the advantages of residual learning.

We implemented model training using Tensorflow 2.11.0 (*137*) and Keras 2.11.0 (*138*) as
programming interfaces and trained on a fifth-generation MacBook Pro (Apple M1 MAX
silicon chips, 10-core CPU, 24-core GPU, 16-core Neural Engine, 64 GB memory). For
regression tasks, selecting an appropriate loss function is crucial for guiding the optimization
process and ensuring accurate predictions. In this study, we opted for the mean squared error
(MSE) loss function due to its suitability for regression problems. To minimize the loss
function, we trained the network using the stochastic gradient descent (SGD) optimizer with a
learning rate of $8e^{-5}$ and a Nesterov momentum of 0.9 (*139*). The number of epochs was set to
100, and the batch size was set to 74. We also added dropout (*140*) after each convolutional
layer, except the first one, with a rate of 0.15.

We conducted training on distinct, identical datasets extracted from DyNAMiC, each
comprising FC maps generated from 120 subjects. To prepare each training dataset, we
randomly shuffled data for training and validation patches, allocating 80% for training and 20%
for validation. For testing, we utilized all FC maps of 60 subjects from the same sample (the
age-heterogeneous DyNAMiC study) enabling us to assess model performance through three-
fold cross-validation (**Fig. 5**). Each cross-validation fold was a new training with unseen
validation set to the model. Based on its performance on the testing dataset, we selected and

employed the winning model for all subsequent analyses. This model, which demonstrated the

best performance on the testing dataset, underwent cross-validation in an independent sample

from the age-homogeneous COBRA study.

To explore and visually represent the crucial features of the deep learning models contributing

to the prediction of cognitive scores, we used the Grad-CAM (Gradient-weighted Class

Activation Mapping) technique (*141*). This method interprets the model's decisions by

highlighting the regions of the input image with the most significant impact on the model's

output. Grad-CAM conducts a backward pass to compute the gradients of the target class score

with respect to the feature maps of the final convolutional layer. We present the average

heatmaps calculated for all input data to the model (*141*).

### *4.7. Positron Emission Tomography (PET)*

The scanning sessions started with a 5-minute low-dose helical CT scan (20 mA, 120 kV, 0.8

s per revolution), obtained for attenuation correction. During scanning, a thermoplastic mask

was attached to the bed surface to minimize head movement.

In DyNAMiC, a 60-minute scan was performed following 350 MBq ($337 \pm 27$ MBq) in list-

mode format. Offline re-binning of list-mode data was conducted to achieve a total of 49 frames

with increasing length. In COBRA, a 55-min, 18-frame dynamic PET scan was acquired during

rest following intravenous bolus injection of approximately 250 MBq 11C-raclopride ($264 \pm$

19 MBq). For both studies, attenuation- and decay-corrected images (47 slices, field of view =

25 cm, $256 \times 256$-pixel transaxial images, voxel size = $0.977 \times 0.97 \times 3.27$ mm$^3$) were

reconstructed with the iterative VUE Point HD-SharpIR algorithm (GE; 6 iterations, 24

subsets, 3.0 mm post filtering; full-width-at-half-maximum (FWHM): 3.2 mm). The estimation

of receptor availability, or binding potential relative to non-displaceable binding ($BP_{ND}$), was

carried out following previously described procedures with the cerebellum as a reference

region (*74*). PET images were motion-corrected and co-registered with the structural T1-

weighted images from the corresponding session using Statistical Parametric Mapping

software (SPM12, Wellcome Department of Imaging Science, Functional Imaging Laboratory,

London, UK). Motion-corrected PET data were resliced to match the spatial dimensions of MR

data (1 mm$^3$ isotropic, $256 \times 256 \times 256$). The mean of the first five frames was used as a source

for co-registration. In DyNAMiC, frame-to-frame head motion correction, with translations

ranging from 0.23 to 4.22 mm (mean $\pm$ sd = $0.95 \pm 0.54$ mm), revealed a trend-level difference

across age-groups (age <40 and age $\geq$ 40 years), as determined by Student's t-test ($t$ =2.0, *p*

=.047; mean ± sd for younger individuals = 1.07 ± 0.52, mean ± sd for older individuals =0.90

± 0.55). Partial-volume-effect (PVE) correction was achieved using the symmetric geometric

transfer matrix (SGTM) method for regional correction, implemented in FreeSurfer (*142*). An

estimated point-spread-function of 2.5 mm FWHM was utilized. Regional estimates of $BP_{ND}$

were calculated within the automated FreeSurfer segmentations employing the simplified

reference tissue model (SRTM (*143*)). In the current study, we focused on the striatal $BP_{ND}$,

calculated as an average of $BP_{ND}$ across the caudate and putamen.

*4.8. The direct impact of dopamine on brain-cognition gaps through mediation analysis*

To evaluate whether functional variability mediates the relationship between D1DR and

prediction gap connectivity, we conducted additional mediation analyses. We first computed

entropy, which estimates within-region signal variability (*84*), and then averaged this measure

across all regions in the parcellation. Following the mediation analysis framework proposed by

Baron and Kenny (*144*), our goal was to determine whether the association between D1/D2

receptors and brain-cognition gaps is mediated by regional variability (entropy) or if the

indirect effect exceeds the direct association between D1/D2 receptors and brain-cognition

gaps. To assess the statistical significance of this mediation effect, we employed the

bootstrapping method as outlined by Preacher and Hayes (*145*).

*4.9. Statistical significance analysis*

Statistical analyses were carried out using SPSS (IBM Corp., V24.0.0, Armonk, NY, USA),

MATLAB (The MathWorks Inc., V9.13.0 (R2022b), Natick, MA, USA), and GraphPad Prism

(GraphPad Software, Inc. V5.01, CA, USA). We performed partial correlations between

predicted and actual scores, as well as linear regression analyses. To investigate the relationship

between generated gap variables and DA receptor availability, we controlled for age (in

DyNAMiC) using partial correlation. The Mann–Whitney *U* test was used to calculate the

mean differences in prediction accuracy. The level of statistical significance was set at $p \leq .05$.

**5. REFERENCES**
1.  B. Biswal, F. Z. Yetkin, V. M. Haughton, J. S. Hyde, Functional connectivity in the motor cortex of resting
human brain using echo-planar MRI. *Magn Reson Med* **34**, 537-541 (1995).
2.  R. L. Buckner, F. M. Krienen, The evolution of distributed association networks in the human brain. *Trends*
*Cogn Sci* **17**, 648-665 (2013).
3.  M. P. van den Heuvel, H. E. Hulshoff Pol, Exploring the brain network: a review on resting-state fMRI
functional connectivity. *European neuropsychopharmacology : the journal of the European College of*
*Neuropsychopharmacology* **20**, 519-534 (2010).
4.  L. K. Ferreira, G. F. Busatto, Resting-state functional connectivity in normal brain aging. *Neurosci Biobehav*
*Rev* **37**, 384-400 (2013).

5. J. S. Damoiseaux, Effects of aging on functional and structural brain connectivity. *Neuroimage* **160**, 32-40
(2017).
6. M. D. Fox, M. Greicius, Clinical applications of resting state functional connectivity. *Front Syst Neurosci* **4**,
19 (2010).
7. J. R. Andrews-Hanna, A. Z. Snyder, J. L. Vincent, C. Lustig, D. Head, M. E. Raichle, R. L. Buckner,
Disruption of large-scale brain systems in advanced aging. *Neuron* **56**, 924-935 (2007).
8. A. Salami, A. Wahlin, N. Kaboodvand, A. Lundquist, L. Nyberg, Longitudinal Evidence for Dissociation of
Anterior and Posterior MTL Resting-State Connectivity in Aging: Links to Perfusion and Memory. *Cereb*
*Cortex* **26**, 3953-3963 (2016).
9. N. Kaboodvand, L. Backman, L. Nyberg, A. Salami, The retrosplenial cortex: A memory gateway between
the cortical default mode network and the medial temporal lobe. *Hum Brain Mapp* **39**, 2020-2034 (2018).
10. W. W. Seeley, V. Menon, A. F. Schatzberg, J. Keller, G. H. Glover, H. Kenna, A. L. Reiss, M. D. Greicius,
Dissociable intrinsic connectivity networks for salience processing and executive control. *The Journal of*
*neuroscience : the official journal of the Society for Neuroscience* **27**, 2349-2356 (2007).
11. B. Avelar-Pereira, L. Backman, A. Wahlin, L. Nyberg, A. Salami, Age-Related Differences in Dynamic
Interactions Among Default Mode, Frontoparietal Control, and Dorsal Attention Networks during Resting-
State and Interference Resolution. *Front Aging Neurosci* **9**, 152 (2017).
12. B. Avelar-Pereira, L. Backman, A. Wahlin, L. Nyberg, A. Salami, Increased functional homotopy of the
prefrontal cortex is associated with corpus callosum degeneration and working memory decline. *Neurobiol*
*Aging* **96**, 68-78 (2020).
13. B. Machner, L. Braun, J. Imholz, P. J. Koch, T. F. Munte, C. Helmchen, A. Sprenger, Resting-State
Functional Connectivity in the Dorsal Attention Network Relates to Behavioral Performance in Spatial
Attention Tasks and May Show Task-Related Adaptation. *Front Hum Neurosci* **15**, 757128 (2021).
14. J. Schimmelpfennig, J. Topczewski, W. Zajkowski, K. Jankowiak-Siuda, The role of the salience network in
cognitive and affective deficits. *Front Hum Neurosci* **17**, 1133367 (2023).
15. L. K. Ferreira, A. C. Regina, N. Kovacevic, G. Martin Mda, P. P. Santos, G. Carneiro Cde, D. S. Kerr, E.
Amaro, Jr., A. R. McIntosh, G. F. Busatto, Aging Effects on Whole-Brain Functional Connectivity in Adults
Free of Cognitive and Psychiatric Disorders. *Cereb Cortex* **26**, 3851-3865 (2016).
16. S. M. Smith, L. T. Elliott, F. Alfaro-Almagro, P. McCarthy, T. E. Nichols, G. Douaud, K. L. Miller, Brain
aging comprises many modes of structural and functional change with distinct genetic and biophysical
associations. *Elife* **9**,  (2020).
17. X. Shen, E. S. Finn, D. Scheinost, M. D. Rosenberg, M. M. Chun, X. Papademetris, R. T. Constable, Using
connectome-based predictive modeling to predict individual behavior from brain connectivity. *Nat Protoc*
**12**, 506-518 (2017).
18. E. Vul, C. Harris, P. Winkielman, H. Pashler, Puzzlingly High Correlations in fMRI Studies of Emotion,
Personality, and Social Cognition. *Perspect Psychol Sci* **4**, 274-290 (2009).
19. D. Bzdok, M. Eickenberg, G. Varoquaux, B. Thirion, Hierarchical Region-Network Sparsity for High-
Dimensional Inference in Brain Imaging. *Inf Process Med Imaging* **10265**, 323-335 (2017).
20. D. Bzdok, A. Meyer-Lindenberg, Machine Learning for Precision Psychiatry: Opportunities and Challenges.
*Biol Psychiatry Cogn Neurosci Neuroimaging* **3**, 223-230 (2018).
21. D. Bzdok, M. Krzywinski, N. Altman, Machine learning: supervised methods. *Nat Methods* **15**, 5-6 (2018).
22. J. Sui, R. Jiang, J. Bustillo, V. Calhoun, Neuroimaging-based Individualized Prediction of Cognition and
Behavior for Mental Disorders and Health: Methods and Promises. *Biol Psychiatry* **88**, 818-828 (2020).
23. S. Vieira, X. Liang, R. Guiomar, A. Mechelli, Can we predict who will benefit from cognitive-behavioural
therapy? A systematic review and meta-analysis of machine learning studies. *Clin Psychol Rev* **97**, 102193
(2022).
24. S. M. Plis, D. R. Hjelm, R. Salakhutdinov, E. A. Allen, H. J. Bockholt, J. D. Long, H. J. Johnson, J. S.
Paulsen, J. A. Turner, V. D. Calhoun, Deep learning for neuroimaging: a validation study. *Front Neurosci* **8**,
229 (2014).
25. H. K. van der Burgh, R. Schmidt, H. J. Westeneng, M. A. de Reus, L. H. van den Berg, M. P. van den
Heuvel, Deep learning predictions of survival based on MRI in amyotrophic lateral sclerosis. *Neuroimage*
*Clin* **13**, 361-369 (2017).
26. H. Nguyen, V. Nguyen, T. Nguyen, M. E. Larsen, B. O'Dea, D. T. Nguyen, T. Le, D. Phung, S. Venkatesh,
H. Christensen. (Springer International Publishing, Cham, 2018), pp. 100-110.
27. H. Li, G. Jiang, J. Zhang, R. Wang, Z. Wang, W. S. Zheng, B. Menze, Fully convolutional network
ensembles for white matter hyperintensities segmentation in MR images. *Neuroimage* **183**, 650-665 (2018).
28. Y. Choi, Y. Kwon, H. Lee, B. J. Kim, M. C. Paik, J.-H. Won, in *Brainlesion: Glioma, Multiple Sclerosis,*
*Stroke and Traumatic Brain Injuries,* A. Crimi, B. Menze, O. Maier, M. Reyes, S. Winzeck, H. Handels,
Eds. (Springer International Publishing, Cham, 2016), pp. 231-243.

Begränsad delning

29. T. He, R. Kong, A. J. Holmes, M. Nguyen, M. R. Sabuncu, S. B. Eickhoff, D. Bzdok, J. Feng, B. T. T. Yeo,
Deep neural networks and kernel regression achieve comparable accuracies for functional connectivity
prediction of behavior and demographics. *Neuroimage* **206**, 116276 (2020).
30. J. Kong, E. Wolcott, Z. Wang, K. Jorgenson, W. F. Harvey, J. Tao, R. Rones, C. Wang, Altered resting state
functional connectivity of the cognitive control network in fibromyalgia and the modulation effect of mind-
body intervention. *Brain Imaging Behav* **13**, 482-492 (2019).
31. C. Sripada, M. Angstadt, A. Taxali, D. A. Clark, T. Greathouse, S. Rutherford, J. R. Dickens, K. Shedden,
A. M. Gard, L. W. Hyde, A. Weigard, M. Heitzeg, Brain-wide functional connectivity patterns support
general cognitive ability and mediate effects of socioeconomic status in youth. *Transl Psychiatry* **11**, 571
(2021).
32. J. Chen, A. Tam, V. Kebets, C. Orban, L. Q. R. Ooi, C. L. Asplund, S. Marek, N. U. F. Dosenbach, S. B.
Eickhoff, D. Bzdok, A. J. Holmes, B. T. T. Yeo, Shared and unique brain network features predict cognitive,
personality, and mental health scores in the ABCD study. *Nat Commun* **13**, 2217 (2022).
33. A. Omidvarnia, L. Sasse, D. I. Larabi, F. Raimondo, F. Hoffstaedter, J. Kasper, J. Dukart, M. Petersen, B.
Cheng, G. Thomalla, S. B. Eickhoff, K. R. Patil, Is resting state fMRI better than individual characteristics at
predicting cognition? *bioRxiv*, 2023.2002.2018.529076 (2023).
34. L. Jiang, K. Qiao, C. Li, Distance-based functional criticality in the human brain: intelligence and emotional
intelligence. *BMC Bioinformatics* **22**, 32 (2021).
35. D. C. Jangraw, J. Gonzalez-Castillo, D. A. Handwerker, M. Ghane, M. D. Rosenberg, P. Panwar, P. A.
Bandettini, A functional connectivity-based neuromarker of sustained attention generalizes to predict recall
in a reading task. *Neuroimage* **166**, 99-109 (2018).
36. Q. Chen, R. E. Beaty, Z. Cui, J. Sun, H. He, K. Zhuang, Z. Ren, G. Liu, J. Qiu, Brain hemispheric
involvement in visuospatial and verbal divergent thinking. *Neuroimage* **202**, 116065 (2019).
37. M. D. Rosenberg, E. S. Finn, D. Scheinost, X. Papademetris, X. Shen, R. T. Constable, M. M. Chun, A
neuromarker of sustained attention from whole-brain functional connectivity. *Nat Neurosci* **19**, 165-171
(2016).
38. R. E. Beaty, Y. N. Kenett, A. P. Christensen, M. D. Rosenberg, M. Benedek, Q. Chen, A. Fink, J. Qiu, T. R.
Kwapil, M. J. Kane, P. J. Silvia, Robust prediction of individual creative ability from brain functional
connectivity. *Proc Natl Acad Sci U S A* **115**, 1087-1092 (2018).
39. W. T. Hsu, M. D. Rosenberg, D. Scheinost, R. T. Constable, M. M. Chun, Resting-state functional
connectivity predicts neuroticism and extraversion in novel individuals. *Soc Cogn Affect Neurosci* **13**, 224-
232 (2018).
40. J. D. E. Gabrieli, S. S. Ghosh, S. Whitfield-Gabrieli, Prediction as a humanitarian and pragmatic contribution
from human cognitive neuroscience. *Neuron* **85**, 11-26 (2015).
41. E. S. Finn, R. Todd Constable, Individual variation in functional brain connectivity: implications for
personalized approaches to psychiatric disease. *Dialogues Clin Neurosci* **18**, 277-287 (2016).
42. C. W. Woo, L. J. Chang, M. A. Lindquist, T. D. Wager, Building better biomarkers: brain models in
translational neuroimaging. *Nat Neurosci* **20**, 365-377 (2017).
43. E. S. Finn, X. Shen, D. Scheinost, M. D. Rosenberg, J. Huang, M. M. Chun, X. Papademetris, R. T.
Constable, Functional connectome fingerprinting: identifying individuals using patterns of brain
connectivity. *Nat Neurosci* **18**, 1664-1671 (2015).
44. E. S. Finn, D. Scheinost, D. M. Finn, X. Shen, X. Papademetris, R. T. Constable, Can brain state be
manipulated to emphasize individual differences in functional connectivity? *Neuroimage* **160**, 140-151
(2017).
45. C. L. Grady, J. D. Ryan, "Age-Related Differences in the Human Hippocampus: Behavioral, Structural and
Functional Measures" in *The Hippocampus from Cells to Systems: Structure, Connectivity, and Functional
Contributions to Memory and Flexible Cognition,* D. E. Hannula, M. C. Duff, Eds. (Springer International
Publishing, Cham, 2017), pp. 167-208.
46. K. L. Campbell, D. L. Schacter, Aging and the Resting State: Cognition is not Obsolete. *Lang Cogn
Neurosci* **32**, 692-694 (2017).
47. L. Geerligs, K. A. Tsvetanov, The use of resting state data in an integrative approach to studying
neurocognitive ageing - Commentary on Campbell and Schacter (2016). *Lang Cogn Neurosci* **32**, 684-691
(2017).
48. I. Tavor, O. Parker Jones, R. B. Mars, S. M. Smith, T. E. Behrens, S. Jbabdi, Task-free MRI predicts
individual differences in brain activity during task performance. *Science* **352**, 216-220 (2016).
49. A. S. Greene, S. Gao, D. Scheinost, R. T. Constable, Task-induced brain state manipulation improves
prediction of individual traits. *Nat Commun* **9**, 2807 (2018).
50. E. S. Finn, P. A. Bandettini, Movie-watching outperforms rest for functional connectivity-based prediction
of behavior. *Neuroimage* **235**, 117963 (2021).

51. E. S. Finn, E. Glerean, U. Hasson, T. Vanderwal, Naturalistic imaging: The use of ecologically valid
conditions to study brain function. *Neuroimage* **247**, 118776 (2022).
52. U. Hasson, Y. Nir, I. Levy, G. Fuhrmann, R. Malach, Intersubject synchronization of cortical activity during
natural vision. *Science* **303**, 1634-1640 (2004).
53. L. Geerligs, M. Rubinov, C. Cam, R. N. Henson, State and Trait Components of Functional Connectivity:
Individual Differences Vary with Mental State. *The Journal of neuroscience : the official journal of the*
*Society for Neuroscience* **35**, 13949-13961 (2015).
54. K. Nordin, T. Gorbach, R. Pedersen, V. Panes Lundmark, J. Johansson, M. Andersson, C. McNulty, K.
Riklund, A. Wahlin, G. Papenberg, G. Kalpouzos, L. Backman, A. Salami, DyNAMiC: A prospective
longitudinal study of dopamine and brain connectomes: A new window into cognitive aging. *J Neurosci Res*
**100**, 1296-1320 (2022).
55. R. B. Cattell, Abilities: Their structure, growth, and action.  (1971).
56. P. C. Kyllonen, "Is working memory capacity Spearman's g?" in *Human abilities* (Psychology Press, 2013),
pp. 49-75.
57. N. Unsworth, K. Fukuda, E. Awh, E. K. Vogel, Working memory and fluid intelligence: capacity, attention
control, and secondary memory retrieval. *Cogn Psychol* **71**, 1-26 (2014).
58. E. Tulving, D. L. Schacter, Priming and human memory systems. *Science* **247**, 301-306 (1990).
59. J. Smallwood, J. W. Schooler, The science of mind wandering: empirically navigating the stream of
consciousness. *Annu Rev Psychol* **66**, 487-518 (2015).
60. N. U. Dosenbach, B. Nardos, A. L. Cohen, D. A. Fair, J. D. Power, J. A. Church, S. M. Nelson, G. S. Wig,
A. C. Vogel, C. N. Lessov-Schlaggar, K. A. Barnes, J. W. Dubis, E. Feczko, R. S. Coalson, J. R. Pruett, Jr.,
D. M. Barch, S. E. Petersen, B. L. Schlaggar, Prediction of individual brain maturity using fMRI. *Science*
**329**, 1358-1361 (2010).
61. J. H. Cole, K. Franke, Predicting Age Using Neuroimaging: Innovative Brain Ageing Biomarkers. *Trends*
*Neurosci* **40**, 681-690 (2017).
62. T. Dunas, A. Wahlin, L. Nyberg, C. J. Boraxbekk, Multimodal Image Analysis of Apparent Brain Age
Identifies Physical Fitness as Predictor of Brain Maintenance. *Cereb Cortex* **31**, 3393-3407 (2021).
63. B. Farnsworth von Cederwald, M. Josefsson, A. Wahlin, L. Nyberg, N. Karalija, Association of
Cardiovascular Risk Trajectory With Cognitive Decline and Incident Dementia. *Neurology* **98**, e2013-e2022
(2022).
64. L. S. Jonasson, L. Nyberg, A. F. Kramer, A. Lundquist, K. Riklund, C. J. Boraxbekk, Aerobic Exercise
Intervention, Cognitive Performance, and Brain Structure: Results from the Physical Influences on Brain in
Aging (PHIBRA) Study. *Front Aging Neurosci* **8**, 336 (2016).
65. A. Tetereva, N. Pat, Brain age has limited utility as a biomarker for capturing fluid cognition in older
individuals. *eLife* **12**, RP87297 (2024).
66. L. Nyberg, M. Lovden, K. Riklund, U. Lindenberger, L. Backman, Memory aging and brain maintenance.
*Trends Cogn Sci* **16**, 292-305 (2012).
67. N. Karalija, A. Wahlin, J. Ek, A. Rieckmann, G. Papenberg, A. Salami, A. M. Brandmaier, Y. Kohncke, J.
Johansson, M. Andersson, J. Axelsson, G. Oradd, K. Riklund, M. Lovden, U. Lindenberger, L. Backman, L.
Nyberg, Cardiovascular factors are related to dopamine integrity and cognition in aging. *Ann Clin Transl*
*Neurol* **6**, 2291-2303 (2019).
68. L. Backman, U. Lindenberger, S. C. Li, L. Nyberg, Linking cognitive aging to alterations in dopamine
neurotransmitter functioning: recent data and future avenues. *Neurosci Biobehav Rev* **34**, 670-677 (2010).
69. L. Bäckman, L. Nyberg, A. Soveri, J. Johansson, M. Andersson, E. Dahlin, A. S. Neely, J. Virta, M. Laine,
J. O. Rinne, Effects of working-memory training on striatal dopamine release. *Science* **333**, 718 (2011).
70. N. D. Volkow, R. C. Gur, G. J. Wang, J. S. Fowler, P. J. Moberg, Y. S. Ding, R. Hitzemann, G. Smith, J.
Logan, Association between decline in brain dopamine activity with age and cognitive and motor
impairment in healthy individuals. *Am J Psychiatry* **155**, 344-349 (1998).
71. E. J. Juarez, G. R. Samanez-Larkin, Exercise, Dopamine, and Cognition in Older Age. *Trends Cogn Sci* **23**,
986-988 (2019).
72. L. de Boer, J. Axelsson, K. Riklund, L. Nyberg, P. Dayan, L. Backman, M. Guitart-Masip, Attenuation of
dopamine-modulated prefrontal value signals underlies probabilistic reward learning deficits in old age. *Elife*
**6**, (2017).
73. L. Nyberg, N. Karalija, A. Salami, M. Andersson, A. Wahlin, N. Kaboovand, Y. Kohncke, J. Axelsson, A.
Rieckmann, G. Papenberg, D. D. Garrett, K. Riklund, M. Lovden, U. Lindenberger, L. Backman, Dopamine
D2 receptor availability is linked to hippocampal-caudate functional connectivity and episodic memory.
*Proc Natl Acad Sci U S A* **113**, 7918-7923 (2016).
74. K. Nordin, L. Nyberg, M. Andersson, N. Karalija, K. Riklund, L. Backman, A. Salami, Distinct and
Common Large-Scale Networks of the Hippocampal Long Axis in Older Age: Links to Episodic Memory
and Dopamine D2 Receptor Availability. *Cereb Cortex* **31**, 3435-3450 (2021).

75. G. Papenberg, N. Karalija, A. Salami, A. Rieckmann, M. Andersson, J. Axelsson, K. Riklund, U.
Lindenberger, M. Lovden, L. Nyberg, L. Backman, Balance between Transmitter Availability and Dopamine
D2 Receptors in Prefrontal Cortex Influences Memory Functioning. *Cereb Cortex* **30**, 989-1000 (2020).
76. R. Cools, M. D'Esposito, Inverted-U-shaped dopamine actions on human working memory and cognitive
control. *Biol Psychiatry* **69**, e113-125 (2011).
77. J. Zahrt, J. R. Taylor, R. G. Mathew, A. F. Arnsten, Supranormal stimulation of D1 dopamine receptors in
the rodent prefrontal cortex impairs spatial working memory performance. *The Journal of neuroscience : the
official journal of the Society for Neuroscience* **17**, 8528-8535 (1997).
78. N. Karalija, G. Papenberg, J. Johansson, A. Wahlin, A. Salami, M. Andersson, J. Axelsson, D. Kuznetsov,
K. Riklund, M. Lovden, U. Lindenberger, L. Backman, L. Nyberg, Longitudinal support for the correlative
triad among aging, dopamine D2-like receptor loss, and memory decline. *Neurobiol Aging* **136**, 125-132
(2024).
79. D. Servan-Schreiber, H. Printz, J. D. Cohen, A network model of catecholamine effects: gain, signal-to-noise
ratio, and behavior. *Science* **249**, 892-895 (1990).
80. J. K. Seamans, C. R. Yang, The principal features and mechanisms of dopamine modulation in the prefrontal
cortex. *Prog Neurobiol* **74**, 1-58 (2004).
81. M. Guitart-Masip, A. Salami, D. Garrett, A. Rieckmann, U. Lindenberger, L. Backman, BOLD Variability is
Related to Dopaminergic Neurotransmission and Cognitive Aging. *Cereb Cortex* **26**, 2074-2083 (2016).
82. S. C. Li, A. Rieckmann, Neuromodulation and aging: implications of aging neuronal gain control on
cognition. *Curr Opin Neurobiol* **29**, 148-158 (2014).
83. R. Pedersen, J. Johansson, A. Salami, Dopamine D1-signaling modulates maintenance of functional network
segregation in aging. *Aging Brain* **3**, 100079 (2023).
84. G. Shafiei, Y. Zeighami, C. A. Clark, J. T. Coull, A. Nagano-Saito, M. Leyton, A. Dagher, B. Misic,
Dopamine Signaling Modulates the Stability and Integration of Intrinsic Brain Networks. *Cereb Cortex* **29**,
397-409 (2019).
85. N. Nevalainen, K. Riklund, M. Andersson, J. Axelsson, M. Ogren, M. Lovden, U. Lindenberger, L.
Backman, L. Nyberg, COBRA: A prospective multimodal imaging study of dopamine, brain structure and
function, and cognition. *Brain Res* **1612**, 83-103 (2015).
86. J. D. Power, A. L. Cohen, S. M. Nelson, G. S. Wig, K. A. Barnes, J. A. Church, A. C. Vogel, T. O.
Laumann, F. M. Miezin, B. L. Schlaggar, S. E. Petersen, Functional network organization of the human
brain. *Neuron* **72**, 665-678 (2011).
87. G. Huang, Z. Liu, L. van der Maaten, K. Q. Weinberger, Densely Connected Convolutional Networks.
*arXiv: 1608.06993*, (2016).
88. W. Zhao, C. Makowski, D. J. Hagler, H. P. Garavan, W. K. Thompson, D. J. Greene, T. L. Jernigan, A. M.
Dale, Task fMRI paradigms may capture more behaviorally relevant information than resting-state
functional connectivity. *Neuroimage* **270**, 119946 (2023).
89. Y. Kohncke, G. Papenberg, L. Jonasson, N. Karalija, A. Wahlin, A. Salami, M. Andersson, J. E. Axelsson,
L. Nyberg, K. Riklund, L. Backman, U. Lindenberger, M. Lovden, Self-rated intensity of habitual physical
activities is positively associated with dopamine D(2/3) receptor availability and cognition. *Neuroimage* **181**,
605-616 (2018).
90. K. Kurkela, M. Ritchey, Intrinsic functional connectivity among memory networks does not predict
individual differences in narrative recall. *Imaging Neuroscience* **2**, 1-17 (2024).
91. E. S. Finn, Is it time to put rest to rest? *Trends in Cognitive Sciences* **25**, 1021-1032 (2021).
92. Y. Zhu, F. Zang, Q. Wang, Q. Zhang, C. Tan, S. Zhang, T. Hu, L. Qi, S. Xu, Q. Ren, C. Xie, Connectome-
based model predicts episodic memory performance in individuals with subjective cognitive decline and
amnestic mild cognitive impairment. *Behav Brain Res* **411**, 113387 (2021).
93. C. N. Wahlheim, A. P. Christensen, Z. M. Reagh, B. S. Cassidy, Intrinsic functional connectivity in the
default mode network predicts mnemonic discrimination: A connectome-based modeling approach.
*Hippocampus* **32**, 21-37 (2022).
94. J. Gonzalez-Castillo, J. W. Y. Kam, C. W. Hoy, P. A. Bandettini, How to Interpret Resting-State fMRI: Ask
Your Participants. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **41**,
1130-1141 (2021).
95. D. Stawarczyk, A. D'Argembeau, Neural correlates of personal goal processing during episodic future
thinking and mind-wandering: An ALE meta-analysis. *Hum Brain Mapp* **36**, 2928-2947 (2015).
96. P. Blonde, M. Sperduti, D. Makowski, P. Piolino, Bored, distracted, and forgetful: The impact of mind
wandering and boredom on memory encoding. *Q J Exp Psychol (Hove)* **75**, 53-69 (2022).
97. T. Karapanagiotidis, B. C. Bernhardt, E. Jefferies, J. Smallwood, Tracking thoughts: Exploring the neural
architecture of mental time travel during mind-wandering. *Neuroimage* **147**, 272-281 (2017).
98. J. Dubois, P. Galdi, L. K. Paul, R. Adolphs, A distributed brain network predicts general intelligence from
resting-state human neuroimaging data. *Philos Trans R Soc Lond B Biol Sci* **373**, (2018).

99. J. Li, B. B. Biswal, Y. Meng, S. Yang, X. Duan, Q. Cui, H. Chen, W. Liao, A neuromarker of individual
general fluid intelligence from the white-matter functional connectome. *Transl Psychiatry* **10**, 147 (2020).
100. U. Pervaiz, D. Vidaurre, M. W. Woolrich, S. M. Smith, Optimising network modelling methods for fMRI.
*Neuroimage* **211**, 116604 (2020).
101. M. Li, D. Wang, J. Ren, G. Langs, S. Stoecklein, B. P. Brennan, J. Lu, H. Chen, H. Liu, Performing group-
level functional image analyses based on homologous functional regions mapped in individuals. *PLoS Biol*
**17**, e2007032 (2019).
102. A. Salami, D. D. Garrett, A. Wahlin, A. Rieckmann, G. Papenberg, N. Karalija, L. Jonasson, M.
Andersson, J. Axelsson, J. Johansson, K. Riklund, M. Lovden, U. Lindenberger, L. Backman, L. Nyberg,
Dopamine D(2/3) Binding Potential Modulates Neural Signatures of Working Memory in a Load-Dependent
Fashion. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **39**, 537-547
(2019).
103. A. Salami, A. Rieckmann, N. Karalija, B. Avelar-Pereira, M. Andersson, A. Wahlin, G. Papenberg, D. D.
Garrett, K. Riklund, M. Lovden, U. Lindenberger, L. Backman, L. Nyberg, Neurocognitive Profiles of Older
Adults with Working-Memory Dysfunction. *Cereb Cortex* **28**, 2525-2539 (2018).
104. J. Eriksson, E. K. Vogel, A. Lansner, F. Bergstrom, L. Nyberg, Neurocognitive Architecture of Working
Memory. *Neuron* **88**, 33-46 (2015).
105. X. Liang, Q. Zou, Y. He, Y. Yang, Topologically Reorganized Connectivity Architecture of Default-Mode,
Executive-Control, and Salience Networks across Working Memory Task Loads. *Cereb Cortex* **26**, 1501-
1511 (2016).
106. M. Gilson, G. Deco, K. J. Friston, P. Hagmann, D. Mantini, V. Betti, G. L. Romani, M. Corbetta, Effective
connectivity inferred from fMRI transition dynamics during movie viewing points to a balanced
reconfiguration of cortical interactions. *Neuroimage* **180**, 534-546 (2018).
107. R. F. Betzel, L. Byrge, F. Z. Esfahlani, D. P. Kennedy, Temporal fluctuations in the brain's modular
architecture during movie-watching. *Neuroimage* **213**, 116687 (2020).
108. H. Zhang, R. Zhao, X. Hu, S. Guan, D. S. Margulies, C. Meng, B. B. Biswal, Cortical connectivity
gradients and local timescales during cognitive states are modulated by cognitive loads. *Brain Struct Funct*
**227**, 2701-2712 (2022).
109.    G. F. Stark, E. W. Avery, M. D. Rosenberg, A. S. Greene, S. Gao, D. Scheinost, R. Todd Constable, M.
M. Chun, K. Yoo, Using functional connectivity models to characterize relationships between working and
episodic memory. *Brain Behav* **11**, e02105 (2021).
110.    K. P. Murphy, *Machine Learning: A Probabilistic Perspective* (MIT Press, 2012).
111.    A. Schaefer, R. Kong, E. M. Gordon, T. O. Laumann, X. N. Zuo, A. J. Holmes, S. B. Eickhoff, B. T. T.
Yeo, Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI.
*Cereb Cortex* **28**, 3095-3114 (2018).
112.    A. Dworetsky, B. A. Seitzman, B. Adeyemo, A. N. Nielsen, A. S. Hatoum, D. M. Smith, T. E. Nichols,
M. Neta, S. E. Petersen, C. Gratton, Two common and distinct forms of variation in human functional brain
networks. *Nat Neurosci* **27**, 1187-1198 (2024).
113.    Y. E. Tian, V. Cropley, A. B. Maier, N. T. Lautenschlager, M. Breakspear, A. Zalesky, Heterogeneous
aging across multiple organ systems and prediction of chronic disease and mortality. *Nat Med* **29**, 1221-1231
(2023).
114.    R. M. Arida, L. Teixeira-Machado, The Contribution of Physical Exercise to Brain Resilience. *Front*
*Behav Neurosci* **14**, 626769 (2020).
115.    S. Klil-Drori, K. Cinalioglu, S. Rej, Brain Health and the Role of Exercise in Maintaining Late-Life
Cognitive Reserve: A Narrative Review Providing the Neuroprotective Mechanisms of Exercise. *The*
*American Journal of Geriatric Psychiatry* **30**, S72 (2022).
116.    J. Nithianantharajah, A. J. Hannan, The neurobiology of brain and cognitive reserve: Mental and
physical activity as modulators of brain disorders. *Progress in Neurobiology* **89**, 369-382 (2009).
117.    R. Cabeza, M. Albert, S. Belleville, F. I. M. Craik, A. Duarte, C. L. Grady, U. Lindenberger, L.
Nyberg, D. C. Park, P. A. Reuter-Lorenz, M. D. Rugg, J. Steffener, M. N. Rajah, Maintenance, reserve and
compensation: the cognitive neuroscience of healthy ageing. *Nat Rev Neurosci* **19**, 701-710 (2018).
118.    S. M. Korkki, J. Johansson, K. Nordin, R. Pedersen, L. Bäckman, A. Rieckmann, A. Salami,
Dedifferentiation of caudate functional organization is linked to reduced D1 dopamine receptor availability
and poorer memory function in aging. *bioRxiv*, 2024.2003.2018.585623 (2024).
119.    L. Nyberg, M. Andersson, A. Lundquist, W. F. C. Baare, D. Bartres-Faz, L. Bertram, C. J. Boraxbekk,
A. M. Brandmaier, N. Demnitz, C. A. Drevon, S. Duezel, K. P. Ebmeier, P. Ghisletta, R. Henson, D. E. A.
Jensen, R. A. Kievit, E. Knights, S. Kuhn, U. Lindenberger, A. Plachti, S. Pudas, J. M. Roe, K. S. Madsen,
C. Sole-Padulles, Y. Sommerer, S. Suri, E. Zsoldos, A. M. Fjell, K. B. Walhovd, Individual differences in
brain aging: heterogeneity in cortico-hippocampal but not caudate atrophy rates. *Cereb Cortex* **33**, 5075-
5081 (2023).

120.    R. B. D'Agostino, Sr., R. S. Vasan, M. J. Pencina, P. A. Wolf, M. Cobain, J. M. Massaro, W. B.
1200           Kannel, General cardiovascular risk profile for use in primary care: the Framingham Heart Study.
1201           *Circulation* **117**, 743-753 (2008).
1202    121.    J. Johansson, K. Nordin, R. Pedersen, N. Karalija, G. Papenberg, M. Andersson, S. M. Korkki, K.
1203           Riklund, M. Guitart-Masip, A. Rieckmann, L. Backman, L. Nyberg, A. Salami, Biphasic patterns of age-
1204           related differences in dopamine D1 receptors across the adult lifespan. *Cell Rep* **42**, 113107 (2023).
1205    122.    M. N. Hallquist, K. Hwang, B. Luna, The nuisance of nuisance regression: spectral misspecification in
1206           a common approach to resting-state fMRI preprocessing reintroduces noise and obscures functional
1207           connectivity. *Neuroimage* **82**, 208-225 (2013).
1208    123.    K. J. Friston, S. Williams, R. Howard, R. S. Frackowiak, R. Turner, Movement-related effects in fMRI
1209           time-series. *Magn Reson Med* **35**, 346-355 (1996).
1210    124.    J. D. Power, K. A. Barnes, A. Z. Snyder, B. L. Schlaggar, S. E. Petersen, Spurious but systematic
1211           correlations in functional connectivity MRI networks arise from subject motion. *Neuroimage* **59**, 2142-2154
1212           (2012).
1213    125.    G. H. Glover, T. Q. Li, D. Ress, Image-based method for retrospective correction of physiological
1214           motion effects in fMRI: RETROICOR. *Magn Reson Med* **44**, 162-167 (2000).
1215    126.    C. Hutton, O. Josephs, J. Stadler, E. Featherstone, A. Reid, O. Speck, J. Bernarding, N. Weiskopf, The
1216           impact of physiological noise correction on fMRI at 7 T. *Neuroimage* **57**, 101-112 (2011).
1217    127.    L. Kasper, S. Bollmann, A. O. Diaconescu, C. Hutton, J. Heinzle, S. Iglesias, T. U. Hauser, M. Sebold,
1218           Z. M. Manjaly, K. P. Pruessmann, K. E. Stephan, The PhysIO Toolbox for Modeling Physiological Noise in
1219           fMRI Data. *J Neurosci Methods* **276**, 56-72 (2017).
1220    128.    D. A. Fair, B. L. Schlaggar, A. L. Cohen, F. M. Miezin, N. U. Dosenbach, K. K. Wenger, M. D. Fox,
1221           A. Z. Snyder, M. E. Raichle, S. E. Petersen, A method for using blocked and event-related fMRI data to
1222           study "resting state" functional connectivity. *Neuroimage* **35**, 396-405 (2007).
1223    129. M. W. Cole, T. Ito, D. Schultz, R. Mill, R. Chen, C. Cocuzza, Task activations produce spurious but
1224           systematic inflation of task functional connectivity estimates. *Neuroimage* **189**, 1-18 (2019).
1225    130. J. Ashburner, A fast diffeomorphic image registration algorithm. *Neuroimage* **38**, 95-113 (2007).
1226    131. V. Fonov, A. C. Evans, K. Botteron, C. R. Almli, R. C. McKinstry, D. L. Collins, G. Brain Development
1227           Cooperative, Unbiased average age-appropriate atlases for pediatric studies. *Neuroimage* **54**, 313-327
1228           (2011).
1229    132. V. S. Fonov, A. C. Evans, R. C. McKinstry, C. R. Almli, D. L. Collins, Unbiased nonlinear average age-
1230           appropriate brain templates from birth to adulthood. *NeuroImage* **47**, S102 (2009).
1231    133. A. Salami, R. Adolfsson, M. Andersson, K. Blennow, A. Lundquist, A. N. Adolfsson, M. Scholl, H.
1232           Zetterberg, L. Nyberg, Association of APOE varepsilon4 and Plasma p-tau181 with Preclinical Alzheimer's
1233           Disease and Longitudinal Change in Hippocampus Function. *J Alzheimers Dis* **85**, 1309-1320 (2022).
1234    134. C. Janiesch, P. Zschech, K. Heinrich, Machine learning and deep learning. *Electronic Markets* **31**, 685-695
1235           (2021).
1236    135. K. He, X. Zhang, S. Ren, J. Sun, Deep Residual Learning for Image Recognition. *arXiv: 1608.06993*,
1237           (2015).
1238    136. V. Nair, G. E. Hinton, paper presented at the Proceedings of the 27th International Conference on
1239           International Conference on Machine Learning, Haifa, Israel,  2010.
1240    137. T. Developers. (https://www.tensorflow.org/, 2022).
1241    138. F. Chollet. (https://keras.io, 2015).
1242    139. I. Sutskever, J. Martens, G. Dahl, G. Hinton, paper presented at the Proceedings of the 30th International
1243           Conference on Machine Learning, Proceedings of Machine Learning Research,  2013.
1244    140. N. Srivastava, G. E. Hinton, A. Krizhevsky, I. Sutskever, R. Salakhutdinov, Dropout: a simple way to
1245           prevent neural networks from overfitting. *Journal of Machine Learning Research* **15**, 1929--1958 (2014).
1246    141. R. R. Selvaraju, M. Cogswell, A. Das, R. Vedantam, D. Parikh, D. Batra, Grad-CAM: Visual Explanations
1247           from Deep Networks via Gradient-Based Localization. *International Journal of Computer Vision* **128**, 336-
1248           359 (2020).
1249    142. D. N. Greve, D. H. Salat, S. L. Bowen, D. Izquierdo-Garcia, A. P. Schultz, C. Catana, J. A. Becker, C.
1250           Svarer, G. M. Knudsen, R. A. Sperling, K. A. Johnson, Different partial volume correction methods lead to
1251           different conclusions: An (18)F-FDG-PET study of aging. *Neuroimage* **132**, 334-343 (2016).
1252    143. A. A. Lammertsma, S. P. Hume, Simplified reference tissue model for PET receptor studies. *Neuroimage*
1253           **4**, 153-158 (1996).
1254    144. R. M. Baron, D. A. Kenny, The moderator-mediator variable distinction in social psychological research:
1255           conceptual, strategic, and statistical considerations. *J Pers Soc Psychol* **51**, 1173-1182 (1986).
1256    145. K. J. Preacher, A. F. Hayes, SPSS and SAS procedures for estimating indirect effects in simple mediation
1257           models. *Behav Res Methods Instrum Comput* **36**, 717-731 (2004).

1258

1259     Supplementary Materials for

1260

1261

1262     **Brain-Cognitive Gaps in relation to Dopamine and Health-related Factors: Insights**

1263     **from AI-Driven Functional Connectome Predictions**

1264

1265     Morteza Esmaeili et al.

1266

1267     Corresponding author: Alireza Salami, Alireza.Salami@kth.se

1268

1269     The supplementary file includes: Figures S1 and S2 and Tables S1 and S2

1270

1271

**Figure S1.** Model trained on functional connectivity maps acquired at rest predicts (**a**) episodic memory (EM) of the test dataset. The models trained on movie-watching (**b**) but not n-back (**c**) datasets predicted EM scores of the test dataset significantly. **Table S1** summarizes the *p* values and correlation power for each model. Test datasets were obtained from the same cohort for rest, movie-watching, and n-back. Bootstrap distributions of correlations between predicted and actual EM scores indicated no significant difference in the predictive power of EM between models trained on resting-state and movie-watching datasets (**d**). Additionally, the bootstrap distribution revealed that models trained on resting-state and movie-watching data yielded higher correlations than those trained on n-back data (**d**).

1280

1286  ***Table S1.*** *Correlation results for episodic memory (EM) score predictions – DyNAMiC dataset with Schaefer300*

1287  *parcellation.*

| Model trained on | rest | | | movie | | | n-back | | |
|---|---|---|---|---|---|---|---|---|---|
| **Model tested on** | rest | movie | n-back | rest | movie | n-back | rest | movie | n-back |
| Correlation | .46 | .38 | .36 | .34 | .39 | .09 | .06 | .15 | .14 |
| Significance | .0002 | .004 | .005 | .007 | .002 | .47 | .67 | .25 | .28 |
| | | | | | | | | | |
| COBRA | | | | | | | | | |
| Correlation | N.A. | | | | | | | | |
| Significance | N.A. | | | | | | | | |

1288

1289

1290

1291

**Figure S2.** Model trained on functional connectivity maps acquired at rest did not significantly predict (**a**) the

working memory (WM) of the test dataset. The corresponding model best performed in WM scores predictions

while trained on the movie-watching data set (**b**). The model trained on the n-back dataset (**c**) was the second-

best, predicting WM scores. **Table S2** summarizes the $p$ values and correlation power for each model. Test

datasets for resting-state, movie-watching, and n-back tasks were derived from the same cohort. Bootstrap

distributions of correlations between predicted and actual WM scores showed no significant difference in

predictive power between models trained on movie-watching and n-back data (**d**). Furthermore, the bootstrap

distribution revealed that models trained on movie-watching and n-back data exhibited higher correlations than

those trained on the resting state dataset (**d**).

1301

1302

1303

1304    ***Table S2.*** *Correlation results for working memory (WM) score predictions – DyNAMiC dataset with*

1305    *Schaefer300 parcellation.*

| Model trained on | rest | | | movie | | | n-back | | |
|---|---|---|---|---|---|---|---|---|---|
| **Model tested on** | rest | movie | n-back | rest | movie | n-back | rest | movie | n-back |
| Correlation | .16 | .12 | .10 | **.33** | **.55** | **.43** | **.38** | **.46** | **.52** |
| Significance | .27 | .36 | .43 | **.01** | **<.0001** | **.006** | **.003** | **.0002** | **<.0001** |
| COBRA | | | | | | | | | |
| Correlation | | | | | | N.A. | | | |
| Significance | | | | | | N.A. | | | |

1306

1307