Title: Actomyosin forces trigger a conformational change in desmoplakin within desmosomes.

Authors: Yinchen Dong [1], Ahmed Elgerbi [3], Bin Xie [2], John S. Choy [3], Sanjeevi Sivasankar [1, 2, *]

Affiliations: [1] Department of Biomedical Engineering, University of California, Davis, CA; [2]

Biophysics Graduate Group, University of California, Davis, CA; [3] Department of Biology, The

Catholic University of America, Washington DC.

*Corresponding author: ssivasankar@ucdavis.edu

Abstract: Desmosomes are essential cell-cell adhesion organelles that enable tension-prone

tissue, like the skin and heart, to withstand mechanical stress. Desmosomal anomalies are

associated with numerous epidermal disorders and cardiomyopathies. Despite their critical role

in maintaining tissue resilience, an understanding of how desmosomes sense and respond to

mechanical stimuli is lacking. Here, we use a combination of super-resolution imaging, FRET-

based tension sensors, atomistic computer simulations, and biochemical assays to demonstrate

that actomyosin forces induce a conformational change in desmoplakin, a critical cytoplasmic

desmosomal protein. We show that in human breast cancer MCF7 cells, actomyosin contractility

reorients keratin intermediate filaments and directs force to desmoplakin along the keratin filament

backbone. These forces induce a conformational change in the N-terminal plakin domain of

desmoplakin, converting this domain from a folded (closed) to an extended (open) conformation.

Our findings establish that desmoplakin is mechanosensitive and responds to changes in cellular

load by undergoing a force-induced conformational change.

Introduction

Desmosomes are critical cell-cell junctions that promote adhesion between cells and preserve tissue integrity. They enable tension-prone tissues like the skin and heart to withstand mechanical stress [1]. Desmosomal anomalies are associated with numerous diseases such as epidermal autoimmune disorders and arrhythmogenic ventricular cardiomyopathy [2-4]. Despite their critical role in maintaining tissue mechanical resilience, an understanding of how desmosomes respond to mechanical stimuli is lacking.

Desmosomes are composed of transmembrane desmosomal cadherins, desmoglein (Dsg) and desmocollin (Dsc), and cytoplasmic plaque proteins: plakophilin, plakoglobin, and desmoplakin (DP). The extracellular regions of the cadherins from opposing cells interact to mediate adhesion, while the plaque proteins bind to the cadherin cytoplasmic tail (Fig. 1A) [5]. DP anchors desmosomes to the keratin intermediate filament (KIF) cytoskeleton, and this linkage is essential for maintaining tissue integrity and facilitating intracellular signal transduction [6, 7]. Several studies show that KIFs work in cooperation with the actomyosin cytoskeleton to modulate the mechanical state of desmosomes [8-10]. Disassociating KIFs from circumferential actin belts alters tension across desmosomes [8]], while disassembling the actin cytoskeleton decreases desmosome resilience [9]. However, the molecular mechanisms by which actomyosin forces alter desmosome structure and function are not understood.

Disruption of the linkage between DP and KIF impairs intercellular adhesion and compromises tissue integrity [10, 11]. Knocking-out keratin leads to elevated phosphorylation of DP, resulting in accelerated endocytosis of desmosomes and rupture of epithelial sheets under rotational stress [12] . Weakening connections between DP and KIF networks dramatically alters the mechanical properties of cells, such as cell stiffness and intercellular forces [9]. While these data suggest that

DP is crucial in regulating the response of desmosomes to mechanical stress, the biophysical

mechanism by which DP mechanically regulates the desmosome is unknown.

The structure of DP offers hints into its mechanoregulatory function. DP is a large, 2871 amino

acid protein composed of an N-terminal region, a central rod domain that facilitates DP

homodimerization [3], and a C-terminal region that is crucial for aligning and binding KIFs [13].

The N-terminal region of DP is subdivided into a globular head domain and a plakin domain, each

playing distinct roles [14]. The globular head domain binds to the C-termini of desmosomal

cadherins and recruits plakophilin and plakoglobin [15]. The plakin domain, characterized by six

spectrin repeat domains interspersed with Src homology domains, exhibits structural flexibility, as

observed in a previous *in vitro*, cell-free small angle X-ray scattering study [16]. That investigation

revealed that the plakin domain adopts either a folded "U" shape or a fully extended "I" shape,

indicating a significant degree of structural adaptability [17]. This finding suggests that actomyosin

forces could induce a conformational change in the N-terminal region of DP, thereby offering a

molecular pathway for DP mechanosensation. However, force-induced conformational changes

in DP, particularly within the cell, have never been demonstrated.

DP exists in two major splice isoforms, I and II, which differ only in the length of their rod domain.

Our study focuses on DP isoform I (DPI) which is expressed in high levels in epithelia and

cardiomyocytes [18]. We demonstrate, in human breast cancer MCF7 cells, that actomyosin

forces induce a conformational change in the N-terminal region of DPI. Since MCF7 cells are

highly enriched in keratin 19 (K19), we interrogated wild type (WT), K19-knock out (K19-KO), and

K19-GFP rescue (K19-GFP) cells with super-resolution microscopy, FRET-based tension

sensors, pharmacological inhibition and activation of actomyosin contractility, co-

immunoprecipitation (co-IP), and atomistic Molecular Dynamics (MD) simulations. We

demonstrate that actomyosin contractility reorients the KIF and directs force to DPI along K19

filaments. We show that these forces induce a conformational change in the N-terminal plakin

domain, from a folded 'closed' to an extended 'open' conformation. Our findings establish, at the

molecular level, that DP is a mechanosensor within the desmosome.

Results

K19 remodels desmosome architecture: Desmosomes are characterized by distinct "railroad"

track structures, represented by discrete parallel fragments of DP, which sandwich the

desmosomal cadherins [19]. To validate this signature structure of desmosomes in MCF7 cells,

we conducted immunostaining targeting the extracellular region of isoform 2 of desmoglein (Dsg2)

and the C-terminal region of DPI (DPC) using confocal and super-resolution Stimulated Emission

Depletion (STED) microscopy. Confocal imaging revealed co-localization of DPC and DSG2

signals, indicating the formation of desmosome puncta along cell–cell junctions (Fig. 1B). STED

imaging in WT cells showed well-defined DP "railroad" tracks, as depicted in Figs. 1C-D. To

investigate the effect of K19 on desmosome architecture, we used the K19-KO cell line [20].

Successful ablation of K19 in these cells was confirmed using western blotting (Fig. S1).

Surprisingly, in the K19-KO cells, STED imaging revealed more compact desmosomes (Fig. 1E).

To quantify the width of the desmosomes, we measured the distance between opposing DP

plaques in the STEAD images by drawing a line scan across the desmosome complex (Fig. 1G)

and measuring the distance between the two peaks of DPC signal (designated as DPC-DPC

distance), as illustrated in the fluorescence intensity plot (Fig. 1H). This quantitative analysis

confirmed that desmosomes in K19-KO cells were significantly narrower than desmosomes in WT

MCF7 cells (334 ± 61 nm for WT cells and 264 ± 65 nm for K19-KO cells; Fig. 1I).

To corroborate that the observed shift in desmosome width was directly related to the absence of

K19, we stably rescued the K19-KO cells with EGFP-tagged K19 (K19-GFP cells). Western blots

validated the successful expression of K19-GFP in the rescued cells (Fig. S2). STED imaging

showed that desmosome width in the K19-GFP cells recovered to a level similar to the WT cell

line (322 ± 56; Figs. 1F & 1I). Together, these findings demonstrate that desmosome architecture

in MCF7 cells is strongly correlated to the presence/absence of K19. In the presence of K19, the

desmosome width is larger, while the desmosome narrows when K19 is knocked out.

The interplay between K19 and the actomyosin networks governs desmosome width: The

cytoplasmic KIF network binds to DP at its C-terminus (illustrated in Fig. 2A) and interactions

between the KIF network and elements of the actomyosin cytoskeleton have been observed at

cell-cell junctions and the cell-substrate interface [8, 21]. To investigate how changes in the KIF

network contribute to desmosome remodeling, we first inspected the intracellular localization of

K19, DP, and F-actin filaments in K19-GFP cells. K19 was found to be colocalized with the DP

"railroad" tracks along the cell border (Fig. 2B). Additionally, actin filaments were observed to

align with the K19 filaments (Fig. 2B). Based on the colocalization of K19 with DP and the

coalignment of K19 with F-actin filaments, we hypothesized that K19 interacts with both DP and

F-actin filaments.

To validate this hypothesis, we conducted Co-IP experiments using IgG as the control and K19

as the bait protein. The results revealed that K19 interacts with both DP and actin, consistent with

our previous immunofluorescence microscopy observations (Fig. 2C). To confirm that the proteins

pulled down with K19 were due to specific interactions, we performed a K19 pulldown in K19-KO

cells. Neither DP nor actin filaments were detected in the pulldown from K19-KO cells (Fig. S3),

supporting that their association with K19 in WT cells reflect specific interactions. To further

confirm the specificity of K19 interactions with DP and actin, we also blotted for the membrane-

associated protein E-cadherin in both WT and K19-KO cells. The absence of E-cadherin in the

pull-down indicated that associations with DP and actin were not due to general membrane

localization but are specific to K19 (Fig. S4). Finally, the Co-IP experiments showed that in

addition to DP, the desmosomal plaque protein plakophilin-3 also interacted with K19 (Fig. S5),

which further supports the interaction between K19 filaments and the desmosome complex.

It has previously been demonstrated that actin stress fiber formation strongly correlates with the

presence of keratin filaments in cells [22]. We therefore used confocal fluorescence microscopy

to image if actin stress fibers, which are a proxy for intracellular forces, are present in WT, K19-

GFP, and K19-KO cells (Fig. S6A). Quantitative data analysis showed that compared to WT and

K19-GFP cells, actin stress fibers in the K19-KO cells were dramatically decreased (Fig. S6B).

This suggests that mechanical tension in the K19-KO cells is lower compared to WT and K19-

GFP cells. We, therefore, hypothesized that actomyosin-generated forces could be transmitted to

the desmosome complex through associations with the KIF network.

To test this hypothesis, we first used Blebbistatin, a myosin-II-specific ATPase inhibitor, to reduce

the actomyosin contractility in the WT cell line (WT+Blebb) and measured the desmosome width.

STED imaging showed more compact desmosomes in the WT+Blebb cells (Fig. 2D); desmosome

width in WT+Blebb cells was reduced to a similar level as K19-KO cells (263 ± 62 for WT+Blebb;

Fig. 2E). At the same time, the control group treated with dimethyl sulfoxide (DMSO, Blebbistatin's

solvent), retained similar desmosome widths as the WT cells (Fig. S7).

Next, to test if stimulating actomyosin contractility increases desmosome width, we treated the

K19-KO cells with Calyculin A, a phosphatase inhibitor (KO+Caly). STED imaging showed that

the width of desmosomes in the KO+Caly cells increased to a level comparable to WT cells

(Fig.2F). Quantitative analysis of the images showed that upon Calyculin A treatment, the

desmosome width in K19-KO cells was 333 ± 83 (Fig. 2G). Taken together, these data

demonstrate that the K19-dependent changes in desmosome architecture are a consequence of

a mechanical pathway that is established due to interactions between K19, F-actin, and DP.

Consequently, contractile actomyosin forces are transmitted to and reorganize the desmosome,

resulting in an increase in desmosome width.

The N-terminal plakin domain of DP transitions from a closed to open conformation when

pulled: Since previous studies have reported structural flexibility in the N-terminal plakin domain

of DP [17], we hypothesized that the increase in desmosome width in WT cells was due to a force-

induced conformational change in this region of DP (Fig. 3A).To test this hypothesis, we

specifically stained both the DPI N-terminal region (DPN) on its globular head domain (which is

upstream of the plakin domain) and the extracellular region of Dsg2. Using super-resolution

imaging, wider desmosomes were observed in WT cells stained with DPC and DSG2 compared

to K19-KO cells (Fig.3B). Interestingly, when stained with DPN and DSG2, both cell lines exhibited

similar desmosome widths (Fig.3C). To gain a clearer understanding of the DP structural

changes in WT and K19-KO cell lines, we measured the distances between Dsg2 and DPC, as

well as Dsg2 and DPN within the half-unit of the desmosome complex. This analysis showed that

while Dsg2-DPC distance was significantly greater in WT compared to the K19-KO cells (167 ±

47 for WT cells and 132 ± 50 for K19-KO cells; Fig. 3D), Dsg2-DPN distance was similar in both

cell lines (101 ± 48 for WT cells and 99 ± 53 for K19-KO cells; Fig. 3E). As seen in Fig. 3G, the

DP length (defined as the distance between DPN and DPC) was 33 nm greater in the WT cells

compared to the K19-KO cells, suggesting that DP was elongated by ~33 nm in the WT cell lines

due to higher mechanical tension in these cells.

To resolve the mechanistic basis by which DP transitions between open and closed structures,

we proceeded to atomistically simulate the effect of force on the conformation of the highly

conserved plakin domain of DP (residues 178-950). We first generated the entire plakin domain

structure using Alphfold2 [23]. The predicted structure exhibited a 'U-shaped' folded conformation,

with spectrin repeat (SR) domains SR3-6 and Src homology (SH) domain SH3 forming a long arm

that interacts with a short arm composed of SR7-8 and SR8-CT domains (Fig. 3F). This prediction

agrees well with the crystal structure of the long arm of the plakin domain [24] (RMSD ~1.7Å, Fig.

S8).

To identify the interactions crucial for stabilizing the folded conformation, we performed three

force-independent molecular dynamics (MD) simulations on the predicted structure. In the MD

simulations, the plakin domain stabilized within 20 ns (Fig. S9). During the simulations, 10-15

hydrogen bonds were observed between the long arm and the short arm, and the interactions

mainly were observed between SR5-6 and SR7-8 regions (Fig. 3H and Fig. S10). Notably, two

consistent salt bridge/hydrogen bond interactions, between four conserved amino acids, were

observed across all three simulations: R592-N744 and K444-E858 (Fig. 3F).

To test if the folded plakin domain structure can be unfolded using pulling forces, we performed

steered molecular dynamics (SMD) simulations (Video S1). In the SMD simulations, we fixed the

position of the N-terminal residues and pulled the C-terminal regions (residues 919-950) using a

constant force. Our simulations showed that the plakin domain's long and short arms separated

due to the pulling force without any secondary structure unfolding (Fig. 3J). Consequently, the

plakin domain transitioned from the initial closed (Fig. 3J, left) to the final open (Fig. 3J, right)

conformation (Video S1). In the open conformation, the distance between the plakin domain's C-

terminus and N-terminus increased by 33 nm compared to the closed structure (Fig. 3I), which is

consistent with the change in DPC-DPC distance observed in the STED images (Fig. 3G).

Keratin intermediate filaments reorient in response to changes in actomyosin forces: Given
that actomyosin forces are transmitted to desmosomes via KIFs, a key question is how this force
transmission is affected by the organization of keratin filaments. A previous study in *Xenopus*
laevis showed that the KIF network reorganizes in response to force transmission across cell-cell
contacts and becomes aligned with the direction of protrusive activity and tissue movement [25].
This raises the question as to whether a similar phenomenon occurs in MCF7 cells (Fig. 4A).

In epithelial cells, KIFs are composed of heterodimers between type I and type II keratins, which
assemble into antiparallel tetramers [26]. A previous RNA-sequencing study revealed that type II
Keratin-8 (K8) was the most highly expressed isoform in the MCF7 cells, followed by roughly
equal amounts of type I Keratin-19 (K19) and type I Keratin-18 (K18) [27]. This suggests that the
primary keratin pairs in MCF7 cells are K8/K19 and K8/K18. Notably, K8 and K18 expression
levels remained unchanged when K19 is knocked out (Fig. S1), which allowed us to stain K8 and
K18 to investigate KIF organization. When we performed Co-IP experiments using K8 and K18
as bait, we observed that both keratin isoforms interacted with DP and actin filaments in WT and
K19-KO cells (Fig. 4B). This suggested that K8/K18 filaments could also transmit actomyosin
forces to desmosomes.

We first imaged the organization of K8 filaments in WT and K19-KO cells by staining both K8 and
DP. In WT cells, the K8 filaments at junctions exhibited a more radial organization (Fig. 4C),
whereas, in K19-KO cells, the K8 filaments were primarily arranged horizontally (Fig. 4D). We
quantified filament organization by determining their local orientation, defined as the angle of the
most dominant filament direction with the primary axis of the cell junction (which was set as 0°),
and aggregated the local orientations into a polar histogram (Fig. 4C-D). Our quantitative analysis
showed that over 60% of WT K8 filaments had a dominant orientation between 40° and 90°,
whereas more than 60% of K19-KO K8 filaments were oriented between 0° and 20° (Fig. 4E).

When actomyosin force was inhibited via Blebbistatin treatment, the K8 filament orientation in the

WT cells changed dramatically with over 70% of filaments oriented between 0° and 20° (Fig. 4F).

In contrast, K19-KO cells treated with Blebbistatin showed no significant change in K8 filament

orientation (Fig. 4F). On the other hand, when actomyosin contractility in the cells was enhanced

using Calyculin A treatment, K8 filament organization in K19-KO cells changed dramatically with

over 60% of filaments having a dominant filament orientation between 40° and 90° (Fig. 4G). In

contrast, K8 filament orientation in WT cells treated with Calyculin A remained largely unchanged

(Fig. 4G). At the same time, the control group treated with Blebbistatin and Calyculin A's solvent,

DMSO, retained similar keratin organizations as the WT and K19-KO cells (Fig. S11).

Next, to distinguish between K8/K19 and K8/K18 filaments, we also imaged K18 filament

organization in WT and K19-KO cells. As observed in our K8 filament images, staining for K18

and DP revealed more radially organized filaments in WT cells and horizontally organized

filaments in K19-KO cells (Fig. 4H-J). Taken together, these findings indicated that KIFs

reorganize in response to changes in actomyosin forces, shifting from predominantly 'poor force-

transmitting' horizontal orientation in a low-tension condition to a more 'efficient force-transmitting'

radial organization under load. Consequently, K19-KO and WT+Blebb cells exhibit horizontally

oriented KIF network and more compact desmosomes while WT cells and KO+Caly display a

radially organized KIF network and wider desmosomes.

Desmoplakin is mechanically loaded in WT cells: Next, we proceeded to directly compare

tension across DP in WT and K19-KO desmosomes using a recently published DPI Förster

Resonance Energy Transfer (FRET) tension sensor [28]. This F40-based tension sensor (TS), is

comprised of a tension-sensing linker peptide $(GPGGA)_8$ flanked by the donor fluorophore mTFP1

and the acceptor fluorophore mEYFP, that is inserted into an unstructured region following the

rod domain, just before Pro1946 in DPI (Fig. 5A) [28]. Additionally, we used a no-tension control

consisting of a truncated DP that lacked the C-terminal keratin-binding domain to account for local

environmental changes that may affect FRET efficiencies (Fig. 5A). While these truncated DP

controls still localized to intercellular junctions, they were unable to interact with keratin filaments

and thus could not bear forces [11, 28, 29]. Lastly, a donor-only control was also included to

determine the FRET efficiency. We transiently expressed the DP constructs in WT and K19-KO

cells and analyzed the fixed samples. DP-TS localized to cell-cell contacts, with distinct DP puncta

observed in both WT and K19-KO cells (Fig. 5B). Similarly, the truncated DP control and donor-

only control exhibited the expected subcellular localization at cell junctions, forming clear

desmosomal puncta (Fig. S12). These results indicate that all three DP constructs were correctly

localized to cell-cell contacts and successfully formed desmosomes, enabling the measurement

of DP tension at the desmosome.

Next, we quantified DP-TS FRET efficiencies using fluorescence lifetime imaging microscopy

(FLIM) in confluent WT and K19-KO monolayers following the workflow illustrated in Fig. 5C.

When imaging each transfected cell, we first isolated individual DP puncta and measured their

lifetime using a bi-exponential fit, accounting for the effects of the instrument response function

(IRF) and autofluorescence, which yielded short lifetime measurements. Finally, FRET efficiency

was determined based on these lifetime measurements. In the WT cells, the FRET efficiency was

significantly lower in the TS (8.80% ± 2.76%) compared to the control (18.5% ± 4.46%), indicating

that DP in these cells experienced substantial mechanical loading (Fig.5D). In contrast, the FRET

efficiency of TS (18.3% ± 3.96%) in K19-KO cells was similar to the control (19.0% ± 3.88%) (Fig.

5E), suggesting that no detectable tension was present across DP in the K19-KO cells. Based on

the 1–6 pN sensitivity of the F40 tension sensor and its averaged FRET–force calibration curve

[28, 30], we estimated that the tension in WT cells was close to the upper range, around 5 pN.

These direct measurements of forces experienced by DP revealed high tension in WT cells and

no detectable tension in K19-KO cells, consistent with our previous analysis of actin stress fibers.

Discussion

Based on our data, we propose a model (Fig. 5F), where in the absence of tension, DP-associated

keratin filaments align horizontally, maintaining a closed DP conformation. Under stress, these

filaments adopt a radial orientation, transmitting forces to the desmosome and inducing a

conformational shift in DP via its flexible plakin domain (Fig. 5F). Several recent studies have

highlighted the potential mechanosensitive role of DP in desmosomes. These studies

demonstrate that that DP experiences tension when cells are subjected to mechanical pulling [28]

and show that the DP-KIF linkage is essential for resisting mechanical strain [9, 31]. Building on

these findings, we demonstrate that DP functions as a mechanosensor by undergoing a

conformational change in response to alterations in cytoskeletal tension. The DP conformational

change described in our study is analogous to conformational changes in α-catenin, associated

with classical cadherins in Adherens Junctions [32-34]. In the case of α-catenin, tension from the

actin cytoskeleton induces a conformational change that opens its central M-domain and exposes

cryptic binding sites for vinculin and other actin-binding proteins [32-34]. It is likely that the force-

induced opening of DP also exposes cryptic binding sites for junctional proteins that signal and

remodel the desmosome and intermediate filaments. The force-induced unfolding of DP could

also serve as a damper of mechanical stress that maintains mechanical homeostasis within

tissue.

Our data suggests that besides actomyosin forces, conformational changes in DP are also

mediated by changes in keratin reorganization. Since Co-IP experiments suggest that both K8

and K18 associate with actin and DP, it is likely that K8/K18 filaments also transmit actomyosin

forces to DP in the K19-KO cells. Why then do the DPs in the K19-KO cells remain in a closed

conformation? Our data suggests that the loss of K19 alters the organization of keratin filaments,

which in turn affects the efficiency of actomyosin force transmission to desmosomes. We show

that in the K19-KO cells, K8/K18 filaments are oriented cortically, almost parallel to the

desmosome, which result in tangentially actomyosin forces which cannot open DP. In contrast,

since the KIFs are oriented radially in WT cells, desmosomes are exposed to normal forces which

can induce a conformational change in DP. Indeed, using FRET tension sensors we demonstrate

that DP in the WT cells are exposed to higher tension than DP in the K19-KO cells. We confirmed

that the force-induced widening of desmosomes was not due to differences in the expression

levels of desmosomal or cytoskeletal proteins between the two cell lines by comparing the

amounts of essential desmosomal and cytoskeletal components in both WT and K19-KO cells.

Immunoblotting results showed no significant differences in the levels of desmosomal cadherins

(Dsg2, Dsc1, and Dsc2-3), plaque proteins (plakophilin-1, plakophilin-3, and DP), or cytoskeletal

proteins, including K8 and K18, F-actin, and myosin heavy chain 10. These results indicate that

the observed changes in desmosome organization under force are not due to altered protein

expression levels but rather the result of modified mechanical conditions driven by the cooperative

interactions between KIF and the actomyosin network.

Earlier studies using FRET-based tension sensors in resting MDCK cells revealed a small amount

of tension across Dsg2 but not across DP [28, 35]. However, the Dsg2 FRET-based tension

sensor showed a higher tension level in the contracting cardiac myocytes, suggesting that

junctional tension depends on both cell type and the specific cellular environment [35]. To validate

our analysis, we also measured DPC-DPC distances in resting MDCK cells. Our measurements

revealed that the desmosome width in MDCK cells is comparable to that observed in K19-KO

cells, implying that DP in resting MDCK cells indeed does not undergo conformational changes

(Fig. S13). Additionally, FRET-based tension measurements in MCF7 cells, using the same DPI

constructs as in previous studies, revealed high levels of tension across DP in WT cells but no

detectable tension in K19-KO cells. These FRET-based tension measurements, along with the

DPC-DPC distance data, support a correlation between desmosomal tension level and

desmosome width, demonstrating that wider desmosomes result from higher tension across the

desmosomes.

Previous rotary shadow imaging of DPI reported an average C-to-N-terminal distance of 162 nm

[36]. In contrast, transmission EM imaging of desmosomes in bovine nasal epidermis reported a

DP length of ~42 nm [37] while super-resolution dSTORM microscopy in human epidermal

keratinocytes measured DP lengths of ~54 nm [19] which suggests that DP is tilted at an angle

rather than being strictly perpendicular to the plasma membrane. In agreement, our STED

measurements reveal a DPI length, based on the antibody-labeling, of approximately 66 nm in

WT cells and 33 nm in K19-KO cells. Importantly, since we measure relative distance changes

between the N- and C- termini of DP, our measurement of force-induced DP conformational

changes remains valid irrespective of the tilted orientation of DP.

Furthermore, based on super-resolution microscopy, a model has been recently proposed where

DP increases its tilt as desmosomes mature and become more adhesive, such that narrower and

longer desmosomes correspond to stronger adhesive states [19, 38]. Since we observe narrower

desmosomes in the K19-KO cells and wider desmosomes in WT cells, we assessed the

intercellular adhesion strength of these cell lines using a dispase assay. In our assay, confluent

sheets of either WT or K19-KO cells were detached using dispase II, a protease that gently

cleaves cells from the substrate. Since cell-cell adhesion is mediated via both Ca^{2+}-dependent

classical cadherins (like E-cadherin) [39] and desmosomes which adhere in both a Ca^{2+}-

independent state and a Ca^{2+} dependent mode [40], we incubated the lifted cell sheets with 4 mM

EGTA, a chelating agent for Ca^{2+}, for 1 hour. Consequently, our assay selectively evaluated

intercellular adhesion mediated mainly through the Ca^{2+}-independent desmosomes. Following

the EGTA treatment, the cell sheets were subjected to mechanical stress, resulting in the

fragmentation of the sheets, as shown in Fig. S14A. Our results showed that the K19-KO cell

sheets exhibited a significantly greater number of fragments than WT cell sheets. In contrast,

K19-GFP cell sheets generated a similar number of fragments compared to WT cell sheets (Fig.

S14B). These results suggest that in MCF7 cells, wider desmosomes do not correspond to

weaker adhesive states. Furthermore, since the proposed model also suggests that desmosomes

lengthen as they become more adhesive, we measured the DP length along the cell borders in in

WT and K19-KO cells (*i.e.* $\frac{desmosome\ length}{cell}$). In contrast to the proposed model, our results showed

a slight decrease in DP length in the weakly adhesive K19-KO cells compared to the strongly

adhesive WT cells (Fig. S15). These results support our findings that changes in desmosome

width are a direct result of DP conformational changes which are driven by actomyosin

contractility and KIF organization. Our findings are also supported by a vast body of literature

demonstrating that keratin deficiency is associated with impaired intercellular adhesion and

increased cellular invasiveness [6, 41, 42].

Previous SMD simulations carried out on the long arm of the plakin domain (i.e., the crystal

structure with PDB code 3R6N [24]) show that the SRs individually unfold when subjected to

pulling force, which exposes the SH3 domain [43, 44]. Since our simulations are carried out on

the entire plakin domain, which includes the flexible hinge between the long and short arms, we

capture the transition of the plakin domain from a closed to an open conformation. The

conformational change we measure is supported by previous SAXS measurements [17]. This

transition of the plakin domain from a closed to an open structure also partially exposes the cryptic

SH3 domain, which is known to act as a signal transduction adapter crucial in cell invasiveness,

migration, and actin reorganization [45].

Many disease-causing mutations in DP are found within the plakin domain of the N-terminal

region, affecting residues both hidden within the protein and exposed to the surrounding

environment [46]. Specifically, mutations within the SR7-8 region of the plakin domain have been

linked to arrhythmogenic ventricular cardiomyopathy (AVC) [16, 46]. Importantly, our simulations

predict that the SR7-8 region participates in several key hydrogen bonds with SR5-6, which in

turn stabilizes DP in its closed conformation. This raises the intriguing possibility that AVC

mutations may interfere with the force-induced structural transition of DP from the closed to the

open conformation.

Materials and Methods

Cell culture and stabilization: MCF7 WT cells (ATCC), MCF7 K19-KO cells, and MCF7 K19-

GFP rescued cells were maintained in low-glucose Dulbecco's Modified Eagle's Medium (DMEM)

(Gibco) supplemented with 10% Fetal Bovine Serum (FBS) (Life Technologies) and 1% penicillin-

streptomycin (Sigma-Aldrich) in 5% CO_2 at 37°C. The rescued cells were additionally treated with

100 ug/ml hygromycin for selection of K19-GFP. MDCK cells (ATCC) were cultured in low glucose

DMEM (Gibco) cell culture media with 10% FBS and 1% PSK.

The K19-KO cells were generated using CRISPR/Cas9 as previously described [47]. We

developed the K19-GFP rescue cell line following protocols described previously [48]. Briefly,

K19-GFP was cloned into *K*19-KO cells from pEGFP-K19 plasmid into pLenti CMV Hygro

(plasmid   #17484.   Addgene)   using   the   following   primers: forward

5'TCAGTCGACTGGATCCATGGTGAGCAAGGGCGAGG3'   and   reverse

5'GAAAGCTGGGTCTAGTCAGAGGACCTTGGAGGC3'. Lentiviral supernatants were generated

using pLenti-K19-GFP plasmid. Lentiviral supernatants collected 24h post-transfection were used

to infect subconfluent K19-KO cells in three sequential 4 h incubations in the presence of 4 ug/ml

polybrene (Sigma-Aldrich). Transductants were selected using a culture medium containing 100

ug/ml hygromycin beginning 48h after infection. The K19-GFP rescue cells were selected for

further stabilization using fluorescence-activated cell sorting.

Preparation of STED and confocal microscopy imaging samples: Approximately 500,000

cells were seeded for all the MCF7 cell lines, and approximately 250,000 cells were seeded for

MDCK cells on collagen-coated coverslips (22 mm, No.1.5) in a single well of a six-well plate and

were kept in 5% CO2 at 37°C for 24 h. A 'calcium switch' was then performed using the protocol

described below to eliminate cell clusters and maintain a similar level of desmosome maturation

across all samples. To perform a calcium switch, the cells were first rinsed with phosphate-

buffered saline (PBS) (Gibco) solution at room temperature, followed by incubation in 4 mM EGTA

(Millipore Sigma) and 10% low Ca^{2+} Fetal Bovine Serum (FBS) in Ca^{2+}-free DMEM for 30 min.

Following EGTA treatment, the cells were switched to a DMEM medium with 2 mM $CaCl_2$ for 24

h to trigger the formation of mature desmosomes.

Next, cells were fixed either with ice-cold 1:1 methanol and acetone for 10 min (for keratin and

DP staining) or with 3% PFA and 0.3% Triton X-100 for 10 min at room temperature (RT) (for F-

actin staining). Samples were rinsed with PBS three times and blocked in 1% Bovine Serum

Albumin (BSA) overnight at 4°C. Samples were then incubated with primary antibody (1:200) for

1 h at RT, followed by incubation with secondary antibody (1:600) for 30 min at RT. The samples

stained with fluorescently labeled phalloidin (1:400) were incubated with phalloidin for 30 min at

RT in the dark. Samples were washed three times with PBS after each antibody incubation. Lastly,

samples were mounted to the glass slides using the ProLong Diamond Antifade Mountant (Life

Technologies) and stored at 4°C in the dark.

Drug treatments

Cells were cultured as previously described. For the blebbistatin treatment, cells were treated with 50 μM (±) blebbistatin (Millipore Sigma) for 24 h. Blebbistatin was dissolved in DMSO and diluted in the low-glucose DMEM (Gibco) supplemented with 10% FBS (Life Technologies) and 1% penicillin-streptomycin (Sigma-Aldrich) media. The control group was prepared with the same protocol but treated with 0.1% DMSO instead. For the Calyculin A treatment, cells were treated with 100 nM Calyculin A (Millipore Sigma) for 15 min. Calyculin A was dissolved in DMSO and diluted in the low-glucose DMEM (Gibco) supplemented with 10% FBS (Life Technologies) and 1% penicillin-streptomycin (Sigma-Aldrich) media. The control group was prepared with the same protocol but treated with 0.1% DMSO instead.

Antibodies: The following primary antibodies were used: human anti-DSG2 (MAB947, R&D systems), rabbit anti-DPC antibody (A303-356A, Bethyl Lab), rabbit anti-DPN antibody (25318-1-AP, Proteintech), chicken anti-GFP (600-901-215, Rockland), mouse anti-K19 antibody (A53-B/A2) (Santa Cruz Biotechnology), mouse anti-K8 antibody (MA1-06318, ThermoFisher), mouse anti-K18 antibody (MA1-19047), control mouse IgG (sc-2025) (Santa Cruz Biotechnology), and mouse anti-actin antibody (66009-1-Ig, Proteintech). The following secondary antibodies are used: anti-chicken AF-488 (Invitrogen), anti-mouse AF-594 (Invitrogen), and anti-rabbit AF-647 (Invitrogen). Phalloidin AF-568 (Invitrogen) was used for F-actin staining.

The following antibodies were used for the Co-IP and western blot experiments: Anti-Keratin 19 antibody (A53-B/A2), anti-plakophilin-3 (E-10), anti-plakophilin-1 (10B2), anti-desmocolin-1 (A-4), anti-desmocolin-2/3 (7G6), anti-myosin-10 (A-3), and control mouse IgG (sc-2025) from Santa

Cruz Biotechnology (Santa Cruz, CA), anti-Keratin 8 (17514-1-AP), ani-Keratin 18 (10830-1-AP),

anti-actin antibody (66009-1-Ig), anti-desmoglein-2 (21880-1-AP), anti-plectin (29170-1-AP), anti-

tubulin (66031-1-Ig), and anti-desmoplakin antibody (25318-1-AP) from Proteintech (Rosemont,

IL).The following secondary HRP conjugated antibodies were used: anti-mouse-HRP (A4416),

and anti-rabbit-HRP (A6154) from Sigma Aldrich (Louis, MO).

STED imaging acquisition and processing: Immunofluorescence images were acquired at RT

using a Leica TCS SP8 STED 3X confocal microscope (Leica Microsystems) equipped with a

100-x oil immersion objective (HC PL APO CS2), white light laser for excitation lines between

470-670 nm, and depletion laser lines at 592, 660, and 775 nm. The acquisition was performed

using LasX software, followed by deconvolution in Huygens Pro software (Scientific Volume

Imaging). Post-acquisition processing, including cropping and resizing, color change in the lookup

table, adjustment of brightness and contrast, and inversion of grayscale images, was performed

using ImageJ software.

Confocal image acquisition and processing: Immunofluorescence images were acquired

using a Leica Stellaris 5 confocal microscope (Leica Microsystems) equipped with a 63x oil

immersion objective (HC PL APO CS2). Image acquisition was performed using LasX software,

followed by post-processing using ImageJ. During post-processing, images were adjusted with

inversion of grayscale, enhanced contrast with 0.3% saturated pixels, and reduced background

noise using the despeckle functionality. All images were acquired using the same procedure.

Distance measurement analysis: We used custom-made ImageJ macros and Python scripts to quantify the desmosome width along the entire cell-cell border. Since Dsg2 signals colocalize between DP "railroad" tracks, they were used as the reference to indicate desmosome location along the membrane. Briefly, Dsg2 signals were selected as regions of interest (ROIs), and an algorithm was used to determine the midline of the ROIs along the membrane axis and draw a segment along the midline every 200 nm to obtain the intensity profile of both Dsg2 and DPC or Dsg2 and DPN. Then, Bar, an ImageJ Plugin, was used to determine the location of peaks of different signals in their respective channels along the drawn segment. After obtaining the raw data, we used custom-made MATLAB code to calculate distances for full desmosome width represented by DPC-DPC or half desmosome width represented by Dsg2-DPC or Dsg2-DPN. To avoid false signals resulting from background noise or lines not fully drawn over the DP "railroad" track, we filtered out the data that showed less or more than two peaks in the DPC signals along the drawn segment or did not have the Dsg2 signal between the two DPC peaks. Furthermore, in certain desmosomes, the Dsg2 signals were observed as two peaks, as shown in Fig. S16A-B, implying that the desmosome had ruptured at the Dsg2 extracellular region. We eliminated these desmosomes from further analysis and only focused on intact desmosomes with unresolvable Dsg2 signals. For the DP line density measurement, we also used a custom-made ImageJ macro script for the quantitative analysis. Briefly, we first subtracted the background noise while retaining the DP signals along the cell borders. Then, we skeletonized these signals into segments to obtain the total amount of DP length along cell borders. Lastly, we obtained the DP line density by normalizing the total DP length using the number of cells in the image (*i.e.* $\frac{desmosome\ length}{cell}$).

Co-immunoprecipitation: Co-immunoprecipitation was performed as described previously [49]. Cells grown on tissue culture plates were washed three times with 1X cold PBS. Cells were lysed

at 4°C using ice-cold 1% triton lysis buffer (1% Triton X-100, 40mM HEPES (pH 7.5), 120mM

NaCl, 1mM EDTA, 1mM phenylmethylsulfonyl fluoride, 10mM sodium pyrophosphate, 1 µg/ml

each of chymostatin, leupeptin and pepstatin, 10 µg/ml each of aprotinin and benzamidine, 2

µg/ml antipain, 1mM sodium orthovanadate, 50mM NaF). Cell lysates were centrifuged to remove

cellular debris and protein concentration was determined using Pierce BCA Protein Assay Kit and

BSA as standards. Cell lysates were pre-cleared using Protein G Plus-Agarose beads (Santa

Cruz Biotechnology) for 40 minutes at 4°C while rotating. While rotating, supernatants were

incubated with the indicated antibody or IgG control for two hours at 4°C. Immuno-complexes

were captured using Protein G Plus-Agarose beads and washed 4X using 1% triton lysis buffer

supplemented with 300 mM NaCl.

Gel electrophoresis and immunoblotting: Cells grown on tissue culture plates were washed

three times with 1X cold PBS. Cells were lysed at 4°C using ice-cold 1% triton lysis buffer (1%

Triton X-100, 40mM HEPES (pH 7.5), 120mM NaCl, 1mM EDTA, 1mM phenylmethylsulfonyl

fluoride, 10mM sodium pyrophosphate, 1 µg/ml each of chymostatin, leupeptin and pepstatin, 10

µg/ml each of aprotinin and benzamidine, 2 µg/ml antipain, 1mM sodium orthovanadate, 50mM

NaF). For immunoblotting, cell lysates were centrifuged at 14,000rpm for 5 minutes at 4°C to

remove cellular debris. Protein concentration was determined using the Pierce BCA Protein Assay

Kit following the manufacturer's instructions (WG327385. Thermo Scientific). BSA (Equitech-Bio)

prepared in 1% lysis buffer was used as the standard. Aliquots of protein lysates were resolved

using SDS-PAGE and then transferred to nitrocellulose membranes (0.45um) (Amersham

Protran; Cytiva) and immunoblotted using the indicated antibodies, followed by horseradish

peroxidase-conjugated goat anti-mouse or goat anti-rabbit IgG (Sigma-Aldrich). Western

Lightning Plus-ECL (Waltham, MA) or Amersham ECL Select Western was used as a blotting

detection reagent. Signals were detected using iBright™ CL1500 Imaging System, Invitrogen

(Thermo Fisher Scientific). The immunoblot result for comparing K19 levels among the three cell

lines was conducted using a different method, as described previously [50].

Structure preparation and MD simulation: Plakin domain structure (residues 178-950) was

predicted using Alphfold2 [23]. Subsequently, the structure was prepared using PDBFixer [51]

and missing hydrogens were added under pH 7.0 conditions. The protein N-terminus was

designated with "NH3+" while "COO-" was assigned to the C-terminus. MD simulations were

conducted on the FARM high-performance computing cluster at the University of California,

Davis, with GROMACS 2022.3, as previously described [52-54]. The simulations were performed

in the OPLS-AA/L force field [55] and TIP4P water model with a 10 Å radius cut-off for Van der

Waals and electrostatic interactions. Electrostatic energy calculations employed the particle mesh

Ewald method with a 0.16-grid spacing. At the beginning of the simulation, the plakin domain

structure was positioned at the center of a dodecahedral box, ensuring a minimum distance of 1

nm from the boundary for every atom. The box was filled with water molecules and neutralized

with charged ions (150 mM NaCl, 4 mM KCl, and 2 mM $CaCl_2$). The system was relaxed with

energy minimization and stabilized with equilibration under isothermal-isochoric and isothermal

isobaric conditions using a modified Berendsen thermostat and Berendsen barostat. Post-

stabilization, a 50 ns MD simulation was performed with 2-fs integration steps, maintaining a

temperature of 300K using a v-rescale thermostat. Equilibration of the protein structure occurred

within 20 ns. The number of hydrogen bonds observed between the long arm and the short arm

was calculated using *gmx hbond.*

Constant-force SMD simulations and analysis: The constant-force SMD simulations were performed as described previously [52-54]. The starting structures for the SMD simulations were the last frame of the corresponding MD simulations. The structures were placed at the side of a rectangular box (23 × 70 × 8 nm box, center at [11.5nm, 65nm, 4nm]). The system, containing ~1,600,000 atoms, was relaxed and equilibrated under isothermal-isobaric conditions using the same protocol as in the MD simulation. The SMD simulations were performed at 310K temperature using a Nose-Hoover thermostat [56]. During each SMD simulation, we fixed the C-terminus of the plakin domain and pulled the N-terminal region (residues 919-950) along the longest axis of the box with a constant force ~830 pN (500 kJ·mol−1·nm−1). The distances between the protein N-terminal and C-terminal were calculated using the *gmx pairdist*.

Keratin filament orientation analysis: Confocal images were acquired as previously described for K8/DP and K18/DP staining. After image acquisition, we utilized a custom ImageJ macro script to determine the global dominant orientation of keratin filaments. Since filament orientation was defined as the angle between the keratin filament and the desmosome axis along the cell-cell junction, we first generated a mask based on DP signals outlining the cell border. This allowed us to define a region of interest (ROI) where the desmosome axis was set to 0°.

For each image, two rectangular ROIs were selected, each extending 2 μm from the cell border into the cytoplasm and having a length matching the desmosome mask. These ROIs were extracted, and the dominant orientation of keratin filaments was determined using the OrientationJ plugin. Specifically, OrientationJ computes the structure tensor, which captures intensity gradients at each pixel and extracts the principal direction of keratin filament orientation, represented as vector fields. The global dominant orientation was then derived by aggregating local orientations across the entire ROI. Consequently, each image provided two data points,

each representing the global dominant orientation of desmoplakin-associated keratin filaments at

the junction on either side of the cell-cell contact.

Fluorescence lifetime imaging microscopy (FLIM): Fluorescence lifetime data were acquired

using a Leica TCS SP8 STED 3X confocal microscope, equipped with a pulsed white light laser

(NKT Photonics) with wavelengths ranging from 470 to 670 nm, a 63x/1.35 NA oil immersion

objective (HC PL APO CS2), and HyD detectors with photon counting capabilities. Images were

captured at a scanning speed of 400 Hz with a resolution of 512 × 512 pixels. For each

experimental condition, 10 images were acquired from 3 biological replicates.

Cells were cultured overnight on a 35 mm dish with a #1.5 glass-like polymer coverslip (Cellvis)

to reach approximately 70% confluency. They were then transiently transfected with the DPI

tension sensor, no-force control, and donor-only control constructs (Addgene), previously

generated in an earlier study [28], using ViaFect Transfection Reagent (Promega) in reduced-

serum Opti-MEM medium (ThermoFisher). After 8-hour incubation at 37°C with 5% CO_2, the

culture medium was replaced with low-glucose DMEM (Gibco) supplemented with 10% FBS (Life

Technologies) and 1% penicillin-streptomycin (Sigma-Aldrich). Cells were then treated with 1

µg/ml doxycycline to induce gene expression of the DPI constructs through a Tet-On system,

which enables conditional gene expression in the presence of doxycycline by activating a

doxycycline-sensitive transcriptional activator. The following day, cells were fixed with 4% PFA

for 10 minutes at room temperature. After fixation, samples were maintained in Dulbecco's PBS

containing $MgCl_2$ and $CaCl_2$ (Sigma-Aldrich) and imaged at RT.

FLIM-FRET analysis: Fluorescence lifetime data were analyzed using the built-in FLIM Wizard

in LAS X software. For each lifetime image, 20–40 ROIs were manually selected, focusing on

individual DP puncta. In each ROI, a bi-exponential decay model was fitted to the photon count

time data to determine the fluorescence lifetime within each ROI. This model was chosen to

mitigate the effects of the instrument response function (IRF) and autofluorescence, which

typically result in small lifetime values. FRET efficiency (E) was then calculated using the donor

lifetime in the presence of an acceptor (τ_{DA}) and the mean donor-only lifetime (τ_D), following the

equation:

590
$$E = \left(1 - \frac{\tau_{DA}}{\tau_D}\right) \times 100$$

The mean donor-only lifetime was measured using donor-only constructs for each cell line, with

WT cells exhibiting a mean lifetime of 2.771 ns and K19-KO cells showing a mean lifetime of

2.784 ns. To ensure reliable lifetime fitting, the minimum required photon count was set to 220

photons per image, as determined by the reduction in the spread of the fitted lifetime values.

Statistical analysis: All statistical tests were performed using GraphPad Prism (GraphPad

Software), and all graphs were plotted using custom-made Python code except those mentioned

specifically. In the boxplots, the box represents the 25th and 75th percentiles with the median

indicated, and whiskers reach 1.5 times the interquartile range (IQR), defined as the difference

between the 25th and 75th percentiles. Data points outside the whiskers are shown as outliers.

The number of independent experiments (N) and data points (n) is indicated in each figure legend.

Data distribution was first tested based on Shapiro-Wilk's test. Since most data was not normally

distributed, non-parametric statistic tests were used for the analysis. Data were analyzed using

Mann-Whitney's U test. A multiple-group comparison was performed using the Kruskal-Wallis

Test, followed by Dunn's multiple comparison Test. Statistical tests used are specified in the figure

legends. Statistical significance was represented at the level of *, $P < 0.05$; **, $P < 0.01$; ***,

$P<0.001$; ns, $P>0.05$.

For the mean FRET efficiency analysis, statistical tests were conducted in R using a linear mixed-

effects model, incorporating a grouped effect to account for puncta derived from the same image

as described previously [28]. This approach corrects for statistical dependence among puncta

within a single image. Specifically, the model assumes that grouped error across images follows

a normal distribution centered around zero. By including this grouped effect, the analysis provides

more conservative and robust statistical confidence compared to methods that treat each puncta

as an independent data point. In R, the applied linear mixed-effects model was structured as

fretEfficiency ~ isTensionSensor + (1|imageNumber), where *isTensionSensor* was assigned a

value of 1 for tension sensor data and 0 for the corresponding truncated control data, while

imageNumber uniquely identified each image.

Acknowledgments: Research in the SS lab was supported by the National Institute of General

Medical Sciences of the National Institutes of Health (R01GM121885). Research in the JSC lab

was supported by the National Institute of General Medical Sciences of the National Institutes of

Health (R15CA267890-01). STED imaging and FRET-FLIM imaging was performed at the UC

Davis Advanced Imaging Facility. FACS was performed at the Flow Cytometry Shared Resource,

funded by the UC Davis Comprehensive Cancer Center Support Grant (CCSG) awarded by the

National Cancer Institute (NCI P30CA093373).

Data availability: All data are made available in the manuscript and supporting information.

Conflict of Interest Statement: The authors do not declare any competing interests.

Figure Legends

Fig. 1: Keratin-19 influences desmosome architecture. (A) Schematic of desmosome structure. **(B)** Representative confocal image of desmosome puncta along cell-cell borders in the WT cells. The desmosomes were immunolabeled for DPC (green) and Dsg2 (red). Scale bar: 50 µm. **(C)** Representative STED image of desmosomes showing a characteristic 'railroad track' pattern. Scale bar: 10 µm. **(D-F)** Representative close-up STED images of desmosomes exhibiting the signature 'railroad track' pattern in **(D)** MCF7 WT cells, **(E)** MCF7 K19-KO cells, and **(F)** MCF7 K19-GFP cells. Scale bar: 500 nm. **(G)** The boxed area from the representative STED images of WT cells shows a closer look at an individual DP railroad track with Dsg2 between the two parallel DP plaques. Scale bar: 200 nm. **(H)** Line-scan analysis of DPC and Dsg2 fluorescence intensity (indicated by the dashed line in E). **(I)** Quantification of DPC-DPC distance from WT, K19-GFP, and K19-KO cells shows that desmosomes in WT and K19-GFP cells are wider than in K19-KO cells. In all boxplots, the box represents the 25th and 75th percentiles with the median indicated, and whiskers reach 1.5 times the interquartile range (IQR), defined as the difference between the 25th and 75th percentiles. Data points outside the whiskers are shown as outliers. Number of data points (n) = 630 (WT), 505 (K19-KO), 716 (K19-GFP) (n represents the number of line scans across the desmosomes); Number of replicates (N) = 3. Kruskal-Wallis Test, followed by Dunn's multiple comparison Test; ***, $P<0.001$.

Fig. 2: Interplay between keratin-19 and actomyosin network determines desmosome width. (A) Schematic of the interactions between the desmosome, keratin intermediate filaments, and actin filaments. Keratin filaments bind to the C-terminal of DP and interacts with actin filaments in the cytoplasm. **(B)** Representative STED images of K19-GFP rescue cells immunolabeled for K19 (magenta), DP (green), and actin (cyan). The individual components are shown in grayscale to highlight the structural features. The merged images between K19 and DP or actin are shown at the right. Scale bar: 500 nm. The images show that K19 filaments colocalize with the DP "railroad" tracks and with cortical actin belts. **(C)** Co-IP in WT cells performed with anti-K19 antibody or IgG control. The Co-Ips show that K19 interacts with both DP and actin. **(D)** Representative STED images of DPC (green) at the cell border of WT cells and Blebbstatin-treated WT cells (WT+Blebb). **(E)** Quantification of desmosome widths from WT, K19-KO, and WT+Blebb cells showing that Blebbistatin treatment reduces the desmosome width to a similar level as in the K19-KO cell line. n = 630 (WT), 505 (K19-KO), 650 (WT+Blebb); N = 3. Kruskal-Wallis Test, followed by Dunn's multiple comparison Test; ***, P<0.001. **(F)** Representative STED images of DPC (green) at the cell border of K19-KO cells and Calyculin A-treated K19-KO cells (KO+Caly). **(G)** Quantification of desmosome widths from WT, K19-KO, and KO+Caly cells showing that Calyculin A increases the desmosome width of K19-KO cells to a similar level as in the WT cell line. n = 630 (WT), 505 (K19-KO), 347 (KO+Caly); N = 3. Kruskal-Wallis Test, followed by Dunn's multiple comparison Test; ***, P<0.001.

Fig. 3: Desmoplakin undergoes force-induced conformational change in the plakin domain. **(A)** Schematic of the desmosome under tension. The N-terminal of DP transitions from a closed conformation to an open conformation. **(B)** Representative STED images of DPC (green) and DSG2 (red) at the cell border for WT and K19-KO cells. Scale bar: 500 nm. **(C)** Representative STED images of DPN (green) and DSG2 (red) at the cell border for WT and K19-KO cells. Scale

bar: 500 nm. **(D)** Quantification of desmosome half-unit widths (Dsg2-DPC distance) from WT

and K19-KO cells. n = 1260 (WT), 1010 (K19-KO); N = 3. Mann-Whitney's U test; ***, P<0.001.

The distances between Dsg2 and DPC are significantly greater in the WT compared to the K19-

KO cells. **(E)** Quantification of desmosome half-unit widths (Dsg2-DPN distance) from WT and

K19-KO cells. n = 692 (WT), 826 (K19-KO); N = 3. Mann-Whitney's U test; ns, P>0.05. The Dsg2-

DPN distance in both cell lines are similar. **(F)** The mean values from the results in D and E are

summarized in the bar chart to compare the DP length (DPN-DPC distance) between the WT and

K19-KO cells. The DP length is 66 nm for the WT and 33 nm for the K19-KO, indicating that DP

extends 33 nm in the WT cells. **(G)** The entire plakin domain structure predicted by Alphafold2

suggests a U-shape conformation in the absence of force. Two common electrostatic interactions

(R592-N744 and K444-E858) observed in all 3 MD simulations (below) are shown. **(H)** During the

MD simulations, 10 to 15 hydrogen bonds were formed between the plakin domain's long arm

(SR3-4 and SR5-6) and short arm (SR7-8 and SR8-CT). **(I)** During the SMD simulations, the

distance between the plakin domain's C-terminus and N-terminus increased by 30~33 nm. This

elongation correlates with the distance change experimentally measured in Fig 3F. **(J)**

Comparison of the plakin domain structure at the start (left) and at the end (right) of the SMD

simulation. The structures show that the plakin domain was elongated by the pulling force without

significant secondary structure unfolding.

Fig. 4: Keratin filaments reorient in response to changes in actomyosin forces: (A)

Schematic representation of DP-associated keratin filaments at the cell-cell junction. Under no-

tension conditions, DP-associated filaments are primarily aligned horizontally. Under tension, they

transition to a radially organized structure that exerts force on the desmosome. **(B)** Co-IP of K8

and K18 in WT and K19-KO cells. Co-IP was performed using anti-K8, anti-K18 antibodies, or an

IgG control. The Co-IPs reveal that both K8 and K18 interact with DP and actin. **(C-D)**

Representative confocal images of WT **(C)** and K19-KO **(D)** cells immunolabeled for K8 (green)

and DP (red). Scale bar: 5 µm (left panel). A magnified grayscale image highlights DP-associated

K8 filaments at the junction with enhanced contrast (middle panel). The local dominant orientation

of K8 filaments represented as a vector field image (right panel). **(E)** Polar histogram of the global

dominant orientation of K8 filaments in WT and K19-KO cells. n = 108 (WT), 96 (K19-KO); N = 3.

Mann-Whitney U test, P < 0.0001. The K8 filaments in WT cells exhibit a more radial organization,

while K19-KO filaments are horizontally aligned. **(F)** Polar histogram of K8 filament orientation in

WT and K19-KO cells treated with Blebbistatin. n = 90 (WT), 90 (K19-KO); N = 3. Mann-Whitney

U test, P > 0.05. Following Blebbistatin treatment, WT filaments shift from a radial to a horizontal

organization, making their global dominant orientation indistinguishable from K19-KO cells. **(G)**

Polar histogram of K8 filament orientation in WT and K19-KO cells treated with Calyculin A. n =

98 (WT), 98 (K19-KO); N = 3. Mann-Whitney U test, P > 0.05. After Calyculin A treatment, K19-

KO K8 filaments shift from a horizontal to a radial organization, making their global dominant

orientation indistinguishable from WT cells. **(H-I)** Representative confocal images of **(H)** WT and

(I) K19-KO cells immunolabeled for K18 (green) and DP (red). Scale bar: 5 µm (left panel).

Magnified images highlighting K18 filaments at the junction and local dominant orientation of K18

filaments are shown in the middle and right panel. **(J)** Polar histogram of the global dominant

orientation of K18 filaments in WT and K19-KO cells. n = 112 (WT), 120 (K19-KO); N = 3. Mann-

Whitney U test, P < 0.0001. The global dominant orientation of K18 filaments in WT cells is

significantly different from that in K19-KO cells, with WT filaments adopting a radial organization

while K19-KO filaments remain horizontal, mirroring the pattern observed for K8 filaments.

Fig. 5: Desmoplakin is mechanically loaded in WT cells but not in K19-KO cells. (A)
Schematic illustration of the DP FRET tension sensor (DP-TS) and no-tension control. The

tension-sensing module consists of a linker peptide flanked by mTFP1 (donor fluorophore) and

mEYFP (acceptor fluorophore). In the absence of tension, the FRET pair stays together in both constructs. Under tension, the FRET pair is pulled apart in DP-TS, but remains together in the no-tension control which lacks the keratin-binding C-terminal region. **(B)** Representative STED images of WT and K19-KO cells transfected with DP-TS. The overlay of fluorescent signals with the bright-field image shows the proper localization of DP-TS at the cell-cell contacts in both WT and K19-KO cells. Scale bar: 10 µm. The zoomed-in images reveal the formation of distinct DP puncta along the cell border. **(C)** Illustration of the workflow for FRET-FLIM analysis. First, lifetime data images were acquired from transfected cells in fixed samples. Scale bar: 10 µm. Next, individual DP puncta were manually selected as regions of interest (ROIs) along the cell border. The fluorescence lifetime for each DP puncta was determined by fitting a bi-exponential model to distinguish true lifetime measurements from the effects of the instrument response function (IRF). Finally, FRET efficiency (E) was calculated from the lifetime measurements as described in the methods. **(D)** Quantification of FRET efficiency for DP-TS and no-tension control constructs in WT cells. The median FRET efficiency of the tension sensor was significantly lower than the control, indicating that DP is mechanically loaded in WT cells. $n = 30$ images, $N = 3$, $P < 0.0001$. **(E)** Quantification of FRET efficiency for DP-TS and no-tension control constructs in K19-KO cells. The median FRET efficiency of DP-TS was indistinguishable from that of the control, suggesting that DP experiences undetectable tension in K19-KO cells. $n = 30$ images, $N = 3$, $P > 0.05$. **(F)** Model for force-induced DP conformational change in desmosomes. When desmosomes experience no tension, DP-associated keratin intermediate filaments align horizontally with the cell membrane, maintaining a closed DP conformation. Under mechanical stress, the keratin filaments rearrange into a more radial orientation, effectively transmitting forces to DP. This results in a conformational change in the flexible DP plakin domain, converting this domain from a folded (closed) to an extended (open) conformation.

References

751   1.   Green, K.J. and C.L. Simpson, *Desmosomes: New Perspectives on a Classic.* Journal of

752   Investigative Dermatology, 2007. **127**(11): p. 2499-2515.

753   2.   Broussard, J.A., S. Getsios, and K.J. Green, *Desmosome regulation and signaling in*

754   *disease.* Cell and Tissue Research, 2015. **360**(3): p. 501-512.

755   3.   Garrod, D. and M. Chidgey, *Desmosome structure, composition and function.* Biochim

756   Biophys Acta, 2008. **1778**(3): p. 572-87.

757   4.   Stahley, S.N. and A.P. Kowalczyk, *Desmosomes in acquired disease.* Cell and Tissue

758   Research, 2015. **360**(3): p. 439-456.

759   5.   Holthofer, B., et al., *Structure and function of desmosomes.* Int Rev Cytol, 2007. **264**: p.

760   65-163.

761   6.   Hatzfeld, M., R. Keil, and T.M. Magin, *Desmosomes and Intermediate Filaments: Their*

762   *Consequences for Tissue Mechanics.* Cold Spring Harbor Perspectives in Biology, 2017.

763   **9**(6): p. a029157.

764   7.   Broussard, J.A., et al., *Scaling up single-cell mechanics to multicellular tissues – the role*

765   *of the intermediate filament–desmosome network.* Journal of Cell Science, 2020. **133**(6):

766   p. jcs228031.

767   8.   Prechova, M., et al., *Plectin-mediated cytoskeletal crosstalk controls cell tension and*

768   *cohesion in epithelial sheets.* Journal of Cell Biology, 2022. **221**(3).

769   9.   Broussard, J.A., et al., *The desmoplakin-intermediate filament linkage regulates cell*

770   *mechanics.* Mol Biol Cell, 2017. **28**(23): p. 3156-3164.

10.   Huen, A.C., et al., *Intermediate filament–membrane attachments function synergistically*

with actin-dependent contacts to regulate intercellular adhesive strength. The Journal of

Cell Biology, 2002. **159**(6): p. 1005-1017.

11.   Bornslaeger, E.A., et al., *Breaking the connection: displacement of the desmosomal*

plaque protein desmoplakin from cell-cell interfaces disrupts anchorage of intermediate

filament bundles and alters intercellular junction assembly. The Journal of cell biology,

1996. **134**(4): p. 985-1001.

12.   Kröger, C., et al., *Keratins control intercellular adhesion involving PKC-α–mediated*

desmoplakin phosphorylation. Journal of Cell Biology, 2013. **201**(5): p. 681-692.

13.   Stappenbeck, T.S., et al., *Phosphorylation of the desmoplakin COOH terminus*

negatively regulates its interaction with keratin intermediate filament networks. Journal of

Biological Chemistry, 1994. **269**(47): p. 29351-29354.

14.   Jefferson, J.J., et al., *Structural analysis of the plakin domain of bullous pemphigoid*

antigen1 (BPAG1) suggests that plakins are members of the spectrin superfamily.

Journal of molecular biology, 2007. **366**(1): p. 244-257.

15.   Smith, E.A. and E. Fuchs, *Defining the interactions between intermediate filaments and*

desmosomes. The Journal of cell biology, 1998. **141**(5): p. 1229-1241.

16.   Al-Jassar, C., et al., *The nonlinear structure of the desmoplakin plakin domain and the*

effects of cardiomyopathy-linked mutations. J Mol Biol, 2011. **411**(5): p. 1049-61.

17.   Al-Jassar, C., et al., *Hinged Plakin Domains Provide Specialized Degrees of Articulation*

in Envoplakin, Periplakin and Desmoplakin. PLoS ONE, 2013. **8**(7): p. e69767.

18.   Green, K.J., et al., *Structure of the human desmoplakins. Implications for function in the*

desmosomal plaque. Journal of Biological Chemistry, 1990. **265**(5): p. 2603-2612.

19.   Stahley, S.N., et al., *Molecular organization of the desmosome as revealed by direct*

stochastic optical reconstruction microscopy. Journal of Cell Science, 2016. **129**(15): p.

2897-2904.

20.   Alsharif, S., et al., *Keratin 19 maintains E-cadherin localization at the cell surface and*

stabilizes cell-cell adhesion of MCF7 cells. Cell Adhesion & Migration, 2021. **15**(1): p. 1-

17.

21.   Leube, R.E., M. Moch, and R. Windoffer, *Intermediate filaments and the regulation of*

focal adhesion. Curr Opin Cell Biol, 2015. **32**: p. 13-20.

22.   Fujiwara, S., et al., *Interplay between Solo and keratin filaments is crucial for mechanical*

force-induced stress fiber reinforcement. Mol Biol Cell, 2016. **27**(6): p. 954-66.

23.   Jumper, J., et al., *Highly accurate protein structure prediction with AlphaFold.* Nature,

2021. **596**(7873): p. 583-589.

24.   Choi, H.-J. and W.I. Weis, *Crystal Structure of a Rigid Four-Spectrin-Repeat Fragment*

of the Human Desmoplakin Plakin Domain. Journal of Molecular Biology, 2011. **409**(5):

p. 800-812.

25.   Weber, G.F., M.A. Bjerke, and D.W. DeSimone, *A mechanoresponsive cadherin-keratin*

complex directs polarized protrusive behavior and collective cell migration.

Developmental Cell, 2012. **22**(1): p. 104-115.

26.   Loschke, F., et al., *Regulation of keratin network organization.* Current opinion in cell
813   biology, 2015. **32**: p. 56-64.

814   27.   Sharma, P., et al., *Keratin 19 regulates cell cycle pathway and sensitivity of breast*
815   *cancer cells to CDK inhibitors.* Scientific reports, 2019. **9**(1): p. 14650.

816   28.   Price, A.J., et al., *Mechanical loading of desmosomes depends on the magnitude and*
817   *orientation of external stress.* Nature Communications, 2018. **9**(1).

818   29.   Stappenbeck, T.S., et al., *Functional analysis of desmoplakin domains: specification of*
819   *the interaction with keratin versus vimentin intermediate filament networks.* The Journal
820   of cell biology, 1993. **123**(3): p. 691-705.

821   30.   Grashoff, C., et al., *Measuring mechanical tension across vinculin reveals regulation of*
822   *focal adhesion dynamics.* Nature, 2010. **466**(7303): p. 263.

823   31.   Sadhanasatish, T., et al., *A molecular optomechanics approach reveals functional*
824   *relevance of force transduction across talin and desmoplakin.* Science Advances, 2023.
825   **9**(25): p. eadg3347.

826   32.   Yao, M., et al., *Force-dependent conformational switch of α-catenin controls vinculin*
827   *binding.* Nature Communications, 2014. **5**(1).

828   33.   Kim, T.-J., et al., *Dynamic Visualization of α-Catenin Reveals Rapid, Reversible*
829   *Conformation Switching between Tension States.* Current Biology, 2015. **25**(2): p. 218-
830   224.

831   34.   Yonemura, S., et al., *α-Catenin as a tension transducer that induces adherens junction*
832   *development.* Nature Cell Biology, 2010. **12**(6): p. 533-542.

833   35.   Baddam, S., et al., *The desmosomal cadherin desmoglein-2 experiences mechanical*

834   *tension as demonstrated by a FRET-based tension biosensor expressed in living cells.*

835   Cells, 2018. **7**(7): p. 66.

836   36.   O'Keefe, E., H. Erickson, and V. Bennett, *Desmoplakin I and desmoplakin II: purification*

837   *and characterization.* Journal of Biological Chemistry, 1989. **264**(14): p. 8310-8318.

838   37.   North, A.J., et al., *Molecular map of the desmosomal plaque.* Journal of cell science,

839   1999. **112**(23): p. 4325-4336.

840   38.   Beggs, R.R., et al., *Desmosomes undergo dynamic architectural changes during*

841   *assembly and maturation.* Tissue Barriers, 2022. **10**(4).

842   39.   Mendonsa, A.M., T.-Y. Na, and B.M. Gumbiner, *E-cadherin in contact inhibition and*

843   *cancer.* Oncogene, 2018. **37**(35): p. 4769-4780.

844   40.   Hobbs, R.P. and K.J. Green, *Desmoplakin Regulates Desmosome Hyperadhesion.*

845   Journal of Investigative Dermatology, 2012. **132**(2): p. 482-485.

846   41.   Seltmann, K., et al., *Keratins significantly contribute to cell stiffness and impact invasive*

847   *behavior.* Proceedings of the National Academy of Sciences, 2013. **110**(46): p. 18507-

848   18512.

849   42.   Cheung, J., Kevin, et al., *Collective Invasion in Breast Cancer Requires a Conserved*

850   *Basal Epithelial Program.* Cell, 2013. **155**(7): p. 1639-1651.

851   43.   Daday, C., K. Kolšek, and F. Gräter, *The mechano-sensing role of the unique SH3*

852   *insertion in plakin domains revealed by Molecular Dynamics simulations.* Scientific

853   Reports, 2017. **7**(1).

44.   Rief, M., J.P. , and M.S. , Hermann E. Gaub, *Single Molecule Force Spectroscopy of*

Spectrin Repeats: Low Unfolding Forces in Helix Bundles. Journal of Molecular Biology,

1999. **286**(2): p. 553-561.

45.   Kurochkina, N. and U. Guha, *SH3 domains: modules of protein–protein interactions.*

Biophysical Reviews, 2013. **5**(1): p. 29-39.

46.   Ortega, E., et al., *The Structure of the Plakin Domain of Plectin Reveals an Extended*

Rod-like Shape. Journal of Biological Chemistry, 2016. **291**(36): p. 18643-18662.

47.   Sharma, P., et al., *Keratin 19 regulates cell cycle pathway and sensitivity of breast*

cancer cells to CDK inhibitors. Scientific Reports, 2019. **9**(1).

48.   Sharma, P., et al., *Keratin 19 interacts with GSK3beta to regulate its nuclear*

accumulation and degradation of cyclin D3. Mol Biol Cell, 2021. **32**(21): p. ar21.

49.   Fallatah, A., et al., *Keratin 19 binds and regulates cytoplasmic HNRNPK mRNA targets*

in triple-negative breast cancer. BMC Molecular and Cell Biology, 2023. **24**(1).

50.   Shafraz, O., C.M.O. Davis, and S. Sivasankar, *Light-activated BioID – an optically*

activated proximity labeling system to study protein–protein interactions. Journal of Cell

Science, 2023. **136**(19).

51.   Eastman, P., et al., *OpenMM 7: Rapid development of high performance algorithms for*

molecular dynamics. PLoS Comput Biol, 2017. **13**(7): p. e1005659.

52.   Xie, B., et al., *Strengthening E-cadherin adhesion via antibody-mediated binding*

stabilization. Structure, 2024. **32**(2): p. 217-227 e3.

53.   Xie, B., et al., *Molecular mechanism for strengthening E-cadherin adhesion using a*

875   *monoclonal antibody.* Proceedings of the National Academy of Sciences, 2022. **119**(32):

876   p. e2204473119.

877   54.   Xie, B., S. Xu, and S. Sivasankar, *Outside-in engineering of cadherin endocytosis using*

878   *a conformation strengthening antibody.* Nature Communications, 2025. **16**(1): p. 1157.

879   55.   Kaminski, G.A., et al., *Evaluation and reparametrization of the OPLS-AA force field for*

880   *proteins via comparison with accurate quantum chemical calculations on peptides.*

881   Journal of Physical Chemistry B, 2001. **105**(28): p. 6474-6487.

882   56.   Lemkul, J.A. and D.R. Bevan, *Assessing the stability of Alzheimer's amyloid protofibrils*

883   *using molecular dynamics.* J Phys Chem B, 2010. **114**(4): p. 1652-60.

A. Tension — No tension

B. WT Cells — K19-KO Cells

C. Image transfected cells → Select puncta → Fit lifetime → Determine FRET

D. WT Cells

E. K19-KO Cells

F. No tension — Force → Tension