Title: Desmoplakin is a desmosomal mechanosensor

Authors: Yinchen Dong [1], Ahmed Elgerbi [3], Bin Xie [2], John S. Choy [3], Sanjeevi Sivasankar [1, 2, *]

Affiliations: [1] Department of Biomedical Engineering, University of California, Davis, CA; [2]

Biophysics Graduate Group, University of California, Davis, CA; [3] Department of Biology, The

Catholic University of America, Washington DC.

*Corresponding author: ssivasankar@ucdavis.edu

Abstract: Desmosomes are essential cell-cell adhesion organelles that enable tension-prone tissue, like the skin and heart, withstand mechanical stress. Desmosomal anomalies are associated with numerous epidermal disorders and cardiomyopathies. Despite their critical role in maintaining tissue resilience, an understanding of how desmosomes respond to mechanical stimuli is lacking. Here we demonstrate, in human breast cancer MCF7 cells, that actomyosin forces induce a conformational change in the N-terminal region of desmoplakin, a critical cytoplasmic desmosomal protein. Using super-resolution fluorescence microscopy, biochemical assays, and atomistic computer simulations, we show that actomyosin forces are directed to desmoplakin along keratin-19 filaments. These forces induce a conformational change in the N-terminal plakin domain of desmoplakin, converting this domain from a folded (closed) to an extended (open) conformation. Functional adhesion assays show that MCF7 cells with desmoplakin in an open conformation are more adhesive than cells with desmoplakin in a closed state. Our findings establish that desmoplakin is mechanosensitive and undergoes force-induced conformational changes which enhance the mechanical resilience of desmosomes.

Significance Statement: Desmosomes are essential cell-cell adhesion junctions that enable

tension-prone tissue, like the skin and heart, withstand mechanical stress. Despite their critical

role in maintaining tissue integrity, it is unclear how desmosomes respond to mechanical forces.

Here we demonstrate that force induces a conformational change in desmoplakin, a critical

intracellular desmosomal protein. This force-induced change in the desmoplakin's structure

enhances the resilience of desmosomes and enables them to withstand mechanical stress.

Introduction

Desmosomes are critical cell-cell junctions that promote adhesion between cells and preserve tissue integrity. They are crucial in enabling tension-prone tissues like the skin and heart to withstand mechanical stresses [1]. Desmosomal anomalies are associated with numerous diseases in these tissues, such as epidermal autoimmune disorders and arrhythmogenic ventricular cardiomyopathy [2-4]. Despite their critical role in maintaining tissue mechanical resilience, an understanding of how desmosomes respond to mechanical stimuli is lacking.

Desmosomes are composed of transmembrane desmosomal cadherins: desmoglein (Dsg) and desmocollin (Dsc), and cytoplasmic plaque proteins: plakophilin, plakoglobin, and desmoplakin (DP). The extracellular regions of the cadherins from opposing cells interact to mediate adhesion while the plaque proteins bind to the cadherin cytoplasmic tail (Fig. 1A) [5]. DP anchors desmosomes to the keratin intermediate filament (KIF) cytoskeleton, and this linkage is essential for maintaining tissue integrity and facilitating intracellular signal transduction [6, 7]. Several studies show that KIFs work in cooperation with the actomyosin cytoskeleton to modulate the mechanical state of desmosomes [8-10]. Disassociation of KIFs from circumferential actin belts alters tension across desmosomes [8] while disassembly of the actin cytoskeleton decreases desmosome resilience [9]. However, the molecular mechanisms by which actomyosin forces alter desmosome structure and function are not understood.

Disruption of the linkage between DP and KIF impairs intercellular adhesion and compromises tissue integrity [10, 11]. Knocking-out keratins leads to elevated phosphorylation of DP, resulting in accelerated endocytosis of desmosomes and rupture of epithelial sheets under rotational stress [12]. Weakening connections between DP and KIF networks dramatically alters the mechanical properties of cells such as cell stiffness and intercellular forces [9]. While these data suggest that

DP is crucial in regulating the response of desmosomes to mechanical stress, the biophysical

mechanism by which DP mechanically regulates the desmosome is unknown.

The structure of DP offers hints into its mechanoregulatory function. DP is a large, 2871 amino

acid protein which is composed of an N-terminal region, a central rod domain which facilitates DP

homodimerization [3], and a C-terminal region which is crucial for aligning and binding KIFs [13].

The N-terminal region of DP is subdivided into a globular head domain and a plakin domain, each

playing distinct roles [14]. The globular head domain binds to the C-termini of desmosomal

cadherins, and recruits plakophilin and plakoglobin [15]. The plakin domain, characterized by six

spectrin repeat (SR) domains interspersed with Src homology (SH) domains, exhibits structural

flexibility, as observed in a previous *in vitro*, cell free small angle X-ray scattering study [16]. That

investigation revealed that the plakin domain adopts either a folded "U" shape or a fully extended

"I" shape, indicating a significant degree of structural adaptability [17]. This suggests that

actomyosin forces could induce a conformational change in the N-terminal region of DP, thereby

offering a molecular pathway for DP mechanosensitivity. However, force induced conformational

changes in DP, particularly within the cell, have never been demonstrated.

Here, we present the first direct evidence for actomyosin force induced conformational changes

in the N-terminal region of DP, in human breast cancer MCF7 cells. These cells are enriched with

keratin 19 (K19), and we use super-resolution microscopy with wild type (WT), K19-knock out

(K19-KO), and K19-GFP rescue (K19-GFP) cells, pharmacological inhibition, co-

immunoprecipitation (co-IP), and atomistic Molecular Dynamics (MD) simulations to show that

actomyosin forces are directed to DP along K19 filaments. We show that these forces induce a

conformational change in the N-terminal plakin domain from a folded 'closed' to an extended

'open' conformation. Functional adhesion assays show that MCF7 cells with DP in an open

conformation are more adhesive than cells with DP in a closed state. Our findings establish, at

the molecular level, that DP is a mechanosensor within the desmosome We show that

desmoplakin undergoes a force-induced conformational change which enhances desmosomal

mechanical resilience.

Results

K19 remodels desmosome architecture: Desmosomes are characterized by distinct "railroad"

track structure, represented by discrete parallel fragments of DP which sandwich the desmosomal

cadherins [18]. To validate this signature structure of desmosomes in MCF7 cells, we conducted

immunostaining targeting the extracellular region of isoform 2 of desmoglein (Dsg2) and the C-

terminal region of DP (DPC) using super-resolution Stimulated Emission Depletion (STED)

microscopy. The results, in WT cells, showed well-defined DP "railroad" tracks, as depicted in Fig

1B. To investigate the effect of K19 on desmosome architecture, we used the K19-KO cell line

[19]. Successful ablation of K19 in these cells was confirmed using western blots (Fig. S1).

Surprisingly, in the K19-KO cells, STED imaging revealed qualitatively more compact

desmosomes (Fig. 1C).

Super-resolution microscopy allowed us to visualize the individual DP plaques and quantify the

desmosome width by measuring the distance between opposing plaques. To measure distance

between opposing plaques, we drew a line scan across the desmosome complex (Fig. 1E) and

measured the distance between the two peaks of DPC signal (designated as DPC-DPC distance),

as illustrated in the fluorescence intensity plot (Fig. 1F). This quantitative analysis of the

desmosome width confirmed our previous observations, with significantly narrower desmosomes

formed in the K19-KO cell line (334 ± 61 nm for WT cells and 264 ± 65 nm for K19-KO cells; Fig.

1G).

To corroborate that the observed shift in desmosome width was directly related to the absence of

K19, we stably rescued the K19-KO cells with EGFP-tagged K19 (K19-GFP cells). Western blots

validated the successful expression of K19-GFP in the rescued cells (Fig. S2). Desmosome width

in the K19-GFP cells recovered to a level similar to the WT cell line (322 ± 56; Fig. 1H). Together,

these findings show that desmosome architecture in MCF7 cells strongly correlates with the

presence/absence of K19. In the presence of K19, the desmosome width is larger, while the

desmosome narrows when K19 is knocked out.

Interplay between K19 and the actomyosin networks governs desmosome width: The

cytoplasmic KIF network has two distinct components: radial KIFs emanating from the nucleus

and anchoring on the desmosomes and the cortical KIFs interconnecting the desmosomes near

the plasma membrane [20] (illustrated in Fig. 2A). Interactions between the KIF network and

elements of the actomyosin network have been observed at cell-cell junctions and at the cell-

substrate interface [8, 21]. To investigate how changes in the KIF network contribute to

desmosome remodeling, we first inspected the intracellular localization of K19, DP, and F-actin

filaments in K19-GFP cells. K19 was found to form circumferential rims that colocalized with the

DP "railroad" tracks along the cell border (Fig. 2B). Additionally, the cortical actin belt was

observed to align with the K19 circumferential rim (Fig. 2B). Based on the colocalization of K19

with DP and the coalignment of K19 with F-actin filaments, we hypothesized that K19 interacts

with both DP and F-actin filaments.

To validate this hypothesis, we conducted Co-IP experiments using IgG as the control and K19

as the bait protein. The results revealed that K19 interacts with both DP and F-actin, consistent

with our previous immunofluorescence microscopy observations (Fig. 2C). In addition,

plakophilin-3, another desmosomal plaque protein, also interacted with K19 (Fig. S3), which

further supports the interaction between K19 filaments and the desmosome complex.

It has been demonstrated in Madin-Darby canine kidney (MDCK) cells that knocking down keratin-

18 (K18) significantly reduces the formation of force-induced actin stress fibers [22]. Since both

K18 and K19 bind to keratin-8 (K8) to form keratin filaments [23], we used confocal fluorescence

microscopy to image actin filaments in WT, K19-GFP and K19-KO cells (Fig. S4A). Quantitative

analysis of the data showed that compared to WT and K19-GFP cells, actin stress fibers in the

K19-KO cells were dramatically decreased (Fig. S4B). This suggests that mechanical tension in

the K19-KO cells is lower compared to WT and K19-GFP cells. We therefore hypothesized that

actomyosin-generated forces could be transmitted to the desmosome complex through

associations with the KIF network.

To test this hypothesis, we used Blebbistatin, a myosin-II-specific ATPase inhibitor, to reduce the

actomyosin contractility in the WT cell line and measured the desmosome widths in these

Blebbistatin-treated cells (WT+Bleb). A quantitative analysis indicated that Blebbistatin treatment

significantly reduced the desmosome width in WT cells to a level that was similar to K19-KO cells

(263 ± 62 for WT+Bleb; Fig. 2D). At the same time, the control group treated with Blebbistatin's

solvent, dimethyl sulfoxide (DMSO), still retained similar desmosome widths as the WT cells (330

± 65 for WT+DMSO; Fig. S5). This demonstrates that the K19 dependent change in desmosome

architecture are a consequence of a mechanical pathway that is established due to interactions

between K19, F-actin, and DP. Consequently, contractile actomyosin forces are transmitted to

and reorganize the desmosome, resulting in an increase in desmosome width.

DP elongates in the presence of mechanical force: Since previous studies have reported a

structural flexibility in the N-terminal plakin domain of DP [17], we hypothesized that the increase

in desmosome width in WT cells was due to a force-induced conformational change in this region

of DP (Fig. 3A). To test this, we specifically stained both the DP N-terminal region (DPN) on its

globular head domain (which is upstream of the plakin domain) and the extracellular region of

Dsg2. We conducted a quantitative comparison of the distances between Dsg2 and DPC, as well

as Dsg2 and DPN (as displayed in Fig. 3A), in in both WT and K19-KO cell lines. We selectively

measured these distances within the half-unit of the desmosome complex to gain a clearer

understanding of the structural changes occurring in a single DP under different conditions. The

distances between Dsg2 and DPC were found to be significantly greater in the WT compared to

the K19-KO cell line (167 ± 47 for WT cells and 132 ± 50 for K19-KO cells; Fig. 3B). In contrast,

no difference was detected in the distances between Dsg2 and DPN in both cell lines (101 ± 48

for WT cells and 99 ± 53 for K19-KO cells; Fig. 3C). As seen in Fig. 3D, the DP length (defined

as the distance between DPN and DPC) was 33 nm greater in the WT cells compared to the K19-

KO cells, suggesting that DP was elongated by higher mechanical tension in the WT cell lines.

The N-terminal plakin domain transitions from a closed to open conformation when pulled:

To resolve the mechanistic basis by which DP transitions between open and closed structures,

we proceeded to atomistically simulate the effect of force on the conformation of the highly

conserved plakin domain of DP (residues 178-950). We first generated the entire plakin domain

structure using Alphfold2 [24]. The predicted structure exhibited a 'U-shaped' folded conformation,

with spectrin repeat (SR) domains SR3-6 and Src homology (SH) domain SH3 forming a long arm

that interacts with a short arm composed of SR7-8 and SR8-CT domains (Fig. 4A). This prediction

agrees well with the crystal structure of the long arm of the plakin domain [25] (RMSD ~1.7Å, Fig.

S6).

To identify the interactions crucial for stabilizing the folded conformation, we performed 3 force-

independent molecular dynamics (MD) simulations on the predicted structure. The structure

stabilized within 20 ns of MD simulation (Fig. S7). During the simulations, 10-15 hydrogen bonds

were observed between the long arm and the short arm, and the interactions were mostly

observed between SR5-6 and SR7-8 regions (Fig. 4B and Fig. S8). Notably, two consistent salt

bridge/hydrogen bond interactions, between four conserved amino acids, were observed across

all three simulations: R592-N744 and K444-E858 (Fig. 4A).

To test if the folded plakin domain structure can be unfolded using pulling forces, we performed

steered molecular dynamics (SMD) simulations (Video S1). In the SMD simulations, we fixed the

position of the N-terminal residues and pulled the C-terminal regions (residues 919-950) using a

constant force. Our simulations showed that the plakin domain's long arm and the short arm

separated due to the pulling force without secondary structure unfolding (Fig. 4D). Consequently,

the plakin domain transitioned from the initial closed (Fig. 4D, left) to the final open (Fig. 4D, right)

conformation (Video S1). In the open conformation, the distance between the plakin domain's C-

terminus and N-terminus increased by 30~33 nm compared to the closed structure (Fig. 4C),

which is consistent with the change in DPC-DPC distance observed in the cell-based experiments

(Fig. 3D).

Desmosome adhesion is enhanced when DP is elongated: To examine if elongated DP

influences desmosome adhesion, we assessed the intercellular adhesion strength in cell sheets

using a dispase assay [26]. In our assay, confluent sheets of cells were detached using dispase

II, a protease that gently cleaves cells from the substrate. Since cell-cell adhesion is mediated via

both Ca^{2+}-dependent classical cadherins (like E-cadherin) [27] and desmosomes which adhere in

both a Ca^{2+}-independent state and a Ca^{2+} dependent mode [6, 28], we incubated the lifted cell

sheets with 4 mM EGTA, a chelating agent for Ca^{2+}, for 1 hour. Consequently, our assay

selectively evaluates intercellular adhesion mediated mainly through the Ca^{2+}-independent

desmosomes. Following the EGTA treatment, the cell sheets were subjected to mechanical

stress, resulting in the fragmentation of the sheets, as shown in Fig. 3E. Our results showed that

the K19-KO cell sheets exhibited a significantly greater number of fragments than WT cell sheets.

In contrast, K19-GFP cell sheets generated a similar number of fragments compared to WT cell

sheets (Fig. 3F). These results indicate that transition of DP from a closed to an open

conformation does not compromise intercellular adhesion but instead strengthens adhesion.

Discussion

Our results demonstrate that, in cells, DP senses mechanical forces that are exerted on

desmosomes. Using super-resolution microscopy, pharmacological inhibition, and co-IPs, we

show that mechanical force generated by the actomyosin cytoskeleton is directed to the

desmosome along the KIF network and elongates DP. Atomistic computer simulations show that

the elongation of DP is due to a structural transition of the N-terminal plakin domain from a closed

to an open conformation. This process is analogous to conformational changes in α-catenin,

which associates with classical cadherins in AJs [29-31]. In the case of α-catenin, tension from

the actin cytoskeleton induces a conformational change that opens its central M-domain and

exposes cryptic binding sites for vinculin and other actin-binding proteins [29-31]. It is likely that

force induced opening of DP also exposes cryptic binding sites for junctional proteins that signal

and remodel the desmosome and intermediate filaments. The force-induced unfolding of DP could

also serve as a damper of mechanical stress that maintains mechanical homeostasis within

tissue.

While desmosomes are known for maintaining the mechanical resilience of tissue, their roles in

force transduction are not known. Previous studies in *C. elegans* have demonstrated the

importance of VAB-10, a spectraplakin family member, in controlling epidermal morphogenesis

[32]. It has been proposed that VAB-10 can change its conformation in response to muscle tension

in *C. elegans*, initiating a signaling cascade that leads to the reorganization of intermediate

filaments and the maturation of junctions [33]. Since DP is part of the spectraplakin family, it has

been proposed that DP may serve a similar role as VAB-10 in mammalian cells [32]. Here, we

report of a potential mechanosensitive function of desmosomes in mammalian cells by showing

that DP undergoes a conformational change in response to changes in cytoskeletal tension.

Using cells with and without K19 and using inhibitors of myosin-II, we show that actomyosin-

generated forces are transmitted to the desmosome via K19 filaments and change desmosome

architecture. To confirm that force induced widening of desmosomes did not occur due to

differences in desmosomal and cytoskeletal protein expression levels in the different cell lines,

we performed a comparative analysis of essential components of the desmosome and

cytoskeleton in both WT and K19-KO cells. The results from immunoblotting showed no significant

differences in the desmosomal cadherin Dsg2, Dsc1, and Dsc2-3 or plaque proteins plakophilin-

1, plakophilin-3, and DP, between the two cell lines (Fig. S1). Likewise, there were no discernable

variations observed in cytoskeletal proteins including keratin-8, which is the type II keratin that

binds to K19, F-actin, and the myosin heavy chain 10 (Fig. S1). Consequently, these results show

that changes in force-induced desmosome organization do not arise from altered protein

expression levels, but instead are a consequence of modified mechanical conditions resulting

from the cooperative interactions between KIF and the actomyosin network.

Earlier studies using FRET-based tension sensors in resting MDCK cells, revealed a small

amount of tension across Dsg2, but not across DP [34, 35]. However, the Dsg2 FRET-based

tension sensor showed a higher tension level in the contracting cardiac myocytes, suggesting that

junctional tension depends on both cell type and the specific cellular environment [34]. To test the

junctional tension in MDCK cells and validate our analysis, we measured DPC-DPC distances in

resting MDCK cells. Our quantitative analysis reveals that the desmosome width in MDCK cells

is comparable to that observed in K19-KO cells implying that DP in MDCK cells indeed does not

undergo conformational changes under low forces (Fig. S9).

Keratin deficiency is known to significantly affect cellular function, leading to impaired intercellular

adhesion and increased invasiveness [6, 36, 37]. In agreement, results of our dispase assay show

that intercellular adhesion strength is considerably reduced in K19-KO cells. It is important to

highlight that our data eliminates the adhesive contribution of Ca^{2+} dependent AJs, since we

chelate out the free Ca^{2+} using EGTA prior to shearing the cell sheets. A previous study which

measured total E-cadherin levels in K19-KO MCF7 cells, reported that E-cadherins are

internalized in these cells, which results in reduced cell-cell adhesion [19]. However, when we

quantitatively compared the E-cadherins expressed on the surface of WT and K19-KO cells using

immunofluorescence, we only found a slight decrease in E-cadherin expression levels (Fig.

S10A). Furthermore, we tested if the 'amount' of desmosomes in the WT and K19-KO cells were

similar. To accomplish this, we measured the desmosome line density (*i.e.* $\frac{desmosome\ length}{cell}$). Line

density is a better metric for quantifying desmosome 'amounts' rather than $\frac{\#\ of\ desmosomes}{cell}$, since

different desmosomes have different lengths. We measured the desmosome line density in WT

and K19-KO cells by measuring the DP length along the cell borders in the respective cell lines.

The results showed a slight decrease in desmosome line density in the K19-KO cells compared

to the WT cells (Fig. S10B). While the desmosome line density in these cell lines were only

marginally different, it is important to recognize that they could also contribute to the impaired

adhesion exhibited by the K19-KO cells. One possible reason for the reduction in desmosome

line density in K19-KO cells is that keratin deficiency has been reported to cause faster

endocytosis of desmosomes through elevated DP phosphorylation [12].

Previous SMD simulations carried out on the long arm of the plakin domain (i.e. the crystal

structure with PDB code 3R6N [25]), show that the SRs individually unfold when subjected to

pulling force, which exposes the SH3 domain [38, 39]. Since our simulations are carried out on

the entire plakin domain, which includes the flexible hinge between the long and short arms, we

capture the transition of the plakin domain from a closed to an open conformation. The

conformational change that we measure is supported by previous SAXS measurements [17].

However the transition of the plakin domain from a closed to an open structure also partially

exposes the cryptic SH3 domain which is known to act as a signal transduction adapter crucial in

cell invasiveness, migration, and actin reorganization [40].

Many disease-causing mutations in DP are found within the plakin domain of the N-terminal

region, affecting residues both hidden within the protein and exposed to the surrounding

environment [41]. Specifically, mutations within the SR7-8 region of the plakin domain have been

linked to arrhythmogenic ventricular cardiomyopathy (AVC) [16, 41]. Importantly, our simulations

predict that the SR7-8 region participates in several key hydrogen bonds with SR5-6, which in

turn stabilizes DP in its closed conformation. This raises the possibility that AVC mutations may

interfere with the force induced structural transition of DP from the closed to the open

conformation.

Materials and Methods

Cell culture and stabilization: MCF7 WT cells (ATCC), MCF7 K19-KO cells, and MCF7 K19-GFP rescued cells were maintained in low-glucose Dulbecco's Modified Eagle's Medium (DMEM) (Gibco) supplemented with 10% Fetal Bovine Serum (FBS) (Life Technologies) and 1% penicillin–streptomycin (Sigma-Aldrich) in 5% CO_2 at 37°C. The rescued cells were additionally treated with 100 ug/ml hygromycin for selection of K19-GFP. MDCK cells (ATCC) were cultured in low glucose DMEM (Gibco) cell culture media with 10% FBS and 1% PSK.

The K19-KO cells were generated using the CRISPR/Cas9 technology as previously described [42].We developed the K19-GFP rescue cell line following protocols described previously [43]. Briefly, K19-GFP was cloned into *K*19-KO cells from pEGFP-K19 plasmid into pLenti CMV Hygro (plasmid #17484. Addgene) using the following primers: forward 5'TCAGTCGACTGGATCCATGGTGAGCAAGGGCGAGG3' and reverse 5'GAAAGCTGGGTCTAGTCAGAGGACCTTGGAGGC3'. Lentiviral supernatants were generated using pLenti-K19-GFP plasmid. Lentiviral supernatants collected 24h post-transfection was used to infect subconfluent K19-KO cells in three sequential 4 h incubations in the presence of 4 ug/ml polybrene (Sigma-Aldrich). Transductants were selected using a culture medium containing 100 ug/ml hygromycin beginning 48h after infection. The K19-GFP rescue cells were further stabilized using fluorescence-activated cell sorting.

Preparation of imaging samples: Approximately 500,000 cells were seeded for all the MCF7 cell lines and approximately 250,000 cells were seeded for MDCK cells on the collagen-coated coverslips (22 mm, No.1.5) in a single well of a six-well plate and were kept in 5% CO_2 at 37°C for 24 h. A 'calcium switch' was then performed using the protocol described below to eliminate cell clusters and maintain a similar level of desmosome maturation across all samples. To perform a calcium switch, the cells were first rinsed with phosphate-buffered saline (PBS) (Gibco) solution

at room temperature, followed by incubation in 4 mM EGTA (Millipore Sigma) and 10% low Ca^{2+}

Fetal Bovine Serum (FBS) in Ca^{2+}-free DMEM for 30 min. Following EGTA treatment, the cells

were switched to a DMEM medium with 2 mM $CaCl_2$ for 24 h to trigger formation of mature

desmosomes.

WT+Bleb cells were treated with 50 µM (±) blebbistatin (Millipore Sigma) for 24 h. The blebbistatin

was dissolved in DMSO and diluted in a complete medium. The control group was prepared with

the same protocol but treated with 0.1% DMSO instead. Next, cells were fixed either with ice-cold

1:1 methanol and acetone for 10 min (for keratin and DP staining) or with 3% PFA and 0.3% Triton

X-100 for 10 min at room temperature (RT) (for F-actin staining). Samples were rinsed with PBS

three times and blocked in 1% Bovine Serum Albumin (BSA) overnight at 4°C. Samples were

then incubated with primary antibody (1:200) for 1 h at RT, followed by incubation with secondary

antibody (1:600) for 30 min at RT. The samples stained with fluorescently labeled phalloidin

(1:400) were incubated with phalloidin for 30 min at RT in the dark. Samples were washed three

times with PBS after each antibody incubation. Lastly, samples were mounted to the glass slides

using the ProLong Diamond Antifade Mountant (Life Technologies) and stored at 4°C in the dark.

Antibodies: The following primary antibodies were used: human anti-DSG2 (MAB947, R&D

systems), rabbit anti-DPC antibody (A303-356A, Bethyl Lab), rabbit anti-DPN antibody (25318-1-

AP, Proteintech), chicken anti-GFP (600-901-215, Rockland), mouse anti-K19 antibody (A53-

B/A2) (Santa Cruz Biotechnology), control mouse IgG (sc-2025) (Santa Cruz Biotechnology), and

mouse anti-actin antibody (66009-1-Ig, Proteintech). The following secondary antibodies are

used: anti-chicken AF-488 (Invitrogen), anti-mouse AF-594 (Invitrogen), and anti-rabbit AF-647

(Invitrogen). Phalloidin AF-568 (Invitrogen) was used for F-actin staining.

STED imaging acquisition and processing: Immunofluorescence images were acquired at RT using a Leica TCS SP8 STED 3X confocal microscope (Leica Microsystems) equipped with a 100-x oil immersion objective (HC PL APO CS2), white light laser for excitation lines between 470-670 nm, and depletion laser lines at 592, 660, and 775 nm. The acquisition was performed using LasX software, followed by deconvolution in Huygens Pro software (Scientific Volume Imaging). Post-acquisition processing, including cropping and resizing, color change in the lookup table, adjustment of brightness and contrast, and inversion of grayscale images, was performed using the ImageJ software.

Confocal image acquisition and processing: Immunofluorescence images were acquired using a Leica Stellaris 5 confocal microscope (Leica Microsystems) equipped with 63x oil immersion objective (HC PL APO CS2). Image acquisition was performed using LasX software, followed by post-processing using ImageJ. During post-processing, images were adjusted with inversion of grayscale, enhanced contrast with 0.3% saturated pixels, and reduced background noise using the despeckle functionality. All images were acquired using the same procedure.

Image analysis: We used custom-made ImageJ macros and Python scripts to quantify the desmosome width along the entire cell-cell border. Since Dsg2 signals colocalize between DP "railroad" tracks, they were used as the reference to indicate desmosome location along the membrane. Briefly, Dsg2 signals were selected as regions of interest (ROIs), and an algorithm was used to determine the midline of the ROIs along the membrane axis and draw a segment along the midline, every 200 nm, to obtain the intensity profile of both Dsg2 and DPC or Dsg2 and

DPN. Then, Bar, an ImageJ Plugin, was used to determine the location of peaks of different

signals in their respective channels along the drawn segment. After obtaining the raw data, we

used custom-made MATLAB code to calculate distances for full desmosome width represented

by DPC-DPC or half desmosome width represented by Dsg2-DPC or Dsg2-DPN. To avoid false

signals resulting from background noise or lines not fully drawn over the DP "railroad" track, we

filtered out the data that showed less or more than two peaks in the DPC signals along the drawn

segment or did not have the Dsg2 signal between the two DPC peaks. Furthermore, in certain

desmosomes, the Dsg2 signals were observed as two peaks, as shown in Fig. S11A-B, implying

that the desmosome had ruptured at the Dsg2 extracellular region. We eliminated these

desmosomes from further analysis and only focused on intact desmosomes with unresolvable

Dsg2 signals, as depicted in Fig. 1F. For the DP line density measurement, we also used a

custom-made ImageJ macro script for the quantitative analysis. Briefly, we first subtracted the

background noise while retaining the DP signals along the cell borders. Then, we skeletonized

these signals into segments to obtain the total amount of DP length along cell borders. Lastly, we

obtained the DP line density by normalizing the total DP length using the number of cells in the

image (*i.e.* $\frac{desmosome\ length}{cell}$).

Co-immunoprecipitation: Co-immunoprecipitation was performed as described previously [44].

Cells grown on tissue culture plates were washed three times with 1X cold PBS. Cells were lysed

at 4°C using ice-cold 1% triton lysis buffer (1% Triton X-100, 40mM HEPES (pH 7.5), 120mM

NaCl, 1mM EDTA, 1mM phenylmethylsulfonyl fluoride, 10mM sodium pyrophosphate, 1 µg/ml

each of chymostatin, leupeptin and pepstatin, 10 µg/ml each of aprotinin and benzamidine, 2

µg/ml antipain, 1mM sodium orthovanadate, 50mM NaF). Cell lysates were centrifuged to remove

cellular debris, and protein concentration was determined using Pierce BCA Protein Assay Kit

and BSA as standards. Cell lysates were pre-cleared using Protein G Plus-Agarose beads (Santa

Cruz Biotechnology) for 40 minutes at 4°C while rotating. While rotating, supernatants were

incubated with the indicated antibody or IgG control for two hours at 4°C. Immuno-complexes

were captured using Protein G Plus-Agarose beads and washed 4X using 1% triton lysis buffer

supplemented with 300 mM NaCl.

Gel electrophoresis and immunoblotting: Cells grown on tissue culture plates were washed

three times with 1X cold PBS. Cells were lysed at 4°C using ice-cold 1% triton lysis buffer (1%

Triton X-100, 40mM HEPES (pH 7.5), 120mM NaCl, 1mM EDTA, 1mM phenylmethylsulfonyl

fluoride, 10mM sodium pyrophosphate, 1 µg/ml each of chymostatin, leupeptin and pepstatin, 10

µg/ml each of aprotinin and benzamidine, 2 µg/ml antipain, 1mM sodium orthovanadate, 50mM

NaF). For immunoblotting, cell lysates were centrifuged at 14,000rpm for 5 minutes at 4°C to

remove cellular debris. Protein concentration was determined using the Pierce BCA Protein Assay

Kit following manufacturer's instructions (WG327385. Thermo Scientific). BSA (Equitech-Bio)

prepared in 1% lysis buffer was used as the standard. Aliquots of protein lysates were resolved

using SDS-PAGE and then transferred to nitrocellulose membranes (0.45um) (Amersham

Protran; Cytiva) and immunoblotted using the indicated antibodies, followed by horseradish

peroxidase-conjugated goat anti-mouse or goat anti-rabbit IgG (Sigma-Aldrich). Western

Lightning Plus-ECL (Waltham, MA) or Amersham ECL Select Western was used as blotting

detection reagent. Signals were detected using iBright™ CL1500 Imaging System, Invitrogen

(Thermo Fisher Scientific). The immunoblot result for comparing K19 levels among the three cell

lines was conducted with a different method, as described previously [45].

Dispase assay: Cells were seeded in the tissue-treated well of a 12-well plate (1000,000

cells/well) and grown for 24 h to form a confluent cell sheet. Then, the cell sheets were detached

using 2.4 U/ml dispase II (Millipore Sigma) buffer for 30 min at 37°C. Detached cell sheets were

treated with 4 mM EGTA for 1h at RT. During the incubation, images of the cell sheet in each well

were obtained using a Leica TCS SP8 confocal microscope with a 10x 0.4 dry objective (HC PL

APO CS2). Since the field of view for each image is small, each well was selected as an ROI. In

the navigation mode, tiles of images were taken individually and stitched together during

acquisition to eventually cover the entire ROI and generate the image of cell sheets in the wells.

After the EGTA incubation, cell sheets were subjected to mechanical stress using an orbital

shaker at 200 rpm for 1 min. Cell fragments in each well were imaged as previously described,

and the images were later used to count the number of fragments.

Statistical analysis: All statistical tests were performed using GraphPad Prism (GraphPad

Software), and all graphs were plotted using custom-made Python code. In the boxplots, the box

represents the 25th and 75th percentiles with the median indicated, and whiskers reach the 1.5

times the interquartile range (IQR), defined as the difference between the 25th and 75th

percentiles. Data points outside the whiskers are shown as outliers. The number of independent

experiments (N) and data points (n) is indicated in each figure legend. Data distribution was first

tested based on Shapiro-Wilk's test. Since most data was not normally distributed, non-parametric

statistic tests were used for the analysis. Data were analyzed using Mann-Whitney's U test. A

multiple-group comparison was performed using the Kruskal-Wallis Test, followed by Dunn's

multiple comparison Test. Statistical tests used are specified in the figure legends. Statistical

significance was represented at the level of *, $P < 0.05$; **, $P < 0.01$; ***, $P<0.001$; ns, $P>0.05$.

Structure preparation and MD simulation: Plakin domain structure (residues 178-950) was predicted using Alphafold2 [24]. Subsequently, the structure was prepared using PDBFixer [46] and missing hydrogens were added under pH 7.0 conditions. The protein N-terminus was designated with "NH3+" while "COO-" was assigned to the C-terminus. MD simulations were conducted on the FARM high-performance computing cluster at University of California, Davis with GROMACS 2022.3 as previously described [47]. The simulations were performed in OPLS-AA/L force field [48] and TIP4P water model with 10 Å radius cut-off for Van der Waals and electrostatic interactions. Electrostatic energy calculations employed the particle mesh Ewald method with a 0.16-grid spacing. At the beginning of the simulation, the plakin domain structure was positioned at the center of a dodecahedral box, ensuring a minimum distance of 1 nm from the boundary for every atom. The box was filled with water molecules and neutralized with charged ions (150 mM NaCl, 4 mM KCl, and 2 mM $CaCl_2$). The system was relaxed with energy minimization and stabilized with equilibration under isothermal-isochoric and isothermal isobaric conditions using a modified Berendsen thermostat and Berendsen barostat. Post-stabilization, a 50 ns MD simulation was performed with 2-fs integration steps maintaining a temperature of 300K using a v-rescale thermostat. Equilibration of the protein structure occurred within 20 ns. The number of hydrogen bonds observed between the long arm and the short arm was calculated using *gmx hbond*.

Constant-force SMD simulations and analysis: The constant force SMD simulations were performed as described previously [47]. The starting structures for the SMD simulations were the last frame of the corresponding MD simulations. The structures were placed at the side of a rectangular box (23 × 70 × 8 nm box, center at [11.5nm, 65nm, 4nm]). The system, containing ~1,600,000 atoms was relaxed and equilibrated under isothermal-isobaric conditions using the same protocol as in the MD simulation. The SMD simulations were performed at 310K

temperature using a Nose-Hoover thermostat [49]. During each SMD simulation, we fixed the C-

terminus of the plakin domain and pulled the N-terminal region (residues 919-950) along the

longest axis of the box with a constant force ~830 pN (500 kJ·mol^{-1}·nm^{-1}). The distances

between protein N-terminal and C-terminal were calculated using *gmx pairdist*.

Acknowledgments: Research in the SS lab was supported by the National Institute of General

Medical Sciences of the National Institutes of Health (R01GM121885). Research in the JSC lab

was supported by the National Institute of General Medical Sciences of the National Institutes of

Health (R15CA267890-01). STED imaging was performed at the UC Davis Advanced Imaging

Facility. FACS was performed at the Flow Cytometry Shared Resource funded by the UC Davis

Comprehensive Cancer Center Support Grant (CCSG) awarded by the National Cancer Institute

(NCI P30CA093373).

Data availability: STED, Co-IP, and MD simulation data are made available in the manuscript

and supporting information.

Conflict of Interest Statement: The authors do not declare any competing interests.

References

1.   Green, K.J. and C.L. Simpson, *Desmosomes: New Perspectives on a Classic.* Journal of

Investigative Dermatology, 2007. **127**(11): p. 2499-2515.

2.   Broussard, J.A., S. Getsios, and K.J. Green, *Desmosome regulation and signaling in*

disease. Cell and Tissue Research, 2015. **360**(3): p. 501-512.

3.   Garrod, D. and M. Chidgey, *Desmosome structure, composition and function.* Biochim

Biophys Acta, 2008. **1778**(3): p. 572-87.

4.   Stahley, S.N. and A.P. Kowalczyk, *Desmosomes in acquired disease.* Cell and Tissue

Research, 2015. **360**(3): p. 439-456.

5.   Holthofer, B., et al., *Structure and function of desmosomes.* Int Rev Cytol, 2007. **264**: p.

65-163.

6.   Hatzfeld, M., R. Keil, and T.M. Magin, *Desmosomes and Intermediate Filaments: Their*

Consequences for Tissue Mechanics. Cold Spring Harbor Perspectives in Biology, 2017.

9(6): p. a029157.

7.   Broussard, J.A., et al., *Scaling up single-cell mechanics to multicellular tissues – the role*

of the intermediate filament–desmosome network. Journal of Cell Science, 2020. **133**(6):

p. jcs228031.

8.   Prechova, M., et al., *Plectin-mediated cytoskeletal crosstalk controls cell tension and*

cohesion in epithelial sheets. Journal of Cell Biology, 2022. **221**(3).

9.   Broussard, J.A., et al., *The desmoplakin-intermediate filament linkage regulates cell*

mechanics. Mol Biol Cell, 2017. **28**(23): p. 3156-3164.

10.   Huen, A.C., et al., *Intermediate filament–membrane attachments function synergistically*

with actin-dependent contacts to regulate intercellular adhesive strength. The Journal of

Cell Biology, 2002. **159**(6): p. 1005-1017.

11.   Bornslaeger, E.A., et al., *Breaking the connection: displacement of the desmosomal*

522   *plaque protein desmoplakin from cell-cell interfaces disrupts anchorage of intermediate*

523   *filament bundles and alters intercellular junction assembly.* The Journal of cell biology,

524   1996. **134**(4): p. 985-1001.

525   12.   Kröger, C., et al., *Keratins control intercellular adhesion involving PKC-α–mediated*

526   *desmoplakin phosphorylation.* Journal of Cell Biology, 2013. **201**(5): p. 681-692.

527   13.   Stappenbeck, T.S., et al., *Phosphorylation of the desmoplakin COOH terminus*

528   *negatively regulates its interaction with keratin intermediate filament networks.* J Biol

529   Chem, 1994. **269**(47): p. 29351-4.

530   14.   Jefferson, J.J., et al., *Structural analysis of the plakin domain of bullous pemphigoid*

531   *antigen1 (BPAG1) suggests that plakins are members of the spectrin superfamily.*

532   Journal of Molecular Biology, 2007. **366**(1): p. 244-257.

533   15.   Smith, E.A. and E. Fuchs, *Defining the interactions between intermediate filaments and*

534   *desmosomes.* Journal of Cell Biology, 1998. **141**(5): p. 1229-1241.

535   16.   Al-Jassar, C., et al., *The nonlinear structure of the desmoplakin plakin domain and the*

536   *effects of cardiomyopathy-linked mutations.* J Mol Biol, 2011. **411**(5): p. 1049-61.

537   17.   Al-Jassar, C., et al., *Hinged Plakin Domains Provide Specialized Degrees of Articulation*

538   *in Envoplakin, Periplakin and Desmoplakin.* PLoS ONE, 2013. **8**(7): p. e69767.

539   18.   Stahley, S.N., et al., *Molecular organization of the desmosome as revealed by direct*

540   *stochastic optical reconstruction microscopy.* Journal of Cell Science, 2016. **129**(15): p.

541   2897-2904.

19.   Alsharif, S., et al., *Keratin 19 maintains E-cadherin localization at the cell surface and*
stabilizes cell-cell adhesion of MCF7 cells. Cell Adhesion & Migration, 2021. **15**(1): p. 1-
17.

20.   Quinlan, R.A., et al., *A rim-and-spoke hypothesis to explain the biomechanical roles for*
cytoplasmic intermediate filament networks. Journal of Cell Science, 2017. **130**(20): p.
3437-3445.

21.   Leube, R.E., M. Moch, and R. Windoffer, *Intermediate filaments and the regulation of*
focal adhesion. Curr Opin Cell Biol, 2015. **32**: p. 13-20.

22.   Fujiwara, S., et al., *Interplay between Solo and keratin filaments is crucial for mechanical*
force-induced stress fiber reinforcement. Mol Biol Cell, 2016. **27**(6): p. 954-66.

23.   Snider, N.T., et al., *Keratin hypersumoylation alters filament dynamics and is a marker*
for human liver disease and keratin mutation. J Biol Chem, 2011. **286**(3): p. 2273-84.

24.   Jumper, J., et al., *Highly accurate protein structure prediction with AlphaFold.* Nature,
2021. **596**(7873): p. 583-589.

25.   Choi, H.-J. and W.I. Weis, *Crystal Structure of a Rigid Four-Spectrin-Repeat Fragment of*
the Human Desmoplakin Plakin Domain. Journal of Molecular Biology, 2011. **409**(5): p.
800-812.

26.   Hobbs, R.P., et al., *The calcium ATPase SERCA2 regulates desmoplakin dynamics and*
intercellular adhesive strength through modulation of PKCα signaling. The FASEB
Journal, 2011. **25**(3): p. 990-1001.

27.   Mendonsa, A.M., T.-Y. Na, and B.M. Gumbiner, *E-cadherin in contact inhibition and*
cancer. Oncogene, 2018. **37**(35): p. 4769-4780.

28.   Thomason, A., Helen, et al., *Desmosomes: adhesive strength and signalling in health*
and disease. Biochemical Journal, 2010. **429**(3): p. 419-433.

29.   Yao, M., et al., *Force-dependent conformational switch of α-catenin controls vinculin*
binding. Nature Communications, 2014. **5**(1).

30.   Kim, T.-J., et al., *Dynamic Visualization of α-Catenin Reveals Rapid, Reversible*
Conformation Switching between Tension States. Current Biology, 2015. **25**(2): p. 218-
224.

31.   Yonemura, S., et al., *α-Catenin as a tension transducer that induces adherens junction*
development. Nature Cell Biology, 2010. **12**(6): p. 533-542.

32.   Gillard, G., et al., *Force Transmission between Three Tissues Controls Bipolar Planar*
Polarity Establishment and Morphogenesis. Curr Biol, 2019. **29**(8): p. 1360-1368 e4.

33.   Zhang, H., et al., *A tension-induced mechanotransduction pathway promotes epithelial*
morphogenesis. Nature, 2011. **471**(7336): p. 99-103.

34.   Baddam, S., et al., *The desmosomal cadherin desmoglein-2 experiences mechanical*
tension as demonstrated by a FRET-based tension biosensor expressed in living cells.
Cells, 2018. **7**(7): p. 66.

35.   Price, A.J., et al., *Mechanical loading of desmosomes depends on the magnitude and*
orientation of external stress. Nature Communications, 2018. **9**(1).

36.   Seltmann, K., et al., *Keratins significantly contribute to cell stiffness and impact invasive*

583   *behavior.* Proceedings of the National Academy of Sciences, 2013. **110**(46): p. 18507-

584   18512.

585   37.   Cheung, J., Kevin, et al., *Collective Invasion in Breast Cancer Requires a Conserved*

586   *Basal Epithelial Program.* Cell, 2013. **155**(7): p. 1639-1651.

587   38.   Daday, C., K. Kolšek, and F. Gräter, *The mechano-sensing role of the unique SH3*

588   *insertion in plakin domains revealed by Molecular Dynamics simulations.* Scientific

589   Reports, 2017. **7**(1).

590   39.   Rief, M., et al., *Single molecule force spectroscopy of spectrin repeats: low unfolding*

591   *forces in helix bundles.* Journal of molecular biology, 1999. **286**(2): p. 553-561.

592   40.   Kurochkina, N. and U. Guha, *SH3 domains: modules of protein–protein interactions.*

593   Biophysical Reviews, 2013. **5**(1): p. 29-39.

594   41.   Ortega, E., et al., *The Structure of the Plakin Domain of Plectin Reveals an Extended*

595   *Rod-like Shape.* Journal of Biological Chemistry, 2016. **291**(36): p. 18643-18662.

596   42.   Sharma, P., et al., *Keratin 19 regulates cell cycle pathway and sensitivity of breast*

597   *cancer cells to CDK inhibitors.* Scientific Reports, 2019. **9**(1).

598   43.   Sharma, P., et al., *Keratin 19 interacts with GSK3beta to regulate its nuclear*

599   *accumulation and degradation of cyclin D3.* Mol Biol Cell, 2021. **32**(21): p. ar21.

600   44.   Fallatah, A., et al., *Keratin 19 binds and regulates cytoplasmic HNRNPK mRNA targets*

601   *in triple-negative breast cancer.* BMC Molecular and Cell Biology, 2023. **24**(1).

602   45.   Shafraz, O., C.M.O. Davis, and S. Sivasankar, *Light-activated BioID – an optically*

603   *activated proximity labeling system to study protein–protein interactions.* Journal of Cell

604   Science, 2023. **136**(19).

605   46.   Eastman, P., et al., *OpenMM 7: Rapid development of high performance algorithms for*

606   *molecular dynamics.* PLoS Comput Biol, 2017. **13**(7): p. e1005659.

607   47.   Xie, B., et al., *Strengthening E-cadherin adhesion via antibody-mediated binding*

608   *stabilization.* Structure, 2024. **32**(2): p. 217-227 e3.

609   48.   Kaminski, G.A., et al., *Evaluation and reparametrization of the OPLS-AA force field for*

610   *proteins via comparison with accurate quantum chemical calculations on peptides.*

611   Journal of Physical Chemistry B, 2001. **105**(28): p. 6474-6487.

612   49.   Lemkul, J.A. and D.R. Bevan, *Assessing the stability of Alzheimer's amyloid protofibrils*

613   *using molecular dynamics.* J Phys Chem B, 2010. **114**(4): p. 1652-60.

Figure Legends

Fig. 1: Keratin-19 influences desmosome architecture. (A) Schematic of desmosome

structure. **(B-D)** Representative STED images of desmosomes showing a characteristic 'railroad

track' pattern in **(B)** MCF7 WT cells, **(C)** MCF7 K19-KO cells, and **(D)** MCF7 K19-GfP cells. The

desmosomes were immunolabeled for DPC (green) and Dsg2 (red). Scale bar: 500 nm. **(E)** The

boxed area from the representative STED images of MCF7 WT cells shows a closer look at an

individual DP railroad track with Dsg2 between the two parallel DP plaques. Scale bar: 200 nm.

(F) Line-scan analysis of DPC and Dsg2 fluorescence intensity (indicated by the dashed line in

E). **(G)** Quantification of DPC-DPC distance from WT and K19-KO cells showing that

desmosomes in WT cells are wider than K19-KO cells. In all boxplots, the box represents the 25th

and 75th percentiles with the median indicated and whiskers reach 1.5 times the interquartile

range (IQR), defined as the difference between the 25th and 75th percentiles. Data points outside

the whiskers are shown as outliers. Number of datapoints (n) = 630 (WT), 505 (K19-KO); Number

of replicates (N) = 3. Mann-Whitney's U test; ***, $P < 0.001$. **(H)** Quantification of DPC-DPC

distance in K19-KO, WT, and K19-GFP cells. Desmosomes in WT and K19-GFP cells are wider

than K19-KO cells. n = 505 (K19-KO), 630 (WT), 716 (K19-GFP); N = 3. Kruskal-Wallis Test,

followed by Dunn's multiple comparison Test; ***, $P < 0.001$.

Fig. 2: Interplay between keratin-19 and actomyosin network determine desmosome width.

(A) Schematic of the interactions between the desmosome, K19 filaments, and actin filaments.

Cortical filaments represent keratin rims and actin belts below the plasma membrane. Radial

filaments represent the keratin filaments that anchor at the desmosome and extend into the cell.

The radial filaments interact with cortical filaments. **(B)** Representative STED images of K19-GFP

rescue cells immunolabeled for K19 (magenta), DP (green), and actin (cyan). The individual

component is shown in grayscale to highlight the structural features. The merged images between

K19 and DP or actin are shown at the right. Scale bar: 500 nm. The images show that K19

circumferential rims colocalize with the DP "railroad" tracks and with cortical actin belts. **(C)** Co-

IP of K19 in WT cells. IP was performed with anti-K19 antibody or IgG control. The Co-Ips show

that K19 interacts with both DP and F-actin. **(D)** Quantification of desmosome widths from WT,

K19-KO, and Blebbistatin-treated WT (WT+Blebb) cells showing that Blebbistatin treatment

reduces the desmosome width to a similar level as in the K19-KO cell line. n = 630 (WT), 505

(K19-KO), 650 (WT+Blebb); N = 3. Kruskal-Wallis Test, followed by Dunn's multiple comparison

Test; ***, $P < 0.001$.

Fig. 3: Desmoplakin undergoes force induced conformational change. (A) Schematic of the desmosome under tension. The N-terminal of DP transitions from a closed conformation to an open conformation. **(B)** Quantification of desmosome half-unit widths (Dsg2-DPC distance) from WT and K19-KO cells. n = 1260 (WT), 1010 (K19-KO); N = 3. Mann-Whitney's U test; ***, P<0.001. The distances between Dsg2 and DPC is significantly greater in the WT compared to the K19-KO cells. **(C)** Quantification of desmosome half-unit widths (Dsg2-DPN distance) from WT and K19-KO cells. n = 692 (WT), 826 (K19-KO); N = 3. Mann-Whitney's U test; ns, P>0.05. The Dsg2-DPN distance in both cell lines are similar. **(D)** The mean values from the results in B and C are summarized in the bar chart to compare the DP length (DPN-DPC distance) between the WT and K19-KO cells. The DP length is 66 nm for the WT and 33 nm for the K19-KO, indicating that DP extends 33 nm in the WT cells. **(E)** Dispase assay after 24 h plating. The confluent cell sheets are treated with 4 mM EGTA for 1 h. The images show the intact cell sheets for WT, K19-KO, and K19-GFP rescued cells before stress and fragmented cell sheets after applying mechanical stress. **(F)** Quantification of the dispase assay from K19-KO, WT, and K19-GFP cells. n = 12 (KO), 12 (WT), 9 (K19-GFP); N = 3. Kruskal-Wallis Test, followed by Dunn's multiple comparison Test; ***, P<0.001. The K19-KO cell sheets show a greater number of fragments compared to WT cell sheets, while K19-GFP cell sheets generate a similar number of fragments as WT. This indicates that elongation of DP strengthens intercellular adhesion in WT and K19-GFP cells.

Fig. 4: Molecular Dynamics simulations reveal force-induced changes in plakin domain conformation. (A) The entire plakin domain structure predicted by Alphafold2, suggesting a U-shape conformation in the absence of force. Two common electrostatic interactions (R592-N744,

and K444-E858) observed in all 3 MD simulations (below) are shown. **(B)** During the MD

simulations, 10 to 15 hydrogen bonds were formed between plakin domain's long arm (SR3-4

and SR5-6) and short arm (SR7-8 and SR8-CT). **(C)** During the SMD simulations, distance

between plakin domain's C-terminus and N-terminus was measured, suggesting the structure has

been elongated by 30~33 nm. This elongation correlates with the distance change experimentally

measured in Fig 3. **(D)** Comparison of the plakin domain structure at the start (left) and at the end

(right) of the SMD simulation. The structures show that the overall plakin domain has been

elongated by the pulling force without significant secondary structure unfolding.

Figure 1

A. DP-C-Terminus(DPC)

Desmoplakin(DP)
Plakophilin(PKP)
Plakoglobin(PG)
Membrane
Desmoglein-2(Dsg2)

DPC-DPC distance

DP-C-Terminus (DPC)

B. DPC/Dsg2
WT

C. KO

D. K19-GFP

E.

F.
DPC
Dsg2
DPC-DPC distance
Intensity (a.u.)
distance (nm)

G.
***
distance (nm)
n = 630   n = 505
WT   KO

H.
***
distance (nm)
n = 505   n = 630   n = 716
KO   WT   K19-GFP

Figure 2

A.

Cortical filaments

Radial filaments

Dsg2
Membrane
PG
PKP
DP
Keratin 19(K19) filament
Actin filament

B.

K19 | DP | Actin | K19/DP | K19/Actin

C.

	Input		IP	
	IgG	K19	IgG	K19

IB:K19

IB:DP

IB:Actin

D.

distance (nm)

***

n = 630   n = 505   n = 650

WT   KO   WT+Blebb

Figure 3

A. Dsg2 Extracellular Region

Dsg2-DPC distance

DSG2-DPN distance

DP-N-Terminus(DPN)

Dsg2
Membrane
PG
PKP
DP
Keratin 19(K19)
filament
Actin filament

DP-C-Terminus(DPC)

B. Dsg2-DPC distance

***

n = 1260   n = 1010

WT   KO

C. Dsg2-DPN distance

n.s.

n = 692   n = 826

WT   KO

D.

Dsg2-DPN
Dsg2-DPC

66 nm

WT

KO

33 nm   Δ = 33 nm

distance (nm)

E.

1 hour 4mM EGTA

WT   KO   K19-GFP

Before Stress

1 minn 200 rpm

F. Dispase assay

***

n = 12

n = 12   n = 9

KO   WT   K19-GFP

Figure 4

A.

SR5-6   SH3   SR3-4

SR7-8   SR8-CT

K444
R592
N744   E858

SR3-4:   178-375
SH3:   444-516
SR5-6:   376-629
Loop: 630-650
SR7-8:   651-883
SR8-CT: 884-950

B. MD

C. SMD

D.

force pulling

~3.5nm

~36.0nm