A single high-zinc activation enhancer can control two genes oriented head-to-head in *C. elegans*

Hanwenheng Liu[1], Brian Earley[1,2], Adelita Mendoza[1,3], Patrick Hunt[1], Sean Teng[1,4], Daniel L.

Schneider[1], Kerry Kornfeld[1,*]

[1] Department of Developmental Biology, Washington University School of Medicine, 660 S.

Euclid Ave, St. Louis, MO 63130, USA

[2] Current address: Blackfoot High School, 870 S. Fisher Avenue, Blackfoot, Idaho 83221, USA

[3] Current address: University of Colorado at Boulder

[4] Current address: Center for Structural Biology, Center for Cancer Research, National Cancer

Institute, Frederick, MD 21702-1201, USA

* Corresponding author. Email: kornfeld@wustl.edu

Running head:

One HZA enhancer controls two head-to-head genes

Keyword:

Zinc metabolism, high-zinc activation element, HZA, enhancer, divergent transcription, *C.*

elegans

Abstract

Enhancers play critical roles in gene expression, but a full understanding of their complex

functions has yet to be defined. The cellular response to excess zinc levels in *C. elegans* requires

the HIZR-1 transcription factor, which binds the high-zinc activation (HZA) enhancer in the

promoters of multiple target genes. Cadmium hijacks the excess zinc response by binding and

activating HIZR-1. By analyzing the genome-wide transcriptional response to excess zinc and

cadmium, we identified two positions in the genome where head-to-head oriented genes are both

induced by metals. In both examples, a single predicted HZA enhancer is positioned between the

two translational start sites. We hypothesized that a single enhancer can control both head-to-

head genes, an arrangement that has not been extensively characterized. To test this hypothesis,

we used CRISPR genome editing to precisely delete the HZA^{mT} enhancer positioned between

mtl-2 and *T08G5.1*, and the HZA^{cF} enhancer positioned between *cdr-1* and *F35E8.10*. In both

examples, deleting the enhancer caused both head-to-head genes to lose metal-activated

transcription, whereas the induction of more distant genes was not strongly affected. These

studies rigorously document that a single HZA enhancer can control two head-to-head genes,

advancing our understanding of the diverse functions of enhancers.

Article Summary

Enhancers are critical for gene expression, but a full understanding their functions has yet to be elucidated. We discovered two positions in the *C. elegans* genome where a pair of head-to-head oriented genes are both transcriptionally activated by the metals zinc and cadmium; in both cases, one high-zinc activation (HZA) enhancer is positioned between the translation start sites. When these HZA enhancers were deleted by genome engineering, both head-to-head genes lost metal-activated transcription. These results demonstrate that a single HZA enhancer can control two head-to-head genes, advancing the understanding of enhancer function.

Introduction

Zinc is essential, but excess zinc is toxic (Coleman 1992; Hambidge 2000; Sandstead 2015;

Zoroddu et al. 2019). Thus, animals evolved homeostatic mechanisms to maintain zinc in an

appropriate concentration range. In *C. elegans*, the HIZR-1 nuclear receptor transcription factor

is the master regulator of high zinc homeostasis. In excess zinc conditions, HIZR-1 directly binds

zinc, translocates to the nucleus, binds the HZA (High Zinc Activation) enhancer, and promotes

transcriptional activation (Roh et al. 2015; Dietrich et al. 2016; Warnhoff et al. 2017; Earley,

Mendoza, et al. 2021). Direct target genes have HZA enhancers, such as *cdf-2* and *ttm-1,* which

encode zinc transporters that reduce cytosolic zinc levels by sequestration in lysosome-related

organelles or excretion into the intestinal lumen, respectively (Davis et al. 2009; Roh et al. 2013;

Roh et al. 2015). Additional direct target genes include the metallothionines, *mtl-1* and *mtl-2*, and

hizr-1 itself (Hall et al. 2012; Roh et al. 2012; Roh et al. 2015; Warnhoff et al. 2017; Mendoza et

al. 2024). Cadmium can directly bind HIZR-1, leading to the activation of genes that contain

HZA enhancers; thus, cadmium-responsive genes like *cdr-1* are also regulated by the HZA

element (Earley, Cubillas, et al. 2021).

Enhancers are *cis*-regulatory DNA sequences that either enhance or repress gene expression

(Banerji et al. 1981; Levine 2010). By recruiting various transcription factors, enhancers drive

cell differentiation and regulate stress response pathways (Panigrahi and O'Malley 2021).

Enhancers function at various locations relative to their target genes. They can be located either

up- or downstream of the target genes, as close as under 1kb or over 100kb away. A single

enhancer may regulate multiple genes (Chen et al. 2013; Mills et al. 2020). Usually, these genes

are encoded on the same DNA strand in a series, and the enhancer is positioned either 5' or 3' of

the series of genes, similar to operons in bacteria (Lercher et al. 2003; Cutter et al. 2009). When

two genes that are encoded on different DNA strands flank a common 5' region, they are called

head-to-head genes. Head-to-head genes are widespread in the *C. elegans* and human genomes;

in these cases, transcription of both genes is initiated from the common 5' region. This is called

"divergent transcription" (Adachi and Lieber 2002; Trinklein et al. 2004; Seila et al. 2009;

Ibrahim et al. 2018). The common 5' regions may contain enhancers that in principle could

regulate one or both transcripts. However, few examples of this situation have been analyzed in

detail. Examples of head-to-head gene pairs that have been analyzed in mice and chordates

indicate that enhancers in the common 5' region regulate one gene or the other, but not both,

which ensures cell type-specific gene expression (Swamynathan and Piatigorsky 2002; Hozumi

et al. 2013). Whether this logic also applies to the HZA enhancer in *C. elegans* is unknown.

The function of the HZA enhancer was previously analyzed using transgenic animals with

extrachromosomal arrays — in this context, the HZA enhancer was demonstrated to be necessary

for the zinc-induced activation of the *mtl-2* gene and sufficient to confer zinc-induced activation

on a *pes-10* basal promoter (Roh et al. 2015). However, the function of endogenous HZA

enhancers has not been reported. Here we used CRISPR genome editing to directly evaluate the

function of endogenous HZA enhancers. We examined two examples of head-to-head genes in

C. elegans that are both activated by cadmium (Earley, Cubillas, et al. 2021). Our results indicate

that a single HZA enhancer can control metal-activated transcription of both head-to-head

oriented genes. Thus, the regulatory function of a single HZA enhancer is divergent and acts on

two direct target genes, elucidating the functional capacity of this important DNA control

element.

Results

<u>Identification of two genomic regions containing adjacent genes that are oriented head-to-head</u>

<u>and regulated by cadmium</u>

Earley et al. (2021) identified many *C. elegans* genes that are activated by cadmium in

wild-type animals. By analyzing the chromosomal locations of these genes, we noticed that a

region of chromosome V (~14,018,911) contains two activated genes oriented head-to-head. One

gene is *mtl-2,* which encodes a *C. elegans* metallothionein protein that has been characterized

extensively for its roles in cadmium and zinc biology (Bofill et al. 2009; Zeitoun-Ghandour et al.

2010; Hall et al. 2012; Essig et al. 2024). The other gene is *T08G5.1,* which encodes an

uncharacterized protein. Roh et al. (2015) reported that the *mtl-2* promoter contains a predicted

HZA enhancer that promotes gene activation by zinc and cadmium. This HZA element, named

"HZA^{mT}" hereafter, is positioned between these two genes: the start codon ATGs of *mtl-2* and

T08G5.1 are 277bp and 288bp from the HZA^{mT} element, respectively (Fig.1A). To quantify the

transcriptional response to cadmium, we exposed WT animals at the L4 stage to 100μM

cadmium for 16 hours, prepared RNA, and determined mRNA levels by quantitative reverse

transcriptase polymerase chain reaction (qRT-PCR) (Earley, Cubillas, et al. 2021). Compared to

control worms not exposed to cadmium, *mtl-2* and *T08G5.1* mRNA levels were significantly

increased by 5.4 log2 fold (~42x) and 7.6 log2 fold (~190x), respectively (Fig.1B-D).

We noticed another example of two genes oriented head-to-head that are both activated by

cadmium on chromosome V (~15,921,253). One gene is *cdr-1*, which was identified and named

after its strong induction by cadmium (Hall et al. 2012). The other gene is *F35E8.10,* which

encodes an uncharacterized protein. By searching the region between these two genes, Roh et al.

(2015) identified a predicted HZA element, named "HZA^{cF}" hereafter: the start codon ATGs of

cdr-1 and *F35E8.10* are 108bp and 908bp from the HZA^cF element, respectively (Fig.1E).

Compared to control worms not exposed to cadmium, *cdr-1* and *F35E8.10* mRNA levels were

significantly increased by 8.1 log2 fold (~270x) and 3.3 log2 fold (~10x), respectively (Fig.1F-

G). For these two genes, the increase in mRNA levels correlated with the distance from the

predicted HZA element (Fig.1H).

<u>*hizr-1* was necessary for cadmium-mediated transcript accumulation of *mtl-2, T08G5.1, cdr-1,*</u>

<u>and *F35E8.10*</u>

Earley et al. (2021) demonstrated that cadmium hijacks the high zinc response by directly

binding and activating HIZR-1. To test the model that HIZR-1 is necessary for activation of

these four genes in response to cadmium, we used qPCR to quantify transcript levels in a strong

loss-of-function mutant, *hizr-1(am286* Q87STOP) (Warnhoff et al. 2017). Indeed, all four genes

were affected. *mtl-2, T08G5.1, F35E8.10,* and *cdr-1* mRNA levels were not significantly

different in *hizr-1* mutants treated with cadmium or untreated. In addition, transcript levels of

mtl-2, *T08G5.1,* and *cdr-1* were significantly lower in *hizr-1* mutants treated with cadmium

compared to wild-type animals treated with cadmium (Fig.1). *F35E8.10* transcript levels were

lower in *hizr-1* mutants treated with cadmium compared to wild-type animals treated with

cadmium, but this trend was not significant with this sample size (Fig.1F). Thus, *hizr-1* is

necessary for the cadmium induced transcript accumulation of these genes.

<u>A single HZA element mediates transcriptional activation of *mtl-2* and *T08G5.1* in response to</u>

<u>metals.</u>

Because HIZR-1 directly binds the HZA element, we hypothesized that the HZA elements

mediated transcript accumulation in response to metal. To evaluate this model, we used

CRISPR/Cas9 to generate the *syb4265* mutation, a 15-bp deletion that removes the HZA^mT

element (Fig.2B; labeled "HZA^{mT}Δ"). Because zinc is the physiological metal that activates

HIZR-1, we first examined the response to excess zinc. WT and *syb4265* animals at the L4 stage

were cultured on noble agar minimal media (NAMM) with either no additional metals (replete

zinc) or 200μM supplemental zinc (excess zinc) for 16 hours, and gene expression levels were

quantified by qPCR. Compared to control worms in zinc replete medium, WT animals in zinc

excess medium displayed *mtl-2* and *T08G5.1* mRNA levels that were significantly increased by

3.7 log2 fold (~13x) and 4.2 log2 fold (~18x), respectively (Fig.2B-C). By contrast, *mtl-2* and

T08G5.1 mRNA levels were not significantly different in *syb4265* mutants cultured in zinc

replete or excess medium. In addition, transcript levels were significantly lower in *syb4265*

mutants treated with excess zinc compared to WT animals treated with excess zinc (Fig.2B-C).

To determine the specificity of the transcriptional response to the deleted HZA^{mT} enhancer,

we examined zinc-regulated genes at distant genomic positions. Similar to WT animals in excess

zinc, *syb4256* mutants in excess zinc displayed significantly higher transcript levels for *mtl-1*,

cdf-2, *hizr-1*, and *cdr-1* and significantly lower transcript levels for *zipt-2.3*, indicating that the

syb4256 mutation specifically affects adjacent, but not distant, genes (Fig.2E-F, S1).

To evaluate the function of the HZA^{mT} enhancer in response to cadmium, we exposed WT

and *syb4256* mutant animals at the L4 stage to NAMM with 0 or 100μM cadmium for 16 hours

and determined mRNA levels by qPCR. In WT animals, both *mtl-2* and *T08G5.1* mRNA levels

were significantly increased by cadmium exposure (Fig.S2A-B). By contrast, *syb4256* mutant

animals displayed much less robust levels of gene activation that were marginally significant for

mtl-2 and not significant for *T08G5.1*. However, similar effects were observed for control genes

positioned far from the mutation, suggesting this strain may have an altered response to cadmium

compared to wild type (Fig.S2C-F).

A single HZA element mediates transcriptional activation of *cdr-1* and *F35E8.10* in response to

metals.

We used CRISPR/Cas9 to delete the HZA element between *cdr-1* and *F35E8.10*; the

syb4134 mutation is a 14-bp deletion that removes 13bp of the HZA^{cF} element (Fig.3A-B;

labeled as "HZA^{cF}Δ"). To evaluate the response of these genes to zinc, we quantified gene

expression levels of the WT and *syb4134* animals as described above. Compared to WT animals

on zinc replete medium, WT animals in zinc excess displayed *cdr-1* mRNA levels that were

significantly increased by 7.5 log2 fold (~180x). *F35E8.10* mRNA levels were increased by 1.7

log2 fold (3.2x), but this trend was not significant with this samples size (Fig.3C-D). By contrast,

in *syb4134* mutants lacking the HZA^{cF} element, *cdr-1* and *F35E8.10* mRNA levels were not

significantly different when treated with excess zinc or untreated. In addition, *cdr-1* transcript

levels were significantly lower in *syb4134* mutants treated with excess zinc compared to

wild-type animals treated with excess zinc (Fig 3D). Thus, the HZA^{cF} element was necessary for

zinc-induced transcription of both genes.

To determine the specificity of the transcriptional response to the deleted HZA^{cF} enhancer,

we examined control genes at distant genomic positions. Similar to WT animals in excess zinc,

syb4134 mutants in excess zinc displayed significantly higher transcript levels for *mtl-1*, *cdf-2*,

hizr-1, *mtl-2*, and *T08G5.1,* and significantly lower transcript levels for *zipt-2.3*, indicating the

syb4134 mutation specifically affects adjacent genes (Fig.3E-F, S3).

To evaluate the effect of the deleted HZA^{cF} enhancer on the response to cadmium, we

analyzed WT and *syb4134* mutant animals as described above. In WT animals, both *cdr-1* and

F35E8.10 mRNA levels were significantly increased by cadmium exposure (Fig.S4A-B). By

contrast, *syb4134* mutant animals displayed much less robust levels of gene activation that were

not significant. For control genes positioned far from the mutation, *mtl-1*, *mtl-2*, and *T08G5.1,*

transcriptional activation by cadmium was similar in WT and *syb4134* mutant animals,

indicating that the HZA mutation affects adjacent, but not distant, genes (Fig.S4C-E).

A deletion of *mtl-2* did not influence transcription of *T08G5.1*

Our results establish that *mtl-2* and *T08G5.1* are regulated by the same HZA enhancer. If

these two genes compete for access to the HZA^{mT} element, then removing one gene might

increase transcription of the other. To evaluate this model, we analyzed *mtl-2(gk125)*, which

deletes a region from 208bp upstream of the *mtl-2* translation start codon (ATG) to 584bp

downstream of the *mtl-2* STOP codon and inserts one adenine (Fig.2A; labeled "*mtl-2Δ*" in

Fig.4) (Hall et al. 2012; The C. elegans Deletion Mutant Consortium 2012). No deletions of

T08G5.1 currently exist. The deleted region includes the predicted TATA box (TATAAAAG)

positioned 47bp upstream of the *mtl-2* ATG. The HZA^{mT} element is not deleted. As expected,

mtl-2 mRNA was not reliably detected in *mtl-2(gk125)* animals. We evaluated the response of

WT and *mtl-2(gk125)* to excess zinc using the aforementioned method. Compared to animals in

zinc replete medium, WT and *mtl-2(gk125)* animals in zinc excess displayed *T08G5.1* mRNA

levels that were significantly increased by 4.6 log2 fold (~24x) and 4.7 log2 fold (~26x),

respectively (Fig 4A). Thus, deletion of *mtl-2* did not affect the transcription of *T08G5.1*.

The *mtl-2(gk125)* deletion results in the ATG start codon of the neighboring gene *srh-308*

becoming 281bp away from the HZA^{mT} element (Fig 2A). To evaluate the possibility that the

HZA^{mT} enhancer now regulates the new adjacent gene, we examined transcription of *srh-308* as

described above. *srh-308* displayed high C_T cycles in WT, indicating a low baseline transcription

activity (Fig.S5E). *srh-308* was not significantly induced by excess zinc in WT or *mtl-2(gk125)*

mutants (Fig.4B). Thus, *srh-308* does not appear to be regulated by the HZA element in this

mutant. One possible explanation for this observation is that the transcription start site of *srh-308*

is deleted in *mtl-2Δ* (Fig.S6) (Gu et al. 2012; Chen et al. 2013; Kruesi et al. 2013; Saito et al.

2013).

Discussion

Based on the analysis of cadmium activated genes reported by Earley et al. (2021), we

identified two examples of adjacent genes that are oriented head-to-head and both activated by

cadmium. In both examples, a predicted HZA enhancer is positioned between the two genes. To

address fundamental questions about the function of these HZA enhancers, we used CRISPR

genome engineering to create small deletions that remove these enhancers from the genome.

While genome editing technology has been used to study *cis*-regulatory elements in many

organisms, including enhancers in *C. elegans*, mice, and mammalian cells, it has not previously

been used to analyze the HZA enhancer (Moorthy and Mitchell 2016; Li et al. 2020; Froehlich et

al. 2021). Deleting the HZA^{mT} enhancer strongly reduced activation of both *mtl-2* and *T08G5.1*

in response to excess zinc and cadmium. Similarly, deleting the HZA^{cF} enhancer strongly

reduced activation of both *F35E8.10* and *cdr-1* in response to excess zinc and cadmium. These

results lead to several conclusions: (1) a single HZA enhancer can regulate two genes oriented

head-to-head, (2) by extension, these enhancers can function in either orientation. Orientation

independence is considered to be a hallmark of enhancers (Banerji et al. 1981); (3) these

enhancers can function at a variable distance; however, the strength of induction is correlated

with distance from the ATG start codon. The *F35E8.10* start codon is 908bp from the HZA^{cF}

enhancer, and the gene is weakly induced; the *mtl-2* and *T08G5.1* start codons are about 280bp

from the HZA^{mT} enhancer, and these genes are induced at an intermediate level; the *cdr-1* start

codon is 108bp from the HZA^{cF} enhancer, and the gene is strongly induced (Fig.1D, H; Fig.S5C-

D). Our analysis using genome engineering to delete endogenous HZA enhancers supports and

extends previous studies of HZA enhancers performed with extrachromosomal arrays (Roh et al.

2015; Earley, Cubillas, et al. 2021).

Enhancers were characterized as orientation-independent when studied on a vector (Banerji

et al. 1981). However, little is understood about the ability of a single enhancer to control two

genes oriented head-to-head. One factor that affects enhancer function is cell type. If the two

head-to-head genes are expressed in different cell types, then the enhancer might exhibit

different cell type-specific functions. Indeed, two groups reported orientation-dependent

enhancer functions in mice and chordate *Ciona intestinalis* in different cell types (Swamynathan

and Piatigorsky 2002; Hozumi et al. 2013). This cell type-specificity may be controlled by other

nearby enhancer elements. In *C. elegans*, the HZA element is often in close proximity to a

GATA element, an intestine-specific enhancer (Roh et al. 2015). At the *mtl-2*/*T08G5.1* locus,

there are two predicted GATA elements flanking the HZA element (Moilanen et al. 1999; Roh et

al. 2015). Another factor that affects enhancer function is chromatin regulators, such as

insulators that block access of RNA polymerase II to one flanking gene (Luan et al. 2022;

Hamamoto et al. 2023). A lack of CTCF homologs in *C. elegans* may explain the ability of the

HZA enhancer to function on both head-to-head genes (Heger et al. 2009).

How does a single enhancer control two genes? A widely supported model for enhancer

function is looping: proteins recruited to the enhancer and the promoter bring the two into a

dynamic enhancer-promoter contact (EPC) (Panigrahi and O'Malley 2021). The dynamic EPC

might be the rate-limiting factor in transcriptional activation of a target gene (Panigrahi et al.

2018). Thus, one possibility is that the HZA enhancer loops to one flanking gene at a time,

forming only one EPC; in this scenario, the two gene promoters might compete for access to the

EPC. To test this model, we deleted the *mtl-2* gene and examined transcription of *T08G5.1*.

Interestingly, transcription of *T08G5.1* was similar in the deletion mutant, indicating that the

EPC may not alternate between the two promoters. Perhaps the EPC interacts with the promoters

of both genes at the same time. MDT-15, a mediator subunit, is a co-activator for HIZR-1, and

may contribute to EPC formation (Shomer et al. 2019).

There are several possible reasons why this orientation of enhancers and promoters may

have evolved: 1) It is beneficial to have co-expression of two functionally related genes in

response to metal stress. If two head-to-head genes both function in the same pathway, a single

enhancer might be an efficient mechanism for mounting a response. 2) The "Sheltered Island

Hypothesis" speculates that functionally unrelated genes remain clustered or nested together

because any changes would be toxic to the essential genes in the cluster (Chen and Stein 2006).

3) One of the head-to-head genes may be a bystander gene that is neither beneficial nor toxic. If

co-regulation of these genes is not deleterious, then this arrangement may have evolved.

Whereas the *mtl-2* gene has been characterized extensively for a function in cadmium resistance,

the function of the *T08G5.1* gene has not been characterized (Almutairi et al. 2024). In a similar

manner, the *cdr-1* gene has been characterized for a function in cadmium resistance, but the

function of the *F35E8.10* gene has not been characterized. Thus, all three models are possible

explanations for this arrangement of genes.

Methods

<u>Worm Handling and Culture</u>

Animals were cultured on nematode growth media (NGM) dishes seeded with *E. coli* OP50

at 20°C (Brenner 1974) unless otherwise noted. We picked three L1-stage larvae onto fresh

NGM dishes weekly to maintain strains.

To minimize precipitation of metals during supplementation experiments, we cultured

animals on noble agar minimal media (NAMM) dishes (Warnhoff et al. 2017; Earley, Cubillas,

et al. 2021). Each NAMM dish contains 0.1% cholesterol and 3.7% noble agar. Zinc replete

NAMM has no supplemental metal. Zinc excess NAMM contains 200μM $ZnSO_4$. Cadmium

NAMM contains 100μM $CdCl_2$.

<u>Strain generation</u>

We used Bristol isolate N2 as wild type (Brenner 1974). The *hizr-1(am286)* strain, WU1958,

was generated by outcrossing the parental strain, WU1500, to N2 three times. The *am286* allele

contains a C259T mutation that changes Q87 to STOP (Warnhoff et al. 2017).

We worked with SunyBiotech (www.sunybiotech.com) to generate CRISPR knockout

strains, PHX4134 and PHX4265. PHX4134 contains *syb4134*, a 15bp deletion that precisely

removes 13bp of the HZA^{cF} element (AAA CAG AAA CTA CA) that is positioned 108bp

upstream of the *cdr-1* translation start site (ATG) (Fig.3). PHX4265 contains *syb4265*, a 15bp

deletion that removes the entire 15bp HZA^{mT} element (TAT CAC AAA CTA GAG) that is

positioned 278bp upstream of the *mtl-2* translation start site (ATG) (Fig.2). The deletions were

validated by Sanger DNA sequencing.

To analyze the *mtl-2(gk125)* mutation, we outcrossed VC128 five times to N2 to obtain the

strain WU946 (The C. elegans Deletion Mutant Consortium 2012). The *gk125* insertion/deletion

removes the region from 208bp upstream of *mtl-2* ATG to 584bp downstream of *mtl-2* STOP

codon (TAA) and inserts a single adenine (Fig.2). The insertion/deletion was validated by Sanger

DNA sequencing. DNA sequencing was performed using GENEWIZ (Azenta Life Sciences,

www.genewiz.com). See Table S2 for oligonucleotide primers.

Data Availability

All strains and primers are available upon request. Supplementary File 1 contains all qPCR data

processing spreadsheet (eg. Organized Data Cd, *hizr-1 am286*, Zn, etc), GraphPad Prism sheets,

and log2 average calculation sheets (for superplot). Corresponding folders to figures: "*hizr-1*

am286" – Fig 1, "Zn" – Fig 2 & 3, Sup Fig S1, S3, & S5, "Cd" – Sup Fig S2, S4, & S5, "*mtl-2*",

"*srh-308*" – Fig 4. Supplementary File 2 contains sequencing files for HZA^{mT} and HZA^{cF} loci.

"*cdr-1*_sequencing.seq" and "*mtl-2*_sequencing.seq" are Sanger sequencing results for the

knockout strains. "*mtl-2 gk125*" folder contains Sanger sequencing results for *gk125* allele

verification. Supplementary File 3 contains sequences of *mtl-2* homologs from *C. breneri*, *C.*

briggsae, and *C. remanei* used for alignment in Fig. 2. Files are made available online at

https://zenodo.org/records/13948425, DOI: 10.5281/zenodo.13948425.

Acknowledgement

We thank Barak Cohen and Mike Nonet for helpful discussions and the *Caenorhabditis* Genetics

Center and SUNY Biotech for providing strains. Brian Egan, Dana Shaw, and Vishnu

Saraswathy provided advice and assistance with qPCR procedures and data analysis.

Funding

Funding was provided by the NIH (grant R01 GM068598 received by K.K. and grant F31

ES030622 received by B.E.).

315   Conflicts of Interest

The authors declare no conflicts of interest.

References

Adachi N, Lieber MR. 2002. Bidirectional Gene Organization. Cell. 109(7):807–809.
doi:10.1016/S0092-8674(02)00758-4.

Almutairi N, Khan N, Harrison-Smith A, Arlt VM, Stürzenbaum SR. 2024. Stage-specific
exposure of *Caenorhabditis elegans* to cadmium identifies unique transcriptomic response
cascades and an uncharacterised cadmium responsive transcript. Metallomics.:mfae016.
doi:10.1093/mtomcs/mfae016.

Banerji J, Rusconi S, Schaffner W. 1981. Expression of a β-globin gene is enhanced by remote
SV40 DNA sequences. Cell. 27(2):299–308. doi:10.1016/0092-8674(81)90413-X.

Bofill R, Orihuela R, Romagosa M, Domènech J, Atrian S, Capdevila M. 2009. *Caenorhabditis*
elegans metallothionein isoform specificity – metal binding abilities and the role of histidine in
CeMT1 and CeMT2. The FEBS Journal. 276(23):7040–7056. doi:10.1111/j.1742-
4658.2009.07417.x.

Brenner S. 1974. The genetics of Caenorhabditis elegans. Genetics. 77(1):71–94.
doi:10.1093/genetics/77.1.71.

Chen N, Stein LD. 2006. Conservation and functional significance of gene topology in the
genome of Caenorhabditis elegans. Genome Res. 16(5):606–617. doi:10.1101/gr.4515306.

Chen RA-J, Down TA, Stempor P, Chen QB, Egelhofer TA, Hillier LW, Jeffers TE, Ahringer J.
2013. The landscape of RNA polymerase II transcription initiation in *C. elegans* reveals
promoter and enhancer architectures. Genome Res. 23(8):1339–1347.
doi:10.1101/gr.153668.112.

Coleman JE. 1992. ZINC PROTEINS: Enzymes, Storage Proteins, Transcription Factors, and
Replication Proteins. Annu Rev Biochem. 61(1):897–946.
doi:10.1146/annurev.bi.61.070192.004341.

Cutter AD, Dey A, Murray RL. 2009. Evolution of the Caenorhabditis elegans Genome.
Molecular Biology and Evolution. 26(6):1199–1234. doi:10.1093/molbev/msp048.

Davis DE, Roh HC, Deshmukh K, Bruinsma JJ, Schneider DL, Guthrie J, Robertson JD,
Kornfeld K. 2009. The Cation Diffusion Facilitator Gene *cdf-2* Mediates Zinc Metabolism in
Caenorhabditis elegans. Genetics. 182(4):1015–1033. doi:10.1534/genetics.109.103614.

Dietrich N, Tan C-H, Cubillas C, Earley BJ, Kornfeld K. 2016. Insights into zinc and cadmium
biology in the nematode Caenorhabditis elegans. Archives of Biochemistry and Biophysics.
611:120–133. doi:10.1016/j.abb.2016.05.021.

Earley BJ, Cubillas C, Warnhoff K, Ahmad R, Alcantar A, Lyon MD, Schneider DL, Kornfeld
K. 2021. Cadmium hijacks the high zinc response by binding and activating the HIZR-1 nuclear
receptor. Proc Natl Acad Sci USA. 118(42):e2022649118. doi:10.1073/pnas.2022649118.

Earley BJ, Mendoza AD, Tan C-H, Kornfeld K. 2021. Zinc homeostasis and signaling in the
roundworm C. elegans. Biochimica et Biophysica Acta (BBA) - Molecular Cell Research.
1868(1):118882. doi:10.1016/j.bbamcr.2020.118882.

Essig YJ, Leszczyszyn OI, Almutairi N, Harrison-Smith A, Blease A, Zeitoun-Ghandour S,
Webb SM, Blindauer CA, Stürzenbaum SR. 2024. Juggling cadmium detoxification and zinc
homeostasis: A division of labour between the two C. elegans metallothioneins. Chemosphere.
350:141021. doi:10.1016/j.chemosphere.2023.141021.

Froehlich JJ, Uyar B, Herzog M, Theil K, Glažar P, Akalin A, Rajewsky N. 2021. Parallel
genetics of regulatory sequences using scalable genome editing in vivo. Cell Reports.
35(2):108988. doi:10.1016/j.celrep.2021.108988.

Gu W, Lee H-C, Chaves D, Youngman EM, Pazour GJ, Conte D, Mello CC. 2012. CapSeq and
CIP-TAP Identify Pol II Start Sites and Reveal Capped Small RNAs as C. elegans piRNA
Precursors. Cell. 151(7):1488–1500. doi:10.1016/j.cell.2012.11.023.

Hall J, Haas KL, Freedman JH. 2012. Role of MTL-1, MTL-2, and CDR-1 in Mediating
Cadmium Sensitivity in Caenorhabditis elegans. Toxicological Sciences. 128(2):418–426.
doi:10.1093/toxsci/kfs166.

Hamamoto K, Umemura Y, Makino S, Fukaya T. 2023. Dynamic interplay between non-coding
enhancer transcription and gene activity in development. Nat Commun. 14(1):826.
doi:10.1038/s41467-023-36485-1.

Hambidge M. 2000. Human Zinc Deficiency. The Journal of Nutrition. 130(5):1344S-1349S.
doi:10.1093/jn/130.5.1344S.

Heger P, Marin B, Schierenberg E. 2009. Loss of the insulator protein CTCF during nematode
evolution. BMC Mol Biol. 10(1):84. doi:10.1186/1471-2199-10-84.

Hozumi A, Yoshida R, Horie T, Sakuma T, Yamamoto T, Sasakura Y. 2013. Enhancer activity
sensitive to the orientation of the gene it regulates in the chordate genome. Developmental
Biology. 375(1):79–91. doi:10.1016/j.ydbio.2012.12.012.

Ibrahim MM, Karabacak A, Glahs A, Kolundzic E, Hirsekorn A, Carda A, Tursun B, Zinzen RP,
Lacadie SA, Ohler U. 2018. Determinants of promoter and enhancer transcription directionality
in metazoans. Nat Commun. 9(1):4472. doi:10.1038/s41467-018-06962-z.

Kruesi WS, Core LJ, Waters CT, Lis JT, Meyer BJ. 2013. Condensin controls recruitment of
RNA polymerase II to achieve nematode X-chromosome dosage compensation. Elife. 2:e00808.
doi:10.7554/eLife.00808.

Lercher MJ, Blumenthal T, Hurst LD. 2003. Coexpression of Neighboring Genes in
Caenorhabditis Elegans Is Mostly Due to Operons and Duplicate Genes. Genome Res.
13(2):238–243. doi:10.1101/gr.553803.

Levine M. 2010. Transcriptional Enhancers in Animal Development and Evolution. Current

Biology. 20(17):R754–R763. doi:10.1016/j.cub.2010.06.070.

Li K, Liu Y, Cao H, Zhang Y, Gu Z, Liu X, Yu A, Kaphle P, Dickerson KE, Ni M, et al. 2020.
Interrogation of enhancer function by enhancer-targeting CRISPR epigenetic editing. Nat
Commun. 11(1):485. doi:10.1038/s41467-020-14362-5.

Luan J, Vermunt MW, Syrett CM, Coté A, Tome JM, Zhang H, Huang A, Luppino JM, Keller
CA, Giardine BM, et al. 2022. CTCF blocks antisense transcription initiation at divergent
promoters. Nat Struct Mol Biol. 29(11):1136–1144. doi:10.1038/s41594-022-00855-y.

Mendoza AD, Dietrich N, Tan C-H, Herrera D, Kasiah J, Payne Z, Cubillas C, Schneider DL,
Kornfeld K. 2024. Lysosome-related organelles contain an expansion compartment that mediates
delivery of zinc transporters to promote homeostasis. Proc Natl Acad Sci USA.
121(7):e2307143121. doi:10.1073/pnas.2307143121.

Mills C, Muruganujan A, Ebert D, Marconett CN, Lewinger JP, Thomas PD, Mi H. 2020.
PEREGRINE: A genome-wide prediction of enhancer to gene relationships supported by
experimental evidence. Prokunina-Olsson L, editor. PLoS ONE. 15(12):e0243791.
doi:10.1371/journal.pone.0243791.

Moilanen LH, Fukushige T, Freedman JH. 1999. Regulation of Metallothionein Gene
Transcription. Journal of Biological Chemistry. 274(42):29655–29665.
doi:10.1074/jbc.274.42.29655.

Moorthy SD, Mitchell JA. 2016. Generating CRISPR/Cas9 Mediated Monoallelic Deletions to
Study Enhancer Function in Mouse Embryonic Stem Cells. JoVE.(110):53552.
doi:10.3791/53552.

Panigrahi A, O'Malley BW. 2021. Mechanisms of enhancer action: the known and the unknown.
Genome Biol. 22(1):108. doi:10.1186/s13059-021-02322-1.

Panigrahi AK, Foulds CE, Lanz RB, Hamilton RA, Yi P, Lonard DM, Tsai M-J, Tsai SY,
O'Malley BW. 2018. SRC-3 Coactivator Governs Dynamic Estrogen-Induced Chromatin
Looping Interactions during Transcription. Molecular Cell. 70(4):679-694.e7.
doi:10.1016/j.molcel.2018.04.014.

Roh HC, Collier S, Deshmukh K, Guthrie J, Robertson JD, Kornfeld K. 2013. ttm-1 Encodes
CDF Transporters That Excrete Zinc from Intestinal Cells of C. elegans and Act in a Parallel
Negative Feedback Circuit That Promotes Homeostasis. Eide DJ, editor. PLoS Genet.
9(5):e1003522. doi:10.1371/journal.pgen.1003522.

Roh HC, Collier S, Guthrie J, Robertson JD, Kornfeld K. 2012. Lysosome-Related Organelles in
Intestinal Cells Are a Zinc Storage Site in C. elegans. Cell Metabolism. 15(1):88–99.
doi:10.1016/j.cmet.2011.12.003.

Roh HC, Dimitrov I, Deshmukh K, Zhao G, Warnhoff K, Cabrera D, Tsai W, Kornfeld K. 2015.
A modular system of DNA enhancer elements mediates tissue-specific activation of transcription
by high dietary zinc in C. elegans. Nucleic Acids Research. 43(2):803–816.

doi:10.1093/nar/gku1360.

Saito TL, Hashimoto S, Gu SG, Morton JJ, Stadler M, Blumenthal T, Fire A, Morishita S. 2013.
The transcription start site landscape of *C. elegans*. Genome Res. 23(8):1348–1361.
doi:10.1101/gr.151571.112.

Sandstead HH. 2015. Zinc. In: Handbook on the Toxicology of Metals. Elsevier. p. 1369–1385.
[accessed 2024 Feb 6]. https://linkinghub.elsevier.com/retrieve/pii/B9780444594532000615.

Seila AC, Core LJ, Lis JT, Sharp PA. 2009. Divergent transcription: A new feature of active
promoters. Cell Cycle. 8(16):2557–2564. doi:10.4161/cc.8.16.9305.

Shomer N, Kadhim AZ, Grants JM, Cheng X, Alhusari D, Bhanshali F, Poon AF-Y, Lee MYY,
Muhuri A, Park JI, et al. 2019. Mediator subunit MDT-15/MED15 and Nuclear Receptor HIZR-
1/HNF4 cooperate to regulate toxic metal stress responses in Caenorhabditis elegans. Blackwell
TK, editor. PLoS Genet. 15(12):e1008508. doi:10.1371/journal.pgen.1008508.

Swamynathan SK, Piatigorsky J. 2002. Orientation-dependent Influence of an Intergenic
Enhancer on the Promoter Activity of the Divergently Transcribed Mouse Shsp/αB-crystallin
andMkbp/HspB2 Genes. Journal of Biological Chemistry. 277(51):49700–49706.
doi:10.1074/jbc.M209700200.

The C. elegans Deletion Mutant Consortium. 2012. Large-Scale Screening for Targeted
Knockouts in the *Caenorhabditis elegans* Genome. G3 Genes|Genomes|Genetics. 2(11):1415–
1425. doi:10.1534/g3.112.003830.

Trinklein ND, Aldred SF, Hartman SJ, Schroeder DI, Otillar RP, Myers RM. 2004. An
Abundance of Bidirectional Promoters in the Human Genome. Genome Res. 14(1):62–66.
doi:10.1101/gr.1982804.

Warnhoff K, Roh HC, Kocsisova Z, Tan C-H, Morrison A, Croswell D, Schneider DL, Kornfeld
K. 2017. The Nuclear Receptor HIZR-1 Uses Zinc as a Ligand to Mediate Homeostasis in
Response to High Zinc. Sengupta P, Tissenbaum H, editors. PLoS Biol. 15(1):e2000094.
doi:10.1371/journal.pbio.2000094.

Zeitoun-Ghandour S, Charnock JM, Hodson ME, Leszczyszyn OI, Blindauer CA, Stürzenbaum
SR. 2010. The two *Caenorhabditis elegans* metallothioneins (CeMT-1 and CeMT-2)
discriminate between essential zinc and toxic cadmium. The FEBS Journal. 277(11):2531–2542.
doi:10.1111/j.1742-4658.2010.07667.x.

Zoroddu MA, Aaseth J, Crisponi G, Medici S, Peana M, Nurchi VM. 2019. The essential metals
for humans: a brief overview. Journal of Inorganic Biochemistry. 195:120–129.
doi:10.1016/j.jinorgbio.2019.03.013.

Fig 1

Fig 2

Fig 3

Figure 1. Two regions of Chromosome V have genes that are oriented head-to-head, induced by

cadmium, and require *hizr-1*. **A, E)** Schematics of regions of Chromosome V are drawn to scale

in base pairs (bp). Numeric chromosome location indicates the first base pair of the HZA. Thick

black line represents DNA; yellow box indicates HZA enhancer; numbers are intervals from

HZA enhancer to ATG initiation codon and end of transcript; arrows above represent pre-mRNA

for indicated genes. Scale bar as shown. **D, H)** Distances from the HZA to the start codons

(shown in panels A and E) plotted against fold change for WT plus cadmium (shown in B, C, F,

and G). **B, C, F, G)** Wild type (WT) and *hizr-1(am286)* mutants at the L4 stage were cultured

with or without 100μM cadmium for 16 hours and analyzed by qPCR. Values for WT and

hizr-1(am286) with no cadmium were set equal to 0, and values with cadmium represent log2

fold change. N = 4 initial biological replicates, but may vary in panels due to outlier removal.

Circles are technical replicates (repeated measurements of the same sample), and triangles are

biological replicates (different samples from different days). Same color denotes the same

experiment trial. Statistical analysis by pairwise one-way ANOVA. Non-significant p-values are

listed for p<0.3; otherwise, "ns". For significant p-values: *<0.05; **<0.01; ***<0.001;

****<0.0001. Error bars represent mean ± 95% confidence intervals. Mean values are listed. All

fold changes are in base 2 logarithm (e.g. 3 fold = 8x). The same approach is used to represent

qPCR data in Figures 2-4. **Methods:** To age-synchronize the animals, we grew each strain on

90mm NGM dishes until the population was recently starved and most animals were adult stage.

We washed the animals off the NGM dishes with M9 buffer into a 14ml falcon tube (M9 buffer:

1ml 1M $MgSO_4$, 3g KH_2PO_4, 6g Na_2HPO_4, 5g NaCl in 1L MilliQ (Stiernagle 2006)). We

centrifuged and washed the worms with M9 buffer three times before one final centrifugation for

a pellet. We added 5ml bleaching solution to the pellet and vortexed for three minutes to release

the eggs (bleaching solution: MilliQ, commercial Clorox bleach, and 1M NaOH at a ratio of

7:2:1). If live worms were still present, we extended vortexing for another 15 seconds. We

washed the eggs with M9 buffer twice and hatched them on 90mm unseeded NGM dishes. After

24 hours, we transferred the hatched L1 larvae onto 90mm seeded NGM dishes and cultured for

28 to 32 hours until most animals were L4 stage. To avoid crowding, which may reduce the

growth rate, we sometimes used two dishes. L4 animals were separated onto NAMM dishes

corresponding to different metal conditions. After 16 hours, we collected the animals for RNA

extraction according to previous publications (Warnhoff et al. 2017; Earley et al. 2021). Briefly,

we washed animals from NAMM dishes to an Eppendorf tube, pelleted to 100μl, and added

400μl TRIzol (Invitrogen Cat# 15596018). We subjected the tube to a liquid nitrogen-37°C

heatblock freeze cycle three times, added an additional 200μl TRIzol and 140μl chloroform

(Sigma-Aldrich Cat#319988), vortexed, and centrifuged. We purified the top aqueous phase

containing crude RNA extract using Qiagen RNeasy Mini Kit (Cat# 74106). For cDNA

synthesis, we used 1μg RNA and followed instruction from Intact Genomics igScript First Strand

cDNA Synthesis Kit (Cat# 4314) or BioRad iScript reverse transcription supermix for RT-qPCR

(Cat# 1708841). We substituted the water with Ambion nuclease-free water (Invitrogen Cat#

AM9937). We performed qPCR using MicroAmp fast optical 96-well reaction plate

(ThermoFisher Cat# 4346906) and BioRad iTaq Universal SYBR Green Supermix (Cat#

1725121) in an Applied Biosystems StepOne Plus thermocycler. The thermocycling program was

95°C for 20 seconds, then 40 cycles of 95°C for 3 seconds, and 60°C for 30 seconds. Primers for

qPCR were designed with the IDT PrimerQuest Tool (www.idtdna.com/PrimerQuest). The

primers are listed in Supplementary Table 2. For each biological replicate, we measured three

technical replicates for each gene (three wells). The results were collected as C_T cycles and

analyzed with $2^{-\Delta\Delta C_T}$ method in Microsoft Excel (Schmittgen and Livak 2008).

GraphPad Prism 9 was used for statistical analysis and graphing. We first transformed the $2^{-\Delta\Delta C_T}$

values of each technical replicate by base 2 logarithm for normal distribution, and then removed

outliers with ROUT method (Q=0.5%). We calculated the biological replicate data points as

averages of the respective outlier-removed technical replicates. We analyzed the biological data

with ANOVA or mixed-effects analysis with Holm-Šídák correction. Changes in expression

levels of any given gene are reported as the average of all biological data in fold change; "3 fold"

means 2^3, or "8 times". We overlaid both technical and biological data into a single SuperPlot in

Adobe Illustrator (Lord et al. 2020).

Figure 2. The HZA^{mT} element is necessary for zinc-induced transcription of both *mtl-2* and

T08G5.1. **A)** Schematic of a region of Chromosome V drawn to scale in base pairs (bp). Thick

black line represents DNA; yellow box indicates HZA enhancer. Arrows above represent

pre-mRNA for indicated genes. Red lines below indicate mutations: *gk125*, a deletion of the

mtl-2 open reading frame; *syb4265*, a 15-bp deletion of the HZA^{mT} (named HZA^{mT}Δ below and

subsequent figures). Scale bar: 200bp. **B)** Alignment of the HZA^{mT} elements and their flanking 6

bp in different *Caenorhabditis* species and the *syb4265* deletion strain. **C-E)** Wild type (WT) and

syb4265 mutants at the L4 stage were cultured with replete (-) or 200μM supplemental zinc (+)

for 16 hours and analyzed by qPCR. N = 15 initial biological replicates, but may vary in panels

due to outlier removal.

Figure 3 The HZA^{cF} element is necessary for zinc-induced transcription of both *F35E8.10* and

cdr-1. **A)** Schematic of a region of Chromosome V drawn to scale in base pairs (bp). Thick black

line represents DNA; yellow box indicates HZA enhancer. Arrows above represent pre-mRNA

for indicated genes. Red line below indicates the *syb4134* mutation, a 14-bp deletion of the

HZA^{cF} enhancer (named HZA^{cF}Δ below and subsequent figures). Scale bar: 200bp. **B)** DNA

sequence of the HZA^{cF} element and flanking 6 bp in WT and *syb4134* mutant. **C-E)** Wild type

(WT) and *syb4134* mutants at the L4 stage were cultured with replete (-) or 200μM supplemental

zinc (+) for 16 hours and analyzed by qPCR. N = 15 initial biological replicates, but may vary in

panels due to outlier removal.

Figure 4. A deletion of *mtl-2* did not influence zinc-activated transcription of *T08G5.1*.

A, B) Wild type (WT) and *mtl-2(gk125)* mutants (shown in Figure 2 and denoted as *mtl-2Δ*) at

the L4 stage were cultured with or without 200μM zinc for 16 hours and analyzed by qPCR.

Biological replicates: *T08G5.1* N = 10, *srh-308* N = 3. **C)** Model for the orientation-independent

enhancer function of the HZA element. Within the intestinal cell, HIZR-1 (red) binds to zinc ions

(cyan circles) during zinc excess conditions and enters the nucleus. HIZR-1 binds the HZA

element (yellow) and upregulates both flanking genes in a head-to-head configuration. Black

arrows indicate movements, and red arrows indicate positive transcriptional regulation. Thick

black lines in the nucleus represent chromosome V, with arrows indicating transcription start

sites of genes. Figure not drawn to scale. **D)** Different mode of enhancer function. Top, a single

enhancer controls multiple genes encoded on the same DNA strand. Middle, a single enhancer

controls multiple genes encoded on different DNA strands under different conditions. Bottom,

the HZA enhancer controls head-to-head genes simultaneously under the same condition.