A single high-zinc activation enhancer can control two genes orientated head-to-head in *C.*

elegans

Hanwenheng Liu[1], Brian Earley[1,2], Adelita Mendoza[1,3], Patrick Hunt[1], Sean Teng[1,4], Daniel L.

Schneider[1], Kerry Kornfeld[1,*]

[1] Department of Developmental Biology, Washington University School of Medicine, 660 S.

Euclid Ave, St. Louis, MO 63130, USA

[2] Current address: Blackfoot High School, 870 S. Fisher Avenue, Blackfoot, Idaho 83221, USA

[3] Current address: University of Colorado at Boulder, 3415 Colorado Ave, Boulder, CO 80303,

USA

[4] Current address: Center for Structural Biology, Center for Cancer Research, National Cancer

Institute, Frederick, MD 21702-1201, USA

* Corresponding author. Email: kornfeld@wustl.edu

Running head:

One HZA enhancer controls two head-to-head genes

Keyword:

Zinc metabolism, high-zinc activation element, HZA, enhancer, divergent transcription, *C.*

elegans

Abstract

Enhancers play critical roles in gene expression, but a full understanding of their complex

functions has yet to be defined. The cellular response to excess zinc levels in *C. elegans* requires

the HIZR-1 transcription factor, which binds the high-zinc activation (HZA) enhancer in the

promoters of multiple target genes. Cadmium hijacks the excess zinc response by binding and

activating HIZR-1. By analyzing the genome-wide transcriptional response to excess zinc and

cadmium, we identified two positions in the genome where head-to-head oriented genes are both

induced by metals. In both examples, a single predicted HZA enhancer is positioned between the

two translational start sites. We hypothesized that a single enhancer can control both

head-to-head genes, an arrangement that has not been extensively characterized. To test this

hypothesis, we used CRISPR genome editing to precisely delete the HZA^{mT} enhancer positioned

between *mtl-2* and *T08G5.1*; in this mutant, both head-to-head genes display severely reduced

zinc-activated transcription, whereas zinc-activated transcription of more distant genes was not

strongly affected. Deleting the HZA^{cF} enhancer positioned between *cdr-1* and *F35E8.10* caused

both head-to-head genes to display reduced cadmium-activated transcription, whereas

cadmium-activated transcription of more distant genes was not strongly affected. These studies

rigorously document that a single HZA enhancer can control two head-to-head genes, advancing

our understanding of the diverse functions of enhancers.

Article Summary

Enhancers are critical for gene expression, but a full understanding their functions has yet to

be elucidated. We discovered two positions in the *C. elegans* genome where a pair of

head-to-head oriented genes are both transcriptionally activated by excess zinc and/or cadmium;

in both cases, one high-zinc activation (HZA) enhancer is positioned between the translation start

sites. When these HZA enhancers were deleted by genome engineering, both head-to-head genes

lost metal-activated transcription. These results demonstrate that a single HZA enhancer can

control two head-to-head genes, advancing the understanding of enhancer function.

Introduction

Zinc is an essential metal. About 6% of the prokaryotic proteome and 10% of the eukaryotic

proteome requires zinc for structural, catalytic, and signaling functions (Andreini et al. 2009). In

humans, zinc is required for normal functioning of multiple systems, such as the nervous,

immune, and reproductive systems, in addition to growth and development (Coleman 1992;

Hambidge 2000; Sandstead 2015; Zoroddu et al. 2019). Both a lack and an excess of zinc lead to

defects and diseases. For *C. elegans*, both deficient and excess zinc in growth media retard

growth (Davis et al. 2009; Mendoza et al. 2024). In humans, zinc deficiency leads to infantile

morbidity and symptoms shown in acrodermatitis enteropathica, a genetic disorder that affects

dermal, digestive, immune, and reproductive systems; excess zinc is neurotoxic and is one of the

causes for metal fume fever (Aggett 1989; Black 2001; Mezzaroba et al. 2019; Schoofs et al.

2024; Brenner and Keyes 2025). To prevent these adverse effects, organisms have evolved

robust mechanisms to maintain appropriate cellular zinc levels. In *C. elegans*, HIZR-1, a nuclear

receptor, functions as a zinc excess sensor and a transcription factor for zinc-inducible genes.

HIZR-1 senses zinc through direct binding to the ligand-binding domain; HIZR-1 activates

transcription through direct binding to the HZA (**H**igh-**Z**inc **A**ctivation) element via the

DNA-binding domain. The HZA enhancer is present in the promoters of multiple genes activated

by zinc, including genes encoding for zinc exporters (*cdf-2*, *ttm-1*), metallothioneins (*mtl-1*,

mtl-2), and itself (*hizr-1*) (Roh et al. 2015; Warnhoff et al. 2017). Increased expression of these

proteins expels, sequesters, or chelates excess zinc to restore zinc homeostasis (Davis et al. 2009;

Hall et al. 2012; Roh et al. 2013; Warnhoff et al. 2017; Essig et al. 2024).

Cadmium is a potent environmental heavy-metal toxicant. Cadmium and zinc have chemical

similarities, since both are divalent d-block elements belonging to group 12 of the periodic table.

Exposure to cadmium causes transcriptional changes in stress-response genes and metal chelator

genes (Jin et al. 2003; Lützen et al. 2004; Cui et al. 2007; Li et al. 2015; Nordberg et al. 2015).

Earley et al (2021) showed that cadmium directly binds the HIZR-1 ligand-binding domain,

similar to zinc. While cadmium has been proposed to displace physiological metals and cause

protein dysfunction, in the case of HIZR-1, cadmium promotes nuclear accumulation in intestinal

cells and activation of HIZR-1-dependent transcription via the HZA enhancer. The dramatic

transcriptional response to cadmium exposure that is mediated by HIZR-1 indicates that

cadmium can function as a zinc mimetic to activate the high zinc homeostasis pathway.

Enhancers are *cis*-regulatory DNA sequences that either enhance or repress gene expression

(Banerji et al. 1981; Levine 2010). By recruiting various transcription factors, enhancers drive

cell differentiation and regulate stress response pathways (Panigrahi and O'Malley 2021).

Enhancers function at various locations relative to their target genes. They can be located either

up- or downstream of the target genes, as close as less than 1kb or as far as more than 100kb

away. A single enhancer may regulate multiple genes (Chen et al. 2013; Mills et al. 2020).

Usually, these genes are encoded on the same DNA strand in a series, and the enhancer is

positioned either 5' or 3' of the series of genes, similar to operons in bacteria (Lercher et al.

2003; Cutter et al. 2009). When two genes that are encoded on different DNA strands flank a

common 5' region, they are called head-to-head genes. Head-to-head genes are widespread in the

C. elegans and human genomes; in these cases, transcription of both genes is initiated from the

common 5' region. This is called "divergent transcription" (Adachi and Lieber 2002; Trinklein et

al. 2004; Seila et al. 2009; Ibrahim et al. 2018). The common 5' regions may contain enhancers

that in principle could regulate one or both transcripts. However, few examples of this situation

have been analyzed in detail. Examples of head-to-head gene pairs that have been analyzed in

mice and chordates indicate that enhancers in the common 5' region regulate one gene or the

other, but not both, which ensures cell type-specific gene expression (Swamynathan and

Piatigorsky 2002; Hozumi et al. 2013). Whether this logic also applies to the HZA enhancer in

C. elegans is unknown.

The function of the HZA enhancer was previously analyzed using transgenic animals with

extrachromosomal arrays — in this context, the HZA enhancer was demonstrated to be necessary

for the zinc-induced activation of the *mtl-2* gene and sufficient to confer zinc-induced activation

on a *pes-10* basal promoter (Roh et al. 2015). However, the function of endogenous HZA

enhancers has not been reported. Here we used CRISPR genome editing to directly evaluate the

function of endogenous HZA enhancers. We examined two examples of head-to-head genes in

C. elegans that are both activated by cadmium (Earley et al. 2021). Our results indicate that a

single HZA enhancer can control metal-activated transcription of both head-to-head oriented

genes. Thus, the regulatory function of a single HZA enhancer is divergent and acts on two direct

target genes, elucidating the functional capacity of this important DNA control element.

Methods and materials

<u>Worm Handling and Culture</u>

Animals were cultured on nematode growth media (NGM) dishes seeded with *E. coli* OP50

at 20°C (Brenner 1974) unless otherwise noted. We picked three L1-stage larvae onto fresh

NGM dishes weekly to maintain strains.

To minimize precipitation of metals during supplementation experiments, we cultured

animals on noble agar minimal media (NAMM) dishes (Warnhoff et al. 2017; Earley et al.

2021). Each NAMM dish contains 0.1% cholesterol and 3.7% noble agar. Zinc replete NAMM

has no supplemental metal. Zinc excess NAMM contains 200μM $ZnSO_4$. Cadmium NAMM

contains 100μM $CdCl_2$.

<u>Strain generation</u>

Strains used in this study are listed in Table S1. We used Bristol isolate N2 as WT (Brenner

1974). The *hizr-1(am286)* strain, WU1958, was generated by outcrossing the parental strain,

WU1500, to N2 three times. The *am286* allele contains a C259T mutation that changes Q87 to

STOP (Warnhoff et al. 2017).

We worked with SunyBiotech (www.sunybiotech.com) to generate CRISPR knockout

strains, PHX4134 and PHX4265. PHX4134 contains *syb4134*, a 14bp deletion that removes

14bp of the HZA[cF] element (AAC AGA AAC TAC AA) that is positioned 108bp upstream of the

cdr-1 translation start site (ATG) (Fig. 3). PHX4265 contains *syb4265*, a 15bp deletion that

removes the entire 15bp HZA[mT] element (ATC ACA AAC TAG AGT) that is positioned 278bp

upstream of the *mtl-2* translation start site (ATG) (Fig. 2). We validated the position of these

deletions by Sanger DNA sequencing.

To analyze the *mtl-2(gk125)* mutation, we outcrossed VC128 five times to N2 to obtain the

strain WU946 (The C. elegans Deletion Mutant Consortium 2012). The *gk125* insertion/deletion

removes the region from 208bp upstream of *mtl-2* ATG to 584bp downstream of *mtl-2* STOP

codon (TAA) and inserts a single adenine (Fig. 4). We validated the position of the

insertion/deletion by Sanger DNA sequencing. DNA sequencing was performed using

GENEWIZ (Azenta Life Sciences, www.genewiz.com). See Table S2 for oligonucleotide

primers.

RNA extraction, qPCR, and statistical analysis

To age-synchronize the animals, we grew each strain on 90mm NGM dishes until the

population was recently starved and most animals were adult stage. We washed the animals off

the NGM dishes with M9 buffer into a 14ml falcon tube (M9 buffer: 1ml 1M $MgSO_4$, 3g

KH_2PO_4, 6g Na_2HPO_4, 5g NaCl in 1L MilliQ (Stiernagle 2006)). We centrifuged and washed the

worms with M9 buffer three times before one final centrifugation for a pellet. We added 5ml

bleaching solution to the pellet and vortexed for three minutes to release the eggs (bleaching

solution: MilliQ, commercial Clorox bleach, and 1M NaOH at a ratio of 7:2:1). If live worms

were still present, we extended vortexing for another 15 seconds. We washed the eggs with M9

buffer twice and hatched them on 90mm unseeded NGM dishes. After 24 hours, we transferred

the hatched L1 larvae onto 90mm seeded NGM dishes and cultured for 28 to 32 hours until most

animals were L4 stage. To avoid crowding, which may reduce the growth rate, we sometimes

used two dishes. L4 animals were separated onto NAMM dishes corresponding to different metal

conditions. After 16 hours, we collected the animals for RNA extraction according to previous

publications (Warnhoff et al. 2017; Earley et al. 2021). Briefly, we washed animals from NAMM

dishes to an Eppendorf tube, pelleted to 100µl, and added 400µl TRIzol (Invitrogen Cat#

15596018). We subjected the tube to a liquid nitrogen-37°C heatblock freeze cycle three times,

added an additional 200µl TRIzol and 140µl chloroform (Sigma-Aldrich Cat#319988), vortexed,

and centrifuged. We purified the top aqueous phase containing crude RNA extract using Qiagen

RNeasy Mini Kit (Cat# 74106). For cDNA synthesis, we used 1µg RNA and followed

instruction from Intact Genomics igScript First Strand cDNA Synthesis Kit (Cat# 4314) or

BioRad iScript reverse transcription supermix for RT-qPCR (Cat# 1708841). We substituted the

water with Ambion nuclease-free water (Invitrogen Cat# AM9937). We performed qPCR using

MicroAmp fast optical 96-well reaction plate (ThermoFisher Cat# 4346906) and BioRad iTaq

Universal SYBR Green Supermix (Cat# 1725121) in an Applied Biosystems StepOne Plus

thermocycler. The thermocycling program was 95°C for 20 seconds, then 40 cycles of 95°C for 3

seconds, and 60°C for 30 seconds. Primers for qPCR were designed with the IDT PrimerQuest

Tool (www.idtdna.com/PrimerQuest). The primers are listed in Table S2. For each biological

replicate, we measured three technical replicates for each gene (three wells). The results were

collected as C_T cycles and analyzed with $2^{-\Delta\Delta C_T}$ method in Microsoft Excel (Schmittgen and

Livak 2008).

GraphPad Prism 9 was used for statistical analysis and graphing. We first transformed the

$2^{-\Delta\Delta C_T}$ values of each technical replicate by base 2 logarithm for normal distribution, and then

removed outliers with the ROUT method (Q=0.5%). We calculated the biological replicate data

points as averages of the respective outlier-removed technical replicates. We analyzed the

biological data with ANOVA or mixed-effects analysis with Holm-Šídák correction. Changes in

expression levels of any given gene are reported as the average of all biological data in fold

change; "3 fold" means 2^3, or "8 times". We overlaid both technical and biological data into a

single SuperPlot in Adobe Illustrator (Lord et al. 2020).

Results

<u>Identification of two genomic regions containing adjacent genes that are oriented head-to-head</u>

<u>and regulated by cadmium</u>

Earley et al. (2021) identified many *C. elegans* genes that are activated by cadmium in

wild-type (WT) animals. While analyzing the chromosomal locations of these genes, we noticed

that a region of chromosome V (14,018,240-14,020,360) contains two activated genes that are

oriented head-to-head. One gene is *mtl-2,* which encodes a *C. elegans* metallothionein protein

that has been characterized extensively for its roles in cadmium and zinc biology (Bofill et al.

2009; Zeitoun-Ghandour et al. 2010; Hall et al. 2012; Essig et al. 2024). The other gene is

T08G5.1, which encodes an uncharacterized protein. Roh et al. (2015) reported that the *mtl-2*

promoter contains a predicted HZA enhancer that promotes gene activation by zinc and

cadmium. This HZA element, named "HZA^{mT}" hereafter, is positioned between these two genes:

the start codon ATGs of *mtl-2* and *T08G5.1* are 277bp and 288bp from the HZA^{mT} element,

respectively (Fig. 1A). To quantify the transcriptional response to cadmium, we exposed WT

animals at the L4 stage to 100μM cadmium for 16 hours, prepared RNA, and determined mRNA

levels by quantitative reverse transcriptase polymerase chain reaction (qRT-PCR) (Earley et al.

2021). Compared to control worms not exposed to cadmium, *mtl-2* and *T08G5.1* mRNA levels

were significantly increased by 5.4 log2 fold (~42x) and 7.6 log2 fold (~190x), respectively (Fig.

1B-D).

We noticed another example of two genes that are oriented head-to-head and both activated

by cadmium on chromosome V (15,919,280-15,922,860). One gene is *cdr-1*, which was

identified and named after its strong induction by cadmium (Hall et al. 2012). The other gene is

F35E8.10, which encodes an uncharacterized protein. By searching the region between these two

genes, Roh et al. (2015) identified a predicted HZA element, named "HZA^{cF}" hereafter: the start

codon ATGs of *F35E8.10* and *cdr-1* are 908bp and 108bp from the HZA^{cF} element, respectively

(Fig. 1E). Compared to control worms not exposed to cadmium, *F35E8.10* and *cdr-1* mRNA

levels were significantly increased by 3.3 log2 fold (~10x) and 8.1 log2 fold (~270x),

respectively (Fig. 1F-G). For these two genes, the increase in mRNA levels correlated with the

distance from the predicted HZA element (Fig. 1H).

hizr-1 was necessary for cadmium-mediated transcript accumulation of *mtl-2, T08G5.1, cdr-1,*

and *F35E8.10*

Warnhoff et al. (2017) showed that the HZA-binding transcription factor, HIZR-1, is

required for zinc-induced expression of multiple genes. To test the model that HIZR-1 is

necessary for activation of these four head-to-head genes in response to cadmium, we used qPCR

to quantify transcript levels in a strong loss-of-function mutant, *hizr-1(am286* Q87STOP)

(Warnhoff et al. 2017). Indeed, all four genes were affected. *mtl-2, T08G5.1, F35E8.10,* and

cdr-1 mRNA levels were not significantly different in *hizr-1* mutants treated with cadmium or

untreated. In addition, transcript levels of *mtl-2*, *T08G5.1,* and *cdr-1* were significantly lower in

hizr-1 mutants treated with cadmium compared to WT animals treated with cadmium (Fig. 1).

F35E8.10 transcript levels were lower in *hizr-1* mutants treated with cadmium compared to WT

animals treated with cadmium, but this trend was not significant with this sample size (Fig. 1F).

Thus, there may be a *hizr-1*-independent mechanism to activate transcription of *F35E8.10* in

response to cadmium. These results are consistent with previous reports (Roh et al. 2015; Earley

et al. 2021) and indicate *hizr-1* is necessary, at least in part, for the cadmium-induced

accumulation of the transcripts these genes.

Earley et al (2021) examined the role of *hizr-1* in the response of three of these genes to

excess zinc: *mtl-2*, *T08G5.1,* and *cdr-1*. RNA-seq analysis showed that mRNA accumulation of

these three genes is significantly increased by excess zinc, and this response is strongly reduced

or abrogated in *hizr-1(lf)* mutants (Earley et al. 2021). The function of *hizr-1* in regulating the

level of *F35E8.10* mRNA in response to excess zinc has not been reported.

<u>A single HZA element mediates transcriptional activation of *mtl-2* and *T08G5.1* in response to</u>

<u>zinc.</u>

Because HIZR-1 directly binds the HZA element, we hypothesized that the HZA elements

mediated transcript accumulation in response to metal. To evaluate this model, we used

CRISPR/Cas9 to generate the *syb4265* mutation, a 15-bp deletion that removes the HZA^{mT}

element on chromosome V at position 14,018,911 (Fig. 2B; labeled "HZA$^{mT}\Delta$"). The sequence

and the approximate position of this HZA enhancer in the *mtl-2* promoter have been conserved in

the *Caenorhabditis* genus (Fig. 2B). Because zinc is the physiological metal that activates

HIZR-1, we first examined the response to excess zinc. WT and HZA$^{mT}\Delta$ mutant animals at the

L4 stage were cultured on noble agar minimal media (NAMM) with either no additional metals

(replete zinc) or 200μM supplemental zinc (excess zinc) for 16 hours, and gene expression levels

were quantified by qPCR. Compared to control worms in zinc replete medium, WT animals in

zinc excess medium displayed *mtl-2* and *T08G5.1* mRNA levels that were significantly increased

by 3.7 log2 fold (~13x) and 4.2 log2 fold (~18x), respectively (Fig. 2C-D). By contrast, *mtl-2*

and *T08G5.1* mRNA levels were not significantly different in HZA$^{mT}\Delta$ mutants cultured in zinc

replete or excess medium. In addition, transcript levels were significantly lower in HZA$^{mT}\Delta$

mutants treated with excess zinc compared to WT animals treated with excess zinc (Fig. 2C-D).

Thus, the HZA^{mT} enhancer was necessary for zinc-activated transcription of both genes.

To determine the specificity of the transcriptional response to the deleted HZA^{mT} enhancer,

we examined zinc-regulated genes at distant genomic positions (Fig. 2E-F, S1). Similar to WT

animals in excess zinc, HZA^{mT}Δ mutants in excess zinc displayed significantly higher transcript

levels for two activated genes on chromosome X: *cdf-2* (chrX:12,462,640-12,468,917) and *hizr-1*

(chrX:11,316,278-11,323,086); and two activated genes on chromosome V: *mtl-1*

(chrV:6,689,370-6,693,863) and *cdr-1* (chrV: 15,921,309-15,922,820) (Fig. S1). Furthermore,

zipt-2.3 (chrII:13,710,905-13,717,893) on chromosome II that is repressed by excess zinc was

not significantly affected by this mutation (Fig. 2F). These results indicate that the HZA^{mT}Δ

mutation specifically affected the response to excess zinc in adjacent genes, but not in genes

positioned distantly on the same chromosome or genes on different chromosomes. Thus, neither

the HZA^{mT} enhancer itself, nor the activation of *mtl-2* and *T08G5.1* by excess zinc mediated by

the HZA^{mT} enhancer, was necessary for the robust regulation by excess zinc of five distantly

positioned genes.

To evaluate the function of the HZA^{mT} enhancer in response to cadmium, we exposed WT

and HZA^{mT}Δ mutant animals at the L4 stage to NAMM with 0 or 100μM cadmium for 16 hours

and determined mRNA levels by qPCR. In WT animals, both *mtl-2* and *T08G5.1* mRNA levels

were significantly increased by cadmium exposure (Fig. S2A-B). By contrast, HZA^{mT}Δ mutant

animals displayed much less robust levels of *mtl-2* and *T08G5.1* activation that were not

significant. Thus, the HZA^{mT} enhancer was necessary for full cadmium-activated transcription of

both flanking genes. However, the residual activation of these two genes by cadmium suggests

there may be an activation mechanism that is independent of the HZA^{mT} enhancer.

Unexpectedly, we observed a partial reduction of activation with four control genes positioned

far from the mutation: *cdf-2*, *mtl-1*, *cdr-1,* and *F35E8.10* (Fig. S2C-F). These results indicate that

the HZA^{mT}Δ mutation affected the response to cadmium in adjacent genes, genes positioned

distantly on the same chromosome, and genes on different chromosomes. Thus, either the

HZA^{mT} enhancer itself, or cadmium activation of *mtl-2* and *T08G5.1* mediated by the HZA^{mT}

enhancer, was necessary for the robust regulation by cadmium of four distantly positioned genes.

Another possible interpretation is that a background mutation in the HZA^{mT}Δ strain reduces the

robustness of cadmium-activated transcription throughout the genome.

A single HZA element mediates transcriptional activation of *cdr-1* and *F35E8.10* in response to

cadmium.

We used CRISPR/Cas9 to delete the HZA element between *cdr-1* and *F35E8.10*; the

syb4134 mutation is a 14-bp deletion that removes 14bp of the HZA^{cF} element (Fig. 3A-B;

labeled as "HZA^{cF}Δ"). To evaluate the response of these genes to zinc, we quantified gene

expression levels of the WT and HZA^{cF}Δ animals as described above. Compared to WT animals

on zinc replete medium, WT animals in zinc excess displayed *cdr-1* mRNA levels that were

significantly increased by 7.5 log2 fold (~180x) (Fig. S3B). In HZA^{cF}Δ mutants, *cdr-1* mRNA

levels were not significantly different when treated with excess zinc or untreated. In addition,

cdr-1 transcript levels were significantly lower in HZA^{cF}Δ mutants treated with excess zinc

compared to WT animals treated with excess zinc (Fig. S3B). Thus, the HZA^{cF} element was

necessary for zinc-induced transcription of *cdr-1*. Compared to WT animals on zinc replete

medium, WT animals in zinc excess displayed *F35E8.10* mRNA levels increased by 1.7 log2

fold (3.2x), but this trend was not significant with this sample size (Fig. S3A). In HZA^{cF}Δ

mutants, *F35E8.10* mRNA levels were not significantly different when treated with excess zinc

or untreated. Because excess zinc did not significantly activate *F35E8.10* in WT, we cannot use

the analysis of the HZA^{cF}Δ mutant to determine if the enhancer is necessary.

To determine the specificity of the transcriptional response to the deleted HZA^{cF} enhancer,

we examined control genes at distant genomic positions (Fig. S3C-H). Similar to WT animals in

excess zinc, HZA^{cF}Δ mutants in excess zinc displayed significantly higher transcript levels for

two activated genes on chromosome X: *cdf-2* and *hizr-1*; and three activated genes on

chromosome V: *mtl-1*, *mtl-2* (chrV: 14,018,270-14,018,673), and *T08G5.1*

(chrV:14,018,766-14,020,388). Furthermore, *zipt-2.3* on chromosome II that is repressed by

excess zinc was not significantly affected by this mutation. These results indicate that the

HZA^{cF}Δ mutation specifically affected the response to excess zinc in the adjacent gene *cdr-1*,

whereas it did not affect genes positioned distantly on the same chromosome or genes on

different chromosomes. Thus, neither the HZA^{cF} enhancer itself, nor the activation of *cdr-1* by

excess zinc mediated by the HZA^{cF} enhancer, was necessary for the robust regulation by excess

zinc of six distantly positioned genes.

To evaluate the effect of the deleted HZA^{cF} enhancer on the response to cadmium, we

analyzed WT and HZA^{cF}Δ mutant animals as described above. Compared to control worms

cultured without cadmium, WT animals cultured with cadmium displayed *F35E8.10* and *cdr-1*

mRNA levels that were significantly increased by 4.9 log2 fold (~30x) and 7.2 log2 fold

(~147x), respectively (Fig. 3C-D). By contrast, in HZA^{cF}Δ mutants *F35E8.10* and *cdr-1* did not

display significant induction, indicating the HZA^{cF} element is necessary for full transcriptional

regulation. However, *cdr-1* and *F35E8.10* did display a trend towards weak induction that was

not significant with this sample size, and the level of induction was not significantly different in

WT and HZA^{cF}Δ mutants. These results suggest the existence of an HZA^{cF}-independent

mechanism for cadmium-mediated induction of *cdr-1* and *F35E8.10.* For control genes

positioned far from the mutation on chromosome V, *mtl-1*, *mtl-2*, and *T08G5.1,* transcriptional

activation by cadmium was similar in WT and HZA^{cF}Δ mutant animals (Fig. 3E-G). Thus,

deleting the HZA^{cF} enhancer significantly reduced cadmium activation of both adjacent genes,

whereas it minimally affected cadmium activation of distant genes on the same chromosome.

A deletion of *mtl-2* did not influence transcription of *T08G5.1*

Our results establish that *mtl-2* and *T08G5.1* are regulated by the same HZA enhancer. If

these two genes compete for access to the HZA^{mT} element, then removing one gene might

increase transcription of the other. No deletions of *T08G5.1* currently exist, but we were able to

take advantage of the existing mutation, *mtl-2(gk125)*, which deletes a region from 208bp

upstream of the *mtl-2* translation start codon (ATG) to 584bp downstream of the *mtl-2* STOP

codon and inserts one adenine (Fig. 4A-B; labeled "*mtl-2Δ*") (Hall et al. 2012; The C. elegans

Deletion Mutant Consortium 2012). The deleted region includes the predicted TATA box

(TATAAAAG) and a GATA element (CTGATAA), positioned 47bp and 87bp upstream of *mtl-2*

ATG, respectively (Fig. 4B). GATA elements are tissue-specific enhancers that promote

expression in the intestine and function with the HZA to drive the transcriptional response to

excess zinc in intestinal cells (Moilanen et al. 1999; Roh et al. 2015). The HZA^{mT} element and a

second GATA element are not deleted. As expected, *mtl-2* mRNA was not reliably detected in

mtl-2Δ mutants. We evaluated the response of WT and *mtl-2Δ* mutants to excess zinc as

described above. Compared to animals in zinc-replete medium, WT and *mtl-2Δ* mutants in zinc

excess medium displayed *T08G5.1* mRNA levels that were significantly increased by 4.6 log2

fold (~24x) and 4.7 log2 fold (~26x) (Fig. 4C). Thus, deletion of *mtl-2* did not affect

transcriptional activation of *T08G5.1* in response to zinc excess.

The *mtl-2Δ* deletion results in the ATG start codon of the neighboring gene *srh-308*

becoming 281bp away from the HZA^{mT} element (Fig. 4A). To evaluate the possibility that the

HZA^{mT} enhancer now regulates the new adjacent gene, we examined transcription of *srh-308* as

described above. *srh-308* displayed high C_T cycles in WT, indicating a low baseline level of

transcripts (Fig. S4C). *srh-308* was not significantly induced by excess zinc in WT or *mtl-2Δ*

mutants (Fig. 4D). Thus, *srh-308* does not appear to be regulated by the HZA^{mT} element in this

mutant. One possible explanation for this observation is that the transcription start site of *srh-308*

is deleted in *mtl-2Δ* mutants (Fig. S5) (Gu et al. 2012; Chen et al. 2013; Kruesi et al. 2013; Saito

et al. 2013).

Discussion

Based on the analysis of cadmium activated genes reported by Earley et al. (2021), we

identified two examples of adjacent genes that are oriented head-to-head and both activated by

cadmium. In both examples, a predicted HZA enhancer is positioned between the two genes. To

address fundamental questions about the function of these HZA enhancers, we used CRISPR

genome engineering to create small deletions that remove these enhancers from the genome.

While genome editing technology has been used to study *cis*-regulatory elements in many

organisms, including enhancers in *C. elegans*, mice, and mammalian cells, it has not previously

been used to analyze the HZA enhancer (Moorthy and Mitchell 2016; Li et al. 2020; Froehlich et

al. 2021).

Deleting the HZA$^{\mathrm{mT}}$ enhancer strongly reduced activation of both adjacent genes *mtl-2* and

T08G5.1 in response to excess zinc. By contrast, control genes positioned at a distance from this

deletion that are regulated by excess zinc were minimally affected. These results lead to several

conclusions: (1) within a reasonable distance, a single HZA enhancer can regulate two genes

oriented head-to-head; (2) by extension, this enhancer can function in either orientation (Fig.

4E-F). Orientation independence is considered a hallmark of enhancers (Banerji et al. 1981). It is

well-established biochemically that HIZR-1 binds the HZA and activates transcription (Warnhoff

et al. 2017; Earley et al. 2021). However, our data do not exclude the model that another factor

also binds the HZA and activates transcription. In addition to removing the HZA$^{\mathrm{mT}}$ enhancer,

this 15 bp deletion mutation also changed the spacing, and potentially the topology, of sequence

elements in the promoter (Kouzine et al. 2014). Thus, an alternative explanation for these results

is that the deletion reduced zinc activated transcription as a result of this topology change. We do

not favor this interpretation, because the HIZR-1 transcription factor that binds HZA enhancers

to stimulate transcription is necessary for transcriptional activation of these two genes in

response to excess zinc, indicating the HZA is important (Warnhoff et al. 2017; Earley et al.

2021). Furthermore, a large deletion of *mtl-2* that would change spacing and topology in the

region but preserves the HZA enhancer did not affect induction of *T08G5.1* by excess zinc.

Future experiments that scramble the sequence of the HZA rather than deleting it would better

preserve topology and directly address this alternative model.

Deleting the HZA$^{\text{mT}}$ enhancer had more complex effects on cadmium activated

transcription. While it reduced activation of both adjacent genes *mtl-2* and *T08G5.1*, these genes

still displayed residual activation in response to cadmium. These results indicate the existence of

mechanisms to activate *mtl-2* and *T08G5.1* that function independently of the HZA$^{\text{mT}}$ enhancer.

This is consistent with the results of Earley et al (2021) showing that a subset of

cadmium-activated genes do not require *hizr-1*. Furthermore, cadmium-activated genes

positioned at a distance from this deletion displayed reduced activation in response to cadmium.

We speculate that the enhancer deletion, by reducing the levels of *mtl-2* and/or *T08G5.1*, may

influence cadmium activated transcription of genes throughout the genome. Another possible

model is that the deletion strain contains a background mutation that affects cadmium-activated

transcription. In this case, the background mutation does not appear to affect transcription

regulated by excess zinc.

Deleting the HZA$^{\text{cF}}$ enhancer partially reduced activation of both adjacent genes *cdr-1* and

F35E8.10 in response to cadmium. By contrast, control genes positioned at a distance from this

deletion that are regulated by cadmium were minimally affected. These results support the

conclusions that a single HZA enhancer can regulate two genes that are oriented head-to-head,

and the HZA$^{\text{cF}}$ enhancer can function in either orientation (Fig. 4E-F). As discussed above, this

14 bp deletion mutation also changed the spacing, and potentially the topology, of sequence

elements in the promoter. Thus, an alternative explanation is that the deletion reduced

cadmium-activated transcription as a result of this topology change. It was more difficult to

interpret the results with excess zinc. Activation of *cdr-1* in response to excess zinc was

significantly reduced by deleting the HZA^{cF} enhancer, demonstrating that the enhancer is

necessary for zinc-activated transcription of the adjacent gene *cdr-1*. However, because

F35E8.10 transcripts in WT animals did not accumulate to a significant level in response to

excess zinc, we were unable to evaluate the function of the HZA^{cF} enhancer in the regulation of

F35E8.10. Our analysis using genome engineering to delete endogenous HZA enhancers

supports and extends previous studies of HZA enhancers performed with extrachromosomal

arrays (Roh et al. 2015; Earley et al. 2021).

Enhancers can function at a distance, but their effectiveness may be reduced by increasing

distance (Quintero-Cadena and Sternberg 2016). We noticed a correlation between distance and

magnitude of transcript accumulation in these two pairs of head-to-head genes. The *F35E8.10*

start codon is 908bp from the HZA^{cF} enhancer, and the gene is weakly induced; the *mtl-2* and

T08G5.1 start codons are about 280bp from the HZA^{mT} enhancer, and these genes are induced at

an intermediate level; the *cdr-1* start codon is 108bp from the HZA^{cF} enhancer, and the gene is

strongly induced (Fig. 1D, H; Fig. S4A-B). These correlations do not rigorously establish a

causal effect, and there may be reasons other than distance that dictate the magnitude of the

response. A further caveat is that the two HZA enhancers are different, and they may have

different relationships between distance and magnitude. Our results establish two experimental

systems that can be used to directly investigate the relationship between distance to enhancer and

magnitude of effect. For example, the HZA^{cF} enhancer could be engineered at different positions

in the deletion mutant background to determine the impact on transcription of the head-to-head

genes.

A single enhancer/suppressor can regulate multiple genes on the same strand, referred to as

"enhancer sharing" (Fig. 4F top) (Quintero-Cadena and Sternberg 2016; Mills et al. 2020).

However, little is understood about the ability of a single enhancer to control two genes that are

oriented head-to-head. One factor that affects enhancer function is cell type. If the two

head-to-head genes are expressed in different cell types, then the enhancer might exhibit

different cell type-specific functions (Fig. 4F middle). Indeed, two groups reported

orientation-dependent enhancer functions in mice and chordate *Ciona intestinalis* in different cell

types (Swamynathan and Piatigorsky 2002; Hozumi et al. 2013). This cell type-specificity may

be controlled by other nearby enhancer elements. In *C. elegans*, the HZA element is often in

close proximity to a GATA element, an intestine-specific enhancer (Roh et al. 2015). At the

mtl-2/*T08G5.1* locus, there are two predicted GATA elements flanking the HZA element (Fig.

4B) (Moilanen et al. 1999; Roh et al. 2015). Another factor that affects enhancer function is

chromatin regulators, such as insulators that block access of RNA polymerase II to one flanking

gene (Luan et al. 2022; Hamamoto et al. 2023). A lack of CTCF homologs in *C. elegans* may

explain the ability of the HZA enhancer to function on both head-to-head genes (Heger et al.

2009). Our results indicate that the HZA^{mT} and HZA^{cF} enhancers can regulate two genes in the

same conditions (Fig. 4F, bottom).

How does a single enhancer control two genes? A widely supported model for enhancer

function is looping: proteins recruited to the enhancer and the promoter bring the two into a

dynamic enhancer-promoter contact (EPC) (Panigrahi and O'Malley 2021). The dynamic EPC

might be the rate-limiting factor in transcriptional activation of a target gene (Panigrahi et al.

2018). Thus, one possibility is that the HZA enhancer loops to one flanking gene at a time,

forming only one EPC; in this scenario, the two gene promoters might compete for access to the

EPC. To test this model, we deleted the *mtl-2* gene and examined transcription of *T08G5.1*.

Interestingly, transcription of *T08G5.1* was similar in the deletion mutant, indicating that the

EPC may not alternate between the two promoters. Perhaps the EPC interacts with the promoters

of both genes at the same time. Future experiments analyzing a deletion of *T08G5.1* would

further test this relationship. MDT-15, a mediator subunit, is a co-activator for HIZR-1, and may

contribute to EPC formation (Shomer et al. 2019).

There are several possible reasons why this orientation of enhancers and promoters may

have evolved: 1) It is beneficial to have co-expression of two functionally related genes in

response to metal stress. If two head-to-head genes function in the same pathway, then a single

enhancer might be an efficient mechanism for mounting a response. 2) The "Sheltered Island

Hypothesis" speculates that functionally unrelated genes remain clustered or nested together

because any changes would be toxic to the essential genes in the cluster (Chen and Stein 2006).

3) One of the head-to-head genes may be a bystander gene that is neither beneficial nor toxic. If

co-regulation of these genes is not deleterious, then this arrangement may have evolved.

Whereas the *mtl-2* gene has been characterized extensively for a function in cadmium resistance,

the function of the *T08G5.1* gene has not been characterized (Almutairi et al. 2024; Essig et al.

2024). In a similar manner, the *cdr-1* gene has been characterized for a function in cadmium

resistance, but the function of the *F35E8.10* gene has not been characterized (Hall et al. 2012).

Thus, all three models are possible explanations for this arrangement of genes.

Data Availability

All strains and primers are available upon request. Supplementary File 1 contains all qPCR data

processing spreadsheet (eg. Organized Data Cd, *hizr-1 am286*, Zn, etc), GraphPad Prism sheets,

and log2 average calculation sheets (for superplot). Corresponding folders to figures: "*hizr-1*

am286" – Fig. 1, "Zn" – Fig. 2, Sup Fig. S1, S3, & S4, "Cd" – Fig. 3, Sup Fig. S2 & S4, "*mtl-2*",

"*srh-308*" – Fig. 4. Supplementary File 2 contains sequencing files for HZA^{mT} and HZA^{cF} loci.

"*cdr-1*_sequencing.seq" and "*mtl-2*_sequencing.seq" are Sanger sequencing results for the

knockout strains. "*mtl-2 gk125*" folder contains Sanger sequencing results for *gk125* allele

verification. Supplementary File 3 contains sequences of *mtl-2* homologs from *C. breneri*, *C.*

briggsae, and *C. remanei* used for alignment in Fig. 2. Files are made available online at

https://zenodo.org/records/13948425, DOI: 10.5281/zenodo.13948425.

Acknowledgement

We thank Barak Cohen and Mike Nonet for helpful discussions and the *Caenorhabditis* Genetics

Center and SUNY Biotech for providing strains. Brian Egan, Dana Shaw, and Vishnu

Saraswathy provided advice and assistance with qPCR procedures and data analysis.

Funding

Funding was provided by the NIH (grant R01 GM068598 received by K.K; grant F31 ES030622

received by B.E.; grant 1K99GM146016-01 received by A.M.).

Conflicts of Interest

The authors declare no conflicts of interest.

References

Adachi N, Lieber MR. 2002. Bidirectional Gene Organization. Cell. 109(7):807–809.
doi:10.1016/S0092-8674(02)00758-4.

Aggett PJ. 1989. Severe Zinc Deficiency. In: Mills CF, editor. Zinc in Human Biology. London:
Springer London. (Macdonald I, editor. ILSI Human Nutrition Reviews). p. 259–279. [accessed
2025 Mar 4]. http://link.springer.com/10.1007/978-1-4471-3879-2_17.

Almutairi N, Khan N, Harrison-Smith A, Arlt VM, Stürzenbaum SR. 2024. Stage-specific
exposure of *Caenorhabditis elegans* to cadmium identifies unique transcriptomic response
cascades and an uncharacterised cadmium responsive transcript. Metallomics.:mfae016.
doi:10.1093/mtomcs/mfae016.

Andreini C, Bertini I, Rosato A. 2009. Metalloproteomes: A Bioinformatic Approach. Acc Chem
Res. 42(10):1471–1479. doi:10.1021/ar900015x.

Banerji J, Rusconi S, Schaffner W. 1981. Expression of a β-globin gene is enhanced by remote
SV40 DNA sequences. Cell. 27(2):299–308. doi:10.1016/0092-8674(81)90413-X.

Black RE. 2001. Zinc Deficiency, Immune Function, and Morbidity and Mortality from
Infectious Disease among Children in Developing Countries. Food Nutr Bull. 22(2):155–162.
doi:10.1177/156482650102200205.

Bofill R, Orihuela R, Romagosa M, Domènech J, Atrian S, Capdevila M. 2009. *Caenorhabditis*
elegans metallothionein isoform specificity – metal binding abilities and the role of histidine in
CeMT1 and CeMT2. The FEBS Journal. 276(23):7040–7056. doi:10.1111/j.1742-
4658.2009.07417.x.

Brenner BE, Keyes D. 2025. Metal Fume Fever. In: StatPearls. Treasure Island (FL): StatPearls
Publishing. [accessed 2025 Mar 5]. http://www.ncbi.nlm.nih.gov/books/NBK583199/.

Brenner S. 1974. The genetics of Caenorhabditis elegans. Genetics. 77(1):71–94.
doi:10.1093/genetics/77.1.71.

Chen N, Stein LD. 2006. Conservation and functional significance of gene topology in the
genome of Caenorhabditis elegans. Genome Res. 16(5):606–617. doi:10.1101/gr.4515306.

Chen RA-J, Down TA, Stempor P, Chen QB, Egelhofer TA, Hillier LW, Jeffers TE, Ahringer J.
2013. The landscape of RNA polymerase II transcription initiation in *C. elegans* reveals
promoter and enhancer architectures. Genome Res. 23(8):1339–1347.
doi:10.1101/gr.153668.112.

Coleman JE. 1992. ZINC PROTEINS: Enzymes, Storage Proteins, Transcription Factors, and
Replication Proteins. Annu Rev Biochem. 61(1):897–946.
doi:10.1146/annurev.bi.61.070192.004341.

Cui Y, McBride SJ, Boyd WA, Alper S, Freedman JH. 2007. Toxicogenomic analysis of

Caenorhabditis elegans reveals novel genes and pathways involved in the resistance to cadmium
toxicity. Genome Biol. 8(6):R122. doi:10.1186/gb-2007-8-6-r122.

Cutter AD, Dey A, Murray RL. 2009. Evolution of the Caenorhabditis elegans Genome.
Molecular Biology and Evolution. 26(6):1199–1234. doi:10.1093/molbev/msp048.

Davis DE, Roh HC, Deshmukh K, Bruinsma JJ, Schneider DL, Guthrie J, Robertson JD,
Kornfeld K. 2009. The Cation Diffusion Facilitator Gene *cdf-2* Mediates Zinc Metabolism in
Caenorhabditis elegans. Genetics. 182(4):1015–1033. doi:10.1534/genetics.109.103614.

Earley BJ, Cubillas C, Warnhoff K, Ahmad R, Alcantar A, Lyon MD, Schneider DL, Kornfeld
K. 2021. Cadmium hijacks the high zinc response by binding and activating the HIZR-1 nuclear
receptor. Proc Natl Acad Sci USA. 118(42):e2022649118. doi:10.1073/pnas.2022649118.

Essig YJ, Leszczyszyn OI, Almutairi N, Harrison-Smith A, Blease A, Zeitoun-Ghandour S,
Webb SM, Blindauer CA, Stürzenbaum SR. 2024. Juggling cadmium detoxification and zinc
homeostasis: A division of labour between the two C. elegans metallothioneins. Chemosphere.
350:141021. doi:10.1016/j.chemosphere.2023.141021.

Froehlich JJ, Uyar B, Herzog M, Theil K, Glažar P, Akalin A, Rajewsky N. 2021. Parallel
genetics of regulatory sequences using scalable genome editing in vivo. Cell Reports.
35(2):108988. doi:10.1016/j.celrep.2021.108988.

Gu W, Lee H-C, Chaves D, Youngman EM, Pazour GJ, Conte D, Mello CC. 2012. CapSeq and
CIP-TAP Identify Pol II Start Sites and Reveal Capped Small RNAs as C. elegans piRNA
Precursors. Cell. 151(7):1488–1500. doi:10.1016/j.cell.2012.11.023.

Hall J, Haas KL, Freedman JH. 2012. Role of MTL-1, MTL-2, and CDR-1 in Mediating
Cadmium Sensitivity in Caenorhabditis elegans. Toxicological Sciences. 128(2):418–426.
doi:10.1093/toxsci/kfs166.

Hamamoto K, Umemura Y, Makino S, Fukaya T. 2023. Dynamic interplay between non-coding
enhancer transcription and gene activity in development. Nat Commun. 14(1):826.
doi:10.1038/s41467-023-36485-1.

Hambidge M. 2000. Human Zinc Deficiency. The Journal of Nutrition. 130(5):1344S-1349S.
doi:10.1093/jn/130.5.1344S.

Heger P, Marin B, Schierenberg E. 2009. Loss of the insulator protein CTCF during nematode
evolution. BMC Mol Biol. 10(1):84. doi:10.1186/1471-2199-10-84.

Hozumi A, Yoshida R, Horie T, Sakuma T, Yamamoto T, Sasakura Y. 2013. Enhancer activity
sensitive to the orientation of the gene it regulates in the chordate genome. Developmental
Biology. 375(1):79–91. doi:10.1016/j.ydbio.2012.12.012.

Ibrahim MM, Karabacak A, Glahs A, Kolundzic E, Hirsekorn A, Carda A, Tursun B, Zinzen RP,
Lacadie SA, Ohler U. 2018. Determinants of promoter and enhancer transcription directionality
in metazoans. Nat Commun. 9(1):4472. doi:10.1038/s41467-018-06962-z.

Jin YH, Clark AB, Slebos RJC, Al-Refai H, Taylor JA, Kunkel TA, Resnick MA, Gordenin DA.
2003. Cadmium is a mutagen that acts by inhibiting mismatch repair. Nat Genet. 34(3):326–329.
doi:10.1038/ng1172.

Kouzine F, Levens D, Baranello L. 2014. DNA topology and transcription. Nucleus. 5(3):195–
202. doi:10.4161/nucl.28909.

Kruesi WS, Core LJ, Waters CT, Lis JT, Meyer BJ. 2013. Condensin controls recruitment of
RNA polymerase II to achieve nematode X-chromosome dosage compensation. Elife. 2:e00808.
doi:10.7554/eLife.00808.

Lercher MJ, Blumenthal T, Hurst LD. 2003. Coexpression of Neighboring Genes in
Caenorhabditis Elegans Is Mostly Due to Operons and Duplicate Genes. Genome Res.
13(2):238–243. doi:10.1101/gr.553803.

Levine M. 2010. Transcriptional Enhancers in Animal Development and Evolution. Current
Biology. 20(17):R754–R763. doi:10.1016/j.cub.2010.06.070.

Li K, Liu Y, Cao H, Zhang Y, Gu Z, Liu X, Yu A, Kaphle P, Dickerson KE, Ni M, et al. 2020.
Interrogation of enhancer function by enhancer-targeting CRISPR epigenetic editing. Nat
Commun. 11(1):485. doi:10.1038/s41467-020-14362-5.

Li W, Gu X, Zhang X, Kong J, Ding N, Qi Y, Zhang Y, Wang J, Huang D. 2015. Cadmium
delays non-homologous end joining (NHEJ) repair via inhibition of DNA-PKcs phosphorylation
and downregulation of XRCC4 and Ligase IV. Mutation Research/Fundamental and Molecular
Mechanisms of Mutagenesis. 779:112–123. doi:10.1016/j.mrfmmm.2015.07.002.

Lord SJ, Velle KB, Mullins RD, Fritz-Laylin LK. 2020. SuperPlots: Communicating
reproducibility and variability in cell biology. Journal of Cell Biology. 219(6):e202001064.
doi:10.1083/jcb.202001064.

Luan J, Vermunt MW, Syrett CM, Coté A, Tome JM, Zhang H, Huang A, Luppino JM, Keller
CA, Giardine BM, et al. 2022. CTCF blocks antisense transcription initiation at divergent
promoters. Nat Struct Mol Biol. 29(11):1136–1144. doi:10.1038/s41594-022-00855-y.

Lützen A, Liberti SE, Rasmussen LJ. 2004. Cadmium inhibits human DNA mismatch repair in
vivo. Biochemical and Biophysical Research Communications. 321(1):21–25.
doi:10.1016/j.bbrc.2004.06.102.

Mendoza AD, Dietrich N, Tan C-H, Herrera D, Kasiah J, Payne Z, Cubillas C, Schneider DL,
Kornfeld K. 2024. Lysosome-related organelles contain an expansion compartment that mediates
delivery of zinc transporters to promote homeostasis. Proc Natl Acad Sci USA.
121(7):e2307143121. doi:10.1073/pnas.2307143121.

Mezzaroba L, Alfieri DF, Colado Simão AN, Vissoci Reiche EM. 2019. The role of zinc, copper,
manganese and iron in neurodegenerative diseases. NeuroToxicology. 74:230–241.
doi:10.1016/j.neuro.2019.07.007.

Mills C, Muruganujan A, Ebert D, Marconett CN, Lewinger JP, Thomas PD, Mi H. 2020.
PEREGRINE: A genome-wide prediction of enhancer to gene relationships supported by
experimental evidence. Prokunina-Olsson L, editor. PLoS ONE. 15(12):e0243791.
doi:10.1371/journal.pone.0243791.

Moilanen LH, Fukushige T, Freedman JH. 1999. Regulation of Metallothionein Gene
Transcription. Journal of Biological Chemistry. 274(42):29655–29665.
doi:10.1074/jbc.274.42.29655.

Moorthy SD, Mitchell JA. 2016. Generating CRISPR/Cas9 Mediated Monoallelic Deletions to
Study Enhancer Function in Mouse Embryonic Stem Cells. JoVE.(110):53552.
doi:10.3791/53552.

Nordberg GF, Nogawa K, Nordberg M. 2015. Cadmium. In: Handbook on the Toxicology of
Metals. Elsevier. p. 667–716. [accessed 2024 Feb 6].
https://linkinghub.elsevier.com/retrieve/pii/B9780444594532000329.

Panigrahi A, O'Malley BW. 2021. Mechanisms of enhancer action: the known and the unknown.
Genome Biol. 22(1):108. doi:10.1186/s13059-021-02322-1.

Panigrahi AK, Foulds CE, Lanz RB, Hamilton RA, Yi P, Lonard DM, Tsai M-J, Tsai SY,
O'Malley BW. 2018. SRC-3 Coactivator Governs Dynamic Estrogen-Induced Chromatin
Looping Interactions during Transcription. Molecular Cell. 70(4):679-694.e7.
doi:10.1016/j.molcel.2018.04.014.

Quintero-Cadena P, Sternberg PW. 2016. Enhancer Sharing Promotes Neighborhoods of
Transcriptional Regulation Across Eukaryotes. G3 Genes|Genomes|Genetics. 6(12):4167–4174.
doi:10.1534/g3.116.036228.

Roh HC, Collier S, Deshmukh K, Guthrie J, Robertson JD, Kornfeld K. 2013. ttm-1 Encodes
CDF Transporters That Excrete Zinc from Intestinal Cells of C. elegans and Act in a Parallel
Negative Feedback Circuit That Promotes Homeostasis. Eide DJ, editor. PLoS Genet.
9(5):e1003522. doi:10.1371/journal.pgen.1003522.

Roh HC, Dimitrov I, Deshmukh K, Zhao G, Warnhoff K, Cabrera D, Tsai W, Kornfeld K. 2015.
A modular system of DNA enhancer elements mediates tissue-specific activation of transcription
by high dietary zinc in C. elegans. Nucleic Acids Research. 43(2):803–816.
doi:10.1093/nar/gku1360.

Saito TL, Hashimoto S, Gu SG, Morton JJ, Stadler M, Blumenthal T, Fire A, Morishita S. 2013.
The transcription start site landscape of *C. elegans*. Genome Res. 23(8):1348–1361.
doi:10.1101/gr.151571.112.

Sandstead HH. 2015. Zinc. In: Handbook on the Toxicology of Metals. Elsevier. p. 1369–1385.
[accessed 2024 Feb 6]. https://linkinghub.elsevier.com/retrieve/pii/B9780444594532000615.

Schmittgen TD, Livak KJ. 2008. Analyzing real-time PCR data by the comparative CT method.
Nat Protoc. 3(6):1101–1108. doi:10.1038/nprot.2008.73.

Schoofs H, Schmit J, Rink L. 2024. Zinc Toxicity: Understanding the Limits. Molecules.
29(13):3130. doi:10.3390/molecules29133130.

Seila AC, Core LJ, Lis JT, Sharp PA. 2009. Divergent transcription: A new feature of active
promoters. Cell Cycle. 8(16):2557–2564. doi:10.4161/cc.8.16.9305.

Shomer N, Kadhim AZ, Grants JM, Cheng X, Alhusari D, Bhanshali F, Poon AF-Y, Lee MYY,
Muhuri A, Park JI, et al. 2019. Mediator subunit MDT-15/MED15 and Nuclear Receptor HIZR-
1/HNF4 cooperate to regulate toxic metal stress responses in Caenorhabditis elegans. Blackwell
TK, editor. PLoS Genet. 15(12):e1008508. doi:10.1371/journal.pgen.1008508.

Stiernagle T. 2006. Maintenance of C. elegans. WormBook. doi:10.1895/wormbook.1.101.1.
[accessed 2023 Jun 14].
http://www.wormbook.org/chapters/www_strainmaintain/strainmaintain.html.

Swamynathan SK, Piatigorsky J. 2002. Orientation-dependent Influence of an Intergenic
Enhancer on the Promoter Activity of the Divergently Transcribed Mouse Shsp/αB-crystallin
andMkbp/HspB2 Genes. Journal of Biological Chemistry. 277(51):49700–49706.
doi:10.1074/jbc.M209700200.

The C. elegans Deletion Mutant Consortium. 2012. Large-Scale Screening for Targeted
Knockouts in the *Caenorhabditis elegans* Genome. G3 Genes|Genomes|Genetics. 2(11):1415–
1425. doi:10.1534/g3.112.003830.

Trinklein ND, Aldred SF, Hartman SJ, Schroeder DI, Otillar RP, Myers RM. 2004. An
Abundance of Bidirectional Promoters in the Human Genome. Genome Res. 14(1):62–66.
doi:10.1101/gr.1982804.

Warnhoff K, Roh HC, Kocsisova Z, Tan C-H, Morrison A, Croswell D, Schneider DL, Kornfeld
K. 2017. The Nuclear Receptor HIZR-1 Uses Zinc as a Ligand to Mediate Homeostasis in
Response to High Zinc. Sengupta P, Tissenbaum H, editors. PLoS Biol. 15(1):e2000094.
doi:10.1371/journal.pbio.2000094.

Zeitoun-Ghandour S, Charnock JM, Hodson ME, Leszczyszyn OI, Blindauer CA, Stürzenbaum
SR. 2010. The two *Caenorhabditis elegans* metallothioneins (CeMT-1 and CeMT-2)
discriminate between essential zinc and toxic cadmium. The FEBS Journal. 277(11):2531–2542.
doi:10.1111/j.1742-4658.2010.07667.x.

Zoroddu MA, Aaseth J, Crisponi G, Medici S, Peana M, Nurchi VM. 2019. The essential metals
for humans: a brief overview. Journal of Inorganic Biochemistry. 195:120–129.
doi:10.1016/j.jinorgbio.2019.03.013.

A

F35E8.10   HZA^{cF}   cdr-1

200bp

syb4134
(HZA^{cF}Δ)

B

Species	Gene	Location	Sequence of 5' UTR
C. elegans	cdr-1(+)	-129(+)	aattgaaacagaaactacaataatttt
C. elegans	cdr-1(syb4134)	-129(+)	aattga-------------taatttt

HZA

C F35E8.10

ns

* ns

Fold Change (log2)

0.0 0.0 4.9 1.9

WT   HZA^{cF}Δ   WT   HZA^{cF}Δ
Cd   -   -   +   +

D cdr-1

ns

** 0.059

7.2 2.2

0.0 0.0

WT   HZA^{cF}Δ   WT   HZA^{cF}Δ
Cd   -   -   +   +

E mtl-1

*

** 0.291

6.0 4.3

0.0 0.0

Fold Change (log2)

WT   HZA^{cF}Δ   WT   HZA^{cF}Δ
Cd   -   -   +   +

F mtl-2

**

** ns

6.1 6.3

0.0 0.0

WT   HZA^{cF}Δ   WT   HZA^{cF}Δ
Cd   -   -   +   +

G *T08G5.1*

Figure 1. Two regions of Chromosome V have genes that are oriented head-to-head, activated by

cadmium, and require *hizr-1* for activation. **A, E)** Schematics of regions of Chromosome V are

drawn to scale in base pairs (bp). Numeric chromosome location indicates the first base pair of

the HZA. Thick black line represents DNA; yellow box indicates HZA enhancer; numbers are

intervals from HZA enhancer to ATG start codon and end of transcript; arrows above represent

pre-mRNA for indicated genes. Scale bar as shown. **B, C, F, G)** Wild type (WT) and

hizr-1(am286) mutants at the L4 stage were cultured with or without 100μM cadmium for 16

hours and analyzed by qPCR. Values for WT and *hizr-1(am286)* with no cadmium were set equal

to 0, and values with cadmium represent log2 fold change. N = 4 initial biological replicates but

may vary in panels due to outlier removal. Circles are technical replicates (repeated

measurements of the same sample), and triangles are biological replicates (different samples

from different days). Same color denotes the same experiment trial. Statistical analysis by

pairwise one-way ANOVA. Non-significant p-values are listed for $p < 0.3$; otherwise, "ns". For

significant p-values: $* < 0.05$; $** < 0.01$; $*** < 0.001$; $**** < 0.0001$. Error bars represent mean ±

95% confidence intervals. Mean values are listed. All fold changes are in base 2 logarithm (e.g. 3

folds = 8x). The same approach is used to represent qPCR data in Figures 2-4, S1-S3. **D, H)**

Distances from the HZA to the start codons (shown in panels A and E) plotted against fold

change for WT plus cadmium (shown in B, C, F, and G).

Figure 2. The HZA^{mT} element is necessary for zinc-induced transcription of both *mtl-2* and

T08G5.1. **A)** Schematic of a region of Chromosome V drawn to scale in base pairs (bp). Thick

black line represents DNA; yellow box indicates HZA enhancer. Arrows above represent

pre-mRNA for indicated genes. Red line below indicates mutation *syb4265*, a 15-bp deletion of

the HZA^{mT} (named HZA$^{mT}\Delta$). Scale bar: 100bp. **B)** Alignment of DNA sequences from the 5'

untranslated regions of *mtl-2* genes, containing the conserved HZA^{mT} element and six flanking

base pairs. Sequences are from four WT *Caenorhabditis* species and the *C. elegans syb4265*

deletion strain. Location indicates the number of base pairs upstream from the ATG start codon

and the strand orientation. **C-E)** Wild type (WT) and *syb4265* mutants at the L4 stage were

cultured with replete (-) or 200μM supplemental zinc (+) for 16 hours and analyzed by qPCR. N

= 15 initial biological replicates but may vary in panels due to outlier removal.

Figure 3. The HZA^{cF} element was necessary for full cadmium-induced transcript accumulation

of both *cdr-1* and *F35E8.10*. **A)** Schematic of a region of Chromosome V drawn to scale in base

pairs (bp). Thick black line represents DNA; yellow box indicates HZA enhancer. Arrows above

represent pre-mRNA for indicated genes. Red line below indicates the *syb4134* mutation, a 14-bp

deletion of the HZA^{cF} enhancer (named HZA^{cF}Δ). Scale bar: 200bp. **B)** Alignment of DNA

sequences from the 5' untranslated regions of *cdr-1* that contain the conserved HZA^{cF} element

and six flanking base pairs in WT and *syb4134* mutant. Location indicates the number of base

pairs upstream from the ATG start codon and the strand orientation. **C-G)** Wild type (WT) and

syb4134 mutants at the L4 stage were cultured with (+) or without (-) 100μM cadmium for 16

hours and analyzed by qPCR. N = 6 initial biological replicates, but may vary in panels due to

outlier removal.

Figure 4. A deletion of *mtl-2* did not influence zinc-activated transcription of *T08G5.1*. **A)**

Schematic of a region of Chromosome V drawn to scale in base pairs (bp). Thick black line

represents DNA; yellow box indicates HZA enhancer. Arrows above represent pre-mRNA for

indicated genes. Red line below indicates the extent of the *gk125* deletion that removes the *mtl-2*

open reading frame (labeled *mtl-2Δ*). Scale bar: 200bp. **B)** An enlarged region of panel A is

indicated by dashed lines and shows the positions of predicted control elements: TATA box,

GATA element 1, HZA^{mT}, and GATA element 2. The *gk125* deletion removes the TATA box and

GATA element 1. Scale bar: 100bp. **C, D)** Wild type (WT) and *mtl-2(gk125)* mutants at the L4

stage were cultured with or without 200μM zinc for 16 hours and analyzed by qPCR. Biological

replicates: *T08G5.1* N = 10, *srh-308* N = 3. **E)** Model for the orientation-independent enhancer

function of the HZA element. In intestinal cells, the ligand-binding domain of HIZR-1 (red)

binds to zinc ions (cyan circles) during zinc excess conditions or cadmium ions (green circles)

during cadmium stress (blue arrows). HIZR-1 then enters the nucleus (blue arrows), the DNA-

binding domain binds the HZA element (yellow), and transcription of both flanking genes is

activated (red arrows). Thick black lines in the nucleus represent chromosome V, and thin black

arrows indicate transcription start sites of genes. Not drawn to scale. **F)** Different modes of

enhancer function. Top, a single enhancer controls gene 1 and gene 2, which are encoded on the

same DNA strand. Middle, a single enhancer controls gene 1 in condition 1 or gene 2 in

condition 2, and these head-to-head genes are encoded on different DNA strands. Bottom, a

single HZA enhancer controls gene 1 and gene 2 in the same condition, and these head-to-head

genes are encoded on different DNA strands.